

Contents

Contents	ii
List of Figures	iv
List of Tables	v
1 Preliminaries	1
1.1 Function Spaces of Differential Forms	1
Appendices	4
A Some results from Calculus on $\mathbb{R}^n$	4
A.1 Results from Differentiation	4
A.2 Results from Integration	5
A.2.1 The Riemann Integral	5
A.2.2 Measure Zero and Content Zero	6
A.2.3 Integrable Functions	7
A.2.4 Fubini's Theorem	10
A.3 Partitions of Unity	10
B Differential Forms	11
B.1 Exterior Algebra: Definitions and Results	11
B.1.1 σf as a group action	11
B.1.2 Tensor Product, Alternating Forms and the Wedge Product	13
B.1.3 The Associativity of the Wedge Product	15
B.1.4 Dimension of Alternating k -forms	17
B.2 Tangent vectors in $\mathbb{R}^n$ and Vector Fields	18
B.2.1 The Directional Derivative and Derivations	19
B.3 Differential Forms	20
B.3.1 Differential Forms in terms of Coordinates	21
B.3.2 Differential k -forms	22
B.3.3 Exterior Derivative	23
B.3.4 Closed and Exact Forms	25
B.4 Vector Calculus	25
B.4.1 Exterior Derivative of forms in $\mathbb{R}^3$	25
B.5 Interior Product	27
B.5.1 The Musical Isomorphism and Interior Product	28
B.6 Hodge Star	32
B.6.1 Hodge Star and Interior	33
B.7 Brief Look at Riemannian Manifolds	34
B.7.1 Natural Inner Product on Forms, Hodge Star and Interior Product	35
B.8 Pullback of Forms	37
B.8.1 Pullback of a 1-form	38
B.8.2 Pullback of a k -form	39
B.9 Integration of 1-forms	42
B.10 The Stokes Theorem	42
B.11 The Codifferential δ	43

B.11.1 Motivation	43
B.11.2 Coordinate formulas for the Codifferential	43
B.12 Boundary Calculus and Tangential and Normal Components	45
B.13 Integration Theorems	45
B.14 A little bit of Riemannian Geometry	45
C Sobolev Spaces and Requisite Measure Theory and Functional Analysis	46
D Manifolds	47
D.1 Topological Manifolds	47
Bibliography	48

List of Figures

B.1 Cylinder around a boundary point $p \in \partial\Omega$ from definition B.10.1	42
--	----

List of Tables

B.1 Correspondence between vector fields and forms.	25
---	----

Chapter 1

Preliminaries

WORK IN PROGRESS!!!

Thus far, we have been speaking about C^∞ differential forms, but we can define differential forms in more general spaces.

1.1 Function Spaces of Differential Forms

Throughout this section, let $\Omega \subset \mathbb{R}^n$ be an open set and let $0 \leq k \leq n$. Since Ω is an open subset of $\mathbb{R}^n$, we identify $T_p\Omega \simeq \mathbb{R}^n$ for every $p \in \Omega$. Consequently, every *differential* k -form may be viewed as a map

$$f : \Omega \longrightarrow \Lambda^k(\mathbb{R}^n).$$

With respect to the standard basis of $\Lambda^k(\mathbb{R}^n)$, every k -form admits a unique expansion of the form

$$f = \sum_{I \in \mathcal{T}_k} f^I dx^I = \sum_{1 \leq i_1 < \dots < i_k \leq n} f^{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k},$$

where each coefficient $f^I : \Omega \rightarrow \mathbb{R}$ is a real-valued function.

Remark 1.1.1. Since $\Lambda^k(\mathbb{R}^n)$ is a finite-dimensional vector space of dimension $\binom{n}{k}$, a differential k -form may equivalently be regarded as a $\binom{n}{k}$ -tuple of scalar-valued functions. All regularity notions are defined coefficient-wise.

Definition 1.1.2 (L^p differential forms). Let $1 \leq p \leq \infty$. We write

$$f \in L^p(\Omega; \Lambda^k)$$

if and only if every coefficient function satisfies

$$f^I \in L^p(\Omega), \quad I \in \mathcal{T}_k.$$

Equivalently,

$$L^p(\Omega; \Lambda^k) = L^p(\Omega; \mathbb{R}^{\binom{n}{k}}).$$

The pointwise inner product on $\Lambda^k(\mathbb{R}^n)$ is defined by

$$\langle f, g \rangle = \sum_{I \in \mathcal{T}_k} f^I g^I,$$

with associated pointwise norm

$$|f|^2 = \langle f, f \rangle = \sum_{I \in \mathcal{T}_k} (f^I)^2.$$

The L^p norm of a differential form is obtained by integrating the pointwise norm:

$$\|f\|_{L^p(\Omega; \Lambda^k)} = \left(\int_{\Omega} |f(x)|^p dx \right)^{1/p},$$

with the usual modification when $p = \infty$. In particular,

$$\|f\|_{L^2(\Omega; \Lambda^k)}^2 = \int_{\Omega} \sum_{I \in \mathcal{T}_k} (f^I)^2 dx.$$

Definition 1.1.3 (Sobolev differential forms). Let $1 \leq p < \infty$. We write

$$f \in W^{1,p}(\Omega; \Lambda^k)$$

if and only if every coefficient satisfies

$$f^I \in W^{1,p}(\Omega), \quad I \in \mathcal{T}_k.$$

Thus a differential form belongs to $W^{1,p}$ precisely when all of its coefficients possess weak first-order partial derivatives lying in $L^p(\Omega)$.

For a differential form $f = \sum_I f^I dx^I$, its gradient is defined coefficientwise by

$$\nabla f = \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \frac{\partial f^I}{\partial x^i} dx^i \otimes dx^I,$$

and its pointwise norm is

$$|\nabla f|^2 = \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \left(\frac{\partial f^I}{\partial x^i} \right)^2.$$

The Sobolev norm is therefore

$$\|f\|_{W^{1,p}(\Omega; \Lambda^k)} = \left(\|f\|_{L^p(\Omega; \Lambda^k)}^p + \|\nabla f\|_{L^p(\Omega; \Lambda^k)}^p \right)^{1/p},$$

and, in particular,

$$\|f\|_{W^{1,2}(\Omega; \Lambda^k)}^2 = \|f\|_{L^2(\Omega; \Lambda^k)}^2 + \|\nabla f\|_{L^2(\Omega; \Lambda^k)}^2.$$

Definition 1.1.4 (Classical regularity spaces). For an integer $r \geq 0$, we write

$$f \in C^r(\Omega; \Lambda^k)$$

if and only if every coefficient function belongs to $C^r(\Omega)$.

Similarly,

$$f \in C^{r,\alpha}(\Omega; \Lambda^k), \quad 0 < \alpha \leq 1,$$

if every coefficient belongs to the Hölder space $C^{r,\alpha}(\Omega)$.

Recall that the Hölder seminorm of a scalar-valued function u is

$$[u]_{C^{0,\alpha}(\Omega)} = \sup_{\substack{x,y \in \Omega \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha},$$

and the $C^{r,\alpha}$ norm is

$$\|u\|_{C^{r,\alpha}(\Omega)} = \sum_{|\beta| \leq r} \sup_{x \in \Omega} |D^\beta u(x)| + \sum_{|\beta|=r} [D^\beta u]_{C^{0,\alpha}(\Omega)}.$$

The corresponding norm for differential forms is again defined componentwise.

Definition 1.1.5 (Spaces on the closure). We write

$$f \in C^r(\overline{\Omega}; \Lambda^k)$$

if every coefficient extends continuously to $\overline{\Omega}$ together with all derivatives of order at most r .

Likewise,

$$f \in C^{r,\alpha}(\overline{\Omega}; \Lambda^k)$$

if every coefficient extends to a function in $C^{r,\alpha}(\overline{\Omega})$.

Remark 1.1.6. The distinction between $C^{r,\alpha}(\Omega)$ and $C^{r,\alpha}(\overline{\Omega})$ is important. Functions in $C^{r,\alpha}(\Omega)$ are only required to satisfy the Hölder condition in the interior of Ω , whereas elements of $C^{r,\alpha}(\overline{\Omega})$ possess continuous extensions, together with their derivatives up to order r , to the boundary $\partial\Omega$. Boundary conditions such as $\nu \wedge f = 0$ or $\nu \lrcorner f = 0$ are naturally stated in the latter setting.

Remark 1.1.7. Since $\Lambda^k(\mathbb{R}^n)$ is finite dimensional, all norms on $\Lambda^k(\mathbb{R}^n)$ are equivalent. Consequently, the particular choice of Euclidean norm used to define the spaces above is immaterial, and all resulting L^p , Sobolev and Hölder norms are equivalent.

Appendices

Appendix A

Some results from Calculus on $\mathbb{R}^n$

A.1 Results from Differentiation

Theorem A.1.1. (Taylor's Theorem) Let $f : U \rightarrow \mathbb{R}$ be C^1 in a neighbourhood U of $p \in \mathbb{R}^n$ that is star shaped. Then there exists $g_1, g_2, \dots, g_n$ such that

$$f(x) = f(p) + \sum_{i=1}^n g_i(x)(x^i - p^i)$$

where $g_i(p) = \frac{\partial f}{\partial x^i}|_p$

Proof. Let $x \in U$. Since U is star shaped, we have $\gamma(t) = p + t(x - p) \in U$ for $t \in [0, 1]$. Consider $\frac{d}{dt}(f \circ \gamma(t)) = Df(\gamma(t)) \cdot \gamma'(t) = \sum_{i=1}^n \frac{\partial f}{\partial x^i}(\gamma(t))(x^i - p^i)$. Integrate both sides to get

$$\int_{t=0}^{t=1} \frac{d}{dt}(f \circ \gamma(t)) dt = f(x) - f(p) = \sum_{i=1}^n (x^i - p^i) \int_{t=0}^{t=1} \frac{\partial f}{\partial x^i}(p + t(x - p)) dt$$

And let $g_i(x) := \int_{t=0}^{t=1} \frac{\partial f}{\partial x^i}(p + t(x - p)) dt$ whence it is easy to verify that $g_i(p) = \frac{\partial f}{\partial x^i}(p)$. ■

Theorem A.1.2. (Inverse Function Theorem) Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be continuously differentiable. Suppose that $\det Df(a) \neq 0$ for a given $a \in \mathbb{R}^n$. Then, there exists U and V open in $\mathbb{R}^n$ such that $a \in U$, $f(a) \in V$ and $f|_U : U \rightarrow V$ is a diffeomorphism.

Theorem A.1.3. (Implicit Function Theorem) Let $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be continuously differentiable in an open set of (a, b) and let $f(a, b) = 0$. Let M be the $m \times m$ matrix

$$D_{n+j} f^i(a, b); 1 \leq i, j \leq m$$

and suppose $\det M \neq 0$. Then there exists open sets $A \subset \mathbb{R}^n$, $B \subset \mathbb{R}^m$ with $a \in A, b \in B$ such that for all $x \in A$ there is a unique $g(x) \in B$ so that $f(x, g(x)) = 0$, and the g so obtained is differentiable.

Proof. Define $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ given by $F(x, y) := (x, f(x, y))$. This function is clearly differentiable, and moreover,

$$DF = \begin{pmatrix} \frac{\partial x^1}{\partial x^1} & \frac{\partial x^1}{\partial x^2} & \cdots & \frac{\partial x^1}{\partial x^n} & \frac{\partial x^1}{\partial y^1} & \frac{\partial x^1}{\partial y^2} & \cdots & \frac{\partial x^1}{\partial y^m} \\ \frac{\partial x^2}{\partial x^1} & \frac{\partial x^2}{\partial x^2} & \cdots & \frac{\partial x^2}{\partial x^n} & \frac{\partial x^2}{\partial y^1} & \frac{\partial x^2}{\partial y^2} & \cdots & \frac{\partial x^2}{\partial y^m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x^n}{\partial x^1} & \frac{\partial x^n}{\partial x^2} & \cdots & \frac{\partial x^n}{\partial x^n} & \frac{\partial x^n}{\partial y^1} & \frac{\partial x^n}{\partial y^2} & \cdots & \frac{\partial x^n}{\partial y^m} \\ \frac{\partial f^1}{\partial x^1} & \frac{\partial f^1}{\partial x^2} & \cdots & \frac{\partial f^1}{\partial x^n} & \frac{\partial f^1}{\partial y^1} & \frac{\partial f^1}{\partial y^2} & \cdots & \frac{\partial f^1}{\partial y^m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f^m}{\partial x^1} & \frac{\partial f^m}{\partial x^2} & \cdots & \frac{\partial f^m}{\partial x^n} & \frac{\partial f^m}{\partial y^1} & \frac{\partial f^m}{\partial y^2} & \cdots & \frac{\partial f^m}{\partial y^m} \end{pmatrix}$$

And it is easy to see that $\det DF(a, b) = \det M \neq 0$. Therefore, from Inverse function theorem (with little modification), there exists an open set $A \times B$ of $\mathbb{R}^n \times \mathbb{R}^m$, and an open set $W \subset \mathbb{R}^n \times \mathbb{R}^m$ such that $(a, b) \in A \times B$, $(a, 0) \in W$ and $F : A \times B \rightarrow W$ is a diffeomorphism. $F^{-1} : W \rightarrow A \times B$ is also a diffeomorphism. Moreover, note that $F^{-1}(p, q)$ is of the form $(p, k(p, q))$ for some differentiable function $k : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$. $\pi \circ F(x, y) = \pi \circ (x, f(x, y)) = f(x, y)$ so $\pi \circ F \equiv f$. Look at $f \circ (p, k(p, q)) = f \circ F^{-1}(p, q) = \pi(p, q) = q$ so $f \circ (p, k(p, 0)) = 0$. Our $g(p)$ is given by $k(p, 0)$, and it is differentiable as a composition of differentiable functions.

Theorem A.1.4. (A corollary of A.1.3) *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^p$ ($p \leq n$) be continuously differentiable in an open set containing $a \in \mathbb{R}^n$. If $f(a) = 0$ and the matrix $D_j(f^i(a))_{p \times n}$ has rank p , then there exists an open neighbourhood A of a and a diffeomorphism $h : A \rightarrow h(A) \subset \mathbb{R}^n$ such that $f \circ h(x_1, x_2, \dots, x_n) = (x_{n-p+1}, x_{n-p+2}, \dots, x_n)$.*

Proof. Consider $f : \mathbb{R}^{n-p} \times \mathbb{R}^p \rightarrow \mathbb{R}^p$. Suppose that the matrix $Q = D_{n-p+i}(f^j); i, j \leq p$ be invertible ($\det Q \neq 0$). Let, as before, $F : \mathbb{R}^{n-p} \times \mathbb{R}^p \rightarrow \mathbb{R}^{n-p} \times \mathbb{R}^p$ given by

$$F(p, q) = (p, f(p, q))$$

Since $\det DF \neq 0$, there exists, around $a = (a^1, a^2, \dots, a^{n-p}, a^{n-p+1}, \dots, a^n)$ a neighbourhood $A_{n-p} \times A_p \subset \mathbb{R}^{n-p} \times \mathbb{R}^p$ and a neighbourhood $W \subset \mathbb{R}^{n-p} \times \mathbb{R}^p$ containing $F(a) = (a_1, a_2, \dots, a_{n-p}, 0)$ such that $F : A_{n-p} \times A_p \rightarrow W$ is invertible and whose inverse is continuously differentiable. Note that, the inverse of F ought to be of the form $F^{-1}(x_1, x_2, \dots, x_{n-p}, y_{n-p+1}, \dots, y_n) = (\vec{x}, b(\vec{x}, \vec{y}))$ (let $\vec{x} = (x_1, x_2, \dots, x_{n-p}) \in \mathbb{R}^{n-p}$ and $\vec{y} = (y_{n-p+1}, y_{n-p+2}, \dots, y_n) \in \mathbb{R}^p$) where it is understood that $b : \mathbb{R}^n \rightarrow \mathbb{R}^p$ is continuously differentiable. Also note that $\pi_2 \circ F \equiv f$. We have that

$$\begin{aligned} f \circ F^{-1}(x_1, x_2, \dots, x_{n-p}, x_{n-p+1}, \dots, x_n) &= \\ \pi_2(x_1, x_2, \dots, x_{n-p}, x_{n-p+1}, \dots, x_n) &= (x_{n-p+1}, x_{n-p+2}, \dots, x_n) \end{aligned}$$

Case 2: If the indices where the matrix $\{(\partial_i f^j(a)); 1 \leq j \leq p, i = i_1, i_2, \dots, i_p\}$ is not the last p indices, we write the linear transformation $\text{Swap} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ that takes these p indices to the last p indices. This is a linear transformation that is invertible since it takes basis elements to basis elements. If we look then, at $f \circ \text{Swap} : \mathbb{R}^n \rightarrow \mathbb{R}^p$, we see that this is a function of the form of Case 1, so we have a h such that $f \circ \text{Swap} \circ h(x_1, x_2, \dots, x_n) = (x_{n-p+1}, x_{n-p+2}, \dots, x_n)$ which means our desired function is $\text{Swap} \circ h$.

A.2 Results from Integration

A.2.1 The Riemann Integral

Definition A.2.1. (Partition) For closed interval $[a, b]$ in $\mathbb{R}$, a partition is simply a collection of sub-intervals $[a_i, a_{i+1}]$ so that $a_0 = a < a_1 < a_2 < \dots < a_k = b$.

For a closed rectangle R in $\mathbb{R}^n$, a partition P , usually written along with its subpartitions (as in, $P \equiv (P_1, P_2, \dots, P_n)$), is a collection of subrectangles R_i (of a particular form) so that $R = \cup_i R_i$ and $R_i^\circ \cap R_j^\circ = \emptyset$. Explicitly, if

$$R = [a^1, b^1] \times [a^2, b^2] \times \dots \times [a^n, b^n]$$

,

$$P_i \equiv [a^i = a_0^i < a_1^i < a_2^i \dots a_{k_i}^i = b^i]$$

and I_j^i denotes the j -th subinterval of the i -th partition, the collection of all possible sub rectangles of R of the form $\{I_{j_1}^1 \times I_{j_2}^2 \times \dots \times I_{j_n}^n : 1 \leq j_q \leq \text{number of intervals}(P_q)\}$ is a partition of R .

Definition A.2.2. (Upper and Lower sums) Let R be a rectangle in $\mathbb{R}^n$, $f : R \rightarrow \mathbb{R}$ be a bounded function and let P be a partition of R .

$$\mathcal{L}(P, f) := \sum_{S \in P (S \text{ is a subrectangle})} \left(\inf_{x \in S} f(x) \right) |S|$$

$$\mathcal{U}(P, f) := \sum_{S \in P (S \text{ is a subrectangle})} \left(\sup_{x \in S} f(x) \right) |S|$$

where $|S|$ is the volume of the subrectangle S . For convenience, we denote $\inf_{x \in S} f(x)$ by $m_S(f)$ and $\sup_{x \in S} f(x) = M_S(f)$.

Lemma A.2.3. For a bounded function $f : R \rightarrow \mathbb{R}$,

$$\mathcal{L}(P, f) \leq \mathcal{U}(P, f)$$

for a given partition P .

Proof. If S_i are subrectangles of S coming from P' , then $m_S(f)|S| = m_S(f)(|S_1| + |S_2| + \cdots + |S_i|) \leq \sum_{S_i \in S} m_{S_i}(f)|S_i|$. This proves the result for lower sums, and the proof for upper sums is similar.

Lemma A.2.4. If P' is a refinement of P (i.e. each subrectangle of P' is contained in some subrectangle of P) then

$$\mathcal{L}(P, f) \leq \mathcal{L}(P', f)$$

and

$$\mathcal{U}(P', f) \leq \mathcal{U}(P, f)$$

Lemma A.2.5. If P and P' are any two partitions and $f : R \rightarrow \mathbb{R}$ a bounded function, then $\mathcal{L}(P', f) \leq \mathcal{U}(P, f)$.

Proof. Let P'' be a partition of both P and P' (i.e. the subrectangles of P'' are contained in some subrectangle of P as well as some subrectangle of P'). Then by the previous lemma,

$$\mathcal{L}(P', f) \leq \mathcal{L}(P'', f) \leq \mathcal{U}(P'', f) \leq \mathcal{U}(P, f)$$

Now we are in a position to define the Riemann Integral.

Definition A.2.6. ($\int_R f$) Let $f : R \rightarrow \mathbb{R}$ be a bounded function from a rectangle R . We say f is *Riemann integrable* if

$$\sup_P \mathcal{L}(P, f) = \inf_P \mathcal{U}(P, f)$$

where $\sup$ and $\inf$ are taken over all partitions P of R . In that case we say $\sup_P \mathcal{L}(P, f) = \inf_P \mathcal{U}(P, f) := \int_R f$.

Lemma A.2.7. A bounded function $f : R \rightarrow \mathbb{R}^n$ is Riemann Integrable if and only if for every $\varepsilon > 0$, there exists a partition P such that $\mathcal{U}(P, f) - \mathcal{L}(P, f) < \varepsilon$.

A.2.2 Measure Zero and Content Zero

Definition A.2.8. (Measure Zero) A set $A \subset \mathbb{R}^n$ is said to be *measure zero* if for every $\varepsilon > 0$, there exists a countable collection of closed rectangles R_i such that $A \subset \bigcup_i R_i$ and $\sum_i |R_i| < \varepsilon$.

Lemma A.2.9.

1. Subsets of Measure Zero sets are Measure Zero.
2. Countable union of Measure Zero sets is Measure Zero.

Proof. (1) Is trivial.

For (2), consider E_i a countable collection of measure zero sets. Let $\varepsilon > 0$. Consider $\varepsilon/2^i$. There exists a collection of closed rectangles $\{R_{ij} : j \in \mathbb{N}\}$ such that $E_i \subseteq \bigcup_j R_{ij}$ and $\sum_j |R_{ij}| < \varepsilon/2^i$. Look at the collection $\{R_{ij} : i, j \leq \mathbb{N}\}$ as a whole. This is a countable collection of closed rectangles that cover $\bigcup_i E_i$, and note that $\sum_{i,j} |R_{ij}| < \varepsilon$.

Definition A.2.10. (Content Zero) A set $A \subset \mathbb{R}^n$ is said to have *content zero* if for every $\varepsilon > 0$, there exists a *finite* collection of closed rectangles R_i such that $A \subseteq \bigcup_{i=1}^n R_i$ and $\sum_{i=1}^n |R_i| < \varepsilon$.

Lemma A.2.11. *If A is compact and has measure zero, then it has content zero.*

A.2.3 Integrable Functions

Definition A.2.12. (Oscillation) Let $f : X \rightarrow \mathbb{R}$ be a bounded function, where X is a metric space. Define

$$M(f, a, \delta) := \sup_{x \in B_\delta(a)} f(x)$$

and

$$m(f, a, \delta) := \inf_{x \in B_\delta(a)} f(x)$$

Let

$$\text{osc}(f, a, \delta) := M(f, a, \delta) - m(f, a, \delta)$$

Note first, that $\text{osc}(f, a, \delta)$ is decreasing for decreasing δ . Now we define

$$\text{osc}(f, a) := \lim_{\delta \rightarrow 0} \text{osc}(f, a, \delta)$$

An alternative characterisation of oscillation is the following:

Lemma A.2.13. *Let f be as in definition A.2.12. The quantity $\sup_{x,y \in B_\delta(a)} |f(y) - f(x)|$ is decreasing for decreasing δ and*

$$\text{osc}(f, a) = \lim_{\delta \rightarrow 0} \left(\sup_{x,y \in B_\delta(a)} |f(x) - f(y)| \right) = \inf_{\delta > 0} \left(\sup_{x,y \in B_\delta(a)} |f(x) - f(y)| \right)$$

Lemma A.2.14. *$f : X \rightarrow \mathbb{R}$ (where X is a metric space) is continuous at a if and only if $\text{osc}(f, a) = 0$*

Proof. $\implies$) Let f be continuous at a . For any given $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x, y \in B_\delta(a)$ we have

$$|f(x) - f(a)| < \varepsilon/2 \text{ and } |f(y) - f(a)| < \varepsilon/2$$

Adding these we get

$$|f(x) - f(y)| < \varepsilon$$

for $x, y \in B_{\delta(\varepsilon)}(a)$. Therefore, $\sup_{x,y \in B_{\delta(\varepsilon)}(a)} |f(x) - f(y)| < \varepsilon$. Since for every $\varepsilon > 0$, there exists some $\delta(\varepsilon) > 0$ such that $\sup_{x,y \in B_{\delta(\varepsilon)}(a)} |f(x) - f(y)| < \varepsilon$, we have $\text{osc}(f, a) = 0$.

$\impliedby$) Let $\text{osc}(f, a) = \inf_{\delta > 0} (\sup_{x,y \in B_\delta(a)} |f(x) - f(y)|) = 0$. This means that for every $\varepsilon > 0$, there exists $\delta > 0$ such that $\sup_{x,y \in B_\delta(a)} |f(x) - f(y)| < \varepsilon$, which means that

$$|f(x) - f(a)| < \varepsilon$$

whenever $x \in B_\delta(a)$.

Lemma A.2.15. (Recharacterisation of oscillation in terms of rectangles in $\mathbb{R}^n$)

$$\text{osc}(f, a) = \inf_{\delta > 0} \left(\sup_{x \in R_\delta(a)} f(x) - \inf_{x \in R_\delta(a)} f(x) \right)$$

Where $R_\delta(a)$ is a cube centred at a with side length δ .

Proof. Note that

$$B_{\delta/\sqrt{n}}(a) \subseteq R_\delta(a) \subseteq B_{\delta\sqrt{n}}(a)$$

From here the proof is trivial. ■

Lemma A.2.16. Let A be a closed rectangle and $f : A \rightarrow \mathbb{R}$ be a bounded function such that $\text{osc}(f, a) < \varepsilon$ for every $a \in A$. Then there exists a partition P of A such that $\mathcal{U}(f, P) - \mathcal{L}(f, P) < \varepsilon \cdot |A|$

Proof. We are given a fixed $\varepsilon_0 > 0$. We have that $\text{osc}(f, a) < \varepsilon_0$ for all $a \in A$. $\inf_{\delta > 0} (\sup_{x \in R_\delta(a)} f(x) - \inf_{y \in R_\delta(a)} f(y)) < \varepsilon_0$ for every $a \in A$. This means that, for every $a \in A$, there exists $\delta_a > 0$ such that $\sup_{x \in R_{\delta_a}(a)} f(x) - \inf_{x \in R_{\delta_a}(a)} f(x) < \varepsilon_0$. Collect all such $R_{\delta_a}(a)$. This is a cover of A , for which a finite subcover $\{R_{\delta_{a_j}}(a_j) : j \leq N\}$ exists. Consider the partition P of A formed by extending the grids of all the cubes $\{R_{\delta_{a_j}}(a_j) : j \leq N\}$. The final partition P 's subrectangles J_i have the property that

$$\sup_{x \in J_i} f(x) - \inf_{x \in J_i} f(x) < \varepsilon_0$$

Essentially,

$$M_S(f) - m_S(f) < \varepsilon_0$$

for S (a subrectangle) $\in P$. This gives us that $\mathcal{U}(f, P) - \mathcal{L}(f, P) = \sum_{S \in P} (M_S(f) - m_S(f))|S| < \varepsilon_0 \sum_{S \in P} |S| = \varepsilon_0 |A|$ ■

Theorem A.2.17. Let A be a closed rectangle and $f : A \rightarrow \mathbb{R}$ be a bounded function. Let $B := \{x \in A : f \text{ not continuous at } x\}$. Then f is integrable if and only if B is a set of measure 0.

Proof. Note that $B := \{x \in A : \text{osc}(f, x) > 0\}$ which could be written as $\bigcup_{n \in \mathbb{N}} \{x \in A : \text{osc}(f, x) \geq 1/n\}$, where we note that $\{x \in A : \text{osc}(f, x) \geq t\}$ is closed for any $t > 0$ (If $\text{osc}(f, a) < \varepsilon$ then there would exist a ball around a such that $\text{osc}(f, x) < \varepsilon$ for x in said ball).

$\Rightarrow$) Let B be measure 0. Consider the closed (and hence compact) set $B_\varepsilon := \{x \in A : \text{osc}(f, x) \geq \varepsilon\}$ whose measure (since $B_\varepsilon \subseteq B$) is 0. We thus have a finite collection of rectangles R_i that cover B_ε and $\sum_{i=1}^N |R_i| < \varepsilon$. Consider the partition P of A generated by extending the grid formed by R_i and collecting all subrectangles K_i . These can be classified into two subcategories (which together partition A):

1. $(\mathbb{S})^1 := \{K_i : K_i \cap B_\varepsilon \neq \emptyset\}$
2. $(\mathbb{S})^2 := \{K_i : K_i \cap B_\varepsilon = \emptyset\}$

We now look at $\mathcal{U}(f, P) - \mathcal{L}(f, P) = \sum_{S \in (\mathbb{S})^1 \cup (\mathbb{S})^2} (M_S(f) - m_S(f))|S|$.

Look at the first sum:

$$\sum_{S \in (\mathbb{S})^1} (M_S(f) - m_S(f))|S|$$

Since f is bounded, $|f(x)| \leq M$ and hence $\sum_{S \in (\mathbb{S})^1} (M_S(f) - m_S(f))|S| \leq 2M \sum_{S \in (\mathbb{S})^1} |S|$. Since we know that S_1 consists of the subrectangles of the cover elements (and only the cover elements) we note that $\sum_{S \in (\mathbb{S})^1} |S| \leq \varepsilon$. Hence we have the first sum $\leq 2M\varepsilon$

Look at the second sum:

$$\sum_{S \in (\mathbb{S})^2} (M_S(f) - m_S(f))|S|$$

For $S \in (\mathbb{S})^2$, we have $\text{osc}(f, x) < \varepsilon$. By A.2.16 we have that there exists a partition of S , call it P_S such that $\mathcal{U}(f, P_S) - \mathcal{L}(f, P_S) \leq \varepsilon|S|$. Extending the grid for each of these subrectangles in P_S forms a new partition P' which is a refinement of P as a whole, such that $\sum_{S' \subset S; S' \in P'} (M_{S'}(f) - m_{S'}(f))|S'| \leq \varepsilon|S|$, so as a whole,

$$\begin{aligned} \mathcal{U}(f, P') - \mathcal{L}(f, P') &= \sum_{S' \in (\mathbb{S})^1} \sum_{S'' \subset S'; S'' \in P'} (M_{S''}(f) - m_{S''}(f))|S''| + \\ &\quad \sum_{S' \in (\mathbb{S})^2} \sum_{S'' \subset S'; S'' \in P'} (M_{S''}(f) - m_{S''}(f))|S''| \\ &\leq 2M\varepsilon + \sum_{S' \in (\mathbb{S})^2} \varepsilon|S'| \leq 2M\varepsilon + \varepsilon|A| \end{aligned}$$

Hence f is integrable by criteria A.2.7 we have that f is integrable.

$\Leftarrow$) Let f be integrable. The set of discontinuities B is $\cup_{n \in \mathbb{N}} B_n$ where $B_n := \{x \in A : \text{osc}(f, x) \geq 1/n\}$ (closed and hence compact). For $a \in B_n$, we have $\text{osc}(f, a) \geq 1/n$ which means that $\inf_{\delta > 0} (M_{R_\delta}(f) - m_{R_\delta}(f)) \geq 1/n$. Let P be the partition of A such that

$$\mathcal{U}(f, P) - \mathcal{L}(f, P) \leq \varepsilon/n$$

and let SS be the collection of subrectangles of P which intersect with B_n . These form a cover for B_n . Therefore, for S such that $S \cap B_n \neq \emptyset$, we have $\exists x \in S$ such that $\inf_{\delta > 0} (M_{R_\delta}(f) - m_{R_\delta}(f)) \geq 1/n$ which implies $M_S(f) - m_S(f) \geq 1/n$ and hence

$$\varepsilon/n \geq \sum_{S \in SS} (M_S(f) - m_S(f))|S| \geq 1/n \sum_{S \in SS} |S|$$

which gives us that $\sum_{S \in SS} |S| \leq \varepsilon$. Therefore, B_n is content zero and hence B is of measure zero.

Definition A.2.18. ($\int_C f$ for arbitrary set C) Let A be any rectangle containing C (in the case where we can find one). Then we define

$$\int_C f := \int_A f \cdot \chi_C$$

where $\chi_C(x) = \begin{cases} 1 & \text{if } x \in C \\ 0 & \text{if } x \notin C \end{cases}$

Theorem A.2.19. Function $\chi_C : A \rightarrow \mathbb{R}$ is integrable if and only if ∂C is a set of measure zero.

Proof. Note that, for $x \in C^\circ$, χ_C is continuous. And likewise for $x \in (\mathbb{R}^n - C)^\circ$. The only place where χ_C is discontinuous is precisely on ∂C . The result follows from A.2.17.

Definition A.2.20. (Jordan-Measurable) A bounded set C whose boundary has measure zero is called **Jordan Measurable**.

Theorem A.2.21. If $f : A \rightarrow \mathbb{R}$ is a non-negative function and $\int_A f = 0$, then $B = \{x : f(x) > 0\}$ is a set of measure zero.

Proof. Let $B_n := \{x \in A : f(x) \geq 1/n\}$ and note that $B = \cup_n B_n$. Since f is said to be integrable, we note that $\inf_P \mathcal{U}(f, P)$ must be equal to the integral $\int_A f = 0$. Let $\varepsilon > 0$ be arbitrary and consider that partition P such that $\mathcal{U}(P, f) < \varepsilon/n$. Look at $\mathbb{J} := \{S \in P : S \cap B_n \neq \emptyset\}$. This covers B_n . Moreover, $1/n \sum_{S \in \mathbb{J}} |S| \leq \sum_{S \in \mathbb{J}} M_S(f) |S| \leq \mathcal{U}(f, P) \leq \varepsilon/n$ (Since $f(x) \geq 1/n$ for elements in B_n , and each S has some intersection with B_n). This implies $\sum_{S \in \mathbb{J}} |S| \leq \varepsilon$ and $\cup_{S \in \mathbb{J}} S$ covers B_n . Therefore, B_n is content zero set, and hence B is a measure zero set. ■

A.2.4 Fubini's Theorem

Definition A.2.22. (Lower and Upper Integrals) If $f : A \rightarrow \mathbb{R}$ is a bounded function on a closed rectangle $A \subset \mathbb{R}^n$, then regardless of whether f is integrable, the least upper bound of the lower sums and the greatest lower bound of the upper sums both exist. They are called **Lower** and **Upper integrals** of f on A , denoted by:

$$\mathbf{L} \int_A f$$

and

$$\mathbf{U} \int_A f$$

Theorem A.2.23. (Fubini) Let $A \subset \mathbb{R}^n$ and $B \subset \mathbb{R}^m$ be closed rectangles and let $f : A \times B \rightarrow \mathbb{R}$ be integrable. For $x \in A$, let $g_x : B \rightarrow \mathbb{R}$ be given by $g_x(y) = f(x, y)$ (and likewise define $h_y : A \rightarrow \mathbb{R}$ as $h_y(x) = f(x, y)$). Let

$$\mathcal{L}(x) := \mathbf{L} \int_B g_x = \mathbf{L} \int_B f(x, y) dy$$

and

$$\mathcal{U}(x) := \mathbf{U} \int_B g_x = \mathbf{U} \int_B f(x, y) dy$$

then $\mathcal{L}(x) : A \rightarrow \mathbb{R}$ and $\mathcal{U}(x) : A \rightarrow \mathbb{R}$ are both integrable on A and

$$\begin{aligned} \int_{A \times B} f &= \int_A \mathcal{L}(x) dx = \int_A \left(\mathbf{L} \int_B f(x, y) dy \right) dx \\ \int_{A \times B} f &= \int_A \mathcal{U}(x) dx = \int_A \left(\mathbf{U} \int_B f(x, y) dy \right) dx \end{aligned}$$

Proof. (sketch) ■

A.3 Partitions of Unity

Lemma A.3.1. There exists a function $f : \mathbb{R} \rightarrow \mathbb{R}$ which is positive on $(-1, 1)$, 0 elsewhere and is C^∞

Proof. Consider

$$f := \begin{cases} e^{-(x-1)^{-2}} e^{-(x+1)^{-2}} & \text{if } x \in (-1, 1) \\ 0 & \text{elsewhere} \end{cases}$$

This function is C^∞ and is positive on $(-1, 1)$ and 0 elsewhere. ■

Lemma A.3.2. Let $a \in \mathbb{R}^n$, $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be given by

$$g(x) = f([x^1 - a^1]/\varepsilon) f([x^2 - a^2]/\varepsilon) \cdots f([x^n - a^n]/\varepsilon)$$

g is then C^∞ , and has positive values only on $(a^1 - \varepsilon, a^1 + \varepsilon) \times (a^2 - \varepsilon, a^2 + \varepsilon) \cdots (a^n - \varepsilon, a^n + \varepsilon)$.

Appendix B

Differential Forms

: !!INCOMPLETE!!

primary sources: Tu Intro manifolds, Csato Dacorogna pullback equation for differential forms, Csato Dacorogna Sil "best constant for gaffney inequality", Do Carmo Differential forms and applications, Lecture notes: <https://sites.google.com/view/reyersjamaar/manifolds-and-differential-forms>

B.1 Exterior Algebra: Definitions and Results

Definition B.1.1. (Dual Space V^*) Let V be a vector space over $\mathbb{R}$. The set of all linear maps $f : V \rightarrow \mathbb{R}$ is called the *dual space* of V and is denoted by V^* or $\Lambda^1(V)$.

Lemma B.1.2. Let $e_1, e_2, \dots, e_n$ be a basis for V . Then the vectors $\xi^1, \xi^2, \dots, \xi^n \in \Lambda^1(V)$ defined by $\xi^i(e_j) = \delta_{ij}$ forms a basis for $\Lambda^1(V)$.

Definition B.1.3. (Multilinear Maps (Tensors))

$$f : \underbrace{V \times V \times \dots \times V}_{k \text{ copies}} \rightarrow \mathbb{R}$$

is a k -tensor if it is linear in each coordinate.

Note: The space of all tensors forms a vector space.

Lemma B.1.4. For the space of all k -tensors from V , $\{dx^{i_1} \otimes dx^{i_2} \otimes \dots \otimes dx^{i_k} : (i_1, i_2, \dots, i_k) \in \{1, 2, \dots, n\}^k\}$ forms a basis (Here we have adopted the notational choice of writing dx^i in place ξ^i).

Definition B.1.5. (σf) Let f be a k -tensor and $\sigma \in \text{Sym}(k)$. We define

$$\sigma f(v_1, v_2, \dots, v_k) := f(v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(k)})$$

B.1.1 σf as a group action

Let X = set of all k -tensors. Define $\rho : \text{sym}(k) \times X \rightarrow X$ by $\rho(\sigma, f) := \sigma f$. We see that ρ is a group action since

1. $\rho(1, f) = f$
2. $\rho(\sigma_1, \rho(\sigma_2, f)) = \rho(\sigma_1 \sigma_2, f)$

Definition B.1.6. (Alternating Tensors) A k -tensor is alternating if for every $\sigma \in \text{sym}(k)$,

$$f(v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(k)}) = \text{sign}(\sigma) f(v_1, v_2, \dots, v_k)$$

or

$$\sigma f \equiv \text{sign}(\sigma) f$$

Definition B.1.7. (Symmetric Tensors) A k -tensor is symmetric if for every $\sigma \in \text{sym}(k)$,

$$f(v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(k)}) = f(v_1, v_2, \dots, v_k)$$

or

$$\sigma f \equiv f$$

Definition B.1.8. (Symmetrization Operator Sf)

$$S : \{k\text{-tensors}\} \rightarrow \{k\text{-tensors}\}$$

given by $Sf(v_1, v_2, \dots, v_k) = \sum_{\sigma \in \text{sym}(k)} f(v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(k)})$ or

$$Sf \equiv \sum_{\sigma \in \text{sym}(k)} \sigma f$$

is called the *symmetrization operator*

Definition B.1.9. (Alternation or Antisymmetrizer Operator Af)

$$A : \{k\text{-tensors}\} \rightarrow \{k\text{-tensors}\}$$

given by $Af(v_1, v_2, \dots, v_k) = \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) f(v_{\sigma(1)}, v_{\sigma(2)}, \dots, v_{\sigma(k)})$ or

$$Af \equiv \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) \sigma f$$

is called the *alternation operator*

Theorem B.1.10. *If f is a k -tensor, then*

1. Sf is a symmetric k -tensor.
2. Af is an alternating k -tensor.

Proof. 1) $Sf := \sum_{\sigma \in \text{sym}(k)} \sigma f$. Let $\tau \in \text{sym}(k)$.

$$\tau Sf = \sum_{\sigma \in \text{sym}(k)} (\tau \sigma) f = Sf$$

(since $\tau \sigma$ runs over all of $\text{sym}(k)$ once and exactly once).

2) $Af := \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) f$. Let $\tau \in \text{sym}(k)$.

$$\begin{aligned} \tau Af &= \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma \tau) \text{sign}(\tau) (\tau \sigma) f = \\ &\quad \text{sign}(\tau) Af \end{aligned}$$

.

Lemma B.1.11.

1. If f is an alternating k -tensor ($\sigma f = \text{sign}(\sigma) f$) then $Af = k! f$.
2. If f is an alternating k -tensor then $Sf = 0$.
3. If f is a symmetric k -tensor ($\sigma f = f$) then $Af = 0$.
4. If f is a symmetric k -tensor then $Sf = k! f$

Proof. 1)

$$Af := \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) \text{sign}(\sigma) f = \sum_{\sigma \in \text{sym}(k)} f = k! f$$

2) If f is alternating, then

$$Sf = \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) f = 0$$

3) If f is symmetric, then

$$A(f) := \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) f = 0$$

4) If f is symmetric, then

$$Sf := \sum_{\sigma \in \text{sym}(k)} \sigma f = k! f$$

B.1.2 Tensor Product, Alternating Forms and the Wedge Product

Definition B.1.12. (Tensor Product) Let f be a k -tensor and g be a l -tensor. We define the **tensor product** $f \otimes g$ to be the $(k+l)$ -tensor given by

$$f \otimes g(v_1, v_2, \dots, v_k, v_{k+1}, v_{k+2}, \dots, v_{k+l}) := f(v_1, v_2, \dots, v_k) \cdot g(v_{k+1}, v_{k+2}, \dots, v_{k+l})$$

Note: The tensor product is *bilinear* (linear in each argument) and *associative* ($(f \otimes g) \otimes h = f \otimes (g \otimes h)$) but not *commutative*.

Definition B.1.13. (Wedge Product) Let f be an alternating k -tensor ($f \in \Lambda^k(V)$) and g an alternating l -tensor ($g \in \Lambda^l(V)$). We define

$$f \wedge g := \frac{1}{k!l!} A(f \otimes g)$$

or written explicitly

$$f \wedge g(v_1, \dots, v_{k+l}) := \frac{1}{k!l!} \sum_{\sigma \in \text{sym}(k+l)} \text{sign}(\sigma) f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) g(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)})$$

Definition B.1.14. ($\text{Sym}(k, l)$) Given natural numbers k and l , we define $\text{Sym}(k, l)$ as follows:

$$\text{Sym}(k, l) := \{\sigma \in \text{sym}(k+l) : \sigma(1) < \sigma(2) < \dots < \sigma(k) \text{ and } \sigma(k+1) < \sigma(k+2) < \dots < \sigma(k+l)\}$$

Theorem B.1.15. (Equivalent Expressions for the Wedge Product) The two expressions

1.

$$f \wedge g(v_1, \dots, v_{k+l}) := \frac{1}{k!l!} \sum_{\sigma \in \text{sym}(k+l)} \text{sign}(\sigma) f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) g(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)})$$

2.

$$f \wedge g(v_1, \dots, v_{k+l}) = \sum_{\sigma \in \text{Sym}(k, l)} \text{sign}(\sigma) f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) g(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)})$$

are equivalent.

Proof. Consider $G = \text{sym}(k+l)$ and consider the subgroup

$$H := \{\sigma \in \text{sym}(k+l) : \sigma|_{\{1, \dots, k\}} \in \{1, \dots, k\} \text{ and } \sigma|_{\{k+1, \dots, k+l\}} \in \{k+1, \dots, k+l\}\}$$

(One can view every element $\sigma \in H$ as $\pi_1 \times \pi_2$ where $\pi_1 \in \text{sym}\{1, \dots, k\}$ and $\pi_2 \in \text{sym}\{k+1, \dots, k+l\}$).

Let $\alpha, \beta \in G$ and define an equivalence relation $\alpha \sim \beta$ if and only if there exists $h \in H$ such that $\alpha h = \beta$

Let S_σ be the expression $\text{sign}(\sigma) \sigma(f \otimes g)$.

Suppose $\alpha \sim \beta$ or $\alpha h = \beta$ where $h = \pi_1 \times \pi_2$.

$$S_\alpha = \text{sign}(\alpha) f(v_{\alpha(1)}, \dots, v_{\alpha(k)}) g(v_{\alpha(k+1)}, \dots, v_{\alpha(k+l)})$$

and

$$S_\beta = \text{sign}(\beta) f(v_{\beta(1)}, \dots, v_{\beta(k)}) g(v_{\beta(k+1)}, \dots, v_{\beta(k+l)})$$

which is equal to

$$\begin{aligned} & \text{sign}(\alpha) \text{sign}(\pi_1) \text{sign}(\pi_2) f(v_{\alpha(h(1))}, \dots, v_{\alpha(h(k))}) g(v_{\alpha(h(k+1))}, \dots, v_{\alpha(h(k+l))}) \\ &= \text{sign}(\alpha) \text{sign}(\pi_1)^2 \text{sign}(\pi_2)^2 f(v_{\alpha(1)}, \dots, v_{\alpha(k)}) g(v_{\alpha(k+1)}, \dots, v_{\alpha(k+l)}) = S_\alpha \end{aligned}$$

So whenever $\alpha \sim \beta$, $S_\alpha = S_\beta$.

We can also see that, via rearrangement, for every $\sigma \in \text{sym}(k+l)$ there exists an $\alpha \in \text{sym}(k,l)$ and a $h \in H$ such that $\sigma h = \alpha$ or $\sigma \sim \alpha$.

The first expression can now be written as

$$\frac{1}{k!l!} \sum_{\sigma \in \text{sym}(k,l)} \sum_{\alpha \in [\sigma]} S_\alpha$$

but for elements of the same equivalence class, S_α are all the same and is equal to S_σ , so we have

$$\frac{1}{k!l!} \sum_{\sigma \in \text{sym}(k,l)} S_\sigma (k!l!) = \sum_{\sigma \in \text{sym}(k,l)} \text{sign}(\sigma) \sigma(f \otimes g)$$

(The $k!l!$ comes from the fact that $|H| = k!l!$ and so the size of each coset is $k!l!$) and the equality is proved. ■

Theorem B.1.16. For $f \in \Lambda^k(V)$ and $g \in \Lambda^l(V)$

$$f \wedge g = (-1)^{kl} g \wedge f$$

Proof.

$$f \wedge g = \sum_{\sigma \in \text{sym}(k,l)} \text{sign}(\sigma) \sigma(f \otimes g)$$

Let τ be the element of $\text{sym}(k+l)$ given by $\tau(1) = k+1, \tau(2) = k+2, \dots, \tau(l) = k+l$ and $\tau(l+1) = 1, \tau(l+2) = 2, \dots, \tau(l+k) = k$ (We are moving the block $\{k+1, k+2, \dots, k+l\}$ back, behind the block $\{1, 2, \dots, k\}$, which amounts to a total of kl swaps/transpositions). Note that τ acts as a correspondence between $\text{sym}(k,l)$ and $\text{sym}(l,k)$ since for every $\sigma \in \text{sym}(k,l)$, we have $\sigma(\tau(l+1)) < \sigma(\tau(l+2)) < \dots < \sigma(\tau(l+k))$ and $\sigma(\tau(1)) < \sigma(\tau(2)) < \dots < \sigma(\tau(l))$ so $\sigma(\tau)$ is in $\text{sym}(l,k)$.

$$\begin{aligned} f \wedge g(v_1, \dots, v_{k+l}) &= \sum_{\sigma \in \text{sym}(k,l)} \text{sign}(\sigma) f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) g(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)}) \\ &= \sum_{\sigma \in \text{sym}(k,l)} \text{sign}(\tau\sigma) \text{sign}(\tau) g(v_{\sigma(\tau(1))}, \dots, v_{\sigma(\tau(l))}) f(v_{\sigma(\tau(l+1))}, \dots, v_{\sigma(\tau(l+k))}) \\ &= \sum_{\gamma \in \text{sym}(l,k)} \text{sign}(\gamma) (-1)^{kl} g(v_{\gamma(1)}, \dots, v_{\gamma(l)}) f(v_{\gamma(l+1)}, \dots, v_{\gamma(l+k)}) = (-1)^{kl} g \wedge f \end{aligned}$$
■

B.1.3 The Associativity of the Wedge Product

Lemma B.1.17. Let $f \in \Lambda^k(V)$ and $g \in \Lambda^l(V)$, then:

1. $A(Af \otimes G) = k!A(f \otimes g)$
2. $A(f \otimes Ag) = l!A(f \otimes g)$

Proof. 1)

$$A(Af \otimes g) = \sum_{\sigma \in \text{sym}(k+l)} \text{sign}(\sigma) \sigma(Af \otimes g)$$

$$Af := \sum_{\substack{\tau \in \text{sym}(k+l) \\ \tau \text{ preserves } \{k+1, \dots, k+l\}}} \text{sign}(\tau) \tau(f)$$

and

$$\begin{aligned} A(Af \otimes g) &:= \sum_{\sigma \in \text{sym}(k+l)} \text{sign}(\sigma) (Af(X_{\sigma(1)}, \dots, X_{\sigma(k)})g(X_{\sigma(k+1)}, \dots, X_{\sigma(k+l)})) \\ &= \sum_{\sigma \in \text{sym}(k+l)} \text{sign}(\sigma) \left(\sum_{\substack{\tau \in \text{sym}(k+l) \\ \tau \text{ preserves } \{k+1, \dots, k+l\}}} \text{sign}(\tau) f(X_{\sigma\tau(1)}, \dots, X_{\sigma\tau(k)}) \right) g(X_{\sigma(k+1)}, \dots, X_{\sigma(k+l)}) \\ &= \sum_{\sigma \in \text{sym}(k+l)} \sum_{\substack{\tau \in \text{sym}(k+l) \\ \tau \text{ fixes } \{k+1, \dots, k+l\}}} \text{sign}(\sigma\tau) f(X_{\sigma\tau(1)}, \dots, X_{\sigma\tau(k)}) g(X_{\sigma\tau(k+1)}, \dots, X_{\sigma\tau(k+l)}) \end{aligned}$$

The number of such τ that fixes $\{k+1, \dots, k+l\}$ are precisely $k!$, and for each such τ , $\sigma\tau$ runs over all of $\text{sym}(k+l)$ once and exactly once. Hence we have

$$A(Af \otimes g) = k! \sum_{\gamma \in \text{sym}(k+l)} \text{sign}(\gamma) \gamma(f \otimes g) = k!A(f \otimes g)$$

2) The 2nd part is proved similarly. ■

Theorem B.1.18. (Associativity of Wedge Product) For any $f \in \Lambda^k(V)$, $g \in \Lambda^l(V)$, $h \in \Lambda^m(V)$

$$f \wedge (g \wedge h) = (f \wedge g) \wedge h$$

Proof.

$$\begin{aligned} f \wedge (g \wedge h) &:= \frac{1}{k!(l+m)!} A(f \otimes (g \wedge h)) = \frac{1}{k!(l+m)!} A(f \otimes (\frac{1}{l!m!} A(g \otimes h))) \\ &= \frac{1}{k!l!m!(l+m)!} A(f \otimes A(g \otimes h)) = \frac{1}{k!l!m!} A(f \otimes g \otimes h) \end{aligned}$$

and likewise

$$\begin{aligned} (f \wedge g) \wedge h &= \frac{1}{(k+l)!m!} A((f \wedge g) \otimes h) = \frac{1}{(k+l)!k!l!m!} A(A(f \otimes g) \otimes h) \\ &= \frac{1}{k!l!m!} (k+l)! A(f \otimes g \otimes h) = \frac{1}{k!l!m!} A(f \otimes g \otimes h) \end{aligned}$$

Note: The proof of B.1.18 shows that if f_{l_1} is an l_1 -form, f_{l_2} is an l_2 -form and so on, $f_{l_1} \wedge f_{l_2} \wedge \dots \wedge f_{l_k}$ is given by

$$f_{l_1} \wedge f_{l_2} \wedge \dots \wedge f_{l_k} = \frac{1}{l_1!l_2! \dots l_k!} A(f_{l_1} \otimes f_{l_2} \otimes \dots \otimes f_{l_k}) \quad (\text{B.1.1})$$

Short Digression on Determinants

Definition B.1.19. (Determinant (*Leibniz Formula*)) Let $\{a_{ij} : 1 \leq i, j \leq n\}$ be a matrix. We define determinant of A as

$$\det(A) := \sum_{\sigma \in \text{sym}(n)} \text{sign}(\sigma) a_{1,\sigma(1)} a_{2,\sigma(2)} \cdots a_{n,\sigma(n)}$$

(To be thought of as multiplying one element from each row chosen from distinct columns, and taking into account the sign of the permutation of the columns).

Definition B.1.20. (Minor of (i, j) -entry) Let A be a matrix and let a_{ij} be the (i, j) -entry of A . The *minor* of a_{ij} is the determinant of the matrix obtained by deleting the i^{th} row and j^{th} column from A . It is denoted m_{ij} . The matrix itself (after deleting i^{th} row and j^{th} column) is denoted M_{ij} .

Definition B.1.21. (Laplace's Expansion) Fix a row i . Then the Laplace's expansion gives

$$\det(B) = \sum_{j=1}^n (-1)^{i+j} b_{ij} \det(M_{ij})$$

Note: From

$$\sum_{\sigma \in \text{sym}(n)} \text{sign}(\sigma) \prod_{q=1}^n a_{q,\sigma(q)} = \sum_{j=1}^n \sum_{\substack{\sigma \in \text{sym}(n) \\ \sigma: i \mapsto j}} \text{sign}(\sigma) a_{i,j} \prod_{\substack{q=1 \\ q \neq i}}^n a_{q,\sigma(q)} \quad (\text{B.1.2})$$

we can see that the Laplace's expansion is a consequence of the Leibniz formula for determinants.

Theorem B.1.22. (Wedge Product of 1-forms or Functionals) Let $f_1, f_2, \dots, f_k$ be in $\Lambda^1(V) \equiv V^*$. Then

$$f_1 \wedge f_2 \wedge \cdots \wedge f_k(x_1, x_2, \dots, x_k) = \det(f_i(x_j))$$

Proof. From B.1.1, we have

$$f_1 \wedge f_2 \wedge \cdots \wedge f_k(x_1, \dots, x_k) = A(f_1 \otimes f_2 \otimes \cdots \otimes f_k)$$

or

$$f_1 \wedge f_2 \wedge \cdots \wedge f_k(x_1, x_2, \dots, x_k) = \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) f_1(x_{\sigma(1)}) f_2(x_{\sigma(2)}) \cdots f_k(x_{\sigma(k)})$$

which is precisely the Leibniz formula for determinants. Hence

$$f_1 \wedge f_2 \wedge \cdots \wedge f_k(x_1, x_2, \dots, x_k) = \det(f_i(x_j))$$

Lemma B.1.23. Any collection $f_1, f_2, \dots, f_k$ of 1-forms is linearly independent if and only if $f_1 \wedge f_2 \wedge \cdots \wedge f_k \neq 0$

Proof. $\implies$) Say $f_1, \dots, f_k$ is a linearly independent collection of functionals. Then it is part of a larger collection that forms a basis for V^* . Let $x_1, x_2, \dots, x_k$ be the vectors in V such that $f_i(x_j) = \delta_{ij}$. Then

$$f_1 \wedge f_2 \wedge \cdots \wedge f_k(x_1, x_2, \dots, x_k) = \det(\delta_{ij}) = 1$$

$\impliedby$) Let $f_1, f_2, \dots, f_k$ be linearly dependent. Then (without loss of generality) $f_1 = c_2 f_2 + c_3 f_3 + \cdots + c_k f_k$ and $f_1 \wedge f_2 \wedge \cdots \wedge f_k = c_2(f_2 \wedge f_2 \wedge f_3 \cdots \wedge f_k) + c_3(f_3 \wedge f_2 \wedge f_3 \wedge \cdots \wedge f_k) + \cdots + c_k(f_k \wedge f_2 \wedge f_3 \wedge \cdots \wedge f_k) = 0$

B.1.4 Dimension of Alternating k-forms

Let $E_1, E_2, \dots, E_n$ be a basis for V . Let $\xi^1, \xi^2, \dots, \xi^n \in V^* = \Lambda^k(V)$ be the dual basis to $E_1, \dots, E_n$.

Theorem B.1.24. (*Basis for $\Lambda^k(V)$*) The set

$$\{\xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k} : 1 \leq i_1 < i_2 < \dots < i_k \leq n\}$$

forms a basis for $\Lambda^k(V)$.

Proof. Linear Independence: Look at

$$\sum_{1 \leq i_1 < \dots < i_k \leq n} c_{i_1 i_2 \dots i_k} \xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k} \equiv 0$$

in $\Lambda^k(V)$. Hence,

$$\begin{aligned} \sum_{1 \leq i_1 < \dots < i_k \leq n} c_{i_1 i_2 \dots i_k} \xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k} (E_{j_1}, E_{j_2}, \dots, E_{j_k}) &= \\ \sum_{1 \leq i_1 < \dots < i_k \leq n} c_{i_1 i_2 \dots i_k} \det(\xi^{i_r}(E_{j_s})) &= 0 \\ \implies c_{j_1 j_2 \dots j_k} &= 0 \end{aligned}$$

for every increasing k-tuple $j_1, j_2, \dots, j_k$ and hence $\{\xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k} : 1 \leq i_1 < \dots < i_k \leq n\}$ is linearly independent.

Span: Let $f \in \Lambda^k(V)$. Let $x_r = \sum_{j_1=1}^n \xi^{j_1}(x_r) E_{j_1}$ for $r = 1, 2, \dots, k$.

$$\begin{aligned} f(x_1, x_2, \dots, x_k) &= f\left(\sum_{j_1=1}^n \xi^{j_1}(x_1) E_{j_1}, \sum_{j_1=1}^n \xi^{j_1}(x_2) E_{j_1}, \dots, \sum_{j_1=1}^n \xi^{j_1}(x_k) E_{j_1}\right) \\ &= \sum_{j_1=1}^n \sum_{j_2=1}^n \dots \sum_{j_k=1}^n \xi^{j_1}(x_1) \xi^{j_2}(x_2) \dots \xi^{j_k}(x_k) f(E_{j_1}, E_{j_2}, \dots, E_{j_k}) \end{aligned}$$

If $j_r = j_s$ for some r and s , note that the summand is 0. Hence we rewrite the above sum as:

$$\sum_{\substack{1 \leq j_1, j_2, \dots, j_k \leq n \\ j_r \text{ distinct}}} \xi^{j_1}(x_1) \xi^{j_2}(x_2) \dots \xi^{j_k}(x_k) f(E_{j_1}, E_{j_2}, \dots, E_{j_k})$$

We now note that each choice of $1 \leq j_1, j_2, \dots, j_k \leq n$ tuple is some $\sigma \in \text{sym}(k)$ of the ordered $1 \leq j_1 < j_2 < \dots < j_k \leq n$ (Simply reorder the elements $j_1, j_2, \dots, j_k$ such that we achieve $\sigma(j_1) < \sigma(j_2) < \dots < \sigma(j_k)$), and hence the above sum can be rewritten as:

$$\sum_{1 \leq j_1 < \dots < j_k \leq n} \sum_{\sigma \in \text{sym}(k)} \xi^{j_{\sigma(1)}}(x_1) \xi^{j_{\sigma(2)}}(x_2) \dots \xi^{j_{\sigma(k)}}(x_k) f(E_{j_{\sigma(1)}}, \dots, E_{j_{\sigma(k)}})$$

which is precisely

$$\begin{aligned} \sum_{1 \leq j_1 < \dots < j_k \leq n} f(E_{j_1}, \dots, E_{j_k}) \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) \xi^{j_{\sigma(1)}}(x_1) \xi^{j_{\sigma(2)}}(x_2) \dots \xi^{j_{\sigma(k)}}(x_k) \\ = \sum_{1 \leq j_1 < \dots < j_k \leq n} f(E_{j_1}, \dots, E_{j_k}) \xi^{j_1} \wedge \xi^{j_2} \wedge \dots \wedge \xi^{j_k}(x_1, x_2, \dots, x_k) \end{aligned}$$

which means f can be written as a linear combination of $\xi^{j_1} \wedge \dots \wedge \xi^{j_k}$.

Lemma B.1.25. Let $\beta^i \in \Lambda^1(V)$ for $1 \leq i \leq k$ and let $\gamma^i \in \Lambda^1(V)$ and if

$$\beta^i = \sum_{j=1}^k \alpha_j^i \gamma^j$$

then

$$\beta^1 \wedge \beta^2 \wedge \cdots \wedge \beta^k = \det(\alpha_j^i) \gamma^1 \wedge \gamma^2 \wedge \cdots \wedge \gamma^k$$

Proof.

$$\begin{aligned} \beta^1 \wedge \cdots \wedge \beta^k &= \sum_{j_1=1}^k \sum_{j_2=1}^k \cdots \sum_{j_k=1}^k (\alpha_{j_1}^1 \alpha_{j_2}^2 \cdots \alpha_{j_k}^k) (\gamma^{j_1} \wedge \gamma^{j_2} \wedge \cdots \wedge \gamma^{j_k}) \\ &= \sum_{\substack{1 \leq j_1 < \cdots < j_k \leq k \\ j_r \text{ distinct}}} (\alpha_{j_1}^1 \alpha_{j_2}^2 \cdots \alpha_{j_k}^k) \gamma^{j_1} \wedge \gamma^{j_2} \cdots \wedge \gamma^{j_k} \\ &= \sum_{\substack{1 \leq j_1 < \cdots < j_k \leq k \\ \sigma \in \text{sym}(k)}} \text{sign}(\sigma) \alpha_{j_1}^1 \cdots \alpha_{j_k}^k \gamma^1 \wedge \gamma^2 \cdots \wedge \gamma^k \\ &= \det(\alpha_j^i) \gamma^1 \wedge \gamma^2 \wedge \cdots \wedge \gamma^k \end{aligned}$$

Lemma B.1.26. Let $f \in \Lambda^k(V)$ and let $u_1, u_2, \dots, u_k$ and $v_1, v_2, \dots, v_k \in V$ be related by

$$u_i = \sum_{j=1}^n a_j^i v_j$$

Then

$$f(u_1, u_2, \dots, u_k) = \det(a_j^i) f(v_1, v_2, \dots, v_k)$$

Proof. $f(\sum_{j_1=1}^n a_{j_1}^1 v_{j_1}, \sum_{j_2=1}^n a_{j_2}^2 v_{j_2}, \dots, \sum_{j_k=1}^n a_{j_k}^k v_{j_k})$

$$\begin{aligned} &= \sum_{\substack{j_1, j_2, \dots, j_k \\ \text{each distinct}}} a_{j_1}^1 a_{j_2}^2 \cdots a_{j_k}^k f(v_{j_1}, v_{j_2}, \dots, v_{j_k}) \\ &= \sum_{\substack{1 \leq j_1 < j_2 < \cdots < j_k \leq k \\ \text{each distinct}}} \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) a_{\sigma(1)}^1 a_{\sigma(2)}^2 \cdots a_{\sigma(k)}^k f(v_1, v_2, \dots, v_k) = \\ &\quad \det(a_j^i) f(v_1, v_2, \dots, v_k) \end{aligned}$$

Notation: Let $\mathcal{T}_k$ denote the set

$$\mathcal{T}_k := \{I \equiv (i_1, i_2, \dots, i_k) : 1 \leq i_1 < i_2 < \cdots < i_k \leq n\} \quad (\text{B.1.3})$$

Suppose that $E_1, E_2, \dots, E_n$ are the basis for V , and $\xi^1, \xi^2, \dots, \xi^n$ be the dual basis for $\Lambda^1(V)$. Then, for $I = (i_1, i_2, \dots, i_k)$ we have the notation

$$\xi^I := \xi^{i_1} \wedge \xi^{i_2} \wedge \cdots \wedge \xi^{i_k} \quad (\text{B.1.4})$$

and hence for any k -form $f : V \times V \times \cdots \times V \rightarrow \mathbb{R}$ we have the notation for the expansion

$$f := \sum_{I \in \mathcal{T}_k} f^I \xi^I \quad (\text{B.1.5})$$

B.2 Tangent vectors in $\mathbb{R}^n$ and Vector Fields

We denote points in $\mathbb{R}^n$ by

$$p = (p^1, p^2, \dots, p^n)$$

and points in $T_p(\mathbb{R}^n)$, the tangent space of $\mathbb{R}^n$ at p (the space of all vectors v emanating from p), by

$$v = [v^1, v^2, \dots, v^n]^T$$

B.2.1 The Directional Derivative and Derivations

Definition B.2.1. (Directional Derivative) Let $p \in \mathbb{R}^n$ be given and let $f \in C_p^\infty$, the space of all functions $f : U \rightarrow \mathbb{R}$ such that $p \in U$ and f is C^∞ in U . Let $v \in T_p(\mathbb{R}^n)$. Let $\gamma(t) := p + tv$ and look at

$$\frac{d}{dt}\bigg|_{t=0}(f \circ \gamma(t)) = Df(\gamma(0)) \cdot \gamma'(0) = \sum_{i=1}^n v^i \frac{\partial}{\partial x^i}\bigg|_p f$$

If we fix a point p (arbitrary) and a direction $v \in T_p(\mathbb{R}^n)$, we have an operator $D_{p,v} : C_p^\infty \rightarrow \mathbb{R}$ given by

$$D_{p,v}(f) := \sum_{i=1}^n v^i \frac{\partial}{\partial x^i}\bigg|_p (f)$$

This operator is *Linear* and follows the property

$$D_{p,v}(f \cdot g) = (D_{p,v}(f))g(p) + (D_{p,v}(g))f(p)$$

The operator $D_{p,v}$ is called the **directional derivative** at p in the direction v .

Definition B.2.2. (Germs of Functions) Fix a point p and look at all pairs (f, U) where $f : U \rightarrow \mathbb{R}$, $p \in U \subseteq \mathbb{R}^n$ open and $f \in C^\infty(U)$. We say $(f, U) \sim (g, V)$ if f and g agree on an open subset of $U \cap V$. This is an equivalence relation and we (by abuse of notation) denote the equivalence class of (f, U) as simply f . The set of all such equivalence classes is denoted by C_p^∞ known as the **Germs of functions at p** .

Defining addition, scalar multiplication and function multiplication on C_p^∞ by the usual operations on representatives, we see that C_p^∞ is a commutative ring with identity. Moreover, it is also a Vector space over $\mathbb{R}$.

Definition B.2.3. (Point Derivation) Fix a point $p \in \mathbb{R}^n$. We say $D : C_p^\infty \rightarrow \mathbb{R}$ is a *point derivation* (at p) if:

1. D is linear.
2. D obeys *Leibniz's Rule*:

$$D(f \cdot g) = D(f)g(p) + D(g)f(p)$$

and the space of all point derivations at p is denoted by $\mathcal{D}_p(\mathbb{R}^n)$

Lemma B.2.4. If $D \in \mathcal{D}_p(\mathbb{R}^n)$, then $D(c) = 0$ for every constant function $c \in C_p^\infty$.

Proof. $D(1) = D(1 \cdot 1) = 1D(1) + 1D(1) = 2D(1)$ which implies $D(1) = 0$. From linearity, we have the result. ■

Lemma B.2.5. The space of all point derivations $\mathcal{D}_p(\mathbb{R}^n)$ is a vector space over $\mathbb{R}$ and $T_p(\mathbb{R}^n) \cong \mathcal{D}_p(\mathbb{R}^n)$. Moreover, $D_{p,(\cdot)} : T_p(\mathbb{R}^n) \rightarrow \mathcal{D}_p(\mathbb{R}^n)$ given by $v \mapsto D_{p,v}$ is the isomorphism.

Proof. That $\mathcal{D}_p(\mathbb{R}^n)$ is a vector space is clear. Look at the map $v \mapsto D_{p,v}$. $v + cw \mapsto D_{p,v+cw}$ Where

$$D_{p,v+cw}(f) = \sum_{i=1}^n (v^i + cw^i) \frac{\partial}{\partial x^i}\bigg|_p (f) = D_{p,v}(f) + cD_{p,w}(f)$$

Which makes $v \mapsto D_{p,v}$ linear.

Injectivity: Let $D_{p,v}(f) = \sum_{i=1}^n v^i \frac{\partial}{\partial x^i}\bigg|_p (f) \equiv 0$ for all $f \in C_p^\infty$. Let $f = x^j : \mathbb{R}^n \rightarrow \mathbb{R}$ that picks out the j -th coordinate. Then $D_{p,v}(x^j) = \sum_{i=1}^n v^i \frac{\partial}{\partial x^i}\bigg|_p (x^j) = v^j = 0$. Therefore, $v = 0$ and hence, the nullspace is trivial.

Surjectivity: Let $f \in C_p^\infty$. We have $f(x) = f(p) + \sum_{i=1}^n g_i(x)(x^i - p^i)$ where $g^i(p) = \frac{\partial f}{\partial x^i}(p)$

(From A.1.1)

Let $D \in \mathcal{D}_p(\mathbb{R}^n)$.

$$\begin{aligned} D(f(x)) &= D(f(p)) + \sum_{i=1}^n D(g_i(x) \cdot (x^i - p^i)) \\ &= 0 + \sum_{i=1}^n D(g_i(x))(0) + D(x^i - p^i) \frac{\partial f}{\partial x^i}(p) = \sum_{i=1}^n D(x^i) \frac{\partial}{\partial x^i} \Big|_p f \end{aligned}$$

Which means

$$D \equiv \sum_{i=1}^n D(x^i) \frac{\partial}{\partial x^i} \Big|_p \equiv D_{p, \langle D(x^1), D(x^2), \dots, D(x^n) \rangle}$$

And hence, the map $v \mapsto D_{p,v}$ is surjective.

■

Definition B.2.6. (Vector Field) A *vector field* on $\mathbb{R}^n$ is a map $X : U \mapsto \Pi_{p \in U} T_p(U)$. For every $p \in U$, $X(p)$ is a vector in the tangent space at p .

$$X(p) = \sum_{i=1}^n a^i(p) \frac{\partial}{\partial x^i} \Big|_p$$

We call X a smooth vector field if $a^i : U \rightarrow \mathbb{R}$ are C^∞ .

Let $f \in C_p^\infty$, that is $f : U \rightarrow \mathbb{R}$, a vector field X to f is

$$Xf(p) := \sum a_i(p) \frac{\partial}{\partial x^i} \Big|_p f$$

Then

$$Xf : U \rightarrow \mathbb{R}$$

is again a C^∞ function.

B.3 Differential Forms

We start with defining what are 1-forms.

Definition B.3.1. (1-forms) A function

$$\omega : U \rightarrow \Pi_{p \in U} \Lambda^1(T_p(\mathbb{R}^n))$$

such that $\omega(p) \in \Lambda^1(T_p(\mathbb{R}^n)) \equiv T_p^*(\mathbb{R}^n)$ is called a 1-form.

Note: Let $f : U \rightarrow \mathbb{R}$ be any C^∞ function. We can construct a 1-form from f as follows:

$$df : U \rightarrow \Pi_{p \in U} \Lambda^1(T_p(U))$$

such that

$$df_p \in \Lambda^1(T_p(U))$$

$$df_p : V \rightarrow \mathbb{R}$$

given by the action (for $v = \sum_{j=1}^n v^j \frac{\partial}{\partial x^j} \Big|_p \in T_p(U)$):

$$df_p(v) = df_p\left(\sum_j v^j \frac{\partial}{\partial x^j} \Big|_p\right) = \sum_j v^j \frac{\partial}{\partial x^j} \Big|_p(f) \in \mathbb{R}$$

Suppose that $v \in T_p(\mathbb{R}^n)$ and $f \in C_p^\infty(\mathbb{R}^n)$.

$$v = \sum_{i=1}^n a^i(p) \frac{\partial}{\partial x^i} \Big|_p \equiv X(p)$$

where $X : U \rightarrow \Pi_{p \in U} T_p(\mathbb{R}^n)$ is formally

$$X(p) := \sum_{i=1}^n a^i(p) \frac{\partial}{\partial x^i} \Big|_p \in T_p(\mathbb{R}^n)$$

We then have a bilinear function

$$\langle, \rangle : T_p(\mathbb{R}^n) \times C_p^\infty(\mathbb{R}^n) \rightarrow \mathbb{R}$$

given by

$$\langle v, f \rangle = \langle \sum_i a^i(p) \frac{\partial}{\partial x^i} \Big|_p, f \rangle := \sum_i a^i(p) \frac{\partial}{\partial x^i} \Big|_p f \in \mathbb{R}$$

One can then think of *tangent vectors* as $\langle v, () \rangle$ functions of the 2nd argument. Likewise, we can think of df as maps on the 1st argument: $df_p() = \langle (), f \rangle$ that takes in tangent vectors and gives a number in $\mathbb{R}$.

Lemma B.3.2. *Let $x^1, x^2, \dots, x^n : \mathbb{R}^n \rightarrow \mathbb{R}$ be the standard coordinate functions on $\mathbb{R}^n$. Then at each point p , $\{dx_p^1, dx_p^2, \dots, dx_p^n\}$ forms a basis for $\Lambda^1(T_p(\mathbb{R}^n))$ dual to $\{\frac{\partial}{\partial x^1} \Big|_p, \frac{\partial}{\partial x^2} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p\} \in T_p(\mathbb{R}^n)$.*

Proof. Note that

$$dx^i : \mathbb{R}^n \rightarrow \cup_{p \in \mathbb{R}^n} \Lambda^1(T_p(\mathbb{R}^n))$$

where $dx_p^i \in \Lambda^1(T_p(\mathbb{R}^n))$ is such that

$$dx_p^i \left(\frac{\partial}{\partial x^j} \Big|_p \right) := \frac{\partial}{\partial x^j} \Big|_p (x^i) = \delta_{ij}$$

and $dx_p^j \in \Lambda_p(T_p(\mathbb{R}^n))$ is precisely the dual basis to $\frac{\partial}{\partial x^i} \Big|_p \in T_p(\mathbb{R}^n)$.

■

Note: If

$$\omega : U \rightarrow \Pi_{p \in U} \Lambda^1(T_p(\mathbb{R}^n))$$

is a 1-form, $\omega(p) \in \Lambda^1(T_p(\mathbb{R}^n))$ and

$$\omega(p) = \sum_i a^i(p) dx_p^i$$

and formally

$$\omega(\cdot) \equiv \sum_i a_i(\cdot) dx_{(\cdot)}^i$$

Where $a^j(p) = w_p(\frac{\partial}{\partial x^j} \Big|_p)$.

B.3.1 Differential Forms in terms of Coordinates

Theorem B.3.3. *Let $f : U \rightarrow \mathbb{R}$ be C^∞ . Then df_p is defined as that linear function on $T_p(\mathbb{R}^n)$ such that*

$$df_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \frac{\partial}{\partial x^i} \Big|_p f$$

Then (as a covector)

$$df_p \equiv \sum_{i=1}^n \frac{\partial f}{\partial x^i} \Big|_p (dx^i)_p$$

and formally (as a map on U)

$$df_{()} \equiv \sum_{j=1}^n \frac{\partial f}{\partial x^j} \Big|_{()} dx_{()}^j$$

Proof. Let $f : U \rightarrow \mathbb{R}$. $df_p \in \Lambda^1(T_p(U))$ and hence we have an expansion of the form

$$df_p = \sum_{j=1}^n a_p^j dx_p^j \quad (\text{B.3.1})$$

where $x^j : \mathbb{R}^n \rightarrow \mathbb{R}$ is the j -th standard coordinate function and dx^j is the dual covector to $\frac{\partial}{\partial x^i} \big|_p \in T_p(U)$. Applying df_p on $\frac{\partial}{\partial x^i} \big|_p$ (in B.3.1) we get

$$df_p\left(\frac{\partial}{\partial x^i} \big|_p\right) := \frac{\partial f}{\partial x^i} \big|_p = \sum_{j=1}^n a_p^j dx_p^j\left(\frac{\partial}{\partial x^i} \big|_p\right) = a_p^i$$

and hence

$$df_p \equiv \sum_{j=1}^n \frac{\partial f}{\partial x^j} \big|_p dx_p^j$$

Note: Let $X_p \in T_p(\mathbb{R}^n)$ be a tangent vector. we have $X_p \equiv \sum_{i=1}^n b^i(X_p) \frac{\partial}{\partial x^i} \big|_p$. Apply dx_p^j on both sides:

$$dx_p^j(X_p) = b^j(X_p)$$

and hence

$$X_p \equiv \sum_{i=1}^n dx_p^i(X_p) \frac{\partial}{\partial x^i} \big|_p$$

B.3.2 Differential k -forms

Definition B.3.4. (Differential k -form) A differential k -form is a function of the kind

$$\begin{aligned} \omega : U &\rightarrow \Pi_{p \in U} \Lambda^k(T_p(\mathbb{R}^n)) \\ \omega_p &\in \Lambda^k(T_p(\mathbb{R}^n)) \end{aligned}$$

Since $\Lambda^k(T_p(\mathbb{R}^n))$ is spanned by vectors of the kind (for $1 \leq i_1 < i_2 < \dots < i_k \leq n$)

$$dx_p^{i_1} \wedge dx_p^{i_2} \wedge \dots \wedge dx_p^{i_k}$$

and hence

$$\omega_p \equiv \sum_{1 \leq i_1 < \dots < i_k \leq n} w_{(p)}^{i_1 i_2 \dots i_k} dx_p^{i_1} \wedge dx_p^{i_2} \wedge \dots \wedge dx_p^{i_k}$$

where $w^{i_1 i_2 \dots i_k} : U \rightarrow \mathbb{R}$ is a C^∞ function. Then we say ω is a C^∞ differential k -form.

Example B.3.5. (In $\mathbb{R}^3$) Let $x, y, z : \mathbb{R}^3 \rightarrow \mathbb{R}$ be the coordinate maps.

1. C^∞ 1-forms are maps

$$\omega : U \rightarrow \Pi_{p \in U} \Lambda^1(T_p(U))$$

and

$$\omega_p \equiv \sum_{i=1}^3 f_p^i dx_p^i = f_p^x dx_p + f_p^y dy_p + f_p^z dz_p \in \Lambda^1(T_p(U))$$

2. C^∞ 2-forms are maps

$$\omega : U \rightarrow \Pi_{p \in U} \Lambda^2(T_p(U))$$

and

$$\omega_p \equiv \sum_{1 \leq i_1 < i_2 \leq 3} f_p^{i_1 i_2} dx_p^{i_1} \wedge dx_p^{i_2} = f_p^{1,2} dx_p \wedge dy_p - f_p^{1,3} dz_p \wedge dx_p + f_p^{2,3} dy_p \wedge dz_p \in \Lambda^2(T_p(U))$$

3. C^∞ 3-forms are maps

$$\omega : U \mapsto \Pi_p \Lambda^3(T_p(U))$$

and

$$\omega_p \equiv \sum_{1 \leq i_1 < i_2 < i_3} f_p^{i_1 i_2 i_3} dx_p^{i_1} \wedge dx_p^{i_2} \wedge dx_p^{i_3} = f_p dx_p \wedge dy_p \wedge dz_p \in \Lambda^3(T_p(U))$$

B.3.3 Exterior Derivative

Consider a C^∞ function $f : U \rightarrow \mathbb{R}$. We have $df : U \rightarrow \Pi_{p \in U} \Lambda^1(T_p(U))$ such that $df_p \in \Lambda^1(T_p(U))$.

$$df_p : T_p(U) \rightarrow \mathbb{R}$$

via

$$df_p(v) = df_p\left(\sum_{i=1}^n dx_p^i(v) \frac{\partial}{\partial x^i} \Big|_p\right) = \sum_{i=1}^n \frac{\partial f}{\partial x^i} \Big|_p dx_p^i(v)$$

and symbolically we have

$$df \equiv \sum_{i=1}^n \frac{\partial f}{\partial x^i} dx^i$$

So d in this case can be viewed as a map that takes 0-forms (functions) and gives us a 1-form df .

Before we define the general exterior derivative for k -forms, we will set up some notation.

Definition B.3.6. (Multi Index Notations) Let $I \equiv \{i_1, i_2, \dots, i_k\} \in \mathcal{T}_k$ and $i \notin I$. For $i \in I$, we write $I_{\hat{i}}$ as the multi index I with i removed. For $i \notin I$, we write $[iI]$ as the multi index I with i added in the correct order.

We also write $\text{sign}[i, I]$ as the sign of the permutation that takes $\{i, I\}$ to $[iI]$. A corollary of this is that

$$dx^{[iI]} = \text{sign}[i, I] dx^i \wedge dx^I$$

We present an important lemma from [CDS18] that will be used in the document.

Lemma B.3.7. *Let $n \geq 2$ and $1 \leq k \leq n-1$ be integers. Then for any $I \in \mathcal{T}_{k+1}$ and $i, j \in I$, with $i \neq j$ we have*

$$\text{sign}[i, I_{\hat{i}j}] \text{sign}[j, I_{\hat{i}j}] = -\text{sign}[i, I_{\hat{i}}] \text{sign}[j, I_{\hat{j}}] \quad (\text{B.3.2})$$

And for any $I \in \mathcal{T}_{k-1}$ and every $i, j \notin I$ with $i \neq j$,

$$\text{sign}[i, [jI]] \text{sign}[j, [iI]] = -\text{sign}[i, I] \text{sign}[j, I] \quad (\text{B.3.3})$$

Proof. We start with the first equation. Suppose that $i = i_r$ and $j = i_s$ for some r and s , such that (without loss of generality) $i < j$. Then we have $I = i_1 < i_2 < \dots < i_r = i < \dots < i_s = j < i_{k+1}$. Then $\text{sign}[i, I_{\hat{i}}] = (-1)^{r-1}$ since to move $i = i_r$ across $r-1$ positions requires $r-1$ transpositions. Likewise $\text{sign}[j, I_{\hat{j}}] = (-1)^{s-1}$. Therefore the right hand side gives $(-1)^{r-1}(-1)^{s-1}$.

Since $I_{\hat{i}j} = i_1 < i_2 < \dots < i_{r-1} < i_{r+1} < \dots < i_{s-1} < i_{s+1} < \dots < i_{k+1}$, $\text{sign}[i, I_{\hat{i}j}] = (-1)^{r-1}$ but on the other hand $\text{sign}[j, I_{\hat{i}j}] = (-1)^{s-2}$ (Since there is one less swap due to the absence of i). Therefore the left hand side reads $(-1)^{r-1}(-1)^{s-1}(-1)$ which gives the identity.

We now prove the second equation. Assume that $i < j$. It is clear that in that case, $\text{sign}[i, [jI]] = \text{sign}[i, I]$ (since we would not have to move i past j in $[i, [jI]]$). But in that case, $\text{sign}[j, [iI]]$ and $\text{sign}[j, I]$ would differ by one extra transposition, since j would have to be moved past i in $[j, [iI]]$. Therefore, $\text{sign}[j, [iI]] = -\text{sign}[j, I]$ and hence the identity follows. ■

Definition B.3.8. (The space of k -forms) Let Ω be an open set in $\mathbb{R}^n$. The space of k -forms is denoted by

$$\{\Omega; \Lambda^k(T\Omega)\}$$

We also write the space of k -forms as

$$\Omega(\Lambda^k)$$

Definition B.3.9. (Exterior derivative d) Let $\omega \in \{\Omega; \Lambda^k(T\Omega)\}$ be a k -form given by (see B.1.3)

$$\omega_p \equiv \sum_{I \in \mathcal{T}_k} \omega_p^I dx_p^I$$

$d : \{\Omega; \Lambda^k(T\Omega)\} \rightarrow \{\Omega; \Lambda^{k+1}\Omega\}$ is the function given by (formally)

$$d\omega_p := \left(\sum_{i=1}^n \frac{\partial}{\partial x^i} \Big|_p dx_p^i \right) \wedge \omega_p \quad (\text{B.3.4})$$

which is

$$\sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \frac{\partial \omega^I}{\partial x^i} \Big|_p dx_p^i \wedge dx_p^I = \sum_{I \in \mathcal{T}_k} d\omega_p^I \wedge dx_p^I \quad (\text{B.3.5})$$

Lemma B.3.10. *Let $\omega = \sum_{I \in \mathcal{T}_k} \omega^I dx^I$ be a k -form. Then*

$$d\omega = \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \frac{\partial \omega^I}{\partial x^i} dx^i \wedge dx^I = \sum_{I \in \mathcal{T}_{k+1}} \sum_{\gamma=1}^{k+1} (-1)^{\gamma-1} \frac{\partial \omega^{I_{i_\gamma}}}{\partial x^{i_\gamma}} dx^I \quad (\text{B.3.6})$$

Definition B.3.11. (Antiderivation) Let $A = \oplus_{k=0}^\infty A^k$ be a graded algebra over a field K . An antiderivation is a map $D : A \rightarrow A$ such that

1. D is K -linear
2. For all $a \in A^k, b \in A^l$,

$$D(ab) = (D(a))b + (-1)^k a(D(b))$$

D is said to be an antiderivation of degree m if $D|_{A^k} : A^k \rightarrow A^{k+m}$

Theorem B.3.12. $d : \oplus_{k=0}^\infty \Omega(\Lambda^k) \rightarrow \oplus_{k=0}^\infty \Omega(\Lambda^k)$ is an antiderivation of degree 1 and

$$d(\omega \wedge \tau) = d\omega \wedge \tau + (-1)^k \omega \wedge d\tau$$

for all $\omega \in \Omega(\Lambda^k)$ and $\tau \in \Omega(\Lambda^l)$

Proof. That d is linear is clear from formula B.3.5. Let $\omega \equiv \sum_{I \in \mathcal{T}_k} \omega^I dx^I$ and $\tau \equiv \sum_{J \in \mathcal{T}_l} \tau^J dx^J$. $\omega \wedge \tau \equiv \sum_{I \in \mathcal{T}_k} \sum_{J \in \mathcal{T}_l} \omega^I \tau^J dx^I \wedge dx^J$ Applying d gives us

$$\begin{aligned} d(\omega \wedge \tau) &= \left(\frac{\partial}{\partial x^i} \Big|_p dx_p^i \right) \wedge \left(\sum_{I \in \mathcal{T}_k} \sum_{J \in \mathcal{T}_l} \omega_p^I \tau_p^J dx_p^I \wedge dx_p^J \right) = \\ &= \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \sum_{J \in \mathcal{T}_l} \left(\frac{\partial \omega^I}{\partial x^i} \Big|_p \tau_p^J + \omega_p^I \frac{\partial \tau^J}{\partial x^i} \Big|_p \right) dx_p^i \wedge dx_p^I \wedge dx_p^J \\ &= d\omega \wedge \tau + \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \sum_{J \in \mathcal{T}_l} \omega_p^I \frac{\partial \tau^J}{\partial x^i} \Big|_p dx_p^i \wedge dx_p^I \wedge dx_p^J \\ &= d\omega \wedge \tau + \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \sum_{J \in \mathcal{T}_l} (-1)^k \omega_p^I \frac{\partial \tau^J}{\partial x^i} \Big|_p dx_p^I \wedge dx_p^i \wedge dx_p^J \\ &= d\omega \wedge \tau + (-1)^k \omega \wedge d\tau \end{aligned}$$

Lemma B.3.13. $d^2 = 0$

Proof.

$$d\omega = \sum_{i=1}^n \sum_{I \in \mathcal{T}_k} \frac{\partial \omega^I}{\partial x^i} dx^i \wedge dx^I$$

and

$$d(d\omega) = \left(\sum_{i=1}^n \frac{\partial}{\partial x^i} dx^i \right) \wedge \left(\sum_{j=1}^n \sum_{I \in \mathcal{T}_k} \frac{\partial \omega^I}{\partial x^j} dx^j \wedge dx^I \right)$$

$$= \sum_{i=1}^n \sum_{j=1}^n \sum_{I \in \mathcal{T}_k} \frac{\partial^2 \omega^I}{\partial x^i \partial x^j} dx^i \wedge dx^j \wedge dx^I$$

Since $dx^i \wedge dx^j = -dx^j \wedge dx^i$ and $\frac{\partial^2 \omega^I}{\partial x^i \partial x^j} = \frac{\partial^2 \omega^I}{\partial x^j \partial x^i}$ we have that $d(d\omega) = 0$.

B.3.4 Closed and Exact Forms

Definition B.3.14. (Closed Form) Let $\omega \in \Omega(\Lambda^k)$ be a k -form. We say ω is *closed* if $d\omega = 0$.

Definition B.3.15. (Exact Form) We say k -form $\omega \in \Omega(\Lambda^k)$ is *exact* if there exists $\lambda \in \Omega(\Lambda^{k-1})$ such that $d\lambda = \omega$.

Lemma B.3.16. Exact forms are closed.

B.4 Vector Calculus

We set $n = 3$, $U \subset \mathbb{R}^3$ and $k \in \{1, 2, 3\}$.

1. **One forms:** $\omega : U \rightarrow \Pi_{p \in U} T_p^*(\mathbb{R}^3)$ with $\omega_{(p)} \in \Lambda^1(T_p(\mathbb{R}^3)) \equiv T_p^*(\mathbb{R}^3)$ given by

$$\omega_{(p)} \equiv \omega_{(p)}^x dx_{(p)} + \omega_{(p)}^y dy_{(p)} + \omega_{(p)}^z dz_{(p)}$$

2. **Two forms:** $\omega : U \rightarrow \Pi_{p \in U} \Lambda^2(T_p(\mathbb{R}^3))$ with $\omega_{(p)} \in \Lambda^2(T_p(\mathbb{R}^3))$ given by

$$\begin{aligned} \omega_{(p)} &\equiv \sum_{1 \leq i_1 < i_2 \leq 3} \omega_{(p)}^{i_1 i_2} dx_{(p)}^{i_1} \wedge dx_{(p)}^{i_2} \\ &= \omega_{(p)}^z dx_{(p)} \wedge dy_{(p)} + \omega_{(p)}^x dy_{(p)} \wedge dz_{(p)} + \omega_{(p)}^y dz_{(p)} \wedge dx_{(p)} \end{aligned}$$

3. **Three forms:** $\omega : U \rightarrow \Pi_{p \in U} \Lambda^3(T_p(\mathbb{R}^3))$ with $\omega_{(p)} \in \Lambda^3(T_p(\mathbb{R}^3))$ given by

$$\omega_{(p)} = \omega_{(p)} dx_{(p)} \wedge dy_{(p)} \wedge dz_{(p)}$$

We immediately see a correspondence in table B.1

Table B.1: Correspondence between vector fields and forms.

Field	Form
Vector field: $[\omega^x, \omega^y, \omega^z]$	1-form: $\omega_{(p)}^x dx_{(p)} + \omega_{(p)}^y dy_{(p)} + \omega_{(p)}^z dz_{(p)}$
Vector field: $[\omega^x, \omega^y, \omega^z]$	2-form: $\omega_{(p)}^x dy_{(p)} \wedge dz_{(p)} + \omega_{(p)}^y dz_{(p)} \wedge dx_{(p)} + \omega_{(p)}^z dx_{(p)} \wedge dy_{(p)}$
Scalar field: ω	3-form: $\omega_{(p)} dx_{(p)} \wedge dy_{(p)} \wedge dz_{(p)}$

B.4.1 Exterior Derivative of forms in $\mathbb{R}^3$

0-forms in $\mathbb{R}^3$

Let $f : U \rightarrow \mathbb{R}$. We know that

$$df : U \rightarrow \Pi_{p \in U} \Lambda^1(T_p(U))$$

given by

$$df_{(p)} = \frac{\partial f}{\partial x} \Big|_{(p)} dx_{(p)} + \frac{\partial f}{\partial y} \Big|_{(p)} dy_{(p)} + \frac{\partial f}{\partial z} \Big|_{(p)} dz_{(p)}$$

Which, from the correspondence B.1, can be identified with the vector field

$$\left[\frac{\partial f}{\partial x} \Big|_{(p)}, \frac{\partial f}{\partial y} \Big|_{(p)}, \frac{\partial f}{\partial z} \Big|_{(p)} \right] \equiv \nabla f(p)$$

1-forms in $\mathbb{R}^3$

Let $\omega \in \Omega(\Lambda^1(T\mathbb{R}^3))$ be a 1-form, i.e.

$$\omega_{(p)} \equiv \omega_{(p)}^x dx_{(p)} + \omega_{(p)}^y dy_{(p)} + \omega_{(p)}^z dz_{(p)}$$

$$\begin{aligned} \text{(Symbolically)} \quad d\omega_p &= \left(\sum_{i=1}^3 \frac{\partial}{\partial x^i} \bigg|_p dx_p^i \right) \wedge \omega_{(p)}^x dx_{(p)} + \omega_{(p)}^y dy_{(p)} + \omega_{(p)}^z dz_{(p)} \\ &= \left(\frac{\partial}{\partial x} \bigg|_p dx_p + \frac{\partial}{\partial y} \bigg|_p dy_p + \frac{\partial}{\partial z} \bigg|_p dz_p \right) \wedge \omega_{(p)}^x dx_{(p)} + \omega_{(p)}^y dy_{(p)} + \omega_{(p)}^z dz_{(p)} \\ &= \left[\frac{\partial \omega^y}{\partial x} \bigg|_p dx_p \wedge dy_p + \frac{\partial \omega^z}{\partial x} \bigg|_p dx_p \wedge dz_p \right] + \left[\frac{\partial \omega^x}{\partial y} \bigg|_p dy_p \wedge dx_p + \frac{\partial \omega^z}{\partial y} \bigg|_p dy_p \wedge dz_p \right] + \left[\frac{\partial \omega^x}{\partial z} \bigg|_p dz_p \wedge dx_p + \frac{\partial \omega^y}{\partial z} \bigg|_p dz_p \wedge dy_p \right] = \\ &= \left(\frac{\partial \omega^z}{\partial y} \bigg|_p - \frac{\partial \omega^y}{\partial z} \bigg|_p \right) dy_p \wedge dz_p + \left(\frac{\partial \omega^y}{\partial x} \bigg|_p - \frac{\partial \omega^x}{\partial y} \bigg|_p \right) dx_p \wedge dy_p + \left(\frac{\partial \omega^x}{\partial z} \bigg|_p - \frac{\partial \omega^z}{\partial x} \bigg|_p \right) dz_p \wedge dx_p \end{aligned}$$

which from the correspondence B.1 can be identified with the vector field

$$\left[\left(\frac{\partial \omega^z}{\partial y} \bigg|_p - \frac{\partial \omega^y}{\partial z} \bigg|_p \right), \left(\frac{\partial \omega^x}{\partial z} \bigg|_p - \frac{\partial \omega^z}{\partial x} \bigg|_p \right), \left(\frac{\partial \omega^y}{\partial x} \bigg|_p - \frac{\partial \omega^x}{\partial y} \bigg|_p \right) \right] \equiv \nabla \times \omega(p)$$

2-forms in $\mathbb{R}^3$

Let $\omega \in \Omega(\Lambda^2(T\mathbb{R}^3))$ be a 2-form i.e.

$$\omega_p \equiv \omega_p^x dy_p \wedge dz_p + \omega_p^y dz_p \wedge dx_p + \omega_p^z dx_p \wedge dy_p \in \Lambda^2(T_p(\mathbb{R}^3))$$

$$d(\omega^x dy \wedge dz)_p = \frac{\partial \omega^x}{\partial x} \bigg|_p dx_p \wedge dy_p \wedge dz_p$$

$$d(\omega^y dz \wedge dx)_p = \frac{\partial \omega^y}{\partial y} \bigg|_p dy_p \wedge dz_p \wedge dx_p$$

$$d(\omega^z dx \wedge dy)_p = \frac{\partial \omega^z}{\partial z} \bigg|_p dz_p \wedge dx_p \wedge dy_p$$

and hence

$$d\omega_p \equiv \left(\frac{\partial \omega^x}{\partial x} \bigg|_p + \frac{\partial \omega^y}{\partial y} \bigg|_p + \frac{\partial \omega^z}{\partial z} \bigg|_p \right) dx_p \wedge dy_p \wedge dz_p$$

Which from the correspondence B.1 can be identified with the scalar field

$$\frac{\partial \omega^x}{\partial x} \bigg|_p + \frac{\partial \omega^y}{\partial y} \bigg|_p + \frac{\partial \omega^z}{\partial z} \bigg|_p \equiv \nabla \cdot \omega(p)$$

From these correspondences, we see that $d : \Omega(\Lambda^0) \rightarrow \Omega(\Lambda^1)$ is the gradient, $d : \Omega(\Lambda^1) \rightarrow \Omega(\Lambda^2)$ the curl and $d : \Omega(\Lambda^2) \rightarrow \Omega(\Lambda^3)$ is the divergence. From this we can see immediately that the well known identities

$$\nabla \times (\nabla f) = 0$$

and

$$\nabla \cdot (\nabla \times \omega) = 0$$

are just special cases of $d^2 = 0$.

B.5 Interior Product

We start with the observation that $\{dx^I : I \in \mathcal{T}_k\}$ not only forms a basis for $\Lambda^k(V)$, but is also an orthonormal set:

$$\langle dx^I, dx^J \rangle = \begin{cases} 1 & \text{if } I = J \\ 0 & \text{if } I \neq J \end{cases}$$

This is a natural consequence of the definition $\langle f, g \rangle = \sum_{I \in \mathcal{T}_k} f^I g^I$. In the case of a Riemannian manifold with a given $g_p : T_p M \times T_p M \rightarrow \mathbb{R}$, there is a natural choice for $\langle f, g \rangle_g$ given by

$$\langle f, g \rangle_g := \sum_{I, J \in \mathcal{T}_k} f^I g^J \det(g_p^{i_a, j_b}) \quad (\text{B.5.1})$$

where $\{g^{pq}\}$ is the inverse of the matrix $\{g_{pq}\}$. We work predominantly in $\mathbb{R}^n$ with $g_{ij} = \delta_{ij}$ unless stated otherwise. The wedge product

$$\wedge : \Lambda^k(V) \times \Lambda^l(V) \rightarrow \Lambda^{k+l}(V)$$

is bilinear. Suppose we have a fixed k form $\alpha \in \Lambda^k(V)$, then

$$\alpha \wedge () : \Lambda^l(V) \rightarrow \Lambda^{k+l}(V)$$

is a linear map that *raises* the degree of the form. For a k -form α , we have the natural dual from the inner product given by:

$$\langle (\alpha \wedge) \omega, \beta \rangle = \langle \omega, (\alpha \lrcorner) \beta \rangle$$

where $\alpha \in \Lambda^k(V)$, $\omega \in \Lambda^{l-k}(V)$ and $\beta \in \Lambda^l$. From this definition it is clear that

$$\alpha \lrcorner : \Lambda^l(V) \rightarrow \Lambda^{l-k}(V)$$

Corollary B.5.1. If $k \geq l$ and $\alpha \in \Lambda^k(V)$ and $\beta \in \Lambda^l(V)$, then $\alpha \lrcorner \beta = 0$.

On *basis elements*, we have for $dx^I \in \Lambda^k(V)$, $dx^J \in \Lambda^l(V)$ and $dx^w \in \Lambda^{l-k}(V)$

$$\langle dx^I \wedge dx^w, dx^J \rangle = \langle dx^w, dx^I \lrcorner dx^J \rangle$$

Proposition B.5.2. For $dx^I \in \Lambda^k(V)$, $dx^J \in \Lambda^l(V)$ and $l > k$, we have

$$dx^I \lrcorner dx^J = \begin{cases} \text{sign}(R(I, J/I)) dx^{J/I} & \text{if } I \subseteq J \\ 0 & \text{if } I \not\subseteq J \end{cases}$$

where J/I denotes the ordered set of indices that are in J but not in I and $R(I, J/I)$ denotes the permutation that reorders the tuple $(I, J/I)$.

Proof. **When $I \not\subseteq J$,** we have some index of dx^I not be in dx^J . Therefore, no matter our choice of dx^w , we must have that

$$\langle dx^I \wedge dx^w, dx^J \rangle = 0 = \langle dx^w, dx^I \lrcorner dx^J \rangle$$

which means that $dx^I \lrcorner dx^J = 0$.

When $I \subseteq J$, If dx^w does not contain all the complementary elements of J/I , we must have that $\langle dx^I \wedge dx^w, dx^J \rangle = \langle dx^w, dx^I \lrcorner dx^J \rangle = 0$. This means that $dx^I \lrcorner dx^J$ contains indices exclusively from J/I (upto reordering). Let $dx^I \lrcorner dx^J = \lambda dx^{J/I}$. Choose $dx^w = dx^{J/I}$. Note in that case that $dx^I \wedge dx^w = \text{sgn}(R(I, J/I)) dx^J$ where $R(I, J/I)$ is the permutation that rearranges $(I, J/I)$ to the canonically ordered tuple J .

We therefore have

$$\begin{aligned} \langle dx^I \wedge dx^w, dx^J \rangle &= \text{sgn}(R(I, J/I)) \langle dx^J, dx^J \rangle = \text{sgn}(R(I, J/I)) \\ &= \langle dx^{J/I}, \lambda dx^{J/I} \rangle = \lambda \end{aligned}$$

This implies $\lambda = \text{sgn}(R(I, J/I))$ and the proof is complete. ■

Let $\alpha \in \Lambda^k(V)$, $\beta \in \Lambda^l(V)$. For $\omega \in \Lambda^m$ $L_{\alpha \wedge \beta}(\omega) = \alpha \wedge (\beta \wedge \omega) = L_\alpha \circ L_\beta(\omega)$. Therefore, the Adjoint must follow

$$L_{\alpha \wedge \beta}^\dagger \equiv (L_\alpha \circ L_\beta)^\dagger = L_\beta^\dagger \circ L_\alpha^\dagger$$

which implies

$$(\alpha \wedge \beta) \lrcorner \gamma = \beta \lrcorner (\alpha \lrcorner \gamma)$$

B.5.1 The Musical Isomorphism and Interior Product

The situation in $\mathbb{R}^n$:

Given a C^∞ vector field $X : U \rightarrow \Pi_{p \in U} T_p(U)$ given by

$$X_{(p)} = \sum_{i=1}^n a^i(p) \frac{\partial}{\partial x^i} \big|_p \in T_p(U)$$

we can find a differential 1-form given by

$$p \mapsto \sum_{i=1}^n a^i(p) dx_{(p)}^i \quad (\text{B.5.2})$$

The $\flat$ map

For any general Manifold, the vector spaces $T_p M$ and $T_p^* M$ are merely of the same dimension, but there is no global way to identify $T_p M$ with $T_p^* M$. A metric changes this.

Let $g_p : T_p M \times T_p M \rightarrow \mathbb{R}$ be an inner product on $T_p M$. For a given vector $X \in T_p M$, define $X^\flat \in \Lambda^1(T_p(M))$ given by

$$X^\flat(Y) = g(X, Y) \quad (\text{B.5.3})$$

for $Y \in T_p(M)$. Therefore, $\flat : T_p(M) \rightarrow T_p^*(M)$. This map is linear, and injective since $X^\flat(Y) = g(X, Y) = 0$, for every $Y \in T_p(M)$ implies $X = 0$. Hence, $\flat : T_p M \rightarrow T_p^* M$ is an isomorphism.

Let $X \equiv \sum_{i=1}^n X^i \frac{\partial}{\partial x^i}$ be a vector field. g being an inner product, is a 2-tensor on $T_p(M)$.

By B.1.4 we see then, that

$$g \equiv \sum_{1 \leq i, j \leq n} g_{ij}(p) dx_p^i \otimes dx_p^j : T_p M \times T_p M \rightarrow \mathbb{R}$$

and B.5.3 tells us that

$$X_p^\flat = \sum_{1 \leq i, j \leq n} g_{ij}(p) \sum_{s=1}^n a^s(p) dx_p^i \left(\frac{\partial}{\partial x^s} \big|_p \right) dx_p^j = \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} g_{ij}(p) a^i(p) dx_p^j \in \Lambda^1(T_p(M))$$

If we assume that $g_{ij} = \delta_{ij}$ we get back B.5.2.

On Basis: If $X \equiv \frac{\partial}{\partial x^q}$, then $X^\flat = g(X, ())$ given by

$$\sum_{i=1}^n \sum_{j=1}^n g_{ij} dx^i \left(\frac{\partial}{\partial x^q} \right) dx^j = \sum_{j=1}^n g_{qj} dx^j$$

Therefore,

$$\left(\frac{\partial}{\partial x^i} \right)^\flat \equiv \sum_{j=1}^n g_{ij} dx^j$$

If our metric g were Euclidean ($g_{ij} = \delta_{ij}$) we would have that

$$\left(\frac{\partial}{\partial x^i} \right)^\flat \equiv dx^i$$

The $\sharp$ map

We know that $\flat : T_p M \rightarrow T_p^* M$ is given by $(\frac{\partial}{\partial x^i}|_p)^\flat = \sum_{j=1}^n g_{ij} dx_p^j$, so we have the matrix representation of $\flat$ as

$$[X^\flat]_{\{dx_p^1, dx_p^2, \dots, dx_p^n\}} = \begin{bmatrix} g_{11} & g_{12} & \cdots & g_{1n} \\ g_{12} & g_{22} & \cdots & g_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ g_{1n} & g_{2n} & \cdots & g_{nn} \end{bmatrix} [X]_{\{\partial_{x^1}, \partial_{x^2}, \dots, \partial_{x^n}\}} = G_p [X]_{\{\partial_{x^1}, \partial_{x^2}, \dots, \partial_{x^n}\}}$$

from which we get the inverse map $\sharp : T_p^* M \rightarrow T_p M$ by

$$(dx_p^i)^\sharp = \sum_{j=1}^n (G_p)^{-1}_{ij} \frac{\partial}{\partial x^j} \Big|_p \in T_p M$$

Interior product of a vector and a k -form (in $\mathbb{R}^n$)

Let $X \equiv \frac{\partial}{\partial x^q}$. In $\mathbb{R}^n$ we have $X^\flat = dx^q$. For a basis k -form $dx^{i_1} \wedge dx^{i_2} \cdots \wedge dx^{i_k}$,

$$\begin{aligned} X^\flat \lrcorner (dx^{i_1} \wedge dx^{i_2} \cdots \wedge dx^{i_k}) &= dx^q \lrcorner (dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_k}) \\ &= \begin{cases} (-1)^{r-1} dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge \widehat{dx^{i_r}} \wedge \cdots \wedge dx^{i_k} & \text{whenever } q = i_r \\ 0 & \text{whenever } q \notin \{i_1, i_2, \dots, i_k\} \end{cases} \end{aligned} \quad (\text{B.5.4})$$

Definition B.5.3. (Interior product of a vector with a form $(\iota_X(\omega))$) Let $\omega \in \Omega(\Lambda^k)$ be a k -form. Let X be a vector field. $\iota_X : \Omega(\Lambda^k) \rightarrow \Omega(\Lambda^{k-1})$ is given by

$$\iota_X(\omega)(X_1, X_2, \dots, X_{k-1}) = \omega(X, X_1, X_2, \dots, X_{k-1})$$

Remark B.5.4. We see that $\iota_X(\omega)$ is a $k-1$ form. Hence, its action on basis vectors $\frac{\partial}{\partial x^i}$ tells us its action on arbitrary vectors by linearity. Moreover, $\iota_{X+cY}(\omega) = \iota_X(\omega) + c\iota_Y(\omega)$. So its action on $X = \frac{\partial}{\partial x^q}$ tells us its action on arbitrary X by linearity.

Theorem B.5.5. In $\mathbb{R}^n$ equipped with the euclidean metric, we have

$$\iota_X(\omega) = X^\flat \lrcorner \omega$$

(B.5.4 and B.5.3 are equivalent.)

Proof. By B.5.4 it is enough to check equality on basis vectors. Let $X \equiv \frac{\partial}{\partial x^q}$ and let $\omega = dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_k}$ be a k -form. Let $X_1 = \frac{\partial}{\partial x^{j_1}}, X_2 = \frac{\partial}{\partial x^{j_2}}, \dots, X_{k-1} = \frac{\partial}{\partial x^{j_{k-1}}}$ be $k-1$ subcollection of basis vectors. From B.5.4 we have

$$\begin{aligned} X^\flat \lrcorner (dx^{i_1} \wedge dx^{i_2} \cdots \wedge dx^{i_k}) &= \begin{cases} (-1)^{r-1} dx^{i_1} \wedge \cdots \wedge \widehat{dx^{i_r}} \wedge \cdots \wedge dx^{i_k} & \text{whenever } q = i_r \\ 0 & \text{whenever } q \notin \{i_1, i_2, \dots, i_k\} \end{cases} \end{aligned}$$

So when evaluated on $\frac{\partial}{\partial x^{j_1}}, \frac{\partial}{\partial x^{j_2}}, \dots, \frac{\partial}{\partial x^{j_{k-1}}}$ we have

$$\begin{aligned} X^\flat \lrcorner (dx^{i_1} \wedge \cdots \wedge dx^{i_k})(\partial_{x^{j_1}}, \partial_{x^{j_2}}, \dots, \partial_{x^{j_{k-1}}}) &= \\ = \begin{cases} (-1)^{r-1} \det(\{dx^{i_s}(\partial_{x^{j_t}}) : s \neq r, 1 \leq t \leq k-1\}) & \text{whenever } q = i_r \\ 0 & \text{whenever } q \notin \{i_1, i_2, \dots, i_k\} \end{cases} \end{aligned}$$

Let us look at $\iota_{\partial_{x^q}}(dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_k})(\partial_{x^{j_1}}, \partial_{x^{j_2}}, \dots, \partial_{x^{j_{k-1}}}) = \det(D)$ where D is the matrix

$$\begin{bmatrix} dx^{i_1}(\partial_{x^q}) & dx^{i_1}(\partial_{x^{j_1}}) & \cdots & dx^{i_1}(\partial_{x^{j_{k-1}}}) \\ dx^{i_2}(\partial_{x^q}) & dx^{i_2}(\partial_{x^{j_1}}) & \cdots & dx^{i_2}(\partial_{x^{j_{k-1}}}) \\ \vdots & \vdots & \ddots & \vdots \\ dx^{i_r}(\partial_{x^q}) & dx^{i_r}(\partial_{x^{j_1}}) & \cdots & dx^{i_r}(\partial_{x^{j_{k-1}}}) \\ \vdots & \vdots & \ddots & \vdots \\ dx^{i_k}(\partial_{x^q}) & dx^{i_k}(\partial_{x^{j_1}}) & \cdots & dx^{i_k}(\partial_{x^{j_{k-1}}}) \end{bmatrix}$$

Assuming that $q = i_r$ we see that D takes the form:

$$\begin{bmatrix} 0 & dx^{i_1}(\partial_{x^{j_1}}) & \cdots & dx^{i_1}(\partial_{x^{j_{k-1}}}) \\ 0 & dx^{i_2}(\partial_{x^{j_1}}) & \cdots & dx^{i_2}(\partial_{x^{j_{k-1}}}) \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & dx^{i_k}(\partial_{x^{j_1}}) & \cdots & dx^{i_k}(\partial_{x^{j_{k-1}}}) \end{bmatrix}$$

Whence we see that the determinant takes the following form:

$$(-1)^{r-1} \det(\{dx^{i_s}(\partial_{x^{j_t}}) : s \neq r, 1 \leq t \leq k-1\})$$

which agrees with $X^{\flat} \lrcorner (dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_k})$.

The case $q \notin \{i_1, i_2, \dots, i_k\}$ follows since $q \neq i_r$ implies that the first column in D full of 0, which makes the determinant 0.

Lemma B.5.6. (Expression for interior product of a 1-form and a k -form) Let $\nu = \sum_{j=1}^n v^j dx^j \in \Lambda^1$ and $\omega \in \Lambda^k$. We have

$$\nu \lrcorner \omega = \sum_{j=1}^n \sum_{I \in \mathcal{T}_k} v^j \omega^I dx^j \lrcorner dx^I = \sum_{I \in \mathcal{T}_k} \sum_{j \in I} v^j \omega^I (\text{sign}[j, I_j]) dx^{I_j} = \sum_{j=1}^n \sum_{I \in \mathcal{T}_{k-1}} v^j \omega^{j i_1 i_2 \dots i_{k-1}} dx^I \quad (\text{B.5.5})$$

Caution: The notation used here is adopted from [CDS18] and [CDK12] wherein, for non-increasing indices, we define

$$\omega^I = \text{sign}([I]) \omega^{[I]} \quad (\text{B.5.6})$$

where I is an arbitrary multi-index set and $[I]$ is (by abuse of notation) both the permutation that rearranges I to be increasing and the increasing index set.

Proof.

$$\nu \lrcorner \omega = \sum_{j=1}^n \sum_{I \in \mathcal{T}_k} v^j \omega^I dx^j \lrcorner dx^I = \sum_{I \in \mathcal{T}_k} \sum_{j \in I} v^j \omega^I (\text{sign}[j, I_j]) dx^{I_j} \quad (\text{B.5.7})$$

Note that, for every $K \in \mathcal{T}_{k-1}$ and $j \notin K$, we have a unique $I \in \mathcal{T}_k$ given by $[jK]$ so that $K = I_j$. We now argue that the expression above is equal to the following:

$$\sum_{j=1}^n \sum_{K \in \mathcal{T}_{k-1}} v^j \omega^{[jK]} \text{sign}[j, K] dx^K$$

For a given $j \in \{1, \dots, n\}$, $dx^j \lrcorner dx^I \neq 0$ only for those $I \in \mathcal{T}_k$ with $j \in I$. And for each such I , there is a unique $K \in \mathcal{T}_{k-1}$ such that $I = [jK]$. Therefore we can write the left hand side of B.5.7 as

$$\sum_{j=1}^n \sum_{K \in \mathcal{T}_{k-1}} v^j \omega^{[jK]} dx^j \lrcorner dx^{[jK]} = \sum_{j=1}^n \sum_{K \in \mathcal{T}_{k-1}} v^j \omega^{[jK]} \text{sign}[j, K] dx^K$$

If we further impose the notation in (B.5.6), we get

$$\sum_{j=1}^n \sum_{K \in \mathcal{T}_{k-1}} v^j \omega^{j, K} dx^K$$

which are the desired results.

Theorem B.5.7. Let X be a vector field, $\omega \in \Omega\Lambda^k$ and $\eta \in \Omega\Lambda^l$. We then have

$$\iota_X(\omega \wedge \eta) = \iota_X(\omega) \wedge \eta + (-1)^k \omega \wedge \iota_X(\eta) \quad (\text{B.5.8})$$

This tells us that $\iota_X : \oplus_{k=0}^{\infty} \Omega(\Lambda^k) \rightarrow \oplus_{k=0}^{\infty} \Omega(\Lambda^k)$ is a rank -1 antiderivation.

Proof. Let $X_{(p)} = \sum_{q=1}^n a^q(p) \frac{\partial}{\partial x^q} \Big|_p$, $\omega = \sum_{I \in \mathcal{I}_k} \omega^I dx^I$ and $\eta = \sum_{J \in \mathcal{I}_l} \eta^J dx^J$.
 $\iota_X(\omega \wedge \eta) = \sum_{q=1}^n a^q \iota_{\partial_{x^q}}(\omega \wedge \eta) = \sum_{q=1}^n a^q \iota_{\partial_{x^q}}(\sum_{I \in \mathcal{I}_k} \sum_{J \in \mathcal{I}_l} \omega^I \eta^J dx^I \wedge dx^J)$ which gives

$$\iota_X(\omega \wedge \eta) = \sum_{q=1}^n \sum_{I \in \mathcal{I}_k} \sum_{J \in \mathcal{I}_l} a^q \omega^I \eta^J \iota_{\partial_{x^q}}(dx^I \wedge dx^J)$$

If we suppose that the identity holds on basis elements i.e. $X = \partial_{x^q}$, $\omega = dx^I$ and $\eta = dx^J$, we then have

$$\iota_X(\omega \wedge \eta) = \sum_{q=1}^n \sum_{I \in \mathcal{I}_k} \sum_{J \in \mathcal{I}_l} a^q \omega^I \eta^J \left(\iota_{\partial_{x^q}}(dx^I) \wedge dx^J + (-1)^k dx^I \wedge \iota_{\partial_{x^q}}(dx^J) \right)$$

Which gives the result. Hence we need only to prove the identity for $X = \partial_{x^q}$, $\omega = dx^I$ and $\eta = dx^J$.

Case 1: $q \in I$ and $I \cap J = \emptyset$

Suppose that $q = i_r \in I$. We have $\iota_{\partial_{x^q}}(dx^I \wedge dx^J) = \det D$ where

$$D = \begin{bmatrix} dx^{i_1}(\partial_{x^q}) & dx^{i_1}(x_1) & \cdots & dx^{i_1}(x_{k+l-1}) \\ dx^{i_2}(\partial_{x^q}) & dx^{i_2}(x_1) & \cdots & dx^{i_2}(x_{k+l-1}) \\ \vdots & \vdots & \ddots & \vdots \\ dx^{i_r}(\partial_{x^q}) & dx^{i_r}(x_1) & \cdots & dx^{i_r}(x_{k+l-1}) \\ \vdots & \vdots & \ddots & \vdots \\ dx^{j_1}(\partial_{x^q}) & dx^{j_1}(x_1) & \cdots & dx^{j_1}(x_{k+l-1}) \\ \vdots & \vdots & \ddots & \vdots \\ dx^{j_l}(\partial_{x^q}) & dx^{j_l}(x_1) & \cdots & dx^{j_l}(x_{k+l-1}) \end{bmatrix}$$

which evaluates to (from Laplace's Formula)

$$(-1)^{r-1} dx^{i_1} \wedge dx^{i_2} \wedge \cdots \widehat{dx^{i_r}} \wedge \cdots \wedge dx^{i_k} \wedge dx^{j_1} \wedge \cdots \wedge dx^{j_l}(x_1, x_2, \dots, x_{k+l-1}) = \iota_{\partial_{x^q}}(dx^I) \wedge dx^J$$

If $J \cap I = \emptyset$ and $q = i_r \in I$, note that $dx^I \wedge \iota_{\partial_{x^q}}(dx^J) = dx^I \wedge (dx^q \lrcorner dx^J) = 0$ since q is not in J (see Formula B.5.1). Therefore, we have the identity in this case.

Case 2: $q \in J$ and $I \cap J = \emptyset$

Let $q = j_s$ for some $j_s \in J$. We have $\iota_{\partial_{x^q}}(dx^I \wedge dx^J) = \det D$ where D is as in case 1, but note that the only non zero element in the first column occurs in the $k+s$ row. The Laplace expansion thus reads

$$(-1)^{k+s-1} dx^{i_1} \wedge dx^{i_2} \wedge \cdots \wedge dx^{i_k} \wedge dx^{j_1} \wedge \cdots \widehat{dx^{j_s}} \wedge \cdots \wedge dx^{j_l}(x_1, x_2, \dots, x_{k+l-1}) = (-1)^k dx^I \wedge \iota_{\partial_{x^q}}(dx^J)$$

If $I \cap J = \emptyset$ and $q = j_s$ for some $j_s \in J$ then note that $\iota_{\partial_{x^q}}(dx^I) \wedge dx^J = (dx^q \lrcorner dx^I) \wedge dx^J = 0$ and likewise we have the identity.

Note that in both case 1 and case 2 we only used the following fact: $q \in I$ (or J) and $q \notin J$ (or $\notin I$).

Case 3: $I \cap J \neq \emptyset$ and $q = i_r = j_s$

Note that the left hand side $\iota_{\partial_{x^q}}(dx^I \wedge dx^J) = 0$. The first expression on the right hand side reads $\iota_{\partial_{x^q}}(dx^I) \wedge dx^J = (dx^q \lrcorner dx^I) \wedge dx^J$ and the second expression reads $(-1)^k dx^I \wedge (dx^q \lrcorner dx^J)$. Note that in case 3, the matrix D we get has two occurrences of 1 (and the rest 0) in the first column: In the r -th and $(k+s)$ -th rows. Laplace's expansion gives us the determinant of D to be

$$(-1)^{r-1} dx^{i_1} \wedge \cdots \widehat{dx^{i_r}} \wedge \cdots \wedge dx^{i_k} \wedge dx^{j_1} \wedge \cdots \wedge dx^{j_l}(x_1, x_2, \dots, x_{k+l-1})$$

$$+(-1)^{k+s-1} dx^{i_1} \wedge dx^{i_2} \wedge \dots \wedge dx^{i_k} \wedge dx^{j_1} \wedge \dots \wedge \widehat{dx^{j_s}} \wedge \dots \wedge dx^{j_l}(x_1, x_2, \dots, x_{k+l-1})$$

which is precisely

$$\iota_{\partial_{x^q}}(dx^I) \wedge dx^J + (-1)^k dx^I \wedge (\iota_{\partial_{x^q}} dx^J)$$

Since the identity is true on basis, it is true in the general case through multilinearity.

Lemma B.5.8. ι is anticommutative, i.e.

$$\iota_X \iota_Y = -\iota_Y \iota_X$$

Proof. This is easy to see since $\iota_X(\omega)(x_1, \dots, x_{k-1}) = \omega(X, x_1, \dots, x_{k-1})$ and $\iota_Y \iota_X(\omega)(x_1, \dots, x_{k-2}) = \iota_X(\omega)(Y, x_1, x_2, \dots, x_{k-2}) = \omega(X, Y, x_1, \dots, x_{k-2}) = -\omega(Y, X, x_1, \dots, x_{k-2})$ and the proof is complete.

Corollary B.5.9. $\iota_X \circ \iota_X = 0$

Theorem B.5.10. Let $\xi \in \Omega(\Lambda^1)$, $\eta \in \Omega(\Lambda^k)$ be 1 and k forms respectively in $\mathbb{R}^n$ (with the Euclidean Metric). Then we have the following identity:

$$\|\xi\|^2 \eta = \xi \wedge (\xi \lrcorner \eta) + \xi \lrcorner (\xi \wedge \eta) \quad (\text{B.5.9})$$

For vector fields we have:

$$|X|^2 \eta = X^\flat \wedge \iota_X(\eta) + \iota_X(X^\flat \wedge \eta)$$

Proof. Using the identity B.5.7 we have $\xi \lrcorner (\xi \wedge \eta) = (\xi \lrcorner \xi) \wedge \eta + (-1)\xi \wedge (\xi \lrcorner \eta)$ which gives

$$\xi \lrcorner (\xi \wedge \eta) + \xi \wedge (\xi \lrcorner \eta) = (\xi \lrcorner \xi) \wedge \eta = \lambda \eta$$

Note that $\xi \wedge 1 = \xi$ and $\langle \xi \wedge 1, \beta \rangle = \langle 1, \xi \lrcorner \beta \rangle$. Choosing $\beta = \xi$ gives

$$\langle 1, \xi \lrcorner \xi \rangle = \xi \lrcorner \xi = \langle \xi, \xi \rangle = \|\xi\|^2$$

So we have the identity.

B.6 Hodge Star

Let f be a k -form given by

$$f \equiv \sum_{I \in \mathcal{T}_k} f^I dx^I$$

We define the operation *Hodge Star* as follows (defined first on basic k -forms, and then extended linearly for any k -form):

$$\star dx^{i_1} \wedge dx^{i_2} \wedge \dots \wedge dx^{i_k} = \text{sign}(R(I, J)) dx^{j_1} \wedge dx^{j_2} \wedge \dots \wedge dx^{j_{n-k}}$$

where

1. $I \equiv \{i_1, i_2, \dots, i_k\} \in \mathcal{T}_k$ and $J \equiv \{j_1, j_2, \dots, j_{n-k}\}$ being the complementary increasing index set to I (i.e. $J \cup I \equiv \{1, 2, \dots, n\}$ where J is also increasingly ordered).
2. $R(I, J) \in \text{sym}(n)$ rearranges $\{i_1, i_2, \dots, i_k, j_1, j_2, \dots, j_{n-k}\}$ in its increasing order.
3. For any $f \equiv \sum_{I \in \mathcal{T}_k} f^I dx^I$ we have

$$\star f := \sum_{I \in \mathcal{T}_k} f^I \star dx^I$$

An alternate way to define the *Hodge Star*: If f and g are k -forms, then $\star g$ is defined as the unique $(n-k)$ -form such that

$$f \wedge \star g = \langle f, g \rangle dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n \quad (\text{B.6.1})$$

where the inner product is

$$\langle f, g \rangle := \sum_{I \in \mathcal{T}_k} f^I g^I$$

Lemma B.6.1. *Let $f \in \Omega(\Lambda^k)$. We then have*

$$\star \star f = (-1)^{k(n-k)} f \quad (\text{B.6.2})$$

Proof. Since $\star$ is linear, it is enough to prove on basis elements.

$$\star dx^I = \text{sign}(R(I, J)) dx^J$$

where J is the complement of I relative to $\{1, 2, \dots, n\}$ and $R(I, J)$ is the permutation that rearranges $\{i_1, i_2, \dots, i_k, j_1, j_2, \dots, j_{n-k}\} \rightarrow \{1, 2, \dots, n\}$.

$$\star \star dx^I = \text{sign}(R(I, J)) \star dx^J = \text{sign}(R(I, J)) \text{sign}(R(J, I)) dx^I$$

To compute the product of the signs we look at the map σ that takes (I, J) , rearranges them to $\{1, 2, \dots, n\}$ and then rearranges them to $\{J, I\}$. This amounts to moving the first k elements behind the last $n - k$ elements. This takes $k(n - k)$ transpositions, and hence the sign is $(-1)^{k(n-k)}$. Therefore,

$$\star \star dx^I = (-1)^{k(n-k)} dx^I$$

and the result follows for a general k -form from linearity. ■

Lemma B.6.2. *For $f, g \in \Omega(\Lambda^k)$*

$$\langle f, g \rangle = \langle \star f, \star g \rangle$$

Proof. Since $f \wedge \star g = \langle f, g \rangle dV$ (where dV is the volume form, in the Euclidean case, $dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n$), we have $\star f \wedge \star \star g = (-1)^{k(n-k)} \star f \wedge g = \langle \star f, \star g \rangle dV$. Recall that $f \wedge g = (-1)^{kl} g \wedge f$ for $f \in \Omega \Lambda^k$ and $g \in \Omega \Lambda^l$. Since $\star f \in \Omega \Lambda^{n-k}$ and $g \in \Omega \Lambda^k$, we have $\star f \wedge g = (-1)^{k(n-k)} g \wedge \star f$. This gives us

$$(-1)^{k(n-k)} \star f \wedge g = g \wedge \star f = \langle \star f, \star g \rangle dV$$

But since $g \wedge \star f := \langle g, f \rangle = \langle f, g \rangle$ (Real inner product), we have the result. ■

B.6.1 Hodge Star and Interior

Lemma B.6.3. *Let ω be a k -form and η an l -form. We have the following identity:*

$$\omega \lrcorner \eta = (-1)^{(n-l)(l-k)} \star (\omega \wedge \star \eta) \quad (\text{B.6.3})$$

If ω were a 1 form, this formula reads:

$$\omega \lrcorner \eta = (-1)^{n(l-1)} \star (\omega \wedge \star \eta) \quad (\text{B.6.4})$$

Proof. Let γ be an arbitrary $(l - k)$ -form.

$$\langle \omega \wedge \gamma, \eta \rangle dV = \omega \wedge \gamma \wedge \star \eta$$

By the anti-commutation rule of the wedge, we have

$$\langle \omega \wedge \gamma, \eta \rangle dV = (-1)^{k(l-k)} \gamma \wedge (\omega \wedge \star \eta)$$

Using the $\star\star$ identity, we can write

$$(\omega \wedge \star\eta) = (-1)^{(k+n-l)(l-k)} \star\star(\omega \wedge \star\eta)$$

which gives us

$$\langle \omega \wedge \gamma, \eta \rangle dV = (-1)^{k(l-k)} (-1)^{(k+n-l)(l-k)} \gamma \wedge \star(\star(\omega \wedge \star\eta))$$

$\star(\omega \wedge \star\eta)$ is a $(l-k)$ form and we can invoke the hodge star definition to get:

$$\langle \omega \wedge \gamma, \eta \rangle dV = (-1)^{k(l-k)} (-1)^{(k+n-l)(l-k)} \langle \gamma, \star(\omega \wedge \star\eta) \rangle dV$$

This gives us

$$\langle \gamma, \omega \lrcorner \eta \rangle = (-1)^{(l-k)(n-l)} \langle \gamma, \star(\omega \wedge \star\eta) \rangle$$

Since γ was arbitrary, we have

$$\omega \lrcorner \eta = (-1)^{(l-k)(n-l)} \star(\omega \wedge \star\eta)$$

■

Corollary B.6.4. Let X be a vector field and ω a k -form.

$$\iota_X(\omega) = (-1)^{(n-k)(k-1)} \star(X^\flat \wedge \star\omega) \equiv (-1)^{n(l-1)} \star(X^\flat \wedge \star\omega)$$

B.7 Brief Look at Riemannian Manifolds

Definition B.7.1. (Riemannian Manifold) A Smooth n -dimensional manifold $(M, \mathcal{A})$ is said to be a *Riemannian Manifold* of dimension n if for every $p \in M$ there exists an inner product $g_p : T_p M \times T_p M \rightarrow \mathbb{R}$ (smoothly depending on p) that is positive definite. Such a g is called the *metric*.

We recall the fact (B.1.4) that the space of covariant-2 tensors has a basis given by $\{dx^i \otimes dx^j : i, j \leq n\}$. This then lets us write the metric in the following useful form:

$$g_p = \sum_{i,j=1}^n g_{ij} dx^i \otimes dx^j \quad (\text{B.7.1})$$

Where $g_{ij} = g(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j})$.

We also recall the musical isomorphisms (B.5.1) here:

The $\flat$ map

Let X be a vector field on M . Define the $\flat$ map that takes vector fields to 1-forms as follows:

$$X^\flat := g(X, \cdot)$$

Let $X \equiv \frac{\partial}{\partial x^q}$, then

$$(\frac{\partial}{\partial x^q})^\flat = \sum_{i \leq n} \sum_{j \leq n} g_{ij} dx^i (\frac{\partial}{\partial x^q}) dx^j = \sum_{j \leq n} g_{qj} dx^j \quad (\text{B.7.2})$$

And for $\omega = \sum_{q=1}^n \omega^q \frac{\partial}{\partial x^q}$ we have

$$(\omega)^\flat = \sum_{j \leq n} (\sum_{q=1}^n g_{qj} \omega^q) dx^j \quad (\text{B.7.3})$$

The $\sharp$ map

Since g was positive-definite, we have that the matrix g_{ij} is invertible. Hence, we have the inverse of the $\flat$ map, called $\sharp$ map that takes 1-forms to vector fields as follows: Let $\alpha = \sum_{i=1}^n \alpha^i dx^i$. Then

$$\alpha^\sharp = \sum_{j=1}^n (\sum_{q=1}^n g^{qj} \alpha^q) \frac{\partial}{\partial x^j} \quad (\text{B.7.4})$$

Where $g^{ij} = ([g_{ij}]^{-1})_{ij}$

B.7.1 Natural Inner Product on Forms, Hodge Star and Interior Product

Let M be an n dimensional Riemannian Manifold with metric g . Let $\alpha = v_1 \wedge v_2 \cdots \wedge v_k$ be a k form generated by wedging 1-forms $v_1, v_2, \dots, v_k$. Likewise let $\beta = w_1 \wedge w_2 \wedge \cdots \wedge w_k$. We define the inner product on k -forms as follows:

$$\langle \alpha, \beta \rangle_g := \det \left(\begin{bmatrix} g(v_1^\sharp, w_1^\sharp) & g(v_1^\sharp, w_2^\sharp) & \cdots & g(v_1^\sharp, w_k^\sharp) \\ g(v_2^\sharp, w_1^\sharp) & g(v_2^\sharp, w_2^\sharp) & \cdots & g(v_2^\sharp, w_k^\sharp) \\ \vdots & \vdots & \ddots & \vdots \\ g(v_k^\sharp, w_1^\sharp) & g(v_k^\sharp, w_2^\sharp) & \cdots & g(v_k^\sharp, w_k^\sharp) \end{bmatrix} \right) \quad (\text{B.7.5})$$

And we extend it bilinearly to all k -forms.

Lemma B.7.2. (A formula for $g(v_1^\sharp, v_2^\sharp)$) Let $v_1 = \sum_{i=1}^n v_1^i dx^i$ and $v_2 = \sum_{j=1}^n v_2^j dx^j$ be 1-forms. We then have that

$$g(v_1^\sharp, v_2^\sharp) = \sum_{i,j=1}^n v_1^i g^{ij} v_2^j \quad (\text{B.7.6})$$

Proof. $v_1^\sharp = \sum_{q=1}^n (\sum_{r_1=1}^n g^{qr_1} v_1^{r_1}) \frac{\partial}{\partial x^q}$ and $v_2^\sharp = \sum_{s=1}^n (\sum_{r_2=1}^n g^{sr_2} v_2^{r_2}) \frac{\partial}{\partial x^s}$. Thus we have

$$g(v_1^\sharp, v_2^\sharp) = \sum_{r_1=1}^n \sum_{r_2=1}^n v_1^{r_1} v_2^{r_2} \left(\sum_{q=1}^n \sum_{s=1}^n g^{qr_1} g^{sr_2} g_{qs} \right) = \sum_{r_1=1}^n \sum_{r_2=1}^n v_1^{r_1} g^{r_1 r_2} v_2^{r_2}$$

Which is the desired result. ■

Orientability

Definition B.7.3. (Orientable Manifold) A differentiable manifold M is said to be orientable if there exists an atlas $\mathcal{A}$ such that for any two charts (U, φ) and (V, ψ) in $\mathcal{A}$, the Jacobian determinant of the transition maps $\psi \circ \varphi^{-1}$ is positive. If we denote our charts as $x = (x^1, x^2, \dots, x^n)$ and $y = (y^1, y^2, \dots, y^n)$ respectively, we can suggestively then write $\det(\frac{\partial x^i}{\partial y^j}) > 0$

Note that, if x and y are coordinate charts around a point p , $T_p M$ has bases $\{\frac{\partial}{\partial x^i}\}_{i \leq n}$ and $\{\frac{\partial}{\partial y^j}\}_{j \leq n}$. Chain rule lets us write $\frac{\partial}{\partial x^i} = \sum_{j=1}^n \frac{\partial y^j}{\partial x^i} \frac{\partial}{\partial y^j}$, and hence the transformation A that takes the basis $\{\frac{\partial}{\partial x^i}\}_{i \leq n}$ to $\{\frac{\partial}{\partial y^j}\}_{j \leq n}$ is of positive determinant.

Orthonormal frames and co-frames

Choose a frame $E_1, E_2, \dots, E_n$ (from Gram-Schmidt process) that forms an orthonormal basis with respect to g , i.e. $g(E_i, E_j) = \delta_{ij}$. Consider the co-frame $\xi^1, \xi^2, \dots, \xi^n$ dual to $E_1, E_2, \dots, E_n$ in $T_p^* M$. $(\xi^i)^\sharp = \sum_j \xi_j^i E_j = \sum_j g(\xi_j^\sharp, E_j) E_j$. Note that $g(\xi_j^\sharp, E_j) = \xi_j^\sharp(E_j) = \xi^j(E_j) = \delta_{ij}$ which gives us that $\xi_j^\sharp = E_j$, and hence we have

$$\langle (\xi^i)^\sharp, (\xi^j)^\sharp \rangle_g = \delta_{ij}$$

Which tells us that the co-frame is also orthonormal with respect to the natural inner product $\langle \cdot, \cdot \rangle_g$. We collect this result in the following lemma:

Lemma B.7.4. If $E_1, E_2, \dots, E_n$ is an orthonormal frame with respect to g , then the dual co-frame $\xi^1, \xi^2, \dots, \xi^n$ is also orthonormal with respect to the natural inner product $\langle \cdot, \cdot \rangle_g$.

These ξ^i in B.7.4 form an orthonormal basis for $T_p^* M$, and hence the collection $\{\xi^{i_1} \wedge \xi^{i_2} \wedge \cdots \wedge \xi^{i_k} : 1 \leq i_1 < i_2 < \cdots < i_k \leq n\}$ forms a basis for $\Lambda^k(T_p M)$. Let $\alpha = \sum_{I \in \mathcal{T}_k} \alpha^I \xi^I$ and $\beta = \sum_{I \in \mathcal{T}_k} \beta^I \xi^I$ be two k -forms written in this basis. Then we have

$$\langle \alpha, \beta \rangle_g = \sum_{I \in \mathcal{T}_k} \sum_{J \in \mathcal{T}_k} \langle \xi^I, \xi^J \rangle_g \alpha^I \beta^J = \sum_{I \in \mathcal{T}_k} \alpha^I \beta^I \quad (\text{B.7.7})$$

The Volume Form

Let M be an orientable manifold and let $E_1, E_2, \dots, E_k$ be a positively oriented orthonormal frame with respect to g . We define the volume form as the unique n form such that:

$$\text{vol}_g(E_1, E_2, \dots, E_n) = 1 \quad (\text{B.7.8})$$

Let $F_1, F_2, \dots, F_n$ be another positively oriented (not necessarily orthonormal) frame. Suppose $\{E_i\}$ are related to $\{F_j\}$ via the orthogonal transformation A , i.e. $F_i = \sum_j A_i^j E_j$. Then we have

$$\text{vol}_g(F_1, F_2, \dots, F_n) = \det(A) \quad (\text{B.7.9})$$

$\det(A)$ can be found as follows:

$$g(F_i, F_k) = g\left(\sum_j A_j^i E_j, \sum_m A_m^k E_m\right) = \sum_j \sum_m A_j^i A_m^k \delta_{jm} = \sum_j A_j^i A_j^k = (A^T A)_{jk}$$

Therefore,

$$\sqrt{\det(\{g(F_i, F_k)\}_{i,k \leq n})} = \det(A)$$

Let $x = (x^1, x^2, \dots, x^n)$ and $y = (y^1, y^2, \dots, y^n)$ be two coordinate charts around $p \in M$, such that $\{\frac{\partial}{\partial x^i}\}_{i \leq n}$ and $\{\frac{\partial}{\partial y^j}\}_{j \leq n}$ are both positively oriented. We then have that

$$\text{vol}_g\left(\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}, \dots, \frac{\partial}{\partial x^n}\right) = \sqrt{\det\left(g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right)\right)} \quad (\text{B.7.10})$$

And

$$\text{vol}_g = \sqrt{\det\left(g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right)\right)} dx^1 \wedge dx^2 \wedge \dots \wedge dx^n \quad (\text{B.7.11})$$

Hodge Star on Riemannian Manifolds

Definition B.7.5. (Hodge Star for Riemannian Manifolds) Let $\beta \in \Lambda^k$ be a k -form on an orientable Riemannian manifold (M, g) . We define the Hodge Star $\star\beta$ as the *unique* $n - k$ form such that for every test k form α the following holds:

$$\alpha \wedge \star\beta = \langle \alpha, \beta \rangle_g \text{vol}_g \quad (\text{B.7.12})$$

Lemma B.7.6. Let $E_1, E_2, \dots, E_n$ be a given positively oriented orthonormal frame. Let $\xi^1, \xi^2, \dots, \xi^n$ be the dual co-frame. Then

$$\star(\xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k}) = \text{sign}(R(I, I^c)) \xi^{j_1} \wedge \xi^{j_2} \wedge \dots \wedge \xi^{j_{n-k}} \quad (\text{B.7.13})$$

Where $R(I, I^c)$ is the permutation that rearranges $\{I, I^c\}$ to its natural order, and $j_1, j_2, \dots, j_{n-k}$ are the canonically ordered indices of I^c .

Proof. The proof follows from the following observations:

1. The volume form is $\xi^1 \wedge \xi^2 \wedge \dots \wedge \xi^n$
2. $\text{sign}(R(I, I^c)) \xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k} \wedge \xi^{j_1} \wedge \dots \wedge \xi^{j_{n-k}} = \text{vol}_g$.
3. $\langle \xi^{i_1} \wedge \dots \wedge \xi^{i_k}, \xi^{i_1} \wedge \dots \wedge \xi^{i_k} \rangle_g = 1$

Theorem B.7.7. (Double Star Identity) Let f be a k -form. Then

$$\star\star f = (-1)^{k(n-k)} f \quad (\text{B.7.14})$$

Proof. Because the star is linear, it is enough to prove the identity on basis elements. Let $E_1, \dots, E_n$ be a positively oriented orthonormal frame and $\xi^1, \dots, \xi^n$ be its dual co-frame. $\star\star(\xi^{i_1} \wedge \xi^{i_2} \wedge \dots \wedge \xi^{i_k}) = \text{sign}(R(I, I^c)) \star(\xi^{I^c}) = \text{sign}(R(I, I^c)) \text{sign}(R(I^c, I)) \xi^I$. The proof is done once we see that the product $\text{sign}(R(I, I^c)) \text{sign}(R(I^c, I))$ is the sign of the permutation that takes swaps the first k terms in $[I, I^c]$ behind the last $n - k$ terms, which amounts to a sign of $(-1)^{k(n-k)}$.

Theorem B.7.8. (Hodge Star is an isometry) The Hodge Star $\star : \Lambda^k \rightarrow \Lambda^{n-k}$ is an isometric isomorphism between Normed Linear Spaces.

Proof. Note that the Hodge Star is linear. Since the dimensions of Λ^k and Λ^{n-k} are the same, it is enough to show that $\star$ is an isometry, i.e. $\langle \star\alpha, \star\beta \rangle_{g, n-k} = \langle \alpha, \beta \rangle_{g, k}$. Let $\xi^1, \xi^2, \dots, \xi^n$ be the co-frame of the orthonormal frame $E_1, E_2, \dots, E_n$. We saw in B.7.4 that these too, are orthonormal with respect to the natural inner product. Let $\alpha = \sum_{I \in \mathcal{T}_k} \alpha^I \xi^I$ and $\beta = \sum_{I \in \mathcal{T}_k} \beta^I \xi^I$. From B.7.7 we have $\langle \alpha, \beta \rangle_g = \sum_{I \in \mathcal{T}_k} \alpha^I \beta^I$. $\langle \star\alpha, \star\beta \rangle = \star\alpha \wedge \star\star\beta = (-1)^{k(n-k)} \star\star\beta \wedge \star\alpha = (-1)^{k(n-k)} \langle \alpha, \star\star\beta \rangle_g = \langle \alpha, \beta \rangle_g$ from the double star identity B.7.7.

Interior Product and Hodge Star

Definition B.7.9. (Interior Product) Let M be an oriented manifold. We define the interior product as the adjoint of the wedge product with respect to the natural inner product, i.e. for every $\alpha \in \Lambda^k$, $\beta \in \Lambda^l$ and $\eta \in \Lambda^{l-k}$

$$\langle \alpha \wedge \eta, \beta \rangle_g = \langle \eta, \alpha \lrcorner \beta \rangle_g \quad (\text{B.7.15})$$

Lemma B.7.10. Let $E_1, E_2, \dots, E_n$ be a positively oriented orthonormal basis and $\xi^1, \xi^2, \dots, \xi^n$ be its dual co-frame. We then have:

$$\xi^I \lrcorner \xi^J = \begin{cases} 0 & \text{if } I \not\subseteq J \\ \text{sign}(R(I, J \setminus I)) \xi^{J \setminus I} & \text{if } I \subseteq J \end{cases}$$

Proof. The proof is the same as the euclidean case B.5.2, since the co-frame is orthonormal with respect to the inner product $\langle \cdot, \cdot \rangle_g$.

Theorem B.7.11. Let $\alpha \in \Lambda^k$, $\beta \in \Lambda^l$. We then have $\alpha \lrcorner \beta = (-1)^{(n-l)(l-k)} \star(\alpha \wedge \star\beta)$

Proof. The proof is the same as the euclidean case B.6.3.

Theorem B.7.12. Let X be a vector field and $\omega \in \Lambda^k$ and $\eta \in \Lambda^l$. We then have

$$\iota_X(\omega \wedge \eta) = \iota_X(\omega) \wedge \eta + (-1)^k \omega \wedge \iota_X(\eta) \quad (\text{B.7.16})$$

Corollary B.7.13. Let $\xi \in \Lambda^1$ and $\eta \in \Lambda^k$. Then we have the following identity:

$$\|\xi\|^2 \eta = \xi \wedge (\xi \lrcorner \eta) + \xi \lrcorner (\xi \wedge \eta) \quad (\text{B.7.17})$$

B.8 Pullback of Forms

Let $F : U \rightarrow V$, where $U \subseteq \mathbb{R}^n$ and $V \subseteq \mathbb{R}^m$. If F is smooth, the Frechet derivative at p is defined as the unique linear map $DF_p : \mathbb{R}^n \rightarrow \mathbb{R}^m$ satisfying

$$\lim_{|h| \rightarrow 0} \frac{\|F(p+h) - F(p) - DF_p(h)\|_m}{\|h\|_n} = 0$$

We have $DF_p : T_p U \rightarrow T_{f(p)} V$ where $DF_p(v) \in T_{f(p)} V$ such that

$$DF_p(v)(g) = v(g \circ F)$$

Where $g \in C_{f(p)}^\infty(V)$. In coordinates we have

$$DF_p\left(\frac{\partial}{\partial x^i}\bigg|_p\right) = \sum_{j=1}^m DF_p\left(\frac{\partial}{\partial x^i}\bigg|_p\right)(y^j) \frac{\partial}{\partial y^j}\bigg|_{F(p)} \quad (\text{B.8.1})$$

which gives

$$DF_p(v) = DF_p\left(\sum_{i=1}^n v^i \frac{\partial}{\partial x^i}\bigg|_p\right) = \sum_{i=1}^n \sum_{j=1}^m DF_p\left(\frac{\partial}{\partial x^i}\bigg|_p\right)(y^j) v^i \frac{\partial}{\partial y^j}\bigg|_{F(p)} = \sum_{i=1}^n \sum_{j=1}^m \frac{\partial}{\partial x^i}\bigg|_p (y^j \circ F) v^i \frac{\partial}{\partial y^j}\bigg|_{F(p)}$$

If we define $F^j = y^j \circ F$ (The j -th coordinate of F) we get

$$DF_p(v) \equiv \sum_{i=1}^n \sum_{j=1}^m \frac{\partial F^j}{\partial x^i}\bigg|_p v^i \frac{\partial}{\partial y^j}\bigg|_{F(p)}$$

Where $\{\frac{\partial F^j}{\partial x^i}\big|_p : 1 \leq i \leq n, 1 \leq j \leq m\}$ is the Jacobian Matrix.

Definition B.8.1. (Pullback of a k -form) Let $\omega : V \rightarrow \Pi_{q \in V} \Lambda^k(T_q(V))$ be a k -form on $V \subset \mathbb{R}^m$. We can get a k -form on $U \subseteq \mathbb{R}^n$ via F as $F^*\omega_p \in \Lambda^k(T_p(U))$ given by

$$F^*\omega_p(v_1, v_2, \dots, v_k) = \omega_{f(p)}(DF_p(v_1), DF_p(v_2), \dots, DF_p(v_k))$$

B.8.1 Pullback of a 1-form

Let $\omega = dy^q$ be a basis 1-form in $V \subset \mathbb{R}^m$.

$$F^*(dy^q)_p(v) = dy_{F(p)}^q(DF_p(v))$$

Which gives

$$\begin{aligned} & dy_{F(p)}^q \left(\sum_{i=1}^n \sum_{j=1}^m \frac{\partial F^j}{\partial x^i}\bigg|_p v^i \frac{\partial}{\partial y^j}\bigg|_{F(p)} \right) \\ &= \sum_{i=1}^n \sum_{j=1}^m \frac{\partial F^j}{\partial x^i}\bigg|_p v^i \delta_{jq} = \sum_{i=1}^n \frac{\partial F^q}{\partial x^i}\bigg|_p v^i = \sum_{i=1}^n \frac{\partial F^q}{\partial x^i}\bigg|_p dx_p^i(v) \end{aligned}$$

Therefore we can set up the equivalence

$$F^*(dy^q)_p \equiv \sum_{i=1}^n \frac{\partial F^q}{\partial x^i}\bigg|_p dx_p^i \quad (\text{B.8.2})$$

Note that $F^*y^q = y^q \circ F \equiv F^q$, and its exterior derivative

$$dF^q = d(F^*(y^q)) = \left(\sum_{i=1}^n \frac{\partial}{\partial x^i}\bigg|_p dx^i \right) \wedge (F^*(y^q)) = \sum_{i=1}^n \frac{\partial y^q \circ F}{\partial x^i}\bigg|_p dx_p^i$$

And hence we see that for basis 1-forms

$$d(F^*(y^q)) = F^*(dy^q)$$

If $\omega : V \rightarrow \Pi_{q \in V} T_q^*(V)$ is any general one form given by

$$\omega \equiv \sum_{j=1}^m \omega^j dy^j$$

then we have

$$F^*(\omega)_p(v) := \omega_{F(p)}(DF_p(v)) = \sum_{j=1}^m \omega^j dy_{F(p)}^j(DF_p(v))$$

which yields

$$\sum_{j=1}^m \omega^j(F(p)) \sum_{i=1}^n \frac{\partial F^j}{\partial x^i}\bigg|_p dx_p^i(v) = \sum_{j=1}^m \omega_{(F(p))}^j dF_p^j(v)$$

which gives us an expansion of the form

$$F^*(\omega)_p \equiv \sum_{i=1}^n \sum_{j=1}^m \frac{\partial F^j}{\partial x^i} \big|_p \omega_{F(p)}^j dx_p^i \equiv \sum_{j=1}^m \omega_{(F(p))}^j dF_p^j$$

Yet another suggestive way to write this would be

$$F^*(\omega)_p = F^*\left(\sum_{j=1}^m \omega^j dy^j\right)_p = \sum_{j=1}^m (\omega^j \circ F)_{(p)} dF_p^j$$

B.8.2 Pullback of a k -form

Let $\omega = dy^I = dy^{i_1} \wedge dy^{i_2} \cdots \wedge dy^{i_k} \in \Omega(V, \Lambda^k)$ be a k form on $V \subset \mathbb{R}^m$.

$$\begin{aligned} F^*(dy^I)_p &= \left(\sum_{j_1=1}^n v_1^{j_1} \frac{\partial}{\partial x^{j_1}} \big|_p, \sum_{j_2=1}^n v_2^{j_2} \frac{\partial}{\partial x^{j_2}} \big|_p, \dots, \sum_{j_k=1}^n v_k^{j_k} \frac{\partial}{\partial x^{j_k}} \big|_p \right) \\ &= \sum_{1 \leq j_1, j_2, \dots, j_k \leq n} (v_1^{j_1} v_2^{j_2} \cdots v_k^{j_k}) F^*(dy^I)_p \left(\frac{\partial}{\partial x^{j_1}} \big|_p, \frac{\partial}{\partial x^{j_2}} \big|_p, \dots, \frac{\partial}{\partial x^{j_k}} \big|_p \right) \end{aligned} \quad (\text{B.8.3})$$

$$F^*(dy^I)_p \left(\frac{\partial}{\partial x^{j_1}} \big|_p, \frac{\partial}{\partial x^{j_2}} \big|_p, \dots, \frac{\partial}{\partial x^{j_k}} \big|_p \right) = dy_{F(p)}^I (DF_p \left(\frac{\partial}{\partial x^{j_1}} \big|_p \right), DF_p \left(\frac{\partial}{\partial x^{j_2}} \big|_p \right), \dots, DF_p \left(\frac{\partial}{\partial x^{j_k}} \big|_p \right))$$

From B.8.1 we see that

$$\begin{aligned} F^*(dy^I)_p \left(\frac{\partial}{\partial x^{j_1}} \big|_p, \frac{\partial}{\partial x^{j_2}} \big|_p, \dots, \frac{\partial}{\partial x^{j_k}} \big|_p \right) &= dy_{F(p)}^I \left(\sum_{q_1=1}^m \frac{\partial F^{q_1}}{\partial x^{j_1}} \big|_p \frac{\partial}{\partial y^{q_1}} \big|_{F(p)}, \sum_{q_2=1}^m \frac{\partial F^{q_2}}{\partial x^{j_2}} \big|_p \frac{\partial}{\partial y^{q_2}} \big|_{F(p)}, \dots, \sum_{q_k=1}^m \frac{\partial F^{q_k}}{\partial x^{j_k}} \big|_p \frac{\partial}{\partial y^{q_k}} \big|_{F(p)} \right) \\ &= \sum_{1 \leq q_1, q_2, \dots, q_k \leq m} \left(\frac{\partial F^{q_1}}{\partial x^{j_1}} \big|_p \cdot \frac{\partial F^{q_2}}{\partial x^{j_2}} \big|_p \cdots \frac{\partial F^{q_k}}{\partial x^{j_k}} \big|_p \right) dy_{F(p)}^I \left(\frac{\partial}{\partial y^{q_1}} \big|_{F(p)}, \frac{\partial}{\partial y^{q_2}} \big|_{F(p)}, \dots, \frac{\partial}{\partial y^{q_k}} \big|_{F(p)} \right) \\ &= \sum_{1 \leq q_1, q_2, \dots, q_k \leq m} \left(\frac{\partial F^{q_1}}{\partial x^{j_1}} \big|_p \cdot \frac{\partial F^{q_2}}{\partial x^{j_2}} \big|_p \cdots \frac{\partial F^{q_k}}{\partial x^{j_k}} \big|_p \right) \det \left((\delta_{i_s}^{q_t}) : 1 \leq s, t \leq k \right) \end{aligned}$$

Combining this with B.8.3 gives us

$$\begin{aligned} F^*(dy^I)(v_1, \dots, v_k) &= \sum_{\substack{1 \leq j_1, j_2, \dots, j_k \leq n \\ 1 \leq q_1, q_2, \dots, q_k \leq m}} (v_1^{j_1} v_2^{j_2} \cdots v_k^{j_k}) \left(\frac{\partial F^{q_1}}{\partial x^{j_1}} \big|_p \cdot \frac{\partial F^{q_2}}{\partial x^{j_2}} \big|_p \cdots \frac{\partial F^{q_k}}{\partial x^{j_k}} \big|_p \right) \det \left((\delta_{i_s}^{q_t})_{1 \leq s, t \leq k} \right) \quad (\text{B.8.4}) \\ &= \sum_{1 \leq j_1, j_2, \dots, j_k \leq n} (v_1^{j_1} v_2^{j_2} \cdots v_k^{j_k}) \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) \left(\frac{\partial F^{i_{\sigma(1)}}}{\partial x^{j_1}} \big|_p \cdot \frac{\partial F^{i_{\sigma(2)}}}{\partial x^{j_2}} \big|_p \cdots \frac{\partial F^{i_{\sigma(k)}}}{\partial x^{j_k}} \big|_p \right) \\ &= \sum_{1 \leq j_1, \dots, j_k \leq n} \left(dx^{j_1}(v_1) dx^{j_2}(v_2) \cdots dx^{j_k}(v_k) \right) \det \left(\left(\frac{\partial F^{i_r}}{\partial x^{j_s}} \big|_p \right)_{1 \leq r, s \leq k} \right) \\ &= \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \sum_{\sigma \in \text{sym}(k)} \left(dx^{j_{\sigma(1)}}(v_1) dx^{j_{\sigma(2)}}(v_2) \cdots dx^{j_{\sigma(k)}}(v_k) \right) \det \left(\left(\frac{\partial F^{i_r}}{\partial x^{j_{\sigma(s)}}} \big|_p \right)_{1 \leq r, s \leq k} \right) \\ &= \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \det \left(\left(\frac{\partial F^{i_r}}{\partial x^{j_s}} \big|_p \right)_{1 \leq r, s \leq k} \right) \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) dx^{j_{\sigma(1)}}(v_1) dx^{j_{\sigma(2)}}(v_2) \cdots dx^{j_{\sigma(k)}}(v_k) \end{aligned}$$

Which finally yields

$$F^*(dy^I)_p = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \det \left(\left(\frac{\partial F^{i_r}}{\partial x^{j_s}} \big|_p \right)_{1 \leq r, s \leq k} \right) dx_p^{j_1} \wedge dx_p^{j_2} \cdots dx_p^{j_k} \quad (\text{B.8.5})$$

We now look at $d(F^*y^{i_1})_p \wedge d(F^*y^{i_2})_p \cdots \wedge d(F^*y^{i_k})_p = dF^{i_1} \wedge dF^{i_2} \cdots dF^{i_k}$.

$$= \left(\sum_{j_1=1}^n \frac{\partial F^{i_1}}{\partial x^{j_1}} \big|_p dx^{j_1} \right) \wedge \left(\sum_{j_2=1}^n \frac{\partial F^{i_2}}{\partial x^{j_2}} \big|_p dx^{j_2} \right) \wedge \cdots \wedge \left(\sum_{j_k=1}^n \frac{\partial F^{i_k}}{\partial x^{j_k}} \big|_p dx^{j_k} \right)$$

$$\begin{aligned}
&= \sum_{1 \leq j_1, j_2, \dots, j_k \leq n} \left(\frac{\partial F^{i_1}}{\partial x^{j_1}} \Big|_p \cdot \frac{\partial F^{i_2}}{\partial x^{j_2}} \Big|_p \cdots \frac{\partial F^{i_k}}{\partial x^{j_k}} \Big|_p \right) dx^{j_1} \wedge \cdots \wedge dx^{j_k} \\
&= \sum_{1 \leq j_1 < j_2 < \cdots < j_k \leq n} dx^{j_1} \wedge dx^{j_2} \cdots \wedge dx^{j_k} \sum_{\sigma \in \text{sym}(k)} \text{sign}(\sigma) \left(\frac{\partial F^{i_1}}{\partial x^{j_{\sigma(1)}}} \Big|_p \cdot \frac{\partial F^{i_2}}{\partial x^{j_{\sigma(2)}}} \Big|_p \cdots \frac{\partial F^{i_k}}{\partial x^{j_{\sigma(k)}}} \Big|_p \right) \\
&= \sum_{1 \leq j_1 < j_2 < \cdots < j_k \leq n} dx^{j_1} \wedge \cdots \wedge dx^{j_k} \det \left(\left(\frac{\partial F^{i_r}}{\partial x^{j_s}} \Big|_p \right)_{1 \leq r, s \leq k} \right)
\end{aligned} \tag{B.8.6}$$

Which agrees with B.8.5. Hence we have the following identity:

$$F^*(dy^{i_1} \wedge dy^{i_2} \wedge \cdots \wedge dy^{i_k}) = d(F^*y^{i_1}) \wedge d(F^*y^{i_2}) \wedge \cdots \wedge d(F^*y^{i_k}) \tag{B.8.7}$$

Corollary B.8.2. If $F : U_{\mathbb{C}\mathbb{R}^n} \rightarrow V_{\mathbb{C}\mathbb{R}^n}$ is a smooth Diffeomorphism, and coordinates for U are given by $x^1, x^2, \dots, x^n$, and those for V given by $y^1, y^2, \dots, y^n$, we have a change of variables formula for k -forms (In view of formula B.8.5) given by:

$$F^*(dy^1 \wedge dy^2 \wedge \cdots \wedge dy^n)_p = \det \left(\left(\frac{\partial F^j}{\partial x^i} \right)_{1 \leq i, j \leq n} \right) dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n$$

Remark B.8.3. The proof of B.8.2 is simply an application of B.8.5. This is the **change of variables formula** for *top forms*.

Theorem B.8.4. (Pullback of Wedge) Let $\alpha \in \Omega(V, \Lambda^k)$ be a k -form on $V \subset \mathbb{R}^m$ and let $\beta \in \Omega(V, \Lambda^l)$ be an l -form on V . For smooth function $F : U \subset \mathbb{R}^n \rightarrow V$ we have

$$F^*(\alpha \wedge \beta) = F^*(\alpha) \wedge F^*(\beta)$$

Proof. Recall $F^*(\alpha \wedge \beta)_p(v_1, v_2, \dots, v_k, v_{k+1}, \dots, v_{k+l}) = (\alpha \wedge \beta)_{F(p)}(DF_p(v_1), \dots, DF_p(v_{k+l})) =$

$$\begin{aligned}
&\sum_{\sigma \in \text{sym}(k, l)} \text{sign}(\sigma) \alpha_{F(p)}(DF_p(v_{\sigma(1)}), \dots, DF_p(v_{\sigma(k)})) \beta_{F(p)}(DF_p(v_{\sigma(k+1)}), \dots, DF_p(v_{\sigma(k+l)})) \\
&= \sum_{\sigma \in \text{sym}(k, l)} \text{sign}(\sigma) F^*(\alpha)_p(v_{\sigma(1)}, \dots, v_{\sigma(k)}) F^*(\beta)_p(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)}) = F^*(\alpha) \wedge F^*(\beta)
\end{aligned}$$

■

If $\omega = \sum_{I \in \mathcal{T}_k} \omega^I dy^I$ is a general k -form on $V \subseteq \mathbb{R}^m$, then we have

$$F^*\left(\sum_I \omega^I dy^I\right)_p = \sum_I \omega^I_{F(p)} dF_p^I \tag{B.8.8}$$

Where $dF_p^I = dF_p^{i_1} \wedge dF_p^{i_2} \wedge \cdots \wedge dF_p^{i_k}$.

Lemma B.8.5. (F^* commutes with d) Let $F : U_{\mathbb{C}\mathbb{R}^n} \rightarrow V_{\mathbb{C}\mathbb{R}^m}$ be smooth, and ω a k -form on V . We then have that

$$F^*(d\omega) = d(F^*\omega)$$

Proof. If $\omega = \sum_{I \in \mathcal{T}_k} \omega^I dy^I$ From B.8.8 we have that

$$F^*(\omega)_p = \sum_I \omega^I_{F(p)} dF_p^{i_1} \wedge dF_p^{i_2} \cdots \wedge dF_p^{i_k}$$

When we apply the exterior derivative to this expression we have (and recalling the antiderivation property)

$$d(F^*(\omega))_p = \sum_I d(\omega^I \circ F)_p \wedge dF_p^{i_1} \wedge dF_p^{i_2} \cdots \wedge dF_p^{i_k} + (\omega^I \circ F) d(dF_p^{i_1} \wedge dF_p^{i_2} \cdots \wedge dF_p^{i_k})_p$$

The second term vanishes because $d^2 = 0$. This gives

$$d(F^*(\omega))_p = \sum_I \sum_{i=1}^n \frac{\partial(\omega^I \circ F)}{\partial x^i} \Big|_p dx_p^i \wedge dF_p^{i_1} \wedge dF_p^{i_2} \wedge \cdots \wedge dF_p^{i_k} = \sum_I d(\omega^I \circ F) \wedge dF_p^I$$

Now, if $\omega = \sum_I \omega^I dy^I$, we have $d\omega = \sum_I d\omega^I \wedge dy^I$. Pulling this back gives us

$$F^*(d\omega) = \sum_I F^*(d\omega^I) \wedge F^*(dy^I) = \sum_I d(\omega^I \circ F) \wedge dF^I$$

Which establishes equality. ■

Pullback and Interior Product

Remark B.8.6. For a smooth map $F: M \rightarrow N$, the identity $i_X(F^*\omega) = F^*(i_Y\omega)$ is ill-defined for generic vector fields since X is a vector field from M and Y is a vector field on N . Naturality under pullbacks holds precisely when X and Y are F -related, which motivates the next definition.

Definition B.8.7. (F -related Vector Fields) Let $F: M \rightarrow N$ be smooth. Vector fields X on M is said to be **F-related** to a vector field Y on N if

$$dF_p(X_p) = Y_{F(p)}$$

for every $p \in M$.

Theorem B.8.8. Let $F: M \rightarrow N$ be a smooth map. If X is a vector field on M and Y on N are F -related, then for any differential form $\omega \in \Omega^k(N)$,

$$i_X(F^*\omega) = F^*(i_Y\omega).$$

Proof. Evaluating the left-hand side at $p \in M$ on vectors $v_1, \dots, v_{k-1} \in T_p M$ yields:

$$\begin{aligned} (i_X F^*\omega)_p(v_1, \dots, v_{k-1}) &= (F^*\omega)_p(X_p, v_1, \dots, v_{k-1}) \\ &= \omega_{F(p)}(dF_p X_p, dF_p v_1, \dots, dF_p v_{k-1}) \\ &= \omega_{F(p)}(Y_{F(p)}, dF_p v_1, \dots, dF_p v_{k-1}) \\ &= (F^* i_Y \omega)_p(v_1, \dots, v_{k-1}). \end{aligned} \quad \blacksquare$$

Lemma B.8.9. Let $U \subseteq \mathbb{R}^n$ be an open connected set and $f: U \rightarrow \mathbb{R}$ be a smooth function. Then $df = 0$ if and only if f is constant on U .

Proof. $\implies$) Let f be constant on U . Then $\frac{\partial f}{\partial x^i} = 0$ for all $i \leq n$, and hence $df_p = \sum_{i=1}^n \frac{\partial f}{\partial x^i} \Big|_p dx_p^i = 0$.

$\impliedby$) Say $df_p = 0$ for all $p \in U$. Let x and y be arbitrary fixed points in U and let $c: [a, b] \rightarrow U$ be any smooth curve connecting x and y (Such a curve exists because U is path-connected, and hence continuity is guaranteed. Smoothing of the polygon edges with smooth curves guarantees smoothness). Note that $c^*f \equiv f \circ c: [a, b] \rightarrow \mathbb{R}$ is a smooth 0 form on $[a, b]$. $d(c^*f) = d(f \circ c) = \frac{d(f \circ c(t))}{dt} dt = c^*df = 0$ (from B.8.5). Therefore $d(c^*f) = 0$, so $f \circ c(a) = f \circ c(b)$ which gives $f(x) = f(y)$. ■

B.9 Integration of 1-forms

Let $U \subseteq \mathbb{R}^n$ be an open set and α be a 1-form.

Definition B.9.1. (Parametrized Paths) Any map $c : [a, b] \rightarrow U \subset \mathbb{R}^n$ that is smooth is said to be a *path* from $c(a)$ to $c(b)$.

The pullback of α given by $c^*\alpha$ is a 1-form on $[a, b]$. Suppose $c^*\alpha = hdt$ where t is the coordinate function of $[a, b]$ and h a real valued smooth function.

Definition B.9.2. (Integral of a 1-form over a curve)

$$\int_c \alpha = \int_{[a,b]} c^*\alpha = \int_a^b h(t)dt \quad (\text{B.9.1})$$

where the right most equality makes sense in the standard way. Moreover, if $\alpha = \sum_{i=1}^n f^i dx^i$, we have $c^*(\sum_{i=1}^n f^i dx^i) = \sum_{i=1}^n c^*(f^i) d(c^*x^i) = \sum_{i=1}^n (f^i \circ c) \frac{dc^i}{dt} dt$ where $c^i = x^i \circ c$. Its integral would then be

$$\int_c \alpha = \int_a^b c^*\alpha = \sum_{i=1}^n \int_a^b f^i \circ c(t) \frac{dc^i}{dt}(t) dt = \int_a^b \sum_{i=1}^n f^i(c(t)) (c^i)'(t) dt \quad (\text{B.9.2})$$

Which can be suggestively written as

$$\int_a^b \alpha^\sharp(c(t)) \cdot c'(t) dt \quad (\text{B.9.3})$$

where $\alpha^\sharp$ is the vector field associated with the 1-form α via the inner product.

B.10 The Stokes Theorem

Definition B.10.1. (Lipschitz and C^k Set) Let $\Omega \subseteq \mathbb{R}^n$ be an open domain and let $\partial\Omega$ denote its boundary. Ω is said to be a *Lipschitz set* (C^k set) if for every point $p \in \partial\Omega$ there exists a hyperplane H of dimension $n-1$ passing through p , a Lipschitz continuous (C^k) function $g : H \rightarrow \mathbb{R}$ and reals $r > 0, h > 0$ such that:

1. $\Omega \cap C = \{x + y\hat{n} : x \in B_r(p) \cap H \text{ and } -h < y < g(x)\}$
2. $\partial\Omega \cap H = \{x + y\hat{n} : x \in B_r(p) \cap H \text{ and } y = g(x)\}$

Where

1. $\hat{n}$ is the unit normal to H .
2. $B_r(p)$ is the ball of radius r at p .
3. $C := \{x + y\hat{n} : x \in B_r(p) \cap H \text{ and } -h < y < h\}$ (Cylinder of height $2h$ and radius r at p)




Figure B.1: Cylinder around a boundary point $p \in \partial\Omega$ from definition B.10.1

B.11 The Codifferential δ

B.11.1 Motivation

Let $\alpha \in \Omega\Lambda^{k-1}$ and $\beta \in \Omega\Lambda^k$. Here we can invoke the definition of the Hodge star (B.6.1) to get

$$\langle d\alpha, \beta \rangle \text{vol} = d\alpha \wedge (\star\beta)$$

$$\int_{\Omega} \langle d\alpha, \beta \rangle \text{vol} = \int_{\Omega} \left(d\alpha \wedge (\star\beta) + (-1)^{k-1} \alpha \wedge d(\star\beta) + (-1)^k \alpha \wedge d(\star\beta) \right)$$

Using the Antiderivation identity B.3.12 we get

$$\int_{\Omega} \langle d\alpha, \beta \rangle \text{vol} = \int_{\Omega} \left(d(\alpha \wedge \star\beta) + (-1)^k \alpha \wedge d(\star\beta) \right) \quad (\text{B.11.1})$$

$$\int_{\Omega} \langle d\alpha, \beta \rangle \text{vol} = \left(\int_{\partial\Omega} \alpha \wedge (\star\beta) + (-1)^k \int_{\Omega} \alpha \wedge d(\star\beta) \right)$$

Since $\star\beta$ is a $n-k$ form and $d(\star\beta)$ is an $n-(k-1)$ form, we apply the definition of Hodge star again to replace $\alpha \wedge d(\star\beta)$ with $(-1)^{(n-(k-1))(k-1)} \langle \alpha, \star d(\star\beta) \rangle \text{vol}$ to get:

$$\int_{\Omega} \langle d\alpha, \beta \rangle \text{vol} = \left(\int_{\partial\Omega} \alpha \wedge (\star\beta) + (-1)^k (-1)^{(n-(k-1))(k-1)} \int_{\Omega} \langle \alpha, \star d(\star\beta) \rangle \text{vol} \right) \quad (\text{B.11.2})$$

Note that (B.11.2) is a formula that is independent of the definition for *Codifferential* that will follow. This is important because different texts (cf. [Jos17] and [CDK12]) opt for different sign conventions for the Codifferential.

Definition B.11.1. (Codifferential) We define a map $\delta : \oplus_{k=0}^{\infty} \Omega\Lambda^k \rightarrow \oplus_{k=0}^{\infty} \Omega\Lambda^k$ given by

$$\delta(\omega) := (-1)^{n(k-1)} \star d \star (\omega) \quad (\text{B.11.3})$$

for a k -form ω .

Using B.11.1 in (B.11.2) we immediately see that

$$\int_{\Omega} \langle d\alpha, \beta \rangle \text{vol} = (-1) \int_{\Omega} \langle \alpha, \delta\beta \rangle \text{vol} + \int_{\partial\Omega} \alpha \wedge \star\beta \quad (\text{B.11.4})$$

This is called the *integration by parts formula* for forms.

Remark B.11.2. 1. Under the conventions of B.11.1, the formula B.11.4 takes the above form. But if we propose that $\delta = (-1)^{nk+n+1} \star d\star$, or $(-1)^{k(n-k+1)} \star d\star$, we see different possible factors in B.11.4.

2. If we assume that our set either has no boundary or that the forms vanish on the boundary, the formula B.11.4 yields

$$\int_{\Omega} \langle d\alpha, \beta \rangle + \int_{\Omega} \langle \alpha, \delta\beta \rangle = 0$$

B.11.2 Coordinate formulas for the Codifferential

Case 1: Let $\omega = dx^I$. Where $I \in \mathcal{T}_k(\mathbb{R}^n)$. Then $\delta(\omega) = (-1)^{n(k-1)} \star d \star (dx^I)$ gives us

$$(-1)^{n(k-1)} \text{sgn}(R(I, I^c)) \star (d(dx^{I^c})) = 0$$

Case 2: Consider the case where $\omega = f dx^I$. For ease sake, let $(-1)^{n(k-1)} := \kappa$. We then have $\delta(\omega) = \kappa \star d \star (f dx^I) = \kappa \star d(f \text{sgn}(R(I, I^c)) dx^{I^c})$ which gives

$$\kappa \text{sgn}(R(I, I^c)) \star d(f dx^{I^c}) = \kappa \text{sgn}(R(I, I^c)) \star \left(\sum_{q=1}^n \frac{df}{dx^q} dx^q \wedge dx^{I^c} \right)$$

since the summand is 0 for when $q \notin I$ we have

$$\delta(f dx^I) = \kappa \text{sgn}(R(I, I^c)) \left(\sum_{\gamma=1}^k \frac{df}{dx^{i_{\gamma}}} \star (dx^{i_{\gamma}} \wedge dx^{I^c}) \right) \quad (\text{B.11.5})$$

We will continue the derivation after the following proposition:

Proposition B.11.3. Let $I = (i_1, i_2, \dots, i_k) \in \mathcal{T}_k(\mathbb{R}^n)$. Then

$$\star(dx^{i_\gamma} \wedge dx^{I^c}) = (-1)^{n-k+\gamma-1+k(n-k)} \text{sgn}(R(I, I^c)) dx^{I_{\hat{i}_\gamma}} \quad (\text{B.11.6})$$

Proof. $\star(dx^{i_\gamma} \wedge dx^{I^c}) = dx^{I_{\hat{i}_\gamma}}(R)$ where R is the coefficient such that $dx^{i_\gamma} \wedge dx^{I^c} \wedge dx^{I_{\hat{i}_\gamma}} = R \text{vol}$. Note that

$$dx^{i_\gamma} \wedge dx^{I^c} \wedge dx^{I_{\hat{i}_\gamma}} = (-1)^{1(n-k)} dx^{I^c} \wedge (-1)^{\gamma-1} dx^I$$

Which gives

$$(-1)^{n-k} (-1)^{\gamma-1} \text{sgn}(I, I^c) (-1)^{k(n-k)} \text{vol}$$

And hence $R = (-1)^{n-k+\gamma-1+k(n-k)} \text{sgn}(R(I, I^c))$

■

From the proposition, we have

$$\begin{aligned} \delta(f dx^I) &= \kappa \text{sgn}(R(I, I^c))^2 (-1)^{n-k+k(n-k)} \left(\sum_{\gamma=1}^k (-1)^{\gamma-1} \frac{\partial f}{\partial x^{i_\gamma}} dx^{I_{\hat{i}_\gamma}} \right) \\ \delta(f dx^I) &= \left(\sum_{\gamma=1}^k (-1)^{\gamma-1} \frac{\partial f}{\partial x^{i_\gamma}} dx^{I_{\hat{i}_\gamma}} \right) \end{aligned} \quad (\text{B.11.7})$$

Remark B.11.4. Note that, if we adopted a different convention for δ (like $\delta = (-1)^{k(n-k+1)}$) we would then have the following formula instead of (B.11.7):

$$\delta(f dx^I) = (-1)^n \left(\sum_{\gamma=1}^k (-1)^{\gamma-1} \frac{\partial f}{\partial x^{i_\gamma}} dx^{I_{\hat{i}_\gamma}} \right) \quad (\text{B.11.8})$$

General Case: Let $\omega = \sum_{I \in \mathcal{T}_k(\mathbb{R}^n)} f^I dx^I$ Then

$$\delta\left(\sum_{I \in \mathcal{T}_k} f^I dx^I\right) = \sum_{I \in \mathcal{T}_k} \delta(f^I dx^I) = \sum_{I \in \mathcal{T}_k} \sum_{\gamma=1}^k (-1)^{\gamma-1} \frac{\partial f^I}{\partial x^{i_\gamma}} dx^{I_{\hat{i}_\gamma}} \quad (\text{B.11.9})$$

Alternative Formula for the Codifferential:

Let $\omega = \sum_{1 \leq i_1 < \dots < i_k \leq n} \omega^{i_1 i_2 \dots i_k} dx^{i_1} \wedge dx^{i_2} \wedge \dots \wedge dx^{i_k}$ and let $\delta(\omega) = \kappa \star d \star (\omega)$.

$$\begin{aligned} \delta(\omega) &= \kappa \star d \left(\sum_{1 \leq i_1 < \dots < i_k \leq n} \omega^{i_1 \dots i_k} \text{sgn}(R(I, I^c)) dx^{I^c} \right) \\ &= \kappa \star \left(\sum_{1 \leq i_1 < \dots < i_k \leq n} \text{sgn}(R(I, I^c)) d\omega^{i_1 \dots i_k} \wedge dx^{I^c} \right) = \kappa \star \left(\sum_{1 \leq i_1 < \dots < i_k \leq n} \text{sgn}(R(I, I^c)) \left(\sum_{q=1}^n \frac{\partial \omega^{i_1 \dots i_k}}{\partial x^q} dx^q \right) \wedge dx^{I^c} \right) \\ \delta(\omega) &= \kappa \left(\sum_{q=1}^n \sum_{I \in \mathcal{T}_{k-1}} \text{sgn}(R([qI], [qI]^c)) \frac{\partial \omega^{[qI]}}{\partial x^q} \star dx^q \wedge dx^{[qI]^c} \right) \end{aligned} \quad (\text{B.11.10})$$

A modification to the proof of B.11.6 yields us the following computation for $\star(dx^q \wedge dx^{[qI]^c})$:

$$\star(dx^q \wedge dx^{[qI]^c}) = (-1)^{n-k+\gamma-1+k(n-k)} \text{sgn}(R([qI], [qI]^c)) dx^I \quad (\text{B.11.11})$$

where $i_{\gamma-1} < q < i_\gamma$. Combining (B.11.11) and (B.11.10) we get:

$$\delta(\omega) = \kappa \left(\sum_{q=1}^n \sum_{I \in \mathcal{T}_{k-1}} \text{sgn}(R([qI], [qI]^c))^2 (-1)^{n-k+\gamma-1+k(n-k)} \frac{\partial \omega^{[qI]}}{\partial x^q} dx^I \right)$$

which gives

$$\delta(\omega) = \kappa (-1)^{n-k+k(n-k)} \left(\sum_{q=1}^n \sum_{I \in \mathcal{T}_{k-1}} \frac{\partial \omega^{q,I}}{\partial x^q} dx^I \right)$$

Which simplifies to

$$\delta(\omega) = \left(\sum_{q=1}^n \sum_{I \in \mathcal{T}_{k-1}} \frac{\partial \omega^{q,I}}{\partial x^q} dx^I \right) \quad (\text{B.11.12})$$

Where the notation $\omega^{q,I}$ is understood from the equation B.5.6.

Remark B.11.5. This is the formula presented in [CDS18] pg. 465, 2 (vii).

B.12 Boundary Calculus and Tangential and Normal Components

!!Not complete!!

1. tangential and normal part proofs
2. admissible boundary coordinates proof
3. subtleties with lipshitz domains and boundaries.

B.13 Integration Theorems

!!Not complete!!

Ideal case scenario– develop the theory of integration of forms– Do Carmo and Frenkel

1. use pullbacks
2. partitions of unity
3. proof of stokes (long shot)
4. corollaries (gauss and green)

B.14 A little bit of Riemannian Geometry

: – Not Complete–

1. To ideally develop metric dependent hodge star and interior product.
2. Define principle curvature, mean curvature, gauss curvature, geodesics etc. 1st and 2nd Fundamental forms etc.

Appendix C

Sobolev Spaces and Requisite Measure Theory and Functional Analysis

Appendix D

Manifolds

D.1 Topological Manifolds

Definition D.1.1. (Topological Manifold of dimension n) A Hausdorff, second countable topological space M is said to be a *Topological Manifold* of dimension n if for every $p \in M$, there exists U open in M with $p \in U$, $\hat{U} \subset \mathbb{R}^n$ and a homeomorphism $\phi : U \rightarrow \hat{U}$. Such a pair (U, ϕ) is called a *coordinate chart*, and a collection of such (U_α, ϕ_α) that covers M is called an *atlas*.

Example D.1.2. (Graph of a continuous function) Let $U \subset \mathbb{R}^n$ be open and $f : U \rightarrow \mathbb{R}^k$ be a continuous mapping. Let $\Gamma \subset \mathbb{R}^n \times \mathbb{R}^k$ be given by

$$\Gamma := \{(x, f(x)) \in \mathbb{R}^k \times \mathbb{R}^n : x \in U\}$$

Let Γ have the subspace topology and let $p \in \Gamma$ be some $(x_p, f(x_p))$. Define $\phi : \Gamma \rightarrow U$ by

$$\phi(x, y) = x$$

This map is continuous, and has a continuous inverse $g(x) = (x, f(x))$. Hence ϕ is a homeomorphism and $\{(\Gamma, \phi)\}$ gives us an atlas.

Bibliography

- [CDK12] Gyula Csató, Bernard Dacorogna, and Olivier Kneuss. *The Pullback Equation for Differential Forms*. Vol. 83. Progress in Nonlinear Differential Equations and Their Applications. New York: Birkhäuser, 2012. ISBN: 978-0-8176-8312-2. DOI: [10.1007/978-0-8176-8313-9](https://doi.org/10.1007/978-0-8176-8313-9).
- [CDS18] G. Csato, B. Dacorogna, and S. Sil. “On the best constant in Gaffney inequality”. In: *Journal of Functional Analysis* 274.2 (Jan. 2018), pp. 461–503. ISSN: 0022-1236. DOI: [10.1016/j.jfa.2017.10.006](https://doi.org/10.1016/j.jfa.2017.10.006). URL: <http://dx.doi.org/10.1016/j.jfa.2017.10.006>.
- [Jos17] Jürgen Jost. *Riemannian Geometry and Geometric Analysis*. 7th ed. Universitext. Cham: Springer, 2017. ISBN: 978-3-319-61859-3.