



The real before-and-after: where the original LOD died

HT115 vs pos-1. Above $|z| = 38.5$ the shipped p column is exactly 0 and $PvalToLOD(0)$ is Inf, so the original LOD is infinite for every marker whose log-space LOD exceeds about 322. The N2 x XZ1516 contrast has 4,765 of them, running to LOD 710; the JU cross has none, peaking at $|z| = 25.4$. A shipped LOD above 322 is itself proof the log-space recomputation was applied.



The conventions differ by $\log_{10}(2)$ ASYMPTOTICALLY, and converge slowly

The gap is 0.2859 at the genome-wide threshold of LOD 3.57, not 0.30103. It reaches 0.290 at LOD 5.2, 0.295 at LOD 10.1 and 0.300 only at LOD 62.7. So for a marker near the threshold the two conventions are 0.286 apart, and treating the offset as a flat 0.301 misstates it by 0.015 LOD.



Where the offset stops being constant, and why cross does not matter here

Below $|z| = 0.6745$ the doubled argument to $qchisq$ exceeds 1, $PvalToLOD()$ returns 0, and the package LOD is pinned there while the chisq LOD is not. Nothing significant lives here. The two crosses lie on ONE curve because the gap is a deterministic function of $|z|$ — colour separates them only in panel A, where the LOD ranges differ.

