



A IV:15.32 Mb admitted

The scan's strongest marker, on the right arm.
-log₁₀ *p* 8.84, over Bonferroni, with 95 eigen-passing neighbours within 100 kb.
Effect allele A, frequency 0.208. Wilcoxon *p* = 1e-07.
Observed gap +0.0514; the model's 2β is +0.0515.




B X:4.88 Mb admitted

The only chromosome X marker over Bonferroni, 28% along the chromosome.
-log₁₀ *p* 7.83, over Bonferroni, with 2 eigen-passing neighbours within 100 kb.
Effect allele G, frequency 0.087. Wilcoxon *p* = 6e-06.
Observed gap +0.0738; the model's 2β is +0.0726.




C III:5.97 Mb discarded

The centre of the chromosome.
-log₁₀ *p* 8.68, over Bonferroni, with 0 eigen-passing neighbours within 100 kb.
Effect allele T, frequency 0.091. Wilcoxon *p* = 0.011.
Observed gap +0.0456; the model's 2β is +0.1216.



D III:12.72 Mb admitted

The right arm, 375 kb proximal to *sid-2*.
-log₁₀ *p* 6.31, under Bonferroni but over the eigen threshold, with 14 eigen-passing neighbours within 100 kb.
Effect allele T, frequency 0.087. Wilcoxon *p* = 2e-04.
Observed gap +0.0637; the model's 2β is +0.0616.



E LD between the four markers

Squared correlation of the dosages, positions in Mb, same 231 strains; agrees exactly with plink's own estimator. Note the scale: the strongest pair is 0.112 and the interval work calls markers linked at r-squared 0.7, so no pair here is close to being one signal. The two chromosome III markers, 6.75 Mb apart, are the least correlated pair of the six.

