

Simple random sampling procedure is used to select three sites from the two coastal belts in India for collection of fish and shellfish. The selected sites are Vishakhapatnam, Chennai and Mandapam from east coast belt as well as Mumbai, Goa and Kochi from west coast belt. All varieties of marine fish and shellfish available at the time of sampling at all the six landing sites are collected. All samples are cleaned, scaled/skinned, eviscerated and deboned at the collection site and only edible portion is composited. Two identical composited samples are collected and transported to NIN by air in frozen condition under dry ice, for analysis.

Upon arrival at the laboratory, the samples are checked for sample integrity. Fresh vegetables are expected to arrive at the laboratory cold and fresh in acceptable conditions, while frozen meat or fish samples are to be without any apparent sign of thawing. Any damaged sample was immediately discarded and logged onto the list for collection in the next sampling cycle. Descriptions of all sample handling, processing, and distribution of composite samples for analysis is maintained. Each sample is assigned an alpha numeric code upon arrival at the laboratory and all subsequent analysis and documentation is linked to that number. All samples are stored at appropriate location or at ambient temperature in the laboratory based on the sample perishability.

2.7 Analytical parameters

Interest in food components is expanding rapidly and the need to access core data on more nutrients and bioactive substances in foods is far out-stripping the amount of data that is available. Special interest database on emerging bioactive components in foods are necessary to meet the increasing demands of health professionals, dieticians, policy makers, epidemiologist studying the relationships between diet and health, as well as consumer interest in what is in the food they eat. In view of the demand for newer components in foods, it was decided to undertake a comprehensive analysis of nutrients and bioactive components in all key foods.

Official methods of analysis of Association of Official Analytical Chemist (AOAC) or if no AOAC method was available, other appropriate and reliable analytical methods that have undergone collaborative evaluations were tested and adopted. Each method was standardized in the laboratory and validated as per the IUPAC/AOAC method validation procedures with respect to precision, accuracy, simplicity, cost-effective, sensitivity and selectivity. About 200 discrete food constituents including sub-components were standardized for each food in the new food composition database representing proximate principles (8), energy (1), water-soluble vitamins (13), fat soluble vitamins (17), carotenoids (11), individual free sugars (5), oligosaccharides (4), minerals (20), organic acids (8), fats and fatty acids (40), amino acid composition (20), sterols (4), polyphenols (39), and bioactive components

(5) (**Table 5**). All the methods adopted here have been fully scrutinized for their robustness in analysing the compound of interest in different food matrices of different foods types.

2.8 Quality assurance and quality control (QAQC) scheme

Analytical Quality Assurance (QA) plan was designed to monitor the performance of the measurement systems to maintain statistical control and provide rapid feedback so that corrective measures could be taken before data quality was compromised. The programme ensures that reported data were sufficiently complete, comparable, representative, unbiased and precise so as to be suitable for their intended use. Certified Reference Material (CRMs) in which one or more properties are well quantified is used for calibration of analytical methods and to ensure data quality. The CRMs are procured from the European Commission Community Bureau of Reference (BCR), Belgium and the National Institute of Standards and Technology (NIST), Gaithersburg, USA.

As far as possible, certified reference materials (CRM) are used in this study, however, due to the prohibitive cost of CRMs, market rice sample is used as in-house quality control material with every batch of analysis. For this, a relatively large amount of the rice was homogenized and kept under controlled conditions. While market rice does not have a certified value, it develops an expected value through repeated analysis. The rationale for the development of control composites is driven by the need for precision and accuracy of data generated for the food samples. Control composites and CRM are included according to the level of validation needed for specific foods and nutrients. Regular participation in external proficiency testing programmes assured quality of the data generated in the laboratory.

3. Conclusions

The current national sampling plan addresses the key foods identified in the Indian food supply but does not include food specific to the sub-population of the country. The Indian national food composition database programme was initiated in India in 2011 with the objective of generating and improving the quantity and quality of food composition data in the country. Adequate infrastructure was developed at the National Institute of Nutrition, Hyderabad to meet the analytical requirements for all the components in foods. The identification of key foods, statistically derived sampling plan and analytical parameters described above provide what may be called an ideal plan. This ideal plan involves sampling all key foods identified as important in the Indian dietary supply from geographically dispersed areas across India covering all agro-ecological zones of the country. The ideal plan also includes the analysis of more than 150 analytical

Table 5. Analytical methods of nutrient and bioactive components.

Analyte	Method	Reference
Moisture	Gravimetric using vacuum /Hot air oven	AOAC 934.01
Protein	Titrimetry by combustion/Kjeldahl/Mixed catalyst	AOAC 2001.11; Jones (1941)
Ash	Gravimetric using Muffle furnace	AOAC 942.05
Total fat	Gravimetry; Ether extraction using Soxhlet apparatus	AOAC 963.15
Total, soluble and insoluble dietary fiber	Enzymatic-gravimetric	AOAC 991.43
Available carbohydrate	Calculation by difference	Greenfield & Southgate (2003)
	Trienzyme extraction and spectrophotometry	AOAC 991.43; AOAC 985.29
Energy	Atwater conversion factors	Merrill & Watt (1973); FAO / INFOODS Guidelines (2012)
Thiamin	Fluorometry	AOAC 942.23; Ramasastri (1975)
Riboflavin	Turbidometric-Microbiological assay	AOAC 940.33; AOAC 960.46
Niacin	Turbidometric-Microbiological assay	AOAC 985.34; 960.46
Pantothenic acid	RP-HPLC with UV detector	David & Harvey (2000)
Vitamin B6	HPLC-fluorometric detection	AOAC 985.32; Ollilainen (2008)
Biotin	U-HPLC and LC-ESI-IT-MS method	Holler et al. (2006)
Vitamin B9	Trienzyme Extraction and U-HPLC - UV or fluorescence detection	Finglas et al. (1999); Vahteristo et al. (1996); AOAC 2011.06; AOAC 2013.13
Vitamin C	RP-HPLC	AOAC 985.33; Keating & Haddad (1982)
Vitamin E	HPLC	AOAC 992.03; AOAC 2012.10
Vitamin A	HPLC	AOAC 974.29; Bieri et al. (1979)
Carotenoids	HPLC	Delia & Mieko (2004)
Vitamin D2 & D3	RP- HPLC-APCI-MS.	AOAC 982.29
Vitamin K1 & K2	RP- HPLC-APCI-MS	AOAC 992.27
Free sugars	HPLC-RID	AOAC 2013.12
Starch	Calculated from the increase in the amount of glucose by using conversion factor	Casterline et al. (1998)
Oligosaccharides (Raffinose, Stachyose, Verbascose and Ajucose)	HPLC	Frias et al. (1994); Johansen et al. (1996); Sanchez-Mata et al. (1998)
Iron, Zinc, Copper, Manganese, Potassium, Magnesium, Sodium, Calcium, Aluminium	Atomic Absorption Spectrophotometry (AAS)	
Lead, Cadmium, Nickel, Cobalt, Chromium, Lithium, Molybdenum	Graphite furnace Atomic Absorption Spectrophotometry	AOAC 968.08; AOAC 999.10; AOAC 999.11
Arsenic, Antimony, Mercury, and Selenium	Atomic Absorption Spectrophotometry (Hydride generation)	
Phosphorus	Colorimetry	AOAC 935.45; AOAC 995.11; Fiske & Subbarow (1925)
Heavy and trace metals Lead, Mercury, Antimony, Cadmium, Molybdenum, Selenium, Arsenic, Cobalt, Nickel, Chromium, Aluminum, Lithium	Inductively Coupled Plasma Mass Spectrometry (ICP-MS)	AOAC 2013.06
Citric acid, Malic acid, Fumaric acid, Succinic acid, Lactic acid, Cis-Aconitic acid, Tartaric acid and Quinic acid	HPLC	AOAC 969.30; AOAC 950.35; AOAC 948.14

Table 5. Continue...

Analyte	Method	Reference
Oxalic acid (Total)		
Soluble oxalic acid	Fast- HPLC	Moreau & Savage (2009)
Insoluble oxalic acid		
Fatty acid profile	Gas Chromatography with Flame Ionization Detector	AOAC 996.06; AOAC 996.01; AOAC 2012.13
Amino acid profile	Amino acid analyzer (Ion exchange chromatography)	AOAC 994.12; AOAC 985.28; Landry & Delhay (1992)
Stigmasterol, β -Sitosterol, Campesterol, Ergosterol, 5- α -Cholestenol	Gas Chromatography (GC)	AOAC 922.06; Sorenson & Sullivan (2007)
Polyphenols	HPLC	Sakakibara et al. (2003)
Phyates	Ion Exchange and Spectrophotometry	AOAC 986.11-1988
Total Saponin	Colorimetry	Dini et al. (2009)

parameters in each food representing six regions of the country to enable region specific nutritional and epidemiological data interpretation. For every food sampled under this design, composite nutrient means are determined for each region and the means of six regions form the country wide average, wherein the

standard deviation provides estimates of the content variability. The ambitious plan could be implemented due to the generous funds received from the Indian Council of Medical Research, Department of Health Research, Government of India, New Delhi.

General information

The new Indian Food Composition Tables (IFCT) 2017 is the major source of food composition data in India, generated, developed, managed and maintained by the National Institute of Nutrition (ICMR), Hyderabad. All data except for poultry and egg pertains to raw food. None of the food sampled for analysis is fortified and it represents only inherent values. The uniqueness of the IFCT is that all nutrient values presented here have been derived from comprehensive national food sampling followed by analysis and contains only analytical data. The data set were carefully scrutinized in detail for consistency and validity, using FAO/INFOODS 'Guidelines for Checking Food Composition Data prior to Publication of a User Tables/Database – version 1.0 (2012)'.

1. Food identification

The Food entity is described by its most common English name, scientific name and photographs of the food sampled for analysis in order to provide as much description as possible. Scientific names consisting of the genus and the species are provided for clarity and to avoid confusion. Foods have been arranged alphabetically by English name for easy identification. The names of foods are also listed in regional official Indian languages namely Assamese, Bengali, Gujarati, Hindi, Kannada, Kashmiri, Konkani, Malayalam, Manipuri, Marathi, Nepali, Oriya, Punjabi, Sanskrit, Telugu and Urdu to make them acquiescent with the local or regional needs. For animal source foods, the specific body part sampled for analysis is also given. Each food tabulated has a unique four character alpha-numeric identification code. The alphabetic food character indicates the food group and the numeric character(s) represent a record position within a category. This allows convenient tracking of the data for each food throughout the food composition database.

2. Food groups

Foods with common characteristics have been placed together and arranged in groups (**Table 1**). Each food group is preceded by text covering points of specific relevance to the foods in that group. All foods have been categorized into 20 food groups and the number in parenthesis indicates the total number of foods present in each group. A total of 528 foods have been analyzed for more than 150 parameters and presented under different nutrient component parameter.

3. Information on nutrients

3.1 Proximate components include moisture, protein, total fat, ash and carbohydrates. The Difference between 100 and the sum of proximate components including total dietary fibre represents carbohydrate 'by difference'.

3.2 Available carbohydrate is the sum of total free sugars and total starch. Free sugars are individual monosaccharides (galactose, glucose and fructose) and disaccharides (sucrose, lactose and maltose). The values for available carbohydrate have generally been obtained from the sum of individually analyzed values for these components and will differ with figures for carbohydrate 'by difference'. Available carbohydrates are those which is digested and absorbed, and are glucogenic in human corresponding to the term 'glycaemic carbohydrates'. For conversion between carbohydrate weights and monosaccharide equivalents, the factors shown in **Table 2** was used.

3.3 Total dietary fibre (TDF) is made of complex and heterogeneous polymeric materials such as soluble and insoluble polysaccharides and non-digestible oligosaccharides, as well as a range of non-swelling, more or less hydrophobic compounds such as cutins, suberins and lignins. Currently three different AOAC approved methods for measuring TDF values are available. TDF was analyzed using enzymatic gravimetric method (AOAC 991.43).

3.4 Protein values were calculated from the estimated total nitrogen in the food using the Jones (1941) conversion factors (**Table 3**). Protein values for soybean and mushrooms contain considerable non-protein nitrogen (NPN) in several form including urea, purines & pyrimidine. Therefore, NPN was estimated and subtracted from total nitrogen of mushrooms and soybean before multiplying with the Jones factor.

Table 1. Food groups in the IFCT.

Code	Food groups	No. of food entries
A	Cereals and Millets	24
B	Grain Legumes	25
C	Green Leafy Vegetables	34
D	Other Vegetables	78
E	Fruits	68
F	Roots and Tubers	19
G	Condiments and Spices	33
H	Nuts and Oil Seeds	21
I	Sugars	2
J	Mushrooms	4
K	Miscellaneous Foods	2
L	Milk and Milk Products	4
M	Egg and Egg Products	15
N	Poultry	19
O	Animal Meat	63
P	Marine Fish	92
Q	Marine Shellfish	8
R	Marine Mollusks	7
S	Fresh Water Fish and Shellfish	10
T	Edible Oils and Fats	14

Table 2. Conversion of carbohydrate weights to monosaccharide equivalents.

S. No.	Carbohydrate	Equivalent after Hydrolysis (g/100g)	Conversion to monosaccharide equivalent
1	Monosaccharides e.g. glucose	100	No conversion necessary
2	Disaccharides e.g. sucrose, lactose, maltose	105	x 1.05 or ÷ 0.95
3	Oligosaccharides		
	a. Raffinose (trisaccharide)	107	x 1.07 or ÷ 0.93
	b. Stachyose (tetrasaccharide)	108	x 1.08 or ÷ 0.93
	c. Verbascose (pentasaccharide)	109	x 1.09 or ÷ 0.92
4	Polysaccharides e.g. starch	110	x 1.10 or ÷ 0.90

Table 3. Jones factors for conversion of nitrogen to protein.

Food	Conversion Factor
Barley and its Flour; Rye and its flour; Oats	5.83
Rice and its flour	5.95
Wheat whole	5.83
Wheat bran	6.31
Refined wheat flour (Maida)	5.70
Almonds	5.18
Peanuts	5.46
Soybean	5.71
Cashew nuts, pistachio nut, Chest nut	5.30
Egg whole	6.25
Meat and Fish	6.25
Milk and milk products	6.38
Sunflower seeds	5.30
Food where specific factor is not listed	6.25

3.5 Total fat content of foods is determined by gravimetric methods, including acid hydrolysis and extraction methods using a mixed solvent system of chloroform and methanol. Total fat includes the triglyceride and other lipid components such as glycerol, sterols and phospholipids.

3.6 Food energy is expressed in kilojoules (kJ) calculated based on the Atwater system for determining energy values (Table 4). The data represent physiologically available energy, which is the energy value remaining after digestive and urinary losses are deducted from gross energy. Food energy in kilojoules (kJ) can be converted using the formula 1 kcal = 4.18 kJ.

3.7 Minerals and trace elements included in the table are Aluminium, Arsenic, Cadmium, Calcium, Chromium, Cobalt, Copper, Iron, Lead, Lithium, Magnesium, Manganese, Mercury, Molybdenum, Nickel, Phosphorus, Potassium, Selenium, Sodium and Zinc. Except phosphorus, levels of all minerals and trace elements are determined by atomic absorption spectroscopy (AAS) or inductively coupled plasma

mass spectrophotometry (ICP-MS). Phosphorus was determined by spectrophotometric method.

3.8 Thiamin, Riboflavin and Niacin: Thiamin was determined chemically by the thiochrome procedure and riboflavin by microbiological method. Niacin is determined by microbiological method. The values represent only preformed niacin and do not include the niacin contributed by tryptophan, a niacin precursor. Niacin equivalent applies to the sum of the preformed niacin and the amount that could be derived from tryptophan can be calculated as

Niacin equivalents (mg) = Niacin (mg) + (Tryptophan, mg / 60).

3.9 Vitamin C data include both ascorbic and L-dehydroascorbic acids, as both forms are biologically active, the sum of which is expressed as total ascorbic acid content.

3.10 Vitamin B6 occurs in foods as pyridoxine, pyridoxal, pyridoxamine and their phosphates which were estimated by HPLC. The sum of all forms represents vitamin B6.

3.11 Pantothenic acid in foods was determined by liquid chromatographic methods.

Table 4. Metabolizable energy conversion factors.

Component	kJ/g	kcal/g
Protein	17	4
Fat	37	9
Available carbohydrate	17	4
Fibre	8	2
Alcohol	29	7

3.12 Food folate values are analyzed using the trienzyme procedure and liquid chromatography. Food folate was expressed as the sum of one or more individual folate vitamers namely 5-methyl tetrahydrofolate, 10-formyl folic acid, 5-formyl tetrahydrofolic acid and tetrahydrofolic acid.

3.13 Vitamin A content is expressed as retinol and provitamin A carotenoid. The groups of plant pigments that are precursors of vitamin A are called carotenoids. Individual carotenoids available in the IFCT includes Lutein, zeaxanthin, lycopene, β -cryptoxanthin, α -carotene, β -carotene, γ -carotene and total carotenoids. The body can utilize these inactive forms once they are converted to the active form, retinol. Vitamin A activity values in Retinol Activity Equivalents (RAE) are calculated from the content of individual carotenoids using appropriate factors. Incidentally, there are many conversion factors for carotenoids and no universal factor exists.

3.14 Vitamin D have been reported in the IFCT which is lacking in many food composition databases. Ergocalciferol or vitamin D2 content in plant foods measured by LC-MS has been reported for the first time in the world. Cholecalciferol or vitamin D3 in animal source foods including 25 hydroxy-D3 measured by LC-MS/MS has also been reported. The availability of new analytical vitamin D data in foods will permit dietary intake assessment. Vitamin D2 (ergocalciferol) made commercially has been reported to have the similar potency in human. Total vitamin D activity has been taken as the sum of vitamin D3 (cholecalciferol) and five times 25-hydroxy vitamin D3 (25-hydroxy cholecalciferol), where data are available.

3.15 Vitamin E in food is presented as tocopherols and tocotrienols and its isomers. The new Dietary Reference Intake (DRI) for vitamin E (Krinsky, 2000) has stated that with the exception of α -tocopherol all other individual isomers of tocopherols and tocotrienols do not contribute to vitamin E activity, nevertheless, they are included in the database.

3.16 Vitamin K data as phyloquinone in plant foods and menaquinone in animal foods estimated using liquid chromatography are presented in the IFCT.

3.17 Fatty acid profile is obtained as the percentage of fatty acid methyl esters determined by gas liquid chromatographic (GLC) analysis. Fatty acids are expressed as the actual quantity of fatty acid in mg per 100g of edible portion of food calculated using Sheppard conversion factor as detailed in Weihrauch et al. (1977). Detailed information on individual fatty acids of each food including values for total saturated, monounsaturated and polyunsaturated fatty acids are reported.

3.18 Amino acid profile of each food is determined by three different analyses. Tryptophan is determined by alkaline hydrolysis, methionine and cystine by performic oxidation and the rest of the amino acids by acid hydrolysis. The amino acid profile of each food is expressed as g/100 g protein.

3.19 Organic acids is naturally present in a wide

variety of foods especially fruits, berries and vegetables. Organic acids *cis*-aconitic acid, citric acid, fumaric acid, mallic acid, quinic acid, succinic acid and tartaric acid were determined in single liquid chromatographic run. Soluble, insoluble and total oxalates were determined separately by HPLC method. The organic acids are energy contributing components, although it varies between the different organic acids. According to the Codex Alimentarius Commission's Guidelines for Nutrition Labeling, the energy conversion factor for organic acids is 13 kJ/g. However, organic acids have not been included in the total energy of foods given in the IFCT.

3.20 Polyphenols exhibit highly diverse structures characterized by the presence in their structure of one or several phenolic groups. The recognition of the antioxidant properties of polyphenols, and their probable role in the prevention of various diseases associated with oxidative stress has generated great interest. Polyphenols are found abundantly in our diet and over 500 different molecules are known to be present in foods. The IFCT presents data on 37 individual and total polyphenols present in key Indian foods.

3.21 Cholesterol values are generated primarily by gas liquid chromatographic method in foods of animal origin assuming that cholesterol is present only in animal origin foods. Cholesterol values are expressed as mg/100g food. To convert to mmol cholesterol, divide the values by 386.6.

3.22 Plant sterols data on campesterol, stigmasterol, and β -sitosterol are obtained by gas-chromatographic procedures and summed to calculate total phytosterols.

3.23 Oligosaccharides are indigestible but the beneficial colonic bacteria break them down in the large intestine. In the recent times there is great interest in the health properties of non-digestible oligosaccharides (NDO) as prebiotic compounds. Data on the oligosaccharide content in foods are very meager, making it difficult to assess oligosaccharide consumption of any population in the world. Therefore, content of raffinose, stachyose, verbascose and ajugoes in key Indian foods is presented.

3.24 Saponins consist of a polycyclic aglycones attached to one or more sugar side chains and are found in most vegetables, beans and herbs. Saponins were not destroyed by processing or cooking and have many health benefits.

3.25 Phytate is commonly known as phytic acid, inositol hexaphosphate and IP-6. It has received considerable attention as anti-nutrient because it binds minerals in the digestive tract, making them less available metabolically to our bodies. However, in the recent years, some novel metabolic effects of phytate

or some of its degradation products have been recognized as antioxidant and may help fight some cancers and other diseases.

4. The expression of nutrient values

All foods are expressed per 100g edible portion. Generally the values have been expressed to a constant number of decimal places for each nutrient. The values of the more variable vitamins or minerals and trace elements have been expressed to less than their usual number of places where greater values render the extra places non-significant.

5. Nutrient identification-Tagnames

A unique abbreviation for a food component developed by International Network of Food Data Systems (INFOODS), have been used to aid in data interchange. Tagnames unambiguously indicate how the nutrient data are obtained indicating the methods of analysis used for the data. In case data was derived by calculation, tagname even indicates the formula used for calculation. INFOODS Tagnames, utilized worldwide, facilitate international data interchange and harmonization.

6. Data documentation

The alpha-numeric code provides a unique food identifier within the IFCTs. The foods are sorted alphabetically in the tables. All analytical data scrutinized against quality control data, unit of expression, conversion factors and calculations applied to each nutrient data and accepted after all criteria were met. The metadata was compiled in the archival database which has all the information on the sampling plan, numbers of samples analyzed, analytical methods used and quality assurance procedures in place.

The data presented in this book is the nutrient mean of all six regions representing the country wide value where the standard deviation represents the variation of the particular nutrient. The second type of sample is the individual sample of rice and wheat collected from the 107 sampling sites and analysed individually which increased variability by increasing the effective sample size. Such samples were drawn only for the main staples in the country and not for other foods due to the high cost associated with such analysis. The third type of samples are those of food within the species which were also analysed according to varieties/cultivars for some foods.

The values of six separate regions namely East, Northeast, North, West, South and Central India obtained from compositing adequate number of similar foods collected from within the region was tabulated. The mean value of six regional data was then used to calculate the national value with standard deviation to represent the variation within each nutrient. The mean $\pm$ SD of all nutrients and bioactive components in foods are tabulated to represent the user database or the Indian Food Composition tables (**Figure 1**). All blank spaces in the nutrient tables represents below detectable limit of the particular method used for analysis.

Electronic format allow capturing higher volume of data, easier to update and can reach wider distribution if they are made publicly available. The region specific database is available in the electronic version at the National Institute of Nutrition (NIN). Regional foods may have a different nutrient profile that may differ from national average and hence individuals seeking local specific nutrient data are encouraged to obtain region specific data from NIN.




Figure 1. Indian food composition tables-data flow

References

- Ahuja, J. K., Moshfegh, A. J., Holden, J. M., & Harris, E. (2013). USDA food and nutrient databases provide the infrastructure for food and nutrition research, policy, and practice. *The Journal of nutrition*, 143(2), 241S-249S.
- Annual Report of GoI (1905). *Annual report of the sanitary commissioner with the Government of India (GoI)*. pp. 126
- AOAC (1930). *Association of Official Agricultural Chemists. Official and tentative methods of analysis*. (3rded) Washington.
- AOAC (2005). Amino acids in feeds performic acid oxidation with acid hydrolysis sodium metabisulfite method. *AOAC Official Method* 994.12.
- AOAC (2005). Loss on Drying (Moisture) at 95°–100°C for Feeds Dry Matter on Oven Drying at 95°–100°C for Feeds. *AOAC Official Method* 934.01.
- AOAC (2005). Acetic and propionic acids in bread chromatographic method. *AOAC Official Method* 950.35.
- AOAC (2005). Determination of Lead, Cadmium, Copper, Iron, and Zinc in Foods. autoanalyzer/ flame atomic absorption spectroscopy, autoanalyzer/ graphite furnace atomic absorption spectroscopy after dry ashing. *AOAC Official Method* 999.11.
- AOAC (2005). Fat in flour. Acid hydrolysis method. *AOAC Official Method* 922.06.
- AOAC (2005). Lead, cadmium, zinc, copper, and iron in foods. autoanalyzer/graphite furnace atomic absorption spectroscopy, autoanalyzer/flame atomic absorption spectroscopy, after microwave digestion. *AOAC Official Method* 999.10.
- AOAC (2005). Minerals in animal feed and pet food, Atomic absorption spectrophotometric method. *AOAC Official Method* 968.08.
- AOAC (2005). Niacin and niacinamide (nicotinic acid and nicotinamide) in ready-to-feed milk-based infant formula microbiological-turbidimetric method. *AOAC Official Method* 985.34.
- AOAC (2005). Organic acids (foreign) in fruit juices chromatography/paper chromatography. *AOAC Official Method* 969.30.
- AOAC (2005). Phosphorus (total) in foods colorimetric method. *AOAC Official Method* 995.11.
- AOAC (2005). Phosphorus in fruits and fruit products colorimetric method. *AOAC Official Method* 935.45.
- AOAC (2005). Phytate in foods anion exchange method. *AOAC Official Method* 986.11.
- AOAC (2005). Protein (Crude) in Animal Feed, Forage (Plant Tissue), Grain, and Oilseeds. *AOAC Official Method* 2001.11.
- AOAC (2005). Quinic, Malic, and citric acids in cranberry juice cocktail and apple juice. *AOAC Official Method* 986.13.
- AOAC (2005). Riboflavin (Vitamin B₂) in vitamin preparations microbiological methods. *AOAC Official Method* 940.33.
- AOAC (2005). Succinic Acid in Eggs. *AOAC Official Method* 948.14.
- AOAC (2005). Sulfur amino acids in food, feed ingredients, and processed foods ion exchange chromatographic method. *AOAC Official Method* 985.28.
- AOAC (2005). Thiamine (Vitamin B₁) in human and pet foods. *AOAC Official Method* 942.23.
- AOAC (2005). Total dietary in foods—enzymatic gravimetric method. *AOAC Official Method* 985.29.
- AOAC (2005). Total, soluble, and insoluble dietary fiber in foods, enzymatic-gravimetric method. *AOAC Official Method* 991.43.
- AOAC (2005). Trans-vitamin K₁ (Phylloquinone) in ready-to-feed milk-based infant formula. *AOAC Official Method* 992.27.
- AOAC (2005). Vitamin A in mixed feeds, premixes, and human and pet foods. *AOAC Official Method* 974.29.
- AOAC (2005). Vitamin assays, Turbidimetric Method. *AOAC Official Method* 960.46.
- AOAC (2005). Vitamin B₆ (Pyridoxine, Pyridoxal, Pyridoxamine) in ready to feed milk based infant formula. 50.1.18. *AOAC Official Method* 985.32.
- AOAC (2005). Vitamin C (Reduced Ascorbic Acid) in ready-to-feed milk-based infant formula. *AOAC Official Method* 985.33.
- AOAC (2005). Vitamin D in mixed feeds, premixes, and pet foods. *AOAC Official Method* 982.29.
- AOAC (2012). Arsenic, Cadmium, mercury, and lead in foods. Pressure Digestion and Inductively Coupled Plasma-Mass Spectrometry. *AOAC Official Method* 2013.06.
- AOAC (2012). Ash of Animal Feed. *AOAC Official Method* 942.05.
- AOAC (2012). Determination of labeled fatty acids content in milk products and infant formula Capillary Gas Chromatography. *AOAC Official Method* 2012.13.
- AOAC (2012). Fat (total, saturated, and unsaturated) in foods hydrolytic extraction gas chromatographic method. *AOAC Official Method* 996.06.
- AOAC (2012). Fat (total, saturated, unsaturated, and monounsaturated) in cereal products acid hydrolysis capillary gas chromatographic method. *AOAC Official Method* 996.01.
- AOAC (2012). Folate in infant formula and adult/pediatric nutritional formula. UHPLC-MS/MS. Chromatography/Liquid Chromatography. *AOAC Official Method* 2013.13.
- AOAC (2012). Simultaneous determination of 13-cis and all-trans vitamin A palmitate (Retinyl Palmitate), vitamin A acetate (Retinyl Acetate), and total vitamin E (DL- α -Tocopherol and DL- α -

- Tocopherol Acetate) in infant formula and adult nutritional. *AOAC Official Method* 2012.10.
- AOAC (2012). Total carbohydrates in wine and wine-like beverages. HPLC with a refractive index detector. *AOAC Official Method* 2013.12.
- AOAC (2012). Total folates in infant formula and adult nutritional trienzyme extraction and UPLC-MS/MS quantitation. *AOAC Official Method* 2011.06
- AOAC (2012). Vitamin E activity (All-rac-alpha-Tocopherol) in milk-based infant formula, Liquid chromatography. *AOAC Official Method* 992.03.
- AOAC (2012). Fat in Cacao Products. Soxhlet extraction method. Gravimetric. *AOAC Official Method* 963.15.
- Archer, E., Hand, G. A., & Blair, S. N. (2013). Validity of US nutritional surveillance: National health and nutrition examination survey caloric energy intake data, 1971–2010. *PloS One*, 8(10), e76632.
- Atwater, W. O. (1889). Food and Health. *Public Health Papers and Reports*, 15, 208-221
- Aykroyd, W. R. & Rajagopal, K. (1936). The State of Nutrition of School Children in South India. Part I. *Indian Journal of Medical Research*, 24, 419-37.
- Aykroyd, W. R., & Swaminathan, M. (1940). The nicotinic acid contents of cereals and pellagra. *Indian Journal of Medical Research*, 27, 667-678
- Aykroyd, W. R. (1932). The effect of parboiling and milling on the antineuritic vitamin (B 1) and phosphate content of rice. *Journal of Hygiene*, 32(2), 184-192.
- Aykroyd, W. R. (1937). *The nutritive value of Indian foods and the planning of satisfactory diets*. Health Bulletin No. 23. Government of India (GoI) press, Simla. pp.48
- Aykroyd, W. R. (1941). Economic aspects of the problem of nutrition in India. *Indian Journal of Social Work*, 2, 1-14.
- Aykroyd, W. R., Patwardhan, V. N., & Ranganathan, S. (1951). *The Nutritive Value of Indian Foods and the Planning of Satisfactory Diets*. Health Bulletin, No. 23. pp. 79
- Aykroyd, W.R. & Patwardhan, S. (1956). *The Nutritive Value of Indian Foods and the Planning of Satisfactory Diets*. (5th ed). Health bulletin No. 23. New Delhi, India.
- Aykroyd, W.R., Gopalan, C., & Balasubramanian, S. C. (1963). *The nutritive value of Indian Foods and the planning of satisfactory diets*. (6th revised ed). Special report No. 42. Indian Council of Medical Research, New Delhi.
- Aykroyd, W.R., Gopalan, C., & Balasubramanian, S. C. (1966). *The nutritive value of Indian Foods and the planning of satisfactory diets*. (6th revised ed). Special report No. 42. Indian Council of Medical Research, New Delhi.
- Ayoya, M. A., Kodio, J., Iknane, A. A., & Sodjinou, R. (2010). Nutritional value of locally produced foods and potential for developing age-appropriate complementary foods for children under 2 years of age in Mali. *Food and Nutrition Bulletin*, 31(3), 391-399.
- Barry, C. (1900). Notes on beri-beri in Rangoon, *Indian Medical Gazette*, 35, 343–345
- Bieri, J. G., Tolliver, T. J., & Catignani, G. L. (1979). Simultaneous determination of alpha-tocopherol and retinol in plasma or red cells by high pressure liquid chromatography. *The American Journal of Clinical Nutrition*, 32(10), 2143-2149.
- Casterline Jr, J. L., Oles, C. J., & Ku, Y. (1998). Measurement of sugars and starches in foods by a modification of the AOAC total dietary fiber method. *Journal of AOAC International*, 82(3), 759-765.
- Census of India (2001). The Registrar General & Census Commissioner, India, New Delhi, Ministry of Home Affairs, Government of India. (www.censusindia.gov.in).
- Chakrabarty, M. M., & Hilditch, T. P. (1951). The fatty acids and glycerides of an Indian sesame oil. *Journal of the Science of Food and Agriculture*, 2(6), 255-259.
- Chaudhuri, A., & Vos, J. W. E. (1988). *Unified theory and Strategies of survey Sampling*. NHC: New York.
- Chiu, T. H., Huang, H. Y., Chiu, Y. F., Pan, W. H., Kao, H. Y., Chiu, J. P., Lin, M. N., & Lin, C. L. (2014). Taiwanese vegetarians and omnivores: dietary composition, prevalence of diabetes and IFG. *PloS One*, 9(2), e88547.
- Cochran, W. G. (1977). *Sampling Techniques*. Wiley, New York, NY.
- Cunningham, J., & Sobolewski, R. (2011). Food composition databases for nutrition labelling: Experience from Australia. *Journal of Food Composition and Analysis*, 24(4), 682-685.
- David, C. W., & Harvey, E. I. (2000). The analysis of pantothenic acid in milk and infant formulas by HPLC. *Food Chemistry*, 69, 201-208.
- Day, C. M. (1944). Nutritional deficiencies and dental caries in northern India. *British Dental Journal*, 76(5), 115-22.
- Deharveng, G., Charrondiere, U. R., Slimani, N., Southgate, D.A., & Riboli, E. (1999). Comparison of nutrients in the food composition tables available in the nine European countries participating in EPIC. European Prospective Investigation into Cancer and Nutrition. *European Journal of Clinical Nutrition*, 53, 60–79.
- Delia, B. R-A., & Mieko, K. (2004). *HarvestPlus Handbook for Carotenoid Analysis*. Harvest plus technical monograph Series 2. International Food Policy Research Institute. Washington, DC USA. pp-58.
- Devadas, R. P., Usha, T. M., Sankarj, L., Rajalakshmi, R. S., Pateath, G., & Babbtiwale, M. (1965). Diet and nutrition survey of a village community in South India. *Indian Journal of Nutrition and Dietetics*, 2, 83-87.

- Dini, I., Tenore, G. C., & Dini, A. (2009). Saponins in *Ipomoea batatas* tubers: Isolation, characterization, quantification and antioxidant properties. *Food Chemistry*, 113(2), 411-419.
- Drewnowski, A., Rehm, C. D., Martin, A., Verger, E. O., Voinnesson, M., & Imbert, P. (2015). Energy and nutrient density of foods in relation to their carbon foot print. *The American Journal of Clinical Nutrition*, 101(1), 184-191.
- Dwyer, J. T. (1994). Future directions in food composition studies. *Journal of Nutrition*, 124, 1783S–1788S
- Ernest, E. W. (1917). *Diabetes, its causation and treatment (with special reference to India)*. Calcutta Thacker, Spink and Co. Agricultural Research Institute Pusa. pp. 170
- FAO/INFOODS (2012). FAO/INFOODS Guidelines for checking food composition data prior to the publication of a user table/database. <http://www.fao.org/docrep/017/ap810e/ap810e.pdf>
- FAO/WHO (1998). *Carbohydrates in Human Nutrition*. Report of a Joint FAO/WHO Expert Consultation. Food and Nutrition Paper No. 66. 14-18 April 1997, Rome, Italy.
- FAO/WHO (1994). *Fats and oils in human nutrition*. Report of a Joint FAO/WHO Expert Consultation. Food and Nutrition Paper No. 91. 19-26 October 1993, Rome, Italy.
- FAOSTAT (2013). Food and Agriculture Organization of the United Nations, FAOSTAT database (2013). <http://faostat.fao.org/> Accessed 12.03.16
- Finglas, P. M., Wigertz, K., Vahteristo, L., Witthoft, C., Southon, S., & de Froidmont-Gortz, I. (1999). Standardisation of HPLC techniques for the determination of naturally-occurring folates in food. *Food Chemistry*, 64 (2), 245-255.
- Fiske, C. H., & Subbarow, Y. (1925). The colorimetric determination of phosphorus. *Journal of Biological Chemistry* 66(2), 375-400
- Foy, H., & Kondi, A. (1957). Anaemias of the tropics: Relation to iron intake, absorption and losses during growth, pregnancy and lactation. *Journal of Tropical Medicine and Hygiene*, 60(5), 105-18.
- Frias, J., Hedley, C. L., Price, K. R., Fenwick, G. R., & Vidal-Valverde, C. (1994). Improved method of oligosaccharide analysis for genetic study of legume seeds. *Journal of Liquid Chromatography*, 17, 2469-2484.
- Gopalan, C., & Ramalingaswami, V. (1955). Kwashiorkor in India. *Indian Journal of Medical Research*, 43(4), 751-73.
- Gopalan, C. (1961). *A report of some recent studies on protein malnutrition in India*. Progress in meeting protein needs of infants and preschool children. Proceedings of an International Conference., Washington D.C., NRC Publication 843, pp. 211-223.
- Gopalan, C. (1970). Some recent studies in the Nutrition Research Laboratories, Hyderabad. *The American Journal of Clinical Nutrition*, 23(1), 35-51.
- Gopalan, C., Rama Sastri, B. V., Balasubramanian, S. C., Revised and updated by Narasinga Rao, B. S., Deosthale, Y. G., & Pant, K.C. (1989). *Nutritive Value of Indian Foods*. Revised and updated Edition. National Institute of Nutrition, Indian Council of Medical Research, Hyderabad.
- Gopalan, C., Ramasastry B. V. & B. V. Balasubramanyam (1971). *Nutritive values of Indian Foods*, National Institute of Nutrition (ICMR), Hyderabad (India).
- Goswami, H., & Basu, U. P. (1938). 'Available' iron in Indian foodstuffs. *Indian Journal of Medical Research*, 25, 893-899.
- Greenfield, H., & Southgate, D. A. T. (2003). Food composition data. Production, management and use. 2nd edition. Rome: *Food and Agricultural Organization of the United Nations*.
- Greenfield, H., Arcot, J., Barnes, J. A., Cunningham, J., Adorno, P., & Stobaus, T., et al. (2009). Nutrient composition of Australian retail pork cuts 2005/2006. *Food Chemistry*, 117, 721–730.
- Harinarayan, C. V., Ramalakshmi, T., Prasad, U. V., Sudhakar, D., Srinivasarao, P. V., Sarma, K. V., & Kumar, E. G. T. (2007). High prevalence of low dietary calcium, high phytate consumption, and vitamin D deficiency in healthy south Indians. *The American Journal of Clinical Nutrition*, 85(4), 1062-1067.
- Haytowitz, D. B., Pehrsson, P. R. & Holden, J. M. (2000). Setting priorities for nutrient analysis in diverse populations. *Journal of Food Composition and Analysis*, 13, 425-433.
- Haytowitz, D. B., Pehrsson, P. R., Smith, J., Gebhardt, S. E., Matthews, R. H., & Anderson, B. A. (1996). Key foods: setting priorities for nutrient analyses. *Journal of Food Composition and Analysis*, 9(4), 331-364.
- Holler, U., Wachter, F., Wehrli, C., & Fizet, C. (2006). Quantification of biotin in feed, food, tablets, and premixes using HPLC–MS/MS. *Journal of Chromatography B*, 831, 8–16.
- IDF (2014). International Diabetes Federation. IDF diabetes atlas. International Diabetes Federation, Executive Office. www.idf.org. Accessed 29.03.16
- Jansen, L., & Roodenburg, A. J. (2016). The use of food composition data in the choices international programme. *Food Chemistry*, 193, 196-202.
- Johansen, H. N., Glitso, V., & Bach-Knudsen, E. K. (1996). Influence of extraction solvent and temperature on the quantitative determination of oligosaccharides from plant materials by high-performance liquid chromatography. *Journal of Agricultural and Food Chemistry*, 44, 1470-1474.
- Jones, D. B. (1941). Factors for converting percentages of nitrogen in foods and feeds into percentages of proteins. Circ No.183 (slight revision). Washington, D.C.: U.S. Department of Agriculture.

- Keating, R. W., & Haddad, P. R. (1982). Simultaneous determination of ascorbic acid and dehydroascorbic acid by reversed-phase ion-pair high-performance liquid chromatography with pre-column derivatisation. *Journal of Chromatography A*, 245(2), 249-255.
- Krinsky, N. I., Beecher, G. R., Burk, R. F., Chan, A. C., Erdman, J. W., Jr., Jacob, R. A., et al. (2000). Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. Washington DC: National Academy Press.
- Landry, J., & Delhay, S. (1992). Simplified procedure for the determination of tryptophan of foods and feed stuffs from barytic hydrolysis. *Journal of Agricultural and Food Chemistry*, 40, 776-779.
- Lewis, T. R. (1880). *Dieteries of laboring prisoners in Indian jails*. Annual report of sanitary commissioner of government of India (GoI), for 1880, pp. 159
- MacIntosh, D. L., Williams, P. L., Hunter, D. J., Sampson, L. A., Morris, S. C., Willett, W. C., & Rimm, E. B. (1997). Evaluation of a food frequency questionnaire-food composition approach for estimating dietary intake of inorganic arsenic and methylmercury. *Cancer Epidemiology Biomarkers & Prevention*, 6(12), 1043-1050.
- Malhotra, S. L. (1970). Dietary factors causing hypertension in India. *The American Journal of Clinical Nutrition*, 23(10), 1353-1363.
- Master, F., & Magar, N. G. (1954). Studies in the nutritive value of Bombay fish. 2. Amino acid composition. *Indian Journal of Medical Research*, 42, 509-513.
- McCance, R. A., & Widdowson, E. M. (1960). The Composition of Foods, Medical Research Council Special Report Series No. 297. London
- McCarrison, R. (1925). The Relationship of Diet to the Physical Efficiency of Indian Races. *Practitioner*, 114, No. 1 (679.), 90-100.
- McCarrison (1929). '*Food*' Macmillan and Co., Madras.
- McCarrison, R., & Norris, R. V. (1924). The Relationship of Rice to Beri-Beri in India. *Indian Medical Research Memoirs*, (Memoir 2), 1-87.
- McCarrison, R. (1932). Problems of Nutrition in India. *Nutrition Abstracts and Reviews*, 2, 1-8.
- McCarrison, R. (1936). Nutrition and national health. *British Medical Journal*, 1, No. 3921, 427-430
- McCay D. (1912). *Protein elements in nutrition*. XVI. pp. 214
- McCay, D. (1910). *Investigations on Bengal jail dietaries with some observations on the influence of dietary on the physical development and well being of the people of Bengal. Scientific memoirs by officers of the medical and sanitary departments of the government of India (GoI)*, New Series, No. 37
- McCay, D. (1911). *Investigations into the jail dietaries of the united provinces with some observations on the physical development and well being of the people of United Provinces. Scientific memoirs by officers of medical and sanitary department of government of India (GoI)*, New Series, No. 48. pp. 200
- McNamara, R. J. (1906). *Notes on Indian jail dietaries with special reference to the Punjab*. (Quoted by McCay, 1910)
- Merrill, A. L., & Watt, B. K. (1973). Energy value of foods, basis and deviation. Human Nutrition Research Branch. Agriculture Research Service, United States Department of Agriculture. Agriculture Handbook No.74.
- Moreau, A. G., & Savage, G. P. (2009). Oxalate content of purslane leaves and the effect of combining them with yoghurt or coconut products. *Journal of Food Composition and Analysis*, 22(4), 303-306.
- Narasinga Rao B. S. (2005a). Progress and development of nutrition research in India. Historical perspective Pre-independence period 1900-1950. In B. S. Narasinga Rao (Ed), *Development of nutritional science in India* (pp.40-43). Allied Publishers (P) Limited, India.
- Narasinga Rao B. S. (2005b). Nutrient and non-nutrient composition of Indian foods. In B. S. Narasinga Rao (Ed), *Development of nutritional science in India* (pp. 274-298). Allied Publishers (P) Limited, India.
- National Family Health Survey (NFHS-3) 2005-06. International Institute for Population Sciences, Mumbai.
- National Nutrition Monitoring Bureau (NNMB) (2006). *Diet and Nutritional status of rural population*, National Institute of Nutrition, Indian Council of Medical Research, Hyderabad.
- NFHS-3 (2007). *National Family Health Survey (2005-06)*. International Institute of Population Sciences IIPS: India. Vol. I. Mumbai.
- Nicholls, L. (1945). *Tropical Nutrition and Dietetics*. (2nd ed). Baillibre, Tindall and Cox, London. pp.370
- NNMB (1991). *Report of Repeat Surveys 1988-90*. National Nutrition Monitoring Bureau. National Institute of Nutrition, Hyderabad. Indian Council of Medical Research
- NNMB (2012). *Diet and nutritional status of rural population, prevalence of hypertension, and diabetes among adults and infants and young child feeding practices. Report of third repeat survey (2011-2012)*. National Nutrition Monitoring Bureau. National Institute of Nutrition, Hyderabad. Indian Council of Medical Research, New Delhi.
- Ollilainen, V. M. (2008). HPLC analysis of vitamin B6 in foods. *Agricultural and Food Science*, 8(6), 515-619.
- Passmore, R. (1948). *Nutritional diseases in India*. U.N. Dhur and Sons Ltd., Calcutta
- Patnaik, M. (1934). The Iodine content of Indian foodstuffs. *Indian Journal of Medical Research*, 22, 249-262.

- Patwardhan, V. N. (1956). Protein element in Indian nutrition. *Indian Journal of Medical Sciences*, 10(1), 1-18.
- Patwardhan, V. N. (1961). *Nutrition in India*. (2nd ed) Indian Journal of Medical Sciences. Bombay.
- Patwardhan, V. N. (1962). Pulses and beans in human nutrition. *American Journal of Clinical Nutrition*, 11, 12-30.
- Pehrsson, P. R., Haytowitz, D. B., Holden, J. M., Perry, C. R., & Beckler, D. G. (2000). USDA's National Food and Nutrient Analysis Program: Food sampling. *Journal of Food Composition and Analysis*, 12, 379-389.
- Raman, T. K. (1940). Pellagra in India. A study of twenty-five cases in Vizagapatam. *Indian Journal of Medical Research*, 27, 743-756.
- Ramasastri, B. V. (1975). A simple method for estimation of thiamine in foods using base-exchange on sand instead of Decalso. International journal for vitamin and nutrition research. Internationale Zeitschrift fur Vitamin-und Ernährungsforschung. *Journal International de Vitaminologie et de Nutrition*, 46(4), 403-411.
- Ranganathan, S., Sundararajan, A. R., & Swaminathan, M. (1937). Survey of the nutritive value of Indian foodstuffs. Part I. The chemical composition of 200 common foods. *Indian Journal of Medical Research*, 24, 689-706
- Rao, K. S., Swaminathan, M. C., Swarup, S., & Patwardhan, V. N. (1959). Protein malnutrition in south India. *Bulletin of the World Health Organization*, 20(4), 603.
- Rao, K. V. (1967). Pattern and trends in food consumption in India. *Indian Journal of Nutrition and Dietetics*, 4, 79-87.
- Reddy, V., M. Shekar, Rao, P., & Gillespie, S. (1992). "Nutrition in India." ACC/SCN case study. National Institute of Nutrition, Hyderabad. Indian Council of Medical Research.
- Rice (1959). *Health Bulletin No. 28*. Delhi. Government of India (GoI) Press
- Sahasrabudhe, D. L. (1925). *The chemical composition of the food grains, vegetables and fruits of western India*. Department of Agriculture, Bombay, Bulletin No. 124 of 1925
- Sakakibara, H., Honda, Y., Nakagawa, S., Ashida, H., & Kanazawa, K. (2003). Simultaneous determination of all polyphenols in vegetables, fruits, and teas. *Journal of Agricultural and Food Chemistry*, 51(3), 571-581.
- Sanchez-Mata, M. C., Penuela-Teruel, M. J., Camara-Hurtado, M., Diez-Marques, C., & Torija-Isasa, M. E. (1998). Determination of mono-, di-, and oligosaccharides in legumes by high-performance liquid chromatography using an amino-bonded silica column. *Journal of Agricultural and Food Chemistry*, 46, 3648-3652.
- Scott, A. C. (1916). A contribution to the study of osteomalacia in India. *Indian Journal of Medical Research*, 4(1), 140-168.
- Seljak, B. K., Stibilj, V., Pograjc, L., Mis, N. F., & Benedik, E. (2013). Food composition databases for effective quality nutritional care. *Food Chemistry*, 140(3), 553-561.
- Sheppard, A. J. (1992). *Lipid Manual: Methodology for appropriate fatty acid - cholesterol analysis*. Dubuque, I.A.: U.S. Food and Drug Administration, William C. Brown Publishers.
- Sorenson, W. R., & Sullivan, D. (2007). Determination of campesterol, stigmasterol, and beta-sitosterol in saw palmetto raw materials and dietary supplements by gas chromatography: collaborative study. *Journal of AOAC International*, 90(3), 670-678.
- Stewart, A. D., Boyd, T. C., & De, D. C. (1931). Analyses and calorific values of some Indian foodstuffs. *Indian Journal of Medical Research*, 19, 675-89.
- Su, C., Jia, X., Wang, Z., Wang, H., & Zhang, B. (2015). Trends in dietary cholesterol intake among Chinese adults: a longitudinal study from the china health and nutrition survey, 1991-2011. *BMJ Open*, 5(6), e007532.
- Subrahmanyam, V., Sreenivasan, A., & Swaminathan, M. (1962). *Protein consumption in India and other countries*. Proceedings. National Institute of Sciences of India. No. 20, pp. 134
- Subrahmanyam, V., Swaminathan, M., Rao, M. N. & Joseph, K. (1961). Protein and amino acid requirements of infants, children and adults. *Indian Journal of Pediatrics*, 28(8), 313-334.
- Swaminathan, M. (1937). The relative value of the proteins of certain foodstuffs in nutrition. Part II The comparative biological values of the proteins of certain cereals, pulses and skimmed milk powder, measured by the growth of young rats. *Indian Journal of Medical Research*, 25(2), 57-79
- Tarley, C. R. T., Visentainer, J. V., Matsushita, M., & De Souza, N. E. (2004). Proximate composition, cholesterol and fatty acids profile of canned sardines (*Sardinella brasiliensis*) in soybean oil and tomato sauce. *Food Chemistry*, 88, 1-6.
- Timmayamma, B. V. S., & Rao, H. D. A (1969). Comparative study of the oral questionnaire method with actual observation of the dietary intake of pre-school children. *Journal of Nutrition & Dietetics*, 6, 177.
- Trainer, D., Pehrsson, P. R., Haytowitz, D. B., Holden, J. M., Phillips, K. M., Rasor, A. S., & Conley, N. A. (2010). Development of sample handling procedures for foods under USDA's National Food and Nutrient Analysis Program. *Journal of Food Composition and Analysis*, 23, 843-851.
- Vaask, S., Pomerleau, J., Pudule, I., Grinberga, D., Abaravicius, A., Robertson, A., & Mckee, M. (2004). Comparison of the Micro-Nutrica Nutritional Analysis program and the Russian Food Composition Database using data from the Baltic Nutrition Surveys. *European Journal of Clinical Nutrition*, 58, 573-579.

- Vahteristo, L., Finglas, P. M., Witthöft, C., Wigertz, K., Seale, R., & de Froidmont-Görtz, I. (1996). Third EU MAT intercomparison study on food folate analysis using HPLC procedures. *Food Chemistry*, 57(1), 109-111.
- Velayutham, M., Mandal, D. K., Mandal, C., & Sehgal, J. (1999). Agro-ecological subregions of India for planning and development. *NBSS Publication*, (35).
- Venkatachalam, P. S. (1968). Iron metabolism and iron deficiency in India. *The American Journal of Clinical Nutrition*, 21(10), 1156-1161.
- Weihrauch, J. L., Posati, L. P., Anderson, B. A., & Exler, J. (1977). Lipid conversion factors for calculating fatty acid contents of foods. *Journal of the American Oil Chemists' Society*, 54(1), 36-40.
- Wilson, D. (1937). Nutrition and diet in Northern India: A comparative dietary survey. *The Lancet*, 230, 1445-1448.
- Wilson, H., Ahmad, B., Ray, G., & Guha, R. (1937). The vitamin-B1 content of some common Indian foodstuffs. *Indian Journal of Medical Research*, 24, 813-16.
- Yusufji, D., Mathan, V. I., & Baker, S. J. (1973). Iron, folate, and vitamin B12 nutrition in pregnancy: a study of 1000 women from southern India. *Bulletin of the World Health Organization*, 48(1), 15.

Abbreviations

A.	Assamese	Kh.	Khasi
AAS	Atomic Absorption Spectroscopy	kJ	kilo Joule
AOAC	Association of Official Analytical Chemists	Kon.	Konkani
APCI	Atmospheric Pressure Chemical Ionization	M.	Manipuri
B.	Bengali	Mal.	Malayalam
BV	Biological Value	Mar.	Marathi
CFTRI	Central Food Technological Research Institute	N	Normality
CHO	Carbohydrate	N.	Nepali
Common	Common name	NAC	Nutritional Advisory Committee
CRM	Certified Reference Material	NDO	Non-digestible oligosaccharides
DRI	Dietary reference intake	NFHS	National Family Health Survey
EAA	Essential Amino Acids	NIN	National Institute of Nutrition
ESI	Electro Spray Ionization	NNMB	National Nutrition Monitoring Bureau
FAO	Food and Agriculture Organization of the United Nations	NRFA	Nutrition Research Funds Association
FAOSTAT	Food and Agriculture Organization Statistics	NRL	Nutrition Research Laboratories
FCT	Food Composition Tables	NVIF	Nutritive Value of Indian Foods
G.	Gujarati	O.	Oriya
GC	Gas Chromatography	P.	Punjabi
GLVs	Green Leafy Vegetables	PEM	Protein Energy Malnutrition
H.	Hindi	PER	Protein Efficiency Ratio
HPLC	High Performance Liquid Chromatography	PSUs	Primary Sampling Units
ICMR	Indian Council of Medical Research	PUFA	Polyunsaturated Fatty Acids
ICP-MS	Inductively Coupled Plasma Mass Spectrometry	QAQC	Quality assurance and quality control
IDA	Iron Deficiency Anemia	RAE	Retinol equivalent activity
IDF	Insoluble Dietary Fibre	RID	Refractive index detector
IDF	International Diabetes Federation	RP	Reversed phase
IFCT	Indian Food Composition Tables	S.	Sanskrit
INFOODS	International Network of Food Data Systems	Sci.	Scientific name
IP-6	Inositol hexaphosphate	SD	Standard deviation
IRFA	Indian Research Fund Association	SDF	Soluble Dietary Fibre
IU	International Units	SSUs	Secondary sampling units
Kan.	Kannada	Tam.	Tamil
Kash.	Kashmiri	TDF	Total Dietary Fibre
		Tel.	Telugu
		U.	Urdu
		UT	Union territory
		UV	Ultra violet
		VAD	Vitamin A deficiency
		WHO	World Health Organization

Table 1

PROXIMATE PRINCIPLES

AND

DIETARY FIBRE



Table 1. PROXIMATE PRINCIPLES AND DIETARY FIBRE

(All values are expressed per 100g edible portion; All blank space in the table represent below detectable limit)

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy	
									Total	Insoluble	Soluble	
			← g →									KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC	
A	CEREALS AND MILLETS											
A001	Amaranth seed, black (<i>Amaranthus cruentus</i>)	1	9.89	14.59	2.78	5.74	7.02	5.76	1.26	59.98	1490	
A002	Amaranth seed, pale brown (<i>Amaranthus cruentus</i>)	6	9.20±0.40	13.27±0.34	3.05±0.30	5.56±0.33	7.47±0.09	5.80±0.17	1.67±0.21	61.46±0.60	1489±10	
A003	Bajra (<i>Pennisetum typhoideum</i>)	6	8.97±0.60	10.96±0.26	1.37±0.17	5.43±0.64	11.49±0.62	9.14±0.58	2.34±0.42	61.78±0.85	1456±18	
A004	Barley (<i>Hordeum vulgare</i>)	6	9.77±0.38	10.94±0.51	1.06±0.22	1.30±0.20	15.64±0.64	9.98±0.62	5.66±0.68	61.29±0.77	1321±19	
A005	Jowar (<i>Sorghum vulgare</i>)	6	9.01±0.77	9.97±0.43	1.39±0.34	1.73±0.31	10.22±0.49	8.49±0.40	1.73±0.40	67.68±1.03	1398±13	
A006	Maize, dry (<i>Zea mays</i>)	6	9.26±0.55	8.80±0.49	1.17±0.16	3.77±0.48	12.24±0.93	11.29±0.85	0.94±0.18	64.77±1.58	1398±25	
A007	Maize, tender, local (<i>Zea mays</i>)	6	68.29±0.52	3.57±0.42	0.38±0.04	1.40±0.30	3.67±0.26	3.23±0.23	0.43±0.07	22.69±0.94	502±7	
A008	Maize, tender, sweet (<i>Zea mays</i>)	4	74.40±0.71	4.16±0.41	0.36±0.06	1.35±0.07	3.30±0.51	2.71±0.53	0.59±0.11	16.42±0.89	405±14	
A009	Quinoa (<i>Chenopodium quinoa</i>)	1	10.43	13.11	2.65	5.50	14.66	10.21	4.46	53.65	1374	
A010	Ragi (<i>Eleusine coracana</i>)	5	10.89±0.61	7.16±0.63	2.04±0.34	1.92±0.14	11.18±1.14	9.51±0.65	1.67±0.55	66.82±0.73	1342±10	
A011	Rice flakes (<i>Oryza sativa</i>)	6	10.36±0.53	7.44±0.35	0.85±0.13	1.14±0.11	3.46±0.32	2.65±0.34	0.81±0.12	76.75±0.96	1480±16	
A012	Rice puffed (<i>Oryza sativa</i>)	6	9.40±0.22	7.47±0.15	1.28±0.10	1.62±0.13	2.56±0.33	1.76±0.13	0.80±0.38	77.68±0.54	1514±4	
A013	Rice, raw, brown (<i>Oryza sativa</i>)	6	9.33±0.39	9.16±0.75	1.04±0.18	1.24±0.08	4.43±0.54	3.60±0.55	0.82±0.15	74.80±0.85	1480±10	
A014	Rice, parboiled, milled (<i>Oryza sativa</i>)	6	10.09±0.43	7.81±0.63	0.65±0.08	0.55±0.08	3.74±0.36	2.98±0.35	0.76±0.09	77.16±0.76	1471±8	
A015	Rice, raw, milled (<i>Oryza sativa</i>)	6	9.93±0.75	7.94±0.58	0.56±0.08	0.52±0.05	2.81±0.42	1.99±0.39	0.82±0.22	78.24±1.07	1491±15	
A016	Samai (<i>Panicum miliare</i>)	6	11.36±0.19	10.13±0.45	1.34±0.16	3.89±0.35	7.72±0.92	5.45±0.48	2.27±0.52	65.55±1.29	1449±19	
A017	Varagu (<i>Paspalum scrobiculatum</i>)	5	14.23±0.45	8.92±1.09	1.72±0.27	2.55±0.13	6.39±0.60	4.29±0.82	2.11±0.34	66.19±1.19	1388±10	
A018	Wheat flour, refined (<i>Triticum aestivum</i>)	6	11.34±0.93	10.36±0.29	0.51±0.07	0.76±0.07	2.76±0.29	2.14±0.30	0.62±0.14	74.27±0.92	1472±16	

Table 1. Proximate Principles and Dietary Fibre

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
A019	Wheat flour, atta (<i>Triticum aestivum</i>)	6	11.10±0.35	10.57±0.37	1.28±0.19	1.53±0.12	11.36±0.29	9.73±0.47	1.63±0.64	64.17±0.32	1340±7
A020	Wheat, whole (<i>Triticum aestivum</i>)	6	10.58±1.11	10.59±0.60	1.42±0.19	1.47±0.05	11.23±0.77	9.63±0.19	1.60±0.75	64.72±1.74	1347±23
A021	Wheat, bulgur (<i>Triticum aestivum</i>)	6	8.61±0.32	10.84±0.75	1.23±0.06	1.45±0.02	8.81±0.45	6.56±0.20	2.25±0.38	69.06±0.74	1430±6
A022	Wheat, semolina (<i>Triticum aestivum</i>)	6	8.94±0.68	11.38±0.37	0.80±0.17	0.74±0.10	9.72±0.74	8.16±0.58	1.55±0.18	68.43±0.99	1396±18
A023	Wheat, vermicelli (<i>Triticum aestivum</i>)	6	9.59±0.37	9.70±0.52	0.60±0.04	0.45±0.03	9.28±0.69	7.53±0.51	1.75±0.24	70.39±0.61	1392±8
A024	Wheat, vermicelli, roasted (<i>Triticum aestivum</i>)	6	7.61±0.47	10.37±0.70	0.56±0.04	0.49±0.05	9.55±0.40	7.79±0.29	1.76±0.18	71.42±0.71	1423±13

B GRAIN LEGUMES

B001	Bengal gram, dal (<i>Cicer arietinum</i>)	6	9.18±0.58	21.55±1.45	2.10±0.10	5.31±0.06	15.15±0.17	12.67±0.22	2.48±0.15	46.72±1.29	1377±10
B002	Bengal gram, whole (<i>Cicer arietinum</i>)	6	8.56±0.37	18.77±0.42	2.78±0.13	5.11±0.11	25.22±0.39	22.70±0.60	2.52±0.87	39.56±0.16	1201±9
B003	Black gram, dal (<i>Phaseolus mungo</i>)	6	9.16±0.35	23.06±0.59	3.17±0.02	1.69±0.12	11.93±0.26	7.58±0.13	4.35±0.15	51.00±0.80	1356±9
B004	Black gram, whole (<i>Phaseolus mungo</i>)	6	8.70±0.33	21.97±0.63	3.35±0.03	1.58±0.06	20.41±0.06	15.47±0.05	4.94±0.07	43.99±0.76	1219±5
B005	Cowpea, brown (<i>Vigna catjang</i>)	6	9.42±0.39	20.36±0.59	2.90±0.11	1.15±0.06	11.54±0.13	8.75±0.09	2.80±0.05	54.62±0.49	1340±7
B006	Cowpea, white (<i>Vigna catjang</i>)	1	9.32	21.25	2.83	1.14	11.70	8.91	2.79	53.77	1340
B007	Field bean, black (<i>Phaseolus vulgaris</i>)	1	9.57	19.93	2.73	0.92	23.40	17.99	5.41	43.46	1155
B008	Field bean, brown (<i>Phaseolus vulgaris</i>)	1	8.74	19.90	2.74	0.98	22.40	17.32	5.08	45.24	1184
B009	Field bean, white (<i>Phaseolus vulgaris</i>)	5	8.61±0.36	19.84±1.04	3.09±0.15	0.94±0.02	22.99±0.83	17.45±2.27	5.54±2.28	44.53±1.42	1173±24
B010	Green gram, dal (<i>Vigna radiata</i>)	6	9.77±0.67	23.88±0.61	3.04±0.03	1.35±0.20	9.37±0.38	7.75±0.39	1.62±0.19	52.59±0.45	1363±10
B011	Green gram, whole (<i>Vigna radiata</i>)	6	9.95±0.42	22.53±0.43	3.22±0.04	1.14±0.17	17.04±0.38	14.59±0.42	2.44±0.15	46.13±0.64	1229±10
B012	Horse gram, whole (<i>Dolichos biflorus</i>)	6	9.28±0.57	21.73±0.29	3.24±0.11	0.62±0.04	7.88±0.02	6.22±0.03	1.66±0.03	57.24±0.50	1379±9
B013	Lentil dal (<i>Lens culinaris</i>)	6	9.71±0.48	24.35±1.10	2.23±0.13	0.75±0.04	10.43±0.39	8.60±0.42	1.83±0.23	52.53±1.05	1349±11

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
B014	Lentil whole, brown (<i>Lens culinaris</i>)	6	9.20±0.77	22.49±0.58	2.39±0.35	0.64±0.02	16.82±1.30	14.16±1.33	2.66±0.42	48.47±1.12	1251±23
B015	Lentil whole, yellowish (<i>Lens culinaris</i>)	2	9.75	22.87	2.20	0.61	16.66	14.15	2.51	47.91	1246
B016	Moth bean (<i>Vigna aconitifolia</i>)	6	8.14±0.49	19.75±0.38	3.14±0.18	1.76±0.09	15.12±0.49	14.50±0.44	0.62±0.10	52.09±0.96	1291±16
B017	Peas, dry (<i>Pisum sativum</i>)	6	9.33±0.61	20.43±0.79	2.41±0.09	1.89±0.08	17.01±0.63	14.55±0.73	2.47±0.17	48.93±0.45	1269±13
B018	Rajmah, black (<i>Phaseolus vulgaris</i>)	2	8.69	19.01	3.35	1.62	17.74	15.16	2.58	49.59	1247
B019	Rajmah, brown (<i>Phaseolus vulgaris</i>)	6	9.68±0.79	19.50±0.84	3.36±0.19	1.68±0.07	16.95±0.27	14.33±0.19	2.62±0.16	48.83±0.59	1245±12
B020	Rajmah, red (<i>Phaseolus vulgaris</i>)	3	9.87±0.30	19.91±1.44	3.28±0.21	1.77±0.04	16.57±0.63	13.86±0.43	2.70±0.20	48.61±0.65	1252±14
B021	Red gram, dal (<i>Cajanus cajan</i>)	6	9.20±0.61	21.70±0.50	3.26±0.03	1.56±0.03	9.06±0.30	6.67±0.23	2.39±0.15	55.23±0.83	1384±10
B022	Red gram, whole (<i>Cajanus cajan</i>)	6	9.30±0.45	20.47±0.72	3.53±0.03	1.38±0.08	22.84±0.43	19.69±0.30	3.15±0.34	42.48±0.77	1146±10
B023	Ricebean (<i>Vigna umbellata</i>)	1	11.12	19.97	3.54	0.74	13.37	10.04	3.33	51.26	1265
B024	Soybean, brown (<i>Glycine max</i>)	6	5.51±0.13	35.58±0.66	4.74±0.31	19.82±0.26	21.55±0.66	16.56±0.30	5.00±0.52	12.79±0.97	1596±11
B025	Soybean, white (<i>Glycine max</i>)	1	5.47	37.80	4.52	19.42	22.63	17.04	5.59	10.16	1579
C GREEN LEAFY VEGETABLES											
C001	Agathi leaves (<i>Sesbania grandiflora</i>)	1	74.43	8.01	2.42	1.35	8.60	6.00	2.60	5.21	295
C002	Amaranth leaves, green (<i>Amaranthus gangeticus</i>)	6	86.85±1.21	3.29±0.57	2.52±0.32	0.65±0.07	4.41±0.10	3.21±0.08	1.20±0.09	2.28±0.62	128±17
C003	Amaranth leaves, red (<i>Amaranthus gangeticus</i>)	1	85.56	3.93	2.61	0.63	4.91	3.72	1.19	2.37	140
C004	Amaranth leaves, red and green mix (<i>Amaranthus gangeticus</i>)	4	86.37±0.38	3.09±0.14	2.55±0.20	0.53±0.03	4.60±0.36	3.23±0.18	1.37±0.23	2.87±0.35	132±6
C005	Amaranth spined, leaves, green (<i>Amaranthus spinosus</i>)	4	86.46±0.48	3.54±0.31	2.94±0.21	0.36±0.02	5.10±0.32	3.89±0.26	1.20±0.07	1.61±0.40	110±5
C006	Amaranth spined, leaves, red and green mix (<i>Amaranthus spinosus</i>)	2	86.64	2.80	3.20	0.34	5.57	3.82	1.75	1.45	99
C007	Basella leaves (<i>Basella alba</i>)	2	92.68	1.57	1.09	0.45	2.21	1.64	0.57	2.01	82

Table 1. Proximate Principles and Dietary Fibre

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
C008	Bathua leaves (<i>Chenopodium album</i>)	2	88.77	2.50	1.71	0.44	4.01	2.34	1.68	2.56	116
C009	Beet greens (<i>Beta vulgaris</i>)	6	86.68±1.35	2.38±0.35	2.69±0.12	0.75±0.02	3.64±0.13	2.20±0.10	1.43±0.19	3.86±1.63	145±23
C010	Betel leaves, big (Kolkata) (<i>Piper betle</i>)	6	84.93±0.85	2.51±0.29	2.33±0.18	0.75±0.02	2.12±0.20	1.32±0.16	0.80±0.05	7.37±0.66	202±14
C011	Betel leaves, small (<i>Piper betle</i>)	4	85.92±0.16	2.62±0.28	2.59±0.18	0.75±0.04	1.97±0.13	1.17±0.12	0.80±0.09	6.16±0.33	183±2
C012	Brussels sprouts (<i>Brassica oleracea</i> var. <i>gemmifera</i>)	1	84.39	4.26	1.47	0.50	4.29	3.35	0.94	5.09	185
C013	Cabbage, Chinese (<i>Brassica rupa</i>)	1	93.19	1.58	0.73	0.13	2.01	1.55	0.45	2.36	75
C014	Cabbage, collard greens (<i>Brassica oleracea</i> var. <i>viridis</i>)	1	89.53	3.63	0.81	0.27	2.98	2.04	0.94	2.79	126
C015	Cabbage, green (<i>Brassica oleracea</i> var. <i>capitata</i> f. <i>alba</i>)	6	91.85±0.97	1.36±0.07	0.67±0.06	0.12±0.01	2.76±0.20	1.91±0.18	0.85±0.10	3.25±0.91	90±16
C016	Cabbage, violet (<i>Brassica oleracea</i> var. <i>capitata</i> f. <i>rubra</i>)	2	91.94	1.39	0.71	0.21	2.21	1.58	0.62	3.54	97
C017	Cauliflower leaves (<i>Brassica oleracea</i> var. <i>botrytis</i>)	6	87.64±1.33	3.90±0.87	1.22±0.12	0.42±0.02	3.43±0.26	2.37±0.20	1.06±0.12	3.39±0.70	148±22
C018	Colocasia leaves, green (<i>Colocasia esculenta</i>)	6	83.61±0.59	3.42±0.29	2.30±0.11	1.38±0.09	5.60±0.18	4.32±0.16	1.29±0.12	3.69±0.49	182±9
C019	Drumstick leaves (<i>Moringa oleifera</i>)	3	75.65±1.49	6.41±0.35	2.46±0.18	1.64±0.12	8.21±0.19	6.12±0.08	2.10±0.11	5.62±1.44	282±27
C020	Fenugreek leaves (<i>Trigonella foenum graecum</i>)	5	86.73±0.66	3.68±0.36	1.69±0.19	0.83±0.02	4.90±0.21	3.20±0.14	1.70±0.09	2.17±0.32	144±10
C021	Garden cress (<i>Lepidium sativum</i>)	2	84.02	5.62	2.48	0.80	2.60	1.77	0.83	4.48	208
C022	Gogu leaves, green (<i>Hibiscus cannabinus</i>)	4	87.42±0.41	1.86±0.16	0.98±0.08	1.09±0.03	4.59±0.39	3.24±0.16	1.35±0.30	4.06±0.18	152±2
C023	Gogu leaves, red (<i>Hibiscus cannabinus</i>)	1	87.98	1.85	0.97	1.07	3.89	2.66	1.23	4.24	153
C024	Knol-Khol, leaves (<i>Brassica oleracea</i> var. <i>gongylodes</i>)	2	86.20	3.12	1.42	0.35	2.76	1.81	0.95	6.16	178
C025	Lettuce (<i>Lactuca sativa</i>)	3	92.27±0.99	1.54±0.48	1.11±0.10	0.27±0.03	1.79±0.24	1.32±0.04	0.47±0.28	3.01±0.56	91±16
C026	Mustard leaves (<i>Brassica juncea</i>)	3	88.17±0.56	3.52±0.21	1.47±0.11	0.51±0.05	3.92±0.06	3.04±0.03	0.87±0.03	2.41±0.46	127±9
C027	Pak Choi leaves (<i>Brassica rapa</i> var. <i>chinensis</i>)	1	93.56	1.41	1.10	0.25	1.91	1.44	0.47	1.78	67

Table 1. Proximate Principles and Dietary Fibre

7

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
C028	Parsley (<i>Petroselinum crispum</i>)	3	77.76±1.13	5.55±0.34	2.25±0.17	1.14±0.12	3.87±0.05	2.79±0.15	1.09±0.12	9.43±1.05	305±18
C029	Ponnaganni (<i>Alternanthera sessilis</i>)	2	79.43	5.29	2.65	0.71	6.74	5.63	1.11	5.17	213
C030	Pumpkin leaves, tender (<i>Cucurbita maxima</i>)	6	85.82±1.04	4.21±0.41	2.24±0.21	0.74±0.03	2.25±0.09	1.56±0.13	0.69±0.15	4.75±1.01	185±15
C031	Radish leaves (<i>Raphanus sativus</i>)	6	91.19±0.53	2.22±0.23	1.50±0.21	0.51±0.05	1.82±0.10	1.18±0.04	0.63±0.07	2.77±0.34	109±9
C032	Rumex leaves (<i>Rumex patientia</i>)	2	93.18	1.62	1.27	0.33	1.27	0.93	0.34	2.33	82
C033	Spinach (<i>Spinacia oleracea</i>)	6	90.31±0.46	2.14±0.14	2.47±0.38	0.64±0.03	2.38±0.24	1.52±0.15	0.86±0.10	2.05±0.31	102±7
C034	Tamarind leaves, tender (<i>Tamarindus indica</i>)	3	71.69±0.06	5.84±0.01	1.25±0.06	0.49±0.01	10.70±0.02	9.34±0.01	1.36±0.01	10.04±0.02	299±1
D OTHER VEGETABLES											
D001	Ash gourd (<i>Benincasa hispida</i>)	6	92.17±0.42	0.79±0.06	0.70±0.08	0.14±0.02	3.37±0.23	2.52±0.26	0.85±0.13	2.84±0.22	73±4
D002	Bamboo shoot, tender (<i>Bambusa vulgaris</i>)	1	94.56	1.33	0.55	0.35	1.55	1.06	0.49	1.67	68
D003	Bean scarlet, tender (<i>Phaseolus coccineus</i>)	5	85.51±1.22	2.86±0.27	0.96±0.03	0.99±0.06	4.50±0.25	3.76±0.14	0.74±0.21	5.16±1.45	179±21
D004	Bitter gourd, jagged, teeth ridges, elongate (<i>Momordica charantia</i>)	6	90.87±0.56	1.44±0.17	0.86±0.04	0.24±0.01	3.78±0.16	3.10±0.19	0.68±0.07	2.82±0.32	87±6
D005	Bitter gourd, jagged, teeth ridges, short (<i>Momordica charantia</i>)	4	91.60±0.28	1.34±0.12	0.81±0.06	0.24±0.01	3.49±0.18	2.96±0.10	0.53±0.11	2.53±0.26	79±5
D006	Bitter gourd, jagged, smooth ridges, elongate (<i>Momordica charantia</i>)	1	91.24	1.61	0.88	0.26	3.72	3.05	0.67	2.29	81
D007	Bottle gourd, elongate, pale green (<i>Lagenaria vulgaris</i>)	6	95.17±0.33	0.53±0.05	0.36±0.04	0.13±0.02	2.12±0.07	1.65±0.05	0.48±0.07	1.68±0.33	46±6
D008	Bottle gourd, round, pale green (<i>Lagenaria vulgaris</i>)	5	94.50±0.32	0.42±0.02	0.34±0.05	0.12±0.03	2.10±0.15	1.72±0.13	0.38±0.05	2.53±0.21	57±3
D009	Bottle gourd, elongate, dark green (<i>Lagenaria vulgaris</i>)	1	94.63	0.49	0.40	0.13	2.11	1.74	0.37	2.25	54
D010	Brinjal-1 (<i>Solanum melongena</i>)	1	89.95	1.77	0.83	0.39	3.57	2.37	1.20	3.49	114
D011	Brinjal-2 (<i>Solanum melongena</i>)	1	90.28	1.82	0.85	0.34	4.01	2.80	1.21	2.71	99
D012	Brinjal-3 (<i>Solanum melongena</i>)	2	90.02	1.36	0.65	0.33	4.26	3.10	1.16	3.38	102

Table 1. Proximate Principles and Dietary Fibre

Food code		No. of Regions	Dietary Fibre								Energy
			Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	
							Total	Insoluble	Soluble		
Food Name			←						→	KJ	
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
D013	Brinjal-4 (<i>Solanum melongena</i>)	4	90.28±1.21	1.51±0.24	0.66±0.09	0.31±0.03	4.04±0.65	3.03±0.82	1.01±0.22	3.19±0.86	100±18
D014	Brinjal-5 (<i>Solanum melongena</i>)	3	89.45±1.73	1.38±0.26	0.72±0.12	0.29±0.06	4.13±0.70	3.10±0.43	1.03±0.29	4.02±2.01	111±38
D015	Brinjal-6 (<i>Solanum melongena</i>)	2	90.57	1.46	0.72	0.29	3.68	3.00	0.68	3.29	97
D016	Brinjal-7 (<i>Solanum melongena</i>)	1	91.25	1.40	0.57	0.37	3.32	2.30	1.02	3.10	98
D017	Brinjal-8 (<i>Solanum melongena</i>)	3	89.28±0.39	1.82±0.16	0.80±0.06	0.33±0.04	4.01±0.48	2.79±0.53	1.22±0.05	3.75±0.18	117±2
D018	Brinjal-9 (<i>Solanum melongena</i>)	2	89.83	1.47	0.83	0.35	4.00	2.94	1.06	3.52	106
D019	Brinjal-10 (<i>Solanum melongena</i>)	2	89.38	1.68	0.72	0.25	3.90	2.74	1.16	4.08	116
D020	Brinjal-11 (<i>Solanum melongena</i>)	6	89.93±0.89	1.43±0.20	0.68±0.07	0.31±0.04	4.12±0.52	2.95±0.56	1.17±0.28	3.53±1.01	105±16
D021	Brinjal-12 (<i>Solanum melongena</i>)	2	90.94	1.60	0.58	0.27	3.89	2.70	1.19	2.73	93
D022	Brinjal-13 (<i>Solanum melongena</i>)	1	89.14	1.49	0.64	0.31	3.92	2.61	1.31	4.51	124
D023	Brinjal-14 (<i>Solanum melongena</i>)	3	90.31±0.06	1.56±0.19	0.64±0.02	0.35±0.03	3.87±0.59	2.50±0.76	1.37±0.24	3.27±0.39	106±9
D024	Brinjal-15 (<i>Solanum melongena</i>)	4	89.44±0.96	1.58±0.27	0.74±0.05	0.29±0.03	3.99±0.79	2.82±0.72	1.17±0.08	3.96±0.19	114±6
D025	Brinjal-16 (<i>Solanum melongena</i>)	1	90.39	1.26	0.80	0.34	3.74	2.54	1.20	3.46	103
D026	Brinjal-17 (<i>Solanum melongena</i>)	1	90.83	1.18	0.81	0.36	3.94	2.80	1.14	2.89	91
D027	Brinjal-18 (<i>Solanum melongena</i>)	3	89.07±0.78	1.47±0.15	0.66±0.02	0.33±0.06	4.38±0.25	3.23±0.15	1.15±0.18	4.10±0.88	116±18
D028	Brinjal-19 (<i>Solanum melongena</i>)	3	90.41±0.55	1.26±0.06	0.66±0.05	0.31±0.02	3.97±0.18	2.81±0.42	1.16±0.27	3.39±0.36	100±8
D029	Brinjal-20 (<i>Solanum melongena</i>)	5	90.65±0.79	1.44±0.21	0.68±0.08	0.34±0.04	3.73±0.21	2.64±0.37	1.09±0.21	3.15±0.60	99±10
D030	Brinjal-21 (<i>Solanum melongena</i>)	2	89.83	1.36	0.69	0.35	3.85	2.57	1.28	3.93	113
D031	Brinjal - all varieties (<i>Solanum melongena</i>)	6	90.00±0.90	1.48±0.22	0.70±0.08	0.32±0.04	3.98±0.49	2.84±0.52	1.14±0.22	3.52±0.80	106±15
D032	Broad beans (<i>Vicia faba</i>)	3	84.20±0.17	3.85±0.13	1.05±0.08	0.15±0.01	8.63±0.15	6.61±0.07	2.03±0.09	2.11±0.11	123±4
D033	Capsicum, green (<i>Capsicum annuum</i>)	6	93.89±0.32	1.11±0.11	0.76±0.05	0.34±0.03	2.06±0.22	1.33±0.29	0.73±0.17	1.84±0.32	68±6

Table 1. Proximate Principles and Dietary Fibre

Food code		No. of Regions	Dietary Fibre								Carbohydrate	Energy
			Moisture	Protein	Ash	Total Fat						
							Total	Insoluble	Soluble			
										g		
Food Name			←						→	KJ		
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC	
D034	Capsicum, red (<i>Capsicum annuum</i>)	4	92.99±1.47	1.47±0.36	0.76±0.14	0.47±0.05	2.19±0.17	1.58±0.13	0.61±0.10	2.14±1.06	83±24	
D035	Capsicum, yellow (<i>Capsicum annuum</i>)	4	93.38±0.37	1.35±0.35	0.72±0.12	0.41±0.02	2.19±0.13	1.39±0.06	0.80±0.08	1.95±0.16	78±4	
D036	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)	6	90.76±0.22	2.15±0.15	0.91±0.10	0.44±0.03	3.71±0.27	2.66±0.35	1.04±0.13	2.03±0.22	96±5	
D037	Celery stalk (<i>Apium graveolens</i>)	3	92.87±0.22	0.98±0.13	1.50±0.16	0.24±0.04	2.09±0.13	1.57±0.07	0.52±0.16	2.33±0.23	69±6	
D038	Cho-cho-marrow (<i>Sechium edule</i>)	4	93.78±0.12	0.66±0.07	0.38±0.05	0.15±0.03	1.55±0.08	1.19±0.07	0.36±0.12	3.47±0.18	79±2	
D039	Cluster beans (<i>Cyamopsis tetragonoloba</i>)	6	84.65±0.83	3.55±0.38	1.68±0.13	0.37±0.04	4.83±0.54	3.55±0.48	1.28±0.13	4.91±0.58	168±15	
D040	Colocasia, stem, black (<i>Colocasia esculenta</i>)	1	91.15	0.76	0.91	0.34	3.01	1.81	1.20	3.83	100	
D041	Colocasia, stem, green (<i>Colocasia esculenta</i>)	5	92.67±0.44	0.91±0.16	1.00±0.04	0.22±0.02	2.33±0.06	1.26±0.03	1.07±0.08	2.86±0.34	81±7	
D042	Corn, baby (<i>Zea mays</i>)	6	75.44±0.98	2.69±0.23	2.79±0.21	1.33±0.23	6.09±0.50	4.47±0.28	1.62±0.31	11.66±1.40	306±18	
D043	Cucumber, green, elongate (<i>Cucumis sativus</i>)	6	92.96±0.48	0.71±0.11	0.54±0.06	0.16±0.04	2.14±0.42	1.52±0.43	0.63±0.11	3.48±0.47	82±8	
D044	Cucumber, green, short (<i>Cucumis sativus</i>)	6	93.52±0.44	0.83±0.14	0.52±0.06	0.18±0.05	2.13±0.26	1.60±0.33	0.54±0.16	2.82±0.49	73±10	
D045	Cucumber, orange, round (<i>Cucumis sativus</i>)	2	92.80	0.98	0.52	0.24	2.46	1.78	0.68	3.01	82	
D046	Drumstick (<i>Moringa oleifera</i>)	6	85.39±0.64	2.62±0.22	1.27±0.09	0.12±0.01	6.83±0.22	5.60±0.26	1.23±0.06	3.76±0.70	123±11	
D047	Field beans, tender, broad (<i>Vicia faba</i>)	2	86.98	3.06	0.94	0.64	5.64	4.84	0.79	2.75	129	
D048	Field beans, tender, lean (<i>Vicia faba</i>)	6	85.57±1.32	3.71±0.62	1.08±0.12	0.60±0.02	6.19±0.47	5.37±0.43	0.82±0.10	2.85±0.32	140±15	
D049	French beans, country (<i>Phaseolus vulgaris</i>)	5	89.14±0.52	2.49±0.21	1.05±0.09	0.26±0.02	4.38±0.41	3.88±0.36	0.51±0.11	2.68±0.48	102±8	
D050	French beans, hybrid (<i>Phaseolus vulgaris</i>)	2	90.11	2.12	0.77	0.19	4.18	3.53	0.65	2.63	93	
D051	Jack fruit, raw (<i>Artocarpus heterophyllus</i>)	5	85.52±0.86	1.98±0.16	0.99±0.10	0.35±0.05	7.69±0.20	7.15±0.13	0.54±0.08	3.48±0.70	110±15	
D052	Jack fruit, seed, mature (<i>Artocarpus heterophyllus</i>)	5	72.32±0.53	5.79±0.14	1.02±0.16	0.44±0.03	8.63±0.09	7.83±0.05	0.80±0.07	11.81±0.74	322±10	
D053	Knol - Khol (<i>Brassica oleracea</i>)	6	93.14±0.27	1.58±0.24	0.79±0.03	0.35±0.02	2.75±0.07	2.31±0.09	0.44±0.04	1.39±0.21	67±4	
D054	Kovai, big (<i>Coccinia cordifolia</i>)	6	92.78±0.61	1.39±0.23	0.58±0.07	0.24±0.08	3.00±0.26	2.19±0.18	0.81±0.11	2.01±0.54	73±10	

Table 1. Proximate Principles and Dietary Fibre

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
D055	Kovai, small (<i>Coccinia cordifolia</i>)	1	92.40	1.22	0.47	0.24	3.25	2.14	1.12	2.41	80
D056	Ladies finger (<i>Abelmoschus esculentus</i>)	6	89.06±0.30	2.08±0.37	0.94±0.09	0.22±0.03	4.08±0.20	2.80±0.29	1.28±0.37	3.62±0.26	115±5
D057	Mango, green, raw (<i>Mangifera indica</i>)	6	85.14±0.59	0.69±0.09	0.49±0.07	0.08±0.02	3.01±0.19	1.68±0.15	1.34±0.11	10.59±0.37	205±8
D058	Onion, stalk (<i>Allium cepa</i>)	6	88.35±0.79	2.07±0.21	1.13±0.09	0.26±0.02	5.21±0.10	3.76±0.07	1.45±0.11	2.99±0.50	107±12
D059	Papaya, raw (<i>Carica papaya</i>)	6	92.04±0.44	0.50±0.06	0.55±0.06	0.23±0.03	2.28±0.05	1.32±0.04	0.96±0.04	4.40±0.37	100±7
D060	Parwar (<i>Trichosanthes dioica</i>)	6	91.57±0.27	1.40±0.10	0.59±0.03	0.30±0.04	2.61±0.07	1.82±0.04	0.79±0.07	3.54±0.32	101±5
D061	Peas, fresh (<i>Pisum sativum</i>)	6	73.37±1.13	7.25±1.03	1.05±0.02	0.13±0.02	6.32±0.26	5.04±0.19	1.28±0.25	11.88±0.67	340±19
D062	Plantain, flower (<i>Musa x paradisiaca</i>)	6	89.14±0.59	1.47±0.12	1.35±0.09	0.63±0.02	5.25±0.39	4.72±0.44	0.52±0.07	2.15±0.43	89±8
D063	Plantain, green (<i>Musa x paradisiaca</i>)	6	76.15±0.42	1.18±0.20	1.27±0.14	0.23±0.02	3.60±0.27	2.79±0.25	0.81±0.05	17.58±0.35	334±6
D064	Plantain, stem (<i>Musa x paradisiaca</i>)	6	87.53±1.01	0.35±0.05	1.20±0.09	0.16±0.07	2.12±0.04	1.33±0.08	0.80±0.09	8.64±0.93	165±15
D065	Pumpkin, green, cylindrical (<i>Cucurbita maxima</i>)	1	91.73	0.87	0.47	0.18	2.53	1.28	1.25	4.22	103
D066	Pumpkin, orange, round (<i>Cucurbita maxima</i>)	6	91.85±0.45	0.84±0.21	0.58±0.09	0.16±0.02	2.56±0.11	1.44±0.14	1.12±0.08	4.00±0.64	97±8
D067	Red gram, tender, fresh (<i>Cajanus cajan</i>)	1	64.00	8.09	1.64	0.92	5.90	3.66	2.25	19.46	520
D068	Ridge gourd (<i>Luffa acutangula</i>)	6	94.99±0.35	0.91±0.08	0.44±0.04	0.14±0.01	1.81±0.13	1.20±0.16	0.61±0.08	1.72±0.46	55±7
D069	Ridge gourd, smooth skin (<i>Luffa acutangula</i>)	3	94.27±0.62	0.98±0.13	0.53±0.06	0.13±0.01	1.85±0.28	1.31±0.17	0.54±0.10	2.24±0.41	64±8
D070	Snake gourd, long, pale green (<i>Trichosanthes anguina</i>)	6	94.81±0.28	0.98±0.33	0.42±0.04	0.25±0.01	2.27±0.11	1.69±0.15	0.58±0.08	1.27±0.15	52±4
D071	Snake gourd, long, dark green (<i>Trichosanthes anguina</i>)	2	94.92	0.89	0.45	0.25	2.27	1.69	0.58	1.23	50
D072	Snake gourd, short (<i>Trichosanthes anguina</i>)	1	94.35	0.54	0.41	0.26	2.29	1.61	0.68	2.15	61
D073	Tinda, tender (<i>Praecitrullus fistulosus</i>)	6	94.41±0.32	1.02±0.04	0.51±0.03	0.17±0.01	2.00±0.04	1.68±0.06	0.32±0.02	1.90±0.37	58±6
D074	Tomato, green (<i>Solanum lycopersicum</i>)	6	93.21±0.46	1.12±0.08	0.60±0.03	0.27±0.07	1.62±0.08	1.05±0.07	0.57±0.04	3.18±0.40	87±7
D075	Tomato, ripe, hybrid ((<i>Solanum lycopersicum</i>)	6	93.79±0.42	0.76±0.03	0.43±0.05	0.25±0.02	1.58±0.16	1.27±0.13	0.30±0.05	3.20±0.30	79±6

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
D076	Tomato, ripe, local (<i>Lycopersicon esculentum</i>)	6	93.62±0.16	0.90±0.07	0.52±0.08	0.47±0.06	1.77±0.04	1.44±0.11	0.33±0.08	2.71±0.08	82±4
D077	Zucchini, green (<i>Cucurbita pepo</i>)	2	92.83	1.10	0.93	0.51	2.30	1.42	0.88	2.33	84
D078	Zucchini, yellow (<i>Cucurbita pepo</i>)	2	93.15	1.31	1.08	0.44	1.84	1.37	0.47	2.20	79
E FRUITS											
E001	Apple, big (<i>Malus domestica</i>)	6	83.01±0.94	0.29±0.08	0.36±0.04	0.64±0.04	2.59±0.15	1.43±0.15	1.16±0.17	13.11±0.76	261±15
E002	Apple, green (<i>Malus domestica</i>)	6	85.54±0.98	0.46±0.07	0.31±0.04	0.50±0.05	2.54±0.26	1.72±0.22	0.81±0.16	10.65±1.00	214±17
E003	Apple, small (<i>Malus domestica</i>)	6	82.91±0.48	0.31±0.05	0.24±0.02	0.53±0.05	2.06±0.12	1.44±0.09	0.62±0.12	13.95±0.48	267±8
E004	Apple, small, Kashmir (<i>Malus domestica</i>)	1	82.79	0.27	0.29	0.60	2.07	1.53	0.54	13.99	269
E005	Apricot, dried (<i>Prunus armeniaca</i>)	6	16.68±0.59	3.17±0.29	3.46±0.14	0.74±0.03	3.32±0.13	2.72±0.11	0.60±0.04	72.63±0.27	1321±9
E006	Apricot, processed (<i>Prunus armeniaca</i>)	3	85.75±0.81	1.47±0.15	0.61±0.16	0.64±0.04	0.59±0.01	0.43±0.02	0.16±0.02	10.93±0.88	236±12
E007	Avocado fruit (<i>Persea</i> sp.)	1	73.56	2.95	1.19	13.86	6.69	5.26	1.42	1.75	604
E008	Bael fruit (<i>Aegle marmelos</i>)	1	61.36	2.63	0.91	0.57	6.31	3.31	3.00	28.21	569
E009	Banana, ripe, montham (<i>Musa x paradisiaca</i>)	1	70.13	1.25	1.15	0.32	2.21	1.43	0.78	24.95	463
E010	Banana, ripe, poovam (<i>Musa x paradisiaca</i>)	2	71.32	1.49	1.09	0.35	2.33	1.29	1.04	23.41	445
E011	Banana, ripe, red (<i>Musa x paradisiaca</i>)	1	70.28	1.29	0.95	0.29	1.98	1.26	0.72	25.21	467
E012	Banana, ripe, robusta (<i>Musa x paradisiaca</i>)	6	71.93±0.85	1.23±0.08	0.94±0.17	0.33±0.01	1.94±0.07	1.23±0.10	0.71±0.07	23.63±0.74	440±14
E013	Black berry (<i>Rubus</i> sp.)	5	82.94±0.42	0.92±0.10	0.52±0.04	0.63±0.02	4.35±0.25	3.44±0.16	0.91±0.13	10.64±0.35	227±6
E014	Cherries, red (<i>Prunus cerasus</i>)	4	83.61±0.21	1.49±0.22	0.45±0.02	0.46±0.05	2.12±0.12	1.35±0.09	0.78±0.06	11.87±0.31	250±4
E015	Currants, black (<i>Ribes nigrum</i>)	1	83.27	1.51	0.69	0.53	4.07	2.43	1.64	9.93	227
E016	Custard apple (<i>Annona squamosa</i>)	1	71.55	1.62	0.68	0.67	5.10	3.17	1.93	20.38	414

Table 1. Proximate Principles and Dietary Fibre

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
E017	Dates, dry, pale brown (<i>Phoenix dactylifera</i>)	6	11.14±0.73	2.45±0.24	2.20±0.07	0.35±0.03	8.95±0.23	7.53±0.22	1.42±0.14	74.91±0.52	1340±12
E018	Dates, dry, dark brown (<i>Phoenix dactylifera</i>)	2	13.13	2.38	2.39	0.35	9.10	7.57	1.53	72.67	1301
E019	Dates, processed (<i>Phoenix dactylifera</i>)	2	22.01	1.18	1.93	0.41	6.52	5.68	0.84	67.95	1197
E020	Fig (<i>Ficus carica</i>)	6	75.60±1.16	2.03±0.22	1.08±0.18	0.37±0.06	4.64±0.41	2.59±0.22	2.05±0.23	16.28±1.13	341±18
E021	Gooseberry (<i>Emblica officinalis</i>)	5	87.02±0.78	0.34±0.03	0.34±0.05	0.16±0.04	7.75±0.64	6.20±0.43	1.55±0.43	4.39±1.09	99±19
E022	Grapes, seeded, round, black (<i>Vitis vinifera</i>)	4	83.88±0.56	0.76±0.13	0.46±0.04	0.32±0.02	1.35±0.14	0.82±0.06	0.52±0.09	13.23±0.60	254±11
E023	Grapes, seeded, round, green (<i>Vitis vinifera</i>)	5	85.07±0.29	0.77±0.12	0.43±0.05	0.29±0.03	1.25±0.04	0.81±0.02	0.44±0.02	12.19±0.20	235±4
E024	Grapes, seeded, round, red (<i>Vitis vinifera</i>)	5	84.46±1.60	0.95±0.11	0.48±0.07	0.29±0.04	1.28±0.10	0.85±0.05	0.43±0.09	12.55±1.38	244±26
E025	Grapes, seedless, oval, black (<i>Vitis vinifera</i>)	5	75.30±1.55	1.41±0.26	0.70±0.13	0.46±0.11	1.64±0.15	0.98±0.19	0.66±0.08	20.48±0.97	395±23
E026	Grapes, seedless, round, green (<i>Vitis vinifera</i>)	5	85.55±1.70	0.62±0.13	0.47±0.08	0.26±0.02	1.28±0.05	0.85±0.05	0.44±0.01	11.81±1.52	224±28
E027	Grapes, seedless, round, black (<i>Vitis vinifera</i>)	5	76.90±1.40	1.24±0.25	0.49±0.05	0.35±0.03	1.15±0.19	0.83±0.04	0.32±0.15	19.86±1.58	374±25
E028	Guava, white flesh (<i>Psidium guajava</i>)	5	83.79±0.30	1.44±0.09	0.72±0.05	0.32±0.03	8.59±0.05	7.14±0.02	1.45±0.04	5.13±0.33	135±5
E029	Guava, pink flesh (<i>Psidium guajava</i>)	5	81.22±0.45	1.19±0.19	0.81±0.06	0.25±0.03	7.39±0.38	6.12±0.29	1.28±0.09	9.14±0.27	195±6
E030	Jack fruit, ripe (<i>Artocarpus heterophyllus</i>)	5	78.56±0.34	2.74±0.47	0.91±0.08	0.15±0.02	3.62±0.17	2.21±0.17	1.41±0.06	14.01±0.56	302±3
E031	Jambu fruit, ripe (<i>Syzygium samarangense</i>)	2	83.33	0.82	0.31	0.17	3.07	2.40	0.67	12.30	235
E032	Karonda fruit (<i>Carissa carandas</i>)	1	86.04	1.15	1.02	1.67	7.25	5.87	1.38	2.87	141
E033	Lemon, juice (<i>Citrus limon</i>)	6	91.59±0.53	0.41±0.05	0.28±0.03	0.75±0.02				6.97±0.56	153±9
E034	Lime, sweet,pulp (<i>Citrus limetta</i>)	6	91.32±0.33	0.76±0.09	0.47±0.03	0.20±0.06	2.07±0.11	1.33±0.04	0.74±0.08	5.18±0.37	114±5
E035	Litchi (<i>Litchi chinensis</i>)	4	85.56±0.27	0.99±0.09	0.44±0.06	0.26±0.02	1.34±0.05	0.81±0.10	0.53±0.06	11.41±0.20	225±5
E036	Mango, ripe, banganapalli (<i>Mangifera indica</i>)	6	88.44±0.55	0.54±0.10	0.42±0.04	0.55±0.03	1.88±0.15	1.01±0.08	0.87±0.11	8.18±0.49	175±9
E037	Mango, ripe, gulabkhas (<i>Mangifera indica</i>)	2	86.65	0.52	0.32	0.53	1.67	1.03	0.64	10.32	209

Food code		No. of Regions	Dietary Fibre								Energy
			Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	
							Total	Insoluble	Soluble		
Food Name			←						→	KJ	
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
E038	Mango, ripe, himsagar (<i>Mangifera indica</i>)	3	88.04±0.10	0.46±0.04	0.38±0.02	0.54±0.02	1.55±0.13	0.82±0.19	0.73±0.06	9.03±0.26	187±5
E039	Mango, ripe, kesar (<i>Mangifera indica</i>)	4	85.07±0.59	0.54±0.05	0.44±0.07	0.57±0.03	2.02±0.20	1.09±0.05	0.93±0.14	11.36±0.49	231±8
E040	Mango, ripe, neelam (<i>Mangifera indica</i>)	2	88.41	0.68	0.38	0.55	1.77	0.97	0.79	8.21	178
E041	Mango, ripe, paheri (<i>Mangifera indica</i>)	2	87.69	0.68	0.42	0.58	1.97	0.99	0.98	8.67	188
E042	Mango, ripe, totapari (<i>Mangifera indica</i>)	1	84.14	0.41	0.47	0.49	1.73	0.95	0.78	12.75	248
E043	Mangosteen (<i>Garcinia mangostana</i>)	1	85.52	0.63	0.33	0.24	1.87	1.23	0.64	11.41	219
E044	Manila tamarind (<i>Pithecellobium dulce</i>)	1	74.54	3.56	2.82	1.14	4.40	3.30	1.10	13.54	342
E045	Musk melon, orange flesh (<i>Cucumis melon</i>)	5	92.97±0.38	0.42±0.06	0.52±0.02	0.35±0.02	1.51±0.28	0.84±0.20	0.67±0.11	4.24±0.40	97±8
E046	Musk melon, yellow flesh (<i>Cucumis melon</i>)	6	91.84±0.93	0.53±0.05	0.48±0.04	0.26±0.02	1.49±0.10	0.79±0.08	0.70±0.05	5.40±1.04	116±18
E047	Orange, pulp (<i>Citrus aurantium</i>)	6	89.61±0.19	0.70±0.12	0.36±0.02	0.13±0.02	1.29±0.05	0.73±0.05	0.56±0.02	7.92±0.27	156±4
E048	Palm fruit, tender (<i>Borassus flabellifer</i>)	1	91.93	0.50	0.13	0.12	2.40	1.87	0.53	4.92	101
E049	Papaya, ripe (<i>Carica papaya</i>)	6	91.47±0.76	0.42±0.05	0.51±0.03	0.16±0.01	2.83±0.26	1.75±0.16	1.08±0.10	4.61±0.48	100±9
E050	Peach (<i>Prunus communis</i>)	1	88.31	0.86	0.51	0.37	2.13	1.22	0.91	7.82	168
E051	Pear (<i>Pyrus</i> sp.)	6	86.49±0.18	0.36±0.04	0.32±0.02	0.27±0.04	4.48±0.08	4.02±0.07	0.46±0.03	8.09±0.23	157±3
E052	Phalsa (<i>Grewia asiatica</i>)	2	77.48	1.66	1.09	0.14	4.54	3.44	1.09	15.09	299
E053	Pineapple (<i>Ananas comosus</i>)	6	86.06±1.19	0.52±0.09	0.38±0.04	0.16±0.02	3.46±0.12	2.88±0.08	0.59±0.06	9.42±1.15	180±19
E054	Plum (<i>Prunus domestica</i>)	3	84.44±1.25	0.64±0.13	0.35±0.16	0.40±0.16	2.07±0.36	1.23±0.22	0.84±0.16	12.10±1.47	238±19
E055	Pomegranate, maroon seeds (<i>Punica granatum</i>)	6	83.55±0.39	1.33±0.07	0.57±0.04	0.15±0.02	2.83±0.06	2.26±0.05	0.57±0.02	11.58±0.43	229±6
E056	Pummelo (<i>Citrus maxima</i>)	3	86.97±1.52	0.68±0.06	0.49±0.03	0.42±0.01	0.80±0.04	0.49±0.03	0.31±0.01	10.64±1.54	210±26
E057	Raisins, dried, black (<i>Vitis vinifera</i>)	6	19.69±0.59	2.57±0.20	2.19±0.14	0.34±0.02	3.92±0.23	2.55±0.30	1.37±0.17	71.29±0.48	1279±12
E058	Raisins, dried, golden (<i>Vitis vinifera</i>)	6	21.50±0.51	2.76±0.26	2.04±0.17	0.35±0.03	4.56±0.50	3.04±0.55	1.53±0.26	68.79±1.00	1241±16

Table 1. Proximate Principles and Dietary Fibre

Food code		No. of Regions	Dietary Fibre								Energy
			Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	
							Total	Insoluble	Soluble		
Food Name			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
E059	Rambutan (<i>Nephelium lappaceum</i>)	1	80.87	0.68	0.43	0.16	1.02	0.71	0.30	16.84	306
E060	Sapota (<i>Achras sapota</i>)	6	73.64±1.01	0.92±0.07	0.68±0.05	1.26±0.02	9.60±0.57	8.46±0.58	1.14±0.04	13.90±1.13	307±18
E061	Soursop (<i>Annona muricata</i>)	1	80.85	0.74	0.58	0.94	4.95	3.79	1.16	11.94	260
E062	Star fruit (<i>Averrhoa carambola</i>)	1	91.18	0.79	0.33	0.39	2.81	2.17	0.64	4.51	110
E063	Strawberry (<i>Fragaria x ananassa</i>)	6	92.03±0.64	0.97±0.10	0.54±0.02	0.56±0.02	2.50±0.03	1.51±0.08	0.99±0.08	3.40±0.64	103±11
E064	Tamarind, pulp (<i>Tamarindus indica</i>)	6	20.83±0.57	2.92±0.22	3.44±0.31	0.15±0.02	5.31±0.27	3.73±0.24	1.58±0.09	67.35±0.45	1207±7
E065	Water melon, dark green (sugar baby) (<i>Citrullus vulgaris</i>)	6	94.54±0.63	0.60±0.06	0.13±0.02	0.16±0.01	0.70±0.14	0.35±0.11	0.34±0.05	3.86±0.59	85±10
E066	Water melon, pale green (<i>Citrullus vulgaris</i>)	6	95.33±0.22	0.59±0.03	0.12±0.01	0.16±0.01	0.78±0.11	0.40±0.07	0.38±0.08	3.02±0.21	70±4
E067	Wood Apple (<i>Limonia acidissima</i>)	3	79.36±0.17	3.14±0.30	1.15±0.05	3.62±0.03	5.21±0.38	3.77±0.08	1.44±0.40	7.52±0.58	327±3
E068	Zizyphus (<i>Zizyphus jujube</i>)	1	84.39	1.34	0.80	0.35	3.73	2.71	1.02	9.40	204

F ROOTS AND TUBERS

F001	Beet root (<i>Beta vulgaris</i>)	6	86.95±0.50	1.95±0.14	1.46±0.21	0.14±0.01	3.31±0.32	2.60±0.30	0.71±0.06	6.18±0.61	149±9
F002	Carrot, orange (<i>Daucus carota</i>)	6	87.69±0.56	0.95±0.15	1.16±0.09	0.47±0.02	4.18±0.30	2.81±0.18	1.37±0.21	5.55±0.48	139±9
F003	Carrot, red (<i>Daucus carota</i>)	4	86.07±1.34	1.04±0.09	1.22±0.10	0.47±0.04	4.49±0.19	3.09±0.26	1.40±0.21	6.71±1.11	160±19
F004	Colocasia (<i>Colocasia esculenta</i>)	6	73.49±0.32	3.31±0.59	1.95±0.27	0.17±0.03	3.22±0.34	2.54±0.30	0.68±0.05	17.85±0.94	372±9
F005	Lotus root (<i>Nelumbium nelumbo</i>)	3	76.26±0.87	1.94±0.32	1.50±0.10	0.93±0.06	4.70±0.04	2.86±0.01	1.84±0.05	14.67±0.45	332±14
F006	Potato, brown skin, big (<i>Solanum tuberosum</i>)	6	80.72±0.40	1.54±0.17	0.92±0.08	0.23±0.02	1.71±0.03	1.13±0.02	0.58±0.04	14.89±0.40	292±7
F007	Potato, brown skin, small (<i>Solanum tuberosum</i>)	1	82.97	1.35	0.87	0.22	1.69	1.15	0.54	12.90	255
F008	Potato, red skin (<i>Solanum tuberosum</i>)	1	79.72	1.83	1.13	0.22	1.68	1.11	0.57	15.43	306
F009	Radish, elongate, red skin (<i>Raphanus sativus</i>)	3	89.32±0.29	0.67±0.03	0.73±0.02	0.13±0.02	2.46±0.16	1.96±0.03	0.49±0.15	6.71±0.12	134±3

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			g								KJ
			←							→	
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
F010	Radish, elongate, white skin (<i>Raphanus sativus</i>)	6	89.05±0.72	0.77±0.08	0.82±0.10	0.15±0.02	2.65±0.15	1.98±0.16	0.67±0.08	6.56±0.70	135±11
F011	Radish, round, red skin (<i>Raphanus sativus</i>)	1	89.68	0.89	0.91	0.16	2.29	1.56	0.73	6.07	130
F012	Radish, round, white skin (<i>Raphanus sativus</i>)	2	89.76	0.80	0.80	0.14	2.37	1.63	0.74	6.13	129
F013	Sweet potato, brown skin (<i>Ipomoea batatas</i>)	4	69.21±0.83	1.33±0.12	0.96±0.07	0.26±0.06	3.99±0.05	2.57±0.07	1.43±0.04	24.25±0.77	456±15
F014	Sweet potato, pink skin (<i>Ipomoea batatas</i>)	3	69.58±0.22	1.27±0.09	0.95±0.01	0.33±0.06	3.94±0.10	2.53±0.04	1.41±0.07	23.93±0.15	452±4
F015	Tapioca (<i>Manihot esculenta</i>)	3	75.23±0.55	1.03±0.10	1.12±0.11	0.20±0.01	4.61±0.12	3.85±0.08	0.76±0.05	17.81±0.57	334±10
F016	Water Chestnut (<i>Eleocharis dulcis</i>)	1	73.34	0.86	0.95	0.37	3.02	2.15	0.87	21.46	400
F017	Yam, elephant (<i>Amorphophallus campanulatus</i>)	6	74.39±0.31	2.56±0.28	1.29±0.10	0.14±0.02	4.17±0.05	3.25±0.03	0.92±0.03	17.46±0.55	353±5
F018	Yam, ordinary (<i>Amorphophallus</i> sp.)	4	74.28±0.63	2.18±0.26	1.64±0.19	0.17±0.02	4.08±0.07	3.32±0.32	0.76±0.25	17.65±0.57	349±12
F019	Yam, wild (<i>Dioscorea villosa</i>)	2	69.35	3.07	1.76	0.30	4.57	3.29	1.29	20.95	430
G CONDIMENTS AND SPICES-FRESH											
G001	Chillies, green-1 (<i>Capsicum annum</i>)	5	84.47±0.29	2.62±0.35	1.01±0.14	0.73±0.02	4.86±0.65	3.31±0.42	1.55±0.35	6.32±0.41	191±5
G002	Chillies, green-2 (<i>Capsicum annum</i>)	3	85.72±0.49	2.28±0.13	0.85±0.06	0.75±0.04	5.13±0.56	3.79±0.33	1.34±0.27	5.27±0.42	167±7
G003	Chillies, green-3 (<i>Capsicum annum</i>)	5	85.59±0.79	2.38±0.40	0.95±0.13	0.75±0.02	5.10±0.10	3.61±0.46	1.50±0.49	5.23±0.71	169±12
G004	Chillies, green-4 (<i>Capsicum annum</i>)	3	85.29±0.53	2.33±0.30	0.87±0.13	0.74±0.01	4.15±0.29	2.84±0.17	1.31±0.26	6.63±0.61	190±9
G005	Chillies, green-5 (<i>Capsicum annum</i>)	2	85.93±0.11	2.01±0.26	0.82±0.08	0.65±0.03	3.95±0.60	2.83±0.37	1.12±0.32	6.64±0.39	180±10
G006	Chillies, green-6 (<i>Capsicum annum</i>)	1	84.93	3.01	0.96	0.64	5.15	3.91	1.24	5.31	175
G007	Chillies, green-7 (<i>Capsicum annum</i>)	1	85.83	2.12	0.90	0.60	4.95	3.75	1.20	5.61	163
G008	Chillies, green - all varieties (<i>Capsicum annum</i>)	6	85.39±0.68	2.36±0.34	0.91±0.12	0.72±0.05	4.77±0.61	3.41±0.52	1.37±0.34	5.86±0.77	177±14
G009	Coriander leaves (<i>Coriandrum sativum</i>)	6	86.99±0.41	3.52±0.26	2.19±0.15	0.70±0.06	4.66±0.24	3.24±0.21	1.42±0.15	1.93±0.27	130±9

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
G010	Curry leaves (<i>Murraya koenigii</i>)	6	65.33±1.57	7.41±0.26	4.86±0.58	1.06±0.04	16.83±0.78	13.81±0.43	3.02±0.44	4.51±1.79	266±32
G011	Garlic, big clove (<i>Allium sativum</i>)	6	64.38±0.56	6.92±0.20	1.40±0.11	0.16±0.02	5.22±0.41	2.56±0.38	2.66±0.17	21.93±0.82	518±12
G012	Garlic, small clove (<i>Allium sativum</i>)	3	64.42±0.30	6.75±0.24	1.38±0.17	0.14±0.01	5.47±0.07	2.61±0.02	2.86±0.07	21.84±0.74	514±8
G013	Garlic, single clove, Kashmir (<i>Allium sativum</i>)	1	64.49	6.12	1.76	0.16	4.01	2.20	1.81	23.46	523
G014	Ginger, fresh (<i>Zingiber officinale</i>)	6	81.27±0.22	2.22±0.16	1.33±0.13	0.85±0.03	5.36±0.40	4.28±0.25	1.08±0.21	8.97±0.28	230±4
G015	Mango ginger (<i>Curcuma amada</i>)	3	84.55±0.16	1.45±0.37	1.57±0.14	0.70±0.03	4.74±0.50	3.79±0.13	0.95±0.62	6.98±0.15	177±3
G016	Mint leaves (<i>Mentha spicata</i>)	4	84.24±0.71	4.66±0.15	2.18±0.17	0.65±0.05	5.89±0.49	4.49±0.54	1.40±0.06	2.39±0.36	155±4
G017	Onion, big (<i>Allium cepa</i>)	6	85.76±0.43	1.50±0.10	0.50±0.04	0.24±0.03	2.45±0.21	1.92±0.15	0.53±0.08	9.56±0.44	201±9
G018	Onion, small (<i>Allium cepa</i>)	5	84.67±0.12	1.82±0.20	0.61±0.07	0.16±0.01	1.16±0.19	0.70±0.18	0.46±0.09	11.58±0.37	237±4
G	CONDIMENTS AND SPICES-DRY										
G019	Asafoetida (<i>Ferula assa-foetida</i>)	6	9.42±0.56	6.34±0.53	5.90±0.12	1.26±0.02	5.13±0.33	3.90±0.37	1.23±0.25	71.95±0.71	1387±9
G020	Cardamom, green (<i>Elettaria cardamomum</i>)	6	11.24±0.51	8.10±0.41	7.21±0.46	2.60±0.15	23.10±0.14	20.46±0.09	2.64±0.10	47.76±0.47	1067±16
G021	Cardamom, black (<i>Elettaria cardamomum</i>)	4	6.69±0.47	6.69±0.31	7.84±0.40	2.80±0.24	23.46±0.58	20.73±0.39	2.74±0.37	52.53±1.51	1132±13
G022	Chillies, red (<i>Capsicum annum</i>)	6	14.57±0.42	12.69±0.22	5.73±0.15	6.40±0.04	31.15±0.04	26.55±0.17	4.60±0.15	29.46±0.62	990±8
G023	Cloves (<i>Syzygium aromaticum</i>)	6	26.49±0.68	5.86±0.21	5.99±0.23	8.41±0.15	34.52±0.48	28.07±0.42	6.46±0.10	18.73±0.39	781±6
G024	Coriander seeds (<i>Coriandrum sativum</i>)	6	8.72±0.47	10.66±0.33	5.36±0.15	17.47±0.22	44.81±2.36	35.27±1.77	9.54±2.42	12.98±2.62	1125±38
G025	Cumin seeds (<i>Cuminum cyminum</i>)	6	10.59±0.51	13.91±0.50	5.90±0.09	16.64±0.15	30.35±0.71	25.73±0.77	4.62±0.27	22.62±1.05	1274±19
G026	Fenugreek seeds (<i>Trigonella foenum graecum</i>)	6	7.82±0.37	25.41±0.24	2.93±0.15	5.72±0.02	47.55±0.54	27.63±0.59	19.92±0.51	10.57±0.57	983±10
G027	Mace (<i>Myristica fragrans</i>)	6	20.06±0.70	6.24±0.27	2.47±0.28	24.41±0.20	20.31±0.23	16.78±0.23	3.54±0.17	26.51±0.81	1488±12
G028	Nutmeg (<i>Myristica fragrans</i>)	6	15.55±0.55	6.30±0.24	1.99±0.11	36.52±0.04	11.99±0.18	10.54±0.18	1.46±0.14	27.64±0.70	1940±11
G029	Omum (<i>Trachyspermum ammi</i>)	6	9.71±0.18	15.89±0.61	8.18±0.23	21.11±0.08	20.58±0.04	17.20±0.02	3.38±0.04	24.53±0.68	1495±7

Food code		No. of Regions	Dietary Fibre								Carbohydrate	Energy
			Moisture	Protein	Ash	Total Fat						
							Total	Insoluble	Soluble			
										← g →		
Food Name			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC	
G030	Pippali (<i>Piper longum</i>)	6	10.95±0.48	10.53±0.45	6.41±0.07	2.27±0.02	34.14±0.60	29.57±0.55	4.57±0.39	35.70±0.45	906±8	
G031	Pepper, black (<i>Piper nigrum</i>)	6	13.18±0.40	10.12±0.40	4.58±0.13	2.74±0.02	33.16±0.29	30.61±0.29	2.54±0.11	36.22±0.45	910±8	
G032	Poppy seeds (<i>Papaver somniferum</i>)	6	4.22±0.32	20.31±0.40	6.04±0.19	30.38±1.87	26.68±3.04	15.62±2.29	11.06±1.75	12.37±1.95	1768±77	
G033	Turmeric powder (<i>Curcuma domestica</i>)	6	10.58±0.49	7.66±0.44	6.13±0.38	5.03±0.07	21.38±0.29	18.79±0.42	2.59±0.29	49.22±0.55	1174±7	
H NUTS AND OIL SEEDS												
H001	Almond (<i>Prunus amygdalus</i>)	6	4.37±0.31	18.41±0.04	2.62±0.13	58.49±0.04	13.06±0.31	10.55±0.22	2.52±0.34	3.04±0.24	2549±4	
H002	Arecanut, dried, brown (<i>Areca catechu</i>)	6	6.63±0.15	5.78±0.45	1.39±0.09	4.35±0.03	11.44±0.64	10.14±0.44	1.30±0.32	70.42±0.81	1467±12	
H003	Arecanut, dried, red color (<i>Areca catechu</i>)	3	6.30±0.32	6.46±0.51	1.41±0.04	4.46±0.03	11.11±0.34	10.09±0.25	1.03±0.27	70.27±0.96	1477±9	
H004	Arecanut, fresh (<i>Areca catechu</i>)	2	37.70	2.73	1.43	5.51	7.63	6.59	1.03	45.00	1024	
H005	Cashew nut (<i>Anacardium occidentale</i>)	6	4.44±0.22	18.78±0.58	2.25±0.08	45.20±0.07	3.86±0.47	2.23±0.35	1.63±0.24	25.46±0.70	2438±11	
H006	Coconut, kernel, dry (<i>Cocos nucifera</i>)	6	3.97±0.73	7.27±0.02	1.61±0.17	63.26±0.30	15.88±0.79	14.55±0.23	1.33±0.99	8.01±1.02	2611±9	
H007	Coconut, kernel, fresh (<i>Cocos nucifera</i>)	6	36.14±0.36	3.84±0.04	1.93±0.15	41.38±0.42	10.42±0.28	9.43±0.24	0.99±0.07	6.30±0.43	1711±13	
H008	Garden cress, seeds (<i>Lepidium sativum</i>)	5	4.60±0.21	23.36±0.32	6.37±0.11	23.74±0.09	8.27±0.04	6.33±0.06	1.95±0.10	33.66±0.50	1863±3	
H009	Gingelly seeds, black (<i>Sesamum indicum</i>)	6	4.51±0.44	19.17±0.64	5.78±0.22	43.10±0.03	17.16±0.19	13.57±0.13	3.59±0.13	10.29±0.74	2124±8	
H010	Gingelly seeds, brown (<i>Sesamum indicum</i>)	4	3.62±0.32	21.61±0.71	4.58±0.13	43.22±0.08	17.21±0.18	13.58±0.18	3.63±0.11	9.76±1.01	2161±8	
H011	Gingelly seeds, white (<i>Sesamum indicum</i>)	5	3.30±0.28	21.70±0.44	4.13±0.16	43.05±0.04	16.99±0.30	13.49±0.19	3.51±0.15	10.83±0.50	2174±9	
H012	Ground nut (<i>Arachis hypogaea</i>)	6	6.97±0.69	23.65±0.85	2.11±0.07	39.63±0.29	10.38±0.18	8.58±0.15	1.79±0.22	17.27±0.33	2176±9	
H013	Mustard seeds (<i>Brassica nigra</i>)	6	5.67±0.48	19.51±0.23	3.73±0.05	40.19±0.21	14.10±0.66	10.63±0.57	3.47±0.20	16.80±0.71	2132±16	
H014	Linseeds (<i>Linum usitatissimum</i>)	6	5.48±0.28	18.55±0.15	3.15±0.10	35.67±0.70	26.17±0.39	21.83±0.60	4.33±0.84	10.99±0.86	1857±16	
H015	Niger seeds, black (<i>Guizotia abyssinica</i>)	4	4.65±0.59	18.92±1.61	3.91±0.26	38.61±0.41	10.93±0.05	10.50±0.03	0.43±0.04	22.98±2.64	2144±11	

Table 1. Proximate Principles and Dietary Fibre

Food code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Dietary Fibre			Carbohydrate	Energy
							Total	Insoluble	Soluble		
			← g →								KJ
			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
H016	Niger seeds, gray (<i>Guizotia abyssinica</i>)	5	5.66±0.29	18.34±2.11	4.92±0.22	39.53±0.26	10.96±0.14	10.54±0.12	0.41±0.03	20.59±1.85	2128±6
H017	Pine seed (<i>Pinus</i> sp.)	5	5.32±0.15	12.55±0.25	2.78±0.10	48.79±0.10	3.79±0.03	2.23±0.03	1.57±0.03	26.77±0.35	2486±3
H018	Pistachio nuts (<i>Pistacia vera</i>)	6	4.68±0.22	23.35±0.35	3.02±0.13	42.49±0.34	10.64±0.16	8.23±0.15	2.41±0.11	15.82±0.52	2257±10
H019	Safflower seeds (<i>Carthamus tinctorius</i>)	5	5.24±0.16	17.66±0.54	2.57±0.15	30.87±0.98	13.49±0.45	10.24±0.27	3.25±0.22	30.18±1.20	1981±20
H020	Sunflower seeds (<i>Helianthus annuus</i>)	5	3.53±0.29	23.53±0.56	3.44±0.26	51.85±0.07	10.80±0.13	8.51±0.17	2.29±0.07	6.85±0.99	2453±8
H021	Walnut (<i>Juglans regia</i>)	6	3.56±0.22	14.92±0.62	1.72±0.08	64.27±0.05	5.39±0.19	4.74±0.17	0.65±0.03	10.14±0.69	2809±3
I SUGARS											
I001	Jaggery, cane (<i>Saccharum officinarum</i>)	6	11.20±0.49	1.85±0.19	1.92±0.17	0.16±0.02				84.87±0.63	1480±9
I002	Sugarcane, juice (<i>Saccharum officinarum</i>)	6	85.54±0.99	0.16±0.03	0.23±0.02	0.40±0.05	0.56±0.09	0.40±0.05	0.16±0.05	13.11±0.93	242±18
J MUSHROOMS											
J001	Button mushroom, fresh (<i>Agaricus</i> sp.)	1	90.09	3.68	0.71	0.42	3.11	2.76	0.35	1.98	115
J002	Chicken mushroom, fresh (<i>Lactiporus</i> sp.)	1	92.43	1.84	0.74	0.25	1.99	1.82	0.18	2.76	89
J003	Shiitake mushroom, fresh (<i>Lentinula</i> sp.)	1	82.94	3.19	1.11	0.76	3.02	2.03	0.99	8.98	243
J004	Oyster mushroom, dried (<i>Pleurotus</i> sp.)	1	4.51	19.04	1.41	2.86	39.12	35.64	3.48	33.07	1019
K MISCELLANEOUS FOODS											
K001	Toddy (<i>Borassus flabellifer</i>)	10	93.86±0.59	0.18±0.05	0.21±0.03	0.03±0.01				5.72±0.55	101±10
K002	Coconut Water (<i>Cocos nucifera</i>)	6	95.77±0.42	0.26±0.04	0.65±0.06	0.16±0.02				3.16±0.39	64±7

Food code		No. of Regions	Dietary Fibre								Energy
			Moisture	Protein	Ash	Total Fat				Carbohydrate	
						Total	Insoluble	Soluble			
			←—————g—————→								
Food Name			WATER	PROTCNT	ASH	FATCE	FIBTG	FIBINS	FIBSOL	CHOAVLDF	ENERC
L	MILK AND MILK PRODUCTS										
L001	Milk, whole, Buffalo	6	80.68±0.66	3.68±0.13	0.67±0.02	6.58±0.20				8.39±0.71	449±9
L002	Milk, whole, Cow	6	86.64±1.10	3.26±0.06	0.68±0.02	4.48±0.29				4.94±1.02	305±23
L003	Paneer	6	51.96±0.76	18.86±0.75	1.98±0.08	24.78±0.17				2.41±0.12	1278±61
L004	Khoa	6	42.51±0.21	16.34±0.61	4.00±0.14	20.62±0.83				16.53±1.26	1322±14

Food Code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			←————— g —————→				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
M	EGG AND EGG PRODUCTS						
M001	Egg, poultry, whole, raw	6	76.51±0.20	13.28±0.29	0.81±0.04	9.15±0.14	564±3
M002	Egg, poultry, white, raw	6	86.68±0.11	10.84±0.07	0.75±0.03	0.06±0.01	187±1
M003	Egg, poultry, yolk, raw	6	53.57±0.40	15.74±0.33	1.05±0.05	26.34±0.17	1242±9
M004	Egg, poultry, whole, boiled	6	73.46±0.22	13.43±0.28	0.86±0.04	10.54±0.19	618±6
M005	Egg, poultry, white, boiled	6	83.54±0.18	12.37±0.16	0.81±0.04	0.26±0.02	220±3
M006	Egg, poultry, yolk, boiled	6	51.42±0.22	16.13±0.45	1.38±0.14	27.46±0.24	1290±8
M007	Egg, poultry, omlet	3	68.48±0.13	16.53±0.43	0.96±0.03	11.60±0.22	710±4
M008	Egg, country hen, whole, raw	1	72.96	13.14	0.86	13.00	704
M009	Egg, country hen, whole, boiled	1	70.43	14.43	0.94	14.10	767
M010	Egg, country hen, omlet	1	67.80	14.80	1.08	16.30	855
M011	Egg, duck, whole, boiled	1	71.61	13.80	0.97	13.61	738
M012	Egg, duck, whole, raw	1	70.53	14.60	0.99	13.83	760
M013	Egg, duck, whole, omlet	1	68.90	15.10	1.18	14.80	804
M014	Egg, quial, whole, raw	1	75.11	12.39	0.94	11.46	635
M015	Egg, quial, whole, boiled	1	74.41	13.03	0.99	11.50	647
N	POULTRY						
N001	Chicken, poultry, leg, skinless	4	67.61±0.30	19.44±0.79	1.14±0.08	12.64±0.53	1605±23
N002	Chicken, poultry, thigh, skinless	6	67.58±0.46	18.18±0.57	1.14±0.10	14.23±0.84	836±38
N003	Chicken, poultry, breast, skinless	5	67.15±0.18	21.81±0.81	1.19±0.09	9.00±0.52	704±24
N004	Chicken, poultry, wing, skinless	5	67.48±0.21	17.42±0.60	1.13±0.05	13.81±0.26	807±17

Food Code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			g				
			KJ				
			WATER	PROTCNT	ASH	FATCE	ENERC
N005	Poultry, chicken, liver	1	73.20	21.57	1.11	4.08	518
N006	Poultry, chicken, gizzard	1	78.05	18.22	1.46	2.07	386
N007	Country hen, leg, with skin	1	70.02	17.01	1.20	11.73	723
N008	Country hen, thigh, with skin	1	67.46	18.29	1.23	12.82	785
N009	Country hen, breast, with skin	1	66.53	22.08	1.14	10.20	753
N010	Country hen, wing, with skin	1	68.01	18.65	1.14	12.09	764
N011	Duck, meat, with skin	1	73.49	19.07	1.22	6.03	547
N012	Emu, meat, skinless	1	71.65	22.65	0.98	4.62	556
N013	Guinea fowl, meat, with skin	1	75.21	20.52	0.92	3.24	469
N014	Pigeon, meat, with skin	1	74.61	17.94	1.32	6.03	528
N015	Quail, meat, skinless	1	71.58	20.92	1.35	5.95	576
N016	Turkey, leg, with skin	1	69.24	20.30	1.21	8.15	647
N017	Turkey, thigh, with skin	1	72.17	20.46	0.97	6.31	581
N018	Turkey, breast, with skin	1	68.44	21.96	1.01	8.04	671
N019	Turkey, wing, with skin	1	66.16	21.91	0.97	10.77	771
O	ANIMAL MEAT						
O001	Goat, shoulder	6	66.39±0.40	20.33±0.50	0.90±0.02	11.94±0.78	787±33
O002	Goat, chops	6	72.46±1.10	20.39±0.83	0.99±0.01	5.98±0.37	568±26
O003	Goat, legs	6	68.85±0.86	22.07±0.63	0.97±0.03	7.94±0.62	669±26
O004	Goat, brain	5	76.74±1.02	13.82±0.80	1.29±0.07	8.06±0.46	533±23
O005	Goat, tongue	4	68.40±1.00	16.63±0.54	1.02±0.04	13.68±1.09	789±39

Table 1. Proximate Principles and Dietary Fibre


Food Code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			←————— g —————→				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
O006	Goat, lungs	4	79.03±1.20	16.86±0.71	0.74±0.01	3.09±0.56	401±32
O007	Goat, heart	5	75.15±1.23	19.38±0.80	0.95±0.09	4.40±0.71	492±33
O008	Goat, liver	6	73.37±1.16	20.32±0.92	1.30±0.07	4.88±0.73	526±31
O009	Goat, tripe	5	80.94±0.84	15.36±0.50	0.23±0.02	3.36±0.29	386±19
O010	Goat, spleen	4	77.93±0.50	18.45±0.15	1.15±0.09	2.37±0.36	401±14
O011	Goat, kidneys	4	80.26±1.07	15.60±1.15	1.10±0.05	2.93±0.18	374±18
O012	Goat, tube (small intestine)	3	78.12±0.63	12.92±0.54	0.54±0.01	8.26±1.08	525±32
O013	Goat, testis	2	84.29	12.32	0.87	2.39	298
O014	Sheep, shoulder	5	66.51±2.51	18.24±1.41	0.86±0.01	14.31±1.37	840±68
O015	Sheep, chops	4	75.66±2.78	18.00±1.22	1.09±0.02	5.15±2.05	496±86
O016	Sheep, leg	5	68.26±1.01	21.42±0.43	0.94±0.01	8.69±0.74	686±25
O017	Sheep, brain	1	78.36	13.05	1.19	7.29	492
O018	Sheep, tongue	1	68.70	16.61	1.06	13.53	783
O019	Sheep, lungs	2	80.41	16.12	0.97	2.41	363
O020	Sheep, heart	1	77.06	18.19	0.97	3.66	445
O021	Sheep, liver	4	69.71±1.12	22.26±1.46	1.26±0.07	4.87±0.12	559±21
O022	Sheep, tripe	2	78.81	16.76	0.26	4.05	435
O023	Sheep, spleen	1	79.66	16.02	1.17	3.02	384
O024	Sheep, kidneys	2	79.77	16.22	1.00	2.92	384
O025	Beef, shoulder	6	63.84±0.59	20.56±1.32	0.95±0.01	14.59±0.86	889±12
O026	Beef, chops	4	72.40±0.34	19.82±0.43	0.96±0.02	6.71±0.45	585±13
O027	Beef, round (leg)	6	68.04±1.36	22.64±1.03	1.19±0.01	7.38±0.71	658±37


Food Code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			g				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
O028	Beef, brain	4	78.71±1.43	10.55±0.50	1.37±0.17	9.28±0.92	523±42
O029	Beef, tongue	4	70.97±1.20	15.63±0.67	0.73±0.00	12.57±0.63	731±32
O030	Beef, lungs	3	80.83±0.18	15.66±0.42	1.13±0.05	2.28±0.21	351±1
O031	Beef, heart	5	77.77±0.72	17.68±0.51	0.87±0.01	3.57±0.46	433±20
O032	Beef, liver	6	74.15±1.31	20.73±0.96	1.07±0.03	3.96±0.45	499±30
O033	Beef, tripe	5	83.74±0.62	13.10±0.29	0.55±0.01	2.51±0.74	316±25
O034	Beef, spleen	6	79.03±0.73	17.42±0.55	1.22±0.04	2.22±0.24	378±16
O035	Beef, kidneys	3	77.59±0.66	17.09±0.30	1.22±0.04	4.01±0.36	439±17
O036	Calf, shoulder	2	70.49	20.98	0.94	7.47	633
O037	Calf, chops	2	72.66	22.47	0.92	3.85	524
O038	Calf, round (leg)	2	71.13	21.10	0.74	6.92	615
O039	Calf, brain	2	81.19	9.84	1.26	7.60	448
O040	Calf, tongue	2	69.44	17.78	0.93	11.76	737
O041	Calf, heart	1	73.44	18.84	0.86	3.75	459
O042	Calf, liver	1	73.66	21.02	1.26	3.95	503
O043	Calf, spleen	1	78.65	17.71	1.41	2.12	379
O044	Calf, kidneys	2	80.23	15.16	1.02	3.50	387
O045	Mithun, shoulder	1	68.86	19.05	0.84	11.14	736
O046	Mithun, chops	1	73.81	18.19	0.79	6.12	536
O047	Mithun, round (leg)	1	72.25	19.61	0.90	4.10	485
O048	Pork, shoulder	6	62.92±0.74	17.41±0.71	0.75±0.12	18.83±0.88	993±28
O049	Pork, chops	6	68.40±1.25	19.41±0.68	0.64±0.05	11.30±1.33	748±51

Table 1. Proximate Principles and Dietary Fibre

Food Code	Food Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			←————— g —————→				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
O050	Pork, ham	6	61.94±0.39	18.83±0.59	0.62±0.02	18.55±0.78	1006±20
O051	Pork, lungs	4	81.12±0.72	15.13±0.52	0.93±0.01	2.74±0.23	358±16
O052	Pork, heart	4	77.74±0.88	16.31±0.16	1.01±0.01	4.87±0.84	457±32
O053	Pork, liver	5	74.88±1.23	19.89±1.16	1.18±0.03	3.94±0.34	484±24
O054	Pork, stomach	1	76.51	15.36	0.23	7.81	550
O055	Pork, spleen	3	80.93±1.08	15.13±0.79	1.13±0.05	2.71±0.71	357±31
O056	Pork, kidneys	5	80.92±0.87	14.35±1.03	0.97±0.10	3.65±0.55	379±21
O057	Pork, tube (small intestine)	1	75.42	14.96	0.55	8.98	587
O058	Hare, shoulder	1	71.06	21.13	1.17	6.58	603
O059	Hare, chops	1	75.47	20.62	1.27	2.55	445
O060	Hare, leg	1	73.00	20.52	1.26	4.16	503
O061	Rabbit, shoulder	1	70.75	20.01	1.19	7.96	635
O062	Rabbit, chops	1	71.03	22.61	1.35	4.88	565
O063	Rabbit, leg	1	70.33	21.31	1.26	5.99	584

Food Code	Fish Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			g				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
P	MARINE FISH						
P001	Allathi (<i>Elops machnata</i>)	1	75.91	21.77	1.06	0.98	406
P002	Aluva (<i>Parastromateus niger</i>)	3	75.71±0.06	21.63±0.28	1.39±0.08	1.80±0.51	434±17
P003	Anchovy (<i>Stolephorus indicus</i>)	2	77.77	19.88	1.62	0.78	367
P004	Ari fish (<i>Aprion virescens</i>)	1	77.09	22.01	1.34	1.11	415
P005	Betki (<i>Lates calcarifer</i>)	1	82.52	15.28	1.12	0.29	284
P006	Black snapper (<i>Macolor niger</i>)	1	78.09	19.50	1.19	1.24	377
P007	Bombay duck (<i>Harpadon nehereus</i>)	2	83.24	13.53	1.07	1.03	287
P008	Bommuralu (<i>Muraenesox cinerius</i>)	3	76.42±0.05	22.30±0.27	1.21±0.20	2.87±0.29	485±14
P009	Cat fish (<i>Tachysurus thalassinus</i>)	1	76.22	22.18	1.32	2.13	456
P010	Chakla (<i>Rachycentron canadum</i>)	5	78.36±0.07	20.21±0.70	1.07±0.09	1.68±1.00	406±36
P011	Chappal (<i>Aluterus monoceros</i>)	1	80.61	17.17	1.23	0.67	317
P012	Chelu (<i>Elagatis bipinnulata</i>)	2	76.36	20.02	1.28	0.70	366
P013	Chembali (<i>Lutjanus quinquelineatus</i>)	1	77.54	20.06	0.86	1.77	406
P014	Eri meen (<i>Pristipomoides filamentosus</i>)	1	76.54	22.30	1.28	2.02	454
P015	Gobro (<i>Epinephelus diacanthus</i>)	2	78.84	19.37	0.98	0.88	362
P016	Guitar fish (<i>Rhinobatus prahli</i>)	1	75.73	22.56	1.32	0.44	400
P017	Hilsa (<i>Tenualosa ilisha</i>)	2	60.09	21.80	1.11	18.49	1083
P018	Jallal (<i>Arius sp.</i>)	1	77.50	21.55	1.17	1.46	420
P019	Jathi vela meen (<i>Lethrinus lentjan</i>)	2	75.66	22.45	1.26	1.92	453
P020	Kadal bral (<i>Synodus indicus</i>)	1	79.66	18.72	1.27	1.34	368
P021	Kadali (<i>Nemipterus mesoprion</i>)	1	73.74	22.07	1.06	4.20	530

Food Code	Fish Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy	
								KJ
			WATER	PROTCNT	ASH	FATCE	ENERC	
P022	Kalamaara (<i>Leptomelanosoma indicum</i>)	2	75.50	20.89	1.08	4.54	523	
P023	Kalava (<i>Epinephelus coioides</i>)	1	80.02	19.38	1.34	1.20	374	
P024	Kanamayya (<i>Lutjanus rivulatus</i>)	2	77.59	20.19	1.07	0.65	367	
P025	Kannadi paarai (<i>Alectis indicus</i>)	3	76.25±0.65	22.09±0.24	1.28±0.19	1.27±0.40	423±11	
P026	Karimeen (<i>Eetroplus suratensis</i>)	1	78.66	19.68	0.97	1.38	386	
P027	Karnagawala (<i>Anchoa hepsetus</i>)	1	79.22	19.52	1.45	0.84	363	
P028	Kayrai (<i>Thunnus albacores</i>)	2	72.69	20.15	1.28	3.00	454	
P029	Kiriyan (<i>Atule mate</i>)	1	72.39	22.49	0.94	4.70	556	
P030	Kite fish (<i>Mobula kuhlii</i>)	1	77.61	23.07	1.50	0.56	413	
P031	Korka (<i>Terapon jarbua</i>)	1	72.25	23.72	1.09	3.36	528	
P032	Kulam paarai (<i>Carangoides fulvoguttatus</i>)	1	73.44	21.62	1.25	3.60	501	
P033	Maagaa (<i>Polynemus plebeius</i>)	1	79.23	20.11	1.01	0.67	366	
P034	Mackerel (<i>Rastrelliger kanagurta</i>)	3	74.55±1.14	21.51±0.34	1.20±0.43	1.53±0.93	423±40	
P035	Manda clathi (<i>Naso reticulatus</i>)	1	76.32	21.27	1.48	1.00	398	
P036	Matha (<i>Acanthurus mata</i>)	2	79.81	21.17	1.35	0.79	389	
P037	Milk fish (<i>Chanos chanos</i>)	1	72.22	23.66	1.12	1.09	442	
P038	Moon fish (<i>Mene maculata</i>)	1	74.21	20.74	1.69	4.69	526	
P039	Mullet (<i>Mugil cephalus</i>)	3	76.12±0.33	20.23±1.30	1.16±0.13	1.32±0.27	393±26	
P040	Mural (<i>Tylosurus crocodilus</i>)	1	78.96	19.04	1.11	0.58	345	
P041	Myil meen (<i>Istiophorus platypterus</i>)	2	75.57	22.89	1.21	0.56	410	
P042	Nalla bontha (<i>Epinephelus</i> sp.)	1	79.32	19.82	1.00	0.72	364	
P043	Narba (<i>Caranx sexfasciatus</i>)	2	76.37	21.95	1.17	1.60	432	

Food Code	Fish Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
							KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
P044	Paarai (<i>Caranx heberi</i>)	1	75.77	21.58	1.21	1.84	435
P045	Padayappa (<i>Canthidermis maculata</i>)	1	79.18	19.70	1.26	0.69	360
P046	Pali kora (<i>Panna microdon</i>)	1	78.63	19.25	1.00	1.81	394
P047	Pambada (<i>Lepturacanthus savala</i>)	2	74.67	21.95	1.41	4.44	537
P048	Pandukopa (<i>Pseudosciaena manchurica</i>)	1	77.46	19.73	1.15	0.65	360
P049	Parava (<i>Lactarius lactarius</i>)	1	77.19	21.50	1.08	2.74	467
P050	Parcus (<i>Psettodes erumei</i>)	1	79.01	19.87	1.12	0.62	361
P051	Parrot fish (<i>Scarus ghobban</i>)	1	76.84	20.82	1.40	0.57	375
P052	Perinkilichai (<i>Pinjalo pinjalo</i>)	1	78.33	20.89	1.02	0.87	387
P053	Phopat (<i>Coryphaena hippurus</i>)	4	76.42±0.14	22.05±0.75	1.24±0.02	1.33±0.47	424±8
P054	Piranha (<i>Pygopritis sp.</i>)	1	76.15	20.46	0.93	5.43	549
P055	Pomfret, black (<i>Parastromateus niger</i>)	1	74.43	18.91	0.97	4.83	515
P056	Pomfret, snub nose (<i>Trachinotus blochii</i>)	2	77.81	21.05	1.29	0.46	375
P057	Pomfret, white (<i>Pampus argenteus</i>)	2	75.91	19.02	1.01	5.12	513
P058	Pranel (<i>Gerres sp.</i>)	1	79.54	19.69	0.76	1.89	405
P059	Pulli paarai (<i>Gnathanodon speciosus</i>)	1	75.99	20.06	1.19	1.55	399
P060	Queen fish (<i>Scomberoides commersonianus</i>)	3	76.63±0.23	20.97±0.31	1.22±0.09	1.17±0.20	400±12
P061	Raai fish (<i>Lobotes surinamensis</i>)	2	77.14	21.61	1.11	1.69	430
P062	Raai vanthu (<i>Epinephelus chlorostigma</i>)	1	79.95	19.39	0.92	2.17	410
P063	Rani (<i>Pink perch</i>)	1	78.55	18.83	1.01	1.49	377
P064	Ray fish, bow head, spotted (<i>Rhina ancylostoma</i>)	1	80.30	19.03	1.15	0.70	349
P065	Red snapper (<i>Lutjanus argentimaculatus</i>)	1	76.28	22.79	1.26	1.34	437

Food Code	Fish Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			g				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
P066	Red snapper, small (<i>Priacanthus hamrur</i>)	1	76.31	21.50	1.10	2.30	451
P067	Sadaya (<i>Platax orbicularis</i>)	1	75.53	20.68	1.32	2.97	462
P068	Salmon (<i>Salmo salar</i>)	1	67.84	20.97	1.13	9.86	721
P069	Sangada (<i>Nemipterus japonicus</i>)	1	78.51	20.26	1.26	2.67	443
P070	Sankata paarai (<i>Caranx ignobilis</i>)	1	74.88	21.87	1.18	1.68	434
P071	Sardine (<i>Sardinella longiceps</i>)	1	72.29	17.91	0.84	8.99	637
P072	Shark (<i>Carcharhinus sorrah</i>)	2	72.82	21.60	1.08	0.83	398
P073	Shark, hammer head (<i>Sphyrna mokarran</i>)	1	74.61	23.40	0.92	0.80	432
P074	Shark, spotted (<i>Stegostoma fasciatum</i>)	1	78.83	20.95	1.04	0.75	384
P075	Shelavu (<i>Sphyaena jello</i>)	4	74.60±1.68	22.46±1.34	1.26±0.07	1.74±0.50	446±41
P076	Silan (<i>Silonia silondia</i>)	1	70.12	22.72	0.77	6.68	633
P077	Silk fish (<i>Beryx</i> sp.)	1	77.61	20.65	1.05	1.53	408
P078	Silver carp (<i>Hypophthalmichthys molitrix</i>)	1	72.47	21.71	0.95	5.17	555
P079	Sole fish (<i>Cynoglossus arel</i>)	1	80.18	19.05	1.06	1.18	367
P080	Stingray (<i>Dasyatis pastinaca</i>)	1	75.67	23.98	1.14	0.68	408
P081	Tarlava (<i>Drepane punctata</i>)	2	76.76	21.72	1.14	1.20	414
P082	Tholam (<i>Plectorhinchus schotaf</i>)	2	76.20	20.78	1.06	2.13	432
P083	Tilapia (<i>Oreochromis niloticus</i>)	1	79.57	18.48	1.18	1.08	349
P084	Tuna (<i>Euthynnus affinis</i>)	5	72.12±0.42	24.50±0.27	1.11±0.12	1.44±0.42	470±16
P085	Tuna, striped (<i>Katsuwonus pelamis</i>)	1	77.76	21.28	0.87	1.12	403
P086	Valava (<i>Chirocentrus nudus</i>)	1	77.25	21.80	1.27	1.06	410
P087	Vanjaram (<i>Scomberomorus commerson</i>)	2	72.30	22.28	1.33	5.18	570

Food Code	Fish Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			← g →				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
P088	Vela meen (<i>Aprion virescens</i>)	1	71.82	22.16	1.34	4.34	537
P089	Vora (<i>Siganus javus</i>)	2	76.94	20.14	1.26	2.14	422
P090	Whale shark (<i>Galeocерdo cuvier</i>)	1	77.59	21.83	1.48	0.82	401
P091	Xiphinis (<i>Xiphias gladius</i>)	1	79.22	19.82	1.15	0.88	370
P092	Eggs, Cat fish (<i>Ompok bimaculatus</i>)	1	69.34	24.68	1.57	5.28	600

Q MARINE SHELLFISH

Q001	Crab (<i>Menippe mercenaria</i>)	1	79.77	10.23	1.74	1.41	343
Q002	Crab, sea (<i>Portunus sanguinolentus</i>)	1	79.67	15.36	0.92	0.60	283
Q003	Lobster, brown (<i>Thenus orientalis</i>)	1	81.48	15.96	1.29	0.56	292
Q004	Lobster, king size (<i>Thenus orientalis</i>)	1	77.77	18.54	1.08	0.78	375
Q005	Mud crab (<i>Scylla tranquebarica</i>)	1	83.10	10.00	2.60	0.55	190
Q006	Oyster (<i>Crassostrea</i> sp.)	1	82.50	9.51	2.47	2.44	252
Q007	Tiger prawns, brown (<i>Solenocera crassicornis</i>)	1	82.41	14.85	0.94	0.56	273
Q008	Tiger Prawns, orange (<i>Penaeus monodon</i>)	1	81.48	14.25	0.83	0.74	270

R MARINE MOLLUSKS

R001	Clam, green shell (<i>Perna viridis</i>)	1	80.07	12.14	2.44	0.98	243
R002	Clam, white shell, ribbed (<i>Meretrix meretrix</i>)	1	80.27	11.82	0.85	1.34	250
R003	Octopus (<i>Octopus vulgaris</i>)	1	80.45	14.71	1.21	1.12	334
R004	Squid, black (<i>Loligo</i> sp.)	1	80.59	16.10	0.99	1.07	335
R005	Squid, hard shell (<i>Sepia pharaonis</i>)	1	80.52	16.82	0.75	0.93	320

Food Code	Fish Name	No. of Regions	Moisture	Protein	Ash	Total Fat	Energy
			←————— g —————→				KJ
			WATER	PROTCNT	ASH	FATCE	ENERC
R006	Squid, red (<i>Loligo duvaucelii</i>)	2	80.85	16.25	1.15	1.44	329
R007	Squid, white, small (<i>Uroteuthis duvauceli</i>)	1	79.38	17.41	1.19	1.15	353
S FRESHWATER FISH AND SHELLFISH							
S001	Cat fish (<i>Tandanus tandanus</i>)	2	77.23±0.91	15.86±3.65	0.93±0.01	6.24±3.03	518±84
S002	Catla (<i>Catla catla</i>)	6	78.43±0.71	17.94±1.36	0.94±0.01	2.15±0.34	394±6
S003	Freshwater Eel (<i>Anguilla anguilla</i>)	1	75.57	20.41	1.03	2.63	451
S004	Gold fish (<i>Carassius auratus</i>)	2	79.07	16.91	1.10	2.94	396
S005	Pangas (<i>Pangasianodon hypophthalmus</i>)	6	68.50±1.01	17.12±1.62	1.12±0.03	16.77±0.91	852±24
S006	Rohu (<i>Labeo rohita</i>)	6	76.34±0.28	19.71±0.57	1.28±0.04	2.39±0.64	428±12
S007	Crab (<i>Pachygrapsus</i> sp.)	1	80.58	13.23	1.23	0.89	327
S008	Prawns, big (<i>Macrobrachium rosenbergii</i>)	1	77.43	19.24	0.84	0.52	380
S009	Prawns, small (<i>Macrobrachium</i> sp.)	3	82.58	13.07	0.86	0.78	297
S010	Tiger prawns (<i>Macrobrachium</i> sp.)	2	83.24	14.24	0.83	0.66	284