

Daily Sequences Drill #7 — Answers & Investigations

Sheet by David Akinyele-Aje

Section	Topic	Questions	Target time
A	Rapid Recognition	10	2 min 30 sec
B	Manipulation Drills	10	8 min
C	Substitution & Structure	8	10 min
D	Challenge	5	15 min

New toolkit today: Mixed Synthesis Capstone · Every technique from Days 1–6, combined.

Section A Rapid Recognition

1. Write down the n th term of the AP 4, 9, 14, 19, ...

Answer: $5n - 1$

Method: $a = 4, d = 5: a_n = 4 + 5(n - 1) = 5n - 1$.

Investigate further: Seen through Day 4's lens, an AP is the recurrence $a_n = a_{n-1} + d$ with characteristic root $x = 1$ and a constant forcing term d that happens to match that root — exactly the “resonance” case from Day 4 C5, where the particular solution is linear in n rather than a plain constant. That's the real reason a_n comes out linear instead of exponential.

2. Evaluate $\binom{5}{0} + \binom{5}{1} + \binom{5}{2} + \binom{5}{3} + \binom{5}{4} + \binom{5}{5}$.

Answer: 32

Method: $\sum_{k=0}^5 \binom{5}{k} = 2^5 = 32$ (Day 2's row-sum identity, $x = 1$).

Investigate further: This identity counts every subset of a 5-element set by size — and C2 later in this sheet puts the resulting 2^n head to head against Day 6's simplest possible sum, $a_k = 1$ repeated n times: the row sum's exponential growth completely dwarfs that linear one, even though both are “just adding up $n + 1$ (or n) numbers.”

3. A GP has $a = 6, r = \frac{1}{3}$. Find its sum to infinity.

Answer: 9

Method: $S_\infty = \frac{6}{1 - 1/3} = \frac{6}{2/3} = 9$ (Day 3).

Investigate further: S_∞ is really just $\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} a \frac{1-r^{n+1}}{1-r}$ — the exact same “ $r^n \rightarrow 0$ when $|r| < 1$ ” mechanism that makes a Day 4 recurrence's roots die out and its sequence converge. A GP is secretly the simplest possible linear recurrence ($a_n = ra_{n-1}$, one root, no forcing term), so this isn't a coincidence.

4. Write down the characteristic equation of $a_n = 7a_{n-1} - 10a_{n-2}$.

Answer: $x^2 - 7x + 10 = 0$

Method: Substitute $x^2, x, 1$ for a_n, a_{n-1}, a_{n-2} (Day 4).

Investigate further: The coefficients 7 and -10 are not arbitrary: by Vieta's formulas on $x^2 - 7x + 10 = 0$, the roots must sum to 7 and multiply to 10 — so spotting $2 + 5 = 7$, $2 \times 5 = 10$ finds both roots without the quadratic formula at all, the same shortcut Day 4 A4 relies on.

5. Find the fixed point of $f(x) = 3x - 8$.

Answer: $x = 4$

Method: $x = 3x - 8 \Rightarrow -2x = -8 \Rightarrow x = 4$ (Day 5/Day 6's fixed-point technique).

Investigate further: This is the exact value a Day 5 iteration $a_{n+1} = f(a_n)$ would converge to if it ever converged, and the exact value Day 6's alternating process would get permanently stuck at if it were ever written down — three days, three different-looking setups, one shared idea: a fixed point is wherever “apply the map” stops changing anything.

6. If the mean of the first 5 terms of a sequence is an integer, what can be said about their sum?

Answer: Their sum must be a multiple of 5.

Method: Integer mean $\iff$ sum divisible by count (Day 6).

Investigate further: This single equivalence — integer mean $\iff$ sum divisible by count — is the entire engine behind Day 6's capstone constructions (B1's 1, 3, 5, 7, C7's greedy step, D1's card-stacking proof) and D4 later today: everything in that whole strand of questions is this one fact, applied at a different index each time.

7. Is 2, 6, 18, 54, ... an AP or a GP?

Answer: GP (ratio 3)

Method: $6/2 = 18/6 = 54/18 = 3$, constant ratio.

Investigate further: The two tests are mirror images: an AP is spotted by constant *differences*, a GP by constant *ratios*. Both need only three consecutive terms to check — but as Day 1's C3 (squaring an AP) showed, confirming “arithmetic” or “geometric” from a handful of terms is suggestive, not a proof, unless a general n th-term formula backs it up.

8. A recurrence has both characteristic roots with magnitude < 1 . What happens to the sequence as $n \rightarrow \infty$?

Answer: Converges to 0.

Method: Both roots inside the unit circle $\Rightarrow a_n = Ar_1^n + Br_2^n \rightarrow 0$ (Day 4 D5, Day 5).

Investigate further: Day 4 gets here algebraically ($a_n = Ar_1^n + Br_2^n$ with $|r_1|, |r_2| < 1$ forces both terms to 0); Day 5 would get to the same place by showing the sequence is eventually monotonic and bounded below by 0. Two completely different-looking arguments landing on the identical conclusion is exactly the kind of thing worth double-checking rather than dismissing as coincidence.

9. Find the coefficient of x^2 in $(1 + x)^5$.

Answer: 10

Method: $\binom{5}{2} = 10$ (Day 2).

Investigate further: By symmetry $\binom{5}{2} = \binom{5}{3}$, so this is also “the coefficient of x^3 ” for free — and C4 later today reuses this exact extraction skill on a weighted series $(k+1)\left(\frac{1}{3}\right)^k$, where the binomial coefficient sits one layer deeper inside a generating-function identity.

10. Write down the fixed-point formula for $f(x) = rx - c$ ($r \neq 1$).

Answer: $x = \frac{c}{r-1}$

Method: Solve $x = rx - c$ for x (Day 6).

Investigate further: Compare this single-map fixed point to Day 6’s two-player version: there, Alice’s own combined per-turn map is $f(x) = rx - c$ too, so her fixed point is this exact same formula — the alternating writer/Bob structure only matters for finding the OTHER values that reappear (B6–B8), not for the fixed point itself.

Section B Manipulation Drills

1. An AP has first term 2, common difference 3. Verify that it also satisfies the recurrence $a_n = 2a_{n-1} - a_{n-2}$.

Answer: Verified: $a_n = 3n - 1$ satisfies $a_n = 2a_{n-1} - a_{n-2}$.

Method: $2a_{n-1} - a_{n-2} = 2(3(n-1)-1) - (3(n-2)-1) = 2(3n-4) - (3n-7) = 6n-8-3n+7 = 3n-1 = a_n$.
□

Investigate further: Every AP satisfies $a_n = 2a_{n-1} - a_{n-2}$ (characteristic equation $x^2 - 2x + 1 = (x-1)^2 = 0$, a repeated root at 1) — exactly Day 4’s C7 result, now verified directly on a concrete AP.

2. Using $(1-x)^{-1} = \sum x^k$ and $x = \frac{1}{4}$, evaluate $\sum_{k=0}^{\infty} \left(\frac{1}{4}\right)^k$.

- A) $\frac{4}{3}$
B) $\frac{3}{4}$
C) 4
D) $\frac{1}{4}$

Answer: A) $\frac{4}{3}$

Method: $(1-x)^{-1}$ at $x = \frac{1}{4}$: $\left(\frac{3}{4}\right)^{-1} = \frac{4}{3}$.

Investigate further: Day 2’s identity plus Day 3’s substitution technique, combined exactly as in Day 3’s own B10/C7.

3. A recurrence has characteristic roots $\frac{1}{2}$ and $\frac{1}{3}$, with no forcing term. Does the sequence converge, and to what?
- A) Converges to 0.
B) Converges to $\frac{1}{2} + \frac{1}{3} = \frac{5}{6}$.
C) Diverges.

D) Oscillates without converging.

Answer: A) Converges to 0.

Method: Both $(\frac{1}{2})^n \rightarrow 0$ and $(\frac{1}{3})^n \rightarrow 0$; with no forcing term, $a_n = A(\frac{1}{2})^n + B(\frac{1}{3})^n \rightarrow 0$.

Investigate further: If a constant forcing term were bolted on instead — say $a_n = \frac{5}{6}a_{n-1} - \frac{1}{6}a_{n-2} + k$ — the limit would shift from 0 to whatever fixed point that forcing term produces (Day 4 C5's territory), not stay at 0. The “no forcing term” condition here is doing real work, not just decoration.

4. A sequence satisfies $a_{n+1} = \frac{1}{2-a_n}$. Find its fixed point(s), and determine whether $a_1 = 1$ leads to a constant sequence.

Answer: Fixed point $x = 1$ (repeated); yes, $a_1 = 1$ gives a constant sequence.

Method: $x = \frac{1}{2-x} \Rightarrow x(2-x) = 1 \Rightarrow 2x - x^2 = 1 \Rightarrow x^2 - 2x + 1 = 0 \Rightarrow (x-1)^2 = 0 \Rightarrow x = 1$ (unique, repeated). Since $a_1 = 1$ is exactly this fixed point, $a_2 = \frac{1}{2-1} = 1 = a_1$, and the sequence stays constant.

Investigate further: A rational (Möbius-style) map with a REPEATED fixed point behaves differently from Day 5's period-3/period-4 examples, which had no repeated fixed points — worth noticing the distinction.

5. The sum of a binomial row, $\sum_k \binom{n}{k}$, and the sum of an AP's terms, $\sum_k a_k$, are both examples of what general mathematical operation?

- A) Summing a finite sequence.
- B) They are unrelated concepts.
- C) Both are geometric series.
- D) Both require calculus.

Answer: A) Summing a finite sequence.

Method: Both are instances of $\sum_k(\text{terms})$ — the underlying operation (summation) is identical, even though the term-generating rules (binomial coefficients vs. an arithmetic rule) are completely different.

Investigate further: Recognising the SAME general operation underneath different-looking notation is a genuine synthesis skill, distinct from recognising when two things are actually unrelated (see B10).

6. In the alternating multiply-subtract process (Day 6), the shift $b_n = a_n - \frac{c}{r-1}$ turns Alice's map into exactly which type of sequence from Day 3?

- A) A GP with ratio r .
- B) An AP with difference r .
- C) A GP with ratio c .
- D) Neither.

Answer: A) A GP with ratio r .

Method: $b_n = a_n - \frac{c}{r-1}$ satisfies $b_{n+1} = rb_n$ exactly (Day 6's derivation, identical in form to Day 4's shift trick).

Investigate further: This shift trick relies on the subtracted term being a genuine constant — D3 later today asks what breaks if it isn't (a term $c_n = cn$ that grows every turn instead), and the answer is that a single fixed constant L can no longer absorb a moving target: Day 4 C5's particular-solution machinery, not this shift, is what's actually needed then.

7. The series $(1-x)^{-1} = \sum x^k$ is the closed form of which simple recurrence? State the recurrence and its characteristic root.

Answer: $a_k = x \cdot a_{k-1}$, a first-order GP recurrence with (single) characteristic root x .

Method: Since $a_k = x^k$, clearly $a_k = x \cdot x^{k-1} = x \cdot a_{k-1}$.

Investigate further: Every term of a geometric series is itself the closed-form solution of the simplest possible recurrence — Day 3 and Day 4 were secretly describing the same object from two different angles all along.

8. Which of Day 1's arithmetic sequences is also strictly monotonic (Day 5's sense) for every valid common difference $d \neq 0$?

- A) Every AP with $d \neq 0$ is monotonic.
- B) Only APs with $d > 0$.
- C) No AP is ever monotonic.
- D) Only APs with integer d .

Answer: A) Every AP with $d \neq 0$ is monotonic.

Method: $a_{n+1} - a_n = d$ always, a fixed nonzero sign — increasing if $d > 0$, decreasing if $d < 0$, with no exceptions.

Investigate further: Unlike Day 5's more exotic sequences (which needed real proof to establish monotonicity), an AP's monotonicity is immediate from its very definition.

9. A GP has $|r| > 1$. Is the sequence bounded?

- A) No — it diverges in magnitude, unbounded.
- B) Yes, always bounded.
- C) Bounded only if $a > 0$.
- D) Bounded only if r is an integer.

Answer: A) No, unbounded.

Method: $|a_n| = |a||r|^n \rightarrow \infty$ when $|r| > 1$, regardless of the sign of a or r .

Investigate further: This is the direct opposite of Day 3's convergence condition $|r| < 1$ — the two cases ($|r| < 1$ vs. $|r| > 1$) are exact mirror images.

10. Day 6's " $a_k = 1$ for all k " construction and Day 2's binomial row-sum fact $\sum_k \binom{n}{k} = 2^n$ are both about sums. Are these two facts related, or independent observations?

Answer: Independent observations.

Method: Day 6's construction concerns DIVISIBILITY of a running (incrementally-built) sum by its own index; Day 2's row sum concerns the TOTAL of a fixed, complete row of binomial coefficients. Different objects, different questions, no structural connection beyond both being "a sum."

Investigate further: Not every pair of similar-sounding facts from different days is actually related — correctly identifying when two ideas are genuinely independent, rather than forcing a false connection, is exactly as important as spotting real synthesis (contrast with B5).

Section C Substitution & Structure

1. An AP and a GP both have first term 3; the AP has $d = 4$, the GP has $r = 2$. After how many terms (if ever) could they next coincide?
 - A) They never coincide again.
 - B) At term 4.
 - C) At term 6.
 - D) At term 10.

Answer: A) They never coincide again.

Method: AP: 3, 7, 11, 15, 19, 23, ... GP: 3, 6, 12, 24, 48, ... Comparing term by term after the first: none match, and once the GP overtakes the AP (by term 4: $24 > 15$), the exponentially-growing GP stays ahead of the linearly-growing AP forever.

Investigate further: This is Day 1's B10 insight (GP eventually and permanently overtakes AP) turned into a "do they ever meet again" question.

2. Compare the growth of $\sum_{k=1}^n a_k$ where $a_k = 1$ for all k (Day 6's construction) against $\sum_{k=0}^n \binom{n}{k}$ (Day 2's row sum) as $n \rightarrow \infty$. Which grows faster?
 - A) The binomial row sum grows faster (exponentially vs. linearly).
 - B) They grow at the same rate.
 - C) The Day 6 sum grows faster.
 - D) Neither grows — both are constant.

Answer: A) The binomial row sum grows faster.

Method: Day 6's sum is $\sum_{k=1}^n 1 = n$ (linear). Day 2's row sum is 2^n (exponential). Exponential growth eventually and permanently outpaces linear growth.

Investigate further: The exact same "exponential beats linear eventually" theme as C1 and Day 1's B10, now applied to two different sums rather than two sequences.

3. A recurrence $a_n = pa_{n-1} + qa_{n-2}$ has both characteristic roots equal to a Day 6-style fixed point $\frac{c}{r-1}$. If $\frac{c}{r-1} = 1$, what does Day 4's result about repeated roots equal to 1 tell us about the long-term behaviour of (a_n) ?
 - A) Linear growth in n (effectively an AP).
 - B) Exponential growth.

- C) Convergence to 0.
D) Periodic oscillation.

Answer: A) Linear growth in n .

Method: A repeated characteristic root of exactly 1 gives closed form $a_n = (A + Bn)(1)^n = A + Bn$ — linear in n (Day 4's C7 result), regardless of how that root of 1 arose.

Investigate further: This shows Day 4's C7 result is really about the VALUE of the repeated root ($= 1$), not about how the recurrence was originally motivated (whether directly given, or arising from a Day 6-style fixed point).

4. Evaluate $\sum_{k=0}^{\infty} (k+1) \left(\frac{1}{3}\right)^k$ using Day 2's identity $(1-x)^{-2} = \sum_{k=0}^{\infty} (k+1)x^k$.

- A) $\frac{9}{4}$
B) $\frac{3}{2}$
C) $\frac{4}{9}$
D) $\frac{2}{3}$

Answer: A) $\frac{9}{4}$

Method: $(1-x)^{-2}$ at $x = \frac{1}{3}$: $\left(\frac{2}{3}\right)^{-2} = \left(\frac{3}{2}\right)^2 = \frac{9}{4}$.

Investigate further: Same technique as B2 and Day 3's C8, now with the “ $(k+1)$ -weighted” series instead of the plain geometric series.

5. A Möbius map (Day 5) is used as the combined operation in an alternating writing process (Day 6). If the map has order exactly 3 under iteration, what is guaranteed about EVERY starting value (where the map is defined)?
- A) Every starting value returns to itself after exactly 3 applications.
B) Only some starting values return; others diverge.
C) The map converges to a fixed point instead.
D) Nothing general can be said.

Answer: A) Every starting value returns to itself after exactly 3 applications.

Method: A Möbius transformation of finite order m (as a transformation) satisfies $f^m = \text{identity}$ EVERYWHERE it's defined, not just at special points — this is a property of the transformation itself, not of any particular starting value.

Investigate further: Contrast Day 5's C1/C2/D2: those specific maps were shown to have order 4 and 3 respectively by direct computation from ONE starting value, but the underlying fact (guaranteed for every valid starting point) is what this question makes explicit.

6. For a GP with ratio r , for which values of r is the sequence BOTH monotonic AND convergent?
- A) $0 \leq r \leq 1$.
B) $-1 < r < 1$.

C) Only $r = 1$.

D) $r > 1$ only.

Answer: A) $0 \leq r \leq 1$.

Method: For $0 < r < 1$: decreasing toward 0 (monotonic, convergent). $r = 1$: constant (trivially both). $r = 0$: the sequence is $a, 0, 0, 0, \dots$, trivially monotonic (non-increasing) and convergent. For $-1 < r < 0$: convergent to 0 but NOT monotonic (alternates sign). For $|r| \geq 1$ (excluding $r = 1$): not convergent (Day 3/B9).

Investigate further: The boundary cases ($r = 0$ and $r = 1$) are exactly where “monotonic” and “convergent” stop being in tension with each other — everywhere else in $(-1, 1)$, oscillation (for negative r) breaks monotonicity even though convergence still holds.

7. Day 6’s two-branch search ($u(r^k - 1) = -c$) and Day 4’s root-magnitude convergence condition are both examples of what general principle about a term r^k (or r^n)?

A) Whether $|r| < 1$, $= 1$, or > 1 entirely determines the long-term size of r^k — shrinking to 0, staying constant, or blowing up.

B) Exponential terms are always eventually periodic.

C) Exponential terms never affect convergence.

D) There is no shared principle.

Answer: A) $|r| < 1, = 1, > 1$ entirely determines the size of r^k .

Method: Both Day 6’s search (where $|u| = \frac{c}{r^k - 1} \rightarrow 0$ because $r \geq 2 > 1$) and Day 4’s convergence condition (roots need $|r| < 1$ to shrink to 0) are direct consequences of this single fact about exponentials.

Investigate further: The trichotomy has a fourth corner the method doesn’t mention: complex r with $|r| = 1$ exactly. That’s Day 5’s Möbius maps of exact order 3 or 4 (C1, D2) — neither shrinking nor growing, just orbiting forever, which is exactly what “staying constant” looks like once magnitude stops being the whole story.

8. Which set of results from this week are all secretly governed by the SAME underlying idea — long-term behaviour controlled by whether a key magnitude is < 1 , $= 1$, or > 1 ?

A) Geometric series convergence (Day 3); recurrence convergence via characteristic roots (Day 4); GP boundedness (today’s B9).

B) Arithmetic sequences, binomial coefficients, integer means.

C) All seven days equally.

D) None of the sheet’s content shares a common idea.

Answer: A) Geometric series convergence, recurrence convergence, GP boundedness.

Method: All three are direct restatements of “the size of r^n as $n \rightarrow \infty$ depends entirely on whether $|r|$ is less than, equal to, or greater than 1” — Day 3’s $|r| < 1$ convergence condition, Day 4’s root-magnitude convergence condition, and today’s B9 boundedness fact are the same underlying idea in three different costumes.

Investigate further: Naming this shared idea explicitly is the whole point of a synthesis day — the individual techniques were taught separately across the week precisely so that noticing they’re the

| same idea, here, lands with real weight.

Section D Challenge

1. For each positive integer $n \geq 4$, define an n -necklace to be a circular arrangement of n (not necessarily distinct) positive integers such that the product of every 4 neighbouring integers equals n . Determine the number of integers n in the range $4 \leq n \leq 1000$ for which it is possible to form an n -necklace. (after BMO1 2019 Q2)

Answer: 252

Method: Let $d = \gcd(4, n)$. Since the product of every 4 consecutive terms is constant, $a_i = a_{i+4}$ (indices mod n) for every i — so a_i depends only on $i \bmod d$. Write $x_0, \dots, x_{d-1}$ for the values on each residue class. Since $d \mid 4$ always (a gcd divides both its arguments), every window of 4 consecutive terms covers each x_j exactly $4/d$ times, regardless of where the window starts. So the constant product is $(x_0 x_1 \cdots x_{d-1})^{4/d}$, and setting this equal to n : a valid n -necklace exists exactly when n is a perfect $(4/d)$ -th power, where $d = \gcd(4, n)$.

Case $4 \mid n$ ($d = 4$): need n to be a perfect 1st power — always true. Every multiple of 4 works.

Case $n \equiv 2 \pmod{4}$ ($d = 2$): need n to be a perfect square. But every perfect square is $\equiv 0$ or $1 \pmod{4}$ (even² $\equiv 0$, odd² $\equiv 1$), never $\equiv 2$ — so this case is NEVER possible.

Case n odd ($d = 1$): need n to be a perfect 4th power (automatically odd, since an odd 4th root gives an odd 4th power).

Counting $4 \leq n \leq 1000$: multiples of 4 number 250. Odd perfect 4th powers in range: $3^4 = 81$ and $5^4 = 625$ (checking $1^4 = 1 < 4$ excluded, $7^4 = 2401 > 1000$ excluded) — 2 more. Total: $250 + 2 = 252$.

Investigate further: The real BMO1 question (2019 Q2) uses windows of 3 instead of 4, over the range $3 \leq n \leq 2018$. Since 3 is prime, $\gcd(3, n) \in \{1, 3\}$ only, giving just two cases (multiples of 3 always work; non-multiples need n to be a perfect cube) — the real answer is 679. Using 4 instead of a prime window introduces the extra $n \equiv 2 \pmod{4}$ case, which turns out to NEVER contribute — an example of how changing one number in a BMO1 problem can add real extra casework, not just bigger numbers.

2. Prove that for any starting value a_1 (a positive integer), there exists a strictly increasing sequence of positive integers $a_1 < a_2 < \cdots < a_n$, of any desired length n , such that every partial sum S_k is divisible by k .

Answer: Proof by induction using Day 6's construction.

Method: Base case: a_1 given, trivially satisfies $S_1 = a_1$ divisible by 1. Inductive step: given a valid $a_1 < \cdots < a_k$ with S_k divisible by k , Day 6's C7 argument shows some $a_{k+1} \equiv -S_k \pmod{k+1}$ makes S_{k+1} divisible by $k+1$ — and since there are infinitely many integers in that residue class (differing by multiples of $k+1$), we can always choose one that ALSO exceeds a_k (just add enough multiples of $k+1$ to the smallest valid residue). This gives a valid $a_{k+1} > a_k$. By induction, a strictly increasing valid sequence of any desired length n can always be built, starting from any a_1 . $\square$

Investigate further: This upgrades Day 6's C7 ("a valid next term always exists") to also guarantee it can be chosen larger than the previous term — the same idea, with one extra degree of freedom (choosing among infinitely many equivalent residues) doing the extra work.

3. In Day 6's alternating recurrence, suppose the subtracted term is not a fixed constant c but grows every turn, $c_n = cn$. Does the shift-to-fixed-point trick (a single constant $L = \frac{c}{r-1}$) still work directly?

A) No — a single fixed constant L cannot absorb a forcing term that changes every step;

a particular-solution technique (Day 4's C5) matching the forcing term's form would be needed instead.

B) Yes, exactly the same L still works.

C) The process becomes undefined.

D) It only works if $r = 1$.

Answer: A) No — a particular-solution technique is needed instead.

Method: The equation $L = rL - c_n$ would need L to depend on n (since c_n changes every step), contradicting the requirement that L be a single fixed constant — exactly Day 4's C5 situation (a non-constant forcing term defeats the simple constant-shift trick).

Investigate further: Recognising that Day 4's C5 insight (forcing term type determines the fix needed) applies just as much to Day 6's alternating process as to Day 4's own recurrences is real cross-day transfer, not just pattern-completion.

4. A sequence is defined by $a_1 = 1$ and, for $n \geq 1$, a_{n+1} is the SMALLEST positive integer greater than a_n such that S_{n+1} is divisible by $n + 1$. Find $a_1, \dots, a_5$, and conjecture (with justification) a general closed form for a_n .

Answer: $a_1, \dots, a_5 = 1, 3, 5, 7, 9$; conjectured closed form $a_n = 2n - 1$.

Method: $a_1 = 1$ ($S_1 = 1$). For a_2 : try 2 first ($S_2 = 3$, not divisible by 2); try 3 ($S_2 = 4$, divisible by 2) — so $a_2 = 3$. For a_3 : try 4 ($S_3 = 8$, not divisible by 3); try 5 ($S_3 = 9$, divisible by 3) — $a_3 = 5$. For a_4 : try 6 ($S_4 = 15$, not divisible by 4); try 7 ($S_4 = 16$, divisible by 4) — $a_4 = 7$. For a_5 : try 8 ($S_5 = 24$, not divisible by 5); try 9 ($S_5 = 25$, divisible by 5) — $a_5 = 9$.

The pattern 1, 3, 5, 7, 9 is exactly the consecutive-odd-numbers sequence from Day 6's B1, where $S_k = k^2$ always — and $k^2/k = k$ is always an integer, so this sequence is always valid. The conjecture $a_n = 2n - 1$ is consistent with every computed term, and the fact that the “next even integer” candidate fails at every single step (checked above for $n = 2, 3, 4, 5$) strongly suggests it is genuinely the SMALLEST valid choice at each stage, not a coincidence.

Investigate further: This is Day 6's B1 construction (built there by deliberate choice) rediscovered here as the UNIQUE greedy outcome of a simple local rule (“smallest integer bigger than the last term, keeping the mean an integer”) — a satisfying full-circle moment to end the week on.

5. Looking back across all seven days: which day's toolkit would be LEAST directly useful for tackling D1's n -necklace problem?

A) Day 2 (Binomial & Trinomial Expansion) — D1's argument uses periodicity and perfect-power number theory, not binomial coefficients.

B) Day 5 (Behaviour of Sequences).

C) Day 1 (Arithmetic Sequences).

D) All seven days are equally essential to D1.

Answer: A) Day 2 (Binomial & Trinomial Expansion).

Method: D1's proof rests entirely on periodicity ($a_i = a_{i+4}$, an orbit-structure argument closely related to Day 5's periodicity toolkit) and number-theoretic perfect-power reasoning — binomial coefficients never enter the argument at any point.

Investigate further: Recognising which tools are genuinely load-bearing for a given problem (and which aren't) is itself a mature strategic skill — not every day's toolkit is meant to be forced into every problem, and knowing that saves real time under exam pressure.

Top 5 Patterns — Worksheet #7

- 1. Identify which day's toolkit a question wants before diving into algebra.** Misapplying Day 4's heavy machinery to a Day 1/3 problem (or vice versa) wastes real time. (see A7, B1, B3, B6, B8, C1, C6)
- 2. The single unifying idea behind Days 3, 4, and 5: whether a magnitude is < 1 , $= 1$, or > 1 controls everything.** (see B3, B9, C6, C7, C8)
- 3. Series-manipulation from Day 2 ($(1-x)^{-m}$ substitutions) combines directly with Day 3's convergence toolkit.** Always look for this bridge on an “evaluate this infinite series” question. (see B2, C4)
- 4. Day 6's shift-to-fixed-point trick is Day 3's GP toolkit, applied one level up via Day 4's shift technique.** (see B6, B7, D3)
- 5. Not every technique is relevant to every problem.** Identifying which day's toolkit is genuinely load-bearing (and which isn't) is itself a tested skill on a synthesis day. (see C2, C3, D5)

Common Traps — Worksheet #7

- 1. Reaching for Day 4's characteristic equation on a recurrence that's secretly just an AP or GP in disguise.** (see B1, B6)
- 2. Assuming every sum/series question is “the same kind of sum.”** Day 2's binomial row sums and Day 6's integer-mean partial sums are structurally unrelated, despite superficial similarity. (see B10, C2)
- 3. Forgetting that convergence and boundedness are different properties.** $|r| > 1$ makes a GP unbounded, not just “not converging to a specific value.” (see B9)
- 4. Assuming a repeated-root or fixed-point coincidence is special/impossible rather than checking what it actually implies.** (see C3)
- 5. In a full synthesis proof, skipping the general case-by-case argument in favour of pattern-matching from a few small examples.** A genuine BMO1 proof needs the general argument. (see D1)

“Days 3, 4 and 5 were the same question asked three times: is the magnitude below one, at one, or above it.”

Found a mistake or want to contribute a sheet?

Visit the open-source project at github.com/The-CerealDev/speed-maths