

# Types of Morphisms in Bicategories

The Clowder Project Authors

July 29, 2025

In this chapter, we study special kinds of morphisms in bicategories:

1. *Monomorphisms and Epimorphisms in Bicategories* ([Sections 14.1](#) and [14.2](#)). There is a large number of different notions capturing the idea of a “monomorphism” or of an “epimorphism” in a bicategory.

Arguably, the notion that best captures these concepts is that of a *pseudomononic morphism* ([Definition 14.1.10.1.1](#)) and of a *pseudoepic morphism* ([Definition 14.2.10.1.1](#)), although the other notions introduced in [Sections 14.1](#) and [14.2](#) are also interesting on their own.

## Contents

|  |           |
|--|-----------|
| <b>14.1 Monomorphisms in Bicategories</b>              | <b>2</b>  |
| 14.1.1 Representably Faithful Morphisms                | 2         |
| 14.1.2 Representably Full Morphisms                    | 3         |
| 14.1.3 Representably Fully Faithful Morphisms          | 4         |
| 14.1.4 Morphisms Representably Faithful on Cores       | 5         |
| 14.1.5 Morphisms Representably Full on Cores           | 6         |
| 14.1.6 Morphisms Representably Fully Faithful on Cores | 6         |
| 14.1.7 Representably Essentially Injective Morphisms   | 8         |
| 14.1.8 Representably Conservative Morphisms            | 8         |
| 14.1.9 Strict Monomorphisms                            | 9         |
| 14.1.10 Pseudomononic Morphisms                        | 9         |
| <b>14.2 Epimorphisms in Bicategories</b>               | <b>11</b> |
| 14.2.1 Corepresentably Faithful Morphisms              | 11        |
| 14.2.2 Corepresentably Full Morphisms                  | 12        |
| 14.2.3 Corepresentably Fully Faithful Morphisms        | 13        |
| 14.2.4 Morphisms Corepresentably Faithful on Cores     | 14        |

|          |  |           |
|----------|--|-----------|
| 14.2.5   | Morphisms Corepresentably Full on Cores.....           | 15        |
| 14.2.6   | Morphisms Corepresentably Fully Faithful on Cores..... | 16        |
| 14.2.7   | Corepresentably Essentially Injective Morphisms.....   | 17        |
| 14.2.8   | Corepresentably Conservative Morphisms.....            | 17        |
| 14.2.9   | Strict Epimorphisms.....                               | 18        |
| 14.2.10  | Pseudoepic Morphisms.....                              | 18        |
| <b>A</b> | <b>Other Chapters.....</b>                             | <b>20</b> |

## 14.1 Monomorphisms in Bicategories

### 14.1.1 Representably Faithful Morphisms

Let  $C$  be a bicategory.

**Definition 14.1.1.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably faithful**<sup>1</sup> if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is faithful.

**Remark 14.1.1.1.2.** In detail,  $f$  is representably faithful if, for all diagrams in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \downarrow \downarrow \downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

**Example 14.1.1.1.3.** Here are some examples of representably faithful morphisms.

1. *Representably Faithful Morphisms in  $\text{Cats}_2$ .* The representably faithful morphisms in  $\text{Cats}_2$  are precisely the faithful functors; see [Categories, Item 2](#) of [Definition 11.6.1.1.2](#).
2. *Representably Faithful Morphisms in  $\mathbf{Rel}$ .* Every morphism of  $\mathbf{Rel}$  is representably faithful; see [Relations](#), ?? of ??.

<sup>1</sup>Further Terminology: Also called simply a **faithful morphism**, based on [Item 1](#) of

### 14.1.2 Representably Full Morphisms

Let  $C$  be a bicategory.

**Definition 14.1.2.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably full**<sup>2</sup> if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is full.

**Remark 14.1.2.1.2.** In detail,  $f$  is representably full if, for each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: f \circ \phi \Rightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

**Example 14.1.2.1.3.** Here are some examples of representably full morphisms.

1. *Representably Full Morphisms in  $\text{Cats}_2$ .* The representably full morphisms in  $\text{Cats}_2$  are precisely the full functors; see **Categories**, ?? of **Definition 11.6.2.1.2**.
2. *Representably Full Morphisms in  $\mathbf{Rel}$ .* The representably full morphisms in  $\mathbf{Rel}$  are characterised in **Relations**, ?? of ??.

---

**Definition 14.1.1.1.3.**

<sup>2</sup>*Further Terminology:* Also called simply a **full morphism**, based on **Item 1** of

### 14.1.3 Representably Fully Faithful Morphisms

Let  $C$  be a bicategory.

**Definition 14.1.3.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably fully faithful**<sup>3</sup> if the following equivalent conditions are satisfied:

1. The 1-morphism  $f$  is representably faithful (Definition 14.1.1.1.1) and representably full (Definition 14.1.2.1.1).
2. For each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is fully faithful.

**Remark 14.1.3.1.2.** In detail,  $f$  is representably fully faithful if the conditions in Definition 14.1.1.1.2 and Definition 14.1.2.1.2 hold:

1. For all diagrams in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: f \circ \phi \Rightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

---

Definition 14.1.2.1.3.

<sup>3</sup>*Further Terminology:* Also called simply a **fully faithful morphism**, based on Item 1 of Definition 14.1.3.1.3.

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

**Example 14.1.3.1.3.** Here are some examples of representably fully faithful morphisms.

1. *Representably Fully Faithful Morphisms in  $\mathbf{Cats}_2$ .* The representably fully faithful morphisms in  $\mathbf{Cats}_2$  are precisely the fully faithful functors; see [Categories, Item 6](#) of [Definition 11.6.3.1.2](#).
2. *Representably Fully Faithful Morphisms in  $\mathbf{Rel}$ .* The representably fully faithful morphisms of  $\mathbf{Rel}$  coincide ([Relations](#), ?? of ??) with the representably full morphisms in  $\mathbf{Rel}$ , which are characterised in [Relations](#), ?? of ??.

#### 14.1.4 Morphisms Representably Faithful on Cores

Let  $C$  be a bicategory.

**Definition 14.1.4.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably faithful on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by  $f$  is faithful.

**Remark 14.1.4.1.2.** In detail,  $f$  is representably faithful on cores if, for all diagrams in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

### 14.1.5 Morphisms Representably Full on Cores

Let  $C$  be a bicategory.

**Definition 14.1.5.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably full on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by  $f$  is full.

**Remark 14.1.5.1.2.** In detail,  $f$  is representably full on cores if, for each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: f \circ \phi \xRightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xRightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

### 14.1.6 Morphisms Representably Fully Faithful on Cores

Let  $C$  be a bicategory.

**Definition 14.1.6.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably fully faithful on cores** if the following equivalent conditions are satisfied:

1. The 1-morphism  $f$  is representably faithful on cores ([Definition 14.1.5.1.1](#)) and representably full on cores ([Definition 14.1.4.1.1](#)).

2. For each  $X \in \text{Obj}(C)$ , the functor

$$f_* : \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by  $f$  is fully faithful.

**Remark 14.1.6.1.2.** In detail,  $f$  is representably fully faithful on cores if the conditions in [Definition 14.1.4.1.2](#) and [Definition 14.1.5.1.2](#) hold:

1. For all diagrams in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta : f \circ \phi \xRightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-isomorphism

$$\alpha : \phi \xRightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

### 14.1.7 Representably Essentially Injective Morphisms

Let  $C$  be a bicategory.

**Definition 14.1.7.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably essentially injective** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is essentially injective.

**Remark 14.1.7.1.2.** In detail,  $f$  is representably essentially injective if, for each pair of morphisms  $\phi, \psi: X \rightrightarrows A$  of  $C$ , the following condition is satisfied:

( $\star$ ) If  $f \circ \phi \cong f \circ \psi$ , then  $\phi \cong \psi$ .

### 14.1.8 Representably Conservative Morphisms

Let  $C$  be a bicategory.

**Definition 14.1.8.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably conservative** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is conservative.

**Remark 14.1.8.1.2.** In detail,  $f$  is representably conservative if, for each pair of morphisms  $\phi, \psi: X \rightrightarrows A$  and each 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of  $C$ , if the 2-morphism

$$\text{id}_f \star \alpha: f \circ \phi \Rightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \parallel \\ \text{id}_f \star \alpha \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

is a 2-isomorphism, then so is  $\alpha$ .



### 14.1.9 Strict Monomorphisms

Let  $C$  be a bicategory.

**Definition 14.1.9.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is a **strict monomorphism** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is injective on objects, i.e. its action on objects

$$f_*: \text{Obj}(\text{Hom}_C(X, A)) \rightarrow \text{Obj}(\text{Hom}_C(X, B))$$

is injective.

**Remark 14.1.9.1.2.** In detail,  $f$  is a strict monomorphism in  $C$  if, for each diagram in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if  $f \circ \phi = f \circ \psi$ , then  $\phi = \psi$ .

**Example 14.1.9.1.3.** Here are some examples of strict monomorphisms.

1. *Strict Monomorphisms in  $\text{Cats}_2$ .* The strict monomorphisms in  $\text{Cats}_2$  are precisely the functors which are injective on objects and injective on morphisms; see [Categories, Item 1](#) of [Definition 11.7.2.1.2](#).
2. *Strict Monomorphisms in  $\mathbf{Rel}$ .* The strict monomorphisms in  $\mathbf{Rel}$  are characterised in [Relations, ??](#).

### 14.1.10 Pseudomonic Morphisms

Let  $C$  be a bicategory.

**Definition 14.1.10.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **pseudomonic** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is pseudomonic.

**Remark 14.1.10.1.2.** In detail, a 1-morphism  $f: A \rightarrow B$  of  $C$  is pseudomonic if it satisfies the following conditions:

1. For all diagrams in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then  $\alpha = \beta$ .

2. For each  $X \in \mathrm{Obj}(C)$  and each 2-isomorphism

$$\beta: f \circ \phi \xRightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xRightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

**Proposition 14.1.10.1.3.** Let  $f: A \rightarrow B$  be a 1-morphism of  $C$ .

1. *Characterisations.* The following conditions are equivalent:

- (a) The morphism  $f$  is pseudomonic.
- (b) The morphism  $f$  is representably full on cores and representably faithful.

(c) We have an isocomma square of the form

$$A \cong^{eq.} A \times_B A, \quad \begin{array}{ccc} A & \xrightarrow{id_A} & A \\ id_A \downarrow & \nearrow \text{dashed} & \downarrow F \\ A & \xrightarrow{F} & B \end{array}$$

in  $C$  up to equivalence.

2. *Interaction With Cotensors.* If  $C$  has cotensors with  $\mathbb{1}$ , then the following conditions are equivalent:

- (a) The morphism  $f$  is pseudomonic.
- (b) We have an isocomma square of the form

$$A \cong^{eq.} A \times_{\mathbb{1} \pitchfork F} B, \quad \begin{array}{ccc} A & \hookrightarrow & \mathbb{1} \pitchfork A \\ F \downarrow & \nearrow \text{dashed} & \downarrow \mathbb{1} \pitchfork F \\ B & \hookrightarrow & \mathbb{1} \pitchfork B \end{array}$$

in  $C$  up to equivalence.

*Proof.* **Item 1**, Characterisations: Omitted.

**Item 2**, Interaction With Cotensors: Omitted. □

## 14.2 Epimorphisms in Bicategories

### 14.2.1 Corepresentably Faithful Morphisms

Let  $C$  be a bicategory.

**Definition 14.2.1.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably faithful** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is faithful.

**Remark 14.2.1.1.2.** In detail,  $f$  is corepresentably faithful if, for all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \parallel \downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

**Example 14.2.1.1.3.** Here are some examples of corepresentably faithful morphisms.

1. *Corepresentably Faithful Morphisms in  $\mathbf{Cats}_2$ .* The corepresentably faithful morphisms in  $\mathbf{Cats}_2$  are characterised in [Categories, Item 5](#) of [Definition 11.6.1.1.2](#).
2. *Corepresentably Faithful Morphisms in  $\mathbf{Rel}$ .* Every morphism of  $\mathbf{Rel}$  is corepresentably faithful; see [Relations](#), ?? of ??.

## 14.2.2 Corepresentably Full Morphisms

Let  $C$  be a bicategory.

**Definition 14.2.2.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably full** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is full.

**Remark 14.2.2.1.2.** In detail,  $f$  is corepresentably full if, for each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: \phi \circ f \Rightarrow \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \Downarrow \beta \\ \xrightarrow{\psi \circ f} \end{array} X$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \Downarrow \alpha \\ \xrightarrow{\psi} \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \Downarrow \alpha \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \Downarrow \beta \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

**Example 14.2.2.1.3.** Here are some examples of corepresentably full morphisms.

1. *Corepresentably Full Morphisms in  $\mathbf{Cats}_2$ .* The corepresentably full morphisms in  $\mathbf{Cats}_2$  are characterised in [Categories](#), [Item 7](#) of [Definition 11.6.2.1.2](#).
2. *Corepresentably Full Morphisms in  $\mathbf{Rel}$ .* The corepresentably full morphisms in  $\mathbf{Rel}$  are characterised in [Relations](#), ?? of ??.

### 14.2.3 Corepresentably Fully Faithful Morphisms

Let  $C$  be a bicategory.

**Definition 14.2.3.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably fully faithful**<sup>4</sup> if the following equivalent conditions are satisfied:

1. The 1-morphism  $f$  is corepresentably full ([Definition 14.2.2.1.1](#)) and corepresentably faithful ([Definition 14.2.1.1.1](#)).
2. For each  $X \in \mathbf{Obj}(C)$ , the functor

$$f^*: \mathbf{Hom}_C(B, X) \rightarrow \mathbf{Hom}_C(A, X)$$

given by precomposition by  $f$  is fully faithful.

**Remark 14.2.3.1.2.** In detail,  $f$  is corepresentably fully faithful if the conditions in [Definition 14.2.1.1.2](#) and [Definition 14.2.2.1.2](#) hold:

1. For all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \downarrow \downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then  $\alpha = \beta$ .

---

<sup>4</sup>*Further Terminology:* Corepresentably fully faithful morphisms have also been called **lax epimorphisms** in the literature (e.g. in [\[Ad  01\]](#)), though we will always use the name “corepresentably fully faithful morphism” instead in this work.

2. For each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: \phi \circ f \Rightarrow \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

**Example 14.2.3.1.3.** Here are some examples of corepresentably fully faithful morphisms.

1. *Corepresentably Fully Faithful Morphisms in  $\mathbf{Cats}_2$ .* The fully faithful epimorphisms in  $\mathbf{Cats}_2$  are characterised in [Categories, Item 10](#) of [Definition 11.6.3.1.2](#).
2. *Corepresentably Fully Faithful Morphisms in  $\mathbf{Rel}$ .* The corepresentably fully faithful morphisms of  $\mathbf{Rel}$  coincide ([Relations, ??](#) of [??](#)) with the corepresentably full morphisms in  $\mathbf{Rel}$ , which are characterised in [Relations, ??](#) of [??](#).

## 14.2.4 Morphisms Corepresentably Faithful on Cores

Let  $C$  be a bicategory.

**Definition 14.2.4.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably faithful on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by  $f$  is faithful.

**Remark 14.2.4.1.2.** In detail,  $f$  is corepresentably faithful on cores if, for all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

### 14.2.5 Morphisms Corepresentably Full on Cores

Let  $C$  be a bicategory.

**Definition 14.2.5.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably full on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by  $f$  is full.

**Remark 14.2.5.1.2.** In detail,  $f$  is corepresentably full on cores if, for each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ f \xRightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xRightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

### 14.2.6 Morphisms Corepresentably Fully Faithful on Cores

Let  $C$  be a bicategory.

**Definition 14.2.6.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably fully faithful on cores** if the following equivalent conditions are satisfied:

1. The 1-morphism  $f$  is corepresentably full on cores (Definition 14.2.5.1.1) and corepresentably faithful on cores (Definition 14.2.1.1.1).
2. For each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by  $f$  is fully faithful.

**Remark 14.2.6.1.2.** In detail,  $f$  is corepresentably fully faithful on cores if the conditions in Definition 14.2.4.1.2 and Definition 14.2.5.1.2 hold:

1. For all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ f \xRightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xRightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$



### 14.2.7 Corepresentably Essentially Injective Morphisms

Let  $C$  be a bicategory.

**Definition 14.2.7.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably essentially injective** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is essentially injective.

**Remark 14.2.7.1.2.** In detail,  $f$  is corepresentably essentially injective if, for each pair of morphisms  $\phi, \psi: B \rightrightarrows X$  of  $C$ , the following condition is satisfied:

( $\star$ ) If  $\phi \circ f \cong \psi \circ f$ , then  $\phi \cong \psi$ .

### 14.2.8 Corepresentably Conservative Morphisms

Let  $C$  be a bicategory.

**Definition 14.2.8.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably conservative** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is conservative.

**Remark 14.2.8.1.2.** In detail,  $f$  is corepresentably conservative if, for each pair of morphisms  $\phi, \psi: B \rightrightarrows X$  and each 2-morphism

$$\alpha: \phi \xRightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of  $C$ , if the 2-morphism

$$\alpha \star \text{id}_f: \phi \circ f \xRightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \parallel \\ \alpha \star \text{id}_f \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

is a 2-isomorphism, then so is  $\alpha$ .

### 14.2.9 Strict Epimorphisms

Let  $C$  be a bicategory.

**Definition 14.2.9.1.1.** A 1-morphism  $f: A \rightarrow B$  is a **strict epimorphism** in  $C$  if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is injective on objects, i.e. its action on objects

$$f_*: \text{Obj}(\text{Hom}_C(B, X)) \rightarrow \text{Obj}(\text{Hom}_C(A, X))$$

is injective.

**Remark 14.2.9.1.2.** In detail,  $f$  is a strict epimorphism if, for each diagram in  $C$  of the form

$$A \xrightarrow{f} B \begin{matrix} \xrightarrow{\phi} \\ \xrightarrow{\psi} \end{matrix} X,$$

if  $\phi \circ f = \psi \circ f$ , then  $\phi = \psi$ .

**Example 14.2.9.1.3.** Here are some examples of strict epimorphisms.

1. *Strict Epimorphisms in  $\mathbf{Cats}_2$ .* The strict epimorphisms in  $\mathbf{Cats}_2$  are characterised in [Categories](#), [Item 1](#) of [Definition 11.7.3.1.2](#).
2. *Strict Epimorphisms in  $\mathbf{Rel}$ .* The strict epimorphisms in  $\mathbf{Rel}$  are characterised in [Relations](#), ??.

### 14.2.10 Pseudoepic Morphisms

Let  $C$  be a bicategory.

**Definition 14.2.10.1.1.** A 1-morphism  $f: A \rightarrow B$  of  $C$  is **pseudoepic** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is pseudomononic.

**Remark 14.2.10.1.2.** In detail, a 1-morphism  $f: A \rightarrow B$  of  $C$  is pseudoepic if it satisfies the following conditions:

1. For all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ f \xRightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xRightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

**Proposition 14.2.10.1.3.** Let  $f: A \rightarrow B$  be a 1-morphism of  $C$ .

1. *Characterisations.* The following conditions are equivalent:

- (a) The morphism  $f$  is pseudoepic.
- (b) The morphism  $f$  is corepresentably full on cores and corepresentably faithful.

(c) We have an isococcomma square of the form

$$B \overset{\text{eq.}}{\cong} B \overset{\leftrightarrow}{\coprod}_A B, \quad \begin{array}{ccc} B & \xleftarrow{\text{id}_B} & B \\ \text{id}_B \uparrow & \nearrow & \uparrow F \\ B & \xleftarrow{F} & A \end{array}$$

in  $\mathcal{C}$  up to equivalence.

*Proof.* **Item 1**, Characterisations: Omitted. □

## Appendices

### A Other Chapters

#### Preliminaries

1. **Introduction**
2. **A Guide to the Literature**

#### Sets

3. **Sets**
4. **Constructions With Sets**
5. **Monoidal Structures on the Category of Sets**
6. **Pointed Sets**
7. **Tensor Products of Pointed Sets**

#### Relations

8. **Relations**
9. **Constructions With Relations**

#### 10. **Conditions on Relations**

#### Categories

11. **Categories**
12. **Presheaves and the Yoneda Lemma**

#### Monoidal Categories

13. **Constructions With Monoidal Categories**

#### Bicategories

14. **Types of Morphisms in Bicategories**

#### Extra Part

15. **Notes**

## References

- [Adá+01] Jiří Adámek, Robert El Bashir, Manuela Sobral, and Jiří Velebil. “On Functors Which Are Lax Epimorphisms”. In: *Theory Appl. Categ.* 8 (2001), pp. 509–521. issn: 1201-561X (cit. on p. 13).