

# Sets

The Clowder Project Authors

July 29, 2025

0000 This chapter (will eventually) contain material on axiomatic set theory, as well as a couple other things.

## Contents

<b>3.1</b>	<b>Sets and Functions.....</b>	<b>1</b>
3.1.1	Functions .....	1
<b>3.2</b>	<b>The Enrichment of Sets in Classical Truth Values.....</b>	<b>4</b>
3.2.1	$(-2)$ -Categories .....	4
3.2.2	$(-1)$ -Categories .....	4
3.2.3	0-Categories.....	10
3.2.4	Tables of Analogies Between Set Theory and Category Theory .....	10
<b>A</b>	<b>Other Chapters .....</b>	<b>13</b>

## 0001 3.1 Sets and Functions

### 0002 3.1.1 Functions

#### 0003 DEFINITION 3.1.1.1.1 ► FUNCTIONS

A **function** is a functional and total relation.

0004

## NOTATION 3.1.1.1.2 ► ADDITIONAL NOTATION FOR FUNCTIONS

Throughout this work, we will sometimes denote a function  $f: X \rightarrow Y$  by

$$f \stackrel{\text{def}}{=} \llbracket x \mapsto f(x) \rrbracket.$$

01Y2

1. For example, given a function

$$\Phi: \text{Hom}_{\text{Sets}}(X, Y) \rightarrow K$$

taking values on a set of functions such as  $\text{Hom}_{\text{Sets}}(X, Y)$ , we will sometimes also write

$$\Phi(f) \stackrel{\text{def}}{=} \Phi(\llbracket x \mapsto f(x) \rrbracket).$$

01Y3

2. This notational choice is based on the lambda notation

$$f \stackrel{\text{def}}{=} (\lambda x. f(x)),$$

but uses a “ $\mapsto$ ” symbol for better spacing and double brackets instead of either:

01Y4

- (a) Square brackets  $[x \mapsto f(x)]$ ;

01Y5

- (b) Parentheses  $(x \mapsto f(x))$ ;

hoping to improve readability when dealing with e.g.:

01Y6

- (a) Equivalence classes, cf.:

01Y7

- i.  $\llbracket [x] \mapsto f([x]) \rrbracket$

01Y8

- ii.  $[ [x] \mapsto f([x]) ]$

01Y9

- iii.  $(\lambda [x]. f([x]))$

01YA

- (b) Function evaluations, cf.:

01YB

- i.  $\Phi(\llbracket x \mapsto f(x) \rrbracket)$

01YC

- ii.  $\Phi((x \mapsto f(x)))$

01YD

- iii.  $\Phi((\lambda x. f(x)))$

01YE

3. We will also sometimes write  $-$ ,  $-_1$ ,  $-_2$ , etc. for the arguments of a function. Some examples include:

01YF

(a) Writing  $f(-_1)$  for a function  $f: A \rightarrow B$ .

01YG

(b) Writing  $f(-_1, -_2)$  for a function  $f: A \times B \rightarrow C$ .

01YH

(c) Given a function  $f: A \times B \rightarrow C$ , writing

$$f(a, -): B \rightarrow C$$

for the function  $\llbracket b \mapsto f(a, b) \rrbracket$ .

01YJ

(d) Denoting a composition of the form

$$A \times B \xrightarrow{\phi \times \text{id}_B} A' \times B \xrightarrow{f} C$$

by  $f(\phi(-_1), -_2)$ .

01YK

4. Finally, given a function  $f: A \rightarrow B$ , we will sometimes write

$$\text{ev}_a(f) \stackrel{\text{def}}{=} f(a)$$

for the value of  $f$  at some  $a \in A$ .

For an example of the above notations being used in practice, see the proof of the adjunction

$$(A \times - \dashv \text{Hom}_{\text{Sets}}(A, -)): \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Hom}_{\text{Sets}}(A, -)} \end{array} \text{Sets},$$

stated in **Constructions With Sets, Item 2** of **Proposition 4.1.3.1.4**.

## 0005 3.2 The Enrichment of Sets in Classical Truth Values

### 0006 3.2.1 $(-2)$ -Categories

#### 0007 DEFINITION 3.2.1.1.1 ► $(-2)$ -CATEGORIES

A  $(-2)$ -**category** is the “necessarily true” truth value.<sup>1,2,3</sup>

<sup>1</sup>Thus, there is only one  $(-2)$ -category.

<sup>2</sup>A  $(-n)$ -category for  $n = 3, 4, \dots$  is also the “necessarily true” truth value, coinciding with a  $(-2)$ -category.

<sup>3</sup>For motivation, see [BS10, p. 13].

### 0008 3.2.2 $(-1)$ -Categories

#### 0009 DEFINITION 3.2.2.1.1 ► $(-1)$ -CATEGORIES

A  $(-1)$ -**category** is a classical truth value.

#### 000A REMARK 3.2.2.1.2 ► MOTIVATION FOR $(-1)$ -CATEGORIES

<sup>1</sup> $(-1)$ -categories should be thought of as being “categories enriched in  $(-2)$ -categories”, having a collection of objects and, for each pair of objects, a Hom-object  $\text{Hom}(x, y)$  that is a  $(-2)$ -category (i.e. trivial).

As a result, a  $(-1)$ -category  $C$  is either:<sup>2</sup>

- 01D2 1. *Empty*, having no objects.
- 01D3 2. *Contractible*, having a collection of objects  $\{a, b, c, \dots\}$ , but with  $\text{Hom}_C(a, b)$  being a  $(-2)$ -category (i.e. trivial) for all  $a, b \in \text{Obj}(C)$ , forcing all objects of  $C$  to be uniquely isomorphic to each other.

Thus there are only two  $(-1)$ -categories up to equivalence:

- 01D4 1. The  $(-1)$ -category false (the empty one);
- 01D5 2. The  $(-1)$ -category true (the contractible one).

<sup>1</sup>For more motivation, see [BS10, p. 13].

<sup>2</sup>See [BS10, pp. 33–34].

000B

**DEFINITION 3.2.2.1.3 ► THE POSET OF TRUTH VALUES**

The **poset of truth values**<sup>1</sup> is the poset  $(\{\text{true}, \text{false}\}, \preceq)$  consisting of:

- *The Underlying Set.* The set  $\{\text{true}, \text{false}\}$  whose elements are the truth values true and false.
- *The Partial Order.* The partial order

$$\preceq: \{\text{true}, \text{false}\} \times \{\text{true}, \text{false}\} \rightarrow \{\text{true}, \text{false}\}$$

on  $\{\text{true}, \text{false}\}$  defined by<sup>2</sup>

$$\begin{aligned} \text{false} &\preceq \text{false} \stackrel{\text{def}}{=} \text{true}, \\ \text{true} &\preceq \text{false} \stackrel{\text{def}}{=} \text{false}, \\ \text{false} &\preceq \text{true} \stackrel{\text{def}}{=} \text{true}, \\ \text{true} &\preceq \text{true} \stackrel{\text{def}}{=} \text{true}. \end{aligned}$$

<sup>1</sup>Further Terminology: Also called the **poset of  $(-1)$ -categories**.

<sup>2</sup>This partial order coincides with logical implication.

000C

**NOTATION 3.2.2.1.4 ► FURTHER NOTATION FOR THE POSET OF TRUTH VALUES**

We also write  $\{t, f\}$  for the poset  $\{\text{true}, \text{false}\}$ .

000D

**PROPOSITION 3.2.2.1.5 ► CARTESIAN CLOSEDNESS OF THE POSET OF TRUTH VALUES**

The poset of truth values  $\{t, f\}$  is Cartesian closed with product given by<sup>1</sup>

$$\begin{aligned} t \times t &= t, & f \times t &= f, \\ t \times f &= f, & f \times f &= f, \end{aligned}$$

$\times$	t	f
t	t	f
f	f	f

and internal Hom  $\mathbf{Hom}_{\{t,f\}}$  given by the partial order of  $\{t, f\}$ , i.e. by

$$\begin{aligned} \mathbf{Hom}_{\{t,f\}}(t, t) &= t, & \mathbf{Hom}_{\{t,f\}}(f, t) &= t, \\ \mathbf{Hom}_{\{t,f\}}(t, f) &= f, & \mathbf{Hom}_{\{t,f\}}(f, f) &= t, \end{aligned}$$

$\mathbf{Hom}_{\{t,f\}}$	t	f
t	t	t
f	t	f


<sup>1</sup>Note that  $\times$  coincides with the “and” operator, while  $\mathbf{Hom}_{\{t,f\}}$  coincides with the logical implication operator.

|


|

## Existence of Products

We claim that the products  $t \times t$ ,  $t \times f$ ,  $f \times t$ , and  $f \times f$  satisfy the universal property of the product in  $\{t, f\}$ . Indeed, suppose we have diagrams of the form



where the  $pr_1$  and  $pr_2$  morphisms are the only possible ones (since  $\{t, f\}$  is posetal). We claim that there are unique morphisms making the diagrams



commute. Indeed:

- 01YL 1. If  $P_1 = t$ , then  $p_1^1 = p_2^1 = \text{id}_t$ , so there's a unique morphism from  $P_1$  to  $t$  making the diagram commute, namely  $\text{id}_t$ .
- 01YM 2. If  $P_1 = f$ , then  $p_1^1 = p_2^1$  are given by the unique morphism from  $f$  to  $t$ , so there's a unique morphism from  $P_1$  to  $t$  making the diagram commute, namely the unique morphism from  $f$  to  $t$ .
- 01YN 3. If  $P_2 = t$ , then there is no morphism  $p_2^2$ .
- 01YP 4. If  $P_2 = f$ , then  $p_1^2$  is the unique morphism from  $f$  to  $t$  while  $p_2^2 = \text{id}_f$ , so there's a unique morphism from  $P_2$  to  $f$  making the diagram commute, namely  $\text{id}_f$ .
- 01YQ 5. The proof for  $P_3$  is similar to the one for  $P_2$ .
- 01YR 6. If  $P_4 = t$ , then there is no morphism  $p_1^4$  or  $p_2^4$ .
- 01YS 7. If  $P_4 = f$ , then  $p_1^4 = p_2^4 = \text{id}_f$ , so there's a unique morphism from  $P_4$  to  $f$  making the diagram commute, namely  $\text{id}_f$ .

This finishes the existence of products part of the proof.

### Cartesian Closedness

We claim there's a bijection

$$\mathrm{Hom}_{\{t,f\}}(A \times B, C) \cong \mathrm{Hom}_{\{t,f\}}(A, \mathbf{Hom}_{\{t,f\}}(B, C)),$$

natural in  $A, B, C \in \{t, f\}$ . Indeed:

- For  $(A, B, C) = (t, t, t)$ , we have

$$\begin{aligned} \mathrm{Hom}_{\{t,f\}}(t \times t, t) &\cong \mathrm{Hom}_{\{t,f\}}(t, t) \\ &= \{\mathrm{id}_{\mathrm{true}}\} \\ &\cong \mathrm{Hom}_{\{t,f\}}(t, t) \\ &\cong \mathrm{Hom}_{\{t,f\}}(t, \mathbf{Hom}_{\{t,f\}}(t, t)). \end{aligned}$$

- For  $(A, B, C) = (t, t, f)$ , we have

$$\begin{aligned} \mathrm{Hom}_{\{t,f\}}(t \times t, f) &\cong \mathrm{Hom}_{\{t,f\}}(t, f) \\ &= \emptyset \\ &\cong \mathrm{Hom}_{\{t,f\}}(t, f) \\ &\cong \mathrm{Hom}_{\{t,f\}}(t, \mathbf{Hom}_{\{t,f\}}(t, f)). \end{aligned}$$

- For  $(A, B, C) = (t, f, t)$ , we have

$$\begin{aligned} \mathrm{Hom}_{\{t,f\}}(t \times f, t) &\cong \mathrm{Hom}_{\{t,f\}}(f, t) \\ &\cong \mathrm{pt} \\ &\cong \mathrm{Hom}_{\{t,f\}}(f, t) \\ &\cong \mathrm{Hom}_{\{t,f\}}(f, \mathbf{Hom}_{\{t,f\}}(f, t)). \end{aligned}$$

- For  $(A, B, C) = (t, f, f)$ , we have

$$\begin{aligned} \mathrm{Hom}_{\{t,f\}}(t \times f, f) &\cong \mathrm{Hom}_{\{t,f\}}(f, f) \\ &\cong \{\mathrm{id}_{\mathrm{false}}\} \\ &\cong \mathrm{Hom}_{\{t,f\}}(f, f) \\ &\cong \mathrm{Hom}_{\{t,f\}}(t, \mathbf{Hom}_{\{t,f\}}(f, f)). \end{aligned}$$



- For  $(A, B, C) = (f, t, t)$ , we have

$$\begin{aligned}
 \text{Hom}_{\{t, f\}}(f \times t, t) &\cong \text{Hom}_{\{t, f\}}(f, t) \\
 &\cong \text{pt} \\
 &\cong \text{Hom}_{\{t, f\}}(f, t) \\
 &\cong \text{Hom}_{\{t, f\}}(f, \mathbf{Hom}_{\{t, f\}}(t, t)).
 \end{aligned}$$

- For  $(A, B, C) = (f, t, f)$ , we have


$$\begin{aligned}
 \text{Hom}_{\{t, f\}}(f \times t, f) &\cong \text{Hom}_{\{t, f\}}(f, f) \\
 &\cong \{\text{id}_{\text{false}}\} \\
 &\cong \text{Hom}_{\{t, f\}}(f, f) \\
 &\cong \text{Hom}_{\{t, f\}}(f, \mathbf{Hom}_{\{t, f\}}(t, f)).
 \end{aligned}$$

- For  $(A, B, C) = (f, f, t)$ , we have

$$\begin{aligned}
 \text{Hom}_{\{t, f\}}(f \times f, t) &\cong \text{Hom}_{\{t, f\}}(f, t) \\
 &\cong \text{pt} \\
 &\cong \text{Hom}_{\{t, f\}}(f, t) \\
 &\cong \text{Hom}_{\{t, f\}}(f, \mathbf{Hom}_{\{t, f\}}(f, t)).
 \end{aligned}$$

- For  $(A, B, C) = (f, f, f)$ , we have

$$\begin{aligned}
 \text{Hom}_{\{t, f\}}(f \times f, f) &\cong \text{Hom}_{\{t, f\}}(f, f) \\
 &= \{\text{id}_{\text{false}}\} \\
 &\cong \text{Hom}_{\{t, f\}}(f, f) \\
 &\cong \text{Hom}_{\{t, f\}}(f, \mathbf{Hom}_{\{t, f\}}(f, f)).
 \end{aligned}$$

Since  $\{t, f\}$  is posetal, naturality is automatic ([Categories, Item 4 of Proposition 11.2.7.1.2](#)). 

### 000E 3.2.3 0-Categories

#### 000F DEFINITION 3.2.3.1.1 ► 0-CATEGORIES

A **0-category** is a poset.<sup>1</sup>

<sup>1</sup>*Motivation:* A 0-category is precisely a category enriched in the poset of  $(-1)$ -categories.

#### 000G DEFINITION 3.2.3.1.2 ► 0-GROUPOIDS

A **0-groupoid** is a 0-category in which every morphism is invertible.<sup>1</sup>

<sup>1</sup>That is, a set.

### 000H 3.2.4 Tables of Analogies Between Set Theory and Category Theory

Here we record some analogies between notions in set theory and category theory. The analogies relating to presheaves relate equally well to copresheaves, as the opposite  $X^{\text{op}}$  of a set  $X$  is just  $X$  again.

#### 01D6 REMARK 3.2.4.1.1 ► BASIC ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY

The basic analogies between set theory and category theory are summarised in the following table:

Set Theory	Category Theory
Enrichment in $\{\text{true}, \text{false}\}$	Enrichment in Sets
Set $X$	Category $\mathcal{C}$
Element $x \in X$	Object $X \in \text{Obj}(\mathcal{C})$
Function $f: X \rightarrow Y$	Functor $F: \mathcal{C} \rightarrow \mathcal{D}$
Function $X \rightarrow \{\text{true}, \text{false}\}$	Copresheaf $\mathcal{C} \rightarrow \text{Sets}$
Function $X \rightarrow \{\text{true}, \text{false}\}$	Presheaf $\mathcal{C}^{\text{op}} \rightarrow \text{Sets}$

01D7

**REMARK 3.2.4.1.2 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: POWERSETS AND CATEGORIES OF PRESHEAVES**

The category of presheaves  $\mathbf{PSh}(C)$  and the category of copresheaves  $\mathbf{CoPSh}(C)$  on a category  $C$  are the 1-categorical counterparts to the powerset  $\mathcal{P}(X)$  of subsets of a set  $X$ . The further analogies built upon this are summarised in the following table:

Set Theory	Category Theory
Powerset $\mathcal{P}(X)$	Presheaf category $\mathbf{PSh}(C)$
Characteristic function $\chi_{\{x\}}: X \rightarrow \{t, f\}$	Representable presheaf $h_X: C^{\text{op}} \hookrightarrow \mathbf{Sets}$
Characteristic embedding $\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$	Yoneda embedding $\mathcal{Y}: C^{\text{op}} \hookrightarrow \mathbf{PSh}(C)$
Characteristic relation $\chi_X(-, -): X \times X \rightarrow \{t, f\}$	Hom profunctor $\text{Hom}_C(-, -): C^{\text{op}} \times C \rightarrow \mathbf{Sets}$
The Yoneda lemma for sets $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$	The Yoneda lemma for categories $\text{Nat}(h_X, \mathcal{F}) \cong \mathcal{F}(X)$
The characteristic embedding is fully faithful, $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$	The Yoneda embedding is fully faithful, $\text{Nat}(h_X, h_Y) \cong \text{Hom}_C(X, Y)$
Subsets are unions of their elements $U = \bigcup_{x \in U} \{x\}$ or $\chi_U = \text{colim}_{\chi_x \in \mathcal{P}(U)} (\chi_x)$	Presheaves are colimits of representables, $\mathcal{F} \cong \text{colim}_{h_X \in \int_C \mathcal{F}} (h_X)$

01D8

**REMARK 3.2.4.1.3 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: CATEGORIES OF ELEMENTS**

We summarise the analogies between un/straightening in set theory and category theory in the following table:

SET THEORY	CATEGORY THEORY
Assignment $U \mapsto \chi_U$	Assignment $\mathcal{F} \mapsto \int_C \mathcal{F}$
Un/straightening isomorphism $\mathcal{P}(X) \cong \text{Sets}(X, \{t, f\})$	Un/straightening equivalence $\text{DFib}(C) \stackrel{\text{eq.}}{\cong} \text{PSh}(C)$

**REMARK 3.2.4.1.4 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: FUNCTIONS BETWEEN POWERSETS AND FUNCTORS BETWEEN PRESHEAF CATEGORIES**

01D9

We summarise the analogies between functions  $\mathcal{P}(X) \rightarrow \mathcal{P}(Y)$  and functors  $\text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$  in the following table:

SET THEORY	CATEGORY THEORY
Direct image function $f_! : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Left Kan extension functor $F_! : \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$
Inverse image function $f^{-1} : \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$	Precomposition functor $F^* : \text{PSh}(\mathcal{D}) \rightarrow \text{PSh}(C)$
Codirect image function $f_* : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Right Kan extension functor $F_* : \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$

**REMARK 3.2.4.1.5 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: RELATIONS AND PROFUNCTORS**

01DA

We summarise the analogies between functions, relations and profunctors in the following table:

SET THEORY	CATEGORY THEORY
Relation $R : X \times Y \rightarrow \{t, f\}$	Profunctor $\mathfrak{p} : \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$
Relation $R : X \rightarrow \mathcal{P}(Y)$	Profunctor $\mathfrak{p} : C \rightarrow \text{PSh}(\mathcal{D})$
Relation as a cocontinuous morphism of posets $R : (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset)$	Profunctor as a colimit-preserving functor $\mathfrak{p} : \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$

# Appendices

## A Other Chapters

### Preliminaries

1. [Introduction](#)
2. [A Guide to the Literature](#)

### Sets

3. [Sets](#)
4. [Constructions With Sets](#)
5. [Monoidal Structures on the Category of Sets](#)
6. [Pointed Sets](#)
7. [Tensor Products of Pointed Sets](#)

### Relations

8. [Relations](#)
9. [Constructions With Relations](#)

10. [Conditions on Relations](#)

### Categories

11. [Categories](#)
12. [Presheaves and the Yoneda Lemma](#)

### Monoidal Categories

13. [Constructions With Monoidal Categories](#)

### Bicategories

14. [Types of Morphisms in Bicategories](#)

### Extra Part

15. [Notes](#)

## References

- [BS10] John C. Baez and Michael Shulman. “Lectures on  $n$ -Categories and Co-homology”. In: *Towards higher categories*. Vol. 152. IMA Vol. Math. Appl. Springer, New York, 2010, pp. 1–68. DOI: [10.1007/978-1-4419-1524-5\\_1](https://doi.org/10.1007/978-1-4419-1524-5_1). URL: [https://doi.org/10.1007/978-1-4419-1524-5\\_1](https://doi.org/10.1007/978-1-4419-1524-5_1) (cit. on p. 4).