

# The Clowder Project

2017–2025



The Clowder Project Authors

Version a3c55e3, compiled on Jul 29, 2025.

# The Clowder Project Contributors

Version a3c55e3, compiled on Jul 29, 2025.

The following people have contributed to this project:

1. Tim Campion
2. Bastiaan Cnossen
3. Naim Camille Favier
4. Valeria de Paiva
5. Lorenzo Riva
6. Emily de Oliveira Santos
7. uni

# **The Clowder Project Supporters**

Version a3c55e3, compiled on Jul 29, 2025.

The following people currently support this project:

1. Estelle Sterrett
2. Sophie Morin
3. Owen Kahn

# Contents

<b>Part I Preliminaries</b>	<b>1</b>
<b>1 Introduction</b>	<b>2</b>
1.1 Introduction.....	2
1.2 Project Overview.....	9
1.3 Lists .....	12
1.A Other Chapters.....	22
<b>2 A Guide to the Literature</b>	<b>24</b>
2.1 Elementary Category Theory .....	24
2.A Other Chapters.....	24
<b>Part II Sets</b>	<b>26</b>
<b>3 Sets</b>	<b>27</b>
3.1 Sets and Functions .....	27
3.2 The Enrichment of Sets in Classical Truth Values.....	29
3.A Other Chapters.....	36
<b>4 Constructions With Sets</b>	<b>38</b>
4.1 Limits of Sets .....	40
4.2 Colimits of Sets.....	74
4.3 Operations With Sets .....	100
4.4 Powersets .....	161
4.5 Characteristic Functions .....	199
4.6 The Adjoint Triple $f_! \dashv f^{-1} \dashv f_*$ .....	209
4.7 Miscellany.....	251
4.A Other Chapters.....	263
<b>5 Monoidal Structures on the Category of Sets</b>	<b>265</b>
5.1 The Monoidal Category of Sets and Products .....	266
5.2 The Monoidal Category of Sets and Coproducts.....	300
5.3 The Bimonoidal Category of Sets, Products, and Coproducts..	318
5.A Other Chapters.....	328

<b>6 Pointed Sets .....</b>	<b>330</b>
6.1 Pointed Sets.....	331
6.2 Limits of Pointed Sets.....	341
6.3 Colimits of Pointed Sets.....	354
6.4 Constructions With Pointed Sets.....	368
6.A Other Chapters.....	383
<b>7 Tensor Products of Pointed Sets .....</b>	<b>384</b>
7.1 Bilinear Morphisms of Pointed Sets.....	386
7.2 Tensors and Cotensors of Pointed Sets by Sets.....	390
7.3 The Left Tensor Product of Pointed Sets.....	405
7.4 The Right Tensor Product of Pointed Sets.....	433
7.5 The Smash Product of Pointed Sets.....	460
7.6 Miscellany.....	519
7.A Other Chapters.....	520
<b>Part III Relations</b>	<b>521</b>
<b>8 Relations.....</b>	<b>522</b>
8.1 Relations.....	526
8.2 Examples of Relations .....	551
8.3 Categories of Relations.....	561
8.4 Categories of Relations With Apartness Composition.....	577
8.5 Properties of the 2-Category of Relations.....	581
8.6 Properties of the 2-Category of Relations With Apartness Composition.....	648
8.7 The Adjoint Pairs $R_! \dashv R_{-1}$ and $R^{-1} \dashv R_*$ .....	654
8.8 The Left Skew Monoidal Structure on $\text{Rel}(A, B)$ .....	680
8.9 The Right Skew Monoidal Structure on $\text{Rel}(A, B)$ .....	685
8.A Other Chapters.....	690
<b>9 Constructions With Relations .....</b>	<b>691</b>
9.1 Co/Limits in the Category of Relations.....	691
9.2 More Constructions With Relations .....	692
9.A Other Chapters.....	702
<b>10 Conditions on Relations .....</b>	<b>704</b>
10.1 Functional and Total Relations.....	704
10.2 Reflexive Relations .....	706
10.3 Symmetric Relations .....	709
10.4 Transitive Relations .....	713
10.5 Equivalence Relations .....	717
10.6 Quotients by Equivalence Relations.....	720
10.A Other Chapters.....	725

<b>Part IV Categories</b>	<b>727</b>
<b>11 Categories.....</b>	<b>728</b>
11.1 Categories .....	731
11.2 Examples of Categories .....	739
11.3 The Quadruple Adjunction With Sets .....	746
11.4 Groupoids .....	754
11.5 Functors.....	762
11.6 Conditions on Functors.....	772
11.7 More Conditions on Functors.....	793
11.8 Even More Conditions on Functors.....	803
11.9 Natural Transformations.....	811
11.10 Categories of Categories .....	828
11.A Other Chapters.....	835
<b>12 Presheaves and the Yoneda Lemma.....</b>	<b>836</b>
12.1 Presheaves.....	837
12.2 Copresheaves .....	849
12.3 Restricted Yoneda Embeddings and Yoneda Extensions.....	854
12.4 Functor Tensor Products .....	861
12.A Other Chapters.....	867
<b>Part V Monoidal Categories</b>	<b>868</b>
<b>13 Constructions With Monoidal Categories .....</b>	<b>869</b>
13.1 Moduli Categories of Monoidal Structures .....	869
13.2 Moduli Categories of Closed Monoidal Structures.....	882
13.3 Moduli Categories of Refinements of Monoidal Structures.....	882
13.A Other Chapters.....	883
<b>Part VI Bicategories</b>	<b>884</b>
<b>14 Types of Morphisms in Bicategories.....</b>	<b>885</b>
14.1 Monomorphisms in Bicategories.....	886
14.2 Epimorphisms in Bicategories .....	896
14.A Other Chapters.....	908
<b>Part VII Extra Part</b>	<b>910</b>
<b>15 Notes.....</b>	<b>911</b>
15.1 TikZ Code for Commutative Diagrams .....	911
15.2 Retired Tags.....	926
15.3 Miscellany .....	932
15.A Other Chapters.....	978

Index of Notation.....	991
Index of Set Theory .....	995
Index of Category Theory.....	999
Index of Higher Category Theory.....	1002

# Contents

<b>Part I Preliminaries</b>	<b>1</b>
<b>1 Introduction</b>	<b>2</b>
1.1 Introduction .....	2
1.1.1 Project Description and Goals.....	2
1.1.1.1 Provide a Unified and Complete Reference for Category Theory ...	3
1.1.1.2 Gather Hard to Find Results.....	3
1.1.1.3 Elaborate on Details That Are Often Left Out.....	3
1.1.1.4 Homogenize Conventions, Notation, and Terminology .....	3
1.1.1.5 Fill Gaps in the Category Theory Literature .....	4
1.1.1.6 Provide a Citable Reference for All Kinds of Results .....	4
1.1.2 Navigating the Clowder Project.....	4
1.1.2.1 Preferences.....	4
1.1.2.2 Large Diagrams and the Zoom in Feature.....	4
1.1.2.3 PDF Styles.....	4
1.1.3 Prerequisites/Assumed Background.....	5
1.1.4 Community Engagement, Contributions and Collaboration.....	5
1.1.4.1 How to Contribute .....	6
1.1.4.2 Ways to Contribute: Missing Proofs.....	6
1.1.4.3 Ways to Contribute: Missing Examples .....	6
1.1.4.4 Ways to Contribute: Questions.....	6
1.1.5 Frequently Asked Questions.....	7
1.1.5.1 How does Clowder differ from the nLab? .....	7
1.1.5.2 Why not just use the nLab instead?.....	7
1.1.6 Goodies.....	8
1.1.6.1 General Utility.....	8
1.1.6.2 Set Theory Through a Categorical Lens .....	8
1.1.6.3 Category Theory .....	9
1.2 Project Overview .....	9
1.2.1 Content and Scope.....	9
1.2.1.1 Elementary Category Theory .....	9
1.2.1.2 Variants of Category Theory .....	9
1.2.1.3 Higher Category Theory .....	9
1.2.1.4 $\infty$ -Categories.....	10
1.2.1.5 Other Topics .....	10
1.2.2 Style.....	10
1.2.2.1 Presentation of Topics .....	10
1.2.2.2 Provable Items Come With Proofs .....	11
1.2.2.3 Proper Justification of Proofs .....	11
1.2.3 Infrastructure and Technical Implementation.....	11
1.2.3.1 Removed Features (in Comparison With the Stacks Project) .....	11

1.2.3.2	Gerby and the Tags System .....	11
1.2.3.3	Placeholder Symbols and Future Style.....	12
1.3	Lists.....	12
1.3.1	List of Omitted Proofs.....	12
1.3.2	List of Missing Examples .....	16
1.3.3	List of Questions.....	17
1.3.4	List of Gaps in the Category Theory Literature.....	18
1.A	Other Chapters.....	22
<b>2</b>	<b>A Guide to the Literature .....</b>	<b>24</b>
2.1	Elementary Category Theory.....	24
2.1.1	Textbooks .....	24
2.A	Other Chapters.....	24
<b>Part II</b>	<b>Sets</b>	<b>26</b>
<b>3</b>	<b>Sets.....</b>	<b>27</b>
3.1	Sets and Functions.....	27
3.1.1	Functions.....	27
3.2	The Enrichment of Sets in Classical Truth Values.....	29
3.2.1	(-2)-Categories.....	29
3.2.2	(-1)-Categories.....	29
3.2.3	0-Categories .....	33
3.2.4	Tables of Analogies Between Set Theory and Category Theory .....	34
3.A	Other Chapters.....	36
<b>4</b>	<b>Constructions With Sets.....</b>	<b>38</b>
4.1	Limits of Sets.....	40
4.1.1	The Terminal Set.....	40
4.1.2	Products of Families of Sets.....	41
4.1.3	Binary Products of Sets .....	45
4.1.4	Pullbacks .....	55
4.1.5	Equalisers .....	67
4.1.6	Inverse Limits .....	71
4.2	Colimits of Sets.....	74
4.2.1	The Initial Set.....	74
4.2.2	Coproducts of Families of Sets .....	75
4.2.3	Binary Coproducts .....	77
4.2.4	Pushouts.....	81
4.2.5	Coequalisers .....	90
4.2.6	Direct Colimits .....	94
4.3	Operations With Sets.....	100
4.3.1	The Empty Set .....	100
4.3.2	Singleton Sets .....	100
4.3.3	Pairings of Sets .....	100
4.3.4	Ordered Pairs.....	100
4.3.5	Sets of Maps .....	101
4.3.6	Unions of Families of Subsets .....	104
4.3.7	Intersections of Families of Subsets.....	119
4.3.8	Binary Unions .....	133
4.3.9	Binary Intersections .....	139
4.3.10	Differences .....	145

---

4.3.11	Complements.....	150
4.3.12	Symmetric Differences.....	153
4.4	Powersets.....	161
4.4.1	Foundations .....	161
4.4.2	Functionality of Powersets.....	164
4.4.3	Adjointness of Powersets I .....	166
4.4.4	Adjointness of Powersets II .....	167
4.4.5	Powersets as Free Cocompletions .....	167
4.4.6	Powersets as Free Completions .....	174
4.4.7	The Internal Hom of a Powerset.....	180
4.4.8	Isbell Duality for Sets .....	197
4.5	Characteristic Functions.....	199
4.5.1	The Characteristic Function of a Subset.....	199
4.5.2	The Characteristic Function of a Point.....	204
4.5.3	The Characteristic Relation of a Set.....	205
4.5.4	The Characteristic Embedding of a Set.....	206
4.5.5	The Yoneda Lemma for Sets.....	207
4.6	The Adjoint Triple $f_! \dashv f^{-1} \dashv f_*$ .....	209
4.6.1	Direct Images.....	209
4.6.2	Inverse Images .....	220
4.6.3	Codirect Images .....	230
4.6.4	A Six-Functor Formalism for Sets .....	243
4.7	Miscellany.....	251
4.7.1	Injective Functions .....	251
4.7.2	Surjective Functions .....	257
4.A	Other Chapters.....	263
<b>5</b>	<b>Monoidal Structures on the Category of Sets .....</b>	<b>265</b>
5.1	The Monoidal Category of Sets and Products .....	266
5.1.1	Products of Sets.....	266
5.1.2	The Internal Hom of Sets .....	266
5.1.3	The Monoidal Unit.....	266
5.1.4	The Associator .....	266
5.1.5	The Left Unitor .....	269
5.1.6	The Right Unitor .....	271
5.1.7	The Symmetry .....	273
5.1.8	The Diagonal .....	275
5.1.9	The Monoidal Category of Sets and Products .....	278
5.1.10	The Universal Property of $(\text{Sets}, \times, \text{pt})$ .....	283
5.2	The Monoidal Category of Sets and Coproducts.....	300
5.2.1	Coproducts of Sets.....	300
5.2.2	The Monoidal Unit .....	300
5.2.3	The Associator .....	300
5.2.4	The Left Unitor .....	303
5.2.5	The Right Unitor .....	306
5.2.6	The Symmetry .....	308
5.2.7	The Monoidal Category of Sets and Coproducts.....	311
5.3	The Bimonoidal Category of Sets, Products, and Coproducts .....	318
5.3.1	The Left Distributor.....	318
5.3.2	The Right Distributor.....	320
5.3.3	The Left Annihilator.....	323

---

5.3.4	The Right Annihilator .....	324
5.3.5	The Bimonoidal Category of Sets, Products, and Coproducts .....	326
5.A	Other Chapters.....	328
<b>6</b>	<b>Pointed Sets .....</b>	<b>330</b>
6.1	Pointed Sets .....	331
6.1.1	Foundations .....	331
6.1.2	Morphisms of Pointed Sets.....	333
6.1.3	The Category of Pointed Sets .....	333
6.1.4	Elementary Properties of Pointed Sets.....	334
6.1.5	Active and Inert Morphisms of Pointed Sets.....	337
6.2	Limits of Pointed Sets .....	341
6.2.1	The Terminal Pointed Set .....	341
6.2.2	Products of Families of Pointed Sets.....	341
6.2.3	Products .....	343
6.2.4	Pullbacks .....	346
6.2.5	Equalisers .....	351
6.3	Colimits of Pointed Sets .....	354
6.3.1	The Initial Pointed Set .....	354
6.3.2	Coproducts of Families of Pointed Sets .....	354
6.3.3	Coproducts.....	357
6.3.4	Pushouts.....	360
6.3.5	Coequalisers .....	366
6.4	Constructions With Pointed Sets .....	368
6.4.1	Free Pointed Sets.....	368
6.4.2	Deleting Basepoints.....	376
6.A	Other Chapters.....	383
<b>7</b>	<b>Tensor Products of Pointed Sets .....</b>	<b>384</b>
7.1	Bilinear Morphisms of Pointed Sets .....	386
7.1.1	Left Bilinear Morphisms of Pointed Sets .....	386
7.1.2	Right Bilinear Morphisms of Pointed Sets .....	387
7.1.3	Bilinear Morphisms of Pointed Sets .....	388
7.2	Tensors and Cotensors of Pointed Sets by Sets .....	390
7.2.1	Tensors of Pointed Sets by Sets .....	390
7.2.2	Cotensors of Pointed Sets by Sets .....	398
7.3	The Left Tensor Product of Pointed Sets.....	405
7.3.1	Foundations .....	405
7.3.2	The Left Internal Hom of Pointed Sets .....	411
7.3.3	The Left Skew Unit .....	414
7.3.4	The Left Skew Associator .....	414
7.3.5	The Left Skew Left Unitor.....	417
7.3.6	The Left Skew Right Unitor .....	420
7.3.7	The Diagonal .....	422
7.3.8	The Left Skew Monoidal Structure on Pointed Sets Associated to $\triangleleft$ .....	424
7.3.9	Monoids With Respect to the Left Tensor Product of Pointed Sets.....	428
7.4	The Right Tensor Product of Pointed Sets.....	433
7.4.1	Foundations .....	433
7.4.2	The Right Internal Hom of Pointed Sets .....	438
7.4.3	The Right Skew Unit .....	441
7.4.4	The Right Skew Associator .....	442
7.4.5	The Right Skew Left Unitor .....	445

---

7.4.6	The Right Skew Right Unitor.....	447
7.4.7	The Diagonal .....	450
7.4.8	The Right Skew Monoidal Structure on Pointed Sets Associated to $\triangleright$ .....	451
7.4.9	Monoids With Respect to the Right Tensor Product of Pointed Sets.....	455
7.5	The Smash Product of Pointed Sets.....	460
7.5.1	Foundations .....	460
7.5.2	The Internal Hom of Pointed Sets.....	470
7.5.3	The Monoidal Unit.....	473
7.5.4	The Associator .....	473
7.5.5	The Left Unitor .....	475
7.5.6	The Right Unitor .....	478
7.5.7	The Symmetry .....	481
7.5.8	The Diagonal .....	483
7.5.9	The Monoidal Structure on Pointed Sets Associated to $\wedge$ .....	486
7.5.10	The Universal Property of $(\text{Sets}_*, \wedge, S^0)$ .....	491
7.5.11	Monoids With Respect to the Smash Product of Pointed Sets .....	516
7.5.12	Comonoids With Respect to the Smash Product of Pointed Sets .....	517
7.6	Miscellany.....	519
7.6.1	The Smash Product of a Family of Pointed Sets .....	519
7.A	Other Chapters.....	520

**Part III Relations** **521**

<b>8</b>	<b>Relations .....</b>	<b>522</b>
8.1	Relations.....	526
8.1.1	Foundations .....	526
8.1.2	Relations as Decategorifications of Profunctors.....	530
8.1.3	Composition of Relations.....	532
8.1.4	Apartness Composition of Relations .....	540
8.1.5	The Converse of a Relation .....	547
8.2	Examples of Relations.....	551
8.2.1	Elementary Examples of Relations.....	551
8.2.2	The Graph of a Function.....	554
8.2.3	The Inverse of a Function .....	558
8.2.4	Representable Relations .....	560
8.3	Categories of Relations .....	561
8.3.1	The Category of Relations Between Two Sets.....	561
8.3.2	The Category of Relations.....	561
8.3.3	The Closed Symmetric Monoidal Category of Relations .....	562
8.3.3.1	The Monoidal Product.....	562
8.3.3.2	The Monoidal Unit .....	563
8.3.3.3	The Associator.....	563
8.3.3.4	The Left Unitor.....	564
8.3.3.5	The Right Unitor .....	564
8.3.3.6	The Symmetry .....	565
8.3.3.7	The Internal Hom .....	566
8.3.3.8	The Closed Symmetric Monoidal Category of Relations .....	567
8.3.4	The 2-Category of Relations .....	568
8.3.5	The Double Category of Relations .....	569
8.3.5.1	The Double Category of Relations .....	569
8.3.5.2	Horizontal Identities.....	571

---

8.3.5.3	Horizontal Composition.....	572
8.3.5.4	Vertical Composition of 2-Morphisms .....	573
8.3.5.5	The Associators .....	575
8.3.5.6	The Left Unitors.....	576
8.3.5.7	The Right Unitors.....	577
8.4	Categories of Relations With Apartness Composition .....	577
8.4.1	The Category of Relations With Apartness Composition .....	577
8.4.2	The 2-Category of Relations With Apartness Composition.....	578
8.4.3	The Linear Bicategory of Relations .....	580
8.4.4	Other Categorical Structures With Apartness Composition.....	580
8.5	Properties of the 2-Category of Relations .....	581
8.5.1	Self-Duality.....	581
8.5.2	Isomorphisms and Equivalences.....	582
8.5.3	Internal Adjunctions.....	584
8.5.4	Internal Monads.....	586
8.5.5	Internal Comonads .....	588
8.5.6	Modules Over Internal Monads.....	590
8.5.7	Comodules Over Internal Comonads .....	591
8.5.8	Eilenberg–Moore and Kleisli Objects.....	593
8.5.9	Co/Monoids .....	593
8.5.10	Monomorphisms and 2-Categorical Monomorphisms.....	594
8.5.11	Epimorphisms and 2-Categorical Epimorphisms .....	612
8.5.12	Co/Limits.....	632
8.5.13	Internal Left Kan Extensions .....	632
8.5.14	Internal Left Kan Lifts.....	635
8.5.15	Internal Right Kan Extensions .....	636
8.5.16	Internal Right Kan Lifts.....	642
8.5.17	Closedness .....	646
8.5.18	Rel as a Category of Free Algebras.....	647
8.6	Properties of the 2-Category of Relations With Apartness Composition.....	648
8.6.1	Self-Duality.....	648
8.6.2	Isomorphisms and Equivalences.....	649
8.6.3	Internal Adjunctions.....	651
8.6.4	Internal Monads.....	651
8.6.5	Internal Comonads .....	652
8.6.6	Modules Over Internal Monads.....	654
8.6.7	Comodules Over Internal Comonads .....	654
8.6.8	Eilenberg–Moore and Kleisli Objects.....	654
8.6.9	Monomorphisms.....	654
8.6.10	2-Categorical Monomorphisms .....	654
8.6.11	Epimorphisms .....	654
8.6.12	2-Categorical Epimorphisms .....	654
8.6.13	Co/Limits.....	654
8.6.14	Internal Left Kan Extensions .....	654
8.6.15	Internal Left Kan Lifts.....	654
8.6.16	Internal Right Kan Extensions .....	654
8.6.17	Internal Right Kan Lifts.....	654
8.6.18	Coclosedness .....	654
8.7	The Adjoint Pairs $R_! \dashv R_{-1}$ and $R^{-1} \dashv R_*$ .....	654
8.7.1	Direct Images.....	654
8.7.2	Coinverse Images.....	660

8.7.3	Inverse Images .....	666
8.7.4	Codirect Images .....	671
8.7.5	Functoriality of Powersets.....	678
8.8	The Left Skew Monoidal Structure on $\text{Rel}(A, B)$ .....	680
8.8.1	The Left Skew Monoidal Product.....	680
8.8.2	The Left Skew Monoidal Unit.....	681
8.8.3	The Left Skew Associators .....	681
8.8.4	The Left Skew Left Unitors .....	682
8.8.5	The Left Skew Right Unitors .....	683
8.8.6	The Left Skew Monoidal Structure on $\text{Rel}(A, B)$ .....	684
8.9	The Right Skew Monoidal Structure on $\text{Rel}(A, B)$ .....	685
8.9.1	The Right Skew Monoidal Product.....	685
8.9.2	The Right Skew Monoidal Unit.....	686
8.9.3	The Right Skew Associators .....	686
8.9.4	The Right Skew Left Unitors .....	687
8.9.5	The Right Skew Right Unitors .....	688
8.9.6	The Right Skew Monoidal Structure on $\text{Rel}(A, B)$ .....	689
8.A	Other Chapters.....	690
<b>9</b>	<b>Constructions With Relations.....</b>	<b>691</b>
9.1	Co/Limits in the Category of Relations .....	691
9.2	More Constructions With Relations.....	692
9.2.1	The Domain and Range of a Relation .....	692
9.2.2	Binary Unions of Relations.....	692
9.2.3	Unions of Families of Relations .....	694
9.2.4	Binary Intersections of Relations .....	695
9.2.5	Intersections of Families of Relations .....	696
9.2.6	Binary Products of Relations .....	697
9.2.7	Products of Families of Relations .....	699
9.2.8	The Collage of a Relation .....	699
9.A	Other Chapters.....	702
<b>10</b>	<b>Conditions on Relations.....</b>	<b>704</b>
10.1	Functional and Total Relations .....	704
10.1.1	Functional Relations.....	704
10.1.2	Total Relations .....	705
10.2	Reflexive Relations .....	706
10.2.1	Foundations .....	706
10.2.2	The Reflexive Closure of a Relation.....	707
10.3	Symmetric Relations.....	709
10.3.1	Foundations .....	709
10.3.2	The Symmetric Closure of a Relation .....	710
10.4	Transitive Relations.....	713
10.4.1	Foundations .....	713
10.4.2	The Transitive Closure of a Relation.....	714
10.5	Equivalence Relations .....	717
10.5.1	Foundations .....	717
10.5.2	The Equivalence Closure of a Relation .....	717
10.6	Quotients by Equivalence Relations .....	720
10.6.1	Equivalence Classes.....	720
10.6.2	Quotients of Sets by Equivalence Relations.....	721
10.A	Other Chapters.....	725

<b>Part IV Categories</b>	<b>727</b>
<b>11 Categories.....</b>	<b>728</b>
11.1 Categories.....	731
11.1.1 Foundations .....	731
11.1.2 Subcategories .....	733
11.1.3 Skeletons of Categories .....	734
11.1.4 Precomposition and Postcomposition .....	735
11.2 Examples of Categories.....	739
11.2.1 The Empty Category .....	739
11.2.2 The Punctual Category.....	740
11.2.3 Monoids as One-Object Categories .....	740
11.2.4 Ordinal Categories.....	741
11.2.5 The Walking Arrow .....	742
11.2.6 More Examples of Categories .....	743
11.2.7 Posetal Categories.....	743
11.3 The Quadruple Adjunction With Sets.....	746
11.3.1 Statement.....	746
11.3.2 Connected Components and Connected Categories.....	747
11.3.2.1 Connected Components of Categories.....	747
11.3.2.2 Sets of Connected Components of Categories.....	748
11.3.2.3 Connected Categories .....	750
11.3.3 Discrete Categories .....	750
11.3.4 Indiscrete Categories .....	752
11.4 Groupoids.....	754
11.4.1 Isomorphisms .....	754
11.4.2 Groupoids.....	754
11.4.3 The Groupoid Completion of a Category.....	755
11.4.4 The Core of a Category .....	759
11.5 Functors .....	762
11.5.1 Foundations .....	762
11.5.2 Contravariant Functors .....	766
11.5.3 Forgetful Functors .....	768
11.5.4 The Natural Transformation Associated to a Functor .....	770
11.6 Conditions on Functors .....	772
11.6.1 Faithful Functors .....	772
11.6.2 Full Functors .....	774
11.6.3 Fully Faithful Functors.....	781
11.6.4 Conservative Functors.....	785
11.6.5 Essentially Injective Functors .....	787
11.6.6 Essentially Surjective Functors.....	788
11.6.7 Equivalences of Categories.....	789
11.6.8 Isomorphisms of Categories .....	791
11.7 More Conditions on Functors .....	793
11.7.1 Dominant Functors .....	793
11.7.2 Monomorphisms of Categories.....	794
11.7.3 Epimorphisms of Categories .....	795
11.7.4 Pseudomonadic Functors .....	797
11.7.5 Pseudoepic Functors.....	800
11.8 Even More Conditions on Functors .....	803
11.8.1 Injective on Objects Functors .....	803

11.8.2 Surjective on Objects Functors.....	804
11.8.3 Bijective on Objects Functors.....	804
11.8.4 Functors Representably Faithful on Cores .....	804
11.8.5 Functors Representably Full on Cores.....	805
11.8.6 Functors Representably Fully Faithful on Cores .....	806
11.8.7 Functors Corepresentably Faithful on Cores.....	807
11.8.8 Functors Corepresentably Full on Cores.....	808
11.8.9 Functors Corepresentably Fully Faithful on Cores .....	809
11.9 Natural Transformations .....	811
11.9.1 Transformations .....	811
11.9.2 Natural Transformations .....	812
11.9.3 Examples of Natural Transformations.....	813
11.9.4 Vertical Composition of Natural Transformations.....	815
11.9.5 Horizontal Composition of Natural Transformations.....	819
11.9.6 Properties of Natural Transformations.....	824
11.9.7 Natural Isomorphisms.....	826
11.10 Categories of Categories.....	828
11.10.1 Functor Categories.....	828
11.10.2 The Category of Categories and Functors.....	832
11.10.3 The 2-Category of Categories, Functors, and Natural Transformations.....	833
11.10.4 The Category of Groupoids.....	834
11.10.5 The 2-Category of Groupoids.....	834
11.A Other Chapters.....	835
<b>12 Presheaves and the Yoneda Lemma.....</b>	<b>836</b>
12.1 Presheaves .....	837
12.1.1 Foundations .....	837
12.1.2 Representable Presheaves.....	838
12.1.3 Representable Natural Transformations .....	840
12.1.4 The Yoneda Embedding .....	840
12.1.5 The Yoneda Lemma.....	843
12.1.6 Properties of Categories of Presheaves .....	848
12.2 Copresheaves.....	849
12.2.1 Foundations .....	849
12.2.2 Corepresentable Copresheaves.....	850
12.2.3 Corepresentable Natural Transformations .....	852
12.2.4 The Contravariant Yoneda Embedding .....	852
12.2.5 The Contravariant Yoneda Lemma.....	854
12.3 Restricted Yoneda Embeddings and Yoneda Extensions .....	854
12.3.1 Foundations .....	854
12.3.2 The Yoneda Extension Functor.....	858
12.4 Functor Tensor Products.....	861
12.4.1 The Tensor Product of Presheaves With Copresheaves.....	861
12.4.2 The Tensor of a Presheaf With a Functor.....	864
12.4.3 The Tensor of a Copresheaf With a Functor .....	865
12.A Other Chapters.....	867
<b>Part V Monoidal Categories</b>	<b>868</b>
<b>13 Constructions With Monoidal Categories.....</b>	<b>869</b>
13.1 Moduli Categories of Monoidal Structures.....	869
13.1.1 The Moduli Category of Monoidal Structures on a Category.....	869

13.1.2 The Moduli Category of Braided Monoidal Structures on a Category.....	882
13.1.3 The Moduli Category of Symmetric Monoidal Structures on a Category.....	882
13.2 Moduli Categories of Closed Monoidal Structures .....	882
13.3 Moduli Categories of Refinements of Monoidal Structures .....	882
13.3.1 The Moduli Category of Braided Refinements of a Monoidal Structure .....	882
13.A Other Chapters.....	883

**Part VI Bicategories** **884**

<b>14 Types of Morphisms in Bicategories.....</b>	<b>885</b>
14.1 Monomorphisms in Bicategories .....	886
14.1.1 Representably Faithful Morphisms.....	886
14.1.2 Representably Full Morphisms.....	887
14.1.3 Representably Fully Faithful Morphisms .....	888
14.1.4 Morphisms Representably Faithful on Cores .....	889
14.1.5 Morphisms Representably Full on Cores .....	890
14.1.6 Morphisms Representably Fully Faithful on Cores.....	891
14.1.7 Representably Essentially Injective Morphisms.....	892
14.1.8 Representably Conservative Morphisms.....	893
14.1.9 Strict Monomorphisms.....	893
14.1.10 Pseudomonic Morphisms.....	894
14.2 Epimorphisms in Bicategories.....	896
14.2.1 Corepresentably Faithful Morphisms.....	896
14.2.2 Corepresentably Full Morphisms.....	897
14.2.3 Corepresentably Fully Faithful Morphisms .....	898
14.2.4 Morphisms Corepresentably Faithful on Cores .....	901
14.2.5 Morphisms Corepresentably Full on Cores .....	901
14.2.6 Morphisms Corepresentably Fully Faithful on Cores .....	902
14.2.7 Corepresentably Essentially Injective Morphisms.....	904
14.2.8 Corepresentably Conservative Morphisms .....	904
14.2.9 Strict Epimorphisms.....	905
14.2.10 Pseudoepic Morphisms .....	906
14.A Other Chapters.....	908

**Part VII Extra Part** **910**

<b>15 Notes .....</b>	<b>911</b>
15.1 TikZ Code for Commutative Diagrams.....	911
15.1.1 Product Diagram With Circular Arrows.....	912
15.1.2 Coproduct Diagram With Circular Arrows.....	914
15.1.3 Cube Diagram.....	916
15.1.4 Cube Diagram With Labelled Faces.....	917
15.1.5 Pentagon Diagram .....	920
15.1.6 Hexagon Diagram.....	921
15.1.7 Double Square Diagram .....	922
15.1.8 Double Hexagon Diagram.....	924
15.2 Retired Tags .....	926
15.2.1 Relations.....	926
15.2.2 Pointed Sets .....	930
15.2.3 Tensor Products of Pointed Sets .....	930
15.2.4 Categories.....	931

15.3 Miscellany.....	932
15.3.1 List of Things To Explore/Add.....	932
15.A Other Chapters.....	978
Index of Notation.....	991
Index of Set Theory.....	995
Index of Category Theory.....	999
Index of Higher Category Theory .....	1002

# **Part I**

# **Preliminaries**

# Chapter 1

## Introduction

**02H3** This chapter contains some general information about the Clowder Project.

### Contents

---

1.1	Introduction.....	2
1.1.1	Project Description and Goals.....	2
1.1.2	Navigating the Clowder Project .....	4
1.1.3	Prerequisites/Assumed Background.....	5
1.1.4	Community Engagement, Contributions and Col- laboration.....	5
1.1.5	Frequently Asked Questions.....	7
1.1.6	Goodies.....	8
1.2	Project Overview.....	9
1.2.1	Content and Scope .....	9
1.2.2	Style.....	10
1.2.3	Infrastructure and Technical Implementation ....	11
1.3	Lists.....	12
1.3.1	List of Omitted Proofs.....	12
1.3.2	List of Missing Examples.....	16
1.3.3	List of Questions.....	17
1.3.4	List of Gaps in the Category Theory Literature....	18
1.A	Other Chapters.....	22

---

**01WL 1.1 Introduction**

**01WM 1.1.1 Project Description and Goals**

In short, the Clowder Project is an online reference work and wiki for category theory and mathematics that aims to essentially become a Stacks Project for category theory.

The project arose from a desire to improve upon a number of issues with the existing category theory literature, as well as fill several gaps in it.

In this section, we list and discuss the goals of the Clowder Project.

**01WN 1.1.1.1 Provide a Unified and Complete Reference for Category Theory**

The category theory literature is at times rather fragmented, and often it takes a long while for book-long treatments on a given subject to appear.

For example, although the theory of bicategories dates back to the late 1960s, it was not until 2020 that the subject would receive its first textbook in the topic, namely [JY21].

The Clowder Project aims to bridge this gap, providing a complete overview of the foundational material on category theory (see also [Section 1.2.1](#)).

**01WP 1.1.1.2 Gather Hard to Find Results**

As an extension of the previous goal, the Clowder Project also aims to gather in a single place results that are hard to find in the literature. These tend to be recorded only on original sources, which often means papers, notes or theses from the 1970s.

Since the Clowder Project is organized as a wiki, it becomes rather easy to search and find such results, as one merely needs to go to the page for a given concept and then look at the properties listed there.

**01WQ 1.1.1.3 Elaborate on Details That Are Often Left Out**

Another goal of the Clowder Project is to include all kinds of details and intuitions that often don't make their way into textbooks, papers, monographs, etc.

For instance, one sometimes finds claims that a given diagram commutes and that it is "easy" to fill in the details. This also tends to happen particularly when the details are rather unwieldy.

One of the goals of the Clowder Project is to provide such proofs in great detail, including discussions of technical results, even when these are indeed "obvious".

**01WR 1.1.1.4 Homogenize Conventions, Notation, and Terminology**

Another issue with practice in the field is that there are often a number of conflicting conventions, notations, and terminology.

Being organized as a comprehensive and encyclopedic wiki, the Clowder Project tries to homogenize these conventions, notations, and terminology.

**01WS 1.1.1.5 Fill Gaps in the Category Theory Literature**

There are quite a few significant gaps in the category theory literature, some of which we hope to fill with the Clowder Project. For a list of (some of) these gaps, see [Section 1.3.4](#).

**01WT 1.1.1.6 Provide a Citable Reference for All Kinds of Results**

It is a common situation to require a well-known result for a paper. Although proving it might be straightforward, it is often more convenient to cite a reference instead. Finding such a reference, however, may be hard and/or time-consuming.

With its encyclopedic nature, the Clowder Project hopes to serve as that convenient reference.

**01WU 1.1.2 Navigating the Clowder Project**

Hopefully, it should be intuitive to navigate through the web version of project. Nevertheless, here we mention a couple things that might be useful to know.

**01WV 1.1.2.1 Preferences**

You can change the font of the site, the style of the PDFs, as well as turn on dark mode by clicking the gear button located at the top right corner of the page.

**01WW 1.1.2.2 Large Diagrams and the Zoom in Feature**

This work features many diagrams that are unfortunately a bit too large to be comfortably legible in their native size.

To compensate for this, it's possible to click on them to expand their size by 200%.

In addition, you may also right-click on diagrams and then select "Open image in new tab" to allow for even higher amounts of zoom.

**01WX 1.1.2.3 PDF Styles**

The PDFs for each chapter as well as for the whole book are generated using twelve different styles, as summarised in the following table:

Typeface	Theorem Environments
Alegreya Sans	tcbthm
Alegreya	tcbthm
EB Garamond	tcbthm
Crimson Pro	tcbthm
XCharter	tcbthm
Computer Modern	tcbthm
Alegreya Sans	amsthm
Alegreya	amsthm
EB Garamond	amsthm
Crimson Pro	amsthm
XCharter	amsthm
Computer Modern	amsthm

The default style uses Alegreya Sans and tcbthm.

### 01WY 1.1.3 Prerequisites/Assumed Background

The Clowder Project assumes at least a background on basic category theory corresponding to e.g. [Rie16], as well as some comfort in working with category-theoretic notions.

In particular, it should be viewed as a reference work/wiki, and *not* as a textbook. This, however, doesn't mean it shouldn't be pedagogical. Indeed, a number of stylistic choices are made aiming to make the material as easily digestible as possible.

For an outline of several introductory references for different topics in category theory, see A Guide to the Literature.

### 01WZ 1.1.4 Community Engagement, Contributions and Collaboration

All kinds of feedback and contributions to the Clowder Project are extremely welcome: pointing out typos, errors, historical remarks, references, layout of webpages, spelling errors, improvements to the overall structure, missing lemmas, etc.

The Clowder Project has an [official Discord server](#) in which people can ask questions, carry out discussions and give feedback. Please join it if you'd like to contribute to the Clowder Project. Alternatively, you may also reach out to the project maintainer at [emily.de.oliveira.santos.tmf@gmail.com](mailto:emily.de.oliveira.santos.tmf@gmail.com).

**02CB 1.1.4.1 How to Contribute**

There's a number of ways to contribute to the Clowder Project, some of which will be detailed a bit below. However, please keep in mind that they are not just examples, and are most definitely not meant to be exhaustive.

If there's another way in which you'd like to contribute, by all means feel free to drop by the project's Discord (or, alternatively, reach out to the project maintainer).

**01X0 1.1.4.2 Ways to Contribute: Missing Proofs**

There is a large number of missing proofs in the project, ranging from trivial proofs to simple lemmas to more involved results.

Missing proofs are listed in [Section 1.3.1](#).

**Note:** The following chapters are undergoing revision. If you're interested in contributing, please disregard them for now:

- Relations
- Constructions With Relations
- Conditions on Relations
- Categories
- Constructions With Monoidal Categories
- Types of Morphisms in Bicategories

**01X1 1.1.4.3 Ways to Contribute: Missing Examples**

New examples to the Clowder Project are always welcome. These could be examples illustrating a new concept, examples showing why certain conditions are necessary in a given proof, counterexamples to be aware of, etc.

Some examples which would be particularly nice to have in Clowder are listed in [Section 1.3.2](#). Please do keep in mind however that *all examples are welcome*, even if they fall outside the examples listed in [Section 1.3.2](#).

**01X2 1.1.4.4 Ways to Contribute: Questions**

A number of questions appear throughout the Clowder Project; tackling these would be an amazing way to contribute to the project.

The questions appearing throughout the Clowder Project are listed in [Section 1.3.3](#).

**02CC 1.1.5 Frequently Asked Questions****02CD 1.1.5.1 How does Clowder differ from the nLab?**

Clowder is meant to be much more comprehensive than the nLab, which includes even filling a number of gaps in the category theory literature. Additionally, it also has a different set of goals and stylistic choices. For a more in-depth explanation, see [Section 1.1.1](#).

**02CE 1.1.5.2 Why not just use the nLab instead?**

There are a number of reasons why Clowder was built as a separate project, instead of e.g. just editing the nLab:

02CF 1. *Curation.* All content on Clowder is personally curated by the project maintainer. This ensures an even quality to everything in the project.

02CG 2. *Cohesion.* As a consequence of [Item 1](#), the Clowder Project ends up being much more cohesive than the nLab, having a clear and coherent organization, consistent notation and conventions, as well as a consistent style.

02CH 3. *Referenceability.* Clowder employs Gerby's Tag system, meaning that every citable statement in Clowder (e.g. definitions, examples, constructions, propositions, remarks, even individual items in lists, etc.) carries a corresponding tag.

This makes the project easy to cite and reference, since although the numbering of e.g. a given definition may change, its associated tag will forever be the same. See also [Clowder — The Tag System](#).

02CJ 4. *Crowdsourcing and Crowdfunding.* Clowder is meant to be a crowd-funded project in which the community can help directly finance its development. As a result, the project has a dedicated project maintainer whose role is to continuously take care of the project, coordinating contributions, developing infrastructure, and expanding the content of the project.

02CK 5. *Infrastructure.* The Clowder Project makes use of several very specific features which simply wouldn't be possible to implement in the nLab. This includes:

02CL (a) An elaborate [fork](#) of [gerby-website](#), implementing a variety of new features and quality-of-life additions.

02CM (b) Another elaborate [fork](#), this time of [Gerby](#) (which is itself a fork of [plasTeX](#)), implement a number of similarly needed features for the website to work as intended.

See [Section 1.2.3.2](#) for a (slightly) more in-depth description of the features and additions that have been created specifically for Clowder.

### 01X3 1.1.6 Goodies

In this section we list a few sample nice results and things from the Clowder Project.

#### 01X4 1.1.6.1 General Utility

- [Section 15.1](#) contains several `tikz-cd` snippets producing somewhat-hard-to-draw diagrams. Examples include cube, pentagon, and hexagon diagrams, as well as e.g. co/product diagrams with perfectly circular arrows.

#### 01X5 1.1.6.2 Set Theory Through a Categorical Lens

Sets:

- [Section 4.4.7](#) contains a discussion of internal Homs in powersets viewed as categories.
- More generally, [Section 4.4](#) discusses several properties of powersets that are analogous to those of presheaf categories.
- ?? discusses the adjoint triple  $f_* \dashv f^{-1} \dashv f_!$  between  $\mathcal{P}(X)$  and  $\mathcal{P}(Y)$  induced by a function  $f: X \rightarrow Y$ .
- [Section 4.6.4](#) constructs a kind of “six functor formalism for (power)sets”.
- Monoidal Structures on the Category of Sets contains explicit proofs that product/coproduct of sets form a monoidal structure.
- [Section 5.1.10](#) gives a completely 1-categorical proof of the universal property of  $(\text{Sets}, \times, \text{pt})$ .

Pointed Sets:

- Tensor Products of Pointed Sets constructs several tensor products of pointed sets, including a few unusual ones giving rise to skew monoidal structures on  $\text{Sets}_*$ .
- [Section 7.5.10](#) gives a completely 1-categorical proof of the universal property of  $(\text{Sets}_*, \wedge, S^0)$ .
- [Proposition 7.5.12.1.1](#) contains a description of comonoids in  $\text{Sets}_*$  with respect to  $\wedge$ .

Relations:

- [Section 8.5](#) contains a discussion of several properties of the 2-category of relations like descriptions of internal adjunctions and internal monads.
- [Sections 8.8 and 8.9](#) contains a discussion of two skew monoidal structures on the category  $\text{Rel}(A, B)$  of relations from a set  $A$  to a set  $B$ .
- [Old Tag 15.2.1.1.8](#) contains a description of left/right Kan extensions and lifts internal to the 2-category of relations.

#### [01X6](#) 1.1.6.3 Category Theory

- Categories contains a description of several properties of functors, including somewhat lesser known ones such as dominant functors or pseudoepic functors.

### [01X7](#) 1.2 Project Overview

#### [01X8](#) 1.2.1 Content and Scope

In this section, we outline what content is expected to be covered in the Clowder Project.

#### [01X9](#) 1.2.1.1 Elementary Category Theory

First and foremost, the Clowder Project aims to cover the foundations of category theory. This comprises all the usual topics treated in basic textbooks in category theory, such as [\[Mac98\]](#) or [\[Rie16\]](#), like adjunctions, co/limits, Kan extensions, co/ends, monoidal categories, etc.

#### [01XA](#) 1.2.1.2 Variants of Category Theory

Second, the Clowder Project aims to cover variants of category theory such as internal, fibred, or enriched category theory. The literature on these topics is often quite scattered and scarce, and so having a comprehensive discussion of them in Clowder aims to fill a large gap in the literature. See also [Gap 1.3.4.1.14](#).

#### [01XB](#) 1.2.1.3 Higher Category Theory

Third, a detailed presentation of the theories of bicategories and double categories is planned, along with *some* material on tricategories.

Bicategories are another topic for which the literature is rather scattered, and, for some topics, scarce. As mentioned in the introduction, only recently has a proper textbook on bicategories appeared, [\[JY21\]](#). Moreover, one

finds several gaps in the literature, with a number of important results missing. As one particular example, one could look at the theory of 2-dimensional co/ends, in which case a comprehensive treatment based upon lax/oplax/pseudo dinatural transformations seems to be missing.

All of the elementary and not-so-elementary topics in the theory of bicategories are planned to appear in Clowder, and the same holds true for the theory of double categories.

#### 01XC 1.2.1.4 $\infty$ -Categories

Lastly, some material on  $\infty$ -categories is planned, although the precise scope of this remains to be defined. Ideally, this would include both model categories as well as synthetic and concrete models for  $\infty$ -categories (e.g. quasi-categories, complete Segal spaces, cubical quasicategories, etc.).

In this way, we view Clowder as a good *complement* to [Lur25].

#### 01XD 1.2.1.5 Other Topics

Occasionally, material on topics not a-priori related to category theory will be included. This may be done for a variety of reasons, including:

- Illustrating general theory.
- Comparison with classical concepts, such as e.g. ionads vs. topological spaces.
- Providing a more consistent and unified treatment of a particular topic, with hyperlinks to relevant concepts or examples.

#### 01XE 1.2.2 Style

The Clowder Project makes several unusual stylistic choices, aligned with its goals.

#### 01XF 1.2.2.1 Presentation of Topics

The presentation of topics is encyclopedic, non-linear, and sometimes idiosyncratic.

In particular, there's some amount of repetition throughout the project. This is a result of simultaneously wanting to cover as much material as possible while still allowing Clowder to be used as an online reference work/wiki.

**01XG 1.2.2.2 Provable Items Come With Proofs**

Every proposition, theorem, lemma, etc. needs to come with a proof. In case a proof has not been written yet, it shall read as “Omitted”. This is to ensure results without proof are clearly labelled as such.

**01XH 1.2.2.3 Proper Justification of Proofs**

Every proof must read either “Omitted” or be properly justified, no matter how trivial the details are.

Expressions like “it is clear that”, “it is straightforward to show that”, “it is obvious”, etc. inside proofs should not be used.

**01XJ 1.2.3 Infrastructure and Technical Implementation****01XK 1.2.3.1 Removed Features (in Comparison With the Stacks Project)**

A few features present in the general infrastructure of the Stacks Project were removed in Clowder, including:

- 02CP     1. The python back-end, in favour of static pages.
- 2. The comment system, as a result of the static nature of the website.

**01XL 1.2.3.2 Gerby and the Tags System**

Clowder is built using [Gerby](#), similarly to the [Stacks Project](#). However, a number of additional features and quality-of-life additions not implemented in plasTeX or Gerby were required by Clowder, including:

- 02CQ     1. Clowder uses `tcbtheorem`-like environments, which affects the placement of footnotes (which are often used).
- 02CR     2. Clowder implements a `dangerous bend` symbol to help visually highlight warnings ([example](#)).
- 02CS     3. There are a few aesthetic changes in Clowder’s HTML/CSS structure, including font selection as well as a dark mode.
- 02CT     4. `tikz-cd` diagrams are very frequently used, and they need to be separately compiled and converted to `svg` files.
- 02CU     5. Code in Clowder can be copied easily using a “Copy” button, with code for bibliography entries also having proper syntax highlighting ([example](#)).
- 02CV     6. Clowder is automatically built using [GitHub actions](#).


**02CW** 7. Non-sectioning tags are rendered differently and shown in context ([example](#)).

These have been implemented using a [fork](#) of Gerby along with a few build scripts.

#### **01XM 1.2.3.3 Placeholder Symbols and Future Style**

Currently, a number of macros have been defined using placeholder symbols, and look very ugly as a result.

They will eventually be replaced with proper symbols coming from the math fonts of [Hundar](#), a free and open-source typeface project currently being worked on.



You can find more details about Hundar at its [GitHub repository](#) or [website](#).

## **01XN 1.3 Lists**

### **01CT 1.3.1 List of Omitted Proofs**

Не так благотворна истина, как зловредна ее видимость.

Даниил Данковский

Truth does not do as much good in the world as the appearance of truth does evil.

*Daniil Dankovsky*

There's a very large number of omitted proofs throughout these notes. In this section we list them in order of decreasing importance.

- If a proof relies on material that has yet to be developed on Clowder, we mark it by a sign. If you're interested in contributing, please disregard those for now.

- The following chapters are undergoing revision. If you're interested in contributing, please disregard them for now:
  - Relations
  - Constructions With Relations
  - Conditions on Relations
  - Categories
  - Constructions With Monoidal Categories
  - Types of Morphisms in Bicategories
- This list is under construction.

01CU

**REMARK 1.3.1.1.1 ► OMITTED PROOFS TO ADD**

Proofs that *need* to be added at some point:

- Extra proof of [Theorem 7.5.10.1.1](#) using the machinery of presentable categories, following Maxime Ranzi's answer to [MO 466593](#) .
- Horizontal composition of natural transformations is associative: [Item 2 of Proposition 11.9.5.1.4](#).
- Fully faithful functors are essentially injective: [Item 4 of Proposition 11.6.3.1.2](#).

Proofs that *would be very nice* to be added at some point:

- Properties of pseudomonadic functors: [Proposition 11.7.4.1.2](#) .
- Characterisation of fully faithful functors: [Item 1 of Proposition 11.6.3.1.2](#).
- The quadruple adjunction between categories and sets: [Proposition 11.3.1.1.1](#).
- $F_*$  faithful iff  $F$  faithful: [Item 2 of Proposition 11.6.1.1.2](#).
- Properties of groupoid completions: [Proposition 11.4.3.1.4](#).
- Properties of cores: [Proposition 11.4.4.1.5](#).
- Rel is isomorphic to the category of free algebras of the powerset monad: [Proposition 8.5.18.1.1](#) .
- Non/existence of left Kan extensions in **Rel**:
  - ?? of ??.
  - ?? of ??.

- Non/existence of left Kan lifts in **Rel**:
  - ?? of ??.
  - ?? of ??.

Proofs that *would be nice* to be added at some point:

- Properties of posetal categories: [Proposition 11.2.7.1.2](#).
- Injective on objects functors are precisely the isocofibrations in  $\text{Cats}_2$ : [Item 1 of Proposition 11.8.1.1.2](#) .
- Characterisations of monomorphisms of categories: [Item 1 of Proposition 11.7.2.1.2](#).
- Epimorphisms of categories are surjective on objects: [Item 2 of Proposition 11.7.3.1.2](#).
- Properties of pseudoepic functors: [Proposition 11.7.5.1.2](#) .

Proofs that *would be nice but not essential* to be added at some point:

- Proof that  $(\text{Sets}, \coprod, \emptyset, \times, \text{pt})$  is a symmetric bimonoidal category: [Item 15 of Proposition 4.1.3.1.4](#) .
- Proof that  $(\text{Sets}, \coprod, \emptyset, \times, \text{pt})$  is a symmetric bimonoidal category: [Proposition 5.3.5.1.1](#) .
- Proof that  $(\text{Sets}, \times_X, X)$  is a symmetric monoidal category: [Item 11 of Proposition 4.1.4.1.7](#) .
- Proof that  $(\text{Sets}, \coprod, \emptyset)$  is a symmetric monoidal category: [Item 6 of Proposition 4.2.3.1.4](#) .
- Proof that  $(\text{Sets}, \coprod_X, X)$  is a symmetric monoidal category: [Item 8 of Proposition 4.2.4.1.8](#) .

Proofs that have been (temporarily) omitted because they are “clear”, “straightforward”, or “tedious”:

- Properties of pushouts of sets:
  - Associativity: [Item 3 of Proposition 4.2.4.1.8](#).
  - Unitality: [Item 5 of Proposition 4.2.4.1.8](#).
  - Commutativity: [Item 6 of Proposition 4.2.4.1.8](#).
  - Pushout of sets over the empty set recovers the coproduct of sets: [Item 7 of Proposition 4.2.4.1.8](#).

- Properties of coequalisers of sets:
  - Associativity: [Item 1 of Proposition 4.2.5.1.7](#).
  - Unitality: [Item 2 of Proposition 4.2.5.1.7](#).
  - Commutativity: [Item 3 of Proposition 4.2.5.1.7](#).
  - Interaction with composition: [Item 4 of Proposition 4.2.5.1.7](#).
- Properties of set differences:
  - [Item 4 of Proposition 4.3.10.1.2](#).
  - [Item 11 of Proposition 4.3.10.1.2](#).
  - [Item 13 of Proposition 4.3.10.1.2](#).
  - [Item 15 of Proposition 4.3.10.1.2](#).
- Complements and characteristic functions: [Item 4 of Proposition 4.3.11.1.2](#).
- Properties of symmetric differences:
  - [Item 1 of Proposition 4.3.12.1.2](#).
  - [Item 16 of Proposition 4.3.12.1.2](#).
- Properties of direct images:
  - Functoriality: [Item 1 of Proposition 4.6.1.1.5](#).
  - Interaction with coproducts: [Item 15 of Proposition 4.6.1.1.5](#).
  - Interaction with products: [Item 16 of Proposition 4.6.1.1.5](#).
- Properties of inverse images:
  - Functoriality: [Item 1 of Proposition 4.6.2.1.3](#).
  - Interaction with coproducts: [Item 15 of Proposition 4.6.2.1.3](#).
  - Interaction with products: [Item 16 of Proposition 4.6.2.1.3](#).
- Properties of codirect images:
  - Functoriality: [Item 1 of Proposition 4.6.3.1.7](#).
  - Lax preservation of colimits: [Item 10 of Proposition 4.6.3.1.7](#).
  - Interaction with coproducts: [Item 14 of Proposition 4.6.3.1.7](#).
  - Interaction with products: [Item 15 of Proposition 4.6.3.1.7](#).

- Left distributor of  $\times$  over  $\coprod$  is a natural isomorphism: [Definition 5.3.1.1.1](#).
- Right distributor of  $\times$  over  $\coprod$  is a natural isomorphism: [Definition 5.3.2.1.1](#).
- Left annihilator of  $\times$  is a natural isomorphism: [Definition 5.3.3.1.1](#).
- Right annihilator of  $\times$  is a natural isomorphism: [Definition 5.3.4.1.1](#).
- Properties of wedge products of pointed sets:
  - Associativity: [Item 2 of Proposition 6.3.3.1.4](#).
  - Unitality: [Item 3 of Proposition 6.3.3.1.4](#).
  - Commutativity: [Item 4 of Proposition 6.3.3.1.4](#).
  - Symmetric monoidality: [Item 5 of Proposition 6.3.3.1.4](#).
- Properties of pushouts of pointed sets:
  - Interaction with coproducts: [Item 5 of Proposition 6.3.4.1.4](#).
  - Symmetric monoidality: [Item 6 of Proposition 6.3.4.1.4](#).

### 01V7 1.3.2 List of Missing Examples

Adding new examples is always welcome! In this section, we list some subjects and sections which could do with more examples:

01V8

#### REMARK 1.3.2.1.1 ► MISSING EXAMPLES TO ADD

Potentially interesting examples to add include, but are definitely not limited to:

- Examples of 2-categorical monomorphisms in **Rel**, following [Old Tag 15.2.1.1.12](#).
- Examples of 2-categorical epimorphisms in **Rel**, following [Old Tag 15.2.1.1.25](#).
- Examples of left Kan extensions and left Kan lifts in **Rel**.
- Examples of functors satisfying the conditions described in Categories.

**01V9 1.3.3 List of Questions**

There's a number of questions listed throughout this project. Here we collect them in a single place.

01VA

**REMARK 1.3.3.1.1 ► QUESTIONS TO ANSWER**

On relations:

- [Old Tag 15.2.1.1.10](#), on better characterisations of corepresentably full morphisms in **Rel**. This question also appears as [[MO 467527](#)].
- ??, seeking a characterisation of which left Kan extensions exist in **Rel**. This question also appears as [[MO 461592](#)].
- ??, seeking a characterisation of which left Kan lifts exist in **Rel**. This question also appears as [[MO 461592](#)].

On categories:

- [Question 11.6.2.1.4](#), seeking a better characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  to always be full. This question also appears as [[MO 468121b](#)].
- [Question 11.6.4.1.4](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be conservative. This question also appears as [[MO 468121a](#)].
- [Question 11.6.5.1.2](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be essentially injective. This question also appears as [[MO 468121a](#)].
- [Question 11.6.6.1.2](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be essentially surjective. This question also appears as [[MO 468121a](#)].
- [Question 11.7.1.1.4](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be dominant. This question also appears as [[MO 468121a](#)].
- [Question 11.7.2.1.4](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be monic. This question also appears as [[MO 468121a](#)].
- [Question 11.7.3.1.4](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be epic. This question also appears as [[MO 468121a](#)].

- [Question 11.7.5.1.6](#), seeking a characterisation of necessary and sufficient conditions on  $F$  for  $F^*$  or  $F_*$  to be pseudoepic. This question also appears as [[MO 468121a](#)].
- [Question 11.7.5.1.4](#), seeking a characterisation of pseudoepic functors. This question also appears as [[MO 321971](#)].
- [Question 11.7.5.1.5](#), which asks whether a pseudomonic and pseudoepic functor must necessarily be an equivalence of categories. This question also appears as [[MO 468334](#)].
- [Question 11.8.4.1.3](#), seeking a characterisation of functors representably faithful on cores.
- [Question 11.8.5.1.3](#), seeking a characterisation of functors representably full on cores.
- [Question 11.8.6.1.3](#), seeking a characterisation of functors representably fully faithful on cores.
- [Question 11.8.7.1.3](#), seeking a characterisation of functors corepresentably faithful on cores.
- [Question 11.8.8.1.3](#), seeking a characterisation of functors corepresentably full on cores.
- [Question 11.8.9.1.3](#), seeking a characterisation of functors corepresentably fully faithful on cores.

#### 01XP 1.3.4 List of Gaps in the Category Theory Literature

The Clowder Project aims to address several significant gaps in the existing literature on category theory, as detailed below. See also [[MO 494959](#)].

01XQ

##### GAP 1.3.4.1.1 ► THE TENSOR PRODUCT OF PRESENTABLE CATEGORIES

Even though its analogue for  $\infty$ -categories has for years been a widely used tool<sup>1</sup>, a comprehensive treatment of the tensor product of presentable categories seems to be currently missing.

<sup>1</sup>See [[MO 490557](#)].

01XR

##### GAP 1.3.4.1.2 ► EXPLICIT DESCRIPTIONS OF CO/LIMITS OF CATEGORIES

An exhaustive concrete description of the various limits and colimits of categories, including 2-dimensional ones, is missing.

01XS

**GAP 1.3.4.1.3 ► DINATURAL TRANSFORMATION CO/CLASSIFIERS**

There seems to be no unified presentation of dinatural transformation co/classifiers in the literature. These are characterised by isomorphisms of the form

$$\text{Nat}(F, G) \cong \text{DiNat}(\Gamma(F), G),$$

and were originally studied in Dubuc–Street’s paper introducing dinatural transformations, [DS06].

Even though these arguably form a fundamental piece of the framework of co/end calculus, it seems that all foundational treatments that followed after ended up not covering this concept.

01XT

**GAP 1.3.4.1.4 ► THE TENSOR PRODUCT OF SYMMETRIC MONOIDAL CATEGORIES**

The tensor product of symmetric monoidal categories had been a missing concept from the literature for years. Recently, [GJO24] covered the case of permutative categories. It would be nice, however, to also have a treatment of the non-strict case available.

01XU

**GAP 1.3.4.1.5 ► A COMPREHENSIVE TREATMENT OF THE THEORY OF PROMONOIDAL CATEGORIES**

A comprehensive and exhaustive treatment of the theory of promonoidal categories is currently missing. There are several important notions undefined, like:

- Promonoidal profunctors.
- Dualisability internal to a promonoidal category.
- Invertibility internal to a promonoidal category.

Moreover, it would be nice to record how promonoidal categories may be viewed as categorifications of “hypermonoids” (i.e. monoids in Rel).

01XV

**GAP 1.3.4.1.6 ► A COMPREHENSIVE TREATMENT OF THE THEORY OF MULTICATEGORIES**

A comprehensive and exhaustive treatment of the theory of multicategories is currently missing. There are several important notions undefined, like:

- Co/limits internal to multicategories.

See [MO 484647].

01XW

**GAP 1.3.4.1.7 ► A COMPENDIUM OF EXAMPLES OF 2-CATEGORICAL NOTIONS**

It would be nice to have an extensive collection of examples of what a given 2-categorical notion looks like in a 2-category. For instance, it would be nice to explicitly list what internal adjunctions look like in **Rel**, **Span**, **Prof**, etc.

See [Section 8.5](#) for a concrete example of what is meant by this gap.

01XX

**GAP 1.3.4.1.8 ► CENTRES AND TRACES OF CATEGORIES**

The literature on centres and traces of categories is really small. There are lots of results missing<sup>1</sup> and very few worked examples<sup>2</sup>.

<sup>1</sup>E.g. There's a certain interaction between traces of categories and Leinster's eventual image.

<sup>2</sup>E.g. what is the trace of Connes's cycle category? Such a computation doesn't seem to be available.

01XY

**GAP 1.3.4.1.9 ► NATURAL COTRANSFORMATIONS**

Natural transformations satisfy an isomorphism of the form

$$\text{Nat}(F, G) \cong \int_{A \in C} \text{Hom}_{\mathcal{D}}(F_A, G_A).$$

It is then exceedingly natural to define *natural cotransformations* via an isomorphism of the form

$$\text{CoNat}(F, G) \cong \int^{A \in C} \text{Hom}_{\mathcal{D}}(F_A, G_A)$$

and study their properties. This generalises traces of categories, since we have

$$\text{Tr}(C) = \text{CoNat}(\text{id}_C, \text{id}_C),$$

much like  $\text{Z}(C) = \text{Nat}(\text{id}_C, \text{id}_C)$ .

01XZ

**GAP 1.3.4.1.10 ► A COMPREHENSIVE TREATMENT OF ISBELL DUALITY**

There are several results, notions, and examples in the theory of Isbell duality missing from the literature, and a truly comprehensive treatment is still lacking.<sup>1</sup>

<sup>1</sup>For instance, there appears to be no mention of the duality pairings

$$\begin{aligned} \text{Spec}(F) \boxtimes F &\rightarrow \text{Tr}(C), \\ \mathcal{F} \boxtimes \mathcal{O}(\mathcal{F}) &\rightarrow \text{Tr}(C) \end{aligned}$$

in the currently available literature.

01Y0

The currently available treatments of 2-dimensional co/ends are unsatisfactory.<sup>1</sup>

<sup>1</sup>For instance, none of them define 2-dimensional co/ends via 2-dimensional dinatural transformations and then go on to develop a general theory from there.

01Y1

**GAP 1.3.4.1.12 ► A COMPREHENSIVE TREATMENT OF FACTORISATION SYSTEMS**

A comprehensive treatment of factorisation systems is currently missing; see [MO 495003].

02C5

**GAP 1.3.4.1.13 ► PROOFS OF COHERENCE THEOREMS FOR STRING DIAGRAMS**

Several proofs of coherence theorems for string diagrams currently have gaps; see [MO 497309].

02CX

**GAP 1.3.4.1.14 ► A COMPREHENSIVE TREATMENT OF VARIANTS OF CATEGORY THEORY**

The currently available treatments of variants of category theory such as fibred category theory, enriched category theory, or internal category theory are unsatisfactory for a number of reasons.

Ideally, there should be a comprehensive and (simultaneously) approachable treatment for these topics. See also [MO 497419].

# Appendices

## 1.A Other Chapters

### Preliminaries

1. Introduction
2. A Guide to the Literature

### Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

### Relations

8. Relations
9. Constructions With Relations
10. Conditions on Relations

### Categories

11. Categories
12. Presheaves and the Yoneda Lemma

### Monoidal Categories

13. Constructions With Monoidal Categories

<b>Bicategories</b>	<b>Extra Part</b>
14. Types of Morphisms in Bicategories	15. Notes

## Chapter 2

# A Guide to the Literature

**02H3** This chapter contains some material about category theory literature.

### Contents

---

2.1	Elementary Category Theory .....	24
2.1.1	Textbooks .....	24
2.A	Other Chapters .....	24

---

**02C9** **2.1 Elementary Category Theory**

**02CA** **2.1.1 Textbooks**

## Appendices

### 2.A Other Chapters

#### Preliminaries

- 1. Introduction
- 2. A Guide to the Literature

#### Sets

- 3. Sets
- 4. Constructions With Sets
- 5. Monoidal Structures on the Category of Sets
- 6. Pointed Sets

#### 7. Tensor Products of Pointed Sets

#### Relations

- 8. Relations
- 9. Constructions With Relations
- 10. Conditions on Relations

#### Categories

- 11. Categories
- 12. Presheaves and the Yoneda Lemma

#### Monoidal Categories

13. Constructions With Monoidal Categories

gories

**Bicategories**

**Extra Part**

14. Types of Morphisms in Bicate-

15. Notes

## **Part II**

# **Sets**

# Chapter 3

## Sets

02H3 This chapter (will eventually) contain material on axiomatic set theory, as well as a couple other things.

### Contents

---

3.1	Sets and Functions.....	27
3.1.1	Functions.....	27
3.2	The Enrichment of Sets in Classical Truth Values.....	29
3.2.1	(-2)-Categories.....	29
3.2.2	(-1)-Categories.....	29
3.2.3	0-Categories.....	33
3.2.4	Tables of Analogies Between Set Theory and Category Theory.....	34
3.A	Other Chapters.....	36

---

0001 **3.1 Sets and Functions**

0002 **3.1.1 Functions**

0003 **DEFINITION 3.1.1.1 ► FUNCTIONS**

A **function** is a functional and total relation.

0004 **NOTATION 3.1.1.2 ► ADDITIONAL NOTATION FOR FUNCTIONS**

Throughout this work, we will sometimes denote a function  $f: X \rightarrow Y$  by

$$f \stackrel{\text{def}}{=} \llbracket x \mapsto f(x) \rrbracket.$$

01Y2 1. For example, given a function

$$\Phi: \text{Hom}_{\text{Sets}}(X, Y) \rightarrow K$$

taking values on a set of functions such as  $\text{Hom}_{\text{Sets}}(X, Y)$ , we will sometimes also write

$$\Phi(f) \stackrel{\text{def}}{=} \Phi(\llbracket x \mapsto f(x) \rrbracket).$$

**01Y3** 2. This notational choice is based on the lambda notation

$$f \stackrel{\text{def}}{=} (\lambda x. f(x)),$$

but uses a “ $\mapsto$ ” symbol for better spacing and double brackets instead of either:

**01Y4** (a) Square brackets  $[x \mapsto f(x)]$ ;

**01Y5** (b) Parentheses  $(x \mapsto f(x))$ ;

hoping to improve readability when dealing with e.g.:

**01Y6** (a) Equivalence classes, cf.:

**01Y7** i.  $\llbracket [x] \mapsto f([x]) \rrbracket$

**01Y8** ii.  $\llbracket [x] \mapsto f([x]) \rrbracket$

**01Y9** iii.  $(\lambda [x]. f([x]))$

**01YA** (b) Function evaluations, cf.:

**01YB** i.  $\Phi(\llbracket x \mapsto f(x) \rrbracket)$

**01YC** ii.  $\Phi((x \mapsto f(x)))$

**01YD** iii.  $\Phi((\lambda x. f(x)))$

**01YE** 3. We will also sometimes write  $-$ ,  $-_1$ ,  $-_2$ , etc. for the arguments of a function. Some examples include:

**01YF** (a) Writing  $f(-_1)$  for a function  $f: A \rightarrow B$ .

**01YG** (b) Writing  $f(-_1, -_2)$  for a function  $f: A \times B \rightarrow C$ .

**01YH** (c) Given a function  $f: A \times B \rightarrow C$ , writing

$$f(a, -): B \rightarrow C$$

for the function  $\llbracket b \mapsto f(a, b) \rrbracket$ .

**01YJ** (d) Denoting a composition of the form

$$A \times B \xrightarrow{\phi \times \text{id}_B} A' \times B \xrightarrow{f} C$$

by  $f(\phi(-_1), -_2)$ .

01YK

4. Finally, given a function  $f: A \rightarrow B$ , we will sometimes write

$$\text{ev}_a(f) \stackrel{\text{def}}{=} f(a)$$

for the value of  $f$  at some  $a \in A$ .

For an example of the above notations being used in practice, see the proof of the adjunction

$$(A \times - \dashv \text{Hom}_{\text{Sets}}(A, -)): \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Hom}_{\text{Sets}}(A, -)} \end{array} \text{Sets},$$

stated in Item 2 of Proposition 4.1.3.1.4.

## 0005 3.2 The Enrichment of Sets in Classical Truth Values

### 0006 3.2.1 (-2)-Categories

0007

#### DEFINITION 3.2.1.1 ▶ (-2)-CATEGORIES

A **(-2)-category** is the “necessarily true” truth value.<sup>1,2,3</sup>

<sup>1</sup>Thus, there is only one (-2)-category.

<sup>2</sup>A  $(-n)$ -category for  $n = 3, 4, \dots$  is also the “necessarily true” truth value, coinciding with a (-2)-category.

<sup>3</sup>For motivation, see [BS10, p. 13].

### 0008 3.2.2 (-1)-Categories

0009

#### DEFINITION 3.2.2.1.1 ▶ (-1)-CATEGORIES

A **(-1)-category** is a classical truth value.

000A

#### REMARK 3.2.2.1.2 ▶ MOTIVATION FOR (-1)-CATEGORIES

<sup>1</sup>(-1)-categories should be thought of as being “categories enriched in (-2)-categories”, having a collection of objects and, for each pair of objects, a Hom-object  $\text{Hom}(x, y)$  that is a (-2)-category (i.e. trivial). As a result, a (-1)-category  $C$  is either:<sup>2</sup>

01D2

1. *Empty*, having no objects.

01D3

2. *Contractible*, having a collection of objects  $\{a, b, c, \dots\}$ , but with  $\text{Hom}_C(a, b)$  being a (-2)-category (i.e. trivial) for all  $a, b \in \text{Obj}(C)$ ,

forcing all objects of  $C$  to be uniquely isomorphic to each other.

Thus there are only two  $(-1)$ -categories up to equivalence:

**01D4** 1. The  $(-1)$ -category false (the empty one);

**01D5** 2. The  $(-1)$ -category true (the contractible one).

<sup>1</sup>For more motivation, see [BS10, p. 13].

<sup>2</sup>See [BS10, pp. 33–34].

#### 000B DEFINITION 3.2.2.1.3 ► THE POSET OF TRUTH VALUES

The **poset of truth values**<sup>1</sup> is the poset  $(\{\text{true}, \text{false}\}, \preceq)$  consisting of:

- *The Underlying Set.* The set  $\{\text{true}, \text{false}\}$  whose elements are the truth values true and false.
- *The Partial Order.* The partial order

$$\preceq: \{\text{true}, \text{false}\} \times \{\text{true}, \text{false}\} \rightarrow \{\text{true}, \text{false}\}$$

on  $\{\text{true}, \text{false}\}$  defined by<sup>2</sup>

$$\begin{aligned} \text{false} \preceq \text{false} &\stackrel{\text{def}}{=} \text{true}, \\ \text{true} \preceq \text{false} &\stackrel{\text{def}}{=} \text{false}, \\ \text{false} \preceq \text{true} &\stackrel{\text{def}}{=} \text{true}, \\ \text{true} \preceq \text{true} &\stackrel{\text{def}}{=} \text{true}. \end{aligned}$$

<sup>1</sup>Further Terminology: Also called the **poset of  $(-1)$ -categories**.

<sup>2</sup>This partial order coincides with logical implication.

#### 000C NOTATION 3.2.2.1.4 ► FURTHER NOTATION FOR THE POSET OF TRUTH VALUES

We also write  $\{t, f\}$  for the poset  $\{\text{true}, \text{false}\}$ .

#### 000D PROPOSITION 3.2.2.1.5 ► CARTESIAN CLOSEDNESS OF THE POSET OF TRUTH VALUES

The poset of truth values  $\{t, f\}$  is Cartesian closed with product given by<sup>1</sup>

$$\begin{array}{l} t \times t = t, \quad f \times t = f, \\ t \times f = f, \quad f \times f = f, \end{array} \quad \begin{array}{c|cc} \times & t & f \\ \hline t & t & f \\ f & f & f \end{array}$$

and internal Hom  $\text{Hom}_{\{t,f\}}$  given by the partial order of  $\{t, f\}$ , i.e. by

$$\begin{aligned}\text{Hom}_{\{t,f\}}(t, t) &= t, & \text{Hom}_{\{t,f\}}(f, t) &= t, \\ \text{Hom}_{\{t,f\}}(t, f) &= f, & \text{Hom}_{\{t,f\}}(f, f) &= t,\end{aligned}$$

$\text{Hom}_{\{t,f\}}$	t	f
t	t	t
f	t	f

<sup>1</sup>Note that  $\times$  coincides with the “and” operator, while  $\text{Hom}_{\{t,f\}}$  coincides with the logical implication operator.

#### PROOF 3.2.2.1.6 ► PROOF OF PROPOSITION 3.2.2.1.5

##### Existence of Products

We claim that the products  $t \times t$ ,  $t \times f$ ,  $f \times t$ , and  $f \times f$  satisfy the universal property of the product in  $\{t, f\}$ . Indeed, suppose we have diagrams of the form

$$\begin{array}{cccc} \begin{array}{c} p_1^1 \\ \swarrow \quad \searrow \\ t \xleftarrow{\text{pr}_1} t \times t \xrightarrow{\text{pr}_2} t \end{array} & \begin{array}{c} p_2^1 \\ \swarrow \quad \searrow \\ t \xleftarrow{\text{pr}_1} t \times f \xrightarrow{\text{pr}_2} f \end{array} & \begin{array}{c} p_3^1 \\ \swarrow \quad \searrow \\ f \xleftarrow{\text{pr}_1} f \times t \xrightarrow{\text{pr}_2} t \end{array} & \begin{array}{c} p_4^1 \\ \swarrow \quad \searrow \\ f \xleftarrow{\text{pr}_1} f \times f \xrightarrow{\text{pr}_2} f \end{array} \end{array}$$

where the  $\text{pr}_1$  and  $\text{pr}_2$  morphisms are the only possible ones (since  $\{t, f\}$  is posetal). We claim that there are unique morphisms making the diagrams

$$\begin{array}{cccc} \begin{array}{c} p_1^1 \\ \swarrow \quad \searrow \\ t \xleftarrow{\text{pr}_1} t \times t \xrightarrow{\text{pr}_2} t \\ \vdots \\ \exists! \end{array} & \begin{array}{c} p_2^1 \\ \swarrow \quad \searrow \\ t \xleftarrow{\text{pr}_1} t \times f \xrightarrow{\text{pr}_2} f \\ \vdots \\ \exists! \end{array} & \begin{array}{c} p_3^1 \\ \swarrow \quad \searrow \\ f \xleftarrow{\text{pr}_1} f \times t \xrightarrow{\text{pr}_2} t \\ \vdots \\ \exists! \end{array} & \begin{array}{c} p_4^1 \\ \swarrow \quad \searrow \\ f \xleftarrow{\text{pr}_1} f \times f \xrightarrow{\text{pr}_2} f \\ \vdots \\ \exists! \end{array} \end{array}$$

commute. Indeed:

- 01YL 1. If  $P_1 = t$ , then  $p_1^1 = p_2^1 = \text{id}_t$ , so there's a unique morphism from  $P_1$  to  $t$  making the diagram commute, namely  $\text{id}_t$ .
- 01YM 2. If  $P_1 = f$ , then  $p_1^1 = p_2^1$  are given by the unique morphism from  $f$  to  $t$ , so there's a unique morphism from  $P_1$  to  $t$  making the diagram commute, namely the unique morphism from  $f$  to  $t$ .
- 01YN 3. If  $P_2 = t$ , then there is no morphism  $p_2^2$ .
- 01YP 4. If  $P_2 = f$ , then  $p_1^2$  is the unique morphism from  $f$  to  $t$  while  $p_2^2 = \text{id}_f$ , so there's a unique morphism from  $P_2$  to  $f$  making the diagram commute, namely  $\text{id}_f$ .
- 01YQ 5. The proof for  $P_3$  is similar to the one for  $P_2$ .

01YR

6. If  $P_4 = t$ , then there is no morphism  $p_1^4$  or  $p_2^4$ .

01YS

7. If  $P_4 = f$ , then  $p_1^4 = p_2^4 = \text{id}_f$ , so there's a unique morphism from  $P_4$  to  $f$  making the diagram commute, namely  $\text{id}_f$ .

This finishes the existence of products part of the proof.

### Cartesian Closedness

We claim there's a bijection

$$\text{Hom}_{\{t,f\}}(A \times B, C) \cong \text{Hom}_{\{t,f\}}(A, \text{Hom}_{\{t,f\}}(B, C)),$$

natural in  $A, B, C \in \{t, f\}$ . Indeed:

- For  $(A, B, C) = (t, t, t)$ , we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times t, t) &\cong \text{Hom}_{\{t,f\}}(t, t) \\ &= \{\text{id}_{\text{true}}\} \\ &\cong \text{Hom}_{\{t,f\}}(t, t) \\ &\cong \text{Hom}_{\{t,f\}}(t, \text{Hom}_{\{t,f\}}(t, t)). \end{aligned}$$

- For  $(A, B, C) = (t, t, f)$ , we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times t, f) &\cong \text{Hom}_{\{t,f\}}(t, f) \\ &= \emptyset \\ &\cong \text{Hom}_{\{t,f\}}(t, f) \\ &\cong \text{Hom}_{\{t,f\}}(t, \text{Hom}_{\{t,f\}}(t, f)). \end{aligned}$$

- For  $(A, B, C) = (t, f, t)$ , we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times f, t) &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{Hom}_{\{t,f\}}(f, \text{Hom}_{\{t,f\}}(f, t)). \end{aligned}$$

- For  $(A, B, C) = (t, f, f)$ , we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times f, f) &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \text{Hom}_{\{t,f\}}(t, \text{Hom}_{\{t,f\}}(f, f)). \end{aligned}$$

- For  $(A, B, C) = (f, t, t)$ , we have

$$\begin{aligned}\text{Hom}_{\{t,f\}}(f \times t, t) &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong pt \\ &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{Hom}_{\{t,f\}}(f, \text{Hom}_{\{t,f\}}(t, t)).\end{aligned}$$

- For  $(A, B, C) = (f, t, f)$ , we have

$$\begin{aligned}\text{Hom}_{\{t,f\}}(f \times t, f) &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \text{Hom}_{\{t,f\}}(f, \text{Hom}_{\{t,f\}}(t, f)).\end{aligned}$$

- For  $(A, B, C) = (f, f, t)$ , we have

$$\begin{aligned}\text{Hom}_{\{t,f\}}(f \times f, t) &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong pt \\ &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{Hom}_{\{t,f\}}(f, \text{Hom}_{\{t,f\}}(f, t)).\end{aligned}$$

- For  $(A, B, C) = (f, f, f)$ , we have

$$\begin{aligned}\text{Hom}_{\{t,f\}}(f \times f, f) &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &= \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \text{Hom}_{\{t,f\}}(f, \text{Hom}_{\{t,f\}}(f, f)).\end{aligned}$$

Since  $\{t, f\}$  is posetal, naturality is automatic (?? of ??).



### 000E 3.2.3 0-Categories

000F

#### DEFINITION 3.2.3.1.1 ► 0-CATEGORIES

A **0-category** is a poset.<sup>1</sup>

---

<sup>1</sup>*Motivation:* A 0-category is precisely a category enriched in the poset of (-1)-categories.

000G

**DEFINITION 3.2.3.1.2 ► 0-GROUPOIDS**

A **0-groupoid** is a 0-category in which every morphism is invertible.<sup>1</sup>

<sup>1</sup>That is, a set.

000H

**3.2.4 Tables of Analogies Between Set Theory and Category Theory**

Here we record some analogies between notions in set theory and category theory. The analogies relating to presheaves relate equally well to copresheaves, as the opposite  $X^{\text{op}}$  of a set  $X$  is just  $X$  again.

01D6

**REMARK 3.2.4.1.1 ► BASIC ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY**

The basic analogies between set theory and category theory are summarised in the following table:

Set Theory	Category Theory
Enrichment in {true, false}	Enrichment in Sets
Set $X$	Category $C$
Element $x \in X$	Object $X \in \text{Obj}(C)$
Function $f: X \rightarrow Y$	Functor $F: C \rightarrow D$
Function $X \rightarrow \{\text{true, false}\}$	Copresheaf $C \rightarrow \text{Sets}$
Function $X \rightarrow \{\text{true, false}\}$	Presheaf $C^{\text{op}} \rightarrow \text{Sets}$

01D7

**REMARK 3.2.4.1.2 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: POWERSETS AND CATEGORIES OF PRESHEAVES**

The category of presheaves  $\text{PSh}(C)$  and the category of copresheaves  $\text{CoPSh}(C)$  on a category  $C$  are the 1-categorical counterparts to the powerset  $\mathcal{P}(X)$  of subsets of a set  $X$ . The further analogies built upon this are summarised in the following table:

Set Theory	Category Theory
Powerset $\mathcal{P}(X)$	Presheaf category $\text{PSh}(\mathcal{C})$
Characteristic function $\chi_{\{x\}} : X \rightarrow \{\text{t, f}\}$	Representable presheaf $h_X : \mathcal{C}^{\text{op}} \hookrightarrow \text{Sets}$
Characteristic embedding $\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$	Yoneda embedding $\underline{\chi} : \mathcal{C}^{\text{op}} \hookrightarrow \text{PSh}(\mathcal{C})$
Characteristic relation $\chi_X(-_1, -_2) : X \times X \rightarrow \{\text{t, f}\}$	Hom profunctor $\text{Hom}_{\mathcal{C}}(-_1, -_2) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \text{Sets}$
The Yoneda lemma for sets $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$	The Yoneda lemma for categories $\text{Nat}(h_X, \mathcal{F}) \cong \mathcal{F}(X)$
The characteristic embedding is fully faithful, $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$	The Yoneda embedding is fully faithful, $\text{Nat}(h_X, h_Y) \cong \text{Hom}_{\mathcal{C}}(X, Y)$
Subsets are unions of their elements $U = \bigcup_{x \in U} \{x\}$ or $\chi_U = \text{colim}_{\chi_x \in \mathcal{P}(U)} (\chi_x)$	Presheaves are colimits of representables, $\mathcal{F} \cong \text{colim}_{h_X \in \int_{\mathcal{C}} \mathcal{F}} (h_X)$

01D8

**REMARK 3.2.4.1.3 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: CATEGORIES OF ELEMENTS**

We summarise the analogies between un/straightening in set theory and category theory in the following table:

SET THEORY	CATEGORY THEORY
Assignment $U \mapsto \chi_U$	Assignment $\mathcal{F} \mapsto \int_{\mathcal{C}} \mathcal{F}$
Un/straightening isomorphism $\mathcal{P}(X) \cong \text{Sets}(X, \{\text{t, f}\})$	Un/straightening equivalence $\text{DFib}(\mathcal{C}) \stackrel{\text{eq.}}{\cong} \text{PSh}(\mathcal{C})$

01D9

**REMARK 3.2.4.1.4 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: FUNCTIONS BETWEEN POWERSETS AND FUNCTORS BETWEEN PRESHEAF CATEGORIES**

We summarise the analogies between functions  $\mathcal{P}(X) \rightarrow \mathcal{P}(Y)$  and functors  $\text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$  in the following table:

SET THEORY	CATEGORY THEORY
Direct image function $f_! : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Left Kan extension functor $F_! : \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$
Inverse image function $f^{-1} : \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$	Precomposition functor $F^* : \text{PSh}(\mathcal{D}) \rightarrow \text{PSh}(C)$
Codirect image function $f_* : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Right Kan extension functor $F_* : \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$

01DA

**REMARK 3.2.4.1.5 ► ANALOGIES BETWEEN SET THEORY AND CATEGORY THEORY: RELATIONS AND PROFUNCTORS**

We summarise the analogies between functions, relations and profunctors in the following table:

SET THEORY	CATEGORY THEORY
Relation $R : X \times Y \rightarrow \{\text{t}, \text{f}\}$	Profunctor $\mathfrak{p} : \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$
Relation $R : X \rightarrow \mathcal{P}(Y)$	Profunctor $\mathfrak{p} : C \rightarrow \text{PSh}(\mathcal{D})$
Relation as a cocontinuous morphism of posets $R : (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset)$	Profunctor as a colimit-preserving functor $\mathfrak{p} : \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$

# Appendices

## 3.A Other Chapters

### Preliminaries

- 1. Introduction
- 2. A Guide to the Literature

### Sets

- 3. Sets

### 4. Constructions With Sets

- 5. Monoidal Structures on the Category of Sets
- 6. Pointed Sets
- 7. Tensor Products of Pointed Sets

**Relations**

- 8. Relations
- 9. Constructions With Relations
- 10. Conditions on Relations

**Categories**

- 11. Categories
- 12. Presheaves and the Yoneda Lemma

**Monoidal Categories**

- 13. Constructions With Monoidal Categories

**Bicategories**

- 14. Types of Morphisms in Bicategories

**Extra Part**

- 15. Notes

## Chapter 4

# Constructions With Sets

**02H3** This chapter develops some material relating to constructions with sets with an eye towards its categorical and higher-categorical counterparts to be introduced later in this work. Of particular interest are perhaps the following:

- 01YT** 1. Explicit descriptions of the major types of co/limits in Sets, including in particular explicit descriptions of pushouts and coequalisers (see [Definitions 4.2.4.1.1](#) and [4.2.5.1.1](#) and [Remarks 4.2.4.1.4](#) and [4.2.5.1.4](#)).
- 01YU** 2. A discussion of powersets as decategorifications of categories of presheaves, including in particular results such as:
  - 01YY** (a) A discussion of the internal Hom of a powerset ([Section 4.4.7](#)).
  - 01YW** (b) A 0-categorical version of the Yoneda lemma ([Theorem 12.1.5.1.1](#)), which we term the *Yoneda lemma for sets* ([Proposition 4.5.5.1.1](#)).
  - 01YX** (c) A characterisation of powersets as free cocompletions ([Section 4.4.5](#)), mimicking the corresponding statement for categories of presheaves [\(??\)](#).
  - 01YY** (d) A characterisation of powersets as free completions ([Section 4.4.6](#)), mimicking the corresponding statement for categories of copresheaves [\(??\)](#).
  - 01YZ** (e) A  $(-1)$ -categorical version of un/straightening ([Item 2 of Proposition 4.5.1.1.4](#) and [Remark 4.5.1.1.6](#)).
  - 01Z0** (f) A 0-categorical form of Isbell duality internal to powersets ([Section 4.4.8](#)).
- 01Z1** 3. A lengthy discussion of the adjoint triple

$$f_! \dashv f^{-1} \dashv f_*: \mathcal{P}(A) \xrightarrow{\cong} \mathcal{P}(B)$$

of functors (i.e. morphisms of posets) between  $\mathcal{P}(A)$  and  $\mathcal{P}(B)$  induced by a map of sets  $f: A \rightarrow B$ , including in particular:

- 
- 01Z2 (a) How  $f^{-1}$  can be described as a precomposition while  $f_!$  and  $f_*$  can be described as Kan extensions ([Remarks 4.6.1.1.4, 4.6.2.1.2 and 4.6.3.1.4](#)).
- 01Z3 (b) An extensive list of the properties of  $f_!$ ,  $f^{-1}$ , and  $f_*$  ([Propositions 4.6.1.1.5, 4.6.1.1.7, 4.6.2.1.3, 4.6.2.1.5, 4.6.3.1.7 and 4.6.3.1.9](#)).
- 01Z4 (c) How the functors  $f_!$ ,  $f^{-1}$ ,  $f_*$ , along with the functors

$$\begin{aligned} -_1 \cap -_2 : \mathcal{P}(X) \times \mathcal{P}(X) &\rightarrow \mathcal{P}(X), \\ [-_1, -_2]_X : \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) &\rightarrow \mathcal{P}(X) \end{aligned}$$

may be viewed as a six-functor formalism with the empty set  $\emptyset$  as the dualising object ([Section 4.6.4](#)).

## Contents

---

<b>4.1 Limits of Sets.....</b>	<b>40</b>
4.1.1 The Terminal Set.....	40
4.1.2 Products of Families of Sets.....	41
4.1.3 Binary Products of Sets.....	45
4.1.4 Pullbacks.....	55
4.1.5 Equalisers.....	67
4.1.6 Inverse Limits.....	71
<b>4.2 Colimits of Sets.....</b>	<b>74</b>
4.2.1 The Initial Set.....	74
4.2.2 Coproducts of Families of Sets.....	75
4.2.3 Binary Coproducts.....	77
4.2.4 Pushouts.....	81
4.2.5 Coequalisers.....	90
4.2.6 Direct Colimits.....	94
<b>4.3 Operations With Sets.....</b>	<b>100</b>
4.3.1 The Empty Set.....	100
4.3.2 Singleton Sets.....	100
4.3.3 Pairings of Sets .....	100
4.3.4 Ordered Pairs .....	100
4.3.5 Sets of Maps.....	101
4.3.6 Unions of Families of Subsets .....	104
4.3.7 Intersections of Families of Subsets.....	119
4.3.8 Binary Unions.....	133
4.3.9 Binary Intersections .....	139
4.3.10 Differences.....	145
4.3.11 Complements .....	150

4.3.12	Symmetric Differences.....	153
<b>4.4</b>	<b>Powersets .....</b>	<b>161</b>
4.4.1	Foundations .....	161
4.4.2	Functionality of Powersets .....	164
4.4.3	Adjointness of Powersets I.....	166
4.4.4	Adjointness of Powersets II.....	167
4.4.5	Powersets as Free Cocompletions .....	167
4.4.6	Powersets as Free Completions .....	174
4.4.7	The Internal Hom of a Powerset.....	180
4.4.8	Isbell Duality for Sets .....	197
<b>4.5</b>	<b>Characteristic Functions.....</b>	<b>199</b>
4.5.1	The Characteristic Function of a Subset.....	199
4.5.2	The Characteristic Function of a Point.....	204
4.5.3	The Characteristic Relation of a Set.....	205
4.5.4	The Characteristic Embedding of a Set.....	206
4.5.5	The Yoneda Lemma for Sets.....	207
<b>4.6</b>	<b>The Adjoint Triple <math>f_! \dashv f^{-1} \dashv f_*</math>.....</b>	<b>209</b>
4.6.1	Direct Images .....	209
4.6.2	Inverse Images.....	220
4.6.3	Codirect Images .....	230
4.6.4	A Six-Functor Formalism for Sets.....	243
<b>4.7</b>	<b>Miscellany .....</b>	<b>251</b>
4.7.1	Injective Functions .....	251
4.7.2	Surjective Functions.....	257
<b>4.A</b>	<b>Other Chapters .....</b>	<b>263</b>

## 000K 4.1 Limits of Sets

### 000L 4.1.1 The Terminal Set

000M

#### DEFINITION 4.1.1.1 ► THE TERMINAL SET

The **terminal set** is the terminal object of Sets as in ??.

01DB

#### CONSTRUCTION 4.1.1.2 ► CONSTRUCTION OF THE TERMINAL SET

Concretely, the terminal set is the pair  $(\text{pt}, \{\text{!}_A\}_{A \in \text{Obj}(\text{Sets})})$  consisting of:

01DC

1. *The Limit.* The punctual set  $\text{pt} \stackrel{\text{def}}{=} \{\star\}$ .

01DD

2. *The Cone.* The collection of maps

$$\{!_A : A \rightarrow \text{pt}\}_{A \in \text{Obj}(\text{Sets})}$$

defined by

$$!_A(a) \stackrel{\text{def}}{=} \star$$

for each  $a \in A$  and each  $A \in \text{Obj}(\text{Sets})$ .

#### PROOF 4.1.1.3 ► PROOF OF CONSTRUCTION 4.1.1.2

We claim that pt is the terminal object of Sets. Indeed, suppose we have a diagram of the form

$$A \quad \text{pt}$$

in Sets. Then there exists a unique map  $\phi : A \rightarrow \text{pt}$  making the diagram

$$A \xrightarrow[\exists!]{} \text{pt}$$

commute, namely  $!_A$ . 

## 000N 4.1.2 Products of Families of Sets

Let  $\{A_i\}_{i \in I}$  be a family of sets.

000P

#### DEFINITION 4.1.2.1.1 ► THE PRODUCT OF A FAMILY OF SETS

The **product**<sup>1</sup> of  $\{A_i\}_{i \in I}$  is the product of  $\{A_i\}_{i \in I}$  in Sets as in ??.

<sup>1</sup>Further Terminology: Also called the **Cartesian product** of  $\{A_i\}_{i \in I}$ .

01DE

#### CONSTRUCTION 4.1.2.1.2 ► CONSTRUCTION OF THE PRODUCT OF A FAMILY OF SETS

Concretely, the product of  $\{A_i\}_{i \in I}$  is the pair  $(\prod_{i \in I} A_i, \{\text{pr}_i\}_{i \in I})$  consisting of:

01DF

1. *The Limit.* The set  $\prod_{i \in I} A_i$  defined by

$$\prod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}\left(I, \bigcup_{i \in I} A_i\right) \middle| \begin{array}{l} \text{for each } i \in I, \text{ we} \\ \text{have } f(i) \in A_i \end{array} \right\}.$$

01DG

2. *The Cone.* The collection

$$\left\{ \text{pr}_i : \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

of maps given by

$$\text{pr}_i(f) \stackrel{\text{def}}{=} f(i)$$

for each  $f \in \prod_{i \in I} A_i$  and each  $i \in I$ .

**PROOF 4.1.2.1.3 ► PROOF OF CONSTRUCTION 4.1.2.1.2**

We claim that  $\prod_{i \in I} A_i$  is the categorical product of  $\{A_i\}_{i \in I}$  in Sets. Indeed, suppose we have, for each  $i \in I$ , a diagram of the form

$$\begin{array}{ccc} P & & \\ & \searrow p_i & \\ & \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} A_i \end{array}$$

in Sets. Then there exists a unique map  $\phi: P \rightarrow \prod_{i \in I} A_i$  making the diagram

$$\begin{array}{ccc} P & & \\ \downarrow \phi & \swarrow \exists! & \searrow p_i \\ \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} & A_i \end{array}$$

commute, being uniquely determined by the condition  $\text{pr}_i \circ \phi = p_i$  for each  $i \in I$  via

$$\phi(x) = (p_i(x))_{i \in I}$$

for each  $x \in P$ . □

01DH

**REMARK 4.1.2.1.4 ► UNWINDING CONSTRUCTION 4.1.2.1.2**

Less formally, we may think of Cartesian products and projection maps as follows:

01DJ

1. We think of  $\prod_{i \in I} A_i$  as the set whose elements are  $I$ -indexed collections  $(a_i)_{i \in I}$  with  $a_i \in A_i$  for each  $i \in I$ .

01DK

2. We view the projection maps

$$\left\{ \text{pr}_i : \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

as being given by

$$\text{pr}_i((a_j)_{j \in I}) \stackrel{\text{def}}{=} a_i$$

for each  $(a_j)_{j \in I} \in \prod_{i \in I} A_i$  and each  $i \in I$ .

000Q

#### PROPOSITION 4.1.2.1.5 ► PROPERTIES OF PRODUCTS OF FAMILIES OF SETS

Let  $\{A_i\}_{i \in I}$  be a family of sets.

000R

1. *Functoriality.* The assignment  $\{A_i\}_{i \in I} \mapsto \prod_{i \in I} A_i$  defines a functor

$$\prod_{i \in I} : \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each  $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$ , we have

$$\left[ \prod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} A_i$$

- *Action on Morphisms.* For each  $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$ , the action on Hom-sets

$$\left( \prod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left( \prod_{i \in I} A_i, \prod_{i \in I} B_i \right)$$

of  $\prod_{i \in I}$  at  $((A_i)_{i \in I}, (B_i)_{i \in I})$  is defined by sending a map

$$\{f_i : A_i \rightarrow B_i\}_{i \in I}$$

in  $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$  to the map of sets

$$\prod_{i \in I} f_i : \prod_{i \in I} A_i \rightarrow \prod_{i \in I} B_i$$

defined by

$$\left[ \prod_{i \in I} f_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} (f_i(a_i))_{i \in I}$$

for each  $(a_i)_{i \in I} \in \prod_{i \in I} A_i$ .



Item 1: Functoriality

This follows from ?? of ??.



### 000S 4.1.3 Binary Products of Sets

Let  $A$  and  $B$  be sets.

#### 000T DEFINITION 4.1.3.1.1 ► BINARY PRODUCTS OF SETS

The **product of  $A$  and  $B$** <sup>1</sup> is the product of  $A$  and  $B$  in Sets as in ??.

<sup>1</sup>Further Terminology: Also called the **Cartesian product of  $A$  and  $B$** .

#### 01DL CONSTRUCTION 4.1.3.1.2 ► CONSTRUCTION OF BINARY PRODUCTS OF SETS

Concretely, the product of  $A$  and  $B$  is the pair  $(A \times B, \{\text{pr}_1, \text{pr}_2\})$  consisting of:

#### 01DM

1. *The Limit.* The set  $A \times B$  defined by

$$\begin{aligned} A \times B &\stackrel{\text{def}}{=} \prod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{f \in \text{Sets}(\{0, 1\}, A \cup B) \mid \text{we have } f(0) \in A \text{ and } f(1) \in B\} \\ &\cong \{\{\{a\}, \{a, b\}\} \in \mathcal{P}(\mathcal{P}(A \cup B)) \mid \text{we have } a \in A \text{ and } b \in B\} \\ &\cong \left\{ \begin{array}{l} \text{ordered pairs } (a, b) \text{ with} \\ a \in A \text{ and } b \in B \end{array} \right\}. \end{aligned}$$

#### 01DN

2. *The Cone.* The maps

$$\begin{aligned} \text{pr}_1 &: A \times B \rightarrow A, \\ \text{pr}_2 &: A \times B \rightarrow B \end{aligned}$$

defined by

$$\begin{aligned} \text{pr}_1(a, b) &\stackrel{\text{def}}{=} a, \\ \text{pr}_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each  $(a, b) \in A \times B$ .

**PROOF 4.1.3.1.3 ► PROOF OF CONSTRUCTION 4.1.3.1.2**

We claim that  $A \times B$  is the categorical product of  $A$  and  $B$  in the category of sets. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & P & & \\ & p_1 \swarrow & & \searrow p_2 & \\ A & \xleftarrow{\text{pr}_1} & A \times B & \xrightarrow{\text{pr}_2} & B \end{array}$$

in Sets. Then there exists a unique map  $\phi: P \rightarrow A \times B$  making the diagram

$$\begin{array}{ccccc} & & P & & \\ & p_1 \swarrow & \downarrow \phi \exists! & \searrow p_2 & \\ A & \xleftarrow{\text{pr}_1} & A \times B & \xrightarrow{\text{pr}_2} & B \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned} \text{pr}_1 \circ \phi &= p_1, \\ \text{pr}_2 \circ \phi &= p_2 \end{aligned}$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each  $x \in P$ .



000U

**PROPOSITION 4.1.3.1.4 ► PROPERTIES OF PRODUCTS OF SETS**

Let  $A, B, C$ , and  $X$  be sets.

000V

- Functoriality.* The assignments  $A, B, (A, B) \mapsto A \times B$  define functors

$$\begin{aligned} A \times -: \quad \text{Sets} &\rightarrow \text{Sets}, \\ - \times B: \quad \text{Sets} &\rightarrow \text{Sets}, \\ -_1 \times -_2: \quad \text{Sets} \times \text{Sets} &\rightarrow \text{Sets}, \end{aligned}$$

where  $-_1 \times -_2$  is the functor where

- Action on Objects.* For each  $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$ , we have

$$[-_1 \times -_2](A, B) \stackrel{\text{def}}{=} A \times B.$$

- Action on Morphisms.* For each  $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$ , the action on Hom-sets

$$\times_{(A,B),(X,Y)}: \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \times B, X \times Y)$$

of  $\times$  at  $((A, B), (X, Y))$  is defined by sending  $(f, g)$  to the function

$$f \times g: A \times B \rightarrow X \times Y$$

defined by

$$[f \times g](a, b) \stackrel{\text{def}}{=} (f(a), g(b))$$

for each  $(a, b) \in A \times B$ .

and where  $A \times -$  and  $- \times B$  are the partial functors of  $-_1 \times -_2$  at  $A, B \in \text{Obj}(\text{Sets})$ .

**000W** 2. *Adjointness I.* We have adjunctions

$$\begin{aligned} (A \times - \dashv \text{Sets}(A, -)): \quad & \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Sets}(A, -)} \end{array} \text{Sets}, \\ (- \times B \dashv \text{Sets}(B, -)): \quad & \text{Sets} \begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Sets}(B, -)} \end{array} \text{Sets}, \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \text{Sets}(A \times B, C) &\cong \text{Sets}(A, \text{Sets}(B, C)), \\ \text{Sets}(A \times B, C) &\cong \text{Sets}(B, \text{Sets}(A, C)), \end{aligned}$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

**01Z5** 3. *Adjointness II.* We have an adjunction

$$(\Delta_{\text{Sets}} \dashv -_1 \times -_2): \quad \text{Sets} \begin{array}{c} \xrightarrow{\Delta_{\text{Sets}}} \\ \perp \\ \xleftarrow{-_1 \times -_2} \end{array} \text{Sets} \times \text{Sets},$$

witnessed by a bijection

$$\text{Hom}_{\text{Sets} \times \text{Sets}}((A, A), (B, C)) \cong \text{Sets}(A, B \times C),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and in  $(B, C) \in \text{Obj}(\text{Sets} \times \text{Sets})$ .

**000X** 4. *Associativity.* We have an isomorphism of sets

$$\alpha_{A,B,C}^{\text{Sets}}: (A \times B) \times C \xrightarrow{\sim} A \times (B \times C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000Y

5. *Unitality.* We have isomorphisms of sets

$$\begin{aligned}\lambda_A^{\text{Sets}} &: \text{pt} \times A \xrightarrow{\sim} A, \\ \rho_A^{\text{Sets}} &: A \times \text{pt} \xrightarrow{\sim} A,\end{aligned}$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

000Z

6. *Commutativity.* We have an isomorphism of sets

$$\sigma_{A,B}^{\text{Sets}} : A \times B \xrightarrow{\sim} B \times A,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$ .

01DP

7. *Distributivity Over Coproducts.* We have isomorphisms of sets

$$\begin{aligned}\delta_\ell^{\text{Sets}} &: A \times (B \coprod C) \xrightarrow{\sim} (A \times B) \coprod (A \times C), \\ \delta_r^{\text{Sets}} &: (A \coprod B) \times C \xrightarrow{\sim} (A \times C) \coprod (B \times C),\end{aligned}$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

0010

8. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{aligned}\zeta_\ell^{\text{Sets}} &: \emptyset \times A \xrightarrow{\sim} \emptyset, \\ \zeta_r^{\text{Sets}} &: A \times \emptyset \xrightarrow{\sim} \emptyset,\end{aligned}$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

0011

9. *Distributivity Over Unions.* Let  $X$  be a set. For each  $U, V, W \in \mathcal{P}(X)$ , we have equalities

$$\begin{aligned}U \times (V \cup W) &= (U \times V) \cup (U \times W), \\ (U \cup V) \times W &= (U \times W) \cup (V \times W)\end{aligned}$$

of subsets of  $\mathcal{P}(X \times X)$ .

0012

10. *Distributivity Over Intersections.* Let  $X$  be a set. For each  $U, V, W \in \mathcal{P}(X)$ , we have equalities

$$\begin{aligned}U \times (V \cap W) &= (U \times V) \cap (U \times W), \\ (U \cap V) \times W &= (U \times W) \cap (V \times W)\end{aligned}$$

of subsets of  $\mathcal{P}(X \times X)$ .

0014

11. *Distributivity Over Differences.* Let  $X$  be a set. For each  $U, V, W \in \mathcal{P}(X)$ , we have equalities

$$\begin{aligned} U \times (V \setminus W) &= (U \times V) \setminus (U \times W), \\ (U \setminus V) \times W &= (U \times W) \setminus (V \times W) \end{aligned}$$

of subsets of  $\mathcal{P}(X \times X)$ .

0015

12. *Distributivity Over Symmetric Differences.* Let  $X$  be a set. For each  $U, V, W \in \mathcal{P}(X)$ , we have equalities

$$\begin{aligned} U \times (V \Delta W) &= (U \times V) \Delta (U \times W), \\ (U \Delta V) \times W &= (U \times W) \Delta (V \times W) \end{aligned}$$

of subsets of  $\mathcal{P}(X \times X)$ .

0013

13. *Middle-Four Exchange with Respect to Intersections.* The diagram

$$\begin{array}{ccc} (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \xrightarrow{\cap \times \cap} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \downarrow \mathcal{P}_{X,X}^{\times} \times \mathcal{P}_{X,X}^{\times} & & \downarrow \mathcal{P}_{X,X}^{\times} \\ \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) & \xrightarrow{\cap} & \mathcal{P}(X \times X) \end{array}$$

commutes, i.e. we have

$$(U \times V) \cap (W \times T) = (U \cap V) \times (W \cap T).$$

for each  $U, V, W, T \in \mathcal{P}(X)$ .

0016

14. *Symmetric Monoidality.* The 8-tuple  $(\text{Sets}, \times, \text{pt}, \text{Sets}(-_1, -_2), \alpha^{\text{Sets}}, \lambda^{\text{Sets}}, \rho^{\text{Sets}}, \sigma^{\text{Sets}})$  is a closed symmetric monoidal category.

0017

15. *Symmetric Bimonoidality.* The 18-tuple

$$\left( \text{Sets}, \coprod, \times, \emptyset, \text{pt}, \text{Sets}(-_1, -_2), \alpha^{\text{Sets}}, \lambda^{\text{Sets}}, \rho^{\text{Sets}}, \sigma^{\text{Sets}}, \alpha^{\text{Sets}, \coprod}, \lambda^{\text{Sets}, \coprod}, \rho^{\text{Sets}, \coprod}, \sigma^{\text{Sets}, \coprod}, \delta_r^{\text{Sets}}, \delta_r^{\text{Sets}}, \zeta_\ell^{\text{Sets}}, \zeta_r^{\text{Sets}} \right),$$

is a symmetric closed bimonoidal category, where  $\alpha^{\text{Sets}, \coprod}, \lambda^{\text{Sets}, \coprod}, \rho^{\text{Sets}, \coprod}$ , and  $\sigma^{\text{Sets}, \coprod}$  are the natural transformations from [Items 3 to 5 of Proposition 4.2.3.1.4](#).

**PROOF 4.1.3.1.5 ► PROOF OF PROPOSITION 4.1.3.1.4****Item 1: Functoriality**

This follows from ?? of ??.

**Item 2: Adjointness**

We prove only that there's an adjunction  $- \times B \dashv \text{Sets}(B, -)$ , witnessed by a bijection

$$\text{Sets}(A \times B, C) \cong \text{Sets}(A, \text{Sets}(B, C)),$$

natural in  $B, C \in \text{Obj}(\text{Sets})$ , as the proof of the existence of the adjunction  $A \times - \dashv \text{Sets}(A, -)$  follows almost exactly in the same way.

01Z6

- *Map I.* We define a map

$$\Phi_{B,C}: \text{Sets}(A \times B, C) \rightarrow \text{Sets}(A, \text{Sets}(B, C)),$$

by sending a function

$$\xi: A \times B \rightarrow C$$

to the function

$$\xi^\dagger: A \longrightarrow \text{Sets}(B, C),$$

$$a \mapsto (\xi_a^\dagger: B \rightarrow C),$$

where we define

$$\xi_a^\dagger(b) \stackrel{\text{def}}{=} \xi(a, b)$$

for each  $b \in B$ . In terms of the  $\llbracket a \mapsto f(a) \rrbracket$  notation of [Notation 3.1.1.2](#), we have

$$\xi^\dagger \stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket.$$

01Z7

- *Map II.* We define a map

$$\Psi_{B,C}: \text{Sets}(A, \text{Sets}(B, C)) \rightarrow \text{Sets}(A \times B, C)$$

given by sending a function

$$\xi: A \longrightarrow \text{Sets}(B, C),$$

$$a \mapsto (\xi_a: B \rightarrow C),$$

to the function

$$\xi^\dagger : A \times B \rightarrow C$$

defined by

$$\begin{aligned}\xi^\dagger(a, b) &\stackrel{\text{def}}{=} \text{ev}_b(\text{ev}_a(\xi)) \\ &\stackrel{\text{def}}{=} \text{ev}_b(\xi_a) \\ &\stackrel{\text{def}}{=} \xi_a(b)\end{aligned}$$

for each  $(a, b) \in A \times B$ .

01Z8

- *Invertibility I.* We claim that

$$\Psi_{A,B} \circ \Phi_{A,B} = \text{id}_{\text{Sets}(A \times B, C)}.$$

Indeed, given a function  $\xi : A \times B \rightarrow C$ , we have

$$\begin{aligned}[\Psi_{A,B} \circ \Phi_{A,B}](\xi) &= \Psi_{A,B}(\Phi_{A,B}(\xi)) \\ &= \Psi_{A,B}(\Phi_{A,B}(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\ &= \Psi_{A,B}(\llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket) \\ &= \Psi_{A,B}(\llbracket a' \mapsto \llbracket b' \mapsto \xi(a', b') \rrbracket \rrbracket) \\ &= \llbracket (a, b) \mapsto \text{ev}_b(\text{ev}_a(\llbracket a' \mapsto \llbracket b' \mapsto \xi(a', b') \rrbracket) \rrbracket) \rrbracket \\ &= \llbracket (a, b) \mapsto \text{ev}_b(\llbracket b' \mapsto \xi(a, b') \rrbracket) \rrbracket \\ &= \llbracket (a, b) \mapsto \xi(a, b) \rrbracket \\ &= \xi.\end{aligned}$$

01Z9

- *Invertibility II.* We claim that

$$\Phi_{A,B} \circ \Psi_{A,B} = \text{id}_{\text{Sets}(A, \text{Sets}(B, C))}.$$

Indeed, given a function

$$\xi : A \longrightarrow \text{Sets}(B, C),$$

$$a \mapsto (\xi_a : B \rightarrow C),$$

we have

$$\begin{aligned}[\Phi_{A,B} \circ \Psi_{A,B}](\xi) &\stackrel{\text{def}}{=} \Phi_{A,B}(\Psi_{A,B}(\xi)) \\ &\stackrel{\text{def}}{=} \Phi_{A,B}(\llbracket (a, b) \mapsto \xi_a(b) \rrbracket) \\ &\stackrel{\text{def}}{=} \Phi_{A,B}(\llbracket (a', b') \mapsto \xi_{a'}(b') \rrbracket) \\ &\stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \text{ev}_{(a,b)}(\llbracket (a', b') \mapsto \xi_{a'}(b') \rrbracket) \rrbracket \rrbracket \\ &\stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \xi_a(b) \rrbracket \rrbracket \\ &\stackrel{\text{def}}{=} \llbracket a \mapsto \xi_a \rrbracket \\ &\stackrel{\text{def}}{=} \xi.\end{aligned}$$

01ZA

- *Naturality for  $\Phi$ , Part I.* We need to show that, given a function  $g: B \rightarrow B'$ , the diagram

$$\begin{array}{ccc} \text{Sets}(A \times B', C) & \xrightarrow{\Phi_{B',C}} & \text{Sets}(A, \text{Sets}(B', C)), \\ \text{id}_A \times g^* \downarrow & & \downarrow (g^*)_! \\ \text{Sets}(A \times B, C) & \xrightarrow{\Phi_{B,C}} & \text{Sets}(A, \text{Sets}(B, C)) \end{array}$$

commutes. Indeed, given a function

$$\xi: A \times B' \rightarrow C,$$

we have

$$\begin{aligned} [\Phi_{B,C} \circ (\text{id}_A \times g^*)](\xi) &= \Phi_{B,C}([\text{id}_A \times g^*](\xi)) \\ &= \Phi_{B,C}(\xi(-_1, g(-_2))) \\ &= [\xi(-_1, g(-_2))]^\dagger \\ &= \xi_{-1}^\dagger(g(-_2)) \\ &= (g^*)_!(\xi^\dagger) \\ &= (g^*)_!(\Phi_{B',C}(\xi)) \\ &= [(g^*)_! \circ \Phi_{B',C}](\xi). \end{aligned}$$

Alternatively, using the  $\llbracket a \mapsto f(a) \rrbracket$  notation of [Notation 3.1.1.1.2](#), we have

$$\begin{aligned} [\Phi_{B,C} \circ (\text{id}_A \times g^*)](\xi) &= \Phi_{B,C}([\text{id}_A \times g^*](\xi)) \\ &= \Phi_{B,C}([\text{id}_A \times g^*](\llbracket (a, b') \mapsto \xi(a, b') \rrbracket)) \\ &= \Phi_{B,C}(\llbracket (a, b) \mapsto \xi(a, g(b)) \rrbracket) \\ &= \llbracket a \mapsto \llbracket b \mapsto \xi(a, g(b)) \rrbracket \rrbracket \\ &= \llbracket a \mapsto g^*(\llbracket b' \mapsto \xi(a, b') \rrbracket) \rrbracket \\ &= (g^*)_!(\llbracket a \mapsto \llbracket b' \mapsto \xi(a, b') \rrbracket \rrbracket) \\ &= (g^*)_!(\Phi_{B',C}(\llbracket (a, b') \mapsto \xi(a, b') \rrbracket)) \\ &= (g^*)_!(\Phi_{B',C}(\xi)) \\ &= [(g^*)_! \circ \Phi_{B',C}](\xi). \end{aligned}$$

01ZB

- *Naturality for  $\Phi$ , Part II.* We need to show that, given a function  $h: C \rightarrow C'$ , the diagram

$$\begin{array}{ccc} \text{Sets}(A \times B, C) & \xrightarrow{\Phi_{B,C}} & \text{Sets}(A, \text{Sets}(B, C)), \\ h_! \downarrow & & \downarrow (h_!)_! \\ \text{Sets}(A \times B, C') & \xrightarrow{\Phi_{B,C'}} & \text{Sets}(A, \text{Sets}(B, C')) \end{array}$$

commutes. Indeed, given a function

$$\xi: A \times B \rightarrow C,$$

we have

$$\begin{aligned} [\Phi_{B,C} \circ h_!](\xi) &= \Phi_{B,C}(h_!(\xi)) \\ &= \Phi_{B,C}(h_!(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\ &= \Phi_{B,C}(\llbracket (a, b) \mapsto h(\xi(a, b)) \rrbracket) \\ &= \llbracket a \mapsto \llbracket b \mapsto h(\xi(a, b)) \rrbracket \rrbracket \\ &= \llbracket a \mapsto h_!(\llbracket b \mapsto \xi(a, b) \rrbracket) \rrbracket \\ &= (h_!)_!(\llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket) \\ &= (h_!)_!(\Phi_{B,C}(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\ &= (h_!)_!(\Phi_{B,C}(\xi)) \\ &= [(h_!)_! \circ \Phi_{B,C}](\xi). \end{aligned}$$

01ZC

- *Naturality for  $\Psi$ .* Since  $\Phi$  is natural in each argument and  $\Phi$  is a componentwise inverse to  $\Psi$  in each argument, it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\Psi$  is also natural in each argument.

This finishes the proof.

**Item 3: Adjointness II**

This follows from the universal property of the product.

**Item 4: Associativity**

This is proved in the proof of [Definition 5.1.4.1.1](#).

**Item 5: Unitality**

This is proved in the proof of [Definitions 5.1.5.1.1](#) and [5.1.6.1.1](#).

**Item 6: Commutativity**

This is proved in the proof of [Definition 5.1.7.1.1](#).

**Item 7: Distributivity Over Coproducts**

This is proved in the proof of [Definitions 5.3.1.1.1](#) and [5.3.2.1.1](#).

**Item 8: Annihilation With the Empty Set**

This is proved in the proof of [Definitions 5.3.3.1.1](#) and [5.3.4.1.1](#).

**Item 9: Distributivity Over Unions**

See [[Pro25c](#)].

**Item 10: Distributivity Over Intersections**

See [[Pro25d](#), Corollary 1].

**Item 11: Distributivity Over Differences**

See [[Pro25a](#)].

**Item 12: Distributivity Over Symmetric Differences**

See [[Pro25b](#)].

**Item 13: Middle-Four Exchange With Respect to Intersections**

See [[Pro25d](#), Corollary 1].

**Item 14: Symmetric Monoidality**

This is a repetition of [Proposition 5.1.9.1.1](#), and is proved there.

**Item 15: Symmetric Bimonoidality**

This is a repetition of [Proposition 5.3.5.1.1](#), and is proved there. 

01DR

#### REMARK 4.1.3.1.6 ▶ THE CARTESIAN PRODUCT OF SETS AS AN $(\mathbb{E}_k, \mathbb{E}_\ell)$ -TENSOR PRODUCT

As shown in [Item 1](#) of [Proposition 4.1.3.1.4](#), the Cartesian product of sets defines a functor

$$-_1 \times -_2 : \text{Sets} \times \text{Sets} \rightarrow \text{Sets}.$$

This functor is the  $(k, \ell) = (-1, -1)$  case of a family of functors


$$\otimes_{k,\ell} : \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_\ell}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_{k+\ell}}(\text{Sets})$$

of tensor products of  $\mathbb{E}_k$ -monoid objects on Sets with  $\mathbb{E}_\ell$ -monoid objects on Sets; see ??.

01DS

#### REMARK 4.1.3.1.7 ▶ DIAGRAMS FOR ITEMS 9 TO 12 OF PROPOSITION 4.1.3.1.4

We may state the equalities in Items 9 to 12 of Proposition 4.1.3.1.4 as the commutativity of the following diagrams:



#### 0018 4.1.4 Pullbacks

Let  $A$ ,  $B$ , and  $C$  be sets and let  $f: A \rightarrow C$  and  $g: B \rightarrow C$  be functions.

0019

## DEFINITION 4.1.4.1.1 ► PULLBACKS OF SETS

The **pullback of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** <sup>1</sup> is the pullback of  $A$  and  $B$  over  $C$  along  $f$  and  $g$  in Sets as in ??.

<sup>1</sup>*Further Terminology:* Also called the **fibre product of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** .

01DT

## CONSTRUCTION 4.1.4.1.2 ► CONSTRUCTION OF PULLBACKS OF SETS

Concretely, the pullback of  $A$  and  $B$  over  $C$  along  $f$  and  $g$  is the pair  $(A \times_C B, \{\text{pr}_1, \text{pr}_2\})$  consisting of:

01DU

1. *The Limit.* The set  $A \times_C B$  defined by

$$A \times_C B \stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\}.$$

01DV

2. *The Cone.* The maps<sup>1</sup>

$$\begin{aligned} \text{pr}_1 &: A \times_C B \rightarrow A, \\ \text{pr}_2 &: A \times_C B \rightarrow B \end{aligned}$$

defined by

$$\begin{aligned} \text{pr}_1(a, b) &\stackrel{\text{def}}{=} a, \\ \text{pr}_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each  $(a, b) \in A \times_C B$ .

<sup>1</sup>*Further Notation:* Also written  $\text{pr}_1^{A \times_C B}$  and  $\text{pr}_2^{A \times_C B}$ .

## PROOF 4.1.4.1.3 ► PROOF OF CONSTRUCTION 4.1.4.1.2

We claim that  $A \times_C B$  is the categorical pullback of  $A$  and  $B$  over  $C$  with respect to  $(f, g)$  in Sets. First we need to check that the relevant pullback diagram commutes, i.e. that we have

$$\begin{array}{ccc} A \times_C B & \xrightarrow{\text{pr}_2} & B \\ f \circ \text{pr}_1 = g \circ \text{pr}_2, & \text{pr}_1 \downarrow & \downarrow g \\ A & \xrightarrow{f} & C. \end{array}$$

Indeed, given  $(a, b) \in A \times_C B$ , we have

$$[f \circ \text{pr}_1](a, b) = f(\text{pr}_1(a, b))$$

$$\begin{aligned}
 &= f(a) \\
 &= g(b) \\
 &= g(\text{pr}_2(a, b)) \\
 &= [g \circ \text{pr}_2](a, b),
 \end{aligned}$$

where  $f(a) = g(b)$  since  $(a, b) \in A \times_C B$ . Next, we prove that  $A \times_C B$  satisfies the universal property of the pullback. Suppose we have a diagram of the form

$$\begin{array}{ccccc}
 P & \xrightarrow{\quad p_2 \quad} & A \times_C B & \xrightarrow{\quad \text{pr}_2 \Rightarrow B \quad} & B \\
 p_1 \swarrow & & \downarrow \text{pr}_1 & & \downarrow g \\
 & & A & \xrightarrow{\quad f \quad} & C
 \end{array}$$

in Sets. Then there exists a unique map  $\phi: P \rightarrow A \times_C B$  making the diagram

$$\begin{array}{ccccc}
 P & \xrightarrow{\quad p_2 \quad} & A \times_C B & \xrightarrow{\quad \text{pr}_2 \Rightarrow B \quad} & B \\
 \exists! \phi \dashv \nearrow & & \downarrow \text{pr}_1 & & \downarrow g \\
 p_1 \swarrow & & A & \xrightarrow{\quad f \quad} & C
 \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned}
 \text{pr}_1 \circ \phi &= p_1, \\
 \text{pr}_2 \circ \phi &= p_2
 \end{aligned}$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each  $x \in P$ , where we note that  $(p_1(x), p_2(x)) \in A \times B$  indeed lies in  $A \times_C B$  by the condition

$$f \circ p_1 = g \circ p_2,$$

which gives

$$f(p_1(x)) = g(p_2(x))$$

for each  $x \in P$ , so that  $(p_1(x), p_2(x)) \in A \times_C B$ . □

01DW

#### REMARK 4.1.4.1.4 ► PULLBACKS OF SETS DEPEND ON THE MAPS

It is common practice to write  $A \times_C B$  for the pullback of  $A$  and  $B$  over  $C$  along  $f$  and  $g$ , omitting the maps  $f$  and  $g$  from the notation and instead leaving them implicit, to be understood from the context. However, the set  $A \times_C B$  depends very much on the maps  $f$  and  $g$ , and sometimes it is necessary or useful to note this dependence explicitly. In such situations, we will write  $A \times_{f,C,g} B$  or  $A \times_C^{f,g} B$  for  $A \times_C B$ .

001A

#### EXAMPLE 4.1.4.1.5 ► EXAMPLES OF PULLBACKS OF SETS

Here are some examples of pullbacks of sets.

001B

1. *Unions via Intersections.* Let  $X$  be a set. We have

$$\begin{array}{ccc} A \cap B & \longrightarrow & B \\ \downarrow \lrcorner & & \downarrow \wr_B \\ A & \xhookrightarrow{\iota_A} & A \cup B \end{array}$$

$A \cap B \cong A \times_{A \cup B} B,$

for each  $A, B \in \mathcal{P}(X)$ .

#### PROOF 4.1.4.1.6 ► PROOF OF EXAMPLE 4.1.4.1.5

##### Item 1: Unions via Intersections

Indeed, we have

$$\begin{aligned} A \times_{A \cup B} B &\cong \{(x, y) \in A \times B \mid x = y\} \\ &\cong A \cap B. \end{aligned}$$

This finishes the proof. □

001C

#### PROPOSITION 4.1.4.1.7 ► PROPERTIES OF PULLBACKS OF SETS


Let  $A, B, C$ , and  $X$  be sets.

001D

1. *Functionality.* The assignment  $(A, B, C, f, g) \mapsto A \times_{f,C,g} B$  defines a functor

$$-_1 \times_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where  $\mathcal{P}$  is the category that looks like this:



In particular, the action on morphisms of  $-_1 \times_{-3} -_1$  is given by sending a morphism

$$\begin{array}{ccccc}
 A \times_C B & \xrightarrow{\quad} & B & & \\
 \downarrow & \lrcorner & \downarrow g & \searrow \psi & \\
 A' \times_{C'} B' & \xrightarrow{\quad} & B' & & \\
 \downarrow & \lrcorner & \downarrow & & \\
 A & \xrightarrow{f} & C & \xrightarrow{g'} & C' \\
 \downarrow & \phi \searrow & \downarrow & \swarrow \chi & \downarrow \\
 A' & \xrightarrow{f'} & C' & &
 \end{array}$$

in  $\text{Fun}(\mathcal{P}, \text{Sets})$  to the map  $\xi: A \times_C B \xrightarrow{\exists!} A' \times_{C'} B'$  given by

$$\xi(a, b) \stackrel{\text{def}}{=} (\phi(a), \psi(b))$$

for each  $(a, b) \in A \times_C B$ , which is the unique map making the diagram

$$\begin{array}{ccccc}
 A \times_C B & \xrightarrow{\quad} & B & & \\
 \downarrow & \lrcorner \searrow & \downarrow g & \searrow \psi & \\
 A' \times_{C'} B' & \xrightarrow{\quad} & B' & & \\
 \downarrow & \lrcorner & \downarrow & & \\
 A & \xrightarrow{f} & C & \xrightarrow{g'} & C' \\
 \downarrow & \phi \searrow & \downarrow & \swarrow \chi & \downarrow \\
 A' & \xrightarrow{f'} & C' & &
 \end{array}$$

commute.

01DX

2. *Adjointness I.* We have adjunctions

$$(A \times_X - \dashv \mathbf{Sets}_{/X}(A, -)): \mathbf{Sets}_{/X} \begin{array}{c} \xrightarrow{\Delta_{\mathbf{Sets}_{/X}}} \\ \perp \\ \xleftarrow{- \times_X A} \end{array} \mathbf{Sets}_{/X},$$

$$(- \times_X B \dashv \mathbf{Sets}_{/X}(B, -)): \mathbf{Sets}_{/X} \begin{array}{c} \xrightarrow{- \times_X B} \\ \perp \\ \xleftarrow{\mathbf{Sets}_{/X}(B, -)} \end{array} \mathbf{Sets}_{/X},$$

witnessed by bijections

$$\mathbf{Sets}_{/X}(A \times_X B, C) \cong \mathbf{Sets}_{/X}(A, \mathbf{Sets}_{/X}(B, C)),$$

$$\mathbf{Sets}_{/X}(A \times_X B, C) \cong \mathbf{Sets}_{/X}(B, \mathbf{Sets}_{/X}(A, C)),$$

natural in  $(A, \phi_A), (B, \phi_B), (C, \phi_C) \in \text{Obj}(\mathbf{Sets}_{/X})$ , where  $\mathbf{Sets}_{/X}(A, B)$  is the object of  $\mathbf{Sets}_{/X}$  consisting of (see ??):

- *The Set.* The set  $\mathbf{Sets}_{/X}(A, B)$  defined by

$$\mathbf{Sets}_{/X}(A, B) \stackrel{\text{def}}{=} \coprod_{x \in X} \mathbf{Sets}(\phi_A^{-1}(x), \phi_B^{-1}(x))$$

- *The Map to X.* The map

$$\phi_{\mathbf{Sets}_{/X}(A, B)}: \mathbf{Sets}_{/X}(A, B) \rightarrow X$$

defined by

$$\phi_{\mathbf{Sets}_{/X}(A, B)}(x, f) \stackrel{\text{def}}{=} x$$

for each  $(x, f) \in \mathbf{Sets}_{/X}(A, B)$ .

01ZD

3. *Adjointness II.* We have an adjunction

$$(\Delta_{\mathbf{Sets}_{/X}} \dashv -_1 \times -_2): \mathbf{Sets}_{/X} \begin{array}{c} \xrightarrow{\Delta_{\mathbf{Sets}_{/X}}} \\ \perp \\ \xleftarrow{-_1 \times -_2} \end{array} \mathbf{Sets}_{/X} \times \mathbf{Sets}_{/X},$$

witnessed by a bijection

$$\text{Hom}_{\mathbf{Sets}_{/X} \times \mathbf{Sets}_{/X}}((A, A), (B, C)) \cong \mathbf{Sets}_{/X}(A, B \times_X C),$$

natural in  $A \in \text{Obj}(\mathbf{Sets}_{/X})$  and in  $(B, C) \in \text{Obj}(\mathbf{Sets}_{/X} \times \mathbf{Sets}_{/X})$ .

001E

4. *Associativity.* Given a diagram

$$\begin{array}{ccccc} A & & B & & C \\ & \searrow f & & \downarrow g & \searrow h \\ & X & & Y & \end{array}$$

in Sets, we have isomorphisms of sets

$$(A \times_X B) \times_Y C \cong (A \times_X B) \times_B (B \times_Y C) \cong A \times_X (B \times_Y C),$$

where these pullbacks are built as in the diagrams

$$\begin{array}{ccc} \begin{array}{c} (A \times_X B) \times_Y C \\ \swarrow \quad \searrow \\ A \times_X B \end{array} & \begin{array}{c} (A \times_X B) \times_B (B \times_Y C) \\ \swarrow \quad \searrow \\ A \times_X B \quad B \times_Y C \\ \downarrow \quad \downarrow \\ A \times_X (B \times_Y C) \end{array} & \begin{array}{c} A \times_X (B \times_Y C) \\ \swarrow \quad \searrow \\ B \times_Y C \end{array} \\ \begin{array}{ccccc} A & & B & & C \\ \downarrow & \swarrow & \downarrow & \searrow & \\ A & & B & & C \\ \downarrow f & \downarrow g & \downarrow h & \downarrow k & \\ X & & Y & & \end{array} & \begin{array}{ccccc} A & & B & & C \\ \downarrow & \swarrow & \downarrow & \searrow & \\ A & & B & & C \\ \downarrow f & \downarrow g & \downarrow h & \downarrow k & \\ X & & Y & & \end{array} & \begin{array}{ccccc} A & & B & & C \\ \downarrow & \swarrow & \downarrow & \searrow & \\ A & & B & & C \\ \downarrow f & \downarrow g & \downarrow h & \downarrow k & \\ X & & Y & & \end{array} \end{array}$$

01DY

5. *Interaction With Composition.* Given a diagram

$$\begin{array}{ccccc} X & & & & Y \\ & \searrow \phi & & \swarrow \psi & \\ & A & & B & \\ & \searrow f & & \swarrow g & \\ & K & & & \end{array}$$

in Sets, we have isomorphisms of sets

$$\begin{aligned} X \times_K^{f \circ \phi, g \circ \psi} Y &\cong (X \times_A^{\phi, q_1} (A \times_K^{f, g} B)) \times_{A \times_K^{f, g} B}^{p_2, p_1} ((A \times_K^{f, g} B) \times_B^{q_2, \psi} Y) \\ &\cong X \times_A^{\phi, p} ((A \times_K^{f, g} B) \times_B^{q_2, \psi} Y) \\ &\cong (X \times_A^{\phi, q_1} (A \times_K^{f, g} B)) \times_B^{q, \psi} Y \end{aligned}$$

where

$$\begin{array}{ll} q_1 = \text{pr}_1^{A \times_K^{f, g} B}, & q_2 = \text{pr}_2^{A \times_K^{f, g} B}, \\ p_1 = \text{pr}_1^{(A \times_K^{f, g} B) \times_B^{q_2, \psi}}, & p_2 = \text{pr}_2^{X \times_{A \times_K^{f, g} B}^{\phi, q_1} (A \times_K^{f, g} B)}, \\ p = q_1 \circ \text{pr}_1^{(A \times_K^{f, g} B) \times_B^{q_2, \psi}}, & q = q_2 \circ \text{pr}_2^{X \times_A^{\phi, q_1} (A \times_K^{f, g} B)}, \end{array}$$

and where these pullbacks are built as in the following diagrams:

$$\begin{array}{ccc}
 \begin{array}{c}
 (X \times_A (A \times_K B)) \times_{A \times_K B} ((A \times_K B) \times_B Y) \\
 \downarrow \quad \downarrow \\
 X \times_A (A \times_K B) \quad (A \times_K B) \times_B Y \\
 \downarrow \quad \downarrow \\
 X \quad A \times_K B \quad Y, \\
 \downarrow \quad \downarrow \quad \downarrow \\
 X \quad A \quad B \quad Y, \\
 \downarrow \phi \quad \downarrow \quad \downarrow \psi \\
 A \quad f \quad g \quad K \\
 \downarrow \quad \downarrow \\
 K
 \end{array}
 & \quad &
 \begin{array}{c}
 X \times_A ((A \times_K B) \times_B Y) \\
 \downarrow \quad \downarrow \\
 (A \times_K B) \times_B Y \\
 \downarrow \quad \downarrow \\
 A \times_K B \quad Y, \\
 \downarrow \quad \downarrow \\
 A \quad B \quad Y, \\
 \downarrow \phi \quad \downarrow \quad \downarrow \psi \\
 A \quad f \quad g \quad K \\
 \downarrow \quad \downarrow \\
 K
 \end{array}
 \end{array}$$
  

$$\begin{array}{ccc}
 \begin{array}{c}
 (X \times_A (A \times_K B)) \times_B Y \\
 \downarrow \quad \searrow \\
 X \times_A (A \times_K B) \quad Y, \\
 \downarrow \quad \downarrow \\
 X \quad A \times_K B \quad Y, \\
 \downarrow \phi \quad \downarrow \quad \downarrow \psi \\
 A \quad f \quad g \quad K \\
 \downarrow \quad \downarrow \\
 K
 \end{array}
 & \quad &
 \begin{array}{c}
 X \times_K Y \\
 \downarrow \quad \searrow \\
 A \times_X X \quad Y, \\
 \downarrow \quad \downarrow \\
 A \quad B \quad Y, \\
 \downarrow \phi \quad \downarrow \quad \downarrow \psi \\
 A \quad f \quad g \quad K \\
 \downarrow \quad \downarrow \\
 K
 \end{array}
 \end{array}$$

001F

6. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc}
 A & \xlongequal{\quad} & A \\
 \downarrow f & \lrcorner & \downarrow f \\
 X & \xlongequal{\quad} & X
 \end{array}
 \quad
 \begin{array}{l}
 \lambda_A^{\text{Sets}/X} : X \times_X A \xrightarrow{\sim} A, \\
 \rho_A^{\text{Sets}/X} : A \times_X X \xrightarrow{\sim} A,
 \end{array}
 \quad
 \begin{array}{ccc}
 A & \xrightarrow{f} & X \\
 \parallel & \lrcorner & \parallel \\
 X & \xrightarrow{f} & X,
 \end{array}$$

natural in  $(A, f) \in \text{Obj}(\text{Sets}/X)$ .

001G

7. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc}
 A \times_X B & \longrightarrow & B \\
 \downarrow \lrcorner & & \downarrow g \\
 A & \xrightarrow{f} & X, \\
 & & \sigma_{A,B}^{\text{Sets}/X} : A \times_X B \xrightarrow{\sim} B \times_X A
 \end{array}
 \quad
 \begin{array}{ccc}
 B \times_X A & \longrightarrow & A \\
 \downarrow \lrcorner & & \downarrow f \\
 B & \xrightarrow{g} & X,
 \end{array}$$

natural in  $(A, f), (B, g) \in \text{Obj}(\text{Sets}/X)$ .

01DZ

8. *Distributivity Over Coproducts.* Let  $A$ ,  $B$ , and  $C$  be sets and let  $\phi_A: A \rightarrow X$ ,  $\phi_B: B \rightarrow X$ , and  $\phi_C: C \rightarrow X$  be morphisms of sets. We have isomorphisms of sets

$$\begin{aligned}\delta_\ell^{\text{Sets}/X}: A \times_X (B \coprod C) &\xrightarrow{\sim} (A \times_X B) \coprod (A \times_X C), \\ \delta_r^{\text{Sets}/X}: (A \coprod B) \times_X C &\xrightarrow{\sim} (A \times_X C) \coprod (B \times_X C),\end{aligned}$$

as in the diagrams

$$\begin{array}{ccc} (A \times_X B) \coprod (A \times_X C) & \longrightarrow & B \coprod C \\ \downarrow & \lrcorner & \downarrow \phi_B \coprod \phi_C \\ A & \xrightarrow{\phi_A} & X \end{array} \quad \begin{array}{ccc} (A \times_X C) \coprod (B \times_X C) & \longrightarrow & C \\ \downarrow & \lrcorner & \downarrow \phi_C \\ A \coprod B & \xrightarrow{\phi_A \coprod \phi_B} & X \end{array}$$

natural in  $A, B, C \in \text{Obj}(\text{Sets}/X)$ .

001H

9. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{array}{ccc} \emptyset & \longrightarrow & \emptyset \\ \downarrow & \lrcorner & \downarrow \\ A & \xrightarrow{f} & X, \end{array} \quad \begin{array}{c} \zeta_\ell^{\text{Sets}/X}: A \times_X \emptyset \xrightarrow{\sim} \emptyset, \\ \zeta_r^{\text{Sets}/X}: \emptyset \times_X A \xrightarrow{\sim} \emptyset, \end{array} \quad \begin{array}{ccc} \emptyset & \longrightarrow & A \\ \downarrow & \lrcorner & \downarrow f \\ \emptyset & \longrightarrow & X, \end{array}$$

natural in  $(A, f) \in \text{Obj}(\text{Sets}/X)$ .

001J

10. *Interaction With Products.* We have an isomorphism of sets

$$\begin{array}{ccc} A \times B & \longrightarrow & B \\ \downarrow & \lrcorner & \downarrow !_B \\ A \times_{\text{pt}} B \cong A \times B, & & \\ \downarrow & & \downarrow \\ A & \xrightarrow{!_A} & \text{pt}. \end{array}$$

001K

11. *Symmetric Monoidality.* The 8-tuple  $(\text{Sets}/X, \times_X, X, \text{Sets}/X, \alpha^{\text{Sets}/X}, \lambda^{\text{Sets}/X}, \rho^{\text{Sets}/X}, \sigma^{\text{Sets}/X})$  is a symmetric closed monoidal category.

**PROOF 4.1.4.1.8 ► PROOF OF PROPOSITION 4.1.4.1.7****Item 1: Functoriality**

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for  $\xi$  following from the commutativity of the cube pullback diagram.

**Item 2: Adjointness I**

This is a repetition of ?? of ??, and is proved there.

**Item 3: Adjointness II**

This follows from the universal property of the product (pullbacks are products in  $\text{Sets}_{/X}$ ).

**Item 4: Associativity**

We have

$$\begin{aligned} (A \times_X B) \times_Y C &\cong \{((a, b), c) \in (A \times_X B) \times C \mid h(b) = k(c)\} \\ &\cong \{((a, b), c) \in (A \times B) \times C \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\ &\cong \{(a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\ &\cong \{(a, (b, c)) \in A \times (B \times_Y C) \mid f(a) = g(b)\} \\ &\cong A \times_X (B \times_Y C) \end{aligned}$$

and

$$\begin{aligned} (A \times_X B) \times_B (B \times_Y C) &\cong \{((a, b), (b', c)) \in (A \times_X B) \times (B \times_Y C) \mid b = b'\} \\ &\cong \left\{ ((a, b), (b', c)) \in (A \times B) \times (B \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\ &\cong \left\{ (a, (b, (b', c))) \in A \times (B \times (B \times C)) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\ &\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times B) \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\ &\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times_B B) \times C) \mid \begin{array}{l} f(a) = g(b) \text{ and} \\ h(b') = k(c) \end{array} \right\} \\ &\cong \{(a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\ &\cong A \times_X (B \times_Y C), \end{aligned}$$

where we have used **Item 6** for the isomorphism  $B \times_B B \cong B$ .

**Item 5: Interaction With Composition**

By **Item 4**, it suffices to construct only the isomorphism

$$X \times_K^{f \circ \phi, g \circ \psi} Y \cong (X \times_A^{\phi, q_1} (A \times_K^{f, g} B)) \times_{A \times_K^{f, g} B}^{p_2, p_1} ((A \times_K^{f, g} B) \times_B^{q_2, \psi} Y).$$

We have

$$\begin{aligned}
 (X \times_A^{\phi, q_1} (A \times_K^{f,g} B)) &\stackrel{\text{def}}{=} \left\{ (x, (a, b)) \in X \times (A \times_K^{f,g} B) \mid \phi(x) = q_1(a, b) \right\} \\
 &\stackrel{\text{def}}{=} \left\{ (x, (a, b)) \in X \times (A \times_K^{f,g} B) \mid \phi(x) = a \right\} \\
 &\cong \{(x, (a, b)) \in X \times (A \times B) \mid \phi(x) = a \text{ and } f(a) = g(b)\}, \\
 ((A \times_K^{f,g} B) \times_B^{q_2, \psi} Y) &\stackrel{\text{def}}{=} \left\{ ((a, b), y) \in (A \times_K^{f,g} B) \times Y \mid q_2(a, b) = \psi(y) \right\} \\
 &\stackrel{\text{def}}{=} \left\{ ((a, b), y) \in (A \times_K^{f,g} B) \times Y \mid b = \psi(y) \right\} \\
 &\cong \{((a, b), y) \in (A \times B) \times Y \mid b = \psi(y) \text{ and } f(a) = g(b)\},
 \end{aligned}$$

so writing

$$\begin{aligned}
 S &= (X \times_A^{\phi, q_1} (A \times_K^{f,g} B)) \\
 S' &= ((A \times_K^{f,g} B) \times_B^{q_2, \psi} Y),
 \end{aligned}$$

we have

$$\begin{aligned}
 S \times_{A \times_K^{f,g} B}^{p_2, p_1} S' &\stackrel{\text{def}}{=} \{((x, (a, b)), ((a', b'), y)) \in S \times S' \mid p_1(x, (a, b)) = p_2((a', b'), y)\} \\
 &\stackrel{\text{def}}{=} \{((x, (a, b)), ((a', b'), y)) \in S \times S' \mid (a, b) = (a', b')\} \\
 &\cong \{((x, a, b, y)) \in X \times A \times B \times Y \mid \phi(x) = a, \psi(y) = b, \text{ and } f(a) = g(b)\} \\
 &\cong \{((x, a, b, y)) \in X \times A \times B \times Y \mid f(\phi(x)) = g(\psi(y))\} \\
 &\stackrel{\text{def}}{=} X \times_K Y.
 \end{aligned}$$

This finishes the proof.

#### Item 6: Unitality

We have

$$\begin{aligned}
 X \times_X A &\cong \{(x, a) \in X \times A \mid f(a) = x\}, \\
 A \times_X X &\cong \{(a, x) \in X \times A \mid f(a) = x\},
 \end{aligned}$$

which are isomorphic to  $A$  via the maps  $(x, a) \mapsto a$  and  $(a, x) \mapsto a$ . The proof of the naturality of  $\lambda^{\text{Sets}/X}$  and  $\rho^{\text{Sets}/X}$  is omitted.

#### Item 7: Commutativity

We have

$$\begin{aligned}
 A \times_C B &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\} \\
 &= \{(a, b) \in A \times B \mid g(b) = f(a)\} \\
 &\cong \{(b, a) \in B \times A \mid g(b) = f(a)\} \\
 &\stackrel{\text{def}}{=} B \times_C A.
 \end{aligned}$$

The proof of the naturality of  $\sigma^{\text{Sets}/X}$  is omitted.

Item 8: Distributivity Over Coproducts

We have

$$\begin{aligned}
 A \times_X (B \coprod C) &\stackrel{\text{def}}{=} \left\{ (a, z) \in A \times (B \coprod C) \mid \phi_A(a) = \phi_{B \coprod C}(z) \right\} \\
 &= \left\{ (a, z) \in A \times (B \coprod C) \mid z = (0, b) \text{ and } \phi_A(a) = \phi_B(z) \right\} \\
 &\quad \cup \left\{ (a, z) \in A \times (B \coprod C) \mid z = (1, c) \text{ and } \phi_A(a) = \phi_C(z) \right\} \\
 &= \{(a, z) \in A \times (B \coprod C) \mid z = (0, b) \text{ and } \phi_A(a) = \phi_B(b)\} \\
 &\quad \cup \{(a, z) \in A \times (B \coprod C) \mid z = (1, c) \text{ and } \phi_A(a) = \phi_C(c)\} \\
 &\cong \{(a, b) \in A \times B \mid \phi_A(a) = \phi_B(b)\} \\
 &\quad \cup \{(a, c) \in A \times C \mid \phi_A(a) = \phi_C(c)\} \\
 &\stackrel{\text{def}}{=} (A \times_X B) \cup (A \times_X C) \\
 &\cong (A \times_X B) \coprod (A \times_X C),
 \end{aligned}$$

with the construction of the isomorphism

$$\delta_r^{\text{Sets}/X} : (A \coprod B) \times_X C \xrightarrow{\sim} (A \times_X C) \coprod (B \times_X C)$$

being similar. The proof of the naturality of  $\delta_\ell^{\text{Sets}/X}$  and  $\delta_r^{\text{Sets}/X}$  is omitted.

Item 9: Annihilation With the Empty Set

We have

$$\begin{aligned}
 A \times_X \emptyset &\stackrel{\text{def}}{=} \{(a, b) \in A \times \emptyset \mid f(a) = g(b)\} \\
 &= \{k \in \emptyset \mid f(a) = g(b)\} \\
 &= \emptyset,
 \end{aligned}$$

and similarly for  $\emptyset \times_X A$ , where we have used Item 8 of Proposition 4.1.3.1.4. The proof of the naturality of  $\zeta_\ell^{\text{Sets}/X}$  and  $\zeta_r^{\text{Sets}/X}$  is omitted.

Item 10: Interaction With Products

We have

$$\begin{aligned}
 A \times_{\text{pt}} B &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid !_A(a) = !_B(b)\} \\
 &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid \star = \star\} \\
 &= \{(a, b) \in A \times B\} \\
 &= A \times B.
 \end{aligned}$$

Item 11: Symmetric Monoidality

Omitted. ☒

### 001L 4.1.5 Equalisers

Let  $A$  and  $B$  be sets and let  $f, g: A \rightrightarrows B$  be functions.

001M

#### DEFINITION 4.1.5.1.1 ► EQUALISERS OF SETS

The **equaliser of  $f$  and  $g$**  is the equaliser of  $f$  and  $g$  in Sets as in ??.

01E0

#### CONSTRUCTION 4.1.5.1.2 ► CONSTRUCTION OF EQUALISERS OF SETS

Concretely, the equaliser of  $f$  and  $g$  is the pair  $(\text{Eq}(f, g), \text{eq}(f, g))$  consisting of:

01E1

1. *The Limit.* The set  $\text{Eq}(f, g)$  defined by

$$\text{Eq}(f, g) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\}.$$

01E2

2. *The Cone.* The inclusion map

$$\text{eq}(f, g): \text{Eq}(f, g) \hookrightarrow A.$$

#### PROOF 4.1.5.1.3 ► PROOF OF CONSTRUCTION 4.1.5.1.2

We claim that  $\text{Eq}(f, g)$  is the categorical equaliser of  $f$  and  $g$  in Sets. First we need to check that the relevant equaliser diagram commutes, i.e. that we have

$$f \circ \text{eq}(f, g) = g \circ \text{eq}(f, g),$$

which indeed holds by the definition of the set  $\text{Eq}(f, g)$ . Next, we prove that  $\text{Eq}(f, g)$  satisfies the universal property of the equaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A \rightrightarrows B \\ & \nearrow e & \\ E & & \end{array}$$

in Sets. Then there exists a unique map  $\phi: E \rightarrow \text{Eq}(f, g)$  making the diagram

$$\begin{array}{ccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A \rightrightarrows B \\ \phi \uparrow \exists! & \nearrow e & \\ E & & \end{array}$$

commute, being uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

via

$$\phi(x) = e(x)$$

for each  $x \in E$ , where we note that  $e(x) \in A$  indeed lies in  $\text{Eq}(f, g)$  by the condition

$$f \circ e = g \circ e,$$

which gives

$$f(e(x)) = g(e(x))$$

for each  $x \in E$ , so that  $e(x) \in \text{Eq}(f, g)$ . □

001N

#### PROPOSITION 4.1.5.1.4 ► PROPERTIES OF EQUALISERS OF SETS

Let  $A$ ,  $B$ , and  $C$  be sets.

001P

1. *Associativity.* We have isomorphisms of sets<sup>1</sup>

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))},$$

where  $\text{Eq}(f, g, h)$  is the limit of the diagram

$$\begin{array}{ccc} & f & \\ A & \xrightarrow{\quad g \rightarrow \quad} & B \\ & h & \end{array}$$

in Sets, being explicitly given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

001Q

2. *Unitality.* We have an isomorphism of sets

$$\text{Eq}(f, f) \cong A.$$

001R

3. *Commutativity.* We have an isomorphism of sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

001S

4. *Interaction With Composition.* Let

$$\begin{array}{ccc} & f & \\ A & \xrightarrow[g]{} & B \xrightarrow[k]{} C \end{array}$$

be functions. We have an inclusion of sets

$$\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) \subset \text{Eq}(h \circ f, k \circ g),$$

where  $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$  is the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \xrightarrow[\substack{g \\ f}]{} B \xrightarrow[\substack{h \\ k}]{} C.$$

<sup>1</sup>That is, the following three ways of forming “the” equaliser of  $(f, g, h)$  agree:

- 01ZE (a) Take the equaliser of  $(f, g, h)$ , i.e. the limit of the diagram

$$\begin{array}{ccc} & f & \\ A & \xrightarrow[\substack{g \\ h}]{} & B \end{array}$$

in Sets.

- 01ZF (b) First take the equaliser of  $f$  and  $g$ , forming a diagram

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \xrightarrow[\substack{f \\ g}]{} B$$

and then take the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \xrightarrow[\substack{f \\ h}]{} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) = \text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))$$

of  $\text{Eq}(f, g)$ .

- 01ZG (c) First take the equaliser of  $g$  and  $h$ , forming a diagram

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \xrightarrow[\substack{g \\ h}]{} B$$

and then take the equaliser of the composition

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \xrightarrow[\substack{f \\ g}]{} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) = \text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))$$

of  $\text{Eq}(g, h)$ .

**PROOF 4.1.5.5 ► PROOF OF PROPOSITION 4.1.5.1.4**
**Item 1: Associativity**

We first prove that  $\text{Eq}(f, g, h)$  is indeed given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} \text{Eq}(f, g, h) & \xrightarrow{\text{eq}(f, g, h)} & A \xrightarrow{\begin{matrix} f \\ -g \\ h \end{matrix}} B \\ & \searrow e & \\ E & & \end{array}$$

in Sets. Then there exists a unique map  $\phi: E \rightarrow \text{Eq}(f, g, h)$ , uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

being necessarily given by

$$\phi(x) = e(x)$$

for each  $x \in E$ , where we note that  $e(x) \in A$  indeed lies in  $\text{Eq}(f, g, h)$  by the condition

$$f \circ e = g \circ e = h \circ e,$$

which gives

$$f(e(x)) = g(e(x)) = h(e(x))$$

for each  $x \in E$ , so that  $e(x) \in \text{Eq}(f, g, h)$ .

We now check the equalities

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) \cong \text{Eq}(f, g, h) \cong \text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)).$$

Indeed, we have

$$\begin{aligned} \text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) &\cong \{x \in \text{Eq}(g, h) \mid [f \circ \text{eq}(g, h)](a) = [g \circ \text{eq}(g, h)](a)\} \\ &\cong \{x \in \text{Eq}(g, h) \mid f(a) = g(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) \text{ and } g(a) = h(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\ &\cong \text{Eq}(f, g, h). \end{aligned}$$

Similarly, we have

$$\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) \cong \{x \in \text{Eq}(f, g) \mid [f \circ \text{eq}(f, g)](a) = [h \circ \text{eq}(f, g)](a)\}$$

$$\begin{aligned}
&\cong \{x \in \text{Eq}(f, g) \mid f(a) = h(a)\} \\
&\cong \{x \in A \mid f(a) = h(a) \text{ and } f(a) = g(a)\} \\
&\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\
&\cong \text{Eq}(f, g, h).
\end{aligned}$$

**Item 2: Unitality**

Indeed, we have

$$\begin{aligned}
\text{Eq}(f, f) &\stackrel{\text{def}}{=} \{a \in A \mid f(a) = f(a)\} \\
&= A.
\end{aligned}$$

**Item 3: Commutativity**

Indeed, we have

$$\begin{aligned}
\text{Eq}(f, g) &\stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\} \\
&= \{a \in A \mid g(a) = f(a)\} \\
&\stackrel{\text{def}}{=} \text{Eq}(g, f).
\end{aligned}$$

**Item 4: Interaction With Composition**

Indeed, we have

$$\begin{aligned}
\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) &\cong \{a \in \text{Eq}(f, g) \mid h(f(a)) = k(g(a))\} \\
&\cong \{a \in A \mid f(a) = g(a) \text{ and } h(f(a)) = k(g(a))\}.
\end{aligned}$$

and

$$\text{Eq}(h \circ f, k \circ g) \cong \{a \in A \mid h(f(a)) = k(g(a))\},$$

and thus there's an inclusion from  $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$  to  $\text{Eq}(h \circ f, k \circ g)$ . □

**01E3 4.1.6 Inverse Limits**

Let  $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}: (I, \preceq) \rightarrow \text{Sets}$  be an inverse system of sets.

**01E4**

**DEFINITION 4.1.6.1.1 ► INVERSE LIMITS OF SETS**

The **inverse limit of**  $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$  is the inverse limit of  $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$  in Sets as in ??.

01E5

## CONSTRUCTION 4.1.6.1.2 ► CONSTRUCTION OF INVERSE LIMITS OF SETS

Concretely, the inverse limit of  $(X_\alpha, f_{\alpha\beta})_{\alpha,\beta \in I}$  is the pair  $\left( \varprojlim_{\alpha \in I} (X_\alpha), \{pr_\alpha\}_{\alpha \in I} \right)$  consisting of:

01E6

1. *The Limit.* The set  $\varprojlim_{\alpha \in I} (X_\alpha)$  defined by

$$\varprojlim_{\alpha \in I} (X_\alpha) \stackrel{\text{def}}{=} \left\{ (x_\alpha)_{\alpha \in I} \in \prod_{\alpha \in I} X_\alpha \mid \begin{array}{l} \text{for each } \alpha, \beta \in I, \text{ if } \alpha \preceq \beta, \\ \text{then we have } x_\alpha = f_{\alpha\beta}(x_\beta) \end{array} \right\}.$$

01E7

2. *The Cone.* The collection

$$\left\{ pr_\gamma : \varprojlim_{\alpha \in I} (X_\alpha) \rightarrow X_\gamma \right\}_{\gamma \in I}$$

of maps of sets defined as the restriction of the maps

$$\left\{ pr_\gamma : \prod_{\alpha \in I} X_\alpha \rightarrow X_\gamma \right\}_{\gamma \in I}$$

of Item 2 of Construction 4.1.2.1.2 to  $\varprojlim_{\alpha \in I} (X_\alpha)$  and hence given by

$$pr_\gamma((x_\alpha)_{\alpha \in I}) \stackrel{\text{def}}{=} x_\gamma$$

for each  $\gamma \in I$  and each  $(x_\alpha)_{\alpha \in I} \in \varprojlim_{\alpha \in I} (X_\alpha)$ .

## PROOF 4.1.6.1.3 ► PROOF OF CONSTRUCTION 4.1.6.1.2

We claim that  $\varprojlim_{\alpha \in I} (X_\alpha)$  is the limit of the inverse system of sets  $(X_\alpha, f_{\alpha\beta})_{\alpha,\beta \in I}$ . First we need to check that the limit diagram defined by it commutes, i.e. that we have

$$\begin{array}{ccc} & \varprojlim_{\alpha \in I} (X_\alpha) & \\ f_{\alpha\beta} \circ pr_\alpha & = pr_\beta, & \\ & \swarrow pr_\alpha \quad \searrow pr_\beta & \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$

for each  $\alpha, \beta \in I$  with  $\alpha \preceq \beta$ . Indeed, given  $(x_\gamma)_{\gamma \in I} \in \lim_{\leftarrow \gamma \in I} (X_\gamma)$ , we have

$$\begin{aligned} [f_{\alpha\beta} \circ \text{pr}_\alpha]((x_\gamma)_{\gamma \in I}) &\stackrel{\text{def}}{=} f_{\alpha\beta}(\text{pr}_\alpha((x_\gamma)_{\gamma \in I})) \\ &\stackrel{\text{def}}{=} f_{\alpha\beta}(x_\alpha) \\ &= x_\beta \\ &\stackrel{\text{def}}{=} \text{pr}_\beta((x_\gamma)_{\gamma \in I}), \end{aligned}$$

where the third equality comes from the definition of  $\lim_{\leftarrow \alpha \in I} (X_\alpha)$ . Next, we prove that  $\lim_{\leftarrow \alpha \in I} (X_\alpha)$  satisfies the universal property of an inverse limit. Suppose that we have, for each  $\alpha, \beta \in I$  with  $\alpha \preceq \beta$ , a diagram of the form

$$\begin{array}{ccc} & L & \\ p_\alpha \swarrow & \downarrow \lim_{\leftarrow \alpha \in I} (X_\alpha) & \searrow p_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \\ \text{pr}_\alpha \swarrow & & \searrow \text{pr}_\beta \\ & & \end{array}$$

in Sets. Then there indeed exists a unique map  $\phi: L \xrightarrow{\exists!} \lim_{\leftarrow \alpha \in I} (X_\alpha)$  making the diagram

$$\begin{array}{ccc} & L & \\ p_\alpha \swarrow & \downarrow \phi \exists! & \searrow p_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \\ \text{pr}_\alpha \swarrow & & \searrow \text{pr}_\beta \\ & & \end{array}$$

commute, being uniquely determined by the family of conditions

$$\{p_\alpha = \text{pr}_\alpha \circ \phi\}_{\alpha \in I}$$

via

$$\phi(\ell) = (p_\alpha(\ell))_{\alpha \in I}$$

for each  $\ell \in L$ , where we note that  $(p_\alpha(\ell))_{\alpha \in I} \in \prod_{\alpha \in I} X_\alpha$  indeed lies in  $\varprojlim_{\alpha \in I} (X_\alpha)$ , as we have

$$\begin{aligned} f_{\alpha\beta}(p_\alpha(\ell)) &\stackrel{\text{def}}{=} [f_{\alpha\beta} \circ p_\alpha](\ell) \\ &\stackrel{\text{def}}{=} p_\beta(\ell) \end{aligned}$$

for each  $\beta \in I$  with  $\alpha \preceq \beta$  by the commutativity of the diagram for  $(L, \{p_\alpha\}_{\alpha \in I})$ . 

01E8

**EXAMPLE 4.1.6.1.4 ► EXAMPLES OF INVERSE LIMITS OF SETS**

Here are some examples of inverse limits of sets.

01E9

1. *The  $p$ -Adic Integers.* The ring of  $p$ -adic integers  $\mathbb{Z}_p$  of ?? is the inverse limit

$$\mathbb{Z}_p \cong \varprojlim_{n \in \mathbb{N}} (\mathbb{Z}/p^n);$$

see ??.

01EA

2. *Rings of Formal Power Series.* The ring  $R[[t]]$  of formal power series in a variable  $t$  is the inverse limit

$$R[[t]] \cong \varprojlim_{n \in \mathbb{N}} (R[t]/t^n R[t]);$$

see ??.

01EB

3. *Profinite Groups.* Profinite groups are inverse limits of finite groups; see ??.

001T

**4.2 Colimits of Sets**

001U

**4.2.1 The Initial Set**

001V

**DEFINITION 4.2.1.1.1 ► THE INITIAL SET**

The **initial set** is the initial object of Sets as in ??.

01EC

## CONSTRUCTION 4.2.1.1.2 ► CONSTRUCTION OF THE INITIAL SET

Concretely, the initial set is the pair  $(\emptyset, \{\iota_A\}_{A \in \text{Obj}(\text{Sets})})$  consisting of:

01ED

1. *The Colimit.* The empty set  $\emptyset$  of [Definition 4.3.1.1.1](#).

01EE

2. *The Cocone.* The collection of maps

$$\{\iota_A : \emptyset \rightarrow A\}_{A \in \text{Obj}(\text{Sets})}$$

given by the inclusion maps from  $\emptyset$  to  $A$ .

## PROOF 4.2.1.1.3 ► PROOF OF CONSTRUCTION 4.2.1.1.2

We claim that  $\emptyset$  is the initial object of Sets. Indeed, suppose we have a diagram of the form

$$\emptyset \quad A$$

in Sets. Then there exists a unique map  $\phi : \emptyset \rightarrow A$  making the diagram

$$\emptyset \xrightarrow[\exists!]{\phi} A$$

commute, namely the inclusion map  $\iota_A$ . 

## 001W 4.2.2 Coproducts of Families of Sets

Let  $\{A_i\}_{i \in I}$  be a family of sets.

001X

## DEFINITION 4.2.2.1.1 ► THE COPRODUCT OF A FAMILY OF SETS

The **coproduct** of  $\{A_i\}_{i \in I}$ <sup>1</sup> is the coproduct of  $\{A_i\}_{i \in I}$  in Sets as in ??.

<sup>1</sup>Further Terminology: Also called the **disjoint union** of the family  $\{A_i\}_{i \in I}$ .

01EF

## CONSTRUCTION 4.2.2.1.2 ► CONSTRUCTION OF THE COPRODUCT OF A FAMILY OF SETS

Concretely, the disjoint union of  $\{A_i\}_{i \in I}$  is the pair  $(\coprod_{i \in I} A_i, \{\text{inj}_i\}_{i \in I})$  consisting of:

01EG

1. *The Colimit.* The set  $\coprod_{i \in I} A_i$  defined by

$$\coprod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ (i, x) \in I \times \left( \bigcup_{i \in I} A_i \right) \mid x \in A_i \right\}.$$

01EH

2. *The Cocone.* The collection

$$\left\{ \text{inj}_i : A_i \rightarrow \coprod_{i \in I} A_i \right\}_{i \in I}$$

of maps given by

$$\text{inj}_i(x) \stackrel{\text{def}}{=} (i, x)$$

for each  $x \in A_i$  and each  $i \in I$ .

#### PROOF 4.2.2.1.3 ► PROOF OF CONSTRUCTION 4.2.2.1.2

We claim that  $\coprod_{i \in I} A_i$  is the categorical coproduct of  $\{A_i\}_{i \in I}$  in Sets. Indeed, suppose we have, for each  $i \in I$ , a diagram of the form

$$\begin{array}{ccc} & & C \\ & \nearrow \iota_i & \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

in Sets. Then there exists a unique map  $\phi : \coprod_{i \in I} A_i \rightarrow C$  making the diagram

$$\begin{array}{ccc} & & C \\ & \nearrow \iota_i & \uparrow \phi \exists! \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

commute, being uniquely determined by the condition  $\phi \circ \text{inj}_i = \iota_i$  for each  $i \in I$  via

$$\phi((i, x)) = \iota_i(x)$$

for each  $(i, x) \in \coprod_{i \in I} A_i$ . □

001Y

#### PROPOSITION 4.2.2.1.4 ► PROPERTIES OF COPRODUCTS OF FAMILIES OF SETS

Let  $\{A_i\}_{i \in I}$  be a family of sets.

001Z

1. *Functionality.* The assignment  $\{A_i\}_{i \in I} \mapsto \coprod_{i \in I} A_i$  defines a functor

$$\coprod_{i \in I} : \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each  $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$ , we have

$$\left[ \coprod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \coprod_{i \in I} A_i$$

- *Action on Morphisms.* For each  $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$ , the action on Hom-sets

$$\left( \coprod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left( \coprod_{i \in I} A_i, \coprod_{i \in I} B_i \right)$$

of  $\coprod_{i \in I}$  at  $((A_i)_{i \in I}, (B_i)_{i \in I})$  is defined by sending a map

$$\{f_i : A_i \rightarrow B_i\}_{i \in I}$$

in  $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$  to the map of sets

$$\coprod_{i \in I} f_i : \coprod_{i \in I} A_i \rightarrow \coprod_{i \in I} B_i$$

defined by

$$\left[ \coprod_{i \in I} f_i \right] (i, a) \stackrel{\text{def}}{=} f_i(a)$$

for each  $(i, a) \in \coprod_{i \in I} A_i$ .

#### PROOF 4.2.2.1.5 ▶ PROOF OF PROPOSITION 4.2.2.1.4

Item 1: Functoriality

This follows from ?? of ??.



### 0020 4.2.3 Binary Coproducts

Let  $A$  and  $B$  be sets.

0021

#### DEFINITION 4.2.3.1.1 ▶ COPRODUCTS OF SETS

The **coproduct of  $A$  and  $B$** <sup>1</sup> is the coproduct of  $A$  and  $B$  in Sets as in ??.

<sup>1</sup>Further Terminology: Also called the **disjoint union of  $A$  and  $B$** .

01EJ

## CONSTRUCTION 4.2.3.1.2 ► CONSTRUCTION OF COPRODUCTS OF SETS

Concretely, the coproduct of  $A$  and  $B$  is the pair  $(A \coprod B, \{\text{inj}_1, \text{inj}_2\})$  consisting of:

01EK

1. *The Colimit.* The set  $A \coprod B$  defined by

$$\begin{aligned} A \coprod B &\stackrel{\text{def}}{=} \coprod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{(0, a) \in S \mid a \in A\} \cup \{(1, b) \in S \mid b \in B\}, \end{aligned}$$

where  $S = \{0, 1\} \times (A \cup B)$ .

01EL

2. *The Cocone.* The maps

$$\begin{aligned} \text{inj}_1 &: A \rightarrow A \coprod B, \\ \text{inj}_2 &: B \rightarrow A \coprod B, \end{aligned}$$

given by

$$\begin{aligned} \text{inj}_1(a) &\stackrel{\text{def}}{=} (0, a), \\ \text{inj}_2(b) &\stackrel{\text{def}}{=} (1, b), \end{aligned}$$

for each  $a \in A$  and each  $b \in B$ .

## PROOF 4.2.3.1.3 ► PROOF OF CONSTRUCTION 4.2.3.1.2

We claim that  $A \coprod B$  is the categorical coproduct of  $A$  and  $B$  in Sets. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & C & & \\ & \nearrow t_1 & & \swarrow t_2 & \\ A & \xhookrightarrow{\text{inj}_1} & A \coprod B & \xhookleftarrow{\text{inj}_2} & B \end{array}$$

in Sets. Then there exists a unique map  $\phi: A \coprod B \rightarrow C$  making the diagram

$$\begin{array}{ccccc} & & C & & \\ & \nearrow t_1 & \uparrow \phi \exists! & \swarrow t_2 & \\ A & \xhookrightarrow{\text{inj}_1} & A \coprod B & \xhookleftarrow{\text{inj}_2} & B \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned}\phi \circ \text{inj}_A &= \iota_A, \\ \phi \circ \text{inj}_B &= \iota_B\end{aligned}$$

via

$$\phi(x) = \begin{cases} \iota_A(a) & \text{if } x = (0, a), \\ \iota_B(b) & \text{if } x = (1, b) \end{cases}$$

for each  $x \in A \coprod B$ . ■

0022

#### PROPOSITION 4.2.3.1.4 ► PROPERTIES OF COPRODUCTS OF SETS

Let  $A, B, C$ , and  $X$  be sets.

0023

1. *Functoriality.* The assignment  $A, B, (A, B) \mapsto A \coprod B$  defines functors

$$\begin{aligned}A \coprod - &: \text{Sets} \rightarrow \text{Sets}, \\ - \coprod B &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \coprod -_2 &: \text{Sets} \times \text{Sets} \rightarrow \text{Sets},\end{aligned}$$

where  $-_1 \coprod -_2$  is the functor where

- *Action on Objects.* For each  $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$ , we have

$$[-_1 \coprod -_2](A, B) \stackrel{\text{def}}{=} A \coprod B.$$

- *Action on Morphisms.* For each  $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$ , the action on Hom-sets

$$\coprod_{(A,B),(X,Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \coprod B, X \coprod Y)$$

of  $\coprod$  at  $((A, B), (X, Y))$  is defined by sending  $(f, g)$  to the function

$$f \coprod g : A \coprod B \rightarrow X \coprod Y$$

defined by

$$[f \coprod g](x) \stackrel{\text{def}}{=} \begin{cases} (0, f(a)) & \text{if } x = (0, a), \\ (1, g(b)) & \text{if } x = (1, b), \end{cases}$$

for each  $x \in A \coprod B$ .

and where  $A \coprod -$  and  $- \coprod B$  are the partial functors of  $-_1 \coprod -_2$  at  $A, B \in \text{Obj}(\text{Sets})$ .

01ZH

2. *Adjointness.* We have an adjunction

$$(-_1 \coprod -_2 \dashv \Delta_{\text{Sets}}) : \text{Sets} \times \text{Sets} \begin{array}{c} \xrightarrow{-_1 \coprod -_2} \\ \perp \\ \xleftarrow{\Delta_{\text{Sets}}} \end{array} \text{Sets},$$

witnessed by a bijection

$$\text{Sets}(A \coprod B, C) \cong \text{Hom}_{\text{Sets} \times \text{Sets}}((A, B), (C, C))$$

natural in  $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$  and in  $C \in \text{Obj}(\text{Sets})$ .

0024

3. *Associativity.* We have an isomorphism of sets

$$\alpha_{X,Y,Z}^{\text{Sets}, \coprod} : (X \coprod Y) \coprod Z \xrightarrow{\sim} X \coprod (Y \coprod Z),$$

natural in  $X, Y, Z \in \text{Obj}(\text{Sets})$ .

0025

4. *Unitality.* We have isomorphisms of sets

$$\lambda_X^{\text{Sets}, \coprod} : \emptyset \coprod X \xrightarrow{\sim} X,$$

$$\rho_X^{\text{Sets}, \coprod} : X \coprod \emptyset \xrightarrow{\sim} X,$$

natural in  $X \in \text{Obj}(\text{Sets})$ .

0026

5. *Commutativity.* We have an isomorphism of sets

$$\sigma_{X,Y}^{\text{Sets}, \coprod} : X \coprod Y \xrightarrow{\sim} Y \coprod X,$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

0027

6. *Symmetric Monoidality.* The 7-tuple  $(\text{Sets}, \coprod, \emptyset, \alpha_{\coprod}^{\text{Sets}}, \lambda_{\coprod}^{\text{Sets}}, \rho_{\coprod}^{\text{Sets}}, \sigma^{\text{Sets}})$  is a symmetric monoidal category.

**PROOF 4.2.3.1.5 ▶ PROOF OF PROPOSITION 4.2.3.1.4**

Item 1: Functoriality

This follows from ?? of ??.

Item 2: Adjointness

This follows from the universal property of the coproduct.

Item 3: Associativity

This is proved in the proof of [Definition 5.2.3.1.1](#).

Item 4: Unitality

This is proved in the proof of [Definitions 5.2.4.1.1](#) and [5.2.5.1.1](#).

Item 5: Commutativity

This is proved in the proof of [Definition 5.2.6.1.1](#).

Item 6: Symmetric Monoidality

This is a repetition of [Proposition 5.2.7.1.1](#), and is proved there. 

#### 0028 4.2.4 Pushouts

Let  $A$ ,  $B$ , and  $C$  be sets and let  $f: C \rightarrow A$  and  $g: C \rightarrow B$  be functions.

0029

##### DEFINITION 4.2.4.1.1 ► PUSHOUTS OF SETS

The **pushout of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** <sup>1</sup> is the pushout of  $A$  and  $B$  over  $C$  along  $f$  and  $g$  in Sets as in ??.

<sup>1</sup>Further Terminology: Also called the **fibre coproduct of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** .

01EM

##### CONSTRUCTION 4.2.4.1.2 ► CONSTRUCTION OF PUSHOUTS OF SETS

Concretely, the pushout of  $A$  and  $B$  over  $C$  along  $f$  and  $g$  is the pair  $(A \coprod_C B, \{\text{inj}_1, \text{inj}_2\})$  consisting of:

01EN

1. *The Colimit.* The set  $A \coprod_C B$  defined by

$$A \coprod_C B \stackrel{\text{def}}{=} A \coprod B / \sim_C,$$

where  $\sim_C$  is the equivalence relation on  $A \coprod B$  generated by  $(0, f(c)) \sim_C (1, g(c))$ .

01EP

2. *The Cocone.* The maps

$$\text{inj}_1: A \rightarrow A \coprod_C B,$$

$$\text{inj}_2: B \rightarrow A \coprod_C B$$

given by

$$\text{inj}_1(a) \stackrel{\text{def}}{=} [(0, a)]$$

$$\text{inj}_2(b) \stackrel{\text{def}}{=} [(1, b)]$$

for each  $a \in A$  and each  $b \in B$ .

## PROOF 4.2.4.1.3 ► PROOF OF CONSTRUCTION 4.2.4.1.2

We claim that  $A \coprod_C B$  is the categorical pushout of  $A$  and  $B$  over  $C$  with respect to  $(f, g)$  in Sets. First we need to check that the relevant pushout diagram commutes, i.e. that we have

$$\begin{array}{ccc} A \coprod_C B & \xleftarrow{\text{inj}_2} & B \\ \text{inj}_1 \circ f = \text{inj}_2 \circ g, & \text{inj}_1 \uparrow & \uparrow g \\ & & A \xleftarrow{f} C. \end{array}$$

Indeed, given  $c \in C$ , we have

$$\begin{aligned} [\text{inj}_1 \circ f](c) &= \text{inj}_1(f(c)) \\ &= [(0, f(c))] \\ &= [(1, g(c))] \\ &= \text{inj}_2(g(c)) \\ &= [\text{inj}_2 \circ g](c), \end{aligned}$$

where  $[(0, f(c))] = [(1, g(c))]$  by the definition of the relation  $\sim$  on  $A \coprod B$ . Next, we prove that  $A \coprod_C B$  satisfies the universal property of the pushout. Suppose we have a diagram of the form

$$\begin{array}{ccccc} P & \xleftarrow{\quad \iota_2 \quad} & A \coprod_C B & \xleftarrow{\text{inj}_2 -} & B \\ \iota_1 \swarrow & & \uparrow \Gamma & & \uparrow g \\ A & \xleftarrow{f} & C & & \end{array}$$

in Sets. Then there exists a unique map  $\phi: A \coprod_C B \rightarrow P$  making the diagram

$$\begin{array}{ccccc}
 & & P & \leftarrow & \\
 & \nearrow \phi & \exists! & & \downarrow \iota_2 \\
 A \coprod_C B & \xleftarrow{\text{inj}_2} & B & & \\
 \uparrow \text{inj}_1 & & \uparrow g & & \\
 A & \xleftarrow{f} & C & &
 \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned}
 \phi \circ \text{inj}_1 &= \iota_1, \\
 \phi \circ \text{inj}_2 &= \iota_2
 \end{aligned}$$

via

$$\phi(x) = \begin{cases} \iota_1(a) & \text{if } x = [(0, a)], \\ \iota_2(b) & \text{if } x = [(1, b)] \end{cases}$$

for each  $x \in A \coprod_C B$ , where the well-definedness of  $\phi$  is guaranteed by the equality  $\iota_1 \circ f = \iota_2 \circ g$  and the definition of the relation  $\sim$  on  $A \coprod B$  as follows:

01EQ

1. Case 1: Suppose we have  $x = [(0, a)] = [(0, a')]$  for some  $a, a' \in A$ . Then, by Remark 4.2.4.1.4, we have a sequence

$$(0, a) \sim' x_1 \sim' \dots \sim' x_n \sim' (0, a').$$

01ER

2. Case 2: Suppose we have  $x = [(1, b)] = [(1, b')]$  for some  $b, b' \in B$ . Then, by Remark 4.2.4.1.4, we have a sequence

$$(1, b) \sim' x_1 \sim' \dots \sim' x_n \sim' (1, b').$$

01ES

3. Case 3: Suppose we have  $x = [(0, a)] = [(1, b)]$  for some  $a \in A$  and  $b \in B$ . Then, by Remark 4.2.4.1.4, we have a sequence

$$(0, a) \sim' x_1 \sim' \dots \sim' x_n \sim' (1, b).$$

In all these cases, we declare  $x \sim' y$  iff there exists some  $c \in C$  such that  $x = (0, f(c))$  and  $y = (1, g(c))$  or  $x = (1, g(c))$  and  $y = (0, f(c))$ . Then, the equality  $\iota_1 \circ f = \iota_2 \circ g$  gives

$$\phi([x]) = \phi([(0, f(c))])$$

$$\begin{aligned}
 &\stackrel{\text{def}}{=} \iota_1(f(c)) \\
 &= \iota_2(g(c)) \\
 &\stackrel{\text{def}}{=} \phi([(1, g(c))]) \\
 &= \phi([y]),
 \end{aligned}$$

with the case where  $x = (1, g(c))$  and  $y = (0, f(c))$  similarly giving  $\phi([x]) = \phi([y])$ . Thus, if  $x \sim' y$ , then  $\phi([x]) = \phi([y])$ . Applying this equality pairwise to the sequences

$$\begin{aligned}
 (0, a) &\sim' x_1 \sim' \cdots \sim' x_n \sim' (0, a'), \\
 (1, b) &\sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b'), \\
 (0, a) &\sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b)
 \end{aligned}$$

gives

$$\begin{aligned}
 \phi([(0, a)]) &= \phi([(0, a')]), \\
 \phi([(1, b)]) &= \phi([(1, b')]), \\
 \phi([(0, a)]) &= \phi([(1, b)]),
 \end{aligned}$$

showing  $\phi$  to be well-defined. □

002A

#### REMARK 4.2.4.1.4 ► UNWINDING DEFINITION 4.2.4.1.1

In detail, by [Construction 10.5.2.1.2](#), the relation  $\sim$  of [Definition 4.2.4.1.1](#) is given by declaring  $a \sim b$  iff one of the following conditions is satisfied:

01ET

1. We have  $a, b \in A$  and  $a = b$ .

01EU

2. We have  $a, b \in B$  and  $a = b$ .

01EV

3. There exist  $x_1, \dots, x_n \in A \coprod B$  such that  $a \sim' x_1 \sim' \cdots \sim' x_n \sim' b$ , where we declare  $x \sim' y$  if one of the following conditions is satisfied:

01EW

- (a) There exists  $c \in C$  such that  $x = (0, f(c))$  and  $y = (1, g(c))$ .

01EX

- (b) There exists  $c \in C$  such that  $x = (1, g(c))$  and  $y = (0, f(c))$ .

In other words, there exist  $x_1, \dots, x_n \in A \coprod B$  satisfying the following conditions:

01EY

- (c) There exists  $c_0 \in C$  satisfying one of the following conditions:

01ZJ

- i. We have  $a = f(c_0)$  and  $x_1 = g(c_0)$ .

01ZK

- ii. We have  $a = g(c_0)$  and  $x_1 = f(c_0)$ .

01EZ

(d) For each  $1 \leq i \leq n - 1$ , there exists  $c_i \in C$  satisfying one of the following conditions:

01ZL

i. We have  $x_i = f(c_i)$  and  $x_{i+1} = g(c_i)$ .

01ZM

ii. We have  $x_i = g(c_i)$  and  $x_{i+1} = f(c_i)$ .

01F0

(e) There exists  $c_n \in C$  satisfying one of the following conditions:

01F1

i. We have  $x_n = f(c_n)$  and  $b = g(c_n)$ .

01F2

ii. We have  $x_n = g(c_n)$  and  $b = f(c_n)$ .

01ZN

#### REMARK 4.2.4.1.5 ► PUSHOUTS OF SETS DEPEND ON THE MAPS

It is common practice to write  $A \coprod_C B$  for the pushout of  $A$  and  $B$  over  $C$  along  $f$  and  $g$ , omitting the maps  $f$  and  $g$  from the notation and instead leaving them implicit, to be understood from the context.

However, the set  $A \coprod_C B$  depends very much on the maps  $f$  and  $g$ , and sometimes it is necessary or useful to note this dependence explicitly. In such situations, we will write  $A \coprod_{f,C,g} B$  or  $A \coprod_C^{f,g} B$  for  $A \coprod_C B$ .

002B

#### EXAMPLE 4.2.4.1.6 ► EXAMPLES OF PUSHOUTS OF SETS

Here are some examples of pushouts of sets.

002C

1. *Wedge Sums of Pointed Sets.* The wedge sum of two pointed sets of [Definition 6.3.3.1.1](#) is an example of a pushout of sets.

002D

2. *Intersections via Unions.* Let  $X$  be a set. We have

$$\begin{array}{ccc} A \cup B & \xleftarrow{\quad} & B \\ \uparrow \lrcorner & & \uparrow \\ A \cong A \coprod_{A \cap B} B, & & A \cap B \\ \uparrow & & \downarrow \\ A & \xleftarrow{\quad} & A \cap B \end{array}$$

for each  $A, B \in \mathcal{P}(X)$ .

#### PROOF 4.2.4.1.7 ► PROOF OF EXAMPLE 4.2.4.1.6

Item 1: Wedge Sums of Pointed Sets

This follows by definition, as the wedge sum of two pointed sets is defined as a pushout.

Item 2: Intersections via Unions

Indeed,  $A \coprod_{A \cap B} B$  is the quotient of  $A \coprod B$  by the equivalence relation obtained by declaring  $(0, a) \sim (1, b)$  iff  $a = b \in A \cap B$ , which is in bijection with  $A \cup B$  via the map with  $[(0, a)] \mapsto a$  and  $[(1, b)] \mapsto b$ . 

002E

**PROPOSITION 4.2.4.1.8 ► PROPERTIES OF PUSHOOTS OF SETS**


Let  $A, B, C$ , and  $X$  be sets.

002F

- Functoriality.* The assignment  $(A, B, C, f, g) \mapsto A \coprod_{f, C, g} B$  defines a functor

$$-_1 \coprod_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where  $\mathcal{P}$  is the category that looks like this:



In particular, the action on morphisms of  $-_1 \coprod_{-3} -_1$  is given by sending a morphism

$$\begin{array}{ccccc} A \coprod_C B & \xleftarrow{\quad \Gamma \quad} & B & & \\ \uparrow & & \uparrow \psi & & \\ A' \coprod_{C'} B' & \xleftarrow{\quad \Gamma \quad} & B' & & \\ \uparrow f & \uparrow g & \uparrow g' & & \\ A & \xleftarrow{\quad \phi \quad} & C & \xrightarrow{\quad \chi \quad} & C' \\ \downarrow & & \downarrow & & \\ A' & \xleftarrow{\quad f' \quad} & & & \end{array}$$

in  $\text{Fun}(\mathcal{P}, \text{Sets})$  to the map  $\xi : A \coprod_C B \xrightarrow{\exists!} A' \coprod_{C'} B'$  given by

$$\xi(x) \stackrel{\text{def}}{=} \begin{cases} \phi(a) & \text{if } x = [(0, a)], \\ \psi(b) & \text{if } x = [(1, b)] \end{cases}$$

for each  $x \in A \coprod_C B$ , which is the unique map making the diagram

$$\begin{array}{ccccc}
 & A \coprod_C B & \xleftarrow{\quad} & B & \\
 \uparrow & \searrow & & \uparrow \psi & \\
 & A' \coprod_{C'} B' & \xleftarrow{\quad} & B' & \\
 \uparrow & \uparrow g & & \uparrow & \\
 A & \xleftarrow{\quad} & C & \xrightarrow{\quad} & \\
 \downarrow \phi & \uparrow f & \downarrow \chi & \uparrow g' & \\
 A' & \xleftarrow{\quad} & C' & \xrightarrow{\quad} &
 \end{array}$$

commute.

01ZP

2. *Adjointness.* We have an adjunction

$$\left( -_1 \coprod_{X-2} \dashv \Delta_{\text{Sets}_{X/}} \right): \text{Sets}_{X/} \times \text{Sets}_{X/} \xrightarrow[\Delta_{\text{Sets}_{X/}}]{\perp} \text{Sets}_{X/},$$

witnessed by a bijection

$$\text{Sets}_{X/}(A \coprod_X B, C) \cong \text{Hom}_{\text{Sets}_{X/} \times \text{Sets}_{X/}}((A, B), (C, C))$$

natural in  $(A, B) \in \text{Obj}(\text{Sets}_{X/} \times \text{Sets}_{X/})$  and in  $C \in \text{Obj}(\text{Sets}_{X/})$ .

002G

3. *Associativity.* Given a diagram

$$\begin{array}{ccccc}
 A & & B & & C \\
 & \swarrow f & \nearrow g & \swarrow h & \nearrow k \\
 X & & Y & &
 \end{array}$$

in  $\text{Sets}$ , we have isomorphisms of sets

$$(A \coprod_X B) \coprod_Y C \cong (A \coprod_X B) \coprod_B (B \coprod_Y C) \cong A \coprod_X (B \coprod_Y C)$$

where these pullbacks are built as in the diagrams

$$\begin{array}{ccc}
 \begin{array}{c}
 (A \coprod_X B) \coprod_Y C \\
 \uparrow \wedge \quad \searrow \\
 A \coprod_X B \quad \quad \quad C, A \\
 \uparrow \wedge \quad \uparrow \wedge \\
 A \quad B \quad \quad \quad X \quad Y \\
 \downarrow f \quad \downarrow g \quad \downarrow h \quad \downarrow k
 \end{array}
 &
 \begin{array}{c}
 (A \coprod_X B) \coprod_B (B \coprod_Y C) \\
 \uparrow \wedge \quad \uparrow \wedge \quad \searrow \\
 A \coprod_X B \quad B \coprod_Y C \quad C, A \\
 \uparrow \wedge \quad \uparrow \wedge \quad \uparrow \wedge \\
 A \quad B \quad \quad \quad X \quad Y \\
 \downarrow f \quad \downarrow g \quad \downarrow h \quad \downarrow k
 \end{array}
 &
 \begin{array}{c}
 A \coprod_X (B \coprod_Y C) \\
 \uparrow \wedge \quad \uparrow \wedge \quad \searrow \\
 B \coprod_Y C \quad C. \\
 \uparrow \wedge \quad \uparrow \wedge \\
 B \quad C. \\
 \downarrow g \quad \downarrow k
 \end{array}
 \end{array}$$

01F4

4. *Interaction With Composition.* Given a diagram

$$\begin{array}{ccc}
 X & & Y \\
 \swarrow \phi & & \searrow \psi \\
 A & & B \\
 \downarrow f & \nearrow g & \\
 K & & 
 \end{array}$$

in Sets, we have isomorphisms of sets


$$\begin{aligned}
 X \coprod_K^{\phi \circ f, \psi \circ g} Y &\cong (X \coprod_A^{\phi, j_1} (A \coprod_K^{f, g} B)) \coprod_{A \coprod_K^{f, g} B}^{i_2, i_1} ((A \coprod_K^{f, g} B) \coprod_B^{j_2, \psi} Y) \\
 &\cong X \coprod_A^{\phi, i} ((A \coprod_K^{f, g} B) \coprod_B^{j_2, \psi} Y) \\
 &\cong (X \coprod_A^{\phi, i_1} (A \coprod_K^{f, g} B)) \coprod_B^{j, \psi} Y
 \end{aligned}$$

where

$$\begin{aligned}
 j_1 &= \text{inj}_1^{A \times_K^{f, g} B}, & j_2 &= \text{inj}_2^{A \times_K^{f, g} B}, \\
 i_1 &= \text{inj}_1^{(A \times_K^{f, g} B) \times_Y^{q_2, \psi}}, & i_2 &= \text{inj}_2^{X \times_{A \times_K^{f, g} B}^{\phi, q_1} (A \times_K^{f, g} B)}, \\
 i &= j_1 \circ \text{inj}_1^{(A \times_K^{f, g} B) \times_B^{q_2, \psi} Y}, & j &= j_2 \circ \text{inj}_2^{X \times_A^{\phi, q_1} (A \times_K^{f, g} B)},
 \end{aligned}$$

and where these pullbacks are built as in the diagrams

$$\begin{array}{ccc}
 (X \coprod_A (A \coprod_K B)) \coprod_{A \coprod_K B} ((A \coprod_K B) \coprod_B Y) & & X \coprod_A ((A \coprod_K B) \coprod_B Y) \\
 \uparrow \wedge \quad \uparrow \wedge & & \uparrow \wedge \\
 X \coprod_A (A \coprod_K B) & (A \coprod_K B) \coprod_B Y & (A \coprod_K B) \coprod_B Y \\
 \uparrow \wedge \quad \uparrow \wedge & & \uparrow \wedge \\
 X & A \coprod_K B & X \\
 \downarrow \phi & \uparrow \wedge & \downarrow \phi \\
 A & B & A \\
 \downarrow f & \uparrow \wedge & \downarrow f \\
 K & & K
 \end{array}$$



002H

5. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc}
 A & \xlongequal{\quad} & A \\
 \uparrow f & & \uparrow f \\
 X & \xlongequal{\quad} & X
 \end{array}
 \quad
 \begin{array}{c}
 \lambda_A^{\text{Sets}_{X/}} : X \coprod_X A \xrightarrow{\sim} A, \\
 \rho_A^{\text{Sets}_{X/}} : A \coprod_X X \xrightarrow{\sim} A,
 \end{array}
 \quad
 \begin{array}{ccc}
 A & \xleftarrow{f} & X \\
 \parallel & & \parallel \\
 X & \xleftarrow{f} & X,
 \end{array}$$

natural in  $(A, f) \in \text{Obj}(\text{Sets}_{X/})$ .

002J

6. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc}
 A \coprod_X B & \longleftarrow & B \\
 \uparrow \lrcorner & & \uparrow g \\
 A & \xleftarrow{f} & X, \\
 & & \uparrow \sigma_A^{\text{Sets}_{X/}} : A \coprod_X B \xrightarrow{\sim} B \coprod_X A
 \end{array}
 \quad
 \begin{array}{ccc}
 B \coprod_X A & \longleftarrow & A \\
 \uparrow \lrcorner & & \uparrow f \\
 B & \xleftarrow{g} & X.
 \end{array}$$

natural in  $(A, f), (B, g) \in \text{Obj}(\text{Sets}_{X/})$ .

002K

7. *Interaction With Coproducts.* We have

$$\begin{array}{ccc}
 A \coprod B & \longleftarrow & B \\
 \uparrow \lrcorner & & \uparrow \iota_B \\
 A \coprod_{\emptyset} B \cong A \coprod B, & & \uparrow \iota_A \\
 & & \uparrow \lrcorner
 \end{array}$$

002L

8. *Symmetric Monoidality.* The triple  $(\text{Sets}_{X/}, \coprod_X, X)$  is a symmetric monoidal category.

**PROOF 4.2.4.1.9 ▶ PROOF OF PROPOSITION 4.2.4.1.8****Item 1: Functoriality**

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for  $\xi$  following from the commutativity of the cube pushout diagram.

**Item 2: : Adjointness**

This follows from the universal property of the coproduct (pushouts are coproducts in  $\text{Sets}_{X/}$ ).

**Item 3: Associativity**

Omitted.

**Item 4: Interaction With Composition**

Omitted.

**Item 5: Unitality**

Omitted.

**Item 6: Commutativity**

Omitted.

**Item 7: Interaction With Coproducts**

Omitted.

**Item 8: Symmetric Monoidality**

Omitted.

**002M 4.2.5 Coequalisers**

Let  $A$  and  $B$  be sets and let  $f, g: A \rightrightarrows B$  be functions.

**002N****DEFINITION 4.2.5.1.1 ▶ COEQUALISERS OF SETS**

The **coequaliser of  $f$  and  $g$**  is the coequaliser of  $f$  and  $g$  in Sets as in ??.

**01F5****CONSTRUCTION 4.2.5.1.2 ▶ CONSTRUCTION OF COEQUALISERS OF SETS**

Concretely, the coequaliser of  $f$  and  $g$  is the pair  $(\text{CoEq}(f, g), \text{coeq}(f, g))$  consisting of:

**01F6**

- 1. The Colimit.* The set  $\text{CoEq}(f, g)$  defined by

$$\text{CoEq}(f, g) \stackrel{\text{def}}{=} B/\sim,$$

01F7

where  $\sim$  is the equivalence relation on  $B$  generated by  $f(a) \sim g(a)$ .

2. *The Cocone.* The map

$$\text{coeq}(f, g) : B \twoheadrightarrow \text{CoEq}(f, g)$$

given by the quotient map  $\pi : B \twoheadrightarrow B/\sim$  with respect to the equivalence relation generated by  $f(a) \sim g(a)$ .

#### PROOF 4.2.5.1.3 ► PROOF OF CONSTRUCTION 4.2.5.1.2

We claim that  $\text{CoEq}(f, g)$  is the categorical coequaliser of  $f$  and  $g$  in Sets. First we need to check that the relevant coequaliser diagram commutes, i.e. that we have

$$\text{coeq}(f, g) \circ f = \text{coeq}(f, g) \circ g.$$

Indeed, we have

$$\begin{aligned} [\text{coeq}(f, g) \circ f](a) &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](f(a)) \\ &\stackrel{\text{def}}{=} [f(a)] \\ &= [g(a)] \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](g(a)) \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g) \circ g](a) \end{aligned}$$

for each  $a \in A$ . Next, we prove that  $\text{CoEq}(f, g)$  satisfies the universal property of the coequaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} A & \xrightleftharpoons[\substack{f \\ g}]{\quad} & B & \xrightarrow{\text{coeq}(f, g)} & \text{CoEq}(f, g) \\ & & & \searrow c & \\ & & & & C \end{array}$$

in Sets. Then, since  $c(f(a)) = c(g(a))$  for each  $a \in A$ , it follows from **Items 4 and 5 of Proposition 10.6.2.1.3** that there exists a unique map  $\text{CoEq}(f, g) \xrightarrow{\exists!} C$  making the diagram

$$\begin{array}{ccc} A & \xrightleftharpoons[\substack{f \\ g}]{\quad} & B & \xrightarrow{\text{coeq}(f, g)} & \text{CoEq}(f, g) \\ & & & \searrow c & \downarrow \exists! \\ & & & & C \end{array}$$

commute. □

002P

**REMARK 4.2.5.1.4 ► UNWINDING DEFINITION 4.2.5.1.1**

In detail, by [Construction 10.5.2.1.2](#), the relation  $\sim$  of [Definition 4.2.5.1.1](#) is given by declaring  $a \sim b$  iff one of the following conditions is satisfied:

01ZQ

1. We have  $a = b$ ;

01ZR

2. There exist  $x_1, \dots, x_n \in B$  such that  $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$ , where we declare  $x \sim' y$  if one of the following conditions is satisfied:

01ZS

- (a) There exists  $z \in A$  such that  $x = f(z)$  and  $y = g(z)$ .

01ZT

- (b) There exists  $z \in A$  such that  $x = g(z)$  and  $y = f(z)$ .

In other words, there exist  $x_1, \dots, x_n \in B$  satisfying the following conditions:

01ZU

- (a) There exists  $z_0 \in A$  satisfying one of the following conditions:

01ZV

- i. We have  $a = f(z_0)$  and  $x_1 = g(z_0)$ .

01ZW

- ii. We have  $a = g(z_0)$  and  $x_1 = f(z_0)$ .

01ZX

- (b) For each  $1 \leq i \leq n - 1$ , there exists  $z_i \in A$  satisfying one of the following conditions:

01ZY

- i. We have  $x_i = f(z_i)$  and  $x_{i+1} = g(z_i)$ .

01ZZ

- ii. We have  $x_i = g(z_i)$  and  $x_{i+1} = f(z_i)$ .

0200

- (c) There exists  $z_n \in A$  satisfying one of the following conditions:

0201

- i. We have  $x_n = f(z_n)$  and  $b = g(z_n)$ .

0202

- ii. We have  $x_n = g(z_n)$  and  $b = f(z_n)$ .

002Q

**EXAMPLE 4.2.5.1.5 ► EXAMPLES OF COEQUALISERS OF SETS**

Here are some examples of coequalisers of sets.

002R

1. *Quotients by Equivalence Relations.* Let  $R$  be an equivalence relation on a set  $X$ . We have a bijection of sets

$$X/\sim_R \cong \text{CoEq}(R \hookrightarrow X \times X \xrightarrow{\begin{smallmatrix} \text{pr}_1 \\ \text{pr}_2 \end{smallmatrix}} X).$$

**PROOF 4.2.5.1.6 ► PROOF OF EXAMPLE 4.2.5.1.5**

Item 1: Quotients by Equivalence Relations

See [Pro25ad].



002S

**PROPOSITION 4.2.5.1.7 ► PROPERTIES OF COEQUALISERS OF SETS**

Let  $A$ ,  $B$ , and  $C$  be sets.

002T

1. *Associativity.* We have isomorphisms of sets<sup>1</sup>

$$\begin{aligned} \text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h) &\cong \text{CoEq}(f, g, h) \cong \underbrace{\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g)}, \\ &= \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h) \\ &= \text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h) \end{aligned}$$

where  $\text{CoEq}(f, g, h)$  is the colimit of the diagram

$$A \xrightarrow[\substack{f \\ g \\ h}]{} B$$

in Sets.

002U

2. *Unitality.* We have an isomorphism of sets

$$\text{CoEq}(f, f) \cong B.$$

002V

3. *Commutativity.* We have an isomorphism of sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

002W

4. *Interaction With Composition.* Let

$$A \xrightarrow[\substack{f \\ g}]{} B \xrightarrow[\substack{h \\ k}]{} C$$

be functions. We have a surjection

$$\text{CoEq}(h \circ f, k \circ g) \twoheadrightarrow \text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$$

exhibiting  $\text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$  as a quotient of  $\text{CoEq}(h \circ f, k \circ g)$  by the relation generated by declaring  $h(y) \sim k(y)$  for each  $y \in B$ .

0203

---

<sup>1</sup>That is, the following three ways of forming “the” coequaliser of  $(f, g, h)$  agree:

- (a) Take the coequaliser of  $(f, g, h)$ , i.e. the colimit of the diagram

$$A \xrightarrow[\substack{f \\ g \\ h}]{} B$$

in Sets.

0204

(b) First take the coequaliser of  $f$  and  $g$ , forming a diagram

$$A \xrightarrow[\underset{g}{\Rightarrow}]{} B \xrightarrow{\text{coeq}(f,g)} \text{CoEq}(f,g)$$

and then take the coequaliser of the composition

$$A \xrightarrow[\underset{h}{\Rightarrow}]{} B \xrightarrow{\text{coeq}(f,g)} \text{CoEq}(f,g),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(f,g) \circ f, \text{coeq}(f,g) \circ h) = \text{CoEq}(\text{coeq}(f,g) \circ g, \text{coeq}(f,g) \circ h)$$

of  $\text{CoEq}(f,g)$

0205

(c) First take the coequaliser of  $g$  and  $h$ , forming a diagram

$$A \xrightarrow[\underset{h}{\Rightarrow}]{} B \xrightarrow{\text{coeq}(g,h)} \text{CoEq}(g,h)$$

and then take the coequaliser of the composition

$$A \xrightarrow[\underset{g}{\Rightarrow}]{} B \xrightarrow{\text{coeq}(g,h)} \text{CoEq}(g,h),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(g,h) \circ f, \text{coeq}(g,h) \circ g) = \text{CoEq}(\text{coeq}(g,h) \circ f, \text{coeq}(g,h) \circ h)$$

of  $\text{CoEq}(g,h)$ .

#### PROOF 4.2.5.1.8 ► PROOF OF PROPOSITION 4.2.5.1.7

Item 1: Associativity

Omitted.

Item 2: Unitality

Omitted.

Item 3: Commutativity

Omitted.

Item 4: Interaction With Composition

Omitted.



#### 01F8 4.2.6 Direct Colimits

Let  $(X_\alpha, f_{\alpha\beta})_{\alpha,\beta \in I}: (I, \preceq) \rightarrow \mathbb{P}$  be a direct system of sets.

01F9

#### DEFINITION 4.2.6.1.1 ► DIRECT COLIMITS OF SETS

The **direct colimit** of  $(X_\alpha, f_{\alpha\beta})_{\alpha,\beta \in I}$  is the direct colimit of  $(X_\alpha, f_{\alpha\beta})_{\alpha,\beta \in I}$  in Sets as in ??.

01FA

**CONSTRUCTION 4.2.6.1.2 ► CONSTRUCTION OF DIRECT COLIMITS OF SETS**

Concretely, the direct colimit of  $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$  is the pair  $\left( \underset{\alpha \in I}{\text{colim}}(X_\alpha), \{ \text{inj}_\alpha \}_{\alpha \in I} \right)$  consisting of:

01FB

- The Colimit.* The set  $\underset{\alpha \in I}{\text{colim}}(X_\alpha)$  defined by

$$\underset{\alpha \in I}{\text{colim}}(X_\alpha) \stackrel{\text{def}}{=} \left( \bigsqcup_{\alpha \in I} X_\alpha \right) / \sim,$$

where  $\sim$  is the equivalence relation on  $\bigsqcup_{\alpha \in I} X_\alpha$  generated by declaring  $(\alpha, x) \sim (\beta, y)$  iff there exists some  $\gamma \in I$  satisfying the following conditions:

01FC

- (a) We have  $\alpha \preceq \gamma$ .
- (b) We have  $\beta \preceq \gamma$ .
- (c) We have  $f_{\alpha\gamma}(x) = f_{\beta\gamma}(y)$ .

01FF

- The Cocone.* The collection

$$\left\{ \text{inj}_\gamma : X_\gamma \rightarrow \underset{\alpha \in I}{\text{colim}}(X_\alpha) \right\}_{\gamma \in I}$$

of maps of sets defined by

$$\text{inj}_\gamma(x) \stackrel{\text{def}}{=} [(\gamma, x)]$$

for each  $\gamma \in I$  and each  $x \in X_\gamma$ .

**PROOF 4.2.6.1.3 ► PROOF OF CONSTRUCTION 4.2.6.1.2**

We will prove **Construction 4.2.6.1.2** below in a bit, but first we need a lemma (which is interesting in its own right). 

01FG

**LEMMA 4.2.6.1.4 ► IDENTIFICATION OF  $x$  WITH  $f_{\alpha\beta}(x)$  IN DIRECT COLIMITS**

For each  $\alpha, \beta \in I$  and each  $x \in X_\alpha$ , if  $\alpha \preceq \beta$ , then we have

$$(\alpha, x) \sim (\beta, f_{\alpha\beta}(x))$$

in  $\underset{\alpha \in I}{\text{colim}}(X_\alpha)$ .

**PROOF 4.2.6.1.5 ▶ PROOF OF LEMMA 4.2.6.1.4**

Taking  $\gamma = \beta$ , we have  $f_{\alpha\gamma} = f_{\alpha\beta}$ , we have  $f_{\beta\gamma} = f_{\beta\beta} \stackrel{\text{def}}{=} \text{id}_{X_\beta}$ , and we have

$$\begin{aligned} f_{\alpha\beta}(x) &= f_{\beta\beta}(f_{\alpha\beta}(x)) \\ &\stackrel{\text{def}}{=} \text{id}_{X_\beta}(f_{\alpha\beta}(x)), \\ &= f_{\alpha\beta}(x). \end{aligned}$$

As a result, since  $\alpha \preceq \beta$  and  $\beta \preceq \beta$  as well, **Items 1a to 1c of Construction 4.2.6.1.2** are met. Thus we have  $(\alpha, x) \sim (\beta, f_{\alpha\beta}(x))$ . ■

We can now prove **Construction 4.2.6.1.2**:

**PROOF 4.2.6.1.6 ▶ PROOF OF CONSTRUCTION 4.2.6.1.2**

We claim that  $\underset{\longrightarrow}{\text{colim}}_{\alpha \in I} (X_\alpha)$  is the colimit of the direct system of sets  $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ .

**Commutativity of the Colimit Diagram**

First, we need to check that the colimit diagram defined by  $\underset{\longrightarrow}{\text{colim}}_{\alpha \in I} (X_\alpha)$  commutes, i.e. that we have

$$\begin{array}{ccc} & \text{colim}(X_\alpha) & \\ & \xrightarrow{\alpha \in I} & \\ \text{inj}_\alpha = \text{inj}_\beta \circ f_{\alpha\beta}, & \text{inj}_\alpha \nearrow & \nwarrow \text{inj}_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$

for each  $\alpha, \beta \in I$  with  $\alpha \preceq \beta$ . Indeed, given  $x \in X_\alpha$ , we have

$$\begin{aligned} [\text{inj}_\beta \circ f_{\alpha\beta}](x) &\stackrel{\text{def}}{=} \text{inj}_\beta(f_{\alpha\beta}(x)) \\ &\stackrel{\text{def}}{=} [(\beta, f_{\alpha\beta}(x))] \\ &= [(\alpha, x)] \\ &\stackrel{\text{def}}{=} \text{inj}_\alpha(x), \end{aligned}$$

where we have used **Lemma 4.2.6.1.4** for the third equality.

**Proof of the Universal Property of the Colimit**

Next, we prove that  $\underset{\longrightarrow}{\text{colim}}_{\alpha \in I} (X_\alpha)$  as constructed in **Construction 4.2.6.1.2** satisfies the universal property of a direct colimit. Suppose

that we have, for each  $\alpha, \beta \in I$  with  $\alpha \preceq \beta$ , a diagram of the form

$$\begin{array}{ccc} & C & \\ i_\alpha \nearrow & \text{colim}(X_\alpha) & \swarrow i_\beta \\ & \xrightarrow[\alpha \in I]{} & \\ \text{inj}_\alpha \nearrow & & \swarrow \text{inj}_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$

in Sets. We claim that there exists a unique map  $\phi: \text{colim}(X_\alpha) \xrightarrow[\alpha \in I]{} C$  making the diagram

$$\begin{array}{ccc} & C & \\ i_\alpha \nearrow & \phi \downarrow \exists! & \swarrow i_\beta \\ & \text{colim}(X_\alpha) & \\ \text{inj}_\alpha \nearrow & & \swarrow \text{inj}_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$

commute. To this end, first consider the diagram

$$\begin{array}{ccc} \coprod_{\alpha \in I} X_\alpha & \xrightarrow{\text{pr}} & \text{colim}(X_\alpha) \\ & \searrow & \\ & \coprod_{\alpha \in I} i_\alpha & \\ & & C. \end{array}$$

**Lemma.** If  $(\alpha, x) \sim (\beta, y)$ , then we have

$$\left[ \coprod_{\alpha \in I} i_\alpha \right](x) = \left[ \coprod_{\alpha \in I} i_\alpha \right](y).$$

*Proof.* Indeed, if  $(\alpha, x) \sim (\beta, y)$ , then there exists some  $\gamma \in I$  satisfying the following conditions:

- 0206 1. We have  $\alpha \preceq \gamma$ .
- 0207 2. We have  $\beta \preceq \gamma$ .
- 0208 3. We have  $f_{\alpha\gamma}(x) = f_{\beta\gamma}(y)$ .

We then have

$$\begin{aligned} \left[ \coprod_{\alpha \in I} i_\alpha \right] (x) &\stackrel{\text{def}}{=} i_\alpha(x) \\ &\stackrel{\text{def}}{=} [i_\gamma \circ f_{\alpha\gamma}](x) \\ &\stackrel{\text{def}}{=} i_\gamma(f_{\alpha\gamma}(x)) \\ &= i_\gamma(f_{\beta\gamma}(x)) \\ &\stackrel{\text{def}}{=} [i_\gamma \circ f_{\beta\gamma}](x) \\ &= i_\beta(y) \\ &\stackrel{\text{def}}{=} \left[ \coprod_{\alpha \in I} i_\alpha \right] (y). \end{aligned}$$

This finishes the proof of the lemma. Continuing, by ?? of **Proposition 10.6.2.1.3**, there then exists a map  $\phi: \underset{\alpha \in I}{\operatorname{colim}}(X_\alpha) \xrightarrow{\exists!} C$  making the diagram

$$\begin{array}{ccc} \coprod_{\alpha \in I} X_\alpha & \xrightarrow{\text{pr}} & \underset{\alpha \in I}{\operatorname{colim}}(X_\alpha) \\ & \searrow & \downarrow \phi \\ \coprod_{\alpha \in I} i_\alpha & \rightarrow & C \end{array}$$

commute. In particular, this implies that the diagram

$$\begin{array}{ccc} X_\alpha & \xrightarrow{\text{inj}_\alpha} & \underset{\alpha \in I}{\operatorname{colim}}(X_\alpha) \\ & \searrow & \downarrow \phi \\ & i_\alpha & \rightarrow C \end{array}$$

also commutes, and thus so does the diagram

$$\begin{array}{ccc}
 & C & \\
 i_\alpha \nearrow & \uparrow \phi \exists! & \swarrow i_\beta \\
 \text{colim}_{\alpha \in I} (X_\alpha) & \xrightarrow{\quad} & \\
 \downarrow \text{inj}_\alpha & & \uparrow \text{inj}_\beta \\
 X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta.
 \end{array}$$

This finishes the proof.<sup>1</sup>

<sup>1</sup>Incidentally, the conditions

$$\{i_\alpha = \phi \circ \text{inj}_\alpha\}_{\alpha \in I}$$

show that  $\phi$  must be given by

$$\phi([(x)]) = (i_\alpha(x))_{\alpha \in I}$$

for each  $[(x)] \in \text{colim}_{\alpha \in I} (X_\alpha)$ , although we would need to show that this assignment is well-defined were we to prove [Construction 4.2.6.1.2](#) in this way. Instead, invoking ?? of [Proposition 10.6.2.1.3](#) gave us a way to avoid having to prove this, leading to a cleaner alternative proof.

01FH

#### EXAMPLE 4.2.6.1.7 ► EXAMPLES OF DIRECT COLIMITS OF SETS

Here are some examples of direct colimits of sets.

01FJ

1. *The Prüfer Group.* The Prüfer group  $\mathbb{Z}(p^\infty)$  is defined as the direct colimit

$$\mathbb{Z}(p^\infty) \stackrel{\text{def}}{=} \text{colim}_{n \in \mathbb{N}} (\mathbb{Z}/p^n);$$

see ??.

**002X 4.3 Operations With Sets****002Y 4.3.1 The Empty Set**

002Z

**DEFINITION 4.3.1.1 ► THE EMPTY SET**

The **empty set** is the set  $\emptyset$  defined by

$$\emptyset \stackrel{\text{def}}{=} \{x \in X \mid x \neq x\},$$

where  $X$  is the set in the set existence axiom, ?? of ??.

**0030 4.3.2 Singleton Sets**

Let  $X$  be a set.

0031

**DEFINITION 4.3.2.1.1 ► SINGLETON SETS**

The **singleton set containing  $X$**  is the set  $\{X\}$  defined by

$$\{X\} \stackrel{\text{def}}{=} \{X, X\},$$

where  $\{X, X\}$  is the pairing of  $X$  with itself of [Definition 4.3.3.1.1](#).

**0032 4.3.3 Pairings of Sets**

Let  $X$  and  $Y$  be sets.

0033

**DEFINITION 4.3.3.1.1 ► PAIRINGS OF SETS**

The **pairing of  $X$  and  $Y$**  is the set  $\{X, Y\}$  defined by

$$\{X, Y\} \stackrel{\text{def}}{=} \{x \in A \mid x = X \text{ or } x = Y\},$$

where  $A$  is the set in the axiom of pairing, ?? of ??.

**0034 4.3.4 Ordered Pairs**

Let  $A$  and  $B$  be sets.

0035

**DEFINITION 4.3.4.1.1 ► ORDERED PAIRS**

The **ordered pair associated to  $A$  and  $B$**  is the set  $(A, B)$  defined by

$$(A, B) \stackrel{\text{def}}{=} \{\{A\}, \{A, B\}\}.$$

0036

**PROPOSITION 4.3.4.1.2 ► PROPERTIES OF ORDERED PAIRS**

Let  $A$  and  $B$  be sets.

0037

1. *Uniqueness.* Let  $A, B, C$ , and  $D$  be sets. The following conditions are equivalent:

0038

(a) We have  $(A, B) = (C, D)$ .

0039

(b) We have  $A = C$  and  $B = D$ .

**PROOF 4.3.4.1.3 ► PROOF OF PROPOSITION 4.3.4.1.2**

Item 1: Uniqueness

See [Cie97, Theorem 1.2.3].

**003A 4.3.5 Sets of Maps**

Let  $A$  and  $B$  be sets.

003B

**DEFINITION 4.3.5.1.1 ► SETS OF MAPS**

The **set of maps from  $A$  to  $B$** <sup>1</sup> is the set  $\text{Sets}(A, B)$ <sup>2</sup> whose elements are the functions from  $A$  to  $B$ .

<sup>1</sup>Further Terminology: Also called the **Hom set from  $A$  to  $B$** .

<sup>2</sup>Further Notation: Also written  $\text{Hom}_{\text{Sets}}(A, B)$ .

003C

**PROPOSITION 4.3.5.1.2 ► PROPERTIES OF SETS OF MAPS**

Let  $A$  and  $B$  be sets.

003D

1. *Functionality.* The assignments  $X, Y, (X, Y) \mapsto \text{Hom}_{\text{Sets}}(X, Y)$  define functors

$$\begin{aligned} \text{Sets}(X, -) : \quad \text{Sets} &\rightarrow \text{Sets}, \\ \text{Sets}(-, Y) : \quad \text{Sets}^{\text{op}} &\rightarrow \text{Sets}, \\ \text{Sets}(-_1, -_2) : \text{Sets}^{\text{op}} \times \text{Sets} &\rightarrow \text{Sets}. \end{aligned}$$

01FK

2. *Adjointness.* We have adjunctions

$$\begin{aligned} (A \times - \dashv \text{Sets}(A, -)) : \quad \text{Sets} &\begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Sets}(A, -)} \end{array} \text{Sets}, \\ (- \times B \dashv \text{Sets}(B, -)) : \quad \text{Sets} &\begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Sets}(B, -)} \end{array} \text{Sets}, \end{aligned}$$

witnessed by bijections

$$\begin{aligned}\text{Sets}(A \times B, C) &\cong \text{Sets}(A, \text{Sets}(B, C)), \\ \text{Sets}(A \times B, C) &\cong \text{Sets}(B, \text{Sets}(A, C)),\end{aligned}$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

**01FL** 3. *Maps From the Punctual Set.* We have a bijection

$$\text{Sets}(\text{pt}, A) \cong A,$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

**01FM** 4. *Maps to the Punctual Set.* We have a bijection

$$\text{Sets}(A, \text{pt}) \cong \text{pt},$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

#### PROOF 4.3.5.1.3 ► PROOF OF PROPOSITION 4.3.5.1.2

Item 1: Functoriality

This follows from **Items 2 and 5** of **Proposition 11.1.4.1.2**.

Item 2: Adjointness

This is a repetition of **Item 2** of **Proposition 4.1.3.1.4** and is proved there.

Item 3: Maps From the Punctual Set

The bijection

$$\Phi_A : \text{Sets}(\text{pt}, A) \xrightarrow{\sim} A$$

is given by

$$\Phi_A(f) \stackrel{\text{def}}{=} f(\star)$$

for each  $f \in \text{Sets}(\text{pt}, A)$ , admitting an inverse

$$\Phi_A^{-1} : A \xrightarrow{\sim} \text{Sets}(\text{pt}, A)$$

given by

$$\Phi_A^{-1}(a) \stackrel{\text{def}}{=} [\star \mapsto a]$$

for each  $a \in A$ . Indeed, we have

$$\begin{aligned}[\Phi_A^{-1} \circ \Phi_A](f) &\stackrel{\text{def}}{=} \Phi_A^{-1}(\Phi_A(f)) \\ &\stackrel{\text{def}}{=} \Phi_A^{-1}(f(\star))\end{aligned}$$

$$\begin{aligned} &\stackrel{\text{def}}{=} [\star \mapsto f(\star)] \\ &\stackrel{\text{def}}{=} f \\ &\stackrel{\text{def}}{=} [\text{id}_{\text{Sets}(\text{pt}, A)}](f) \end{aligned}$$

for each  $f \in \text{Sets}(\text{pt}, A)$  and

$$\begin{aligned} [\Phi_A \circ \Phi_A^{-1}](a) &\stackrel{\text{def}}{=} \Phi_A(\Phi_A^{-1}(a)) \\ &\stackrel{\text{def}}{=} \Phi_A([\star \mapsto a]) \\ &\stackrel{\text{def}}{=} \text{ev}_\star([\star \mapsto a]) \\ &\stackrel{\text{def}}{=} a \\ &\stackrel{\text{def}}{=} [\text{id}_A](a) \end{aligned}$$

for each  $a \in A$ , and thus we have

$$\begin{aligned} \Phi_A^{-1} \circ \Phi_A &= \text{id}_{\text{Sets}(\text{pt}, A)} \\ \Phi_A \circ \Phi_A^{-1} &= \text{id}_A. \end{aligned}$$

To prove naturality, we need to show that the diagram

$$\begin{array}{ccc} \text{Sets}(\text{pt}, A) & \xrightarrow{f_!} & \text{Sets}(\text{pt}, B) \\ \Phi_A \downarrow & & \downarrow \Phi_B \\ A & \xrightarrow{f} & B \end{array}$$

commutes. Indeed, we have

$$\begin{aligned} [f \circ \Phi_A](\phi) &\stackrel{\text{def}}{=} f(\Phi_A(\phi)) \\ &\stackrel{\text{def}}{=} f(\phi(\star)) \\ &\stackrel{\text{def}}{=} [f \circ \phi](\star) \\ &\stackrel{\text{def}}{=} \Phi_B(f \circ \phi) \\ &\stackrel{\text{def}}{=} \Phi_B(f_!(\phi)) \\ &\stackrel{\text{def}}{=} [\Phi_B \circ f_!](\phi) \end{aligned}$$

for each  $\phi \in \text{Sets}(\text{pt}, A)$ . This finishes the proof.

**Item 4: Maps to the Punctual Set**

This follows from the universal property of pt as the terminal set, [Definition 4.1.1.1.1](#).

**003E 4.3.6 Unions of Families of Subsets**

Let  $X$  be a set and let  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

003F

**DEFINITION 4.3.6.1.1 ► UNIONS OF FAMILIES OF SUBSETS**

The **union** of  $\mathcal{U}$  is the set  $\bigcup_{U \in \mathcal{U}} U$  defined by

$$\bigcup_{U \in \mathcal{U}} U \stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } U \in \mathcal{U} \\ \text{such that we have } x \in U \end{array} \right\}.$$

01FN

**PROPOSITION 4.3.6.1.2 ► PROPERTIES OF UNIONS OF FAMILIES OF SUBSETS**

Let  $X$  be a set.

01FP

- Functoriality.* The assignment  $\mathcal{U} \mapsto \bigcup_{U \in \mathcal{U}} U$  defines a functor

$$\bigcup: (\mathcal{P}(\mathcal{P}(X)), \subset) \rightarrow (\mathcal{P}(X), \subset).$$

In particular, for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ , the following condition is satisfied:

$$(\star) \text{ If } \mathcal{U} \subset \mathcal{V}, \text{ then } \bigcup_{U \in \mathcal{U}} U \subset \bigcup_{V \in \mathcal{V}} V.$$

01FQ

- Associativity.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(\mathcal{P}(X))) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \bigcup} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \bigcup \star \text{id}_{\mathcal{P}(X)} & & \downarrow \bigcup \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow[\bigcup]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \bigcup_{A \in \mathcal{A}} A} U = \bigcup_{A \in \mathcal{A}} (\bigcup_{U \in A} U)$$

for each  $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$ .

01FR

- Left Unitality.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \downarrow \chi_{\mathcal{P}(X)} & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow[\bigcup]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{V \in \{U\}} V = U$$

for each  $U \in \mathcal{P}(X)$ .

01FS

4. *Right Unitality.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \downarrow \mathcal{P}(\chi_X) & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{\{u\} \in \chi_X(U)} \{u\} = U$$

for each  $U \in \mathcal{P}(X)$ .

01FT

5. *Interaction With Unions I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup \times \cup & & \downarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\cup]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{W \in \mathcal{U} \cup \mathcal{V}} W = \left( \bigcup_{U \in \mathcal{U}} U \right) \cup \left( \bigcup_{V \in \mathcal{V}} V \right)$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01FU

6. *Interaction With Unions II.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \setminus \{\emptyset\} & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (\cup \cup -)} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow[\cup \cup -]{} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \setminus \{\emptyset\} & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cup V)} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow[- \cup V]{} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$U \cup \left( \bigcup_{V \in \mathcal{V}} V \right) = \bigcup_{V \in \mathcal{V}} (U \cup V),$$

$$\left(\bigcup_{U \in \mathcal{U}} U\right) \cup V = \bigcup_{U \in \mathcal{U}} (U \cup V)$$

for each nonempty  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$  and each  $U, V \in \mathcal{P}(X)$ .

**01FV**

7. *Interaction With Intersections I.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \times \cup \downarrow & \curvearrowright & \downarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X), \end{array}$$

with components

$$\bigcup_{w \in \mathcal{U} \cap \mathcal{V}} w \subset \left( \bigcup_{U \in \mathcal{U}} U \right) \cap \left( \bigcup_{V \in \mathcal{V}} V \right)$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

**01FW**

8. *Interaction With Intersections II.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cap -)} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow[U \cap -]{} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cap V)} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow[- \cap V]{} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$\begin{aligned} U \cap \left( \bigcup_{V \in \mathcal{V}} V \right) &= \bigcup_{V \in \mathcal{V}} (U \cap V), \\ \left( \bigcup_{U \in \mathcal{U}} U \right) \cap V &= \bigcup_{U \in \mathcal{U}} (U \cap V) \end{aligned}$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$  and each  $U, V \in \mathcal{P}(X)$ .

**01FX**

9. *Interaction With Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\setminus} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \times \cup \downarrow & \text{X} & \downarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\setminus]{} & \mathcal{P}(X), \end{array}$$

*does not commute* in general, i.e. we may have

$$\bigcup_{W \in \mathcal{U} \setminus \mathcal{V}} W \neq \left( \bigcup_{U \in \mathcal{U}} U \right) \setminus \left( \bigcup_{V \in \mathcal{V}} V \right)$$

in general, where  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01FY

10. *Interaction With Complements I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(\mathcal{P}(X)) \\ \cup^{\text{op}} \downarrow & \times & \downarrow \cup \\ \mathcal{P}(X)^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(X), \end{array}$$

*does not commute* in general, i.e. we may have

$$\bigcup_{U \in \mathcal{U}^c} U \neq \bigcup_{U \in \mathcal{U}} U^c$$

in general, where  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

01FZ

11. *Interaction With Complements II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & & \searrow \sim \\ \mathcal{P}(\mathcal{P}(X)) & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\ \cup \searrow & & \swarrow \cap^{\text{op}} \\ \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}}, \end{array}$$

commutes, i.e. we have

$$\left( \bigcup_{U \in \mathcal{U}} U \right)^c = \bigcap_{U \in \mathcal{U}} U^c$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

01G0

12. *Interaction With Complements III.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & \\
 \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & \searrow \text{dashed} & \\
 \mathcal{P}(\mathcal{P}(X)) & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\
 \downarrow \cap & & \downarrow \cup^{\text{op}} \\
 \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}},
 \end{array}$$

commutes, i.e. we have

$$\left( \bigcap_{U \in \mathcal{U}} U \right)^c = \bigcup_{U \in \mathcal{U}} U^c$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

01G1

13. *Interaction With Symmetric Differences.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\Delta} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cup \times \cup & \text{X} & \downarrow \cup \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\Delta]{} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in \mathcal{U} \Delta \mathcal{V}} W \neq \left( \bigcup_{U \in \mathcal{U}} U \right) \Delta \left( \bigcup_{V \in \mathcal{V}} V \right)$$

in general, where  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01G2

14. *Interaction With Internal Hom I.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-1, -2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cup^{\text{op}} \times \cup^{\text{op}} & \text{X} & \downarrow \cup \\
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[ \bigcup_{U \in \mathcal{U}} U, \bigcup_{V \in \mathcal{V}} V \right]_X$$

in general, where  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

01G3

15. *Interaction With Internal Hom II.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\
 & \nearrow \dashv \quad \searrow \cup^{\text{op}} & \\
 \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\
 \downarrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & & \downarrow [-, V]_X \\
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$\left[ \bigcup_{U \in \mathcal{U}} U, V \right]_X = \bigcap_{U \in \mathcal{U}} [U, V]_X$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$  and each  $V \in \mathcal{P}(X)$ .

01G4

16. *Interaction With Internal Hom III.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \\
 \downarrow \text{id}_{\mathcal{P}(X)} \star [U, -]_X & & \downarrow [-, -]_X \\
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$\left[ U, \bigcup_{V \in \mathcal{V}} V \right]_X = \bigcup_{V \in \mathcal{V}} [U, V]_X$$

for each  $U \in \mathcal{P}(X)$  and each  $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01G5

17. *Interaction With Direct Images.* Let  $f: X \rightarrow Y$  be a map of sets.  
The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_!)!} & \mathcal{P}(\mathcal{P}(Y)) \\
 \downarrow \cup & & \downarrow \cup \\
 \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_!(U) = \bigcup_{V \in f_!(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f_!)_!(\mathcal{U})$ .

01G6

18. *Interaction With Inverse Images.* Let  $f: X \rightarrow Y$  be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{V \in \mathcal{V}} f^{-1}(V) = \bigcup_{U \in f^{-1}(\mathcal{U})} U$$

for each  $\mathcal{V} \in \mathcal{P}(Y)$ , where  $f^{-1}(\mathcal{V}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{V})$ .

01G7

19. *Interaction With Codirect Images.* Let  $f: X \rightarrow Y$  be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_*(U) = \bigcup_{V \in f_*(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$ .

01G8

20. *Interaction With Intersections of Families I.* The diagram


$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \cap} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \star \text{id}_{\mathcal{P}(X)} \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{\cap} & X \end{array}$$

commutes, i.e. we have

$$\bigcap_{\substack{U \in \bigcup_{A \in \mathcal{A}} A}} U = \bigcap_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right)$$

for each  $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$ .

- 01G9** 21. *Interaction With Intersections of Families II.* Let  $X$  be a set and consider the compositions



given by

$$\begin{aligned} \mathcal{A} &\mapsto \bigcup_{\substack{U \in \bigcap_{A \in \mathcal{A}} A}} U, & \mathcal{A} &\mapsto \bigcap_{\substack{U \in \bigcup_{A \in \mathcal{A}} A}} U, \\ \mathcal{A} &\mapsto \bigcup_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right), & \mathcal{A} &\mapsto \bigcap_{A \in \mathcal{A}} \left( \bigcup_{U \in A} U \right) \end{aligned}$$

for each  $\mathcal{A} \in \mathcal{P}(\mathcal{P}(\mathcal{P}(X)))$ . We have the following inclusions:

$$\begin{array}{ccc}
 & \bigcup_{U \in \bigcap_{A \in \mathcal{A}} A} U & \\
 & \swarrow \quad \searrow & \\
 \bigcup_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right) & & \bigcap_{A \in \mathcal{A}} \left( \bigcup_{U \in A} U \right) \\
 & \swarrow \quad \searrow & \\
 & \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U &
 \end{array}$$

All other possible inclusions fail to hold in general.

#### PROOF 4.3.6.1.3 ► PROOF OF PROPOSITION 4.3.6.1.2

##### Item 1: Functoriality

Since  $\mathcal{P}(X)$  is posetal, it suffices to prove the condition  $(\star)$ . So let  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$  with  $\mathcal{U} \subset \mathcal{V}$ . We claim that

$$\bigcup_{U \in \mathcal{U}} U \subset \bigcup_{V \in \mathcal{V}} V.$$

Indeed, given  $x \in \bigcup_{U \in \mathcal{U}} U$ , there exists some  $U \in \mathcal{U}$  such that  $x \in U$ , but since  $\mathcal{U} \subset \mathcal{V}$ , we have  $U \in \mathcal{V}$  as well, and thus  $x \in \bigcup_{V \in \mathcal{V}} V$ , which gives our desired inclusion.

##### Item 2: Associativity

We have

$$\begin{aligned}
 \bigcup_{U \in \bigcup_{A \in \mathcal{A}} A} U &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } U \in \bigcup_{A \in \mathcal{A}} A \\ \text{such that we have } x \in U \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } A \in \mathcal{A} \\ \text{and some } U \in A \text{ such that} \\ \text{we have } x \in U \end{array} \right\}
 \end{aligned}$$

$$\begin{aligned}
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } A \in \mathcal{A} \\ \text{such that we have } x \in \bigcup_{U \in A} U \end{array} \right\} \\
 &\stackrel{\text{def}}{=} \bigcup_{A \in \mathcal{A}} \left( \bigcup_{U \in A} U \right).
 \end{aligned}$$

This finishes the proof.

#### Item 3: Left Unitality

We have

$$\begin{aligned}
 \bigcup_{V \in \{U\}} V &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } V \in \{U\} \\ \text{such that we have } x \in U \end{array} \right\} \\
 &= \{x \in X \mid x \in U\} \\
 &= U.
 \end{aligned}$$

This finishes the proof.

#### Item 4: Right Unitality

We have

$$\begin{aligned}
 \bigcup_{\{u\} \in \chi_X(U)} \{u\} &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } \{u\} \in \chi_X(U) \\ \text{such that we have } x \in \{u\} \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } \{u\} \in \chi_X(U) \\ \text{such that we have } x = u \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } u \in U \\ \text{such that we have } x = u \end{array} \right\} \\
 &= \{x \in X \mid x \in U\} \\
 &= U.
 \end{aligned}$$

This finishes the proof.

#### Item 5: Interaction With Unions I

We have

$$\begin{aligned}
 \bigcup_{W \in \mathcal{U} \cup \mathcal{V}} W &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } W \in \mathcal{U} \cup \mathcal{V} \\ \text{such that we have } x \in W \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } W \in \mathcal{U} \text{ or some } \\ W \in \mathcal{V} \text{ such that we have } x \in W \end{array} \right\}
 \end{aligned}$$

$$\begin{aligned}
& \stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } W \in \mathcal{U} \\ \text{such that we have } x \in W \end{array} \right\} \\
& \quad \cup \left\{ x \in X \mid \begin{array}{l} \text{there exists some } W \in \mathcal{V} \\ \text{such that we have } x \in W \end{array} \right\} \\
& \stackrel{\text{def}}{=} \left( \bigcup_{W \in \mathcal{U}} W \right) \cup \left( \bigcup_{W \in \mathcal{V}} W \right) \\
& = \left( \bigcup_{U \in \mathcal{U}} U \right) \cup \left( \bigcup_{V \in \mathcal{V}} V \right).
\end{aligned}$$

This finishes the proof.

#### Item 6: Interaction With Unions II

Assume  $\mathcal{V}$  is nonempty. We have

$$\begin{aligned}
U \cup \bigcup_{V \in \mathcal{V}} V & \stackrel{\text{def}}{=} \left\{ x \in X \mid x \in U \text{ or } x \in \bigcup_{V \in \mathcal{V}} V \right\} \\
& = \left\{ x \in X \mid \begin{array}{l} x \in U \text{ or there exists some} \\ V \in \mathcal{V} \text{ such that } x \in V \end{array} \right\} \\
& = \left\{ x \in X \mid \begin{array}{l} \text{there exists some } V \in \mathcal{V} \\ \text{such that } x \in U \text{ or } x \in V \end{array} \right\} \\
& = \left\{ x \in X \mid \begin{array}{l} \text{there exists some } V \in \mathcal{V} \\ \text{such that } x \in U \cup V \end{array} \right\} \\
& \stackrel{\text{def}}{=} \bigcup_{V \in \mathcal{V}} U \cup V.
\end{aligned}$$

This concludes the proof of the first statement. For the second statement, use [Item 4 of Proposition 4.3.8.1.2](#) to rewrite

$$\begin{aligned}
\left( \bigcup_{U \in \mathcal{U}} U \right) \cup V & = V \cup \left( \bigcup_{U \in \mathcal{U}} U \right), \\
\bigcup_{U \in \mathcal{U}} (U \cup V) & = \bigcup_{U \in \mathcal{U}} (V \cup U).
\end{aligned}$$

But these two sets are equal by the first statement.

#### Item 7: Interaction With Intersections I

We have

$$\begin{aligned}
 \bigcup_{W \in \mathcal{U} \cap \mathcal{V}} W &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{there exists some } W \in \mathcal{U} \cap \mathcal{V} \\ \text{such that we have } x \in W \end{array} \right\} \\
 &\subset \left\{ x \in X \mid \begin{array}{l} \text{there exists some } U \in \mathcal{U} \text{ and some } V \in \mathcal{V} \\ \text{such that we have } x \in U \text{ and } x \in V \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } U \in \mathcal{U} \\ \text{such that we have } x \in U \end{array} \right\} \\
 &\quad \cup \left\{ x \in X \mid \begin{array}{l} \text{there exists some } V \in \mathcal{V} \\ \text{such that we have } x \in V \end{array} \right\} \\
 &\stackrel{\text{def}}{=} \left( \bigcup_{U \in \mathcal{U}} U \right) \cap \left( \bigcup_{V \in \mathcal{V}} V \right).
 \end{aligned}$$

This finishes the proof.

#### Item 8: Interaction With Intersections II

We have

$$\begin{aligned}
 U \cap \bigcup_{V \in \mathcal{V}} V &\stackrel{\text{def}}{=} \left\{ x \in X \mid x \in U \text{ and } x \in \bigcup_{V \in \mathcal{V}} V \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} x \in U \text{ and there exists some} \\ V \in \mathcal{V} \text{ such that } x \in V \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } V \in \mathcal{V} \\ \text{such that } x \in U \text{ and } x \in V \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } V \in \mathcal{V} \\ \text{such that } x \in U \cap V \end{array} \right\} \\
 &\stackrel{\text{def}}{=} \bigcup_{V \in \mathcal{V}} U \cap V.
 \end{aligned}$$

This concludes the proof of the first statement. For the second statement, use [Item 5 of Proposition 4.3.9.1.2](#) to rewrite

$$\begin{aligned}
 \left( \bigcup_{U \in \mathcal{U}} U \right) \cap V &= V \cap \left( \bigcup_{U \in \mathcal{U}} U \right), \\
 \bigcup_{U \in \mathcal{U}} (U \cap V) &= \bigcup_{U \in \mathcal{U}} (V \cap U).
 \end{aligned}$$

But these two sets are equal by the first statement.

**Item 9: Interaction With Differences**

Let  $X = \{0, 1\}$ , let  $\mathcal{U} = \{\{0, 1\}\}$ , and let  $\mathcal{V} = \{\{0\}\}$ . We have

$$\begin{aligned}\bigcup_{W \in \mathcal{U} \setminus \mathcal{V}} U &= \bigcup_{W \in \{\{0, 1\}\}} W \\ &= \{0, 1\},\end{aligned}$$

whereas

$$\begin{aligned}\left(\bigcup_{U \in \mathcal{U}} U\right) \setminus \left(\bigcup_{V \in \mathcal{V}} V\right) &= \{0, 1\} \setminus \{0\} \\ &= \{1\}.\end{aligned}$$

Thus we have

$$\bigcup_{W \in \mathcal{U} \setminus \mathcal{V}} W = \{0, 1\} \neq \{1\} = \left(\bigcup_{U \in \mathcal{U}} U\right) \setminus \left(\bigcup_{V \in \mathcal{V}} V\right).$$

This finishes the proof.

**Item 10: Interaction With Complements I**

Let  $X = \{0, 1\}$  and let  $\mathcal{U} = \{0\}$ . We have

$$\begin{aligned}\bigcup_{U \in \mathcal{U}^c} U &= \bigcup_{U \in \{\emptyset, \{1\}, \{0, 1\}\}} U \\ &= \{0, 1\},\end{aligned}$$

whereas

$$\begin{aligned}\bigcup_{U \in \mathcal{U}} U^c &= \{0\}^c \\ &= \{1\}.\end{aligned}$$

Thus we have

$$\bigcup_{U \in \mathcal{U}^c} U = \{0, 1\} \neq \{1\} = \bigcup_{U \in \mathcal{U}} U^c.$$

This finishes the proof.

**Item 11: Interaction With Complements II**

We have

$$\left(\bigcup_{U \in \mathcal{U}} U\right)^c \stackrel{\text{def}}{=} \left\{x \in X \mid \begin{array}{l} \text{there exists no } U \in \mathcal{U} \\ \text{such that we have } x \in U\end{array}\right\}$$

$$\begin{aligned}
 &= \left\{ x \in X \mid \begin{array}{l} \text{for all } U \in \mathcal{U} \\ \text{we have } x \notin U \end{array} \right\} \\
 &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for all } U \in \mathcal{U} \\ \text{we have } x \in U^c \end{array} \right\} \\
 &\stackrel{\text{def}}{=} \bigcap_{U \in \mathcal{U}} U^c.
 \end{aligned}$$

## Item 12: Interaction With Complements III

By Item 11 Item 3 of Proposition 4.3.11.1.2, we have

$$\begin{aligned}
 \left( \bigcap_{U \in \mathcal{U}} U \right)^c &= \left( \bigcap_{U \in \mathcal{U}} (U^c)^c \right)^c \\
 &= \left( \left( \bigcup_{U \in \mathcal{U}} U^c \right)^c \right)^c \\
 &= \bigcup_{U \in \mathcal{U}} U^c.
 \end{aligned}$$

## Item 13: Interaction With Symmetric Differences

Let  $X = \{0, 1\}$ , let  $\mathcal{U} = \{\{0, 1\}\}$ , and let  $\mathcal{V} = \{\{0\}, \{0, 1\}\}$ . We have

$$\begin{aligned}
 \bigcup_{W \in \mathcal{U} \Delta \mathcal{V}} W &= \bigcup_{W \in \{\{0\}\}} W \\
 &= \{0\},
 \end{aligned}$$

whereas

$$\begin{aligned}
 \left( \bigcup_{U \in \mathcal{U}} U \right) \Delta \left( \bigcup_{V \in \mathcal{V}} V \right) &= \{0, 1\} \Delta \{0, 1\} \\
 &= \emptyset,
 \end{aligned}$$

Thus we have

$$\bigcup_{W \in \mathcal{U} \Delta \mathcal{V}} W = \{0\} \neq \emptyset = \left( \bigcup_{U \in \mathcal{U}} U \right) \Delta \left( \bigcup_{V \in \mathcal{V}} V \right).$$

This finishes the proof.

## Item 14: Interaction With Internal Homs I

This is a repetition of [Item 7 of Proposition 4.4.7.1.4](#) and is proved there.

[Item 15: Interaction With Internal Homs II](#)

This is a repetition of [Item 8 of Proposition 4.4.7.1.4](#) and is proved there.

[Item 16: Interaction With Internal Homs III](#)

This is a repetition of [Item 9 of Proposition 4.4.7.1.4](#) and is proved there.

[Item 17: Interaction With Direct Images](#)

This is a repetition of [Item 3 of Proposition 4.6.1.1.5](#) and is proved there.

[Item 18: Interaction With Inverse Images](#)

This is a repetition of [Item 3 of Proposition 4.6.2.1.3](#) and is proved there.

[Item 19: Interaction With Codirect Images](#)

This is a repetition of [Item 3 of Proposition 4.6.3.1.7](#) and is proved there.

[Item 20: Interaction With Intersections of Families I](#)

We have

$$\begin{aligned} \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } U \in \bigcup_{A \in \mathcal{A}} A, \\ \text{we have } x \in U \end{array} \right\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{for each } A \in \mathcal{A} \text{ and each} \\ U \in A, \text{ we have } x \in U \end{array} \right\} \\ &\stackrel{\text{def}}{=} \bigcap_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right). \end{aligned}$$

This finishes the proof.

[Item 21: Interaction With Intersections of Families II](#)

Omitted. 

### 003V 4.3.7 Intersections of Families of Subsets

Let  $X$  be a set and let  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

003W

#### DEFINITION 4.3.7.1.1 ► INTERSECTIONS OF FAMILIES OF SUBSETS

The **intersection** of  $\mathcal{U}$  is the set  $\bigcap_{U \in \mathcal{U}} U$  defined by

$$\bigcap_{U \in \mathcal{U}} U \stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } U \in \mathcal{U}, \\ \text{we have } x \in U \end{array} \right\}.$$

01GA

**PROPOSITION 4.3.7.1.2 ► PROPERTIES OF INTERSECTIONS OF FAMILIES OF SUBSETS**

Let  $X$  be a set.

01GB

- Functoriality.* The assignment  $\mathcal{U} \mapsto \bigcap_{U \in \mathcal{U}} U$  defines a functor

$$\bigcap: (\mathcal{P}(\mathcal{P}(X)), \supset) \rightarrow (\mathcal{P}(X), \subset).$$

In particular, for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ , the following condition is satisfied:

$$(\star) \text{ If } \mathcal{U} \subset \mathcal{V}, \text{ then } \bigcap_{V \in \mathcal{V}} V \subset \bigcap_{U \in \mathcal{U}} U.$$

01GC

- Oplax Associativity.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(\mathcal{P}(X))) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \bigcap} & \mathcal{P}(\mathcal{P}(X)) \\ \bigcap \star \text{id}_{\mathcal{P}(X)} \downarrow & \curvearrowright & \downarrow \bigcap \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow[\bigcap]{} & \mathcal{P}(X) \end{array}$$

with components

$$\bigcap_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right) \subset \bigcap_{U \in \bigcap_{A \in \mathcal{A}} A} U$$

for each  $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$ .

01GD

- Left Unitality.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \downarrow \chi_{\mathcal{P}(X)} & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow[\bigcap]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcap_{V \in \{U\}} V = U.$$

for each  $U \in \mathcal{P}(X)$ .

01GE

4. *Oplax Right Unitality.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \downarrow \mathcal{P}(\chi_X) & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

*does not commute in general, i.e. we may have*

$$\bigcap_{\{x\} \in \chi_X(U)} \{x\} \neq U$$

in general, where  $U \in \mathcal{P}(X)$ . However, when  $U$  is nonempty, we have

$$\bigcap_{\{x\} \in \chi_X(U)} \{x\} \subset U.$$

01GF

5. *Interaction With Unions I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cap \times \cap & & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\cap]{} & \mathcal{P}(X) \end{array}$$

*commutes, i.e. we have*

$$\bigcap_{W \in \mathcal{U} \cup \mathcal{V}} W = \left( \bigcap_{U \in \mathcal{U}} U \right) \cap \left( \bigcap_{V \in \mathcal{V}} V \right)$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01GG

6. *Interaction With Unions II.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cup -)} & \mathcal{P}(\mathcal{P}(X)) & \quad \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cup V)} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cap & & \downarrow \cap & \quad \downarrow \cap & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow[U \cup -]{} & \mathcal{P}(X) & \quad \mathcal{P}(X) & \xrightarrow[- \cup V]{} & \mathcal{P}(X) \end{array}$$

*commute, i.e. we have*

$$\begin{aligned} U \cup \left( \bigcap_{V \in \mathcal{V}} V \right) &= \bigcap_{V \in \mathcal{V}} (U \cup V), \\ \left( \bigcap_{U \in \mathcal{U}} U \right) \cup V &= \bigcap_{U \in \mathcal{U}} (U \cup V) \end{aligned}$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$  and each  $U, V \in \mathcal{P}(X)$ .

01GH

7. *Interaction With Intersections I.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \times \cap \downarrow & \curvearrowright & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\cap]{} & \mathcal{P}(X), \end{array}$$

with components

$$\left( \bigcap_{U \in \mathcal{U}} U \right) \cap \left( \bigcap_{V \in \mathcal{V}} V \right) \subset \bigcap_{W \in \mathcal{U} \cap \mathcal{V}} W$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01GJ

8. *Interaction With Intersections II.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cap -)} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow[U \cap -]{} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cap V)} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow[- \cap V]{} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$\begin{aligned} U \cup \left( \bigcap_{V \in \mathcal{V}} V \right) &= \bigcap_{V \in \mathcal{V}} (U \cup V), \\ \left( \bigcap_{U \in \mathcal{U}} U \right) \cup V &= \bigcap_{U \in \mathcal{U}} (U \cup V) \end{aligned}$$

for each  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$  and each  $U, V \in \mathcal{P}(X)$ .

01GK

9. *Interaction With Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\setminus} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \times \cap \downarrow & \text{X} & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\setminus]{} & \mathcal{P}(X), \end{array}$$

*does not commute* in general, i.e. we may have

$$\bigcap_{W \in \mathcal{U} \setminus \mathcal{V}} W \neq \left( \bigcap_{U \in \mathcal{U}} U \right) \setminus \left( \bigcap_{V \in \mathcal{V}} V \right)$$

in general, where  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01GL

10. *Interaction With Complements I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(\mathcal{P}(X)) \\ \cap^{\text{op}} \downarrow & \text{X} & \downarrow \cap \\ \mathcal{P}(X)^{\text{op}} & \xrightarrow[(-)^c]{} & \mathcal{P}(X), \end{array}$$

*does not commute in general, i.e. we may have*

$$\bigcap_{W \in \mathcal{U}^c} W \neq \bigcap_{U \in \mathcal{U}} U^c$$

*in general, where  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .*

01GM

11. *Interaction With Complements II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & & \searrow \sim \\ \mathcal{P}(\mathcal{P}(X)) & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\ \searrow \cap & & \swarrow \cup^{\text{op}} \\ \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}}, \end{array}$$

*commutes, i.e. we have*

$$\left( \bigcap_{U \in \mathcal{U}} U \right)^c = \bigcup_{U \in \mathcal{U}} U^c$$

*for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .*

01GN

12. *Interaction With Complements III.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & & \searrow \sim \\ \mathcal{P}(\mathcal{P}(X)) & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\ \searrow \cup & & \swarrow \cap^{\text{op}} \\ \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}}, \end{array}$$

commutes, i.e. we have

$$\left( \bigcup_{U \in \mathcal{U}} U \right)^c = \bigcap_{U \in \mathcal{U}} U^c$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

**01GP** 13. *Interaction With Symmetric Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\Delta} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \times \cap \downarrow & \times & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\Delta]{} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in \mathcal{U} \Delta \mathcal{V}} W \neq \left( \bigcap_{U \in \mathcal{U}} U \right) \Delta \left( \bigcap_{V \in \mathcal{V}} V \right)$$

in general, where  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

**01GQ** 14. *Interaction With Internal Hom I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-_1, -_2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\ \cap^{\text{op}} \times \cap^{\text{op}} \downarrow & \times & \downarrow \cap \\ \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-_1, -_2]_X} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[ \bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X$$

in general, where  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

**01GR** 15. *Interaction With Internal Hom II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ & \nearrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & \searrow \cap^{\text{op}} \\ \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\ & \swarrow id_{\mathcal{P}(X)} \star [-, V]_X & \searrow [-, V]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow[\cup]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[ \bigcap_{U \in \mathcal{U}} U, V \right]_X = \bigcup_{U \in \mathcal{U}} [U, V]_X$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$  and each  $V \in \mathcal{P}(X)$ .

- 01GS** 16. *Interaction With Internal Hom III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \\ \text{id}_{\mathcal{P}(X)} \star [U, -]_X \downarrow & & \downarrow [U, -]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[ U, \bigcap_{V \in \mathcal{V}} V \right]_X = \bigcap_{V \in \mathcal{V}} [U, V]_X$$

for each  $U \in \mathcal{P}(X)$  and each  $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

- 01GT** 17. *Interaction With Direct Images.* Let  $f: X \rightarrow Y$  be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_!)!} & \mathcal{P}(\mathcal{P}(Y)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_!(U) = \bigcap_{V \in f_!(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f_!)_!(\mathcal{U})$ .

- 01GU** 18. *Interaction With Inverse Images.* Let  $f: X \rightarrow Y$  be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcap_{V \in \mathcal{V}} f^{-1}(V) = \bigcap_{U \in f^{-1}(\mathcal{U})} U$$

for each  $\mathcal{V} \in \mathcal{P}(Y)$ , where  $f^{-1}(\mathcal{V}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{V})$ .

01GV

19. *Interaction With Codirect Images.* Let  $f: X \rightarrow Y$  be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_*(U) = \bigcap_{V \in f_*(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$ .

01GW

20. *Interaction With Unions of Families I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \cap} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \star \text{id}_{\mathcal{P}(X)} \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{\cap} & X \end{array}$$

commutes, i.e. we have


$$\bigcup_{U \in \bigcup_{A \in \mathcal{A}} A} U = \bigcap_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right)$$

for each  $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$ .

01GX

21. *Interaction With Unions of Families II.* Let  $X$  be a set and consider

the compositions



given by

$$\begin{aligned}
 \mathcal{A} &\mapsto \bigcup_{U \in \bigcap_{A \in \mathcal{A}} A} U, & \mathcal{A} &\mapsto \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U, \\
 \mathcal{A} &\mapsto \bigcup_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right), & \mathcal{A} &\mapsto \bigcap_{A \in \mathcal{A}} \left( \bigcup_{U \in A} U \right)
 \end{aligned}$$

for each  $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$ . We have the following inclusions:

$$\begin{array}{ccc}
 \bigcup_{U \in \bigcap_{A \in \mathcal{A}} A} U & & \\
 \uparrow & \nearrow & \\
 \bigcup_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right) & & \bigcap_{A \in \mathcal{A}} \left( \bigcup_{U \in A} U \right) \\
 \downarrow & & \uparrow \\
 \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U & &
 \end{array}$$

All other possible inclusions fail to hold in general.



**Item 1: Functoriality**

Since  $\mathcal{P}(X)$  is posetal, it suffices to prove the condition  $(\star)$ . So let  $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$  with  $\mathcal{U} \subset \mathcal{V}$ . We claim that

$$\bigcap_{V \in \mathcal{V}} V \subset \bigcap_{U \in \mathcal{U}} U.$$

Indeed, if  $x \in \bigcap_{V \in \mathcal{V}} V$ , then  $x \in V$  for all  $V \in \mathcal{V}$ . But since  $\mathcal{U} \subset \mathcal{V}$ , it follows that  $x \in U$  for all  $U \in \mathcal{U}$  as well. Thus  $x \in \bigcap_{U \in \mathcal{U}} U$ , which gives our desired inclusion.

**Item 2: Oplax Associativity**

We have

$$\begin{aligned} \bigcap_{A \in \mathcal{A}} \left( \bigcap_{U \in A} U \right) &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } A \in \mathcal{A}, \\ \text{we have } x \in \bigcap_{U \in A} U \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } A \in \mathcal{A} \text{ and each} \\ U \in A, \text{ we have } x \in U \end{array} \right\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{for each } U \in \bigcup_{A \in \mathcal{A}} A, \\ \text{we have } x \in U \end{array} \right\} \\ &\subset \left\{ x \in X \mid \begin{array}{l} \text{for each } U \in \bigcap_{A \in \mathcal{A}} A, \\ \text{we have } x \in U \end{array} \right\} \\ &\stackrel{\text{def}}{=} \bigcap_{U \in \bigcap_{A \in \mathcal{A}} A} U. \end{aligned}$$

Since  $\mathcal{P}(X)$  is posetal, naturality is automatic (?? of ??). This finishes the proof.

**Item 3: Left Unitality**

We have

$$\begin{aligned} \bigcap_{V \in \{U\}} V &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } V \in \{U\}, \\ \text{we have } x \in U \end{array} \right\} \\ &= \{x \in X \mid x \in U\} \\ &= U. \end{aligned}$$

This finishes the proof.

## Item 4: Oplax Right Unitality

If  $U = \emptyset$ , then we have

$$\begin{aligned} \bigcap_{\{u\} \in \chi_X(U)} \{u\} &= \bigcap_{\{u\} \in \emptyset} \{u\} \\ &= X, \end{aligned}$$

so  $\bigcap_{\{u\} \in \chi_X(U)} \{u\} = X \neq \emptyset = U$ . When  $U$  is nonempty, we have two cases:

020B

1. If  $U$  is a singleton, say  $U = \{u\}$ , we have

$$\begin{aligned} \bigcap_{\{u\} \in \chi_X(U)} \{u\} &= \{u\} \\ &\stackrel{\text{def}}{=} U. \end{aligned}$$

020C

2. If  $U$  contains at least two elements, we have

$$\begin{aligned} \bigcap_{\{u\} \in \chi_X(U)} \{u\} &= \emptyset \\ &\subset U. \end{aligned}$$

This finishes the proof.

## Item 5: Interaction With Unions I

We have

$$\begin{aligned} \bigcap_{W \in \mathcal{U} \cup \mathcal{V}} W &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } W \in \mathcal{U} \cup \mathcal{V}, \\ \text{we have } x \in W \end{array} \right\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{for each } W \in \mathcal{U} \text{ and each} \\ W \in \mathcal{V}, \text{ we have } x \in W \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } W \in \mathcal{U}, \\ \text{we have } x \in W \end{array} \right\} \\ &\quad \cap \left\{ x \in X \mid \begin{array}{l} \text{for each } W \in \mathcal{V}, \\ \text{we have } x \in W \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left( \bigcap_{W \in \mathcal{U}} W \right) \cap \left( \bigcap_{W \in \mathcal{V}} W \right) \\ &= \left( \bigcap_{U \in \mathcal{U}} U \right) \cap \left( \bigcap_{V \in \mathcal{V}} V \right). \end{aligned}$$

This finishes the proof.

**Item 6: Interaction With Unions II**

Omitted.

**Item 7: Interaction With Intersections I**

We have

$$\begin{aligned}
 \left( \bigcap_{U \in \mathcal{U}} U \right) \cap \left( \bigcap_{V \in \mathcal{V}} V \right) &\stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } U \in \mathcal{U}, \\ \text{we have } x \in U \end{array} \right\} \\
 &\quad \cup \left\{ x \in X \mid \begin{array}{l} \text{for each } V \in \mathcal{V}, \\ \text{we have } x \in V \end{array} \right\} \\
 &= \left\{ x \in X \mid \begin{array}{l} \text{for each } W \in \mathcal{U} \cap \mathcal{V}, \\ \text{we have } x \in W \end{array} \right\} \\
 &\subset \left\{ x \in X \mid \begin{array}{l} \text{for each } W \in \mathcal{U} \cup \mathcal{V}, \\ \text{we have } x \in W \end{array} \right\} \\
 &\stackrel{\text{def}}{=} \bigcap_{W \in \mathcal{U} \cup \mathcal{V}} W.
 \end{aligned}$$

Since  $\mathcal{P}(X)$  is posetal, naturality is automatic (?? of ??). This finishes the proof.

**Item 8: Interaction With Intersections II**

Omitted.

**Item 9: Interaction With Differences**

Let  $X = \{0, 1\}$ , let  $\mathcal{U} = \{\{0\}, \{0, 1\}\}$ , and let  $\mathcal{V} = \{\{0\}\}$ . We have

$$\begin{aligned}
 \bigcap_{W \in \mathcal{U} \setminus \mathcal{V}} W &= \bigcap_{W \in \{\{0, 1\}\}} W \\
 &= \{0, 1\},
 \end{aligned}$$

whereas

$$\begin{aligned}
 \left( \bigcap_{U \in \mathcal{U}} U \right) \setminus \left( \bigcap_{V \in \mathcal{V}} V \right) &= \{0\} \setminus \{0\} \\
 &= \emptyset.
 \end{aligned}$$

Thus we have

$$\bigcap_{W \in \mathcal{U} \setminus \mathcal{V}} W = \{0, 1\} \neq \emptyset = \left( \bigcap_{U \in \mathcal{U}} U \right) \setminus \left( \bigcap_{V \in \mathcal{V}} V \right).$$

This finishes the proof.

**Item 10: Interaction With Complements I**

Let  $X = \{0, 1\}$  and let  $\mathcal{U} = \{\{0\}\}$ . We have

$$\begin{aligned}\bigcap_{W \in \mathcal{U}^c} U &= \bigcap_{W \in \{\emptyset, \{1\}, \{0, 1\}\}} W \\ &= \emptyset,\end{aligned}$$

whereas

$$\begin{aligned}\bigcap_{U \in \mathcal{U}} U^c &= \{0\}^c \\ &= \{1\}.\end{aligned}$$

Thus we have

$$\bigcap_{W \in \mathcal{U}^c} U = \emptyset \neq \{1\} = \bigcap_{U \in \mathcal{U}} U^c.$$

This finishes the proof.

**Item 11: Interaction With Complements II**

This is a repetition of [Item 12 of Proposition 4.3.6.1.2](#) and is proved there.

**Item 12: Interaction With Complements III**

This is a repetition of [Item 11 of Proposition 4.3.6.1.2](#) and is proved there.

**Item 13: Interaction With Symmetric Differences**

Let  $X = \{0, 1\}$ , let  $\mathcal{U} = \{\{0, 1\}\}$ , and let  $\mathcal{V} = \{\{0\}, \{0, 1\}\}$ . We have

$$\begin{aligned}\bigcap_{W \in \mathcal{U} \Delta \mathcal{V}} W &= \bigcap_{W \in \{\{0\}\}} W \\ &= \{0\},\end{aligned}$$

whereas

$$\begin{aligned}\left(\bigcap_{U \in \mathcal{U}} U\right) \Delta \left(\bigcap_{V \in \mathcal{V}} V\right) &= \{0, 1\} \Delta \{0\} \\ &= \emptyset,\end{aligned}$$

Thus we have

$$\bigcap_{W \in \mathcal{U} \Delta \mathcal{V}} W = \{0\} \neq \emptyset = \left(\bigcap_{U \in \mathcal{U}} U\right) \Delta \left(\bigcap_{V \in \mathcal{V}} V\right).$$

This finishes the proof.

**Item 14: Interaction With Internal Homs I**

This is a repetition of [Item 10](#) of [Proposition 4.4.7.1.4](#) and is proved there.

**Item 15: Interaction With Internal Homs II**

This is a repetition of [Item 11](#) of [Proposition 4.4.7.1.4](#) and is proved there.

**Item 16: Interaction With Internal Homs III**

This is a repetition of [Item 12](#) of [Proposition 4.4.7.1.4](#) and is proved there.

**Item 17: Interaction With Direct Images**

This is a repetition of [Item 4](#) of [Proposition 4.6.1.1.5](#) and is proved there.

**Item 18: Interaction With Inverse Images**

This is a repetition of [Item 4](#) of [Proposition 4.6.2.1.3](#) and is proved there.

**Item 19: Interaction With Codirect Images**

This is a repetition of [Item 4](#) of [Proposition 4.6.3.1.7](#) and is proved there.

**Item 20: Interaction With Unions of Families I**

This is a repetition of [Item 20](#) of [Proposition 4.3.6.1.2](#) and is proved there.

**Item 21: Interaction With Unions of Families II**

This is a repetition of [Item 21](#) of [Proposition 4.3.6.1.2](#) and is proved there. 

### 003G 4.3.8 Binary Unions

Let  $X$  be a set and let  $U, V \in \mathcal{P}(X)$ .

003H

#### DEFINITION 4.3.8.1.1 ► BINARY UNIONS

The **union of  $U$  and  $V$**  is the set  $U \cup V$  defined by

$$\begin{aligned} U \cup V &\stackrel{\text{def}}{=} \bigcup_{z \in \{U, V\}} z \\ &\stackrel{\text{def}}{=} \{x \in X \mid x \in U \text{ or } x \in V\}. \end{aligned}$$

003J

#### PROPOSITION 4.3.8.1.2 ► PROPERTIES OF BINARY UNIONS

Let  $X$  be a set.

003K

1. *Functionality.* The assignments  $U, V, (U, V) \mapsto U \cup V$  define functors

$$\begin{aligned} U \cup -: & (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ - \cup V: & (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cup -_2: & (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each  $U, V, A, B \in \mathcal{P}(X)$ :

01GY

(a) If  $U \subset A$ , then  $U \cup V \subset A \cup V$ .

01GZ

(b) If  $V \subset B$ , then  $U \cup V \subset U \cup B$ .

01H0

(c) If  $U \subset A$  and  $V \subset B$ , then  $U \cup V \subset A \cup B$ .

003M

2. *Associativity.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\ \alpha_{\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \nearrow & \swarrow \text{id}_{\mathcal{P}(X)} \times \cup & \\ (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cup \times \text{id}_{\mathcal{P}(X)} \searrow & & \swarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\cup]{} & \mathcal{P}(X), \end{array}$$

commutes, i.e. we have an equality of sets

$$(U \cup V) \cup W = U \cup (V \cup W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

003N

3. *Unitality.* The diagrams

$$\begin{array}{ccc} \text{pt} \times \mathcal{P}(X) & \xrightarrow{[\emptyset] \times \text{id}_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) & \quad \mathcal{P}(X) \times \text{pt} & \xrightarrow{\text{id}_{\mathcal{P}(X)} \times [\emptyset]} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \lambda_{\mathcal{P}(X)}^{\text{Sets}} \sim \searrow & & \downarrow \cup & \rho_{\mathcal{P}(X)}^{\text{Sets}} \sim \searrow & & \downarrow \cup \\ & & \mathcal{P}(X) & & & \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned} \emptyset \cup U &= U, \\ U \cup \emptyset &= U \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

003P

## 4. Commutativity. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\sigma_{\mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \quad} & \mathcal{P}(X) \times \mathcal{P}(X) \\ & \searrow \cup & \downarrow \cup \\ & & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cup V = V \cup U$$

for each  $U, V \in \mathcal{P}(X)$ .

01H1

5. Annihilation With  $X$ . The diagrams

$$\begin{array}{ccccc} \text{id}_{\text{pt}} \times \epsilon_{\mathcal{P}(X)}^{\text{Sets}} & \nearrow \text{pt} \times \text{pt} & & \epsilon_{\mathcal{P}(X)}^{\text{Sets}} \times \text{id}_{\text{pt}} & \nearrow \text{pt} \times \text{pt} \\ \text{pt} \times \mathcal{P}(X) & & \text{pt} & & \mathcal{P}(X) \times \text{pt} \\ \downarrow [X] \times \text{id}_{\mathcal{P}(X)} & & \downarrow [X] & & \downarrow \text{id}_{\mathcal{P}(X)} \times [X] \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cup} & \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X) \xrightarrow{\cup} \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$U \cup X = X,$$

$$X \cup V = X$$

for each  $U, V \in \mathcal{P}(X)$ .

003R

## 6. Distributivity of Unions Over Intersections. The diagrams

$$\begin{array}{ccc} (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) \\ \Delta_{\mathcal{P}(X)} \times (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \nearrow & \searrow \mu_{4|(\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X))}^{\text{Sets}} & \nearrow (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \times \Delta_{\mathcal{P}(X)} & \searrow \mu_{4|(\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X))}^{\text{Sets}} \\ \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) \\ \downarrow \text{id}_{\mathcal{P}(X)} \times \cap & & \downarrow \cup \times \cup & & \downarrow \cap \times \text{id}_{\mathcal{P}(X)} \\ \mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X), & & \mathcal{P}(X) \times \mathcal{P}(X) \\ \searrow \cup & \nearrow \cap & & \searrow \cup & \nearrow \cap \\ & \mathcal{P}(X) & & & \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$U \cup (V \cap W) = (U \cup V) \cap (U \cup W),$$

$$(U \cap V) \cup W = (U \cup W) \cap (V \cup W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

01H2

## 7. Distributivity of Intersections Over Unions. The diagrams

$$\begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \Delta_{\mathcal{P}(X)} \times (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \nearrow & \searrow p_{4(\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X))}^{\text{Sets}} & \\
 \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \downarrow \text{id}_{\mathcal{P}(X)} \times \cup & \downarrow \cap \times \cap & \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \mathcal{P}(X) \times \mathcal{P}(X), & \\
 \searrow \cap & \swarrow \cup & \\
 & \mathcal{P}(X) &
 \end{array}
 \quad
 \begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \times \Delta_{\mathcal{P}(X)} \nearrow & \searrow p_{4(\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X))}^{\text{Sets}} & \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \downarrow \cup \times \text{id}_{\mathcal{P}(X)} & \downarrow \cap \times \cap & \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \mathcal{P}(X) \times \mathcal{P}(X), & \\
 \searrow \cap & \swarrow \cup & \\
 & \mathcal{P}(X) &
 \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned}
 U \cap (V \cup W) &= (U \cap V) \cup (U \cap W), \\
 (U \cup V) \cap W &= (U \cap W) \cup (V \cap W)
 \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

003Q

## 8. Idempotency. The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X) & \xrightarrow{\Delta_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 & \searrow \text{id}_{\mathcal{P}(X)} & \downarrow \cup \\
 & & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cup U = U$$

for each  $U \in \mathcal{P}(X)$ .

003L

## 9. Via Intersections and Symmetric Differences. The diagram

$$\begin{array}{ccc}
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \xrightarrow{\Delta \times \cap} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \Delta_{\mathcal{P}(X) \times \mathcal{P}(X)} \nearrow & & \searrow \Delta \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\cup]{} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cup V = (U \Delta V) \Delta (U \cap V)$$

for each  $U, V \in \mathcal{P}(X)$ .

003S

10. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each  $U, V \in \mathcal{P}(X)$ .

003T

11. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each  $U, V \in \mathcal{P}(X)$ .

01H3

12. *Interaction With Direct Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U \cup V) = f_!(U) \cup f_!(V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01H4

13. *Interaction With Inverse Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cup V) = f^{-1}(U) \cup f^{-1}(V)$$

for each  $U, V \in \mathcal{P}(Y)$ .

01H5

14. *Interaction With Codirect Images.* Let  $f: X \rightarrow Y$  be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \downarrow \cup & \curvearrowright & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U) \cup f_*(V) \subset f_*(U \cup V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

- 003U** 15. *Interaction With Powersets and Semirings.* The quintuple  $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$  is an idempotent commutative semiring.

#### PROOF 4.3.8.1.3 ► PROOF OF PROPOSITION 4.3.8.1.2

Item 1: Functoriality

See [Pro25ar].

Item 2: Associativity

See [Pro25be].

Item 3: Unitality

This follows from [Pro25bh] and Item 4.

Item 4: Commutativity

See [Pro25bf].

Item 5: Annihilation With  $X$

We have

$$\begin{aligned} U \cup X &\stackrel{\text{def}}{=} \{x \in X \mid x \in U \text{ or } x \in X\} \\ &= \{x \in X \mid x \in X\}, \\ &= X \end{aligned}$$

and

$$\begin{aligned} X \cup V &\stackrel{\text{def}}{=} \{x \in X \mid x \in X \text{ or } x \in V\} \\ &= \{x \in X \mid x \in X\} \\ &= X. \end{aligned}$$

This finishes the proof.

Item 6: Distributivity of Unions Over Intersections

See [Pro25bd].

Item 7: Distributivity of Intersections Over Unions

See [Pro25an].

Item 8: Idempotency

See [Pro25aq].

Item 9: Via Intersections and Symmetric Differences

See [Pro25bc].

Item 10: Interaction With Characteristic Functions I

See [Pro25h].

Item 11: Interaction With Characteristic Functions II

See [Pro25h].

Item 12: Interaction With Direct Images

See [Pro25s].

Item 13: Interaction With Inverse Images

See [Pro25ac].

Item 14: Interaction With Codirect Images

This is a repetition of Item 5 of Proposition 4.6.3.1.7 and is proved there.

Item 15: Interaction With Powersets and Semirings

This follows from Items 2 to 4 and 8 of this proposition and Items 3 to 6 and 8 of Proposition 4.3.9.1.2. 

### 003X 4.3.9 Binary Intersections

Let  $X$  be a set and let  $U, V \in \mathcal{P}(X)$ .

003Y

#### DEFINITION 4.3.9.1.1 ► BINARY INTERSECTIONS

The **intersection of  $U$  and  $V$**  is the set  $U \cap V$  defined by

$$\begin{aligned} U \cap V &\stackrel{\text{def}}{=} \bigcap_{z \in \{U, V\}} z \\ &\stackrel{\text{def}}{=} \{x \in X \mid x \in U \text{ or } x \in V\}. \end{aligned}$$

003Z

#### PROPOSITION 4.3.9.1.2 ► PROPERTIES OF BINARY INTERSECTIONS

Let  $X$  be a set.

0040

1. *Functionality.* The assignments  $U, V, (U, V) \mapsto U \cap V$  define functors

$$\begin{aligned} U \cap -: & (\mathcal{P}(X), \subset) & \rightarrow (\mathcal{P}(X), \subset), \\ - \cap V: & (\mathcal{P}(X), \subset) & \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cap -_2: & (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) & \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each  $U, V, A, B \in \mathcal{P}(X)$ :

01H6 (a) If  $U \subset A$ , then  $U \cap V \subset A \cap V$ .

01H7 (b) If  $V \subset B$ , then  $U \cap V \subset U \cap B$ .

01H8 (c) If  $U \subset A$  and  $V \subset B$ , then  $U \cap V \subset A \cap B$ .

0041 2. *Adjointness.* We have adjunctions

$$(U \cap - \dashv [U, -]_X): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{\quad U \cap - \quad} \\ \perp \\ \xleftarrow{\quad [U, -]_X \quad} \end{array} \mathcal{P}(X),$$

$$(- \cap V \dashv [V, -]_X): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{\quad - \cap V \quad} \\ \perp \\ \xleftarrow{\quad [V, -]_X \quad} \end{array} \mathcal{P}(X),$$

witnessed by bijections

$$\text{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \text{Hom}_{\mathcal{P}(X)}(U, [V, W]_X),$$

$$\text{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \text{Hom}_{\mathcal{P}(X)}(V, [U, W]_X),$$

natural in  $U, V, W \in \mathcal{P}(X)$ , where

$$[-_1, -_2]_X: \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

is the bifunctor of [Section 4.4.7](#). In particular, the following statements hold for each  $U, V, W \in \mathcal{P}(X)$ :

01H9 (a) The following conditions are equivalent:

01HA i. We have  $U \cap V \subset W$ .

01HB ii. We have  $U \subset [V, W]_X$ .

01HC (b) The following conditions are equivalent:

01HD i. We have  $U \cap V \subset W$ .

01HE ii. We have  $V \subset [U, W]_X$ .

0042 3. *Associativity.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\ \alpha_{\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \swarrow & \nearrow \text{id}_{\mathcal{P}(X)} \times \cap & \\ (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cap \times \text{id}_{\mathcal{P}(X)} \searrow & & \nearrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\quad \cap \quad} & \mathcal{P}(X), \end{array}$$

commutes, i.e. we have an equality of sets

$$(U \cap V) \cap W = U \cap (V \cap W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

0043

4. *Unitality.* The diagrams

$$\begin{array}{ccc} \text{pt} \times \mathcal{P}(X) & \xrightarrow{[X] \times \text{id}_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \searrow \lambda_{\mathcal{P}(X)}^{\text{Sets}} \sim \curvearrowright & & \downarrow \cap \\ & & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(X) \times \text{pt} & \xrightarrow{\text{id}_{\mathcal{P}(X)} \times [X]} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \swarrow \rho_{\mathcal{P}(X)}^{\text{Sets}} \sim \curvearrowright & & \downarrow \cap \\ & & \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$X \cap U = U,$$

$$U \cap X = U$$

for each  $U \in \mathcal{P}(X)$ .

0044

5. *Commutativity.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\sigma_{\mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ & \searrow \cap & \downarrow \cap \\ & & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cap V = V \cap U$$

for each  $U, V \in \mathcal{P}(X)$ .

0047

6. *Annihilation With the Empty Set.* The diagrams

$$\begin{array}{ccccc} & \text{pt} \times \text{pt} & & \text{pt} \times \text{pt} & \\ \text{id}_{\text{pt}} \times \epsilon_{\mathcal{P}(X)}^{\text{Sets}} \nearrow & \swarrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} & & \epsilon_{\mathcal{P}(X)}^{\text{Sets}} \times \text{id}_{\text{pt}} \nearrow & \swarrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \\ \text{pt} \times \mathcal{P}(X) & & \text{pt} & & \text{pt} \\ \searrow [\emptyset] \times \text{id}_{\mathcal{P}(X)} & & \swarrow [\emptyset] & & \searrow [\emptyset] \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X) & & \mathcal{P}(X) \\ & & \text{id}_{\mathcal{P}(X)} \times [\emptyset] & & \\ & & \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$\emptyset \cap X = \emptyset,$$

$$X \cap \emptyset = \emptyset$$

for each  $U \in \mathcal{P}(X)$ .

01HF

## 7. Distributivity of Unions Over Intersections. The diagrams

$$\begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \Delta_{\mathcal{P}(X)} \times (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \nearrow & \searrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} & \\
 \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 & \downarrow \cup \cup & \\
 & \mathcal{P}(X) \times \mathcal{P}(X), & \\
 \text{id}_{\mathcal{P}(X)} \times \cap \downarrow & & \\
 \mathcal{P}(X) \times \mathcal{P}(X) & & \\
 & \swarrow \cup \quad \searrow \cap & \\
 & \mathcal{P}(X) & 
 \end{array}
 \quad
 \begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \times \Delta_{\mathcal{P}(X)} \nearrow & \searrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} & \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 & \downarrow \cap \times \text{id}_{\mathcal{P}(X)} & \\
 \mathcal{P}(X) \times \mathcal{P}(X) & & \\
 & \swarrow \cup \quad \searrow \cap & \\
 & \mathcal{P}(X) & \\
 & \downarrow \cup \cup & \\
 & \mathcal{P}(X) \times \mathcal{P}(X), & 
 \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned}
 U \cup (V \cap W) &= (U \cup V) \cap (U \cup W), \\
 (U \cap V) \cup W &= (U \cup W) \cap (V \cup W)
 \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

0046

## 8. Distributivity of Intersections Over Unions. The diagrams

$$\begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \Delta_{\mathcal{P}(X)} \times (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \nearrow & \searrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} & \\
 \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 & \downarrow \cap \cap & \\
 & \mathcal{P}(X) \times \mathcal{P}(X), & \\
 \text{id}_{\mathcal{P}(X)} \times \cup \downarrow & & \\
 \mathcal{P}(X) \times \mathcal{P}(X) & & \\
 & \swarrow \cap \quad \searrow \cup & \\
 & \mathcal{P}(X) & 
 \end{array}
 \quad
 \begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 (\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)}) \times \Delta_{\mathcal{P}(X)} \nearrow & \searrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} & \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 & \downarrow \cup \times \text{id}_{\mathcal{P}(X)} & \\
 \mathcal{P}(X) \times \mathcal{P}(X) & & \\
 & \swarrow \cap \quad \searrow \cup & \\
 & \mathcal{P}(X) & \\
 & \downarrow \cap \cap & \\
 & \mathcal{P}(X) \times \mathcal{P}(X), & 
 \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned}
 U \cap (V \cup W) &= (U \cap V) \cup (U \cap W), \\
 (U \cup V) \cap W &= (U \cap W) \cup (V \cap W)
 \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

0045

## 9. Idempotency. The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X) & \xrightarrow{\Delta_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 & \searrow \text{id}_{\mathcal{P}(X)} & \downarrow \cap \\
 & & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cap U = U$$

for each  $U \in \mathcal{P}(X)$ .

0048

10. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each  $U, V \in \mathcal{P}(X)$ .

0049

11. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01HG

12. *Interaction With Direct Images.* Let  $f: X \rightarrow Y$  be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & \curvearrowleft & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow[f_!]{} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U \cap V) \subset f_!(U) \cap f_!(V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01HH

13. *Interaction With Inverse Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow[f^{-1}]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cap V) = f^{-1}(U) \cap f^{-1}(V)$$

for each  $U, V \in \mathcal{P}(Y)$ .

01HJ

14. *Interaction With Codirect Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow[f_*]{} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U) \cap f_*(V) = f_*(U \cap V)$$

for each  $U, V \in \mathcal{P}(X)$ .

- 004A 15. *Interaction With Powersets and Monoids With Zero.* The quadruple  $((\mathcal{P}(X), \emptyset), \cap, X)$  is a commutative monoid with zero.
- 004B 16. *Interaction With Powersets and Semirings.* The quintuple  $((\mathcal{P}(X), \cup, \cap, \emptyset, X)$  is an idempotent commutative semiring.

#### PROOF 4.3.9.1.3 ► PROOF OF PROPOSITION 4.3.9.1.2

Item 1: Functoriality

See [Pro25ap].

Item 2: Adjointness

See [MSE 267469].

Item 3: Associativity

See [Pro25u].

Item 4: Unitality

This follows from [Pro25y] and Item 5.

Item 5: Commutativity

See [Pro25v].

Item 6: Annihilation With the Empty Set

This follows from [Pro25w] and Item 5.

Item 7: Distributivity of Unions Over Intersections

See [Pro25bd].

Item 8: Distributivity of Intersections Over Unions

See [Pro25an].

Item 9: Idempotency

See [Pro25ao].

Item 10: Interaction With Characteristic Functions I

See [Pro25e].

Item 11: Interaction With Characteristic Functions II

See [Pro25e].

Item 12: Interaction With Direct Images

See [Pro25q].

Item 13: Interaction With Inverse Images

See [Pro25aa].

Item 14: Interaction With Codirect Images

This is a repetition of [Item 6 of Proposition 4.6.3.1.7](#) and is proved there.

Item 15: Interaction With Powersets and Monoids With Zero

This follows from [Items 3 to 6](#).

Item 16: Interaction With Powersets and Semirings

This follows from [Items 2 to 4](#) and [8](#) and [Items 3 to 6](#) and [8](#) of [Proposition 4.3.9.1.2](#). 

#### 004D 4.3.10 Differences

Let  $X$  and  $Y$  be sets.

004E

##### DEFINITION 4.3.10.1.1 ► DIFFERENCES

The **difference of  $X$  and  $Y$**  is the set  $X \setminus Y$  defined by

$$X \setminus Y \stackrel{\text{def}}{=} \{a \in X \mid a \notin Y\}.$$

004F

##### PROPOSITION 4.3.10.1.2 ► PROPERTIES OF DIFFERENCES

Let  $X$  be a set.

004G

- Functoriality.* The assignments  $U, V, (U, V) \mapsto U \cap V$  define functors

$$\begin{aligned} U \setminus - &: (\mathcal{P}(X), \supset) \rightarrow (\mathcal{P}(X), \subset), \\ - \setminus V &: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \setminus -_2 &: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each  $U, V, A, B \in \mathcal{P}(X)$ :

01HK

(a) If  $U \subset A$ , then  $U \setminus V \subset A \setminus V$ .

01HL

(b) If  $V \subset B$ , then  $U \setminus B \subset U \setminus V$ .

01HM

(c) If  $U \subset A$  and  $V \subset B$ , then  $U \setminus B \subset A \setminus V$ .

004H

- De Morgan's Laws.* We have equalities of sets

$$X \setminus (U \cup V) = (X \setminus U) \cap (X \setminus V),$$

$$X \setminus (U \cap V) = (X \setminus U) \cup (X \setminus V)$$

for each  $U, V \in \mathcal{P}(X)$ .

**004J** 3. *Interaction With Unions I.* We have equalities of sets

$$U \setminus (V \cup W) = (U \setminus V) \cap (U \setminus W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

**004K** 4. *Interaction With Unions II.* We have equalities of sets

$$(U \setminus V) \cup W = (U \cup W) \setminus (V \setminus W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

**004L** 5. *Interaction With Unions III.* We have equalities of sets

$$\begin{aligned} U \setminus (V \cup W) &= (U \cup W) \setminus (V \cup W) \\ &= (U \setminus V) \setminus W \\ &= (U \setminus W) \setminus V \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

**004M** 6. *Interaction With Unions IV.* We have equalities of sets

$$(U \cup V) \setminus W = (U \setminus W) \cup (V \setminus W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

**004N** 7. *Interaction With Intersections.* We have equalities of sets

$$\begin{aligned} (U \setminus V) \cap W &= (U \cap W) \setminus V \\ &= U \cap (W \setminus V) \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

**004P** 8. *Interaction With Complements.* We have an equality of sets

$$U \setminus V = U \cap V^c$$

for each  $U, V \in \mathcal{P}(X)$ .

**004Q** 9. *Interaction With Symmetric Differences.* We have an equality of sets

$$U \setminus V = U \Delta (U \cap V)$$

for each  $U, V \in \mathcal{P}(X)$ .

004R

10. *Triple Differences.* We have

$$U \setminus (V \setminus W) = (U \cap W) \cup (U \setminus V)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

004S

11. *Left Annihilation.* We have

$$\emptyset \setminus U = \emptyset$$

for each  $U \in \mathcal{P}(X)$ .

004T

12. *Right Unitality.* We have

$$U \setminus \emptyset = U$$

for each  $U \in \mathcal{P}(X)$ .

01HN

13. *Right Annihilation.* We have

$$U \setminus X = \emptyset$$

for each  $U \in \mathcal{P}(X)$ .

004U

14. *Invertibility.* We have

$$U \setminus U = \emptyset$$

for each  $U \in \mathcal{P}(X)$ .

004V

15. *Interaction With Containment.* The following conditions are equivalent:

004W

(a) We have  $V \setminus U \subset W$ .

004X

(b) We have  $V \setminus W \subset U$ .

004Y

16. *Interaction With Characteristic Functions.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each  $U, V \in \mathcal{P}(X)$ .

01HP

17. *Interaction With Direct Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow & \curvearrowright & \downarrow \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \setminus f_!(V) \subset f_!(U \setminus V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01HQ

18. *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \downarrow \backslash & & \downarrow \backslash \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \setminus V) = f^{-1}(U) \setminus f^{-1}(V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01HR

19. *Interaction With Codirect Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow \backslash & \curvearrowright & \downarrow \backslash \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \setminus f_!(V) \subset f_!(U \setminus V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

#### PROOF 4.3.10.1.3 ► PROOF OF PROPOSITION 4.3.10.1.2

Item 1: Functoriality

See [Pro25ah] and [Pro25al].

Item 2: De Morgan's Laws

See [Pro25n].

**Item 3: Interaction With Unions I**

See [Pro25o].

**Item 4: Interaction With Unions II**

We have

$$\begin{aligned}
 (U \setminus V) \cup W &\stackrel{\text{def}}{=} \{x \in X \mid (x \in U \text{ and } x \notin V) \text{ or } x \in W\} \\
 &= \{x \in X \mid (x \in U \text{ or } x \in W) \text{ and } (x \notin V \text{ or } x \in W)\} \\
 &= \{x \in X \mid (x \in U \cup W) \text{ and not } (x \in V \text{ and } x \notin W)\} \\
 &= \{x \in X \mid (x \in U \cup W) \text{ and not } (x \in V \setminus W)\} \\
 &= \{x \in X \mid (x \in (U \cup W) \setminus (V \setminus W))\} \\
 &= (U \cup W) \setminus (V \setminus W).
 \end{aligned}$$

**Item 5: Interaction With Unions III**

See [Pro25am].

**Item 6: Interaction With Unions IV**

See [Pro25ag].

**Item 7: Interaction With Intersections**

See [Pro25x].

**Item 8: Interaction With Complements**

See [Pro25ae].

**Item 9: Interaction With Symmetric Differences**

See [Pro25af].

**Item 10: Triple Differences**

See [Pro25ak].

**Item 11: Left Annihilation**

The direction  $\emptyset \subset \emptyset \setminus U$  always holds. Now assume  $x \in \emptyset \setminus U$ . Then,  $x \in \emptyset$  and  $x \notin U$ . Hence  $\emptyset \setminus U \subset \emptyset$  must hold and the sets are equal.

**Item 12: Right Unitality**

See [Pro25ai].

**Item 13: Right Annihilation**

It suffices to show that no  $x \in X$  can be an element of  $U \setminus X$ . Assume  $x \in U \setminus X$ . Then  $x \notin X$ , contradicting  $x \in X$ . This completes the proof.

**Item 14: Invertibility**

See [Pro25aj].

**Item 15: Interaction With Containment**

The conditions are symmetric in  $U, W$ , hence it suffices to show that  $V \setminus U \subset W$  implies  $V \setminus W \subset U$ . So assume  $V \setminus U \subset W, x \in V \setminus W$ . Then  $x \in V, x \notin W$ . So by contraposition,  $x \notin V \setminus U$ . But  $x \in V$ , so we must have  $x \in U$ , completing the proof.

**Item 16: Interaction With Characteristic Functions**

See [Pro25f].

**Item 17: Interaction With Direct Images**

See [Pro25r].

**Item 18: Interaction With Inverse Images**

See [Pro25ab].

**004Z 4.3.11 Complements**

Let  $X$  be a set and let  $U \in \mathcal{P}(X)$ .

0050

**DEFINITION 4.3.11.1 ► COMPLEMENTS**

The **complement** of  $U$  is the set  $U^c$  defined by

$$\begin{aligned} U^c &\stackrel{\text{def}}{=} X \setminus U \\ &\stackrel{\text{def}}{=} \{a \in X \mid a \notin U\}. \end{aligned}$$

0051

**PROPOSITION 4.3.11.2 ► PROPERTIES OF COMPLEMENTS**

Let  $X$  be a set.

0052

1. *Functoriality.* The assignment  $U \mapsto U^c$  defines a functor

$$(-)^c : \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X).$$

In particular, the following statements hold for each  $U, V \in \mathcal{P}(X)$ :

(★) If  $U \subset V$ , then  $V^c \subset U^c$ .

0053

2. *De Morgan's Laws.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} & \xrightarrow{\cup^{\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ (-)^c \times (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} & \xrightarrow{\cap^{\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ (-)^c \times (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned}(U \cup V)^c &= U^c \cap V^c, \\ (U \cap V)^c &= U^c \cup V^c\end{aligned}$$

for each  $U, V \in \mathcal{P}(X)$ .

0054

3. *Involutority.* The diagram

$$\begin{array}{ccc}\mathcal{P}(X)^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(X) \\ & \searrow \text{id}_{\mathcal{P}(X)^{\text{op}}} & \downarrow (-)^{c,\text{op}} \\ & & \mathcal{P}(X)^{\text{op}}\end{array}$$

commutes, i.e. we have

$$(U^c)^c = U$$

for each  $U \in \mathcal{P}(X)$ .

0055

4. *Interaction With Characteristic Functions.* We have

$$\chi_{U^c} = 1 - \chi_U$$

for each  $U \in \mathcal{P}(X)$ .

01HS

5. *Interaction With Direct Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc}\mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)\end{array}$$

commutes, i.e. we have

$$f_!(U^c) = f_*(U)^c$$

for each  $U \in \mathcal{P}(X)$ .

01HT

6. *Interaction With Inverse Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc}\mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1,\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)\end{array}$$

commutes, i.e. we have

$$f^{-1}(U^c) = f^{-1}(U)^c$$

for each  $U \in \mathcal{P}(X)$ .

01HU

7. *Interaction With Codirect Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U^c) = f_!(U)^c$$

for each  $U \in \mathcal{P}(X)$ .

#### PROOF 4.3.11.1.3 ► PROOF OF PROPOSITION 4.3.11.1.2

Item 1: Functoriality

This follows from [Item 1 of Proposition 4.3.10.1.2](#).

Item 2: De Morgan's Laws

See [\[Pro25n\]](#).

Item 3: Involutority

See [\[Pro25i\]](#).

Item 4: Interaction With Characteristic Functions

We consider the two cases  $x \in U, x \notin U$ .

1. If  $x \in U$ , then  $x \notin U^c$ . So  $\chi_U(x) = 1$  and

$$\begin{aligned} \chi_{U^c}(x) &= 0 \\ &= 1 - \chi_U(x). \end{aligned}$$

2. If  $x \notin U$ , then  $x \in U^c$ . So  $\chi_U(x) = 0$  and

$$\begin{aligned} \chi_{U^c}(x) &= 1 \\ &= 1 - \chi_U(x). \end{aligned}$$

Hence, the equation holds for all  $x \in X$ .

**Item 5: Interaction With Direct Images**

This is a repetition of **Item 8** of [Proposition 4.6.1.1.5](#) and is proved there.

**Item 6: Interaction With Inverse Images**

This is a repetition of **Item 8** of [Proposition 4.6.2.1.3](#) and is proved there.

**Item 7: Interaction With Codirect Images**

This is a repetition of **Item 7** of [Proposition 4.6.3.1.7](#) and is proved there.



### 0056 4.3.12 Symmetric Differences

Let  $X$  be a set and let  $U, V \in \mathcal{P}(X)$ .

0057

#### DEFINITION 4.3.12.1.1 ► SYMMETRIC DIFFERENCES

The **symmetric difference of  $U$  and  $V$**  is the set  $U \Delta V$  defined by<sup>1</sup>

$$U \Delta V \stackrel{\text{def}}{=} (U \setminus V) \cup (V \setminus U).$$

<sup>1</sup>Illustration:



0058

#### PROPOSITION 4.3.12.1.2 ► PROPERTIES OF SYMMETRIC DIFFERENCES

Let  $X$  be a set.

0059

1. *Lack of Functoriality.* The assignment  $(U, V) \mapsto U \Delta V$  **does not** in general define functors


$$\begin{aligned} U \Delta -: & (\mathcal{P}(X), \subset) & \rightarrow (\mathcal{P}(X), \subset), \\ - \Delta V: & (\mathcal{P}(X), \subset) & \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \Delta -_2: & (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) & \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

005A

2. *Via Unions and Intersections.* We have

$$U \Delta V = (U \cup V) \setminus (U \cap V)$$

for each  $U, V \in \mathcal{P}(X)$ , as in the Venn diagram



01HV

3. *Symmetric Differences of Disjoint Sets.* If  $U$  and  $V$  are disjoint, then we have

$$U \Delta V = U \cup V.$$

005B


4. *Associativity.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 & \alpha_{\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \curvearrowleft & \searrow \text{id}_{\mathcal{P}(X)} \times \Delta \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \downarrow \Delta \times \text{id}_{\mathcal{P}(X)} & & \downarrow \Delta \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\Delta]{} & \mathcal{P}(X),
 \end{array}$$

commutes, i.e. we have

$$(U \Delta V) \Delta W = U \Delta (V \Delta W)$$

for each  $U, V, W \in \mathcal{P}(X)$ , as in the Venn diagram



005D

5. *Unitality.* The diagrams

$$\begin{array}{ccc}
 \text{pt} \times \mathcal{P}(X) & \xrightarrow{[\emptyset] \times \text{id}_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \downarrow \lambda_{\mathcal{P}(X)}^{\text{Sets}} \curvearrowright & & \downarrow \Delta \\
 & & \mathcal{P}(X)
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{P}(X) \times \text{pt} & \xrightarrow{\text{id}_{\mathcal{P}(X)} \times [\emptyset]} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \downarrow \rho_{\mathcal{P}(X)}^{\text{Sets}} \curvearrowright & & \downarrow \Delta \\
 & & \mathcal{P}(X)
 \end{array}$$

commute, i.e. we have

$$\begin{aligned}
 U \Delta \emptyset &= U, \\
 \emptyset \Delta U &= U
 \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

005C

6. *Commutativity.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\sigma_{\mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \downarrow \Delta & & \downarrow \Delta \\
 & & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$U \Delta V = V \Delta U$$

for each  $U, V \in \mathcal{P}(X)$ .

**005E** 7. *Invertibility.* We have

$$U \Delta U = \emptyset$$

for each  $U \in \mathcal{P}(X)$ .

**005F** 8. *Interaction With Unions.* We have

$$(U \Delta V) \cup (V \Delta W) = (U \cup V \cup W) \setminus (U \cap V \cap W)$$

for each  $U, V, W \in \mathcal{P}(X)$ .

**005G** 9. *Interaction With Complements I.* We have

$$U \Delta U^c = X$$

for each  $U \in \mathcal{P}(X)$ .

**005H** 10. *Interaction With Complements II.* We have

$$\begin{aligned} U \Delta X &= U^c, \\ X \Delta U &= U^c \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

**005J** 11. *Interaction With Complements III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\Delta} & \mathcal{P}(X) \\ (-)^c \times (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\Delta} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$U^c \Delta V^c = U \Delta V$$

for each  $U, V \in \mathcal{P}(X)$ .

**005K** 12. “*Transitivity*”. We have

$$(U \Delta V) \Delta (V \Delta W) = U \Delta W$$

for each  $U, V, W \in \mathcal{P}(X)$ .

005L

13. *The Triangle Inequality for Symmetric Differences.* We have

$$U \Delta W \subset U \Delta V \cup V \Delta W$$

for each  $U, V, W \in \mathcal{P}(X)$ .

005M

14. *Distributivity Over Intersections.* We have

$$\begin{aligned} U \cap (V \Delta W) &= (U \cap V) \Delta (U \cap W), \\ (U \Delta V) \cap W &= (U \cap W) \Delta (V \cap W) \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

005N

15. *Interaction With Characteristic Functions.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each  $U, V \in \mathcal{P}(X)$ .

005P

16. *Bijectivity.* Given  $U, V \in \mathcal{P}(X)$ , the maps

$$\begin{aligned} U \Delta -: \mathcal{P}(X) &\rightarrow \mathcal{P}(X), \\ - \Delta V: \mathcal{P}(X) &\rightarrow \mathcal{P}(X) \end{aligned}$$

are self-inverse bijections. Moreover, the map

$$\begin{aligned} \mathcal{P}(X) &\longrightarrow \mathcal{P}(X) \\ C &\longmapsto C \Delta (U \Delta V) \end{aligned}$$

is a bijection of  $\mathcal{P}(X)$  onto itself sending  $U$  to  $V$  and  $V$  to  $U$ .

005Q

17. *Interaction With Powersets and Groups.* Let  $X$  be a set.

005R

(a) The quadruple  $(\mathcal{P}(X), \Delta, \emptyset, \text{id}_{\mathcal{P}(X)})$  is an abelian group.<sup>1</sup>

005S

(b) Every element of  $\mathcal{P}(X)$  has order 2 with respect to  $\Delta$ , and thus  $\mathcal{P}(X)$  is a *Boolean group* (i.e. an abelian 2-group).

005T

18. *Interaction With Powersets and Vector Spaces I.* The pair  $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$  consisting of

- The group  $\mathcal{P}(X)$  of Item 17;

- The map  $\alpha_{\mathcal{P}(X)} : \mathbb{F}_2 \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$  defined by

$$\begin{aligned} 0 \cdot U &\stackrel{\text{def}}{=} \emptyset, \\ 1 \cdot U &\stackrel{\text{def}}{=} U; \end{aligned}$$

is an  $\mathbb{F}_2$ -vector space.

**005U** 19. *Interaction With Powersets and Vector Spaces II.* If  $X$  is finite, then:

**020L** (a) The set of singletons sets on the elements of  $X$  forms a basis for the  $\mathbb{F}_2$ -vector space  $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$  of [Item 18](#).

**020M** (b) We have

$$\dim(\mathcal{P}(X)) = \#X.$$

**005V** 20. *Interaction With Powersets and Rings.* The quintuple  $(\mathcal{P}(X), \Delta, \cap, \emptyset, X)$  is a commutative ring.<sup>2</sup>

**01HW** 21. *Interaction With Direct Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow \Delta & \curvearrowright & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \Delta f_!(V) \subset f_!(U \Delta V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

**01HX** 22. *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \downarrow \Delta & & \downarrow \Delta \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

i.e. we have

$$f^{-1}(U) \Delta f^{-1}(V) = f^{-1}(U \Delta V)$$

for each  $U, V \in \mathcal{P}(Y)$ .

01HY

23. *Interaction With Codirect Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \curvearrowright & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U \Delta V) \subset f_*(U) \Delta f_*(V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

<sup>1</sup>Here are some examples:

020H

- i. When  $X = \emptyset$ , we have an isomorphism of groups between  $\mathcal{P}(\emptyset)$  and the trivial group:

$$(\mathcal{P}(\emptyset), \Delta, \emptyset, \text{id}_{\mathcal{P}(\emptyset)}) \cong \text{pt}.$$

020J

- ii. When  $X = \text{pt}$ , we have an isomorphism of groups between  $\mathcal{P}(\text{pt})$  and  $\mathbb{Z}/2$ :

$$(\mathcal{P}(\text{pt}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\text{pt})}) \cong \mathbb{Z}/2.$$

020K

- iii. When  $X = \{0, 1\}$ , we have an isomorphism of groups between  $\mathcal{P}(\{0, 1\})$  and  $\mathbb{Z}/2 \times \mathbb{Z}/2$ :

$$(\mathcal{P}(\{0, 1\}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\{0, 1\})}) \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

<sup>2</sup> Warning:

The analogous statement replacing intersections by unions (i.e. that the quintuple  $(\mathcal{P}(X), \Delta, \cup, \emptyset, X)$  is a ring) is false, however. See [Pro25ba] for a proof.

#### PROOF 4.3.12.1.3 ► PROOF OF PROPOSITION 4.3.12.1.2

##### Item 1: Lack of Functoriality

Let  $X = \{0, 1\}$ ,  $U = \{0\}$ . Then  $\emptyset \subset U$ , but  $U \Delta \emptyset = U \not\subset \emptyset = U \Delta U$  from Item 5 and Item 7. This gives a counterexample to the first statement. By using Item 6, we can adapt it to the second and third statement.

##### Item 2: Via Unions and Intersections

See [Pro25p].

##### Item 3: Symmetric Differences of Disjoint Sets

Since  $U$  and  $V$  are disjoint, we have  $U \cap V = \emptyset$ , and therefore we have

$$\begin{aligned} U \Delta V &= (U \cup V) \setminus (U \cap V) \\ &= (U \cup V) \setminus \emptyset \\ &= U \cup V, \end{aligned}$$

where we've used [Item 2](#) and [Item 12](#) of Proposition 4.3.10.1.2.

**Item 4:** Associativity

See [[Pro25as](#)].

**Item 5:** Unitality

This follows from [Item 6](#) and [[Pro25ax](#)].

**Item 6:** Commutativity

See [[Pro25at](#)].

**Item 7:** Invertibility

See [[Pro25az](#)].

**Item 8:** Interaction With Unions

See [[Pro25bg](#)].

**Item 9:** Interaction With Complements I

See [[Pro25aw](#)].

**Item 10:** Interaction With Complements II

This follows from [Item 6](#) and [[Pro25bb](#)].

**Item 11:** Interaction With Complements III

See [[Pro25au](#)].

**Item 12:** “Transitivity”

We have

$$\begin{aligned}
 (U \Delta V) \Delta (V \Delta W) &= U \Delta (V \Delta (V \Delta W)) && (\text{by } \text{Item 4}) \\
 &= U \Delta ((V \Delta V) \Delta W) && (\text{by } \text{Item 4}) \\
 &= U \Delta (\emptyset \Delta W) && (\text{by } \text{Item 7}) \\
 &= U \Delta W. && (\text{by } \text{Item 5})
 \end{aligned}$$

This finishes the proof.

**Item 13:** The Triangle Inequality for Symmetric Differences

This follows from [Items 2](#) and [12](#).

**Item 14:** Distributivity Over Intersections

See [[Pro25t](#)].

**Item 15:** Interaction With Characteristic Functions

See [[Pro25g](#)].

**Item 16:** Bijectivity

- We show that

$$(U \Delta -) : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

is self-inverse.

Let  $W \in \mathcal{P}(X)$ . Then,

$$\begin{aligned} U \Delta (U \Delta W) &= (U \Delta U) \Delta W && (\text{by Item 4}) \\ &= \emptyset \Delta W && (\text{by Item 7}) \\ &= W. && (\text{by Item 5}) \end{aligned}$$

- By Item 6,  $(- \Delta V) = (V \Delta -)$ , hence the former is also self-inverse by the first point.
- The map  $- \Delta (U \Delta V)$  is a bijection as a special case of the second point. From the first two points and Item 6, we get

$$U \Delta (U \Delta V) = V, \quad V \Delta (U \Delta V) = V \Delta (V \Delta U) = U.$$

Hence the function maps  $U$  to  $V$  and  $V$  to  $U$ .

**Item 17: Interaction With Powersets and Groups**

Item 17a follows from Items 4 to 7, while Item 17b follows from Item 7.<sup>1</sup>

**Item 18: Interaction With Powersets and Vector Spaces I**

See [MSE 2719059].

**Item 19: Interaction With Powersets and Vector Spaces II**

See [MSE 2719059].

**Item 20: Interaction With Powersets and Rings**

This follows from Items 6 and 15 of Proposition 4.3.9.1.2 and Items 14 and 17.<sup>2</sup>

**Item 21: Interaction With Direct Images**

This is a repetition of Item 9 of Proposition 4.6.1.1.5 and is proved there.

**Item 22: Interaction With Inverse Images**

This is a repetition of Item 9 of Proposition 4.6.2.1.3 and is proved there.

**Item 23: Interaction With Codirect Images**

This is a repetition of Item 8 of Proposition 4.6.3.1.7 and is proved there.



<sup>1</sup>Reference: [Pro25av].

<sup>2</sup>Reference: [Pro25ay].

**005W 4.4 Powersets****01HZ 4.4.1 Foundations**

Let  $X$  be a set.

006P

**DEFINITION 4.4.1.1.1 ► POWERSETS**

The **powerset** of  $X$  is the set  $\mathcal{P}(X)$  defined by

$$\mathcal{P}(X) \stackrel{\text{def}}{=} \{U \in P \mid U \subset X\},$$

where  $P$  is the set in the axiom of powerset, ?? of ??.

006Q

**REMARK 4.4.1.1.2 ► POWERSETS AS DECATEGORIFICATIONS OF CO/PRESHEAF CATEGORIES**

Under the analogy that  $\{t, f\}$  should be the  $(-1)$ -categorical analogue of Sets, we may view the powerset of a set as a decategorification of the category of presheaves of a category (or of the category of copresheaves):

- The powerset of a set  $X$  is equivalently (Item 2 of Proposition 4.5.1.1.4) the set

$$\text{Sets}(X, \{t, f\})$$

of functions from  $X$  to the set  $\{t, f\}$  of classical truth values.

- The category of presheaves on a category  $C$  is the category

$$\text{Fun}(C^{\text{op}}, \text{Sets})$$

of functors from  $C^{\text{op}}$  to the category Sets of sets.

01J0

**NOTATION 4.4.1.1.3 ► FURTHER NOTATION FOR POWERSETS**

Let  $X$  be a set.

01J1

1. We write  $\mathcal{P}_0(X)$  for the set of nonempty subsets of  $X$ .

01J2

2. We write  $\mathcal{P}_{\text{fin}}(X)$  for the set of finite subsets of  $X$ .

01J3

**PROPOSITION 4.4.1.1.4 ► ELEMENTARY PROPERTIES OF POWERSETS**

Let  $X$  be a set.

01J4

1. *Co/Completeness.* The (posetal) category (associated to)  $(\mathcal{P}(X), \subset)$  is complete and cocomplete:

020P

(a) *Products.* The products in  $\mathcal{P}(X)$  are given by intersection of subsets.

020Q

(b) *Coproducts.* The coproducts in  $\mathcal{P}(X)$  are given by union of subsets.

020R

(c) *Co/Equalisers.* Being a posetal category,  $\mathcal{P}(X)$  only has at most one morphisms between any two objects, so co/equalisers are trivial.

01J5

2. *Cartesian Closedness.* The category  $\mathcal{P}(X)$  is Cartesian closed.

01J6

3. *Powersets as Sets of Relations.* We have bijections

$$\begin{aligned}\mathcal{P}(X) &\cong \text{Rel}(\text{pt}, X), \\ \mathcal{P}(X) &\cong \text{Rel}(X, \text{pt}),\end{aligned}$$

natural in  $X \in \text{Obj}(\text{Sets})$ .

01J7

4. *Interaction With Products I.* The map

$$\begin{aligned}\mathcal{P}(X) \times \mathcal{P}(Y) &\longrightarrow \mathcal{P}(X \sqcup Y) \\ (U, V) &\longmapsto U \cup V\end{aligned}$$

is an isomorphism of sets, natural in  $X, Y \in \text{Obj}(\text{Sets})$  with respect to each of the functor structures  $\mathcal{P}_!$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  on  $\mathcal{P}$  of [Proposition 4.4.2.1.1](#). Moreover, this makes each of  $\mathcal{P}_!$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  into a symmetric monoidal functor.

01J8

5. *Interaction With Products II.* The map

$$\begin{aligned}\mathcal{P}(X) \times \mathcal{P}(Y) &\longrightarrow \mathcal{P}(X \sqcup Y) \\ (U, V) &\longmapsto U \boxtimes_{X \times Y} V,\end{aligned}$$

where<sup>1</sup>

$$U \boxtimes_{X \times Y} V \stackrel{\text{def}}{=} \{(u, v) \in X \times Y \mid u \in U \text{ and } v \in V\}$$

is an inclusion of sets, natural in  $X, Y \in \text{Obj}(\text{Sets})$  with respect to each of the functor structures  $\mathcal{P}_!$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  on  $\mathcal{P}$  of [Proposition 4.4.2.1.1](#). Moreover, this makes each of  $\mathcal{P}_!$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  into a symmetric monoidal functor.

01J9

6. *Interaction With Products III.* We have an isomorphism

$$\mathcal{P}(X) \otimes \mathcal{P}(Y) \cong \mathcal{P}(X \times Y),$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$  with respect to each of the functor structures  $\mathcal{P}_!$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  on  $\mathcal{P}$  of [Proposition 4.4.2.1.1](#), where  $\otimes$  denotes the tensor product of suplattices of  $\mathbb{P}$ . Moreover, this makes each of  $\mathcal{P}_!$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  into a symmetric monoidal functor.

<sup>1</sup>The set  $U \boxtimes_{X \times Y} V$  is usually denoted simply  $U \times V$ . Here we denote it in this somewhat weird way to highlight the similarity to external tensor products in six-functor formalisms (see also [Section 4.6.4](#)).

#### PROOF 4.4.1.1.5 ► PROOF OF PROPOSITION 4.4.1.1.4

Item 1: Co/Completeness

Omitted.

Item 2: Cartesian Closedness

See [Section 4.4.7](#).

Item 3: Powersets as Sets of Relations

Indeed, we have

$$\begin{aligned}\text{Rel}(\text{pt}, X) &\stackrel{\text{def}}{=} \mathcal{P}(\text{pt} \times X) \\ &\cong \mathcal{P}(X)\end{aligned}$$

and

$$\begin{aligned}\text{Rel}(X, \text{pt}) &\stackrel{\text{def}}{=} \mathcal{P}(X \times \text{pt}) \\ &\cong \mathcal{P}(X),\end{aligned}$$

where we have used [Item 5 of Proposition 4.1.3.1.4](#).

Item 4: Interaction With Products I

The inverse of the map in the statement is the map

$$\Phi: \mathcal{P}(X \sqcup Y) \rightarrow \mathcal{P}(X) \times \mathcal{P}(Y)$$

defined by

$$\Phi(S) \stackrel{\text{def}}{=} (S_X, S_Y)$$

for each  $S \in \mathcal{P}(X \sqcup Y)$ , where

$$\begin{aligned}S_X &\stackrel{\text{def}}{=} \{x \in X \mid (0, x) \in S\} \\ S_Y &\stackrel{\text{def}}{=} \{y \in Y \mid (1, y) \in S\}.\end{aligned}$$

The rest of the proof is omitted.

Item 5: Interaction With Products II

Omitted.

Item 6: Interaction With Products III

Omitted.



### 01JA 4.4.2 Functoriality of Powersets

01JB

#### PROPOSITION 4.4.2.1.1 ► FUNCTORIALITY OF POWERSETS

Let  $X$  be a set.

01JC

1. *Functoriality I.* The assignment  $X \mapsto \mathcal{P}(X)$  defines a functor

$$\mathcal{P}_! : \text{Sets} \rightarrow \text{Sets},$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$\mathcal{P}_!(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(\text{Sets})$ , the action on morphisms

$$\mathcal{P}_{*|A,B} : \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B))$$

of  $\mathcal{P}_!$  at  $(A, B)$  is the map defined by sending a map of sets  $f : A \rightarrow B$  to the map

$$\mathcal{P}_!(f) : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\mathcal{P}_!(f) \stackrel{\text{def}}{=} f_!,$$

as in [Definition 4.6.1.1.1](#).

01JD

2. *Functoriality II.* The assignment  $X \mapsto \mathcal{P}(X)$  defines a functor

$$\mathcal{P}^{-1} : \text{Sets}^{\text{op}} \rightarrow \text{Sets},$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$\mathcal{P}^{-1}(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(\text{Sets})$ , the action on morphisms

$$\mathcal{P}_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(B), \mathcal{P}(A))$$

of  $\mathcal{P}^{-1}$  at  $(A, B)$  is the map defined by sending a map of sets  $f: A \rightarrow B$  to the map

$$\mathcal{P}^{-1}(f): \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by

$$\mathcal{P}^{-1}(f) \stackrel{\text{def}}{=} f^{-1},$$

as in [Definition 4.6.2.1.1](#).

01JE

3. *Functionality III.* The assignment  $X \mapsto \mathcal{P}(X)$  defines a functor

$$\mathcal{P}_*: \text{Sets} \rightarrow \text{Sets},$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$\mathcal{P}_*(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(\text{Sets})$ , the action on morphisms

$$\mathcal{P}_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B))$$

of  $\mathcal{P}_*$  at  $(A, B)$  is the map defined by sending a map of sets  $f: A \rightarrow B$  to the map

$$\mathcal{P}_*(f): \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\mathcal{P}_*(f) \stackrel{\text{def}}{=} f_*,$$

as in [Definition 4.6.3.1.1](#).

#### PROOF 4.4.2.1.2 ► PROOF OF PROPOSITION 4.4.2.1.1

Item 1: Functoriality I

This follows from [Items 3 and 4 of Proposition 4.6.1.1.7](#).

Item 2: Functoriality II

This follows from [Items 3 and 4 of Proposition 4.6.2.1.5](#).

Item 3: Functoriality III

This follows from [Items 3 and 4 of Proposition 4.6.3.1.9](#). 

### 01JF 4.4.3 Adjointness of Powersets I

01JG

#### PROPOSITION 4.4.3.1.1 ► ADJOINTNESS OF POWERSETS I

We have an adjunction

$$\left( \mathcal{P}^{-1} \dashv \mathcal{P}^{-1,\text{op}} \right): \quad \text{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{\mathcal{P}^{-1}} \\ \perp \\ \xleftarrow{\mathcal{P}^{-1,\text{op}}} \end{array} \text{Sets},$$

witnessed by a bijection

$$\underbrace{\text{Sets}^{\text{op}}(\mathcal{P}(X), Y)}_{\stackrel{\text{def}}{=} \text{Sets}(Y, \mathcal{P}(X))} \cong \text{Sets}(X, \mathcal{P}(Y)),$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $Y \in \text{Obj}(\text{Sets}^{\text{op}})$ .

#### PROOF 4.4.3.1.2 ► PROOF OF PROPOSITION 4.4.3.1.1

We have

$$\begin{aligned} \text{Sets}^{\text{op}}(\mathcal{P}(A), B) &\stackrel{\text{def}}{=} \text{Sets}(B, \mathcal{P}(A)) \\ &\cong \text{Sets}(B, \text{Sets}(A, \{t, f\})) \quad (\text{by Item 2 of Proposition 4.5.1.1.4}) \\ &\cong \text{Sets}(A \times B, \{t, f\}) \quad (\text{by Item 2 of Proposition 4.1.3.1.4}) \\ &\cong \text{Sets}(A, \text{Sets}(B, \{t, f\})) \quad (\text{by Item 2 of Proposition 4.1.3.1.4}) \\ &\cong \text{Sets}(A, \mathcal{P}(B)), \quad (\text{by Item 2 of Proposition 4.5.1.1.4}) \end{aligned}$$

where all bijections are natural in  $A$  and  $B$ .<sup>1</sup> 

<sup>1</sup>Here we are using [Item 3 of Proposition 4.5.1.1.4](#).

**01JH 4.4.4 Adjointness of Powersets II**

01JJ

**PROPOSITION 4.4.4.1.1 ► ADJOINTNESS OF POWERSETS II**

We have an adjunction

$$(Gr \dashv \mathcal{P}_!): \text{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_!} \end{array} \text{Rel},$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(X), Y) \cong \text{Sets}(X, \mathcal{P}(Y))$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $Y \in \text{Obj}(\text{Rel})$ , where  $\text{Gr}$  is the graph functor of Item 1 of Proposition 8.2.2.1.2 and  $\mathcal{P}_!$  is the functor of Proposition 8.7.5.1.1.

**PROOF 4.4.4.1.2 ► PROOF OF PROPOSITION 4.4.4.1.1**

We have

$$\begin{aligned} \text{Rel}(\text{Gr}(A), B) &\cong \mathcal{P}(A \times B) \\ &\cong \text{Sets}(A \times B, \{t, f\}) \quad (\text{by Item 2 of Proposition 4.5.1.1.4}) \\ &\cong \text{Sets}(A, \text{Sets}(B, \{t, f\})) \quad (\text{by Item 2 of Proposition 4.1.3.1.4}) \\ &\cong \text{Sets}(A, \mathcal{P}(B)), \quad (\text{by Item 2 of Proposition 4.5.1.1.4}) \end{aligned}$$

where all bijections are natural in  $A$ , (where we are using Item 3 of Proposition 4.5.1.1.4). Explicitly, this isomorphism is given by sending a relation  $R: \text{Gr}(A) \rightarrow B$  to the map  $R^\dagger: A \rightarrow \mathcal{P}(B)$  sending  $a$  to the subset  $R(a)$  of  $B$ , as in Definition 8.1.1.1.1.

Naturality in  $B$  is then the statement that given a relation  $R: B \rightarrow B'$ , the diagram

$$\begin{array}{ccc} \text{Rel}(\text{Gr}(A), B) & \xrightarrow{R^\dagger} & \text{Rel}(\text{Gr}(A), B') \\ \downarrow & & \downarrow \\ \text{Sets}(A, \mathcal{P}(B)) & \xrightarrow{R_!} & \text{Sets}(A, \mathcal{P}(B')) \end{array}$$

commutes, which follows from Remark 8.7.1.1.3. 

**01JK 4.4.5 Powersets as Free Cocompletions**

Let  $X$  be a set.

01JL

**PROPOSITION 4.4.5.1.1 ► POWERSETS AS FREE Cocompletions: UNIVERSAL PROPERTY**

The pair  $(\mathcal{P}(X), \chi_{(-)})$  consisting of

- The powerset  $(\mathcal{P}(X), \subset)$  of  $X$  of [Definition 4.4.1.1.1](#);
- The characteristic embedding  $\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$  of  $X$  into  $\mathcal{P}(X)$  of [Definition 4.5.4.1.1](#);

satisfies the following universal property:

(★) Given another pair  $(Y, f)$  consisting of

- A suplattice  $(Y, \preceq)$ ;
- A function  $f: X \rightarrow Y$ ;

there exists a unique morphism of suplattices

$$(\mathcal{P}(X), \subset) \xrightarrow{\exists!} (Y, \preceq)$$

making the diagram

$$\begin{array}{ccc} & \mathcal{P}(X) & \\ \chi_X \nearrow & \downarrow \exists! & \\ X & \xrightarrow{f} & Y \end{array}$$

commute.

**PROOF 4.4.5.1.2 ► PROOF OF PROPOSITION 4.4.5.1.1**

This is a rephrasing of [Proposition 4.4.5.1.3](#), which we prove below.<sup>1</sup> 

<sup>1</sup>Here we only remark that the unique morphism of suplattices in the statement is given by the left Kan extension  $\text{Lan}_{\chi_X}(f)$  of  $f$  along  $\chi_X$ .

01JM

**PROPOSITION 4.4.5.1.3 ► POWERSETS AS FREE Cocompletions: ADJOINTNESS**

We have an adjunction

$$(\mathcal{P} \dashv \overline{\text{忘}}): \text{Sets} \begin{array}{c} \xrightarrow{\quad \mathcal{P} \quad} \\ \perp \\ \xleftarrow{\quad \text{忘} \quad} \end{array} \text{SupLat},$$

witnessed by a bijection

$$\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) \cong \text{Sets}(X, Y),$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $(Y, \preceq) \in \text{Obj}(\text{SupLat})$ , where:

- The category  $\text{SupLat}$  is the category of suplattices of ??.
- The map

$$\chi_X^*: \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

witnessing the above bijection is defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a morphism of suplattices  $f: \mathcal{P}(X) \rightarrow Y$  to the composition

$$X \xrightarrow{\chi_X} \mathcal{P}(X) \xrightarrow{f} Y.$$

- The map

$$\text{Lan}_{\chi_X}: \text{Sets}(X, Y) \rightarrow \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$$

witnessing the above bijection is given by sending a function  $f: X \rightarrow Y$  to its left Kan extension along  $\chi_X$ ,

$$\begin{array}{ccc} & \mathcal{P}(X) & \\ \text{Lan}_{\chi_X}(f): \mathcal{P}(X) & \xrightarrow{\chi_X} & Y, \\ & \downarrow \text{Lan}_{\chi_X}(f) & \\ X & \xrightarrow{f} & Y. \end{array}$$

Moreover, invoking the bijection  $\mathcal{P}(X) \cong \text{Sets}(X, \{t, f\})$  of [Item 2 of Proposition 4.5.1.1.4](#),  $\text{Lan}_{\chi_X}(f)$  can be explicitly computed by

$$\begin{aligned} [\text{Lan}_{\chi_X}(f)](U) &= \int^{x \in X} \chi_{\mathcal{P}(X)}(\chi_x, U) \odot f(x) \\ &= \int^{x \in X} \chi_U(x) \odot f(x) \\ &= \bigvee_{x \in X} (\chi_U(x) \odot f(x)) \\ &= \left( \bigvee_{x \in U} (\chi_U(x) \odot f(x)) \right) \vee \left( \bigvee_{x \in U^c} (\chi_U(x) \odot f(x)) \right) \\ &= \left( \bigvee_{x \in U} f(x) \right) \vee \left( \bigvee_{x \in U^c} \emptyset_Y \right) \\ &= \bigvee_{x \in U} f(x) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , where:

- We have used ?? for the first equality.
- We have used **Proposition 4.5.5.1.1** for the second equality.
- We have used ?? for the third equality.
- The symbol  $\vee$  denotes the join in  $(Y, \preceq)$ .
- The symbol  $\odot$  denotes the tensor of an element of  $Y$  by a truth value as in ???. In particular, we have

$$\begin{aligned}\text{true} \odot f(x) &\stackrel{\text{def}}{=} f(x), \\ \text{false} \odot f(x) &\stackrel{\text{def}}{=} \emptyset_Y,\end{aligned}$$

where  $\emptyset_Y$  is the bottom element of  $(Y, \preceq)$ .

In particular, when  $(Y, \preceq_Y) = (\mathcal{P}(B), \subset)$  for some set  $B$ , the Kan extension  $\text{Lan}_{\chi_X}(f)$  is given by

$$\begin{aligned}[\text{Lan}_{\chi_X}(f)](U) &= \bigvee_{x \in U} f(x) \\ &= \bigcup_{x \in U} f(x)\end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

#### PROOF 4.4.5.1.4 ► PROOF OF PROPOSITION 4.4.5.1.3

##### Map I

We define a map

$$\Phi_{X,Y}: \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

as in the statement, i.e. by

$$\Phi_{X,Y}(f) \stackrel{\text{def}}{=} f \circ \chi_X$$

for each  $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$ .

##### Map II

We define a map

$$\Psi_{X,Y}: \text{Sets}(X, Y) \rightarrow \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$$

as in the statement, i.e. by

$$\Psi_{X,Y}(f) \stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f), \quad \begin{array}{ccc} & \mathcal{P}(X) & \\ & \nearrow \chi_X & \downarrow \text{Lan}_{\chi_X}(f) \\ X & \xrightarrow{f} & Y, \end{array}$$

for each  $f \in \text{Sets}(X, Y)$ .

**Invertibility I**

We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))}.$$

We have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](f) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}(f \circ \chi_X) \\ &\stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f \circ \chi_X) \end{aligned}$$

for each  $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$ . We now claim that

$$\text{Lan}_{\chi_X}(f \circ \chi_X) = f$$

for each  $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$ . Indeed, we have

$$\begin{aligned} [\text{Lan}_{\chi_X}(f \circ \chi_X)](U) &= \bigvee_{x \in U} f(\chi_X(x)) \\ &= f\left(\bigvee_{x \in U} \chi_X(x)\right) \\ &= f\left(\bigcup_{x \in U} \{x\}\right) \\ &= f(U) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , where we have used that  $f$  is a morphism of suplattices and hence preserves joins for the second equality. This proves our claim. Since we have shown that

$$[\Psi_{X,Y} \circ \Phi_{X,Y}](f) = f$$

for each  $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$ , it follows that  $\Psi_{X,Y} \circ \Phi_{X,Y}$  must be equal to the identity map  $\text{id}_{\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))}$  of  $\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$ .

### Invertibility II

We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X,Y)}.$$

We have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](f) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\text{Lan}_{\chi_X}(f)) \\ &\stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f) \circ \chi_X \end{aligned}$$

for each  $f \in \text{Sets}(X, Y)$ . We now claim that

$$\text{Lan}_{\chi_X}(f) \circ \chi_X = f$$

for each  $f \in \text{Sets}(X, Y)$ . Indeed, we have

$$\begin{aligned} [\text{Lan}_{\chi_X}(f) \circ \chi_X](x) &= \bigvee_{y \in \{x\}} f(y) \\ &= f(x) \end{aligned}$$

for each  $x \in X$ . This proves our claim. Since we have shown that

$$[\Phi_{X,Y} \circ \Psi_{X,Y}](f) = f$$

for each  $f \in \text{Sets}(X, Y)$ , it follows that  $\Phi_{X,Y} \circ \Psi_{X,Y}$  must be equal to the identity map  $\text{id}_{\text{Sets}(X,Y)}$  of  $\text{Sets}(X, Y)$ .

### Naturality for $\Phi$ , Part I

We need to show that, given a function  $f: X \rightarrow X'$ , the diagram

$$\begin{array}{ccc} \text{SupLat}((\mathcal{P}(X'), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}(X', Y) \\ \mathcal{P}_!(f)^* \downarrow & & \downarrow f^* \\ \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \end{array}$$

commutes. Indeed, we have

$$[\Phi_{X,Y} \circ \mathcal{P}_!(f)^*](\xi) \stackrel{\text{def}}{=} \Phi_{X,Y}(\mathcal{P}_!(f)^*(\xi))$$

$$\begin{aligned}
&\stackrel{\text{def}}{=} \Phi_{X,Y}(\xi \circ f_!) \\
&\stackrel{\text{def}}{=} (\xi \circ f_!) \circ \chi_X \\
&= \xi \circ (f_! \circ \chi_X) \\
&\stackrel{(\dagger)}{=} \xi \circ (\chi_{X'} \circ f) \\
&= (\xi \circ \chi_{X'}) \circ f \\
&\stackrel{\text{def}}{=} \Phi_{X',Y}(\xi) \circ f \\
&\stackrel{\text{def}}{=} f^*(\Phi_{X',Y}(\xi)) \\
&\stackrel{\text{def}}{=} [f^* \circ \Phi_{X',Y}](\xi),
\end{aligned}$$

for each  $\xi \in \text{SupLat}((\mathcal{P}(X'), \subset), (Y, \preceq))$ , where we have used [Item 1](#) of [Proposition 4.5.4.1.3](#) for the fifth equality above.

### Naturality for $\Phi$ , Part II

We need to show that, given a morphism of suplattices

$$g: (Y, \preceq_Y) \rightarrow (Y', \preceq_{Y'}),$$

the diagram

$$\begin{array}{ccc}
\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\
g_! \downarrow & & \downarrow g_! \\
\text{SupLat}((\mathcal{P}(X), \subset), (Y', \preceq')) & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y')
\end{array}$$

commutes. Indeed, we have

$$\begin{aligned}
[\Phi_{X,Y'} \circ g_!](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y'}(g_!(\xi)) \\
&\stackrel{\text{def}}{=} \Phi_{X,Y'}(g \circ \xi) \\
&\stackrel{\text{def}}{=} (g \circ \xi) \circ \chi_X \\
&= g \circ (\xi \circ \chi_X) \\
&\stackrel{\text{def}}{=} g \circ (\Phi_{X,Y}(\xi)) \\
&\stackrel{\text{def}}{=} g_!(\Phi_{X,Y}(\xi)) \\
&\stackrel{\text{def}}{=} [g_! \circ \Phi_{X,Y}](\xi).
\end{aligned}$$

for each  $\xi \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$ .

### Naturality for $\Psi$

Since  $\Phi$  is natural in each argument and  $\Phi$  is a componentwise inverse to  $\Psi$  in each argument, it follows from [Item 2](#) of [Proposition 11.9.7.1.2](#) that  $\Psi$  is also natural in each argument. ■


**WARNING 4.4.5.1.5 ► FREE COCOMPLETION IS NOT AN IDEMPOTENT OPERATION**

Although the assignment  $X \mapsto \mathcal{P}(X)$  is called the *free cocompletion* of  $X$ , it is not an idempotent operation, i.e. we have  $\mathcal{P}(\mathcal{P}(X)) \neq \mathcal{P}(X)$ .

**01JP 4.4.6 Powersets as Free Completions**

Let  $X$  be a set.

**01JQ**
**PROPOSITION 4.4.6.1.1 ► POWERSETS AS FREE COMPLETIONS: UNIVERSAL PROPERTY**

The pair  $(\mathcal{P}(X), \chi_{(-)})$  consisting of

- The powerset of  $X$  together with reverse inclusion  $\mathcal{P}(X)^{\text{op}} = (\mathcal{P}(X), \supset)$  of [Definition 4.4.1.1.1](#);
- The characteristic embedding  $\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$  of  $X$  into  $\mathcal{P}(X)$  of [Definition 4.5.4.1.1](#);

satisfies the following universal property:

- (★) Given another pair  $(Y, f)$  consisting of
- An inflattice  $(Y, \preceq)$ ;
  - A function  $f: X \rightarrow Y$ ;

there exists a unique morphism of inflattices

$$(\mathcal{P}(X), \supset) \xrightarrow{\exists!} (Y, \preceq)$$

making the diagram

$$\begin{array}{ccc} & \mathcal{P}(X)^{\text{op}} & \\ \chi_X \nearrow & \downarrow \exists! & \\ X & \xrightarrow{f} & Y \end{array}$$

commute.

## PROOF 4.4.6.1.2 ► PROOF OF PROPOSITION 4.4.6.1.1

This is a rephrasing of [Proposition 4.4.6.1.3](#), which we prove below.<sup>1</sup>

<sup>1</sup>Here we only remark that the unique morphism of inflattices in the statement is given by the right Kan extension  $\text{Ran}_{\chi_X}(f)$  of  $f$  along  $\chi_X$ .

01JR

## PROPOSITION 4.4.6.1.3 ► POWERSETS AS FREE COMPLETIONS: ADJOINTNESS

We have an adjunction

$$(\mathcal{P} \dashv \text{InfLat}): \text{Sets} \begin{array}{c} \xrightarrow{\quad \mathcal{P} \quad} \\ \perp \\ \xleftarrow{\quad \text{InfLat} \quad} \end{array} \text{InfLat},$$

witnessed by a bijection

$$\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) \cong \text{Sets}(X, Y),$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $(Y, \preceq) \in \text{Obj}(\text{InfLat})$ , where:

- The category  $\text{InfLat}$  is the category of inflattices of [??](#).
- The map

$$\chi_X^*: \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

witnessing the above bijection is defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a morphism of inflattices  $f: \mathcal{P}(X)^{\text{op}} \rightarrow Y$  to the composition

$$X \xrightarrow{\chi_X} \mathcal{P}(X)^{\text{op}} \xrightarrow{f} Y.$$

- The map

$$\text{Ran}_{\chi_X}: \text{Sets}(X, Y) \rightarrow \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$$

witnessing the above bijection is given by sending a function  $f: X \rightarrow Y$  to its right Kan extension along  $\chi_X$ ,

$$\text{Ran}_{\chi_X}(f): \mathcal{P}(X)^{\text{op}} \rightarrow Y, \quad \begin{array}{ccc} & \mathcal{P}(X)^{\text{op}} & \\ & \nearrow \chi_X & \downarrow \text{Ran}_{\chi_X}(f) \\ X & \xrightarrow{f} & Y. \end{array}$$

Moreover, invoking the bijection  $\mathcal{P}(X) \cong \text{Sets}(X, \{\text{t}, \text{f}\})$  of [Item 2](#) of [Proposition 4.5.1.1.4](#),  $\text{Ran}_{\chi_X}(f)$  can be explicitly computed by

$$\begin{aligned} [\text{Ran}_{\chi_X}(f)](U) &= \int_{x \in X} \chi_{\mathcal{P}(X)^{\text{op}}}(\chi_x, U) \pitchfork f(x) \\ &= \int_{x \in X} \chi_{\mathcal{P}(X)}(U, \chi_x) \pitchfork f(x) \\ &= \int_{x \in X} \chi_U(x) \pitchfork f(x) \\ &= \bigwedge_{x \in X} \chi_U(x) \pitchfork f(x) \\ &= \left( \bigwedge_{x \in U} \chi_U(x) \pitchfork f(x) \right) \wedge \left( \bigwedge_{x \in U^c} \chi_U(x) \pitchfork f(x) \right) \\ &= \left( \bigwedge_{x \in U} f(x) \right) \wedge \left( \bigwedge_{x \in U^c} \infty_Y \right) \\ &= \left( \bigwedge_{x \in U} f(x) \right) \wedge \infty_Y \\ &= \bigwedge_{x \in U} f(x) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , where:

- We have used ?? for the first equality.
- We have used [Proposition 4.5.1.1](#) for the second equality.
- We have used ?? for the third equality.
- The symbol  $\wedge$  denotes the meet in  $(Y, \preceq)$ .
- The symbol  $\pitchfork$  denotes the cotensor of an element of  $Y$  by a truth value as in ?. In particular, we have

$$\begin{aligned} \text{true} \pitchfork f(x) &\stackrel{\text{def}}{=} f(x), \\ \text{false} \pitchfork f(x) &\stackrel{\text{def}}{=} \infty_Y, \end{aligned}$$

where  $\infty_Y$  is the top element of  $(Y, \preceq)$ .

In particular, when  $(Y, \preceq_Y) = (\mathcal{P}(B), \subset)$  for some set  $B$ , the Kan extension  $\text{Ran}_{\chi_X}(f)$  is given by

$$\begin{aligned} [\text{Ran}_{\chi_X}(f)](U) &= \bigwedge_{x \in U} f(x) \\ &= \bigcap_{x \in U} f(x) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

**PROOF 4.4.6.1.4 ► PROOF OF PROPOSITION 4.4.5.1.3**
**Map I**

We define a map

$$\Phi_{X,Y} : \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

as in the statement, i.e. by

$$\Phi_{X,Y}(f) \stackrel{\text{def}}{=} f \circ \chi_X$$

for each  $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$ .

**Map II**

We define a map

$$\Psi_{X,Y} : \text{Sets}(X, Y) \rightarrow \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$$

as in the statement, i.e. by

$$\begin{array}{c} \mathcal{P}(X)^{\text{op}} \\ \Psi_{X,Y}(f) \stackrel{\text{def}}{=} \text{Ran}_{\chi_X}(f), \\ X \xrightarrow[f]{\quad} Y, \\ \downarrow \quad \uparrow \\ \text{Ran}_{\chi_X}(f) \end{array}$$

for each  $f \in \text{Sets}(X, Y)$ .

**Invertibility I**

We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))}.$$

We have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](f) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}(f \circ \chi_X) \\ &\stackrel{\text{def}}{=} \text{Ran}_{\chi_X}(f \circ \chi_X) \end{aligned}$$

for each  $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$ . We now claim that

$$\text{Ran}_{\chi_X}(f \circ \chi_X) = f$$

for each  $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$ . Indeed, we have

$$\begin{aligned} [\text{Ran}_{\chi_X}(f \circ \chi_X)](U) &= \bigwedge_{x \in U} f(\chi_X(x)) \\ &= f\left(\bigwedge_{x \in U} \chi_X(x)\right) \\ &= f\left(\bigcup_{x \in U} \{x\}\right) \\ &= f(U) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , where we have used that  $f$  is a morphism of inflattices and hence preserves meets in  $(\mathcal{P}(X), \supset)$  (i.e. joins in  $(\mathcal{P}(X), \subset)$ ) for the second equality. This proves our claim. Since we have shown that

$$[\Psi_{X,Y} \circ \Phi_{X,Y}](f) = f$$

for each  $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$ , it follows that  $\Psi_{X,Y} \circ \Phi_{X,Y}$  must be equal to the identity map  $\text{id}_{\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))}$  of  $\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$ .

### Invertibility II

We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X,Y)}.$$

We have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](f) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\text{Ran}_{\chi_X}(f)) \\ &\stackrel{\text{def}}{=} \text{Ran}_{\chi_X}(f) \circ \chi_X \end{aligned}$$

for each  $f \in \text{Sets}(X, Y)$ . We now claim that

$$\text{Ran}_{\chi_X}(f) \circ \chi_X = f$$

for each  $f \in \text{Sets}(X, Y)$ . Indeed, we have

$$\begin{aligned} [\text{Ran}_{\chi_X}(f) \circ \chi_X](x) &= \bigwedge_{y \in \{x\}} f(y) \\ &= f(x) \end{aligned}$$

for each  $x \in X$ . This proves our claim. Since we have shown that

$$[\Phi_{X,Y} \circ \Psi_{X,Y}](f) = f$$

for each  $f \in \text{Sets}(X, Y)$ , it follows that  $\Phi_{X,Y} \circ \Psi_{X,Y}$  must be equal to the identity map  $\text{id}_{\text{Sets}(X,Y)}$  of  $\text{Sets}(X, Y)$ .

### Naturality for $\Phi$ , Part I

We need to show that, given a function  $f: X \rightarrow X'$ , the diagram

$$\begin{array}{ccc} \text{InfLat}((\mathcal{P}(X'), \supset), (Y, \preceq)) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}(X', Y) \\ \mathcal{P}_!(f)^* \downarrow & & \downarrow f^* \\ \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \end{array}$$

commutes. Indeed, we have

$$\begin{aligned} [\Phi_{X,Y} \circ \mathcal{P}_!(f)^*](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\mathcal{P}_!(f)^*(\xi)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\xi \circ f_!) \\ &\stackrel{\text{def}}{=} (\xi \circ f_!) \circ \chi_X \\ &= \xi \circ (f_! \circ \chi_X) \\ &\stackrel{(\dagger)}{=} \xi \circ (\chi_{X'} \circ f) \\ &= (\xi \circ \chi_{X'}) \circ f \\ &\stackrel{\text{def}}{=} \Phi_{X',Y}(\xi) \circ f \\ &\stackrel{\text{def}}{=} f^*(\Phi_{X',Y}(\xi)) \\ &\stackrel{\text{def}}{=} [f^* \circ \Phi_{X',Y}](\xi), \end{aligned}$$

for each  $\xi \in \text{InfLat}((\mathcal{P}(X'), \supset), (Y, \preceq))$ , where we have used **Item 1** of **Proposition 4.5.4.1.3** for the fifth equality above.

### Naturality for $\Phi$ , Part II

We need to show that, given a cocontinuous morphism of posets

$$g: (Y, \preceq_Y) \rightarrow (Y', \preceq_{Y'}),$$

the diagram

$$\begin{array}{ccc} \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\ g_! \downarrow & & \downarrow g^* \\ \text{InfLat}((\mathcal{P}(X), \supset), (Y', \preceq)) & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y') \end{array}$$

commutes. Indeed, we have

$$[\Phi_{X,Y'} \circ g_!](\xi) \stackrel{\text{def}}{=} \Phi_{X,Y'}(g_!(\xi))$$

$$\begin{aligned}
 &\stackrel{\text{def}}{=} \Phi_{X,Y'}(g \circ \xi) \\
 &\stackrel{\text{def}}{=} (g \circ \xi) \circ \chi_X \\
 &= g \circ (\xi \circ \chi_X) \\
 &\stackrel{\text{def}}{=} g \circ (\Phi_{X,Y}(\xi)) \\
 &\stackrel{\text{def}}{=} g_!(\Phi_{X,Y}(\xi)) \\
 &\stackrel{\text{def}}{=} [g_! \circ \Phi_{X,Y}](\xi).
 \end{aligned}$$


for each  $\xi \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$ .

#### Naturality for $\Psi$

Since  $\Phi$  is natural in each argument and  $\Phi$  is a componentwise inverse to  $\Psi$  in each argument, it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\Psi$  is also natural in each argument. 

#### WARNING 4.4.6.1.5 ► FREE COMPLETION IS NOT AN IDEMPOTENT OPERATION



Although the assignment  $X \mapsto \mathcal{P}(X)^{\text{op}}$  is called the *free completion* of  $X$ , it is not an idempotent operation, i.e. we have  $\mathcal{P}(\mathcal{P}(X)^{\text{op}})^{\text{op}} \neq \mathcal{P}(X)^{\text{op}}$ . 

### 01JT 4.4.7 The Internal Hom of a Powerset

Let  $X$  be a set and let  $U, V \in \mathcal{P}(X)$ .

01JU

#### PROPOSITION 4.4.7.1.1 ► THE INTERNAL HOM OF A POWERSET

The **internal Hom** of  $\mathcal{P}(X)$  from  $U$  to  $V$  is the subset  $[U, V]_X^1$  of  $X$  given by

$$\begin{aligned}
 [U, V]_X &= U^c \cup V \\
 &= (U \setminus V)^c
 \end{aligned}$$

where  $U^c$  is the complement of  $U$  of [Definition 4.3.11.1.1](#).

<sup>1</sup>Further Notation: Also written  $\text{Hom}_{\mathcal{P}(X)}(U, V)$ .

#### PROOF 4.4.7.1.2 ► PROOF OF PROPOSITION 4.4.7.1.1

Proof of the Equality  $U^c \cup V = (U \setminus V)^c$

We have

$$\begin{aligned}
 (U \setminus V)^c &\stackrel{\text{def}}{=} X \setminus (U \setminus V) \\
 &= (X \cap V) \cup (X \setminus U) \\
 &= V \cup (X \setminus U) \\
 &\stackrel{\text{def}}{=} V \cup U^c \\
 &= U^c \cup V,
 \end{aligned}$$

where we have used:

- 020S 1. Item 10 of Proposition 4.3.10.1.2 for the second equality.
- 020T 2. Item 4 of Proposition 4.3.9.1.2 for the third equality.
- 020U 3. Item 4 of Proposition 4.3.8.1.2 for the last equality.

This finishes the proof.

Proof that  $U^c \cup V$  Is Indeed the Internal Hom

This follows from Item 2 of Proposition 4.3.9.1.2. 

004C

#### REMARK 4.4.7.1.3 ► INTUITION FOR THE INTERNAL HOM OF $\mathcal{P}(X)$

Henning Makholm suggests the following heuristic intuition for the internal Hom of  $\mathcal{P}(X)$  from  $U$  to  $V$  ([MSE 267365]):

01JV

1. Since products in  $\mathcal{P}(X)$  are given by binary intersections (Item 1 of Proposition 4.4.1.1.4), the right adjoint  $\text{Hom}_{\mathcal{P}(X)}(U, -)$  of  $U \cap -$  may be thought of as a function type  $[U, V]$ .

01JW

2. Under the Curry–Howard correspondence (??), the function type  $[U, V]$  corresponds to implication  $U \Rightarrow V$ .

01JX

3. Implication  $U \Rightarrow V$  is logically equivalent to  $\neg U \vee V$ .

01JY

4. The expression  $\neg U \vee V$  then corresponds to the set  $U^c \cup V$  in  $\mathcal{P}(X)$ .

01JZ

5. The set  $U^c \cup V$  turns out to indeed be the internal Hom of  $\mathcal{P}(X)$ .

01K0

#### PROPOSITION 4.4.7.1.4 ► PROPERTIES OF INTERNAL HOMS OF POWERSETS

Let  $X$  be a set.

01K1

1. *Functionality.* The assignments  $U, V, (U, V) \mapsto \mathbf{Hom}_{\mathcal{P}(X)}$  define functors

$$\begin{aligned} [U, -]_X : (\mathcal{P}(X), \supset) &\rightarrow (\mathcal{P}(X), \subset), \\ [-, V]_X : (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ [-_1, -_2]_X : (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) &\rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each  $U, V, A, B \in \mathcal{P}(X)$ :

01K2

(a) If  $U \subset A$ , then  $[A, V]_X \subset [U, V]_X$ .

01K3

(b) If  $V \subset B$ , then  $[U, V]_X \subset [U, B]_X$ .

01K4

(c) If  $U \subset A$  and  $V \subset B$ , then  $[A, V]_X \subset [U, B]_X$ .

01K5

2. *Adjointness.* We have adjunctions

$$\begin{aligned} (U \cap - \dashv [U, -]_X) : \mathcal{P}(X) &\xrightleftharpoons[\substack{[U, -]_X}]{} \mathcal{P}(X), \\ (- \cap V \dashv [V, -]_X) : \mathcal{P}(X) &\xrightleftharpoons[\substack{[V, -]_X}]{} \mathcal{P}(X), \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \mathbf{Hom}_{\mathcal{P}(X)}(U \cap V, W) &\cong \mathbf{Hom}_{\mathcal{P}(X)}(U, [V, W]_X), \\ \mathbf{Hom}_{\mathcal{P}(X)}(U \cap V, W) &\cong \mathbf{Hom}_{\mathcal{P}(X)}(V, [U, W]_X). \end{aligned}$$

In particular, the following statements hold for each  $U, V, W \in \mathcal{P}(X)$ :

01K6

- (a) The following conditions are equivalent:

01K7

i. We have  $U \cap V \subset W$ .

01K8

ii. We have  $U \subset [V, W]_X$ .

01K9

- (b) The following conditions are equivalent:

01KA

i. We have  $U \cap V \subset W$ .

01KB

ii. We have  $V \subset [U, W]_X$ .

01KC

3. *Interaction With the Empty Set I.* We have

$$\begin{aligned} [U, \emptyset]_X &= U^c, \\ [\emptyset, V]_X &= X, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

01KD

4. *Interaction With X.* We have

$$\begin{aligned}[U, X]_X &= X, \\ [X, V]_X &= V,\end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

01KE

5. *Interaction With the Empty Set II.* The functor

$$D_X: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X)$$

defined by

$$\begin{aligned}D_X &\stackrel{\text{def}}{=} [-, \emptyset]_X \\ &= (-)^c\end{aligned}$$

is an involutory isomorphism of categories, making  $\emptyset$  into a dualising object for  $(\mathcal{P}(X), \cap, X, [-, -]_X)$  in the sense of [??](#). In particular:

01KF

(a) The diagram

$$\begin{array}{ccc}\mathcal{P}(X)^{\text{op}} & \xrightarrow{D_X} & \mathcal{P}(X) \\ & \searrow \text{id}_{\mathcal{P}(X)^{\text{op}}} & \downarrow D_X \\ & & \mathcal{P}(X)^{\text{op}}\end{array}$$

commutes, i.e. we have

$$\underbrace{D_X(D_X(U))}_{\stackrel{\text{def}}{=} [[U, \emptyset]_X, \emptyset]_X} = U$$

for each  $U \in \mathcal{P}(X)$ .

01KG

(b) The diagram

$$\begin{array}{ccc}\mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} & \xrightarrow{\cap^{\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ \text{id}_{\mathcal{P}(X)^{\text{op}}} \times D_X \nearrow & & \searrow D_X \\ \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-_1, -_2]_X} & \mathcal{P}(X)\end{array}$$

commutes, i.e. we have

$$\underbrace{D_X(U \cap D_X(V))}_{\stackrel{\text{def}}{=} [U \cap [V, \emptyset]_X, \emptyset]_X} = [U, V]_X$$

for each  $U, V \in \mathcal{P}(X)$ .

01KH

6. *Interaction With the Empty Set III.* Let  $f: X \rightarrow Y$  be a function.

01KJ

(a) *Interaction With Direct Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ D_X \downarrow & & \downarrow D_Y \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(D_X(U)) = D_Y(f_*(U))$$

for each  $U \in \mathcal{P}(X)$ .

01KK

(b) *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1,\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ D_Y \downarrow & & \downarrow D_X \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(D_Y(U)) = D_X(f^{-1}(U))$$

for each  $U \in \mathcal{P}(X)$ .

01KL

(c) *Interaction With Codirect Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ D_X \downarrow & & \downarrow D_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(D_X(U)) = D_Y(f_!(U))$$

for each  $U \in \mathcal{P}(X)$ .

01KM

## 7. Interaction With Unions of Families of Subsets I. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-_1, -_2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\ \cup^{\text{op}} \times \cup^{\text{op}} \downarrow & \times & \downarrow \cup \\ \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-_1, -_2]_X} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[ \bigcup_{U \in \mathcal{U}} U, \bigcup_{V \in \mathcal{V}} V \right]_X$$

in general, where  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

01KN

## 8. Interaction With Unions of Families of Subsets II. The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ & \searrow \cup^{\text{op}} & \\ \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & \nearrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & \mathcal{P}(X)^{\text{op}} \\ & \downarrow & \downarrow [-, V]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[ \bigcup_{U \in \mathcal{U}} U, V \right]_X = \bigcap_{U \in \mathcal{U}} [U, V]_X$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$  and each  $V \in \mathcal{P}(X)$ .

01KP

## 9. Interaction With Unions of Families of Subsets III. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \\ \text{id}_{\mathcal{P}(X)} \star [U, -]_X \downarrow & & \downarrow [U, -]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[ U, \bigcup_{V \in \mathcal{V}} V \right]_X = \bigcup_{V \in \mathcal{V}} [U, V]_X$$

for each  $U \in \mathcal{P}(X)$  and each  $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01KQ

10. *Interaction With Intersections of Families of Subsets I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-1, -2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\ \cap^{\text{op}} \times \cap^{\text{op}} \downarrow & \times & \downarrow \cap \\ \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[ \bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X$$

in general, where  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ .

01KR

11. *Interaction With Intersections of Families of Subsets II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ & \nearrow \text{dotted} \quad \searrow \cap^{\text{op}} & \\ \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\ \text{id}_{\mathcal{P}(X)} \star [-, V]_X \swarrow & & \searrow [-, V]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[ \bigcap_{U \in \mathcal{U}} U, V \right]_X = \bigcup_{U \in \mathcal{U}} [U, V]_X$$

for each  $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$  and each  $V \in \mathcal{P}(X)$ .

01KS

12. *Interaction With Intersections of Families of Subsets III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \\ \text{id}_{\mathcal{P}(X)} \star [U, -]_X \downarrow & & \downarrow [U, -]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[ U, \bigcap_{V \in \mathcal{V}} V \right]_X = \bigcap_{V \in \mathcal{V}} [U, V]_X$$

for each  $U \in \mathcal{P}(X)$  and each  $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ .

01KT

13. *Interaction With Binary Unions.* We have equalities of sets

$$\begin{aligned} [U \cap V, W]_X &= [U, W]_X \cup [V, W]_X, \\ [U, V \cap W]_X &= [U, V]_X \cap [U, W]_X \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

01KU

14. *Interaction With Binary Intersections.* We have equalities of sets

$$\begin{aligned} [U \cup V, W]_X &= [U, W]_X \cap [V, W]_X, \\ [U, V \cup W]_X &= [U, V]_X \cup [U, W]_X \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

01KV

15. *Interaction With Differences.* We have equalities of sets

$$\begin{aligned} [U \setminus V, W]_X &= [U, W]_X \cup [V^c, W]_X \\ &= [U, W]_X \cup [U, V]_X, \\ [U, V \setminus W]_X &= [U, V]_X \setminus (U \cap W) \end{aligned}$$

for each  $U, V, W \in \mathcal{P}(X)$ .

01KW

16. *Interaction With Complements.* We have equalities of sets

$$\begin{aligned} [U^c, V]_X &= U \cup V, \\ [U, V^c]_X &= U \cap V, \\ [U, V]_X^c &= U \setminus V \end{aligned}$$

for each  $U, V \in \mathcal{P}(X)$ .

01KX

17. *Interaction With Characteristic Functions.* We have

$$\chi_{[U,V]_{\mathcal{P}(X)}}(x) = \max(1 - \chi_U(x) \pmod{2}, \chi_V(x))$$

for each  $U, V \in \mathcal{P}(X)$ .

01KY

18. *Interaction With Direct Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ [-1,-2]_X \downarrow & & \downarrow [-1,-2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have an equality of sets

$$f_!([U, V]_X) = [f_*(U), f_!(V)]_Y,$$

natural in  $U, V \in \mathcal{P}(X)$ .

01KZ

19. *Interaction With Inverse Images.* Let  $f: X \rightarrow Y$  be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{-1,\text{op}} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ [-1,-2]_Y \downarrow & & \downarrow [-1,-2]_X \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$f^{-1}([U, V]_Y) = [f^{-1}(U), f^{-1}(V)]_X,$$

natural in  $U, V \in \mathcal{P}(X)$ .

01L0

20. *Interaction With Codirect Images.* Let  $f: X \rightarrow Y$  be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ [-1,-2]_X \downarrow & \circlearrowleft & \downarrow [-1,-2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$[f_!(U), f_*(V)]_Y \subset f_*([U, V]_X)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

## PROOF 4.4.7.1.5 ► PROOF OF PROPOSITION 4.4.7.1.4

## Item 1: Functoriality

Since  $\mathcal{P}(X)$  is posetal, it suffices to prove **Items 1a** to **1c**.

020W

1. *Proof of Item 1a:* We have

$$\begin{aligned} [A, V]_X &\stackrel{\text{def}}{=} A^c \cup V \\ &\subset U^c \cup V \\ &\stackrel{\text{def}}{=} [U, V]_X, \end{aligned}$$

where we have used:

020X

(a) **Item 1** of **Proposition 4.3.11.1.2**, which states that if  $U \subset A$ , then  $A^c \subset U^c$ .

020Y

(b) **Item 1a** of **Item 1** of **Proposition 4.3.11.1.2**, which states that if  $A^c \subset U^c$ , then  $A^c \cup K \subset U^c \cup K$  for any  $K \in \mathcal{P}(X)$ .

020Z

2. *Proof of Item 1b:* We have

$$\begin{aligned} [U, V]_X &\stackrel{\text{def}}{=} U^c \cup V \\ &\subset U^c \cup B \\ &\stackrel{\text{def}}{=} [U, B]_X, \end{aligned}$$

where we have used **Item 1b** of **Item 1** of **Proposition 4.3.11.1.2**, which states that if  $V \subset B$ , then  $K \cup V \subset K \cup B$  for any  $K \in \mathcal{P}(X)$ .

0210

3. *Proof of Item 1c:* We have

$$\begin{aligned} [A, V]_X &\subset [U, V]_X \\ &\subset [U, B]_X, \end{aligned}$$

where we have used **Items 1a** and **1b**.

This finishes the proof.

## Item 2: Adjointness

This is a repetition of **Item 2** of **Proposition 4.3.9.1.2** and is proved there.

## Item 3: Interaction With the Empty Set I

We have

$$\begin{aligned} [U, \emptyset]_X &\stackrel{\text{def}}{=} U^c \cup \emptyset \\ &= U^c, \end{aligned}$$

where we have used [Item 3 of Proposition 4.3.8.1.2](#), and we have

$$\begin{aligned} [\emptyset, V]_X &\stackrel{\text{def}}{=} \emptyset^c \cup V \\ &\stackrel{\text{def}}{=} (X \setminus \emptyset) \cup V \\ &= X \cup V \\ &= X, \end{aligned}$$

where we have used:

- 0211 1. [Item 12 of Proposition 4.3.10.1.2](#) for the first equality.  
 0212 2. [Item 5 of Proposition 4.3.8.1.2](#) for the last equality.

Since  $\mathcal{P}(X)$  is posetal, naturality is automatic ([??](#) of [??](#)).

#### Item 4: Interaction With $X$

We have

$$\begin{aligned} [U, X]_X &\stackrel{\text{def}}{=} U^c \cup X \\ &= X, \end{aligned}$$

where we have used [Item 5 of Proposition 4.3.8.1.2](#), and we have

$$\begin{aligned} [X, V]_X &\stackrel{\text{def}}{=} X^c \cup V \\ &\stackrel{\text{def}}{=} (X \setminus X) \cup V \\ &= \emptyset \cup V \\ &= V, \end{aligned}$$

where we have used [Item 3 of Proposition 4.3.8.1.2](#) for the last equality.  
 Since  $\mathcal{P}(X)$  is posetal, naturality is automatic ([??](#) of [??](#)).

#### Item 5: Interaction With the Empty Set II

We have

$$\begin{aligned} D_X(D_X(U)) &\stackrel{\text{def}}{=} [[U, \emptyset]_X, \emptyset]_X \\ &= [U^c, \emptyset]_X \\ &= (U^c)^c \\ &= U, \end{aligned}$$

where we have used:

- 0213 1. [Item 3](#) for the second and third equalities.  
 0214 2. [Item 3 of Proposition 4.3.11.1.2](#) for the fourth equality.

Since  $\mathcal{P}(X)$  is posetal, naturality is automatic (?? of ??), and thus we have

$$[[-, \emptyset]_X, \emptyset]_X \cong \text{id}_{\mathcal{P}(X)}$$

This finishes the proof.

**Item 6: Interaction With the Empty Set III**

Since  $D_X = (-)^c$ , this is essentially a repetition of the corresponding results for  $(-)^c$ , namely **Items 5 to 7 of Proposition 4.3.11.1.2**.

**Item 7: Interaction With Unions of Families of Subsets I**

By **Item 3 of Proposition 4.4.7.1.4**, we have

$$\begin{aligned} [\mathcal{U}, \emptyset]_{\mathcal{P}(X)} &= \mathcal{U}^c, \\ [U, \emptyset]_X &= U^c. \end{aligned}$$

With this, the counterexample given in the proof of **Item 10 of Proposition 4.3.6.1.2** then applies.

**Item 8: Interaction With Unions of Families of Subsets II**

We have

$$\begin{aligned} \left[ \bigcup_{U \in \mathcal{U}} U, V \right]_X &\stackrel{\text{def}}{=} \left( \bigcup_{U \in \mathcal{U}} U \right)^c \cup V \\ &= \left( \bigcap_{U \in \mathcal{U}} U^c \right) \cup V \\ &= \bigcap_{U \in \mathcal{U}} (U^c \cup V) \\ &\stackrel{\text{def}}{=} \bigcap_{U \in \mathcal{U}} [U, V]_X, \end{aligned}$$

where we have used:

0215 1. **Item 11 of Proposition 4.3.6.1.2** for the second equality.

0216 2. **Item 6 of Proposition 4.3.7.1.2** for the third equality.

This finishes the proof.

**Item 9: Interaction With Unions of Families of Subsets III**

We have

$$\bigcup_{V \in \mathcal{V}} [U, V]_X \stackrel{\text{def}}{=} \bigcup_{V \in \mathcal{V}} (U^c \cup V)$$

$$\begin{aligned}
&= U^c \cup \left( \bigcup_{V \in \mathcal{V}} V \right) \\
&\stackrel{\text{def}}{=} \left[ U, \bigcup_{V \in \mathcal{V}} V \right]_X.
\end{aligned}$$

where we have used **Item 6**. This finishes the proof.

**Item 10: Interaction With Intersections of Families of Subsets I**

Let  $X = \{0, 1\}$ , let  $\mathcal{U} = \{\{0, 1\}\}$ , and let  $\mathcal{V} = \{\{0\}, \{0, 1\}\}$ . We have

$$\begin{aligned}
\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W &= \bigcap_{W \in \mathcal{P}(X)} W \\
&= \{0, 1\},
\end{aligned}$$

whereas

$$\begin{aligned}
\left[ \bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X &= [\{0, 1\}, \{0\}] \\
&= \{0\},
\end{aligned}$$

Thus we have

$$\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W = \{0, 1\} \neq \{0\} = \left[ \bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X.$$

This finishes the proof.

**Item 11: Interaction With Intersections of Families of Subsets II**

We have

$$\begin{aligned}
\left[ \bigcap_{U \in \mathcal{U}} U, V \right]_X &\stackrel{\text{def}}{=} \left( \bigcap_{U \in \mathcal{U}} U \right)^c \cup V \\
&= \left( \bigcup_{U \in \mathcal{U}} U^c \right) \cup V \\
&= \bigcup_{U \in \mathcal{U}} (U^c \cup V) \\
&\stackrel{\text{def}}{=} \bigcup_{U \in \mathcal{U}} [U, V]_X,
\end{aligned}$$

where we have used:

- 0217     1. Item 12 of Proposition 4.3.6.1.2 for the second equality.  
 0218     2. Item 6 of Proposition 4.3.7.1.2 for the third equality.

This finishes the proof.

**Item 12: Interaction With Intersections of Families of Subsets III**

We have

$$\begin{aligned} \bigcap_{V \in \mathcal{V}} [U, V]_X &\stackrel{\text{def}}{=} \bigcap_{V \in \mathcal{V}} (U^c \cup V) \\ &= U^c \cup \left( \bigcap_{V \in \mathcal{V}} V \right) \\ &\stackrel{\text{def}}{=} \left[ U, \bigcap_{V \in \mathcal{V}} V \right]_X. \end{aligned}$$

where we have used Item 6. This finishes the proof.

**Item 13: Interaction With Binary Unions**

We have

$$\begin{aligned} [U \cap V, W]_X &\stackrel{\text{def}}{=} (U \cap V)^c \cup W \\ &= (U^c \cup V^c) \cup W \\ &= (U^c \cup V^c) \cup (W \cup W) \\ &= (U^c \cup W) \cup (V^c \cup W) \\ &\stackrel{\text{def}}{=} [U, W]_X \cup [V, W]_X, \end{aligned}$$

where we have used:

- 0219     1. Item 2 of Proposition 4.3.11.1.2 for the second equality.  
 021A     2. Item 8 of Proposition 4.3.8.1.2 for the third equality.  
 021B     3. Several applications of Items 2 and 4 of Proposition 4.3.8.1.2 and for the fourth equality.

For the second equality in the statement, we have

$$\begin{aligned} [U, V \cap W]_X &\stackrel{\text{def}}{=} U^c \cup (V \cap W) \\ &= (U^c \cup V) \cap (U^c \cap W) \\ &\stackrel{\text{def}}{=} [U, V]_X \cap [U, W]_X, \end{aligned}$$

where we have used Item 6 of Proposition 4.3.8.1.2 for the second equality.

Item 14: Interaction With Binary Intersections

We have

$$\begin{aligned} [U \cup V, W]_X &\stackrel{\text{def}}{=} (U \cup V)^c \cup W \\ &= (U^c \cap V^c) \cup W \\ &= (U^c \cup W) \cap (V^c \cup W) \\ &\stackrel{\text{def}}{=} [U, W]_X \cap [V, W]_X, \end{aligned}$$

where we have used:

- 021C 1. Item 2 of Proposition 4.3.11.1.2 for the second equality.  
 021D 2. Item 6 of Proposition 4.3.8.1.2 for the third equality.

Now, for the second equality in the statement, we have

$$\begin{aligned} [U, V \cup W]_X &\stackrel{\text{def}}{=} U^c \cup (V \cup W) \\ &= (U^c \cup U^c) \cup (V \cup W) \\ &= (U^c \cup V) \cup (U^c \cup W) \\ &\stackrel{\text{def}}{=} [U, V]_X \cup [U, W]_X, \end{aligned}$$

where we have used:

- 021E 1. Item 8 of Proposition 4.3.8.1.2 for the second equality.  
 021F 2. Several applications of Items 2 and 4 of Proposition 4.3.8.1.2 and for the third equality.

This finishes the proof.

Item 15: Interaction With Differences

We have

$$\begin{aligned} [U \setminus V, W]_X &\stackrel{\text{def}}{=} (U \setminus V)^c \cup W \\ &\stackrel{\text{def}}{=} (X \setminus (U \setminus V)) \cup W \\ &= ((X \cap V) \cup (X \setminus U)) \cup W \\ &= (V \cup (X \setminus U)) \cup W \\ &\stackrel{\text{def}}{=} (V \cup U^c) \cup W \\ &= (V \cup (U^c \cup U^c)) \cup W \\ &= (U^c \cup W) \cup (U^c \cup V) \\ &\stackrel{\text{def}}{=} [U, W]_X \cup [U, V]_X, \end{aligned}$$

where we have used:

- 021G 1. Item 10 of Proposition 4.3.10.1.2 for the third equality.
- 021H 2. Item 4 of Proposition 4.3.9.1.2 for the fourth equality.
- 021J 3. Item 8 of Proposition 4.3.8.1.2 for the sixth equality.
- 021K 4. Several applications of Items 2 and 4 of Proposition 4.3.8.1.2 and for the seventh equality.

We also have

$$\begin{aligned}
 [U \setminus V, W]_X &\stackrel{\text{def}}{=} (U \setminus V)^c \cup W \\
 &\stackrel{\text{def}}{=} (X \setminus (U \setminus V)) \cup W \\
 &= ((X \cap V) \cup (X \setminus U)) \cup W \\
 &= (V \cup (X \setminus U)) \cup W \\
 &\stackrel{\text{def}}{=} (V \cup U^c) \cup W \\
 &= (V \cup U^c) \cup (W \cup W) \\
 &= (U^c \cup W) \cup (V \cup W) \\
 &= (U^c \cup W) \cup ((V^c)^c \cup W) \\
 &\stackrel{\text{def}}{=} [U, W]_X \cup [V^c, W]_X,
 \end{aligned}$$

where we have used:

- 021L 1. Item 10 of Proposition 4.3.10.1.2 for the third equality.
- 021M 2. Item 4 of Proposition 4.3.9.1.2 for the fourth equality.
- 021N 3. Item 8 of Proposition 4.3.8.1.2 for the sixth equality.
- 021P 4. Several applications of Items 2 and 4 of Proposition 4.3.8.1.2 and for the seventh equality.
- 021Q 5. Item 3 of Proposition 4.3.11.1.2 for the eighth equality.

Now, for the second equality in the statement, we have

$$\begin{aligned}
 [U, V \setminus W]_X &\stackrel{\text{def}}{=} U^c \cup (V \setminus W) \\
 &= (V \setminus W) \cup U^c \\
 &= (V \cup U^c) \setminus (W \setminus U^c) \\
 &\stackrel{\text{def}}{=} (V \cup U^c) \setminus (W \setminus (X \setminus U)) \\
 &= (V \cup U^c) \setminus ((W \cap U) \cup (W \setminus X)) \\
 &= (V \cup U^c) \setminus ((W \cap U) \cup \emptyset)
 \end{aligned}$$

$$\begin{aligned}
 &= (V \cup U^c) \setminus (W \cap U) \\
 &= (V \cup U^c) \setminus (U \cap W) \\
 &\stackrel{\text{def}}{=} [U, V]_X \setminus (U \cap W)
 \end{aligned}$$

where we have used:

- 021R 1. Item 4 of Proposition 4.3.8.1.2 for the second equality.
- 021S 2. Item 4 of Proposition 4.3.10.1.2 for the third equality.
- 021T 3. Item 10 of Proposition 4.3.10.1.2 for the fifth equality.
- 021U 4. Item 13 of Proposition 4.3.10.1.2 for the sixth equality.
- 021V 5. Item 3 of Proposition 4.3.8.1.2 for the seventh equality.
- 021W 6. Item 5 of Proposition 4.3.9.1.2 for the eighth equality.

This finishes the proof.

#### Item 16: Interaction With Complements

We have

$$\begin{aligned}
 [U^c, V]_X &\stackrel{\text{def}}{=} (U^c)^c \cup V, \\
 &= U \cup V,
 \end{aligned}$$

where we have used Item 3 of Proposition 4.3.11.1.2. We also have

$$\begin{aligned}
 [U, V^c]_X &\stackrel{\text{def}}{=} U^c \cup V^c \\
 &= U \cap V
 \end{aligned}$$

where we have used Item 2 of Proposition 4.3.11.1.2. Finally, we have

$$\begin{aligned}
 [U, V]^c_X &= ((U \setminus V)^c)^c \\
 &= U \setminus V,
 \end{aligned}$$

where we have used Item 2 of Proposition 4.3.11.1.2.

#### Item 17: Interaction With Characteristic Functions

We have

$$\begin{aligned}
 \chi_{[U, V]_{\mathcal{P}(X)}}(x) &\stackrel{\text{def}}{=} \chi_{U^c \cup V}(x) \\
 &= \max(\chi_{U^c}, \chi_V) \\
 &= \max(1 - \chi_U \pmod{2}, \chi_V),
 \end{aligned}$$

where we have used:

021X

1. Item 10 of Proposition 4.3.8.1.2 for the second equality.
2. Item 4 of Proposition 4.3.11.1.2 for the third equality.

This finishes the proof.

**Item 18: Interaction With Direct Images**

This is a repetition of Item 10 of Proposition 4.6.1.1.5 and is proved there.

**Item 19: Interaction With Inverse Images**

This is a repetition of Item 10 of Proposition 4.6.2.1.3 and is proved there.

**Item 20: Interaction With Codirect Images**

This is a repetition of Item 9 of Proposition 4.6.3.1.7 and is proved there.

■

### 01L1 4.4.8 Isbell Duality for Sets

Let  $X$  be a set.

01L2

#### DEFINITION 4.4.8.1.1 ► THE ISBELL FUNCTION

The **Isbell function** of  $X$  is the map

$$! : \mathcal{P}(X) \rightarrow \text{Sets}(X, \mathcal{P}(X))$$

defined by

$$!(U) \stackrel{\text{def}}{=} [\![x \mapsto [U, \{x\}]_X]\!]$$

for each  $U \in \mathcal{P}(X)$ .

01L3

#### REMARK 4.4.8.1.2 ► MOTIVATION FOR THE ISBELL FUNCTION

Recall from Remark 4.4.1.1.2 that we may view the powerset  $\mathcal{P}(X)$  of a set  $X$  as the decategorification of the category of presheaves  $\text{PSh}(C)$  of a category  $C$ . Building upon this analogy, we want to mimic the definition of the Isbell Spec functor, which is given on objects by

$$\text{Spec}(\mathcal{F}) \stackrel{\text{def}}{=} \text{Nat}(\mathcal{F}, h_{(-)})$$

for each  $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$ . To this end, we could define

$$!(U) \stackrel{\text{def}}{=} [U, \chi_{(-)}]_X,$$

replacing:

- The Yoneda embedding  $X \mapsto h_X$  of  $C$  into  $\text{PSh}(C)$  with the characteristic embedding  $x \mapsto \chi_x$  of  $X$  into  $\mathcal{P}(X)$  of [Definition 4.5.4.1.1](#).
- The internal Hom Nat of  $\text{PSh}(C)$  with the internal Hom  $[-, -]_X$  of  $\mathcal{P}(X)$  of [Proposition 4.4.7.1.1](#).

However, since  $[U, \chi_x]_X$  is a subset of  $U$  instead of a truth value, we get a function

$$l: \mathcal{P}(X) \rightarrow \text{Sets}(X, \mathcal{P}(X))$$

instead of a function

$$l: \mathcal{P}(X) \rightarrow \mathcal{P}(X).$$

This makes some of the properties involving  $l$  a bit more cumbersome to state, although we still have an analogue of Isbell duality in that  $l_l \circ l$  evaluates to  $\text{id}_{\mathcal{P}(X)}$  in the sense of [Proposition 4.4.8.1.3](#).

01L4

#### PROPOSITION 4.4.8.1.3 ► ISBELL DUALITY FOR SETS

The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{l} & \text{Sets}(X, \mathcal{P}(X)) \\ & \searrow \Delta_{\text{id}_{\mathcal{P}(X)}} & \downarrow l_l \\ & & \text{Sets}(X, \text{Sets}(X, \mathcal{P}(X))) \end{array}$$

commutes, i.e. we have

$$l_l(l(U)) = [[x \mapsto [y \mapsto U]]]$$

for each  $U \in \mathcal{P}(X)$ .

#### PROOF 4.4.8.1.4 ► PROOF OF PROPOSITION 4.4.8.1.3

We have

$$\begin{aligned} l_l(l(U)) &\stackrel{\text{def}}{=} l_l([x \mapsto U^c \cup \{x\}]) \\ &\stackrel{\text{def}}{=} [x \mapsto l(U^c \cup \{x\})] \\ &\stackrel{\text{def}}{=} [x \mapsto [y \mapsto (U^c \cup \{x\})^c \cup \{x\}]] \\ &= [x \mapsto [y \mapsto (U \cap (X \setminus \{x\})) \cup \{x\}]] \\ &= [x \mapsto [y \mapsto (U \setminus \{x\}) \cup \{x\}]] \\ &= [x \mapsto [y \mapsto U]], \end{aligned}$$

where we have used Item 2 of Proposition 4.3.11.1.2 for the fourth equality above.

## 01L5 4.5 Characteristic Functions

### 005X 4.5.1 The Characteristic Function of a Subset

Let  $X$  be a set and let  $U \in \mathcal{P}(X)$ .

#### 005Z DEFINITION 4.5.1.1.1 ► THE CHARACTERISTIC FUNCTION OF A SUBSET

The **characteristic function** of  $U^1$  is the function  $\chi_U: X \rightarrow \{\text{t, f}\}^2$  defined by

$$\chi_U(x) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in U, \\ \text{false} & \text{if } x \notin U \end{cases}$$

for each  $x \in X$ .

<sup>1</sup>Further Terminology: Also called the **indicator function** of  $U$ .

<sup>2</sup>Further Notation: Also written  $\chi_X(U, -)$  or  $\chi_X(-, U)$ .

#### 01L6 REMARK 4.5.1.1.2 ► CHARACTERISTIC FUNCTIONS OF SUBSETS AS DECATEGORIFICATIONS OF PRESHEAVES

Under the analogy that  $\{\text{t, f}\}$  should be the  $(-1)$ -categorical analogue of Sets, we may view a function

$$f: X \rightarrow \{\text{t, f}\}$$

as a decategorification of presheaves and copresheaves

$$\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets},$$

$$F: C \rightarrow \text{Sets}.$$

The characteristic functions  $\chi_U$  of the subsets of  $X$  are then the primordial examples of such functions (and, in fact, all of them).

#### 01L7 NOTATION 4.5.1.3 ► FURTHER NOTATION FOR CHARACTERISTIC FUNCTIONS

We will often employ the bijection  $\{\text{t, f}\} \cong \{0, 1\}$  to make use of the arithmetical operations defined on  $\{0, 1\}$  when discussing characteristic functions.

Examples of this include Items 4 to 11 of Proposition 4.5.1.4 below.

0069

**PROPOSITION 4.5.1.4 ► PROPERTIES OF CHARACTERISTIC FUNCTIONS OF SUBSETS**

01L8

Let  $X$  be a set.

01L9

- Functionality.* The assignment  $U \mapsto \chi_U$  defines a function

$$\chi_{(-)} : \mathcal{P}(X) \rightarrow \text{Sets}(X, \{\text{t}, \text{f}\}).$$

01LA

- Bijectivity.* The function  $\chi_{(-)}$  from **Item 1** is bijective.

- Naturality.* The collection

$$\{\chi_{(-)} : \mathcal{P}(X) \rightarrow \text{Sets}(X, \{\text{t}, \text{f}\})\}_{X \in \text{Obj}(\text{Sets})}$$

defines a natural isomorphism between  $\mathcal{P}^{-1}$  and  $\text{Sets}(-, \{\text{t}, \text{f}\})$ .

In particular, given a function  $f : X \rightarrow Y$ , the diagram

$$\begin{array}{ccc} \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \\ \chi_{(-)} \downarrow \lrcorner & & \downarrow \lrcorner \chi_{(-)} \\ \text{Sets}(Y, \{\text{t}, \text{f}\}) & \xrightarrow{f^*} & \text{Sets}(X, \{\text{t}, \text{f}\}) \end{array}$$

commutes, i.e. we have

$$\chi_V \circ f = \chi_{f^{-1}(V)}$$

for each  $V \in \mathcal{P}(Y)$ .

006B

- Interaction With Unions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each  $U, V \in \mathcal{P}(X)$ .

006C

- Interaction With Unions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each  $U, V \in \mathcal{P}(X)$ .

006D

- Interaction With Intersections I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each  $U, V \in \mathcal{P}(X)$ .

006E

7. *Interaction With Intersections II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each  $U, V \in \mathcal{P}(X)$ .

006F

8. *Interaction With Differences.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each  $U, V \in \mathcal{P}(X)$ .

006G

9. *Interaction With Complements.* We have

$$\chi_{U^c} = 1 - \chi_U$$

for each  $U \in \mathcal{P}(X)$ .

006H

10. *Interaction With Symmetric Differences.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each  $U, V \in \mathcal{P}(X)$ .

01LB

11. *Interaction With Internal Hom.* We have

$$\chi_{[U,V]_{\mathcal{P}(X)}} = \max(1 - \chi_U \pmod{2}, \chi_V)$$

for each  $U, V \in \mathcal{P}(X)$ .

#### PROOF 4.5.1.1.5 ► PROOF OF PROPOSITION 4.5.1.1.4

**Item 1: Functionality**

There is nothing to prove.

**Item 2: Bijectivity**

We proceed in three steps:

021Z

1. *The Inverse of  $\chi_{(-)}$ .* The inverse of  $\chi_{(-)}$  is the map

$$\Phi: \text{Sets}(X, \{\text{t, f}\}) \xrightarrow{\sim} \mathcal{P}(X),$$

defined by

$$\begin{aligned}\Phi(f) &\stackrel{\text{def}}{=} U_f \\ &\stackrel{\text{def}}{=} f^{-1}(\text{true}) \\ &\stackrel{\text{def}}{=} \{x \in X \mid f(x) = \text{true}\}\end{aligned}$$

for each  $f \in \text{Sets}(X, \{\text{t}, \text{f}\})$ .

**0220** 2. *Invertibility I.* We have

$$\begin{aligned}[\Phi \circ \chi_{(-)}](U) &\stackrel{\text{def}}{=} \Phi(\chi_U) \\ &\stackrel{\text{def}}{=} \chi_U^{-1}(\text{true}) \\ &\stackrel{\text{def}}{=} \{x \in X \mid \chi_U(x) = \text{true}\} \\ &\stackrel{\text{def}}{=} \{x \in X \mid x \in U\} \\ &= U \\ &\stackrel{\text{def}}{=} [\text{id}_{\mathcal{P}(X)}](U)\end{aligned}$$

for each  $U \in \mathcal{P}(X)$ . Thus, we have

$$\Phi \circ \chi_{(-)} = \text{id}_{\mathcal{P}(X)}.$$

**0221** 3. *Invertibility II.* We have

$$\begin{aligned}[\chi_{(-)} \circ \Phi](U) &\stackrel{\text{def}}{=} \chi_{\Phi(f)} \\ &\stackrel{\text{def}}{=} \chi_{f^{-1}(\text{true})} \\ &\stackrel{\text{def}}{=} \llbracket x \mapsto \begin{cases} \text{true} & \text{if } x \in f^{-1}(\text{true}) \\ \text{false} & \text{otherwise} \end{cases} \rrbracket \\ &= \llbracket x \mapsto f(x) \rrbracket \\ &= f \\ &\stackrel{\text{def}}{=} [\text{id}_{\text{Sets}(X, \{\text{t}, \text{f}\})}](f)\end{aligned}$$

for each  $f \in \text{Sets}(X, \{\text{t}, \text{f}\})$ . Thus, we have

$$\chi_{(-)} \circ \Phi = \text{id}_{\text{Sets}(X, \{\text{t}, \text{f}\})}.$$

This finishes the proof.

Item 3: Naturality

We proceed in two steps:

0222

1. *Naturality of  $\chi_{(-)}$ .* We have

$$\begin{aligned} [\chi_V \circ f](v) &\stackrel{\text{def}}{=} \chi_V(f(v)) \\ &= \begin{cases} \text{true} & \text{if } f(v) \in V, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if } v \in f^{-1}(V), \\ \text{false} & \text{otherwise} \end{cases} \\ &\stackrel{\text{def}}{=} \chi_{f^{-1}(V)}(v) \end{aligned}$$

for each  $v \in V$ .

0223

2. *Naturality of  $\Phi$ .* Since  $\chi_{(-)}$  is natural and a componentwise inverse to  $\Phi$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\Phi$  is also natural in each argument.

This finishes the proof.

#### Item 4: Interaction With Unions I

This is a repetition of Item 10 of Proposition 4.3.8.1.2 and is proved there.

#### Item 5: Interaction With Unions II

This is a repetition of Item 11 of Proposition 4.3.8.1.2 and is proved there.

#### Item 6: Interaction With Intersections I

This is a repetition of Item 10 of Proposition 4.3.9.1.2 and is proved there.

#### Item 7: Interaction With Intersections II

This is a repetition of Item 11 of Proposition 4.3.9.1.2 and is proved there.

#### Item 8: Interaction With Differences

This is a repetition of Item 16 of Proposition 4.3.10.1.2 and is proved there.

#### Item 9: Interaction With Complements

This is a repetition of Item 4 of Proposition 4.3.11.1.2 and is proved there.

#### Item 10: Interaction With Symmetric Differences

This is a repetition of Item 15 of Proposition 4.3.12.1.2 and is proved there.

**Item 11: Interaction With Internal Hom**

This is a repetition of [Item 17](#) of [Proposition 4.4.7.1.4](#) and is proved there.

0224

**REMARK 4.5.1.1.6 ► POWERSETS AS SETS OF FUNCTIONS AND UN/STRAIGHTENING**

The bijection

$$\mathcal{P}(X) \cong \text{Sets}(X, \{\text{t, f}\})$$

of [Item 2](#) of [Proposition 4.5.1.1.4](#), which

- Takes a subset  $U \hookrightarrow X$  of  $X$  and *straightens* it to a function  $\chi_U: X \rightarrow \{\text{true, false}\}$ ;
- Takes a function  $f: X \rightarrow \{\text{true, false}\}$  and *unstraightens* it to a subset  $f^{-1}(\text{true}) \hookrightarrow X$  of  $X$ ;

may be viewed as the  $(-1)$ -categorical version of the  $0$ -categorical un/sstraightening isomorphism between indexed and fibred sets

$$\overbrace{\underline{\text{Sets}}_{/X}}^{\text{def}} \cong \overbrace{\underline{\text{ISets}}_X}^{\text{def}} \underline{\text{Fun}}(X_{\text{disc}}, \text{Sets})$$

of [??](#). Here we view:

- Subsets  $U \hookrightarrow X$  as being analogous to  $X$ -fibred sets  $\phi_X: A \rightarrow X$ .
- Functions  $f: X \rightarrow \{\text{t, f}\}$  as being analogous to  $X$ -indexed sets  $A: X_{\text{disc}} \rightarrow \text{Sets}$ .

**01LC 4.5.2 The Characteristic Function of a Point**

Let  $X$  be a set and let  $x \in X$ .

0060

**DEFINITION 4.5.2.1.1 ► THE CHARACTERISTIC FUNCTION OF A POINT**

The **characteristic function** of  $x$  is the function<sup>1</sup>

$$\chi_x: X \rightarrow \{\text{t, f}\}$$

defined by

$$\chi_x \stackrel{\text{def}}{=} \chi_{\{x\}},$$

i.e. by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each  $y \in X$ .

<sup>1</sup>Further Notation: Also written  $\chi^x$ ,  $\chi_X(x, -)$ , or  $\chi_X(-, x)$ .

01LD

#### REMARK 4.5.2.1.2 ► CHARACTERISTIC FUNCTIONS OF POINTS AS DECATEGORIFICATIONS OF REPRESENTABLE PRESHEAVES

Expanding upon Remark 4.5.1.1.2, we may think of the characteristic function

$$\chi_x: X \rightarrow \{\text{t, f}\}$$

of an *element*  $x$  of  $X$  as a decategorification of the representable presheaf and of the representable copresheaf

$$\begin{aligned} h_X: C^{\text{op}} &\rightarrow \text{Sets}, \\ h^X: C &\rightarrow \text{Sets} \end{aligned}$$

associated of an *object*  $X$  of a category  $C$ .

01LE 4.5.3 The Characteristic Relation of a Set

Let  $X$  be a set.

0061

#### DEFINITION 4.5.3.1.1 ► THE CHARACTERISTIC RELATION OF A SET

The **characteristic relation on  $X$** <sup>1</sup> is the relation<sup>2</sup>

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\text{t, f}\}$$

on  $X$  defined by<sup>3</sup>

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each  $x, y \in X$ .

<sup>1</sup>Further Terminology: Also called the **identity relation on  $X$** .

<sup>2</sup>Further Notation: Also written  $\chi_{-2}^{-1}$ , or  $\sim_{\text{id}}$  in the context of relations.

<sup>3</sup>Under the bijection  $\text{Sets}(X \times X, \{\text{t, f}\}) \cong \mathcal{P}(X \times X)$  of Item 2 of Proposition 4.5.1.1.4, the relation  $\chi_X$  corresponds to the diagonal  $\Delta_X \subset X \times X$  of  $X$ .

01LF

#### REMARK 4.5.3.1.2 ► THE CHARACTERISTIC RELATION OF A SET AS A DECATEGORIFICATION OF THE HOM PROFUNCTOR

Expanding upon Remarks 4.5.1.1.2 and 4.5.2.1.2, we may view the characteristic relation

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\text{t, f}\}$$

of  $X$  as a decategorification of the Hom profunctor

$$\text{Hom}_C(-_1, -_2) : C^{\text{op}} \times C \rightarrow \text{Sets}$$

of a category  $C$ .

01LG

#### PROPOSITION 4.5.3.1.3 ► PROPERTIES OF CHARACTERISTIC RELATIONS

Let  $f : X \rightarrow Y$  be a function.

006A

##### 1. The Inclusion of Characteristic Relations Associated to a Function.

Let  $f : A \rightarrow B$  be a function. We have an inclusion<sup>1</sup>

$$\begin{array}{ccc} A \times A & \xrightarrow{f \times f} & B \times B \\ \chi_B \circ (f \times f) \subset \chi_A, & \swarrow \curvearrowright \searrow & \chi_B \\ & \{t, f\}. & \end{array}$$

<sup>1</sup>Note: This is the 0-categorical version of Definition 11.5.4.1.1.

#### PROOF 4.5.3.1.4 ► PROOF OF PROPOSITION 4.5.3.1.3

Item 1: The Inclusion of Characteristic Relations Associated to a Function.

The inclusion  $\chi_B(f(a), f(b)) \subset \chi_A(a, b)$  is equivalent to the statement “if  $a = b$ , then  $f(a) = f(b)$ ”, which is true. 

## 01LH 4.5.4 The Characteristic Embedding of a Set

Let  $X$  be a set.

0062

#### DEFINITION 4.5.4.1.1 ► THE CHARACTERISTIC EMBEDDING OF A SET

The **characteristic embedding**<sup>1</sup> of  $X$  into  $\mathcal{P}(X)$  is the function

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

defined by<sup>2</sup>

$$\begin{aligned} \chi_{(-)}(x) &\stackrel{\text{def}}{=} \chi_x \\ &= \{x\} \end{aligned}$$

for each  $x \in X$ .

<sup>1</sup>The name “characteristic embedding” is justified by Corollary 4.5.5.1.3, which gives an analogue of fully faithfulness for  $\chi_{(-)}$ .

<sup>2</sup>Here we are identifying  $\mathcal{P}(X)$  with  $\text{Sets}(X, \{t, f\})$  as per Item 2 of Proposition 4.5.1.1.4.

01LJ

**REMARK 4.5.4.1.2 ► THE CHARACTERISTIC EMBEDDING OF A SET AS A DECATEGORIFICATION OF THE YONEDA EMBEDDING**

Expanding upon [Remarks 4.5.1.1.2](#), [4.5.2.1.2](#) and [4.5.3.1.2](#), we may view the characteristic embedding

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

of  $X$  into  $\mathcal{P}(X)$  as a decategorification of the Yoneda embedding

$$\flat : C^{\text{op}} \hookrightarrow \text{PSh}(C)$$

of a category  $C$  into  $\text{PSh}(C)$ .

01LK

**PROPOSITION 4.5.4.1.3 ► PROPERTIES OF CHARACTERISTIC EMBEDDINGS**

Let  $f : X \rightarrow Y$  be a map of sets.

01LL

1. *Interaction With Functions.* We have

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ f_! \circ \chi_X = \chi_Y \circ f, & \chi_X \downarrow & \downarrow \chi_Y \\ \mathcal{P}(X) & \xrightarrow[f_!]{} & \mathcal{P}(Y). \end{array}$$

**PROOF 4.5.4.1.4 ► PROOF OF PROPOSITION 4.5.4.1.3**

Item 1: Interaction With Functions

Indeed, we have

$$\begin{aligned} [f_! \circ \chi_X](x) &\stackrel{\text{def}}{=} f_!(\chi_X(x)) \\ &\stackrel{\text{def}}{=} f_!(\{x\}) \\ &= \{f(x)\} \\ &\stackrel{\text{def}}{=} \chi_{X'}(f(x)) \\ &\stackrel{\text{def}}{=} [\chi_{X'} \circ f](x), \end{aligned}$$

for each  $x \in X$ , showing the desired equality. 

**006K 4.5.5 The Yoneda Lemma for Sets**

Let  $X$  be a set and let  $U \subset X$  be a subset of  $X$ .

006L

**PROPOSITION 4.5.5.1.1 ► THE YONEDA LEMMA FOR SETS**

We have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$$

for each  $x \in X$ , giving an equality of functions

$$\chi_{\mathcal{P}(X)}(\chi(-), \chi_U) = \chi_U,$$

where

$$\chi_{\mathcal{P}(X)}(U, V) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } U \subset V, \\ \text{false} & \text{otherwise.} \end{cases}$$

**PROOF 4.5.5.1.2 ► PROOF OF PROPOSITION 4.5.5.1.1**

We have

$$\begin{aligned} \chi_{\mathcal{P}(X)}(\chi_x, \chi_U) &\stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } \{x\} \subset U, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if } x \in U \\ \text{false} & \text{otherwise} \end{cases} \\ &\stackrel{\text{def}}{=} \chi_U(x). \end{aligned}$$

This finishes the proof. 

006M

**COROLLARY 4.5.5.1.3 ► THE CHARACTERISTIC EMBEDDING IS FULLY FAITHFUL**

The characteristic embedding is fully faithful, i.e., we have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_y) \cong \chi_X(x, y)$$

for each  $x, y \in X$ .

**PROOF 4.5.5.1.4 ► PROOF OF COROLLARY 4.5.5.1.3**

We have

$$\begin{aligned} \chi_{\mathcal{P}(X)}(\chi_x, \chi_y) &= \chi_y(x) \\ &\stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in \{y\} \\ \text{false} & \text{otherwise} \end{cases} \end{aligned}$$

$$= \begin{cases} \text{true} & \text{if } x = y \\ \text{false} & \text{otherwise} \end{cases} \\ \stackrel{\text{def}}{=} \chi_X(x, y).$$

where we have used [Proposition 4.5.5.1.1](#) for the first equality. 

## 01LM 4.6 The Adjoint Triple $f_! \dashv f^{-1} \dashv f_*$

### 007F 4.6.1 Direct Images

Let  $f: X \rightarrow Y$  be a function.

007G

#### DEFINITION 4.6.1.1.1 ► DIRECT IMAGES

The **direct image function associated to  $f$**  is the function<sup>1</sup>

$$f_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by<sup>2</sup>

$$\begin{aligned} f_!(U) &\stackrel{\text{def}}{=} \left\{ y \in Y \mid \begin{array}{l} \text{there exists some } x \in U \\ \text{such that } y = f(x) \end{array} \right\} \\ &= \{f(x) \in Y \mid x \in U\} \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

<sup>1</sup>Further Notation: Also written simply  $f: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ .

<sup>2</sup>Further Terminology: The set  $f(U)$  is called the **direct image of  $U$  by  $f$** .

007H

#### NOTATION 4.6.1.1.2 ► FURTHER NOTATION FOR DIRECT IMAGES

Sometimes one finds the notation

$$\exists_f: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

for  $f_!$ . This notation comes from the fact that the following statements are equivalent, where  $y \in Y$  and  $U \in \mathcal{P}(X)$ :

- We have  $y \in \exists_f(U)$ .
- There exists some  $x \in U$  such that  $f(x) = y$ .

We will not make use of this notation elsewhere in Clowder.

0225

## WARNING 4.6.1.3 ► NOTATION FOR DIRECT IMAGES IS CONFUSING

0226

Notation for direct images between powersets is tricky:

0227

1. Direct images for powersets and presheaves are both adjoint to their corresponding inverse image functors. However, the direct image functor for powersets is a *left* adjoint, while the direct image functor for presheaves is a *right* adjoint:

- (a) *Powersets*. Given a function  $f: X \rightarrow Y$ , we have an inverse image functor

$$f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X).$$

The *left* adjoint of this functor is the usual direct image, defined above in [Definition 4.6.1.1.1](#).

0228

- (b) *Presheaves*. Given a morphism of topological spaces  $f: X \rightarrow Y$ , we have an inverse image functor

$$f^{-1}: \mathbf{PSh}(Y) \rightarrow \mathbf{PSh}(X).$$

The *right* adjoint of this functor is the direct image functor of presheaves, defined in ??.

0229

2. The presheaf direct image functor is denoted  $f_*$ , but the direct image functor for powersets is denoted  $f_!$  (as it's a left adjoint).

022A

3. Adding to the confusion, it's somewhat common for  $f_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$  to be denoted  $f_*$ .

We chose to write  $f_!$  for the direct image to keep the notation aligned with the following similar adjoint situations:

SITUATION	ADJOINT STRING
Functionality of Powersets	$(f_! \dashv f^{-1} \dashv f_*): \mathcal{P}(X) \xrightarrow{\sim} \mathcal{P}(Y)$
Functionality of Presheaf Categories	$(f_! \dashv f^{-1} \dashv f_*): \mathbf{PSh}(X) \xrightarrow{\sim} \mathbf{PSh}(Y)$
Base Change	$(f_! \dashv f^* \dashv f_*): \mathcal{C}_{/X} \xrightarrow{\sim} \mathcal{C}_{/Y}$
Kan Extensions	$(F_! \dashv F^* \dashv F_*): \mathbf{Fun}(\mathcal{C}, \mathcal{E}) \xrightarrow{\sim} \mathbf{Fun}(\mathcal{D}, \mathcal{E})$

007J

## REMARK 4.6.1.1.4 ► UNWINDING DEFINITION 4.6.1.1.1

Identifying  $\mathcal{P}(X)$  with  $\text{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$  via Item 2 of Proposition 4.5.1.1.4, we see that the direct image function associated to  $f$  is equivalently the function

$$f_! : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by

$$\begin{aligned} f_!(\chi_U) &\stackrel{\text{def}}{=} \text{Lan}_f(\chi_U) \\ &= \text{colim}((f \vec{\times} \underline{(-_1)}) \xrightarrow{\text{pr}} A \xrightarrow{\chi_U} \{\mathbf{t}, \mathbf{f}\}) \\ &= \text{colim}_{\substack{x \in X \\ f(x) = -1}} (\chi_U(x)) \\ &= \bigvee_{\substack{x \in X \\ f(x) = -1}} (\chi_U(x)), \end{aligned}$$

where we have used ?? for the second equality. In other words, we have

$$\begin{aligned} [f_!(\chi_U)](y) &= \bigvee_{\substack{x \in X \\ f(x) = y}} (\chi_U(x)) \\ &= \begin{cases} \text{true} & \text{if there exists some } x \in X \text{ such} \\ & \text{that } f(x) = y \text{ and } x \in U, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if there exists some } x \in U \\ & \text{such that } f(x) = y, \\ \text{false} & \text{otherwise} \end{cases} \end{aligned}$$

for each  $y \in Y$ .

007K

## PROPOSITION 4.6.1.5 ► PROPERTIES OF DIRECT IMAGES I

Let  $f: X \rightarrow Y$  be a function.

007L

1. *Functoriality.* The assignment  $U \mapsto f_!(U)$  defines a functor

$$f_! : (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset).$$

In particular, for each  $U, V \in \mathcal{P}(X)$ , the following condition is satisfied:

- (★) If  $U \subset V$ , then  $f_!(U) \subset f_!(V)$ .

007M

2. *Triple Adjunction*. We have a triple adjunction

$$(f_! \dashv f^{-1} \dashv f_*): \quad \mathcal{P}(X) \begin{array}{c} \xleftarrow{\quad f_! \quad} \\[-1ex] \perp \\[-1ex] \xrightarrow{\quad f_* \quad} \end{array} \mathcal{P}(Y),$$

witnessed by:

01LN

(a) Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\hookrightarrow f^{-1} \circ f_!, & \text{id}_{\mathcal{P}(Y)} &\hookrightarrow f_* \circ f^{-1}, \\ f_! \circ f^{-1} &\hookrightarrow \text{id}_{\mathcal{P}(Y)}, & f^{-1} \circ f_* &\hookrightarrow \text{id}_{\mathcal{P}(X)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset f^{-1}(f_!(U)), & V &\subset f_*(f^{-1}(V)), \\ f_!(f^{-1}(V)) &\subset V, & f^{-1}(f_*(U)) &\subset U \end{aligned}$$

indexed by  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ .

01LP

(b) Bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(Y)}(f_!(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(X)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f_*(V)), \end{aligned}$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$  and (respectively)  $V \in \mathcal{P}(X)$  and  $U \in \mathcal{P}(Y)$ . In particular:

01LQ

i. The following conditions are equivalent:

A. We have  $f_!(U) \subset V$ .

B. We have  $U \subset f^{-1}(V)$ .

01LR

ii. The following conditions are equivalent:

A. We have  $f^{-1}(U) \subset V$ .

B. We have  $U \subset f_*(V)$ .

01LS

01LT

01LU

01LV

01LW

3. *Interaction With Unions of Families of Subsets*. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_!)!} & \mathcal{P}(\mathcal{P}(Y)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_!(U) = \bigcup_{V \in f_!(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f_!)_!(\mathcal{U})$ .

01LX

4. *Interaction With Intersections of Families of Subsets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_!)_!} & \mathcal{P}(\mathcal{P}(Y)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_!(U) = \bigcap_{V \in f_!(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f_!)_!(\mathcal{U})$ .

01LY

5. *Interaction With Binary Unions.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U \cup V) = f_!(U) \cup f_!(V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01LZ

6. *Interaction With Binary Intersections.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & \curvearrowleft & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U \cap V) \subset f_!(U) \cap f_!(V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01M0

7. *Interaction With Differences.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow \backslash & \curvearrowright & \downarrow \backslash \\ \mathcal{P}(X) & \xrightarrow[f_!]{} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \setminus f_!(V) \subset f_!(U \setminus V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01M1

8. *Interaction With Complements.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow[f_!]{} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U^c) = f_*(U)^c$$

for each  $U \in \mathcal{P}(X)$ .

01M2

9. *Interaction With Symmetric Differences.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \curvearrowright & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow[f_!]{} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \Delta f_!(V) \subset f_!(U \Delta V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01M3

10. *Interaction With Internal Homs of Powersets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ [-1,-2]_X \downarrow & & \downarrow [-1,-2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have an equality of sets

$$f_!([U, V]_X) = [f_*(U), f_!(V)]_Y,$$

natural in  $U, V \in \mathcal{P}(X)$ .

007N

11. *Preservation of Colimits.* We have an equality of sets

$$f_! \left( \bigcup_{i \in I} U_i \right) = \bigcup_{i \in I} f_!(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f_!(U) \cup f_!(V) &= f_!(U \cup V), \\ f_!(\emptyset) &= \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

007P

12. *Oplax Preservation of Limits.* We have an inclusion of sets

$$f_! \left( \bigcap_{i \in I} U_i \right) \subset \bigcap_{i \in I} f_!(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} f_!(U \cap V) &\subset f_!(U) \cap f_!(V), \\ f_!(X) &\subset Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

007Q

13. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of Item 1 has a symmetric strict monoidal structure

$$(f_!, f_!^\otimes, f_{!|1}^\otimes): (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U) \cup f_!(V) &\xrightarrow{=} f_!(U \cup V), \\ f_{!|1}^\otimes: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

007R

14. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of Item 1 has a symmetric oplax monoidal structure

$$(f_!, f_!^\otimes, f_{!|1}^\otimes): (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with inclusions

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U \cap V) &\hookrightarrow f_!(U) \cap f_!(V), \\ f_{!|1}^\otimes: f_!(X) &\hookrightarrow Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

007S

15. *Interaction With Coproducts.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be maps of sets. We have

$$(f \sqcup g)_!(U \sqcup V) = f_!(U) \sqcup g_!(V)$$

for each  $U \in \mathcal{P}(X)$  and each  $V \in \mathcal{P}(Y)$ .

007T

16. *Interaction With Products.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be maps of sets. We have

$$(f \boxtimes_{X \times Y} g)_!(U \boxtimes_{X \times Y} V) = f_!(U) \boxtimes_{X' \times Y'} g_!(V)$$

for each  $U \in \mathcal{P}(X)$  and each  $V \in \mathcal{P}(Y)$ .

007U

17. *Relation to Codirect Images.* We have

$$\begin{aligned} f_!(U) &= f_*(U^c)^c \\ &\stackrel{\text{def}}{=} Y \setminus f_*(X \setminus U) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

#### PROOF 4.6.1.1.6 ► PROOF OF PROPOSITION 4.6.1.1.5

Item 1: Functoriality

Omitted.

Item 2: Triple Adjointness

This follows from Remark 4.6.1.1.4, Remark 4.6.2.1.2, Remark 4.6.3.1.4, and ?? of ??.

**Item 3: Interaction With Unions of Families of Subsets**

We have

$$\begin{aligned}\bigcup_{V \in f_!(\mathcal{U})} V &= \bigcup_{V \in \{f_!(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcup_{U \in \mathcal{U}} f_!(U).\end{aligned}$$

This finishes the proof.

**Item 4: Interaction With Intersections of Families of Subsets**

We have

$$\begin{aligned}\bigcap_{V \in f_!(\mathcal{U})} V &= \bigcap_{V \in \{f_!(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcap_{U \in \mathcal{U}} f_!(U).\end{aligned}$$

This finishes the proof.

**Item 5: Interaction With Binary Unions**

See [Pro25s].

**Item 6: Interaction With Binary Intersections**

See [Pro25q].

**Item 7: Interaction With Differences**

See [Pro25r].

**Item 8: Interaction With Complements**

Applying Item 17 to  $X \setminus U$ , we have

$$\begin{aligned}f_!(U^c) &= f_!(X \setminus U) \\ &= Y \setminus f_*(X \setminus (X \setminus U)) \\ &= Y \setminus f_*(U) \\ &= f_*(U)^c.\end{aligned}$$

This finishes the proof.

**Item 9: Interaction With Symmetric Differences**

We have

$$\begin{aligned}f_!(U) \Delta f_!(V) &= (f_!(U) \cup f_!(V)) \setminus (f_!(U) \cap f_!(V)) \\ &\subset (f_!(U) \cup f_!(V)) \setminus (f_!(U \cap V))\end{aligned}$$

$$\begin{aligned}
&= (f_!(U \cup V)) \setminus (f_!(U \cap V)) \\
&\subset f_!((U \cup V) \setminus (U \cap V)) \\
&= f_!(U \Delta V),
\end{aligned}$$

where we have used:

- 022C 1. Item 2 of Proposition 4.3.12.1.2 for the first equality.
- 022D 2. Item 6 of this proposition together with Item 1 of Proposition 4.3.10.1.2 for the first inclusion.
- 022E 3. Item 5 for the second equality.
- 022F 4. Item 7 for the second inclusion.
- 022G 5. Item 2 of Proposition 4.3.12.1.2 for the third equality.

Since  $\mathcal{P}(Y)$  is posetal, naturality is automatic (?? of ??). This finishes the proof.

#### Item 10: Interaction With Internal Homs of Powersets

We have

$$\begin{aligned}
f_!([U, V]_X) &\stackrel{\text{def}}{=} f_!(U^c \cup V) \\
&= f_!(U^c) \cup f_!(V) \\
&= f_*(U)^c \cup f_!(V) \\
&\stackrel{\text{def}}{=} [f_*(U), f_!(V)]_Y,
\end{aligned}$$

where we have used:

- 022H 1. Item 5 for the second equality.
- 022J 2. Item 17 for the third equality.

Since  $\mathcal{P}(Y)$  is posetal, naturality is automatic (?? of ??). This finishes the proof.

#### Item 11: Preservation of Colimits

This follows from Item 2 and ?? of ??.<sup>1</sup>

#### Item 12: Oplax Preservation of Limits

The inclusion  $f_!(X) \subset Y$  is automatic. See [Pro25q] for the other inclusions.

#### Item 13: Symmetric Strict Monoidality With Respect to Unions

This follows from Item 11.

**Item 14: Symmetric Oplax Monoidality With Respect to Intersections**

The inclusions in the statement follow from [Item 12](#). Since  $\mathcal{P}(Y)$  is posetal, the commutativity of the diagrams in the definition of a symmetric oplax monoidal functor is automatic ([??](#) of [??](#)).

**Item 15: Interaction With Coproducts**

Omitted.

**Item 16: Interaction With Products**

Omitted.

**Item 17: Relation to Codirect Images**

Applying [Item 16](#) of [Proposition 4.6.3.1.7](#) to  $X \setminus U$ , we have

$$\begin{aligned} f_*(X \setminus U) &= B \setminus f_!(X \setminus (X \setminus U)) \\ &= B \setminus f_!(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} f_!(U) &= B \setminus (B \setminus f_!(U)), \\ &= B \setminus f_*(X \setminus U), \end{aligned}$$

which finishes the proof. 

<sup>1</sup>Reference: [Pro25s].

007V

**PROPOSITION 4.6.1.7 ► PROPERTIES OF DIRECT IMAGES II**

Let  $f: X \rightarrow Y$  be a function.

007W

1. *Functionality I.* The assignment  $f \mapsto f_!$  defines a function

$$(-)_{*|X,Y}: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

007X

2. *Functionality II.* The assignment  $f \mapsto f_!$  defines a function

$$(-)_{*|X,Y}: \text{Sets}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

007Y

3. *Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have

$$(\text{id}_X)_! = \text{id}_{\mathcal{P}(X)}.$$

007Z

4. *Interaction With Composition.* For each pair of composable functions  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$ , we have

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \\ (g \circ f)_! = g_! \circ f_!, & \searrow^{(g \circ f)_!} & \downarrow g_! \\ & & \mathcal{P}(Z). \end{array}$$

#### PROOF 4.6.1.1.8 ► PROOF OF PROPOSITION 4.6.1.1.7

Item 1: Functionality I

There is nothing to prove.

Item 2: Functionality II

This follows from Item 1 of Proposition 4.6.1.1.5.

Item 3: Interaction With Identities

This follows from Remark 4.6.1.1.4 and ?? of ??.

Item 4: Interaction With Composition

This follows from Remark 4.6.1.1.4 and ?? of ??.



## 00SA 4.6.2 Inverse Images

Let  $f: X \rightarrow Y$  be a function.

0081

#### DEFINITION 4.6.2.1.1 ► INVERSE IMAGES

The **inverse image function associated to  $f$**  is the function<sup>1</sup>

$$f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$$

defined by<sup>2</sup>

$$f^{-1}(V) \stackrel{\text{def}}{=} \{x \in X \mid \text{we have } f(x) \in V\}$$

for each  $V \in \mathcal{P}(Y)$ .

<sup>1</sup>Further Notation: Also written  $f^*: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$ .

<sup>2</sup>Further Terminology: The set  $f^{-1}(V)$  is called the **inverse image of  $V$  by  $f$** .

0082

#### REMARK 4.6.2.1.2 ► UNWINDING DEFINITION 4.6.2.1.1

Identifying  $\mathcal{P}(Y)$  with  $\text{Sets}(Y, \{t, f\})$  via Item 2 of Proposition 4.5.1.1.4, we see that the inverse image function associated to  $f$  is equivalently

the function

$$f^*: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$$

defined by

$$f^*(\chi_V) \stackrel{\text{def}}{=} \chi_V \circ f$$

for each  $\chi_V \in \mathcal{P}(Y)$ , where  $\chi_V \circ f$  is the composition

$$X \xrightarrow{f} Y \xrightarrow{\chi_V} \{\text{true, false}\}$$

in Sets.

0083

#### PROPOSITION 4.6.2.1.3 ► PROPERTIES OF INVERSE IMAGES I

Let  $f: X \rightarrow Y$  be a function.

0084

1. *Functionality.* The assignment  $V \mapsto f^{-1}(V)$  defines a functor

$$f^{-1}: (\mathcal{P}(Y), \subset) \rightarrow (\mathcal{P}(X), \subset).$$

In particular, for each  $U, V \in \mathcal{P}(Y)$ , the following condition is satisfied:

(★) If  $U \subset V$ , then  $f^{-1}(U) \subset f^{-1}(V)$ .

0085

2. *Triple Adjunction.* We have a triple adjunction

$$(f_! \dashv f^{-1} \dashv f_*): \quad \mathcal{P}(X) \begin{array}{c} \xleftarrow{f_!} \\[-1ex] \perp \\[-1ex] \xleftarrow{f^{-1}} \end{array} \mathcal{P}(Y), \quad \mathcal{P}(Y) \begin{array}{c} \xrightarrow{f_*} \\[-1ex] \perp \\[-1ex] \xrightarrow{f^{-1}} \end{array} \mathcal{P}(X),$$

witnessed by:

01M4

- (a) Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\hookrightarrow f^{-1} \circ f_!, & \text{id}_{\mathcal{P}(Y)} &\hookrightarrow f_* \circ f^{-1}, \\ f_! \circ f^{-1} &\hookrightarrow \text{id}_{\mathcal{P}(Y)}, & f^{-1} \circ f_* &\hookrightarrow \text{id}_{\mathcal{P}(X)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset f^{-1}(f_!(U)), & V &\subset f_*(f^{-1}(V)), \\ f_!(f^{-1}(V)) &\subset V, & f^{-1}(f_*(U)) &\subset U \end{aligned}$$

indexed by  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ .

01M5

(b) *Bijections of sets*

$$\begin{aligned}\text{Hom}_{\mathcal{P}(Y)}(f_!(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(X)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f_*(V)),\end{aligned}$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$  and (respectively)  $V \in \mathcal{P}(X)$  and  $U \in \mathcal{P}(Y)$ . In particular:

01M6

- i. The following conditions are equivalent:

A. We have  $f_!(U) \subset V$ .

B. We have  $U \subset f^{-1}(V)$ .

01M7

- ii. The following conditions are equivalent:

A. We have  $f^{-1}(U) \subset V$ .

01M8

B. We have  $U \subset f_*(V)$ .

01M9

01MA

01MB

01MC

3. *Interaction With Unions of Families of Subsets.* The diagram

$$\begin{array}{ccc}\mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)\end{array}$$

commutes, i.e. we have

$$\bigcup_{V \in \mathcal{V}} f^{-1}(V) = \bigcup_{U \in f^{-1}(\mathcal{U})} U$$

for each  $\mathcal{V} \in \mathcal{P}(Y)$ , where  $f^{-1}(\mathcal{V}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{V})$ .

01MD

4. *Interaction With Intersections of Families of Subsets.* The diagram

$$\begin{array}{ccc}\mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)\end{array}$$

commutes, i.e. we have

$$\bigcap_{V \in \mathcal{V}} f^{-1}(V) = \bigcap_{U \in f^{-1}(\mathcal{U})} U$$

for each  $\mathcal{V} \in \mathcal{P}(Y)$ , where  $f^{-1}(\mathcal{V}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{V})$ .

01ME

5. *Interaction With Binary Unions.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cup V) = f^{-1}(U) \cup f^{-1}(V)$$

for each  $U, V \in \mathcal{P}(Y)$ .

01MF

6. *Interaction With Binary Intersections.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cap V) = f^{-1}(U) \cap f^{-1}(V)$$

for each  $U, V \in \mathcal{P}(Y)$ .

01MG

7. *Interaction With Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \setminus \downarrow & & \downarrow \setminus \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \setminus V) = f^{-1}(U) \setminus f^{-1}(V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01MH

8. *Interaction With Complements.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1,\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U^c) = f^{-1}(U)^c$$

for each  $U \in \mathcal{P}(X)$ .

01MJ

9. *Interaction With Symmetric Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \Delta \downarrow & & \downarrow \Delta \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

i.e. we have

$$f^{-1}(U) \Delta f^{-1}(V) = f^{-1}(U \Delta V)$$

for each  $U, V \in \mathcal{P}(Y)$ .

01MK

10. *Interaction With Internal Hom of Powersets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{-1,\text{op}} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ [-1, -2]_Y \downarrow & & \downarrow [-1, -2]_X \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$f^{-1}([U, V]_Y) = [f^{-1}(U), f^{-1}(V)]_X,$$

natural in  $U, V \in \mathcal{P}(X)$ .

0086

11. *Preservation of Colimits.* We have an equality of sets

$$f^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f^{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cup f^{-1}(V) &= f^{-1}(U \cup V), \\ f^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

0087

12. *Preservation of Limits.* We have an equality of sets

$$f^{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f^{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cap f^{-1}(V) &= f^{-1}(U \cap V), \\ f^{-1}(Y) &= X, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

0088

13. *Symmetric Strict Monoidality With Respect to Unions.* The inverse image function of [Item 1](#) has a symmetric strict monoidal structure

$$(f^{-1}, f^{-1, \otimes}, f_{\mathbb{1}}^{-1, \otimes}): (\mathcal{P}(Y), \cup, \emptyset) \rightarrow (\mathcal{P}(X), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{U,V}^{-1, \otimes}: f^{-1}(U) \cup f^{-1}(V) &\xrightarrow{\equiv} f^{-1}(U \cup V), \\ f_{\mathbb{1}}^{-1, \otimes}: \emptyset &\xrightarrow{\equiv} f^{-1}(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

0089

14. *Symmetric Strict Monoidality With Respect to Intersections.* The inverse image function of [Item 1](#) has a symmetric strict monoidal structure

$$(f^{-1}, f^{-1, \otimes}, f_{\mathbb{1}}^{-1, \otimes}): (\mathcal{P}(Y), \cap, Y) \rightarrow (\mathcal{P}(X), \cap, X),$$

being equipped with equalities

$$\begin{aligned} f_{U,V}^{-1, \otimes}: f^{-1}(U) \cap f^{-1}(V) &\xrightarrow{\equiv} f^{-1}(U \cap V), \\ f_{\mathbb{1}}^{-1, \otimes}: X &\xrightarrow{\equiv} f^{-1}(Y), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

008A

15. *Interaction With Coproducts.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be maps of sets. We have

$$(f \coprod g)^{-1}(U' \coprod V') = f^{-1}(U') \coprod g^{-1}(V')$$

for each  $U' \in \mathcal{P}(X')$  and each  $V' \in \mathcal{P}(Y')$ .

008B

16. *Interaction With Products.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be maps of sets. We have

$$(f \boxtimes_{X' \times Y'} g)^{-1}(U' \boxtimes_{X' \times Y'} V') = f^{-1}(U') \boxtimes_{X \times Y} g^{-1}(V')$$

for each  $U' \in \mathcal{P}(X')$  and each  $V' \in \mathcal{P}(Y')$ .

#### PROOF 4.6.2.1.4 ► PROOF OF PROPOSITION 4.6.2.1.3

**Item 1: Functoriality**

Omitted.

**Item 2: Triple Adjointness**

This follows from Remark 4.6.1.1.4, Remark 4.6.2.1.2, Remark 4.6.3.1.4, and ?? of ??.

**Item 3: Interaction With Unions of Families of Subsets**

We have

$$\begin{aligned} \bigcup_{U \in f^{-1}(\mathcal{V})} U &= \bigcup_{U \in \{f^{-1}(V) \in \mathcal{P}(X) \mid v \in \mathcal{V}\}} U \\ &= \bigcup_{V \in \mathcal{V}} f^{-1}(V). \end{aligned}$$

This finishes the proof.

**Item 4: Interaction With Intersections of Families of Subsets**

We have

$$\begin{aligned} \bigcap_{U \in f^{-1}(\mathcal{V})} U &= \bigcap_{U \in \{f^{-1}(V) \in \mathcal{P}(X) \mid v \in \mathcal{V}\}} U \\ &= \bigcap_{V \in \mathcal{V}} f^{-1}(V). \end{aligned}$$

This finishes the proof.

**Item 5: Interaction With Binary Unions**

See [Pro25ac].

Item 6: Interaction With Binary Intersections

See [Pro25aa].

Item 7: Interaction With Differences

See [Pro25ab].

Item 8: Interaction With Complements

See [Pro25j].

Item 9: Interaction With Symmetric Differences

We have

$$\begin{aligned} f^{-1}(U \Delta V) &= f^{-1}((U \cup V) \setminus (U \cap V)) \\ &= f^{-1}(U \cup V) \setminus f^{-1}(U \cap V) \\ &= f^{-1}(U) \cup f^{-1}(V) \setminus f^{-1}(U \cap V) \\ &= f^{-1}(U) \cup f^{-1}(V) \setminus f^{-1}(U) \cap f^{-1}(V) \\ &= f^{-1}(U) \Delta f^{-1}(V), \end{aligned}$$

where we have used:

022K 1. Item 2 of Proposition 4.3.12.1.2 for the first equality.

022L 2. Item 7 for the second equality.

022M 3. Item 5 for the third equality.

022N 4. Item 6 for the fourth equality.

022P 5. Item 2 of Proposition 4.3.12.1.2 for the fifth equality.

This finishes the proof.

Item 10: Interaction With Internal Homs of Powersets

We have

$$\begin{aligned} f^{-1}([U, V]_Y) &\stackrel{\text{def}}{=} f^{-1}(U^c \cup V) \\ &= f^{-1}(U^c) \cup f^{-1}(V) \\ &= f^{-1}(U)^c \cup f^{-1}(V) \\ &\stackrel{\text{def}}{=} [f^{-1}(U), f^{-1}(V)]_X, \end{aligned}$$

where we have used:

022Q 1. Item 8 for the second equality.

022R

2. Item 5 for the third equality.

Since  $\mathcal{P}(Y)$  is posetal, naturality is automatic (?? of ??). This finishes the proof.

Item 11: Preservation of Colimits

This follows from Item 2 and ?? of ??.<sup>1</sup>

Item 12: Preservation of Limits

This follows from Item 2 and ?? of ??.<sup>2</sup>

Item 13: Symmetric Strict Monoidality With Respect to Unions

This follows from Item 11.

Item 14: Symmetric Strict Monoidality With Respect to Intersections

This follows from Item 12.

Item 15: Interaction With Coproducts

Omitted.

Item 16: Interaction With Products

Omitted. 

<sup>1</sup>Reference: [Pro25ac].

<sup>2</sup>Reference: [Pro25aa].

008C

#### PROPOSITION 4.6.2.1.5 ► PROPERTIES OF INVERSE IMAGES II

Let  $f: X \rightarrow Y$  be a function.

008D

1. *Functionality I.* The assignment  $f \mapsto f^{-1}$  defines a function

$$(-)_{X,Y}^{-1}: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(Y), \mathcal{P}(X)).$$

008E

2. *Functionality II.* The assignment  $f \mapsto f^{-1}$  defines a function

$$(-)_{X,Y}^{-1}: \text{Sets}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(Y), \subset), (\mathcal{P}(X), \subset)).$$

008F

3. *Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have

$$\text{id}_X^{-1} = \text{id}_{\mathcal{P}(X)}.$$

008G

4. *Interaction With Composition.* For each pair of composable functions  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$ , we have

$$(g \circ f)^{-1} = f^{-1} \circ g^{-1}, \quad \begin{array}{ccc} \mathcal{P}(Z) & \xrightarrow{g^{-1}} & \mathcal{P}(Y) \\ & \searrow^{(g \circ f)^{-1}} & \downarrow f^{-1} \\ & & \mathcal{P}(X). \end{array}$$



Item 1: Functionality I

There is nothing to prove.

Item 2: Functionality II

This follows from Item 1 of Proposition 4.6.2.1.3.

Item 3: Interaction With Identities

This follows from Remark 4.6.2.1.2 and Item 5 of Proposition 11.1.4.1.2.

Item 4: Interaction With Composition

This follows from Remark 4.6.2.1.2 and Item 2 of Proposition 11.1.4.1.2.



### 008H 4.6.3 Codirect Images

Let  $f: X \rightarrow Y$  be a function.

008J

#### DEFINITION 4.6.3.1.1 ► CODIRECT IMAGES

The **codirect image function associated to  $f$**  is the function

$$f_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by<sup>1,2</sup>

$$\begin{aligned} f_*(U) &\stackrel{\text{def}}{=} \left\{ y \in Y \mid \begin{array}{l} \text{for each } x \in X, \text{ if we have} \\ f(x) = y, \text{ then } x \in U \end{array} \right\} \\ &= \{y \in Y \mid \text{we have } f^{-1}(y) \subset U\} \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

<sup>1</sup>Further Terminology: The set  $f_*(U)$  is called the **codirect image of  $U$  by  $f$** .

<sup>2</sup>We also have

$$\begin{aligned} f_*(U) &= f_!(U^c)^c \\ &\stackrel{\text{def}}{=} Y \setminus f_!(X \setminus U); \end{aligned}$$

see Item 16 of Proposition 4.6.3.1.7.

008K

#### NOTATION 4.6.3.1.2 ► FURTHER NOTATION FOR CODIRECT IMAGES

Sometimes one finds the notation

$$\forall_f: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

for  $f_!$ . This notation comes from the fact that the following statements are equivalent, where  $y \in Y$  and  $U \in \mathcal{P}(X)$ :

- We have  $y \in \forall_f(U)$ .
- For each  $x \in X$ , if  $y = f(x)$ , then  $x \in U$ .

We will not make use of this notation elsewhere in Clowder.

**WARNING 4.6.3.1.3 ► NOTATION FOR CODIRECT IMAGES IS CONFUSING**



See [Warning 4.6.1.1.3](#).

008L

**REMARK 4.6.3.1.4 ► UNWINDING DEFINITION 4.6.3.1.1**

Identifying  $\mathcal{P}(X)$  with  $\text{Sets}(X, \{\text{t}, \text{f}\})$  via [Item 2 of Proposition 4.5.1.1.4](#), we see that the codirect image function associated to  $f$  is equivalently the function

$$f_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by

$$\begin{aligned} f_*(\chi_U) &\stackrel{\text{def}}{=} \text{Ran}_f(\chi_U) \\ &= \lim((\underline{(-1)} \xrightarrow{\rightarrow} f) \xrightarrow{\text{pr}} X \xrightarrow{\chi_U} \{\text{true, false}\}) \\ &= \lim_{\substack{x \in X \\ f(x)=-1}} (\chi_U(x)) \\ &= \bigwedge_{\substack{x \in X \\ f(x)=-1}} (\chi_U(x)). \end{aligned}$$

where we have used ?? for the second equality. In other words, we have

$$\begin{aligned} [f_*(\chi_U)](y) &= \bigwedge_{\substack{x \in X \\ f(x)=y}} (\chi_U(x)) \\ &= \begin{cases} \text{true} & \text{if, for each } x \in X \text{ such that} \\ & f(x)=y, \text{ we have } x \in U, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if } f^{-1}(y) \subset U \\ \text{false} & \text{otherwise} \end{cases} \end{aligned}$$

for each  $y \in Y$ .

008M

DEFINITION 4.6.3.1.5 ► THE IMAGE AND COMPLEMENT PARTS OF  $f_*$ 

Let  $U$  be a subset of  $X$ .<sup>1,2</sup>

008N

1. The **image part of the codirect image**  $f_*(U)$  of  $U$  is the set  $f_{*,\text{im}}(U)$  defined by

$$\begin{aligned} f_{*,\text{im}}(U) &\stackrel{\text{def}}{=} f_*(U) \cap \text{Im}(f) \\ &= \left\{ y \in Y \mid \begin{array}{l} \text{we have } f^{-1}(y) \subset U \\ \text{and } f^{-1}(y) \neq \emptyset. \end{array} \right\}. \end{aligned}$$

008P

2. The **complement part of the codirect image**  $f_*(U)$  of  $U$  is the set  $f_{*,\text{cp}}(U)$  defined by

$$\begin{aligned} f_{*,\text{cp}}(U) &\stackrel{\text{def}}{=} f_*(U) \cap (Y \setminus \text{Im}(f)) \\ &= Y \setminus \text{Im}(f) \\ &= \left\{ y \in Y \mid \begin{array}{l} \text{we have } f^{-1}(y) \subset U \\ \text{and } f^{-1}(y) = \emptyset. \end{array} \right\} \\ &= \{y \in Y \mid f^{-1}(y) = \emptyset\}. \end{aligned}$$

---

<sup>1</sup>Note that we have

$$f_*(U) = f_{*,\text{im}}(U) \cup f_{*,\text{cp}}(U),$$

as

$$\begin{aligned} f_*(U) &= f_*(U) \cap Y \\ &= f_*(U) \cap (\text{Im}(f) \cup (Y \setminus \text{Im}(f))) \\ &= (f_*(U) \cap \text{Im}(f)) \cup (f_*(U) \cap (Y \setminus \text{Im}(f))) \\ &\stackrel{\text{def}}{=} f_{*,\text{im}}(U) \cup f_{*,\text{cp}}(U). \end{aligned}$$

<sup>2</sup>In terms of the meet computation of  $f_*(U)$  of Remark 4.6.3.1.4, namely

$$f_*(\chi_U) = \bigwedge_{\substack{x \in X \\ f(x)=-1}} (\chi_U(x)),$$

we see that  $f_{*,\text{im}}$  corresponds to meets indexed over nonempty sets, while  $f_{*,\text{cp}}$  corresponds to meets indexed over the empty set.

008Q

## EXAMPLE 4.6.3.1.6 ► EXAMPLES OF CODIRECT IMAGES

Here are some examples of codirect images.

0231

1. *Multiplication by Two.* Consider the function  $f: \mathbb{N} \rightarrow \mathbb{N}$  given by

$$f(n) \stackrel{\text{def}}{=} 2n$$

for each  $n \in \mathbb{N}$ . Since  $f$  is injective, we have

$$\begin{aligned} f_{*,\text{im}}(U) &= f_!(U) \\ f_{*,\text{cp}}(U) &= \{\text{odd natural numbers}\} \end{aligned}$$

for any  $U \subset \mathbb{N}$ . In particular, we have

$$f_*(\{\text{even natural numbers}\}) = \mathbb{N}.$$

0232

2. *Parabolas.* Consider the function  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by

$$f(x) \stackrel{\text{def}}{=} x^2$$

for each  $x \in \mathbb{R}$ . We have

$$f_{*,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any  $U \subset \mathbb{R}$ . Moreover, since  $f^{-1}(x) = \{-\sqrt{x}, \sqrt{x}\}$ , we have e.g.:

$$\begin{aligned} f_{*,\text{im}}([0, 1]) &= \{0\}, \\ f_{*,\text{im}}([-1, 1]) &= [0, 1], \\ f_{*,\text{im}}([1, 2]) &= \emptyset, \\ f_{*,\text{im}}([-2, -1] \cup [1, 2]) &= [1, 4]. \end{aligned}$$

0233

3. *Circles.* Consider the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  given by

$$f(x, y) \stackrel{\text{def}}{=} x^2 + y^2$$

for each  $(x, y) \in \mathbb{R}^2$ . We have

$$f_{*,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any  $U \subset \mathbb{R}^2$ , and since

$$f^{-1}(r) = \begin{cases} \text{a circle of radius } r \text{ about the origin} & \text{if } r > 0, \\ \{(0, 0)\} & \text{if } r = 0, \\ \emptyset & \text{if } r < 0, \end{cases}$$

we have e.g.:

$$\begin{aligned} f_{*,\text{im}}([-1, 1] \times [-1, 1]) &= [0, 1], \\ f_{*,\text{im}}(([ -1, 1] \times [-1, 1]) \setminus [-1, 1] \times \{0\}) &= \emptyset. \end{aligned}$$

008R

**PROPOSITION 4.6.3.1.7 ► PROPERTIES OF CODIRECT IMAGES I**

008S

Let  $f: X \rightarrow Y$  be a function.

008T

- Functoriality.* The assignment  $U \mapsto f_*(U)$  defines a functor

$$f_*: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset).$$

In particular, for each  $U, V \in \mathcal{P}(X)$ , the following condition is satisfied:

(★) If  $U \subset V$ , then  $f_*(U) \subset f_*(V)$ .

008T

- Triple Adjunction.* We have a triple adjunction

$$(f_! \dashv f^{-1} \dashv f_*): \quad \mathcal{P}(X) \begin{array}{c} \xleftarrow{\quad f_! \quad} \\[-1ex] \perp \\[-1ex] \xrightarrow{\quad f^{-1} \quad} \\[-1ex] \perp \\[-1ex] \xrightarrow{\quad f_* \quad} \end{array} \mathcal{P}(Y),$$

witnessed by:

01ML

- Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\hookrightarrow f^{-1} \circ f_!, & \text{id}_{\mathcal{P}(Y)} &\hookrightarrow f_* \circ f^{-1}, \\ f_! \circ f^{-1} &\hookrightarrow \text{id}_{\mathcal{P}(Y)}, & f^{-1} \circ f_* &\hookrightarrow \text{id}_{\mathcal{P}(X)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset f^{-1}(f_!(U)), & V &\subset f_*(f^{-1}(V)), \\ f_!(f^{-1}(V)) &\subset V, & f^{-1}(f_*(U)) &\subset U \end{aligned}$$

indexed by  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ .

01MM

- Bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(Y)}(f_!(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(X)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(Y)}(U, f_*(V)), \end{aligned}$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$  and (respectively)  $V \in \mathcal{P}(X)$  and  $U \in \mathcal{P}(Y)$ . In particular:

01MN

- The following conditions are equivalent:

A. We have  $f_!(U) \subset V$ .

B. We have  $U \subset f^{-1}(V)$ .

01MP

- The following conditions are equivalent:

A. We have  $f^{-1}(U) \subset V$ .

01MQ

01MR

01MS

01MT

B. We have  $U \subset f_*(V)$ .

01MU

3. *Interaction With Unions of Families of Subsets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_*(U) = \bigcup_{V \in f_*(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$ .

01MV

4. *Interaction With Intersections of Families of Subsets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_*(U) = \bigcap_{V \in f_*(\mathcal{U})} V$$

for each  $\mathcal{U} \in \mathcal{P}(X)$ , where  $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$ .

01MW

5. *Interaction With Binary Unions.* Let  $f: X \rightarrow Y$  be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cup \downarrow & \curvearrowleft & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U) \cup f_*(V) \subset f_*(U \cup V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01MX

6. *Interaction With Binary Intersections.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U) \cap f_*(V) = f_*(U \cap V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01MY

7. *Interaction With Complements.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U^c) = f_!(U)^c$$

for each  $U \in \mathcal{P}(X)$ .

01MZ

8. *Interaction With Symmetric Differences.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \curvearrowright & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U \Delta V) \subset f_*(U) \Delta f_*(V)$$

indexed by  $U, V \in \mathcal{P}(X)$ .

01N0

9. *Interaction With Internal Hom of Powersets.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ [-1,-2]_X \downarrow & \curvearrowright & \downarrow [-1,-2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$[f_!(U), f_*(V)]_Y \subset f_*( [U, V]_X )$$

indexed by  $U, V \in \mathcal{P}(X)$ .

008U

10. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} f_*(U_i) \subset f_* \left( \bigcup_{i \in I} U_i \right),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} f_*(U) \cup f_*(V) &\hookrightarrow f_*(U \cup V), \\ \emptyset &\hookrightarrow f_*(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

008V

11. *Preservation of Limits.* We have an equality of sets

$$f_* \left( \bigcap_{i \in I} U_i \right) = \bigcap_{i \in I} f_*(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f^{-1}(U \cap V) &= f_*(U) \cap f^{-1}(V), \\ f_*(X) &= Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

008W

12. *Symmetric Lax Monoidality With Respect to Unions.* The codirect image function of [Item 1](#) has a symmetric lax monoidal structure

$$(f_*, f_*^\otimes, f_{*|1}^\otimes): (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U) \cup f_*(V) &\hookrightarrow f_*(U \cup V), \\ f_{*|1}^\otimes: \emptyset &\hookrightarrow f_*(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

008X

13. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$(f_*, f_*^\otimes, f_{*\mid\mathbb{1}}^\otimes): (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with equalities

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U \cap V) &\xrightarrow{\cong} f_*(U) \cap f_*(V), \\ f_{*\mid\mathbb{1}}^\otimes: f_*(X) &\xrightarrow{\cong} Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

008Y

14. *Interaction With Coproducts.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be maps of sets. We have

$$(f \coprod g)_*(U \coprod V) = f_*(U) \coprod g_*(V)$$

for each  $U \in \mathcal{P}(X)$  and each  $V \in \mathcal{P}(Y)$ .

008Z

15. *Interaction With Products.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be maps of sets. We have

$$(f \boxtimes_{X \times Y} g)_*(U \boxtimes_{X \times Y} V) = f_*(U) \boxtimes_{X' \times Y'} g_*(V)$$

for each  $U \in \mathcal{P}(X)$  and each  $V \in \mathcal{P}(Y)$ .

0090

16. *Relation to Direct Images.* We have

$$\begin{aligned} f_*(U) &= f_!(U^c)^c \\ &= Y \setminus f_!(X \setminus U) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

0091

17. *Interaction With Injections.* If  $f$  is injective, then we have

$$\begin{aligned} f_{*,\text{im}}(U) &= f_!(U), \\ f_{*,\text{cp}}(U) &= Y \setminus \text{Im}(f), \end{aligned}$$

and so

$$\begin{aligned} f_*(U) &= f_{*,\text{im}}(U) \cup f_{*,\text{cp}}(U) \\ &= f_!(U) \cup (Y \setminus \text{Im}(f)) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

0092

18. *Interaction With Surjections.* If  $f$  is surjective, then we have

$$\begin{aligned} f_{*,\text{im}}(U) &\subset f_!(U), \\ f_{*,\text{cp}}(U) &= \emptyset, \end{aligned}$$

and so

$$f_*(U) \subset f_!(U)$$

for each  $U \in \mathcal{P}(X)$ .

#### PROOF 4.6.3.1.8 ► PROOF OF PROPOSITION 4.6.3.1.7

**Item 1: Functoriality**

Omitted.

**Item 2: Triple Adjunctness**

This follows from Remark 4.6.1.1.4, Remark 4.6.2.1.2, Remark 4.6.3.1.4, and ?? of ??.

**Item 3: Interaction With Unions of Families of Subsets**

We have

$$\begin{aligned} \bigcup_{V \in f_*(\mathcal{U})} V &= \bigcup_{V \in \{f_*(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcup_{U \in \mathcal{U}} f_*(U). \end{aligned}$$

This finishes the proof.

**Item 4: Interaction With Intersections of Families of Subsets**

We have

$$\begin{aligned} \bigcap_{V \in f_*(\mathcal{U})} V &= \bigcap_{V \in \{f_*(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcap_{U \in \mathcal{U}} f_*(U). \end{aligned}$$

This finishes the proof.

**Item 5: Interaction With Binary Unions**

We have

$$f_*(U) \cup f_*(V) = f_!(U^c)^c \cup f_!(V^c)^c$$

$$\begin{aligned}
&= (f_!(U^c) \cap f_!(V^c))^c \\
&\subset (f_!(U^c \cap V^c))^c \\
&= f_!((U \cup V)^c)^c \\
&= f_*(U \cup V),
\end{aligned}$$

where:

- 023X 1. We have used [Item 16](#) for the first equality.
- 023Y 2. We have used [Item 2](#) of [Proposition 4.3.11.1.2](#) for the second equality.
- 023Z 3. We have used [Item 6](#) of [Proposition 4.6.1.1.5](#) for the third equality.
- 0240 4. We have used [Item 2](#) of [Proposition 4.3.11.1.2](#) for the fourth equality.
- 0241 5. We have used [Item 16](#) for the last equality.

This finishes the proof.

[Item 6: Interaction With Binary Intersections](#)

This follows from [Item 11](#).

[Item 7: Interaction With Complements](#)

Omitted.

[Item 8: Interaction With Symmetric Differences](#)

Omitted.

[Item 9: Interaction With Internal Homs of Powersets](#)

We have

$$\begin{aligned}
[f_!(U), f_!(V)]_X &\stackrel{\text{def}}{=} f_!(U)^c \cup f_*(V) \\
&= f_*(U^c) \cup f_*(V) \\
&\subset f_*(U^c \cup V) \\
&\stackrel{\text{def}}{=} f_*( [U, V]_X ),
\end{aligned}$$

where we have used:

- 0242 1. [Item 7 of Proposition 4.6.3.1.7](#) for the second equality.
- 0243 2. [Item 5 of Proposition 4.6.3.1.7](#) for the inclusion.

Since  $\mathcal{P}(X)$  is posetal, naturality is automatic (?? of ??). This finishes the proof.

**Item 10: Lax Preservation of Colimits**

Omitted.

**Item 11: Preservation of Limits**

This follows from **Item 2** and ?? of ??.

**Item 12: Symmetric Lax Monoidality With Respect to Unions**

This follows from **Item 10**.

**Item 13: Symmetric Strict Monoidality With Respect to Intersections**

This follows from **Item 11**.

**Item 14: Interaction With Coproducts**

Omitted.

**Item 15: Interaction With Products**

Omitted.

**Item 16: Relation to Direct Images**

We claim that  $f_*(U) = Y \setminus f_!(X \setminus U)$ .

- *The First Implication.* We claim that

$$f_*(U) \subset Y \setminus f_!(X \setminus U).$$

Let  $y \in f_*(U)$ . We need to show that  $y \notin f_!(X \setminus U)$ , i.e. that there is no  $x \in X \setminus U$  such that  $f(x) = y$ .

This is indeed the case, as otherwise we would have  $x \in f^{-1}(y)$  and  $x \notin U$ , contradicting  $f^{-1}(y) \subset U$  (which holds since  $y \in f_*(U)$ ).

Thus  $y \in Y \setminus f_!(X \setminus U)$ .

- *The Second Implication.* We claim that

$$Y \setminus f_!(X \setminus U) \subset f_*(U).$$

Let  $y \in Y \setminus f_!(X \setminus U)$ . We need to show that  $y \in f_*(U)$ , i.e. that  $f^{-1}(y) \subset U$ .

Since  $y \notin f_!(X \setminus U)$ , there exists no  $x \in X \setminus U$  such that  $y = f(x)$ , and hence  $f^{-1}(y) \subset U$ .

Thus  $y \in f_*(U)$ .

This finishes the proof of **Item 16**.

Item 17: Interaction With Injections

Omitted.

Item 18: Interaction With Surjections

Omitted. 

0093

#### PROPOSITION 4.6.3.1.9 ► PROPERTIES OF CODIRECT IMAGES II

Let  $f: X \rightarrow B$  be a function.

0094

1. *Functionality I.* The assignment  $f \mapsto f_*$  defines a function

$$(-)_{!|X,Y}: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

0095

2. *Functionality II.* The assignment  $f \mapsto f_*$  defines a function

$$(-)_{!|X,Y}: \text{Sets}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

0096

3. *Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have

$$(\text{id}_X)_* = \text{id}_{\mathcal{P}(X)}.$$

0097

4. *Interaction With Composition.* For each pair of composable functions  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$ , we have

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \\ (g \circ f)_* = g_* \circ f_*, & \searrow & \downarrow g_* \\ & & \mathcal{P}(Z). \end{array}$$

#### PROOF 4.6.3.1.10 ► PROOF OF PROPOSITION 4.6.3.1.9

Item 1: Functionality I

There is nothing to prove.

Item 2: Functionality II

This follows from Item 1 of Proposition 4.6.3.1.7.

Item 3: Interaction With Identities

This follows from Remark 4.6.3.1.4 and ?? of ??.

## Item 4: Interaction With Composition

This follows from Remark 4.6.3.1.4 and ?? of ??.



## 01N1 4.6.4 A Six-Functor Formalism for Sets

01N2

## REMARK 4.6.4.1.1 ► A SIX-FUNCTOR FORMALISM FOR SETS

The assignment  $X \mapsto \mathcal{P}(X)$  together with the functors  $f_*$ ,  $f^{-1}$ , and  $f_!$  of Item 1 of Proposition 4.6.1.1.5, Item 1 of Proposition 4.6.2.1.3, and Item 1 of Proposition 4.6.3.1.7, and the functors

$$\begin{aligned} -_1 \cap -_2 : \mathcal{P}(X) \times \mathcal{P}(X) &\rightarrow \mathcal{P}(X), \\ [-_1, -_2]_X : \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) &\rightarrow \mathcal{P}(X) \end{aligned}$$

of Item 1 of Proposition 4.3.9.1.2 and Item 1 of Proposition 4.4.7.1.4 satisfy several properties reminiscent of a six functor formalism in the sense of ??.

We collect these properties in Proposition 4.6.4.1.2 below.<sup>1</sup>

<sup>1</sup>See also [nLa25b].

01N3

## PROPOSITION 4.6.4.1.2 ► A SIX-FUNCTOR FORMALISM FOR SETS

Let  $X$  be a set.

01N4

1. *The Beck–Chevalley Condition.* Let

$$\begin{array}{ccc} X \times_Z Y & \xrightarrow{\text{pr}_2} & Y \\ \text{pr}_1 \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

be a pullback diagram in Sets. We have

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{\text{pr}_1^{-1}} & \mathcal{P}(X \times_Z Y) \\ f_! \downarrow & & \downarrow (\text{pr}_2)_! \\ \mathcal{P}(Z) & \xrightarrow[g^{-1}]{} & \mathcal{P}(Y), \end{array} \quad \begin{array}{ccc} g^{-1} \circ f_! = (\text{pr}_2)_! \circ \text{pr}_1^{-1}, & & \mathcal{P}(X) \xrightarrow{\text{pr}_2^{-1}} \mathcal{P}(X \times_Z Y) \\ f^{-1} \circ g_! = (\text{pr}_1)_! \circ \text{pr}_2^{-1}, & & g_! \downarrow \quad \downarrow (\text{pr}_1)_! \\ \mathcal{P}(Z) & \xrightarrow[f^{-1}]{} & \mathcal{P}(Y). \end{array}$$

01N5

## 2. The Projection Formula I. The diagram

$$\begin{array}{ccc}
& \mathcal{P}(X) \times \mathcal{P}(X) & \\
\text{id}_{\mathcal{P}(X)} \times f^{-1} \nearrow & & \searrow \cap \\
\mathcal{P}(X) \times \mathcal{P}(Y) & & \mathcal{P}(X) \\
\downarrow f_! \times \text{id}_{\mathcal{P}(Y)} & & \downarrow f_! \\
\mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{\cap} & \mathcal{P}(Y),
\end{array}$$

commutes, i.e. we have

$$f_!(U \cap f^{-1}(V)) = f_!(U) \cap V$$

for each  $U \in \mathcal{P}(X)$  and each  $V \in \mathcal{P}(Y)$ .

01N6

## 3. The Projection Formula II. We have a natural transformation

$$\begin{array}{ccc}
& \mathcal{P}(X) \times \mathcal{P}(X) & \\
\text{id}_{\mathcal{P}(X)} \times f^{-1} \nearrow & & \searrow \cap \\
\mathcal{P}(X) \times \mathcal{P}(Y) & & \mathcal{P}(X) \\
\downarrow f_* \times \text{id}_{\mathcal{P}(Y)} & \curvearrowleft & \downarrow f_* \\
\mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{\cap} & \mathcal{P}(Y),
\end{array}$$

with components

$$f_*(U) \cap V \subset f_*(U \cap f^{-1}(V))$$

indexed by  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ .

01N7

## 4. Strong Closed Monoidality. The diagram

$$\begin{array}{ccc}
\mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{-1,\text{op}} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\
[-1,-2]_Y \downarrow & & \downarrow [-1,-2]_X \\
\mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)
\end{array}$$

commutes, i.e. we have an equality of sets

$$f^{-1}([U, V]_Y) = [f^{-1}(U), f^{-1}(V)]_X,$$

natural in  $U, V \in \mathcal{P}(X)$ .

01N8

5. *The External Tensor Product.* We have an external tensor product

$$-_1 \boxtimes_{X \times Y} -_2 : \mathcal{P}(X) \times \mathcal{P}(Y) \rightarrow \mathcal{P}(X \times Y)$$

given by

$$\begin{aligned} U \boxtimes_{X \times Y} V &\stackrel{\text{def}}{=} \text{pr}_1^{-1}(U) \cap \text{pr}_2^{-1}(V) \\ &= \{(u, v) \in X \times Y \mid u \in U \text{ and } v \in V\}. \end{aligned}$$

This is the same map as the one in [Item 5 of Proposition 4.4.1.1.4](#). Moreover, the following conditions are satisfied:

01N9

(a) *Interaction With Direct Images.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be functions. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(Y) & \xrightarrow{f_! \times g_!} & \mathcal{P}(X') \times \mathcal{P}(Y') \\ \boxtimes_{X \times Y} \downarrow & & \downarrow \boxtimes_{X' \times Y'} \\ \mathcal{P}(X \times Y) & \xrightarrow{f_! \times g_!} & \mathcal{P}(X' \times Y') \end{array}$$

commutes, i.e. we have

$$[f_! \times g_!](U \boxtimes_{X \times Y} V) = f_!(U) \boxtimes_{X' \times Y'} g_!(V)$$

for each  $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$ .

01NA

(b) *Interaction With Inverse Images.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be functions. The diagram

$$\begin{array}{ccc} \mathcal{P}(X') \times \mathcal{P}(Y') & \xrightarrow{f^{-1} \times g^{-1}} & \mathcal{P}(X) \times \mathcal{P}(Y) \\ \boxtimes_{X' \times Y'} \downarrow & & \downarrow \boxtimes_{X \times Y} \\ \mathcal{P}(X' \times Y') & \xrightarrow{f^{-1} \times g^{-1}} & \mathcal{P}(X \times Y) \end{array}$$

commutes, i.e. we have

$$[f^{-1} \times g^{-1}](U \boxtimes_{X' \times Y'} V) = f^{-1}(U) \boxtimes_{X \times Y} g^{-1}(V)$$

for each  $(U, V) \in \mathcal{P}(X') \times \mathcal{P}(Y')$ .

01NB

(c) *Interaction With Codirect Images.* Let  $f: X \rightarrow X'$  and  $g: Y \rightarrow Y'$  be functions. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(Y) & \xrightarrow{f_* \times g_*} & \mathcal{P}(X') \times \mathcal{P}(Y') \\ \boxtimes_{X \times Y} \downarrow & & \downarrow \boxtimes_{X' \times Y'} \\ \mathcal{P}(X \times Y) & \xrightarrow{f_* \times g_*} & \mathcal{P}(X' \times Y') \end{array}$$

commutes, i.e. we have

$$[f_* \times g_*](U \boxtimes_{X \times Y} V) = f_*(U) \boxtimes_{X' \times Y'} g_*(V)$$

for each  $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$ .

01NC

(d) *Interaction With Diagonals.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\boxtimes_{X \times X}} & \mathcal{P}(X \times X) \\ & \searrow \cap & \downarrow \Delta_X^{-1} \\ & & \mathcal{P}(X), \end{array}$$

i.e. we have

$$U \cap V = \Delta_X^{-1}(U \boxtimes_{X \times X} V)$$

for each  $U, V \in \mathcal{P}(X)$ .

01ND

6. *The Dualisation Functor.* We have a functor

$$D_X: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X)$$

given by

$$\begin{aligned} D_X(U) &\stackrel{\text{def}}{=} [U, \emptyset]_X \\ &\stackrel{\text{def}}{=} U^c \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , as in Item 5 of Proposition 4.4.7.1.4, satisfying the following conditions:

01NE

(a) *Duality.* We have

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{D_X} & \mathcal{P}(X) \\ D_X(D_X(U)) = U, & \searrow \text{id}_{\mathcal{P}(X)} & \downarrow D_X \\ & & \mathcal{P}(X). \end{array}$$

01NF

(b) *Duality.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} & \xrightarrow{\cap^{\text{op}}} & \mathcal{P}(X)^{\text{op}} \\
 \text{id}_{\mathcal{P}(X)^{\text{op}}} \times D_X \swarrow & & \searrow D_X \\
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-_1, -_2]_X} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$\underbrace{D_X(U \cap D_X(V))}_{\stackrel{\text{def}}{=} [U \cap [V, \emptyset]_X, \emptyset]_X} = [U, V]_X$$

for each  $U, V \in \mathcal{P}(X)$ .

01NG

(c) *Interaction With Direct Images.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\
 D_X \downarrow & & \downarrow D_Y \\
 \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$f_!(D_X(U)) = D_Y(f_*(U))$$

for each  $U \in \mathcal{P}(X)$ .

01NH

(d) *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1,\text{op}}} & \mathcal{P}(X)^{\text{op}} \\
 D_Y \downarrow & & \downarrow D_X \\
 \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$f^{-1}(D_Y(U)) = D_X(f^{-1}(U))$$

for each  $U \in \mathcal{P}(X)$ .

01NJ

(e) *Interaction With Codirect Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ D_X \downarrow & & \downarrow D_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(D_X(U)) = D_Y(f_!(U))$$

for each  $U \in \mathcal{P}(X)$ .**PROOF 4.6.4.1.3 ► PROOF OF PROPOSITION 4.6.4.1.2****Item 1: The Beck–Chevalley Condition**

We have

$$\begin{aligned} [g^{-1} \circ f_!](U) &\stackrel{\text{def}}{=} g^{-1}(f_!(U)) \\ &\stackrel{\text{def}}{=} \{y \in Y \mid g(y) \in f_!(U)\} \\ &= \left\{ y \in Y \mid \begin{array}{l} \text{there exists some } x \in U \\ \text{such that } f(x) = g(y) \end{array} \right\} \\ &= \left\{ y \in Y \mid \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid x \in U\} \end{array} \right\} \\ &= \left\{ y \in Y \mid \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid x \in U\} \\ \text{such that } y = y \end{array} \right\} \\ &= \left\{ y \in Y \mid \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid x \in U\} \\ \text{such that } \text{pr}_2(x, y) = y \end{array} \right\} \\ &\stackrel{\text{def}}{=} (\text{pr}_2)_!(\{(x, y) \in X \times_Z Y \mid x \in U\}) \\ &= (\text{pr}_2)_!(\{(x, y) \in X \times_Z Y \mid \text{pr}_1(x, y) \in U\}) \\ &\stackrel{\text{def}}{=} (\text{pr}_2)_!(\text{pr}_1^{-1}(U)) \\ &\stackrel{\text{def}}{=} [(\text{pr}_2)_! \circ \text{pr}_1^{-1}](U) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ . Therefore, we have

$$g^{-1} \circ f_! = (\text{pr}_2)_! \circ \text{pr}_1^{-1}.$$

For the second equality, we have

$$\begin{aligned} [f^{-1} \circ g_!](U) &\stackrel{\text{def}}{=} f^{-1}(g_!(U)) \\ &\stackrel{\text{def}}{=} \{x \in X \mid f(x) \in g_!(V)\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } y \in V \\ \text{such that } f(x) = g(y) \end{array} \right\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid y \in V\} \end{array} \right\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid y \in V\} \\ \text{such that } x = x \end{array} \right\} \\ &= \left\{ x \in X \mid \begin{array}{l} \text{there exists some } \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid y \in V\} \\ \text{such that } \text{pr}_1(x, y) = x \end{array} \right\} \\ &\stackrel{\text{def}}{=} (\text{pr}_1)_!(\{(x, y) \in X \times_Z Y \mid y \in V\}) \\ &= (\text{pr}_1)_!(\{(x, y) \in X \times_Z Y \mid \text{pr}_2(x, y) \in V\}) \\ &\stackrel{\text{def}}{=} (\text{pr}_1)_!(\text{pr}_2^{-1}(V)) \\ &\stackrel{\text{def}}{=} [(\text{pr}_1)_! \circ \text{pr}_2^{-1}](V) \end{aligned}$$

for each  $V \in \mathcal{P}(Y)$ . Therefore, we have

$$f^{-1} \circ g_! = (\text{pr}_1)_! \circ \text{pr}_2^{-1}.$$

This finishes the proof.

### Item 2: The Projection Formula I

We claim that

$$f_!(U) \cap V \subset f_!(U \cap f^{-1}(V)).$$

Indeed, we have

$$\begin{aligned} f_!(U) \cap V &\subset f_!(U) \cap f_!(f^{-1}(V)) \\ &= f_!(U \cap f^{-1}(V)), \end{aligned}$$

where we have used:

024B

1. Item 2 of Proposition 4.6.1.1.5 for the inclusion.

024C

2. Item 6 of Proposition 4.6.1.1.5 for the equality.

Conversely, we claim that

$$f_!(U \cap f^{-1}(V)) \subset f_!(U) \cap V.$$

Indeed:

024D

1. Let  $y \in f_!(U \cap f^{-1}(V))$ .

024E

2. Since  $y \in f_!(U \cap f^{-1}(V))$ , there exists some  $x \in U \cap f^{-1}(V)$  such that  $f(x) = y$ .

024F

3. Since  $x \in U \cap f^{-1}(V)$ , we have  $x \in U$ , and thus  $f(x) \in f_!(U)$ .

024G

4. Since  $x \in U \cap f^{-1}(V)$ , we have  $x \in f^{-1}(V)$ , and thus  $f(x) \in V$ .

024H

5. Since  $f(x) \in f_!(U)$  and  $f(x) \in V$ , we have  $f(x) \in f_!(U) \cap V$ .

024J

6. But  $y = f(x)$ , so  $y \in f_!(U) \cap V$ .

024K

7. Thus  $f_!(U \cap f^{-1}(V)) \subset f_!(U) \cap V$ .

This finishes the proof.

**Item 3: The Projection Formula II**

We have

$$\begin{aligned} f_*(U) \cap V &\subset f_*(U) \cap f_*(f^{-1}(V)) \\ &= f_*(U \cap f^{-1}(V)), \end{aligned}$$

where we have used:

024L

1. Item 2 of Proposition 4.6.3.1.7 for the inclusion.

024M

2. Item 6 of Proposition 4.6.3.1.7 for the equality.

Since  $\mathcal{P}(Y)$  is posetal, naturality is automatic (?? of ??).

**Item 4: Strong Closed Monoidality**

This is a repetition of Item 19 of Proposition 4.4.7.1.4 and is proved there.

**Item 5: The External Tensor Product**

We have

$$U \boxtimes_{X \times Y} V \stackrel{\text{def}}{=} \text{pr}_1^{-1}(U) \cap \text{pr}_2^{-1}(V)$$

$$\begin{aligned}
&\stackrel{\text{def}}{=} \{(x, y) \in X \times Y \mid \text{pr}_1(x, y) \in U\} \\
&\quad \cup \{(x, y) \in X \times Y \mid \text{pr}_2(x, y) \in V\} \\
&= \{(x, y) \in X \times Y \mid x \in U\} \\
&\quad \cup \{(x, y) \in X \times Y \mid y \in V\} \\
&= \{(x, y) \in X \times Y \mid x \in U \text{ and } y \in V\} \\
&\stackrel{\text{def}}{=} U \times V.
\end{aligned}$$

Next, we claim that **Items 5a** to **5d** are indeed true:

- 024N 1. *Proof of Item 5a:* This is a repetition of **Item 16** of **Proposition 4.6.1.1.5** and is proved there.
- 024P 2. *Proof of Item 5b:* This is a repetition of **Item 16** of **Proposition 4.6.2.1.3** and is proved there.
- 024Q 3. *Proof of Item 5c:* This is a repetition of **Item 15** of **Proposition 4.6.3.1.7** and is proved there.
- 024R 4. *Proof of Item 5d:* We have

$$\begin{aligned}
\Delta_X^{-1}(U \boxtimes_{X \times X} V) &\stackrel{\text{def}}{=} \{x \in X \mid (x, x) \in U \boxtimes_{X \times X} V\} \\
&= \{x \in X \mid (x, x) \in \{(u, v) \in X \times X \mid u \in U \text{ and } v \in V\}\} \\
&= U \cap V.
\end{aligned}$$

This finishes the proof.

**Item 6: The Dualisation Functor**

This is a repetition of **Items 5** and **6** of **Proposition 4.4.7.1.4** and is proved there. 

## 02RG 4.7 Miscellany

### 02RH 4.7.1 Injective Functions

Let  $A$  and  $B$  be sets.

#### 02RJ DEFINITION 4.7.1.1.1 ► INJECTIVE FUNCTIONS

A function  $f: A \rightarrow B$  is **injective** if it satisfies the following condition:

- (★) For each  $a, a' \in A$ , if  $f(a) = f(a')$ , then  $a = a'$ .

02RK

**PROPOSITION 4.7.1.1.2 ► PROPERTIES OF INJECTIVE FUNCTIONS**

Let  $f: A \rightarrow B$  be a function.

02RL

1. *Characterisations.* The following conditions are equivalent:<sup>1</sup>

02RM

(a) The function  $f$  is injective.

02RN

(b) The function  $f$  is a monomorphism in Sets.

02RP

(c) The direct image function

$$f_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to  $f$  is injective.

02RQ

(d) The codirect image function

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to  $f$  is injective.

02RR

(e) The direct image functor

$$f_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

associated to  $f$  is full.

02RS

(f) The codirect image function

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to  $f$  is full.

02RT

(g) The diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow \chi_A & \downarrow f^{-1} \\ & & \mathcal{P}(A) \end{array}$$

commutes. That is, we have

$$f^{-1}(f(a)) = \{a\}$$

for each  $a \in A$ .

02RU

(h) We have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_!} & \mathcal{P}(B) \\ f^{-1} \circ f_! = \text{id}_{\mathcal{P}(A)} & \swarrow & \downarrow f^{-1} \\ & & \mathcal{P}(A). \end{array}$$

In other words, we have

$$\{a \in A \mid f(a) \in f(U)\} = U$$

for each  $U \in \mathcal{P}(A)$ .

02RV

(i) We have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_*} & \mathcal{P}(B) \\ f^{-1} \circ f_* = \text{id}_{\mathcal{P}(A)} & \searrow & \downarrow f^{-1} \\ & & \mathcal{P}(A). \end{array}$$

In other words, we have

$$\{a \in A \mid f^{-1}(f(a)) \subset U\} = U$$

for each  $U \in \mathcal{P}(A)$ .

<sup>1</sup>Items 1c to 1f unwind respectively to the following statements:

- For each  $U, V \in \mathcal{P}(A)$ , if  $f_!(U) = f_!(V)$ , then  $U = V$ .
- For each  $U, V \in \mathcal{P}(A)$ , if  $f_*(U) = f_*(V)$ , then  $U = V$ .
- For each  $U, V \in \mathcal{P}(A)$ , if  $f_!(U) \subset f_!(V)$ , then  $U \subset V$ .
- For each  $U, V \in \mathcal{P}(A)$ , if  $f_*(U) \subset f_*(V)$ , then  $U \subset V$ .

#### PROOF 4.7.1.1.3 ► PROOF OF PROPOSITION 4.7.1.1.2

##### Item 1: Characterisations

We will proceed by showing:

- Step 1: Item 1a  $\iff$  Item 1b.
- Step 2: Item 1a  $\iff$  Item 1c.
- Step 3: Item 1a  $\iff$  Item 1d.
- Step 4: Item 1c  $\iff$  Item 1e.
- Step 5: Item 1e  $\iff$  Item 1f.
- Step 6: Item 1a  $\iff$  Item 1g.
- Step 7: Item 1g  $\iff$  Item 1h.

- Step 8: Item 1a  $\iff$  Item 1i.

### Step 1: Item 1a $\iff$ Item 1b

We claim that Items 1a and 1b are equivalent:

- *Item 1a  $\implies$  Item 1b*: We proceed in a few steps:
  - Proceeding by contrapositive, we claim that given a pair of maps  $g, h: C \rightrightarrows A$  such that  $g \neq h$ , we have  $f \circ g \neq f \circ h$ .
  - Indeed, as  $g$  and  $h$  are different maps, there must exist at least one element  $x \in C$  such that  $g(x) \neq h(x)$ .
  - But then we have  $f(g(x)) \neq f(h(x))$ , as  $f$  is injective.
  - Thus  $f \circ g \neq f \circ h$ , and we are done.
- *Item 1b  $\implies$  Item 1a*: We proceed in a few steps:
  - Consider the diagram

$$\text{pt} \xrightarrow{\quad [x] \quad} A \xrightarrow{f} B,$$

where  $[x]$  and  $[y]$  are the morphisms picking the elements  $x$  and  $y$  of  $A$ .

- Note that we have  $f(x) = f(y)$  iff  $f \circ [x] = f \circ [y]$ .
- Since  $f$  is assumed to be a monomorphism, if  $f(x) = f(y)$ , then  $f \circ [x] = f \circ [y]$  and therefore  $[x] = [y]$ .
- This shows that if  $f(x) = f(y)$ , then  $x = y$ , so  $f$  is injective.

### Step 2: Item 1a $\iff$ Item 1c

We claim that Items 1a and 1c are indeed equivalent:

- *Item 1a  $\implies$  Item 1c*: We proceed in a few steps:
  - Assume that  $f$  is injective and let  $U, V \in \mathcal{P}(A)$  such that  $f_!(U) = f_!(V)$ . We wish to show that  $U = V$ .
  - To show that  $U \subset V$ , let  $u \in U$ .
    - By the definition of the direct image, we have  $f(u) \in f_!(U)$ .
    - Since  $f_!(U) = f_!(V)$ , it follows that  $f(u) \in f_!(V)$ .
    - Thus, there exists some  $v \in V$  such that  $f(v) = f(u)$ .

- Since  $f$  is injective, the equality  $f(v) = f(u)$  implies that  $v = u$ .
  - Thus  $u \in V$  and  $U \subset V$ .
  - A symmetric argument shows that  $V \subset U$ .
  - Therefore  $U = V$ , showing  $f_!$  to be injective.
- *Item 1c  $\implies$  Item 1a:* We proceed in a few steps:
    - Assume that the direct image function  $f_!$  is injective and let  $a, a' \in A$  such that  $f(a) = f(a')$ . We wish to show that  $a = a'$ .
    - Since

$$\begin{aligned} f_!(\{a\}) &= \{f(a)\} \\ &= \{f(a')\} \\ &= f_!(\{a'\}), \end{aligned}$$

we must have  $\{a\} = \{a'\}$ , as  $f_!$  is injective, so  $a = a'$ , showing  $f$  to be injective.

#### Step 3: Item 1c $\iff$ Item 1d

This follows from [Item 17](#) of [Proposition 4.6.1.1.5](#).

#### Step 4: Item 1c $\iff$ Item 1e

We claim that [Items 1c](#) and [1e](#) are equivalent:

- *Item 1c  $\implies$  Item 1e:* We proceed in a few steps:
  - Let  $U, V \in \mathcal{P}(A)$  such that  $f_!(U) \subset f_!(V)$ , assume  $f_!$  to be injective, and consider the set  $U \cup V$ .
  - Since  $f_!(U) \subset f_!(V)$ , we have

$$\begin{aligned} f_!(U \cup V) &= f_!(U) \cup f_!(V) \\ &= f_!(V), \end{aligned}$$

where we have used [Item 5](#) of [Proposition 4.6.1.1.5](#) for the first equality.

- Since  $f_!$  is injective, this implies  $U \cup V = V$ .
- Thus  $U \subset V$ , as we wished to show.

- *Item 1c*  $\implies$  *Item 1e*: We proceed in a few steps:
  - Suppose *Item 1e* holds, and let  $U, V \in \mathcal{P}(A)$  such that  $f_!(U) = f_!(V)$ .
  - Since  $f_!(U) = f_!(V)$ , we have  $f_!(U) \subset f_!(V)$  and  $f_!(V) \subset f_!(U)$ .
  - By assumption, this implies  $U \subset V$  and  $V \subset U$ .
  - Thus  $U = V$ , showing  $f_!$  to be injective.

### Step 5: Item 1e $\iff$ Item 1f

This follows from *Item 17* of [Proposition 4.6.1.1.5](#).

### Step 6: Item 1a $\iff$ Item 1g

We have

$$f^{-1}(f(a)) = \{a' \in A \mid f(a') = f(a)\}$$

so the condition  $f^{-1}(f(a)) = \{a\}$  states precisely that if  $f(a') = f(a)$ , then  $a' = a$ .

### Step 7: Item 1g $\iff$ Item 1h

We claim that *Items 1g* and *1h* are indeed equivalent:

- *Item 1g*  $\implies$  *Item 1h*: We have

$$\begin{aligned} [f^{-1} \circ f_!](U) &\stackrel{\text{def}}{=} f^{-1}(f_!(U)) \\ &= f^{-1}\left(f_!\left(\bigcup_{u \in U} \{u\}\right)\right) \\ &= f^{-1}\left(\bigcup_{u \in U} f_!(\{u\})\right) \\ &= \bigcup_{u \in U} f^{-1}(f_!(\{u\})) \\ &= \bigcup_{u \in U} f^{-1}(f_!(u)) \\ &= \bigcup_{u \in U} \{u\} \\ &= U \end{aligned}$$

for each  $U \in \mathcal{P}(A)$ , where we have used [Item 5 of Proposition 4.6.1.1.5](#) for the third equality and [Item 5 of Proposition 4.6.2.1.3](#) for the fourth equality.

- [Item 1h](#)  $\implies$  [Item 1g](#): Applying the condition  $f^{-1} \circ f_! = \text{id}_{\mathcal{P}(A)}$  to  $U = \{a\}$  gives

$$f^{-1}(f_!(\{a\})) = \{a\}.$$

### Step 8: Item 1a $\iff$ Item 1i

We claim that [Items 1a](#) and [1i](#) are equivalent:

- [Item 1a](#)  $\implies$  [Item 1i](#): If  $f$  is injective, then  $f^{-1}(f(a)) = \{a\}$ , so we have

$$\begin{aligned} f^{-1}(f_*(a)) &= \{a \in A \mid \{a\} \subset U\} \\ &= U. \end{aligned}$$

- [Item 1i](#)  $\implies$  [Item 1a](#): For  $U = \{a\}$ , the condition  $f^{-1}(f_*(U)) = U$  becomes

$$\{a' \in A \mid f^{-1}(f(a')) \subset \{a\}\} = \{a\}.$$

Since the set  $f^{-1}(f(a'))$  is given by

$$\{a \in A \mid f(a) = f(a')\},$$

it follows that  $f$  is injective.

This finishes the proof. ■

## 02RW 4.7.2 Surjective Functions

Let  $A$  and  $B$  be sets.

02RX

### DEFINITION 4.7.2.1.1 ► SURJECTIVE FUNCTIONS

A function  $f: A \rightarrow B$  is **surjective** if it satisfies the following condition:

- (★) For each  $b \in B$ , there exists some  $a \in A$  such that  $f(a) = b$ .

02RY

### PROPOSITION 4.7.2.1.2 ► PROPERTIES OF SURJECTIVE FUNCTIONS

Let  $f: A \rightarrow B$  be a function.

02RZ

1. *Characterisations.* The following conditions are equivalent:

02S0

(a) The function  $f$  is surjective.

02S1

(b) The function  $f$  is an epimorphism in Sets.

02S2

(c) The inverse image function

$$f^{-1} : \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to  $f$  is injective.

02S3

(d) The inverse image functor

$$f^{-1} : (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

associated to  $f$  is full.

02S4

(e) The diagram

$$\begin{array}{ccc} B & \xrightarrow{f^{-1}} & \mathcal{P}(A) \\ & \searrow \chi_B & \downarrow f_! \\ & & \mathcal{P}(B) \end{array}$$

commutes. That is, we have

$$f_!(f^{-1}(b)) = \{b\}$$

for each  $b \in B$ .

02S5

(f) We have

$$\begin{array}{ccc} \mathcal{P}(B) & \xrightarrow{f^{-1}} & \mathcal{P}(A) \\ f_! \circ f^{-1} = \text{id}_{\mathcal{P}(B)} & \swarrow & \downarrow f_! \\ & & \mathcal{P}(B). \end{array}$$

In other words, we have

$$\left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in f^{-1}(U) \\ \text{such that } f(a) = b \end{array} \right\} = U$$

for each  $U \in \mathcal{P}(A)$ .

02S6

(g) We have

$$\begin{array}{ccc} \mathcal{P}(B) & \xrightarrow{f^{-1}} & \mathcal{P}(A) \\ f_* \circ f^{-1} = \text{id}_{\mathcal{P}(B)} & \searrow & \downarrow f_* \\ & & \mathcal{P}(B). \end{array}$$

In other words, we have

$$\{b \in B \mid f^{-1}(b) \subset f^{-1}(U)\} = U$$

for each  $U \in \mathcal{P}(B)$ .

## PROOF 4.7.2.1.3 ► PROOF OF PROPOSITION 4.7.2.1.2

## Item 1: Characterisations

We will proceed by showing:

- Step 1: Item 1a  $\iff$  Item 1b.
- Step 2: Item 1a  $\iff$  Item 1c.
- Step 3: Item 1c  $\iff$  Item 1d.
- Step 4: Item 1a  $\iff$  Item 1e.
- Step 5: Item 1e  $\iff$  Item 1f.
- Step 6: Item 1a  $\iff$  Item 1g.

Step 1: Item 1a  $\iff$  Item 1b

We claim Items 1a and 1b are indeed equivalent:

- Item 1a  $\implies$  Item 1b: We proceed in a few steps:

- Let  $g, h: B \rightrightarrows C$  be morphisms such that  $g \circ f = h \circ f$ .
- For each  $a \in A$ , we have

$$g(f(a)) = h(f(a)).$$

- However, this implies that

$$g(b) = h(b)$$

for each  $b \in B$ , as  $f$  is surjective.

– Thus  $g = h$  and  $f$  is an epimorphism.

- *Item 1b  $\implies$  Item 1a:* We proceed by contrapositive. Consider the diagram

$$A \xrightarrow{f} B \xrightarrow[h]{g} C,$$

where  $h$  is the map defined by  $h(b) = 0$  for each  $b \in B$  and  $g$  is the map defined by

$$g(b) = \begin{cases} 1 & \text{if } b \in \text{Im}(f), \\ 0 & \text{otherwise.} \end{cases}$$

Then  $h \circ f = g \circ f$ , as  $h(f(a)) = 1 = g(f(a))$  for each  $a \in A$ . However, for any  $b \in B \setminus \text{Im}(f)$ , we have

$$g(b) = 0 \neq 1 = h(b).$$

Therefore  $g \neq h$  and  $f$  is not an epimorphism.

### Step 2: Item 1a $\iff$ Item 1c

We claim **Items 1a** and **1c** are indeed equivalent:

- *Item 1a  $\implies$  Item 1c:* We proceed in a few steps:
  - Assume that  $f$  is surjective. Let  $U, V \in \mathcal{P}(B)$  such that  $f^{-1}(U) = f^{-1}(V)$ . We wish to show that  $U = V$ .
  - To show that  $U \subset V$ , let  $b \in U$ .
  - Since  $f$  is surjective, there must exist some  $a \in A$  such that  $f(a) = b$ .
  - By the definition of the inverse image, since  $f(a) = b$  and  $b \in U$ , we have  $a \in f^{-1}(U)$ .
  - By our initial assumption,  $f^{-1}(U) = f^{-1}(V)$ , so it follows that  $a \in f^{-1}(V)$ .
  - Again, by the definition of the inverse image,  $a \in f^{-1}(V)$  means that  $f(a) \in V$ .
  - Since  $f(a) = b$ , we have shown that  $b \in V$ .
  - This establishes that  $U \subset V$ . A symmetric argument shows that  $V \subset U$ .
  - Thus  $U = V$ , proving that  $f^{-1}$  is injective.

- *Item 1c  $\implies$  Item 1a:* We proceed in a few steps:
  - Assume that the inverse image function  $f^{-1}$  is injective. Suppose, for the sake of contradiction, that  $f$  is not surjective.
  - The assumption that  $f$  is not surjective means there exists some  $b_0 \in B$  such that for all  $a \in A$ , we have  $f(a) \neq b_0$ .
  - By the definition of the inverse image, this is equivalent to stating that  $f^{-1}(\{b_0\}) = \emptyset$ .
  - Since  $f^{-1}(\emptyset) = \emptyset$ , we have  $f^{-1}(\{b_0\}) = f^{-1}(\emptyset)$ .
  - Since  $f^{-1}$  is injective, this implies that  $\{b_0\} = \emptyset$ .
  - This is a contradiction, as the singleton set  $\{b_0\}$  is non-empty.
  - Therefore,  $f$  is surjective.

### Step 3: Item 1c $\iff$ Item 1d

We claim that **Items 1c** and **1d** are equivalent:

- *Item 1c  $\implies$  Item 1d:* We proceed in a few steps:
  - Let  $U, V \in \mathcal{P}(B)$  such that  $f^{-1}(U) \subset f^{-1}(V)$ , assume  $f^{-1}$  to be injective, and consider the set  $U \cup V$ .
  - Since  $f^{-1}(U) \subset f^{-1}(V)$ , we have

$$\begin{aligned} f^{-1}(U \cup V) &= f^{-1}(U) \cup f^{-1}(V) \\ &= f^{-1}(V), \end{aligned}$$

where we have used **Item 5 of Proposition 4.6.2.1.3** for the first equality.

- Since  $f^{-1}$  is injective, this implies  $U \cup V = V$ .
- Thus  $U \subset V$ , as we wished to show.
- *Item 1d  $\implies$  Item 1c:* We proceed in a few steps:

- Suppose **Item 1d** holds, and let  $U, V \in \mathcal{P}(B)$  such that  $f^{-1}(U) = f^{-1}(V)$ .
- Since  $f^{-1}(U) = f^{-1}(V)$ , we have  $f^{-1}(U) \subset f^{-1}(V)$  and  $f^{-1}(V) \subset f^{-1}(U)$ .
- By assumption, this implies  $U \subset V$  and  $V \subset U$ .
- Thus  $U = V$ , showing  $f^{-1}$  to be injective.

### Step 4: Item 1a $\iff$ Item 1e

We have

$$f_!(f^{-1}(b)) = \left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in f^{-1}(b) \\ \text{such that } f(a) = b \end{array} \right\},$$

so the condition  $f_!(f^{-1}(b)) = \{b\}$  holds iff  $f$  is surjective.

### Step 5: Item 1e $\iff$ Item 1f

We claim that **Items 1e** and **1f** are indeed equivalent:

- **Item 1e  $\implies$  Item 1f:** We have

$$\begin{aligned} [f_! \circ f^{-1}](U) &\stackrel{\text{def}}{=} f_!(f^{-1}(U)) \\ &= f_!\left(f^{-1}\left(\bigcup_{u \in U} \{u\}\right)\right) \\ &= f_!\left(\bigcup_{u \in U} f^{-1}(\{u\})\right) \\ &= \bigcup_{u \in U} f_!(f^{-1}(\{u\})) \\ &= \bigcup_{u \in U} f_!(f^{-1}(u)) \\ &= \bigcup_{u \in U} \{u\} \\ &= U \end{aligned}$$

for each  $U \in \mathcal{P}(B)$ , where we have used **Item 5** of **Proposition 4.6.1.1.5** for the third equality and **Item 5** of **Proposition 4.6.2.1.3** for the fourth equality.

- **Item 1f  $\implies$  Item 1e:** Applying the condition  $f_! \circ f^{-1} = \text{id}_{\mathcal{P}(B)}$  to  $U = \{b\}$  gives

$$f_!(f^{-1}(\{b\})) = \{b\}.$$

### Step 6: Item 1a $\iff$ Item 1g

First, note that for the condition  $f^{-1}(b) \subset f^{-1}(U)$  to hold, we must have  $b \in U$  or  $f^{-1}(b) = \emptyset$ . Thus

$$f_*(f^{-1}(U)) = (U \cap \text{Im}(f)) \cup (B \setminus \text{Im}(f)).$$

We now claim that **Items 1a** and **1g** are indeed equivalent:

- **Item 1a**  $\implies$  **Item 1g**: If  $f$  is surjective, we have

$$\begin{aligned} (U \cap \text{Im}(f)) \cup (B \setminus \text{Im}(f)) &= U \cup \emptyset \\ &= U, \end{aligned}$$

so  $f_* \circ f^{-1} = \text{id}_{\mathcal{P}(B)}$ .

- **Item 1g**  $\implies$  **Item 1a**: Taking  $U = \emptyset$  gives

$$\begin{aligned} f_*(f^{-1}(\emptyset)) &= (\emptyset \cap \text{Im}(f)) \cup (B \setminus \text{Im}(f)) \\ &= B \setminus \text{Im}(f), \end{aligned}$$

so the condition  $f_*(f^{-1}(\emptyset)) = \emptyset$  implies  $B \setminus \text{Im}(f) = \emptyset$ . Thus  $\text{Im}(f) = B$  and  $f$  is surjective.

This finishes the proof. 

# Appendices

## 4.A Other Chapters

### Preliminaries

1. Introduction
2. A Guide to the Literature

### Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

### Relations

8. Relations
9. Constructions With Relations
10. Conditions on Relations

### Categories

11. Categories
12. Presheaves and the Yoneda Lemma

### Monoidal Categories

13. Constructions With Monoidal Categories

**Bicategories**

14. Types of Morphisms in Bicategories

**Extra Part**

15. Notes

## Chapter 5

# Monoidal Structures on the Category of Sets

**02H3** This chapter contains some material on monoidal structures on Sets.

### Contents

---

5.1	The Monoidal Category of Sets and Products .....	266
5.1.1	Products of Sets .....	266
5.1.2	The Internal Hom of Sets .....	266
5.1.3	The Monoidal Unit .....	266
5.1.4	The Associator .....	266
5.1.5	The Left Unitor .....	269
5.1.6	The Right Unitor .....	271
5.1.7	The Symmetry .....	273
5.1.8	The Diagonal .....	275
5.1.9	The Monoidal Category of Sets and Products .....	278
5.1.10	The Universal Property of (Sets, $\times$ , pt) .....	283
5.2	The Monoidal Category of Sets and Coproducts.....	300
5.2.1	Coproducts of Sets .....	300
5.2.2	The Monoidal Unit .....	300
5.2.3	The Associator .....	300
5.2.4	The Left Unitor .....	303
5.2.5	The Right Unitor .....	306
5.2.6	The Symmetry .....	308
5.2.7	The Monoidal Category of Sets and Coproducts ..	311
5.3	The Bimonoidal Category of Sets, Products, and Co- products .....	318
5.3.1	The Left Distributor .....	318
5.3.2	The Right Distributor .....	320
5.3.3	The Left Annihilator .....	323

5.3.4	The Right Annihilator.....	324
5.3.5	The Bimonoidal Category of Sets, Products, and Coproducts.....	326
5.A	Other Chapters.....	328

---

## 01NL 5.1 The Monoidal Category of Sets and Products

### 01NM 5.1.1 Products of Sets

See [Section 4.1.3](#).

### 01NN 5.1.2 The Internal Hom of Sets

See [Section 4.3.5](#).

### 01NP 5.1.3 The Monoidal Unit

#### 01NQ DEFINITION 5.1.3.1.1 ► THE MONOIDAL UNIT OF $\times$

The **monoidal unit of the product of sets** is the functor

$$\mathbb{1}^{\text{Sets}}: \text{pt} \rightarrow \text{Sets}$$

defined by

$$\mathbb{1}_{\text{Sets}} \stackrel{\text{def}}{=} \text{pt},$$

where  $\text{pt}$  is the terminal set of [Definition 4.1.1.1.1](#).

### 01NR 5.1.4 The Associator

#### 01NS DEFINITION 5.1.4.1.1 ► THE ASSOCIATOR OF $\times$

The **associator of the product of sets** is the natural isomorphism

$$\alpha^{\text{Sets}}: \times \circ (\times \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Sets}} \times \times) \circ \alpha_{\text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2},$$

as in the diagram

$$\begin{array}{ccc}
 & \text{Sets} \times (\text{Sets} \times \text{Sets}) & \\
 \alpha_{\text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} \nearrow & \swarrow id \times \times & \\
 (\text{Sets} \times \text{Sets}) \times \text{Sets} & & \text{Sets} \times \text{Sets} \\
 \times \times id \searrow & \swarrow \alpha^{\text{Sets}} & \downarrow \times \\
 \text{Sets} \times \text{Sets} & \xrightarrow{x} & \text{Sets},
 \end{array}$$

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}} : (X \times Y) \times Z \xrightarrow{\sim} X \times (Y \times Z)$$

at  $(X, Y, Z)$  is given by

$$\alpha_{X,Y,Z}^{\text{Sets}}((x, y), z) \stackrel{\text{def}}{=} (x, (y, z))$$

for each  $((x, y), z) \in (X \times Y) \times Z$ .

#### PROOF 5.1.4.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.1.4.1.1

##### Invertibility

The inverse of  $\alpha_{X,Y,Z}^{\text{Sets}}$  is the morphism

$$\alpha_{X,Y,Z}^{\text{Sets}, -1} : X \times (Y \times Z) \xrightarrow{\sim} (X \times Y) \times Z$$

defined by

$$\alpha_{X,Y,Z}^{\text{Sets}, -1}(x, (y, z)) \stackrel{\text{def}}{=} ((x, y), z)$$

for each  $(x, (y, z)) \in X \times (Y \times Z)$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned}
 [\alpha_{X,Y,Z}^{\text{Sets}, -1} \circ \alpha_{X,Y,Z}^{\text{Sets}}]((x, y), z) &\stackrel{\text{def}}{=} \alpha_{X,Y,Z}^{\text{Sets}, -1}(\alpha_{X,Y,Z}^{\text{Sets}}((x, y), z)) \\
 &\stackrel{\text{def}}{=} \alpha_{X,Y,Z}^{\text{Sets}, -1}(x, (y, z)) \\
 &\stackrel{\text{def}}{=} ((x, y), z) \\
 &\stackrel{\text{def}}{=} [\text{id}_{(X \times Y) \times Z}]((x, y), z)
 \end{aligned}$$

for each  $((x, y), z) \in (X \times Y) \times Z$ , and therefore we have

$$\alpha_{X,Y,Z}^{\text{Sets}, -1} \circ \alpha_{X,Y,Z}^{\text{Sets}} = \text{id}_{(X \times Y) \times Z}.$$

- *Invertibility II.* We have

$$\begin{aligned}
 [\alpha_{X,Y,Z}^{\text{Sets}} \circ \alpha_{X,Y,Z}^{\text{Sets},-1}](x, (y, z)) &\stackrel{\text{def}}{=} \alpha_{X,Y,Z}^{\text{Sets}}(\alpha_{X,Y,Z}^{\text{Sets},-1}(x, (y, z))) \\
 &\stackrel{\text{def}}{=} \alpha_{X,Y,Z}^{\text{Sets}}((x, y), z) \\
 &\stackrel{\text{def}}{=} (x, (y, z)) \\
 &\stackrel{\text{def}}{=} [\text{id}_{(X \times Y) \times Z}](x, (y, z))
 \end{aligned}$$

for each  $(x, (y, z)) \in X \times (Y \times Z)$ , and therefore we have

$$\alpha_{X,Y,Z}^{\text{Sets},-1} \circ \alpha_{X,Y,Z}^{\text{Sets}} = \text{id}_{X \times (Y \times Z)}.$$

Therefore  $\alpha_{X,Y,Z}^{\text{Sets}}$  is indeed an isomorphism.

### Naturality

We need to show that, given functions

$$\begin{aligned}
 f: X &\rightarrow X', \\
 g: Y &\rightarrow Y', \\
 h: Z &\rightarrow Z'
 \end{aligned}$$

the diagram

$$\begin{array}{ccc}
 (X \times Y) \times Z & \xrightarrow{(f \times g) \times h} & (X' \times Y') \times Z' \\
 \alpha_{X,Y,Z}^{\text{Sets}} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}} \\
 X \times (Y \times Z) & \xrightarrow[f \times (g \times h)]{} & X' \times (Y' \times Z')
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 ((x, y), z) & & ((x, y), z) \mapsto ((f(x), g(y)), h(z)) \\
 \downarrow & & \downarrow \\
 (x, (y, z)) \mapsto (f(x), (g(y), h(z))) & & (f(x), (g(y), h(z)))
 \end{array}$$

and hence indeed commutes, showing  $\alpha^{\text{Sets}}$  to be a natural transformation.

### Being a Natural Isomorphism

Since  $\alpha^{\text{Sets}}$  is natural and  $\alpha^{\text{Sets},-1}$  is a componentwise inverse to  $\alpha^{\text{Sets}}$ , it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\alpha^{\text{Sets},-1}$  is also natural. Thus  $\alpha^{\text{Sets}}$  is a natural isomorphism. 

**01NT 5.1.5 The Left Unit**

**01NU**

**DEFINITION 5.1.5.1.1 ► THE LEFT UNIT OF  $\times$**

The **left unit of the product of sets** is the natural isomorphism

$$\begin{array}{ccc}
 \text{pt} \times \text{Sets} & \xrightarrow{\mathbb{1}^{\text{Sets}} \times \text{id}} & \text{Sets} \times \text{Sets} \\
 \lambda^{\text{Sets}} : \times \circ (\mathbb{1}^{\text{Sets}} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \lambda^{\text{Cats}_2}_{\text{Sets}} & \swarrow \quad \searrow & \downarrow \times \\
 & \lambda^{\text{Cats}_2}_{\text{Sets}} & \text{Sets}, \\
 & & \end{array}$$

whose component

$$\lambda_X^{\text{Sets}} : \text{pt} \times X \xrightarrow{\sim} X$$

at  $X \in \text{Obj}(\text{Sets})$  is given by

$$\lambda_X^{\text{Sets}}(\star, x) \stackrel{\text{def}}{=} x$$

for each  $(\star, x) \in \text{pt} \times X$ .

**PROOF 5.1.5.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.1.5.1**

**Invertibility**

The inverse of  $\lambda_X^{\text{Sets}}$  is the morphism

$$\lambda_X^{\text{Sets}, -1} : X \xrightarrow{\sim} \text{pt} \times X$$

defined by

$$\lambda_X^{\text{Sets}, -1}(x) \stackrel{\text{def}}{=} (\star, x)$$

for each  $x \in X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned}
 [\lambda_X^{\text{Sets}, -1} \circ \lambda_X^{\text{Sets}}](\text{pt}, x) &= \lambda_X^{\text{Sets}, -1}(\lambda_X^{\text{Sets}}(\text{pt}, x)) \\
 &= \lambda_X^{\text{Sets}, -1}(x) \\
 &= (\text{pt}, x) \\
 &= [\text{id}_{\text{pt} \times X}](\text{pt}, x)
 \end{aligned}$$

for each  $(\text{pt}, x) \in \text{pt} \times X$ , and therefore we have

$$\lambda_X^{\text{Sets}, -1} \circ \lambda_X^{\text{Sets}} = \text{id}_{\text{pt} \times X}.$$

- *Invertibility II.* We have

$$\begin{aligned} [\lambda_X^{\text{Sets}} \circ \lambda_X^{\text{Sets}, -1}](x) &= \lambda_X^{\text{Sets}}(\lambda_X^{\text{Sets}, -1}(x)) \\ &= \lambda_X^{\text{Sets}, -1}(\text{pt}, x) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

for each  $x \in X$ , and therefore we have

$$\lambda_X^{\text{Sets}} \circ \lambda_X^{\text{Sets}, -1} = \text{id}_X.$$

Therefore  $\lambda_X^{\text{Sets}}$  is indeed an isomorphism.

### Naturality

We need to show that, given a function  $f: X \rightarrow Y$ , the diagram

$$\begin{array}{ccc} \text{pt} \times X & \xrightarrow{\text{id}_{\text{pt}} \times f} & \text{pt} \times Y \\ \lambda_X^{\text{Sets}} \downarrow & & \downarrow \lambda_Y^{\text{Sets}} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (\star, x) & & (\star, x) \mapsto (\star, f(x)) \\ \downarrow & & \downarrow \\ x & \longmapsto & f(x) \end{array}$$

and hence indeed commutes. Therefore  $\lambda^{\text{Sets}}$  is a natural transformation.

### Being a Natural Isomorphism

Since  $\lambda^{\text{Sets}}$  is natural and  $\lambda^{\text{Sets}, -1}$  is a componentwise inverse to  $\lambda^{\text{Sets}}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\lambda^{\text{Sets}, -1}$  is also natural. Thus  $\lambda^{\text{Sets}}$  is a natural isomorphism. □

**01NV 5.1.6 The Right Unitor**

01NW

DEFINITION 5.1.6.1.1 ► THE RIGHT UNITOR OF  $\times$

The **right unitor of the product of sets** is the natural isomorphism

$$\begin{array}{ccc}
 \text{Sets} \times \text{pt} & \xrightarrow{\text{id} \times \mathbb{1}_{\text{Sets}}} & \text{Sets} \times \text{Sets} \\
 \rho^{\text{Sets}} : \times \circ (\text{id} \times \mathbb{1}_{\text{Sets}}) \xrightarrow{\sim} \rho_{\text{Sets}}^{\text{Cats}_2}, & \swarrow \quad \searrow & \downarrow \times \\
 \rho_{\text{Sets}}^{\text{Cats}_2} & & \text{Sets},
 \end{array}$$

whose component

$$\rho_X^{\text{Sets}} : X \times \text{pt} \xrightarrow{\sim} X$$

at  $X \in \text{Obj}(\text{Sets})$  is given by

$$\rho_X^{\text{Sets}}(x, \star) \stackrel{\text{def}}{=} x$$

for each  $(x, \star) \in X \times \text{pt}$ .

PROOF 5.1.6.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.1.6.1.1

Invertibility

The inverse of  $\rho_X^{\text{Sets}}$  is the morphism

$$\rho_X^{\text{Sets}, -1} : X \xrightarrow{\sim} X \times \text{pt}$$

defined by

$$\rho_X^{\text{Sets}, -1}(x) \stackrel{\text{def}}{=} (x, \star)$$

for each  $x \in X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned}
 [\rho_X^{\text{Sets}, -1} \circ \rho_X^{\text{Sets}}](x, \star) &= \rho_X^{\text{Sets}, -1}(\rho_X^{\text{Sets}}(x, \star)) \\
 &= \rho_X^{\text{Sets}, -1}(x) \\
 &= (x, \star) \\
 &= [\text{id}_{X \times \text{pt}}](x, \star)
 \end{aligned}$$

for each  $(x, \star) \in X \times \text{pt}$ , and therefore we have

$$\rho_X^{\text{Sets}, -1} \circ \rho_X^{\text{Sets}} = \text{id}_{X \times \text{pt}}.$$

- *Invertibility II.* We have

$$\begin{aligned} [\rho_X^{\text{Sets}} \circ \rho_X^{\text{Sets}, -1}](x) &= \rho_X^{\text{Sets}}(\rho_X^{\text{Sets}, -1}(x)) \\ &= \rho_X^{\text{Sets}, -1}(x, \star) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

for each  $x \in X$ , and therefore we have

$$\rho_X^{\text{Sets}} \circ \rho_X^{\text{Sets}, -1} = \text{id}_X.$$

Therefore  $\rho_X^{\text{Sets}}$  is indeed an isomorphism.

### Naturality

We need to show that, given a function  $f: X \rightarrow Y$ , the diagram

$$\begin{array}{ccc} X \times \text{pt} & \xrightarrow{f \times \text{id}_{\text{pt}}} & Y \times \text{pt} \\ \rho_X^{\text{Sets}} \downarrow & & \downarrow \rho_Y^{\text{Sets}} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x, \star) & & (x, \star) \mapsto (f(x), \star) \\ \downarrow & & \downarrow \\ x & \longmapsto & f(x) \end{array}$$

and hence indeed commutes. Therefore  $\rho^{\text{Sets}}$  is a natural transformation.

### Being a Natural Isomorphism

Since  $\rho^{\text{Sets}}$  is natural and  $\rho^{\text{Sets}, -1}$  is a componentwise inverse to  $\rho^{\text{Sets}}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\rho^{\text{Sets}, -1}$  is also natural. Thus  $\rho^{\text{Sets}}$  is a natural isomorphism. 

**01NX 5.1.7 The Symmetry**

**01NY**

**DEFINITION 5.1.7.1.1 ► THE SYMMETRY OF  $\times$**

The **symmetry of the product of sets** is the natural isomorphism

$$\sigma^{\text{Sets}} : \times \xrightarrow{\sim} \times \circ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2}$$

$$\begin{array}{ccc} \text{Sets} \times \text{Sets} & \xrightarrow{\times} & \text{Sets}, \\ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2} \curvearrowleft & \downarrow \sigma^{\text{Sets}} & \curvearrowright \times \\ \text{Sets} \times \text{Sets} & & \end{array}$$

whose component

$$\sigma_{X,Y}^{\text{Sets}} : X \times Y \xrightarrow{\sim} Y \times X$$

at  $X, Y \in \text{Obj}(\text{Sets})$  is defined by

$$\sigma_{X,Y}^{\text{Sets}}(x, y) \stackrel{\text{def}}{=} (y, x)$$

for each  $(x, y) \in X \times Y$ .

**PROOF 5.1.7.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.1.7.1.1**

**Invertibility**

The inverse of  $\sigma_{X,Y}^{\text{Sets}}$  is the morphism

$$\sigma_{X,Y}^{\text{Sets}, -1} : Y \times X \xrightarrow{\sim} X \times Y$$

defined by

$$\sigma_{X,Y}^{\text{Sets}, -1}(y, x) \stackrel{\text{def}}{=} (x, y)$$

for each  $(y, x) \in Y \times X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned} [\sigma_{X,Y}^{\text{Sets}, -1} \circ \sigma_{X,Y}^{\text{Sets}}](x, y) &\stackrel{\text{def}}{=} \sigma_{X,Y}^{\text{Sets}, -1}(\sigma_{X,Y}^{\text{Sets}}(x, y)) \\ &\stackrel{\text{def}}{=} \sigma_{X,Y}^{\text{Sets}, -1}(y, x) \\ &\stackrel{\text{def}}{=} (x, y) \\ &\stackrel{\text{def}}{=} [\text{id}_{X \times Y}](x, y) \end{aligned}$$

for each  $(x, y) \in X \times Y$ , and therefore we have

$$\sigma_{X,Y}^{\text{Sets}, -1} \circ \sigma_{X,Y}^{\text{Sets}} = \text{id}_{X \times Y}.$$

- *Invertibility II.* We have

$$\begin{aligned} [\sigma_{X,Y}^{\text{Sets}} \circ \sigma_{X,Y}^{\text{Sets},-1}](y, x) &\stackrel{\text{def}}{=} \sigma_{X,Y}^{\text{Sets},-1}(\sigma_{X,Y}^{\text{Sets}}(y, x)) \\ &\stackrel{\text{def}}{=} \sigma_{X,Y}^{\text{Sets},-1}(x, y) \\ &\stackrel{\text{def}}{=} (y, x) \\ &\stackrel{\text{def}}{=} [\text{id}_{Y \times X}](y, x) \end{aligned}$$

for each  $(y, x) \in Y \times X$ , and therefore we have

$$\sigma_{X,Y}^{\text{Sets}} \circ \sigma_{X,Y}^{\text{Sets},-1} = \text{id}_{Y \times X}.$$

Therefore  $\sigma_{X,Y}^{\text{Sets}}$  is indeed an isomorphism.

### Naturality

We need to show that, given functions

$$\begin{aligned} f: X &\rightarrow A, \\ g: Y &\rightarrow B \end{aligned}$$

the diagram

$$\begin{array}{ccc} X \times Y & \xrightarrow{f \times g} & A \times B \\ \sigma_{X,Y}^{\text{Sets}} \downarrow & & \downarrow \sigma_{A,B}^{\text{Sets}} \\ Y \times X & \xrightarrow{g \times f} & B \times A \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x, y) & & (x, y) \mapsto (f(x), g(y)) \\ \downarrow & & \downarrow \\ (y, x) \mapsto (g(y), f(x)) & & (g(y), f(x)) \end{array}$$

and hence indeed commutes, showing  $\sigma^{\text{Sets}}$  to be a natural transformation.

### Being a Natural Isomorphism

Since  $\sigma^{\text{Sets}}$  is natural and  $\sigma^{\text{Sets},-1}$  is a componentwise inverse to  $\sigma^{\text{Sets}}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\sigma^{\text{Sets},-1}$  is also natural. Thus  $\sigma^{\text{Sets}}$  is a natural isomorphism. 

**01NZ 5.1.8 The Diagonal**

01P0

**DEFINITION 5.1.8.1.1 ► THE DIAGONAL OF  $\times$**

The **diagonal of the product of sets** is the natural transformation

$$\Delta: \text{id}_{\text{Sets}} \Rightarrow \times \circ \Delta_{\text{Sets}}^{\text{Cats}_2}, \quad \begin{array}{ccc} \text{Sets} & \xrightarrow{\text{id}_{\text{Sets}}} & \text{Sets} \\ \Delta_{\text{Sets}}^{\text{Cats}_2} \searrow & \Downarrow \Delta & \swarrow \times \\ & \text{Sets} \times \text{Sets}, & \end{array}$$

whose component

$$\Delta_X: X \rightarrow X \times X$$

at  $X \in \text{Obj}(\text{Sets})$  is given by

$$\Delta_X(x) \stackrel{\text{def}}{=} (x, x)$$

for each  $x \in X$ .

**PROOF 5.1.8.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.1.8.1.1**

We need to show that, given a function  $f: X \rightarrow Y$ , the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X \downarrow & & \downarrow \Delta_Y \\ X \times X & \xrightarrow{f \times f} & Y \times Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ (x, x) & \longmapsto & (f(x), f(x)) \end{array}$$

and hence indeed commutes, showing  $\Delta$  to be natural.



01P1

**PROPOSITION 5.1.8.1.3 ► PROPERTIES OF THE DIAGONAL MAP**

Let  $X$  be a set.

01P2

1. *Monoidality.* The diagonal map

$$\Delta: \text{id}_{\text{Sets}} \Longrightarrow \times \circ \Delta_{\text{Sets}}^{\text{Cats}_2},$$

is a monoidal natural transformation:

01P3

- (a) *Compatibility With Strong Monoidality Constraints.* For each  $X, Y \in \text{Obj}(\text{Sets})$ , the diagram

$$\begin{array}{ccc} X \times Y & \xrightarrow{\Delta_X \times \Delta_Y} & (X \times X) \times (Y \times Y) \\ & \searrow \Delta_{X \times Y} & \downarrow \\ & & (X \times Y) \times (X \times Y) \end{array}$$

commutes.

01P4

- (b) *Compatibility With Strong Unitality Constraints.* The diagrams

$$\begin{array}{ccc} \text{pt} & \xrightarrow{\Delta_{\text{pt}}} & \text{pt} \times \text{pt} \\ & \swarrow \quad \searrow & \downarrow \lambda_{\text{pt}}^{\text{Sets}} \\ & & \text{pt} \end{array} \quad \begin{array}{ccc} \text{pt} & \xrightarrow{\Delta_{\text{pt}}} & \text{pt} \times \text{pt} \\ & \swarrow \quad \searrow & \downarrow \rho_{\text{pt}}^{\text{Sets}} \\ & & \text{pt} \end{array}$$

commute, i.e. we have

$$\begin{aligned} \Delta_{\text{pt}} &= \lambda_{\text{pt}}^{\text{Sets}, -1} \\ &= \rho_{\text{pt}}^{\text{Sets}, -1}, \end{aligned}$$

where we recall that the equalities

$$\begin{aligned} \lambda_{\text{pt}}^{\text{Sets}} &= \rho_{\text{pt}}^{\text{Sets}}, \\ \lambda_{\text{pt}}^{\text{Sets}, -1} &= \rho_{\text{pt}}^{\text{Sets}, -1} \end{aligned}$$

are always true in any monoidal category by ?? of ??.

01P5

2. *The Diagonal of the Unit.* The component

$$\Delta_{\text{pt}}: \text{pt} \xrightarrow{\sim} \text{pt} \times \text{pt}$$

of  $\Delta$  at pt is an isomorphism.

## PROOF 5.1.8.1.4 ► PROOF OF PROPOSITION 5.1.8.1.3

## Item 1: Monoidality

We claim that  $\Delta$  is indeed monoidal:

024S

- Item 1a: Compatibility With Strong Monoidality Constraints:** We need to show that the diagram

$$\begin{array}{ccc} X \times Y & \xrightarrow{\Delta_X \times \Delta_Y} & (X \times X) \times (Y \times Y) \\ & \searrow \Delta_{X \times Y} & \downarrow \\ & & (X \times Y) \times (X \times Y) \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x, y) & \longmapsto & ((x, x), (y, y)) \\ & \downarrow & \swarrow \\ & ((x, y), (x, y)) & ((x, y), (x, y)) \end{array}$$

and hence indeed commutes.

024T

- Item 1b: Compatibility With Strong Unitality Constraints:** As shown in the proof of [Definition 5.1.5.1.1](#), the inverse of the left unit of Sets with respect to the product at  $X \in \text{Obj}(\text{Sets})$  is given by

$$\lambda_X^{\text{Sets}, -1}(x) \stackrel{\text{def}}{=} (\star, x)$$

for each  $x \in X$ , so when  $X = \text{pt}$ , we have

$$\lambda_{\text{pt}}^{\text{Sets}, -1}(\star) \stackrel{\text{def}}{=} (\star, \star),$$

and also

$$\Delta_{\text{pt}}^{\text{Sets}}(\star) \stackrel{\text{def}}{=} (\star, \star),$$

so we have  $\Delta_{\text{pt}} = \lambda_{\text{pt}}^{\text{Sets}, -1}$ .

This finishes the proof.

## Item 2: The Diagonal of the Unit

This follows from [Item 1](#) and the invertibility of the left/right unit of Sets with respect to  $\times$ , proved in the proof of [Definition 5.1.5.1.1](#) for the left unit or the proof of [Definition 5.1.6.1.1](#) for the right unitor. ☐

**01P6 5.1.9 The Monoidal Category of Sets and Products**

01P7

**PROPOSITION 5.1.9.1.1 ► THE MONOIDAL STRUCTURE ON SETS ASSOCIATED TO THE PRODUCT**

The category Sets admits a closed symmetric monoidal category with diagonals structure consisting of:

- *The Underlying Category.* The category Sets of pointed sets.
- *The Monoidal Product.* The product functor

$$\times: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}$$

of Item 1 of Proposition 4.1.3.1.4.

- *The Internal Hom.* The internal Hom functor

$$\text{Sets}: \text{Sets}^{\text{op}} \times \text{Sets} \rightarrow \text{Sets}$$

of Item 1 of Proposition 4.3.5.1.2.

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\text{Sets}}: \text{pt} \rightarrow \text{Sets}$$

of Definition 5.1.3.1.1.

- *The Associators.* The natural isomorphism

$$\alpha^{\text{Sets}}: \times \circ (\times \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Sets}} \times \times) \circ \alpha_{\text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}}$$

of Definition 5.1.4.1.1.

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\text{Sets}}: \times \circ (\mathbb{1}^{\text{Sets}} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \lambda_{\text{Sets}}^{\text{Cats}_2}$$

of Definition 5.1.5.1.1.

- *The Right Unitors.* The natural isomorphism

$$\rho^{\text{Sets}}: \times \circ (\text{id} \times \mathbb{1}^{\text{Sets}}) \xrightarrow{\sim} \rho_{\text{Sets}}^{\text{Cats}_2}$$

of Definition 5.1.6.1.1.

- *The Symmetry.* The natural isomorphism

$$\sigma^{\text{Sets}}: \times \xrightarrow{\sim} \times \circ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2}$$

of Definition 5.1.7.1.1.

- *The Diagonals.* The monoidal natural transformation

$$\Delta: \text{id}_{\text{Sets}} \Longrightarrow \times \circ \Delta_{\text{Sets}}^{\text{Cats}_2}$$

of Definition 5.1.8.1.1.

## PROOF 5.1.9.1.2 ► PROOF OF PROPOSITION 5.1.9.1.1

## The Pentagon Identity

Let  $W, X, Y$  and  $Z$  be sets. We have to show that the diagram

$$\begin{array}{ccccc}
 & & (W \times (X \times Y)) \times Z & & \\
 & \swarrow \alpha_{W,X,Y}^{\text{Sets}} \times \text{id}_Z & & \searrow \alpha_{W,X \times Y,Z}^{\text{Sets}} & \\
 ((W \times X) \times Y) \times Z & & & & W \times ((X \times Y) \times Z) \\
 & \swarrow \alpha_{W \times X,Y,Z}^{\text{Sets}} & & & \searrow \text{id}_W \times \alpha_{X,Y,Z}^{\text{Sets}} \\
 & (W \times X) \times (Y \times Z) & \xrightarrow{\alpha_{W,X,Y \times Z}^{\text{Sets}}} & & W \times (X \times (Y \times Z))
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 & ((w, (x, y)), z) & \\
 & \nearrow & \searrow \\
 (((w, x), y), z) & & (((w, x), y), z) \\
 & \nearrow & \searrow \\
 & ((w, x), (y, z)) \mapsto (w, (x, (y, z))) & \\
 & \nearrow & \searrow \\
 & (w, (x, (y, z))), &
 \end{array}$$

and thus the pentagon identity is satisfied.

## The Triangle Identity

Let  $X$  and  $Y$  be sets. We have to show that the diagram

$$\begin{array}{ccc}
 (X \times \text{pt}) \times Y & \xrightarrow{\alpha_{X,\text{pt},Y}^{\text{Sets}}} & X \times (\text{pt} \times Y) \\
 & \searrow \rho_X^{\text{Sets}} \times \text{id}_Y & \swarrow \text{id}_X \times \lambda_Y^{\text{Sets}} \\
 & X \times Y &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} ((x, \star), y) & & ((x, \star), y) \mapsto (x, (\star, y)) \\ \swarrow & & \searrow \\ (x, y) & & (x, y) \end{array}$$

and thus the triangle identity is satisfied.

### The Left Hexagon Identity

Let  $X$ ,  $Y$ , and  $Z$  be sets. We have to show that the diagram

$$\begin{array}{ccccc} & & (X \times Y) \times Z & & \\ & \alpha_{X,Y,Z}^{\text{Sets}} \swarrow & & \searrow \sigma_{X,Y}^{\text{Sets}} \times \text{id}_Z & \\ X \times (Y \times Z) & & & & (Y \times X) \times Z \\ \downarrow \sigma_{X,Y \times Z}^{\text{Sets}} & & & & \downarrow \alpha_{Y,X,Z}^{\text{Sets}} \\ (Y \times Z) \times X & & & & Y \times (X \times Z) \\ \downarrow \alpha_{Y,Z,X}^{\text{Sets}} & \searrow & & \swarrow \text{id}_Y \times \sigma_{X,Z}^{\text{Sets}} & \\ Y \times (Z \times X) & & & & \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} ((x, y), z) & & ((x, y), z) \\ \swarrow & & \searrow \\ (x, (y, z)) & & ((y, x), z) \\ \downarrow & & \downarrow \\ ((y, z), x) & & (y, (x, z)) \\ \swarrow & & \searrow \\ (y, (z, x)) & & (y, (z, x)) \end{array}$$

and thus the left hexagon identity is satisfied.

### The Right Hexagon Identity

Let  $X$ ,  $Y$ , and  $Z$  be sets. We have to show that the diagram

$$\begin{array}{ccc}
 & X \times (Y \times Z) & \\
 (\alpha_{X,Y,Z}^{\text{Sets}})^{-1} \swarrow & & \searrow \text{id}_X \times \sigma_{Y,Z}^{\text{Sets}} \\
 (X \times Y) \times Z & & X \times (Z \times Y) \\
 \downarrow \sigma_{X \times Y, Z}^{\text{Sets}} & & \downarrow (\alpha_{X,Z,Y}^{\text{Sets}})^{-1} \\
 Z \times (X \times Y) & & (X \times Z) \times Y \\
 \downarrow (\alpha_{Z,X,Y}^{\text{Sets}})^{-1} & & \downarrow \sigma_{X,Z}^{\text{Sets}} \times \text{id}_Y \\
 (Z \times X) \times Y & &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (x, (y, z)) & & (x, (y, z)) \\
 \nearrow & & \swarrow \\
 ((x, y), z) & & (x, (z, y)) \\
 \downarrow & & \downarrow \\
 (z, (x, y)) & & ((x, z), y) \\
 \nearrow & & \swarrow \\
 ((z, x), y) & & ((z, x), y)
 \end{array}$$

and thus the right hexagon identity is satisfied.

Monoidal Closedness

This follows from [Item 2 of Proposition 4.3.5.1.2](#)

Existence of Monoidal Diagonals

This follows from [Items 1 and 2 of Proposition 5.1.8.1.3](#).



**01P8 5.1.10 The Universal Property of  $(\text{Sets}, \times, \text{pt})$**

01P9

**THEOREM 5.1.10.1.1 ► THE UNIVERSAL PROPERTY OF  $(\text{Sets}, \times, \text{pt})$**

The symmetric monoidal structure on the category  $\text{Sets}$  of [Proposition 5.1.9.1.1](#) is uniquely determined by the following requirements:

01PA

1. *Existence of an Internal Hom.* The tensor product

$$\otimes_{\text{Sets}} : \text{Sets} \times \text{Sets} \rightarrow \text{Sets}$$

of  $\text{Sets}$  admits an internal Hom  $[-_1, -_2]_{\text{Sets}}$ .

01PB

2. *The Unit Object Is pt.* We have  $\mathbb{1}_{\text{Sets}} \cong \text{pt}$ .

More precisely, the full subcategory of the category  $\mathcal{M}_{\mathbb{E}_\infty}^{\text{cld}}(\text{Sets})$  of ?? spanned by the closed symmetric monoidal categories  $(\text{Sets}, \otimes_{\text{Sets}}, [-_1, -_2]_{\text{Sets}}, \mathbb{1}_{\text{Sets}}, \lambda^{\text{Sets}}, \rho^{\text{Sets}}, \sigma^{\text{Sets}})$  satisfying [Items 1](#) and [2](#) is contractible (i.e. equivalent to the punctual category).

**PROOF 5.1.10.1.2 ► PROOF OF THEOREM 5.1.10.1.1**

**Unwinding the Statement**

Let  $(\text{Sets}, \otimes_{\text{Sets}}, [-_1, -_2]_{\text{Sets}}, \mathbb{1}_{\text{Sets}}, \lambda', \rho', \sigma')$  be a closed symmetric monoidal category satisfying [Items 1](#) and [2](#). We need to show that the identity functor

$$\text{id}_{\text{Sets}} : \text{Sets} \rightarrow \text{Sets}$$

admits a *unique* closed symmetric monoidal functor structure

$$\begin{aligned} \text{id}_{\text{Sets}}^\otimes &: A \otimes_{\text{Sets}} B \xrightarrow{\sim} A \times B, \\ \text{id}_{\text{Sets}}^{\text{Hom}} &: [A, B]_{\text{Sets}} \xrightarrow{\sim} \text{Sets}(A, B), \\ \text{id}_{\mathbb{1}_{\text{Sets}}}^\otimes &: \mathbb{1}_{\text{Sets}} \xrightarrow{\sim} \text{pt}, \end{aligned}$$

making it into a symmetric monoidal strongly closed isomorphism of categories from  $(\text{Sets}, \otimes_{\text{Sets}}, [-_1, -_2]_{\text{Sets}}, \mathbb{1}_{\text{Sets}}, \lambda', \rho', \sigma')$  to the closed symmetric monoidal category  $(\text{Sets}, \times, \text{Sets}(-_1, -_2), \mathbb{1}_{\text{Sets}}, \lambda^{\text{Sets}}, \rho^{\text{Sets}}, \sigma^{\text{Sets}})$  of [Proposition 5.1.9.1.1](#).

**Constructing an Isomorphism  $[-_1, -_2]_{\text{Sets}} \cong \text{Sets}(-_1, -_2)$**

By ??, we have a natural isomorphism

$$\text{Sets}(\text{pt}, [-_1, -_2]_{\text{Sets}}) \cong \text{Sets}(-_1, -_2).$$

By Item 3 of Proposition 4.3.5.1.2, we also have a natural isomorphism

$$\text{Sets}(\text{pt}, [-_1, -_2]_{\text{Sets}}) \cong [-_1, -_2]_{\text{Sets}}.$$

Composing both natural isomorphisms, we obtain a natural isomorphism

$$\text{Sets}(-_1, -_2) \cong [-_1, -_2]_{\text{Sets}}.$$

Given  $A, B \in \text{Obj}(\text{Sets})$ , we will write

$$\text{id}_{A,B}^{\text{Hom}} : \text{Sets}(A, B) \xrightarrow{\sim} [A, B]_{\text{Sets}}$$

for the component of this isomorphism at  $(A, B)$ .

Constructing an Isomorphism  $\otimes_{\text{Sets}} \cong \times$

Since  $\otimes_{\text{Sets}}$  is adjoint in each variable to  $[-_1, -_2]_{\text{Sets}}$  by assumption and  $\times$  is adjoint in each variable to  $\text{Sets}(-_1, -_2)$  by Item 2 of Proposition 4.3.5.1.2, uniqueness of adjoints (??) gives us natural isomorphisms

$$\begin{aligned} A \otimes_{\text{Sets}} - &\cong A \times -, \\ - \otimes_{\text{Sets}} B &\cong B \times -. \end{aligned}$$

By ??, we then have  $\otimes_{\text{Sets}} \cong \times$ . We will write

$$\text{id}_{\text{Sets}|A,B}^{\otimes} : A \otimes_{\text{Sets}} B \xrightarrow{\sim} A \times B$$

for the component of this isomorphism at  $(A, B)$ .

Alternative Construction of an Isomorphism  $\otimes_{\text{Sets}} \cong \times$

Alternatively, we may construct a natural isomorphism  $\otimes_{\text{Sets}} \cong \times$  as follows:

01W2 1. Let  $A \in \text{Obj}(\text{Sets})$ .

01W3 2. Since  $\otimes_{\text{Sets}}$  is part of a closed monoidal structure, it preserves colimits in each variable by ??.

01W4 3. Since  $A \cong \coprod_{a \in A} \text{pt}$  and  $\otimes_{\text{Sets}}$  preserves colimits in each variable, we have

$$\begin{aligned} A \otimes_{\text{Sets}} B &\cong (\coprod_{a \in A} \text{pt}) \otimes_{\text{Sets}} B \\ &\cong \coprod_{a \in A} (\text{pt} \otimes_{\text{Sets}} B) \\ &\cong \coprod_{a \in A} B \end{aligned}$$

$$\cong A \times B,$$

naturally in  $B \in \text{Obj}(\text{Sets})$ , where we have used that  $\text{pt}$  is the monoidal unit for  $\otimes_{\text{Sets}}$ . Thus  $A \otimes_{\text{Sets}} - \cong A \times -$  for each  $A \in \text{Obj}(\text{Sets})$ .

01W5 4. Similarly,  $- \otimes_{\text{Sets}} B \cong - \times B$  for each  $B \in \text{Obj}(\text{Sets})$ .

01W6 5. By ??, we then have  $\otimes_{\text{Sets}} \cong \times$ .

Below, we'll show that if a natural isomorphism  $\otimes_{\text{Sets}} \cong \times$  exists, then it must be unique. This will show that the isomorphism constructed above is equal to the isomorphism  $\text{id}_{\text{Sets}|A,B}^{\otimes}: A \otimes_{\text{Sets}} B \rightarrow A \times B$  from before.

#### Constructing an Isomorphism $\text{id}_{\mathbb{1}}^{\otimes}: \mathbb{1}_{\text{Sets}} \rightarrow \text{pt}$

We define an isomorphism  $\text{id}_{\mathbb{1}}^{\otimes}: \mathbb{1}_{\text{Sets}} \rightarrow \text{pt}$  as the composition

$$\begin{array}{ccccc} \mathbb{1}_{\text{Sets}} & \xrightarrow[\sim]{\rho_{\mathbb{1}_{\text{Sets}}}^{S_{\text{Sets}},-1}} & \mathbb{1}_{\text{Sets}} \times \text{pt} & \xrightarrow[\sim]{\text{id}_{\text{Sets}|1_{\text{Sets}}}^{\otimes}} & \mathbb{1}_{\text{Sets}} \otimes_{\text{Sets}} \text{pt} \xrightarrow[\sim]{\lambda'_{\text{pt}}} \text{pt} \end{array}$$

in  $\text{Sets}$ .

#### Monoidal Left Unity of the Isomorphism $\otimes_{\text{Sets}} \cong \times$

We have to show that the diagram

$$\begin{array}{ccc} \text{pt} \otimes_{\text{Sets}} A & \xrightarrow{\text{id}_{\text{Sets}|pt,A}^{\otimes}} & \text{pt} \times A \\ \text{id}_{\mathbb{1}_{\text{Sets}} \otimes_{\text{Sets}} A}^{\otimes} \text{id}_A \nearrow & & \searrow \lambda_A^{\text{Sets}} \\ \mathbb{1}_{\text{Sets}} \otimes_{\text{Sets}} A & \xrightarrow{\lambda'_A} & A \end{array}$$

commutes. First, note that the diagram

$$\begin{array}{ccc} \text{pt} \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|pt,pt}^{\otimes}} & \text{pt} \times \text{pt} \\ \text{id}_{\mathbb{1}_{\text{Sets}} \otimes_{\text{Sets}} \text{pt}}^{\otimes} \text{id}_{\text{pt}} \nearrow & & \searrow \lambda_{\text{pt}}^{\text{Sets}} \\ \mathbb{1}_{\text{Sets}} \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\lambda'_{\text{pt}}} & \text{pt}, \end{array}$$

corresponding to the case  $A = \text{pt}$ , commutes by the terminality of  $\text{pt}$  (Construction 4.1.1.1.2). Since this diagram commutes, so does the

diagram

$$\begin{array}{ccc}
 pt \times pt & \xrightarrow{id_{Sets|pt,pt}^{\otimes,-1}} & pt \otimes_{Sets} pt \\
 \lambda_{pt}^{Sets,-1} \nearrow & & \searrow id_{\mathbb{1}|Sets}^{\otimes,-1} \otimes_{Sets} id_{pt} \\
 pt & \xrightarrow{\lambda'^{-1}_{pt}} & \mathbb{1}_{Sets} \otimes_{Sets} pt.
 \end{array}$$

Now, let  $A \in Obj(Sets)$ , let  $a \in A$ , and consider the diagram

$$\begin{array}{ccccc}
 & pt \times pt & \xrightarrow{id_{Sets|pt,pt}^{\otimes,-1}} & pt \otimes_{Sets} pt & \\
 \lambda_{pt}^{Sets,-1} \nearrow & | & & | & id_{\mathbb{1}|Sets}^{\otimes,-1} \times id_{pt} \\
 pt & \xrightarrow{\lambda'^{-1}_{pt}} & & & \mathbb{1}_{Sets} \otimes_{Sets} pt \\
 & id_{pt} \times [a] & (1) & id_{pt} \otimes_{Sets} [a] & \\
 \downarrow [a] & & (3) & & \downarrow id_{\mathbb{1} Sets} \times [a] \\
 & pt \times A & \xrightarrow{id_{Sets|pt,A}^{\otimes,-1}} & pt \otimes_{Sets} A & \\
 \lambda_A^{Sets,-1} \nearrow & & (2) & & id_{\mathbb{1}|Sets}^{\otimes,-1} \times id_A \\
 A & \xrightarrow{\lambda'^{-1}_A} & & & \mathbb{1}_{Sets} \otimes_{Sets} A
 \end{array}$$

Since:

- Subdiagram (5) commutes by the naturality of  $\lambda'^{-1}$ .
- Subdiagram ( $\dagger$ ) commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $id_{\mathbb{1}|Sets}^{\otimes,-1}$ .
- Subdiagram (1) commutes by the naturality of  $id_{Sets}^{\otimes,-1}$ .
- Subdiagram (3) commutes by the naturality of  $\lambda^{Sets,-1}$ .

it follows that the diagram

$$\begin{array}{ccc}
 pt \times A & \xrightarrow{id_{Sets|pt,A}^{\otimes,-1}} & pt \otimes_{Sets} A \\
 \lambda_A^{Sets,-1} \nearrow & & \searrow id_{\mathbb{1}|Sets}^{\otimes,-1} \otimes_{Sets} id_A \\
 pt \xrightarrow{[a]} A & \xrightarrow{\lambda'^{-1}_A} & \mathbb{1}_{Sets} \otimes_{Sets} A
 \end{array}$$

Here's a step-by-step showcase of this argument: [Link]. We then have

$$\begin{aligned}\lambda_A'^{-1}(a) &= [\lambda_A'^{-1} \circ [a]](\star) \\ &= [(\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes, -1} \times \text{id}_A) \circ \text{id}_{\text{Sets}|\text{pt}, A}^{\otimes, -1} \circ \lambda_A^{\text{Sets}, -1} \circ [a]](\star) \\ &= [(\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes, -1} \times \text{id}_A) \circ \text{id}_{\text{Sets}|\text{pt}, A}^{\otimes, -1} \circ \lambda_A^{\text{Sets}, -1}](a)\end{aligned}$$

for each  $a \in A$ , and thus we have

$$\lambda_A'^{-1} = (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes, -1} \times \text{id}_A) \circ \text{id}_{\text{Sets}|\text{pt}, A}^{\otimes, -1} \circ \lambda_A^{\text{Sets}, -1}.$$

Taking inverses then gives

$$\lambda_A' = \lambda_A^{\text{Sets}} \circ \text{id}_{\text{Sets}|\text{pt}, A}^{\otimes} \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \times \text{id}_A),$$

showing that the diagram

$$\begin{array}{ccc} \text{pt} \otimes_{\text{Sets}} A & \xrightarrow{\text{id}_{\text{Sets}|\text{pt}, A}^{\otimes}} & \text{pt} \times A \\ \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \otimes_{\text{Sets}} \text{id}_A \nearrow & & \searrow \lambda_A^{\text{Sets}} \\ \mathbb{1}_{\text{Sets}} \otimes_{\text{Sets}} A & \xrightarrow{\lambda_A'} & A \end{array}$$

indeed commutes.

### Monoidal Right Unity of the Isomorphism $\otimes_{\text{Sets}} \cong \times$

We can use the same argument we used to prove the monoidal left unity of the isomorphism  $\otimes_{\text{Sets}} \cong \times$  above. For completeness, we repeat it below.

We have to show that the diagram

$$\begin{array}{ccc} A \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|A, \text{pt}}^{\otimes}} & A \times \text{pt} \\ \text{id}_A \otimes_{\text{Sets}} \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \nearrow & & \searrow \rho_A^{\text{Sets}} \\ A \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}} & \xrightarrow{\rho_A'} & A \end{array}$$

commutes. First, note that the diagram

$$\begin{array}{ccc} \text{pt} \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|\text{pt}, \text{pt}}^{\otimes}} & \text{pt} \times \text{pt} \\ \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \nearrow & & \searrow \rho_{\text{pt}}^{\text{Sets}} \\ \text{pt} \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}} & \xrightarrow{\rho'_{\text{pt}}} & \text{pt}, \end{array}$$

corresponding to the case  $A = \text{pt}$ , commutes by the terminality of  $\text{pt}$  ([Construction 4.1.1.1.2](#)). Since this diagram commutes, so does the diagram

$$\begin{array}{ccc}
 \text{pt} \times \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes,-1}} & \text{pt} \otimes_{\text{Sets}} \text{pt} \\
 \rho_{\text{pt}}^{\text{Sets},-1} \nearrow & & \searrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1} \\
 \text{pt} & \xrightarrow{\rho'^{-1}_{\text{pt}}} & \text{pt} \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}}.
 \end{array}$$

Now, let  $A \in \text{Obj}(\text{Sets})$ , let  $a \in A$ , and consider the diagram

$$\begin{array}{ccccc}
 & \text{pt} \times \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes,-1}} & \text{pt} \otimes_{\text{Sets}} \text{pt} & \\
 \rho_{\text{pt}}^{\text{Sets},-1} \nearrow & \downarrow & & \downarrow & \text{id}_{\text{pt}} \times \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1} \searrow \\
 \text{pt} & \xrightarrow{\rho'^{-1}_{\text{pt}}} & & \text{pt} \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}} & \\
 & \downarrow \text{id}_{\text{pt}} \times [a] & \text{id}_{\text{pt}} \times [a] \downarrow & \downarrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} [a] & \\
 & (3) & (1) & (4) & \\
 & \downarrow & \downarrow & \downarrow & \downarrow \text{id}_{\mathbb{1}_{\text{Sets}}} \times [a] \\
 A \times \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|A,\text{pt}}^{\otimes,-1}} & A \otimes_{\text{Sets}} \text{pt} & & \\
 \rho_A^{\text{Sets},-1} \nearrow & & \text{id}_A \times \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1} \searrow & & \\
 A & \xrightarrow{\rho'^{-1}_A} & A \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}} & &
 \end{array}$$

Since:

- Subdiagram (5) commutes by the naturality of  $\rho'^{-1}$ .
- Subdiagram  $(\dagger)$  commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1}$ .
- Subdiagram (1) commutes by the naturality of  $\text{id}_{\text{Sets}}^{\otimes,-1}$ .
- Subdiagram (3) commutes by the naturality of  $\rho^{\text{Sets},-1}$ .

it follows that the diagram

$$\begin{array}{ccccc}
 & & A \times \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|A,\text{pt}}^{\otimes,-1}} & A \otimes_{\text{Sets}} \text{pt} \\
 & \swarrow \rho_A^{\text{Sets},-1} & & & \searrow \text{id}_A \otimes_{\text{Sets}} \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1} \\
 \text{pt} & \xrightarrow{[a]} & A & \xrightarrow{\rho_A'^{-1}} & A \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}}
 \end{array}$$

Here's a step-by-step showcase of this argument: [\[Link\]](#). We then have

$$\begin{aligned}
 \rho_A'^{-1}(a) &= [\rho_A'^{-1} \circ [a]](\star) \\
 &= [(\text{id}_A \times \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1}) \circ \text{id}_{\text{Sets}|\text{pt},A}^{\otimes,-1} \circ \rho_A^{\text{Sets},-1} \circ [a]](\star) \\
 &= [(\text{id}_A \times \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1}) \circ \text{id}_{\text{Sets}|\text{pt},A}^{\otimes,-1} \circ \rho_A^{\text{Sets},-1}](a)
 \end{aligned}$$

for each  $a \in A$ , and thus we have

$$\rho_A'^{-1} = (\text{id}_A \times \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1}) \circ \text{id}_{\text{Sets}|\text{pt},A}^{\otimes,-1} \circ \rho_A^{\text{Sets},-1}.$$

Taking inverses then gives

$$\rho_A' = \rho_A^{\text{Sets}} \circ \text{id}_{\text{Sets}|\text{pt},A}^{\otimes} \circ (\text{id}_A \times \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes}),$$

showing that the diagram

$$\begin{array}{ccc}
 A \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|A,\text{pt}}^{\otimes}} & A \times \text{pt} \\
 \swarrow \text{id}_A \otimes_{\text{Sets}} \text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} & & \searrow \rho_A^{\text{Sets}} \\
 A \otimes_{\text{Sets}} \mathbb{1}_{\text{Sets}} & \xrightarrow{\rho_A'} & A
 \end{array}$$

indeed commutes.

Monoidality of the Isomorphism  $\otimes_{\text{Sets}} \cong \times$

We have to show that the diagram

$$\begin{array}{ccc}
 & (A \otimes_{\text{Sets}} B) \otimes_{\text{Sets}} C & \\
 \text{id}_{\text{Sets}|A,B}^{\otimes} \otimes_{\text{Sets}} \text{id}_C & \swarrow & \searrow \alpha'_{A,B,C} \\
 (A \times B) \otimes_{\text{Sets}} C & & A \otimes_{\text{Sets}} (B \otimes_{\text{Sets}} C) \\
 \downarrow \text{id}_{\text{Sets}|A \times B,C}^{\otimes} & & \downarrow \text{id}_A \otimes_{\text{Sets}} \text{id}_{\text{Sets}|B,C}^{\otimes} \\
 (A \times B) \times C & & A \otimes_{\text{Sets}} (B \times C) \\
 \searrow \alpha_{A,B,C}^{\text{Sets}} & & \swarrow \text{id}_{\text{Sets}|A,B \times C}^{\otimes} \\
 A \times (B \times C) & &
 \end{array}$$

commutes. First, note that the diagram

$$\begin{array}{ccc}
 & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} \text{pt} & \\
 \text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes} \otimes_{\text{Sets}} \text{id}_{\text{pt}} & \swarrow & \searrow \alpha'_{\text{pt},\text{pt},\text{pt}} \\
 (\text{pt} \times \text{pt}) \otimes_{\text{Sets}} \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) \\
 \downarrow \text{id}_{\text{Sets}|\text{pt} \times \text{pt},\text{pt}}^{\otimes} & & \downarrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes} \\
 (\text{pt} \times \text{pt}) \times \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \times \text{pt}) \\
 \searrow \alpha_{\text{pt},\text{pt},\text{pt}}^{\text{Sets}} & & \swarrow \text{id}_{\text{Sets}|\text{pt},\text{pt} \times \text{pt}}^{\otimes} \\
 \text{pt} \times (\text{pt} \times \text{pt}) & & \\
 \downarrow !_{\text{pt} \times (\text{pt} \times \text{pt})} & & \\
 \text{pt} & &
 \end{array}$$

commutes by the terminality of  $\text{pt}$  (Construction 4.1.1.1.2). Since the map  $!_{\text{pt} \times (\text{pt} \times \text{pt})} : \text{pt} \times (\text{pt} \times \text{pt}) \rightarrow \text{pt}$  is an isomorphism (e.g. having inverse

$\lambda_{\text{pt}}^{\text{Sets}, -1} \circ \lambda_{\text{pt}}^{\text{Sets}, -1}$ , it follows that the diagram

$$\begin{array}{ccc}
 & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} \text{pt} & \\
 \text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes} \otimes_{\text{Sets}} \text{id}_{\text{pt}} & \swarrow & \searrow \alpha'_{\text{pt},\text{pt},\text{pt}} \\
 (\text{pt} \times \text{pt}) \otimes_{\text{Sets}} \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) \\
 \downarrow & & \downarrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes} \\
 (\text{pt} \times \text{pt}) \times \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \times \text{pt}) \\
 \downarrow \alpha_{\text{pt},\text{pt},\text{pt}}^{\text{Sets}} & \searrow & \swarrow \text{id}_{\text{Sets}|\text{pt},\text{pt} \times \text{pt}}^{\otimes} \\
 \text{pt} \times (\text{pt} \times \text{pt}) & & 
 \end{array}$$

also commutes. Taking inverses, we see that the diagram

$$\begin{array}{ccc}
 & \text{pt} \times (\text{pt} \times \text{pt}) & \\
 \alpha_{\text{pt},\text{pt},\text{pt}}^{\text{Sets}, -1} & \swarrow & \searrow \text{id}_{\text{Sets}|\text{pt},\text{pt} \times \text{pt}}^{\otimes, -1} \\
 (\text{pt} \times \text{pt}) \times \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \times \text{pt}) \\
 \downarrow \text{id}_{\text{Sets}|\text{pt} \times \text{pt},\text{pt}}^{\otimes, -1} & \text{(\dagger)} & \downarrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes, -1} \\
 (\text{pt} \times \text{pt}) \otimes_{\text{Sets}} \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) \\
 \downarrow \text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_{\text{pt}} & & \swarrow \alpha'_{\text{pt},\text{pt},\text{pt}}^{\otimes, -1} \\
 (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} \text{pt} & & 
 \end{array}$$


commutes as well. Now, let  $A, B, C \in \text{Obj}(\text{Sets})$ , let  $a \in A$ , let  $b \in B$ , let

$c \in C$ , and consider the diagram

$$\begin{array}{ccccc}
& \text{pt} \times (\text{pt} \times \text{pt}) & & A \times (B \times C) & \\
\alpha_{\text{pt},\text{pt} \times \text{pt}}^{-1} \swarrow & \downarrow \text{id}_{\text{pt} \times \text{pt} \times \text{pt}}^{\otimes,-1} & \searrow [a] \times ([b] \times [c]) & & \\
(\text{pt} \times \text{pt}) \times \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \times \text{pt}) & & A \times (B \times C) \\
\downarrow \text{id}_{\text{pt} \times \text{pt},\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_{\text{Sets}(A,B \times C)}^{\otimes,-1} \\
(\text{pt} \times \text{pt}) \otimes_{\text{Sets}} \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) & & A \otimes_{\text{Sets}} (B \times C) \\
\downarrow \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \otimes_{\text{Sets}} \text{id}_{\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_{\text{pt} \otimes_{\text{Sets}} \text{pt},\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_A \otimes_{\text{Sets}} \text{id}_{\text{Sets}(B,C)}^{\otimes,-1} \\
(\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} \text{pt} & & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) & & A \otimes_{\text{Sets}} (B \otimes_{\text{Sets}} C) \\
& \downarrow \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \otimes_{\text{Sets}} \text{id}_{\text{pt}}^{\otimes,-1} & \downarrow \text{id}_{\text{pt} \otimes_{\text{Sets}} \text{pt},\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \\
& & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) & & \\
& & \downarrow \text{id}_{\text{pt} \otimes_{\text{Sets}} \text{pt},\text{pt}}^{\otimes,-1} & & \\
& & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) & & \\
& & \downarrow \text{id}_{\text{pt} \otimes_{\text{Sets}} \text{pt},\text{pt}}^{\otimes,-1} & & \\
& & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) & & 
\end{array}$$

which we partition into subdiagrams as follows:

$$\begin{array}{ccc}
& \text{pt} \times (\text{pt} \times \text{pt}) & \\
\alpha_{\text{pt},\text{pt} \times \text{pt}}^{-1} \swarrow & \downarrow \text{id}_{\text{pt} \times \text{pt} \times \text{pt}}^{\otimes,-1} & \\
(\text{pt} \times \text{pt}) \times \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \times \text{pt}) \\
\downarrow \text{id}_{\text{pt} \times \text{pt},\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_{\text{pt}} \otimes_{\text{Sets}} \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \\
(\text{pt} \times \text{pt}) \otimes_{\text{Sets}} \text{pt} & & \text{pt} \otimes_{\text{Sets}} (\text{pt} \otimes_{\text{Sets}} \text{pt}) \\
\downarrow \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \otimes_{\text{Sets}} \text{id}_{\text{pt}}^{\otimes,-1} & & \downarrow \text{id}_{\text{pt} \otimes_{\text{Sets}} \text{pt},\text{pt}}^{\otimes,-1} \\
(\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} \text{pt} & & (\dagger) \\
& \downarrow \text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \otimes_{\text{Sets}} \text{id}_{\text{pt}}^{\otimes,-1} & \downarrow \text{id}_{\text{pt} \otimes_{\text{Sets}} \text{pt},\text{pt}}^{\otimes,-1} \\
& & (\text{pt} \otimes_{\text{Sets}} \text{pt}) \otimes_{\text{Sets}} \text{pt}
\end{array}$$



it follows that the diagram

$$\begin{array}{ccccc}
 & & \text{pt} \times (\text{pt} \times \text{pt}) & & \\
 & & \downarrow & & \\
 & & [a] \times ([b] \times [c]) & & \\
 & & \downarrow & & \\
 A \times (B \times C) & & & & \\
 \alpha_{A,B,C}^{\text{Sets}, -1} \swarrow & & \searrow \text{id}_{\text{Sets}|A,B \times C}^{\otimes, -1} \\
 (A \times B) \times C & & & & A \otimes_{\text{Sets}} (B \times C) \\
 \downarrow \text{id}_{\text{Sets}|A \times B, C}^{\otimes, -1} & & & & \downarrow \text{id}_A \times \text{id}_{\text{Sets}|B, C}^{\otimes, -1} \\
 (A \times B) \otimes_{\text{Sets}} C & & & & A \otimes_{\text{Sets}} (B \otimes_{\text{Sets}} C) \\
 \downarrow \text{id}_{\text{Sets}|A, B}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_C & & & & \downarrow \alpha'_{A,B,C}^{-1} \\
 (A \otimes_{\text{Sets}} B) \otimes_{\text{Sets}} C & & & &
 \end{array}$$

also commutes. We then have

$$\begin{aligned}
 & \left[ (\text{id}_{\text{Sets}|A, B}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_C) \circ \text{id}_{\text{Sets}|A \times B, C}^{\otimes, -1} \right. \\
 & \quad \left. \circ \alpha_{A, B, C}^{\text{Sets}, -1} \right] (a, (b, c)) = \left[ (\text{id}_{\text{Sets}|A, B}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_C) \circ \text{id}_{\text{Sets}|A \times B, C}^{\otimes, -1} \right. \\
 & \quad \left. \circ \alpha_{A, B, C}^{\text{Sets}, -1} \circ ([a] \times ([b] \times [c])) \right] (\star, (\star, \star)) \\
 & = \left[ \alpha'_{A, B, C}^{-1} \circ (\text{id}_A \times \text{id}_{\text{Sets}|B, C}^{\otimes, -1}) \right. \\
 & \quad \left. \circ \text{id}_{\text{Sets}|A, B \times C}^{\otimes, -1} \circ ([a] \times ([b] \times [c])) \right] (\star, (\star, \star)) \\
 & = [\alpha'_{A, B, C}^{-1} \circ (\text{id}_A \times \text{id}_{\text{Sets}|B, C}^{\otimes, -1}) \circ \text{id}_{\text{Sets}|A, B \times C}^{\otimes, -1}] (a, (b, c))
 \end{aligned}$$

for each  $(a, (b, c)) \in A \times (B \times C)$ , and thus we have

$$(\text{id}_{\text{Sets}|A, B}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_C) \circ \text{id}_{\text{Sets}|A \times B, C}^{\otimes, -1} \circ \alpha_{A, B, C}^{\text{Sets}, -1} = \alpha'_{A, B, C}^{-1} \circ (\text{id}_A \times \text{id}_{\text{Sets}|B, C}^{\otimes, -1}) \circ \text{id}_{\text{Sets}|A, B \times C}^{\otimes, -1}.$$

Taking inverses then gives

$$\alpha_{A, B, C}^{\text{Sets}} \circ \text{id}_{\text{Sets}|A \times B, C}^{\otimes} \circ (\text{id}_{\text{Sets}|A, B}^{\otimes} \otimes_{\text{Sets}} \text{id}_C) = \text{id}_{\text{Sets}|A, B \times C}^{\otimes} \circ (\text{id}_A \times \text{id}_{\text{Sets}|B, C}^{\otimes}) \circ \alpha'_{A, B, C},$$

showing that the diagram

$$\begin{array}{ccc}
 & (A \otimes_{\text{Sets}} B) \otimes_{\text{Sets}} C & \\
 \text{id}_{\text{Sets}|A,B}^{\otimes} \otimes_{\text{Sets}} \text{id}_C & \swarrow & \searrow \alpha'_{A,B,C} \\
 (A \times B) \otimes_{\text{Sets}} C & & A \otimes_{\text{Sets}} (B \otimes_{\text{Sets}} C) \\
 \downarrow \text{id}_{\text{Sets}|A \times B,C}^{\otimes} & & \downarrow \text{id}_A \otimes_{\text{Sets}} \text{id}_{\text{Sets}|B,C}^{\otimes} \\
 (A \times B) \times C & & A \otimes_{\text{Sets}} (B \times C) \\
 \searrow \alpha_{A,B,C}^{\text{Sets}} & & \swarrow \text{id}_{\text{Sets}|A,B \times C}^{\otimes} \\
 A \times (B \times C) & &
 \end{array}$$

indeed commutes.

**Braidedness of the Isomorphism  $\otimes_{\text{Sets}} \cong \times$**

We have to show that the diagram

$$\begin{array}{ccc}
 A \otimes_{\text{Sets}} B & \xrightarrow{\text{id}_{\text{Sets}|A,B}^{\otimes}} & A \times B \\
 \sigma'_{A,B} \downarrow & & \downarrow \sigma_{A,B}^{\text{Sets}} \\
 B \otimes_{\text{Sets}} A & \xrightarrow{\text{id}_{\text{Sets}|B,A}^{\otimes}} & B \times A
 \end{array}$$

commutes. First, note that the diagram

$$\begin{array}{ccc}
 \text{pt} \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes}} & \text{pt} \times \text{pt} \\
 \sigma'_{\text{pt},\text{pt}} \downarrow & & \downarrow \sigma_{\text{pt},\text{pt}}^{\text{Sets}} \\
 \text{pt} \otimes_{\text{Sets}} \text{pt} & \xrightarrow{\text{id}_{\text{Sets}|\text{pt},\text{pt}}^{\otimes}} & \text{pt} \times \text{pt} \\
 & & \searrow !_{\text{pt} \times \text{pt}}
 \end{array}$$

commutes by the terminality of  $\text{pt}$  ([Construction 4.1.1.1.2](#)). Since the map  $!_{\text{pt} \times \text{pt}}: \text{pt} \times \text{pt} \rightarrow \text{pt}$  is invertible (e.g. with inverse  $\lambda_{\text{pt}}^{\text{Sets}, -1}$ ), the

diagram

$$\begin{array}{ccc}
 pt \otimes_{\text{Sets}} pt & \xrightarrow{\text{id}_{\text{Sets}|pt,pt}^{\otimes}} & pt \times pt \\
 \sigma'_{pt,pt} \downarrow & & \downarrow \sigma_{pt,pt}^{\text{Sets}} \\
 pt \otimes_{\text{Sets}} pt & \xrightarrow{\text{id}_{\text{Sets}|pt,pt}^{\otimes}} & pt \times pt
 \end{array}$$

also commutes. Taking inverses, we see that the diagram

$$\begin{array}{ccc}
 pt \times pt & \xrightarrow{\text{id}_{\text{Sets}|pt,pt}^{\otimes,-1}} & pt \otimes_{\text{Sets}} pt \\
 \sigma_{pt,pt}^{\text{Sets},-1} \downarrow & (\dagger) & \downarrow \sigma_{pt,pt}^{\prime,-1} \\
 pt \times pt & \xrightarrow{\text{id}_{\text{Sets}|pt,pt}^{\otimes,-1}} & pt \otimes_{\text{Sets}} pt
 \end{array}$$

commutes as well. Now, let  $A, B \in \text{Obj}(\text{Sets})$ , let  $a \in A$ , let  $b \in B$ , and consider the diagram

$$\begin{array}{ccccc}
 & id_{pt,pt}^{\otimes,-1} & & & \\
 pt \times pt & \xrightarrow{\quad} & pt \otimes_{\text{Sets}} pt & \xrightarrow{\quad} & \\
 \downarrow \sigma_{pt,pt}^{\text{Sets},-1} & \searrow [b] \times [a] & \downarrow \sigma_{pt,pt}^{\prime,-1} & \swarrow [b] \otimes_{\text{Sets}} [a] & \\
 & B \times A & \xrightarrow{\text{id}_{\text{Sets}|B,A}^{\otimes,-1}} & B \otimes_{\text{Sets}} A & \\
 \downarrow \sigma_{A,B}^{\text{Sets},-1} & & \downarrow & & \downarrow \sigma_{A,B}^{\prime,-1} \\
 pt \times pt & \xrightarrow{\text{id}_{pt,pt}^{\otimes,-1}} & pt \otimes_{\text{Sets}} pt & \xrightarrow{\quad} & \\
 \downarrow [a] \times [b] & \searrow & \downarrow [a] \otimes_{\text{Sets}} [b] & \swarrow & \\
 A \times B & \xrightarrow{\text{id}_{\text{Sets}|A,B}^{\otimes,-1}} & A \otimes_{\text{Sets}} B & &
 \end{array}$$

which we partition into subdiagrams as follows:

$$\begin{array}{ccc}
 \begin{array}{c}
 \text{pt} \times \text{pt} \xrightarrow{\text{id}_{\text{pt},\text{pt}}^{\otimes,-1}} \text{pt} \otimes_{\text{Sets}} \text{pt} \\
 \downarrow \sigma_{\text{pt},\text{pt}}^{\text{Sets},-1} \quad \swarrow [b] \times [a] \quad \searrow [b] \otimes_{\text{Sets}} [a] \\
 \text{pt} \times \text{pt} \xrightarrow{\text{id}_{\text{Sets}|B,A}^{\otimes,-1}} B \otimes_{\text{Sets}} A \\
 \downarrow \sigma_{A,B}^{\text{Sets},-1} \quad \text{(3)} \quad \downarrow \sigma'_{A,B}^{-1} \\
 \text{pt} \times \text{pt} \xrightarrow{\text{id}_{\text{Sets}|A,B}^{\otimes,-1}} A \otimes_{\text{Sets}} B
 \end{array} & \quad & 
 \begin{array}{c}
 \text{pt} \times \text{pt} \xrightarrow{\text{id}_{\text{pt},\text{pt}}^{\otimes,-1}} \text{pt} \otimes_{\text{Sets}} \text{pt} \\
 \downarrow \sigma_{\text{pt},\text{pt}}^{\text{Sets},-1} \quad \text{(\dagger)} \quad \downarrow \sigma'_{\text{pt},\text{pt}}^{-1} \\
 \text{pt} \times \text{pt} \xrightarrow{\text{id}_{\text{pt},\text{pt}}^{\otimes,-1}} \text{pt} \otimes_{\text{Sets}} \text{pt} \\
 \downarrow \sigma'_{A,B}^{-1} \quad \text{(4)} \quad \downarrow \sigma'_{A,B}^{-1} \\
 \text{pt} \times \text{pt} \xrightarrow{\text{id}_{\text{Sets}|A,B}^{\otimes,-1}} A \otimes_{\text{Sets}} B
 \end{array}
 \end{array}$$

Since:

- Subdiagram (2) commutes by the naturality of  $\sigma^{\text{Sets},-1}$ .
- Subdiagram (5) commutes by the naturality of  $\text{id}^{\otimes,-1}$ .
- Subdiagram ( $\dagger$ ) commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $\sigma'^{-1}$ .
- Subdiagram (1) commutes by the naturality of  $\text{id}^{\otimes,-1}$ .

it follows that the diagram

$$\begin{array}{ccc}
 \text{pt} \times \text{pt} & \xrightarrow{\text{id}_{\text{pt},\text{pt}}^{\otimes,-1} \circ \sigma_{A,B}^{\text{Sets},-1}} & B \times A \xrightarrow{\text{id}_{\text{Sets}|B,A}^{\otimes,-1}} B \otimes_{\text{Sets}} A \\
 & \swarrow [b] \times [a] & \downarrow \sigma_{A,B}^{\text{Sets}} \\
 & & A \times B \xrightarrow{\text{id}_{\text{Sets}|A,B}^{\otimes,-1}} A \otimes_{\text{Sets}} B
 \end{array}$$

commutes. We then have

$$\begin{aligned}
 [\text{id}_{\text{Sets}|A,B}^{\otimes,-1} \circ \sigma_{A,B}^{\text{Sets},-1}](b, a) &= [\text{id}_{\text{Sets}|A,B}^{\otimes,-1} \circ \sigma_{A,B}^{\text{Sets},-1} \circ ([b] \times [a])](\star, \star) \\
 &= [\sigma_{A,B}^{\text{Sets}}^{-1} \circ \text{id}_{\text{Sets}|B,A}^{\otimes,-1} \circ ([b] \times [a])](\star, \star) \\
 &= [\sigma_{A,B}^{\text{Sets}}^{-1} \circ \text{id}_{\text{Sets}|B,A}^{\otimes,-1}](b, a)
 \end{aligned}$$

for each  $(b, a) \in B \times A$ , and thus we have

$$\text{id}_{\text{Sets}|A,B}^{\otimes,-1} \circ \sigma_{A,B}^{\text{Sets},-1} = \sigma'_{A,B}^{-1} \circ \text{id}_{\text{Sets}|B,A}^{\otimes,-1}.$$

Taking inverses then gives

$$\sigma_{A,B}^{\text{Sets}} \circ \text{id}_{\text{Sets}|A,B}^{\otimes} = \text{id}_{\text{Sets}|B,A}^{\otimes} \circ \sigma'_{A,B},$$

showing that the diagram

$$\begin{array}{ccc} A \otimes_{\text{Sets}} B & \xrightarrow{\text{id}_{\text{Sets}|A,B}^{\otimes}} & A \times B \\ \sigma'_{A,B} \downarrow & & \downarrow \sigma_{A,B}^{\text{Sets}} \\ B \otimes_{\text{Sets}} A & \xrightarrow{\text{id}_{\text{Sets}|B,A}^{\otimes}} & B \times A \end{array}$$

indeed commutes.

Uniqueness of the Isomorphism  $\otimes_{\text{Sets}} \cong \times$

Let  $\phi, \psi: -_1 \otimes_{\text{Sets}} -_2 \Rightarrow -_1 \times -_2$  be natural isomorphisms. Since these isomorphisms are compatible with the unitors of  $\text{Sets}$  with respect to  $\times$  and  $\otimes$  (as shown above), we have

$$\begin{aligned} \lambda'_B &= \lambda_B^{\text{Sets}} \circ \phi_{\text{pt},B} \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \otimes_{\text{Sets}} \text{id}_Y), \\ \lambda'_B &= \lambda_B^{\text{Sets}} \circ \psi_{\text{pt},B} \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \otimes_{\text{Sets}} \text{id}_Y). \end{aligned}$$

Postcomposing both sides with  $\lambda_B^{\text{Sets},-1}$  gives

$$\begin{aligned} \lambda_B^{\text{Sets},-1} \circ \lambda'_B \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes,-1} \otimes_{\text{Sets}} \text{id}_Y) &= \phi_{\text{pt},B}, \\ \lambda_B^{\text{Sets},-1} \circ \lambda'_B \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \otimes_{\text{Sets}} \text{id}_Y) &= \psi_{\text{pt},B}, \end{aligned}$$

and thus we have

$$\phi_{\text{pt},B} = \psi_{\text{pt},B}$$

for each  $B \in \text{Obj}(\text{Sets})$ . Now, let  $a \in A$  and consider the naturality diagrams

$$\begin{array}{ccc} \text{pt} \times B & \xrightarrow{[a] \times \text{id}_B} & A \times B \\ \phi_{\text{pt},B} \downarrow & & \downarrow \phi_{A,B} \\ \text{pt} \otimes_{\text{Sets}} B & \xrightarrow{[a] \otimes_{\text{Sets}} \text{id}_B} & A \otimes_{\text{Sets}} B \end{array} \quad \begin{array}{ccc} \text{pt} \times B & \xrightarrow{[a] \times \text{id}_B} & A \times B \\ \psi_{\text{pt},B} \downarrow & & \downarrow \psi_{A,B} \\ \text{pt} \otimes_{\text{Sets}} B & \xrightarrow{[a] \otimes_{\text{Sets}} \text{id}_B} & A \otimes_{\text{Sets}} B \end{array}$$

for  $\phi$  and  $\psi$  with respect to the morphisms  $[a]$  and  $\text{id}_B$ . Having shown that  $\phi_{\text{pt},B} = \psi_{\text{pt},B}$ , we have

$$\begin{aligned}\phi_{A,B}(a,b) &= [\phi_{A,B} \circ ([a] \times \text{id}_B)](\star, b) \\ &= [([a] \otimes_{\text{Sets}} \text{id}_B) \circ \phi_{\text{pt},B}](\star, b) \\ &= [([a] \otimes_{\text{Sets}} \text{id}_B) \circ \psi_{\text{pt},B}](\star, b) \\ &= [\psi_{A,B} \circ ([a] \times \text{id}_B)](\star, b) \\ &= \psi_{A,B}(a, b)\end{aligned}$$

for each  $(a, b) \in A \times B$ . Therefore we have

$$\phi_{A,B} = \psi_{A,B}$$

for each  $A, B \in \text{Obj}(\text{Sets})$  and thus  $\phi = \psi$ , showing the isomorphism  $\otimes_{\text{Sets}} \cong \times$  to be unique. 

01PH

#### COROLLARY 5.1.10.1.3 ► A SECOND UNIVERSAL PROPERTY FOR $(\text{Sets}, \times, \text{pt})$

The symmetric monoidal structure on the category  $\text{Sets}$  of [Proposition 5.1.9.1.1](#) is uniquely determined by the following requirements:

01PJ

1. *Two-Sided Preservation of Colimits.* The tensor product

$$\otimes_{\text{Sets}} : \text{Sets} \times \text{Sets} \rightarrow \text{Sets}$$

of  $\text{Sets}$  preserves colimits separately in each variable.

01PK

2. *The Unit Object Is pt.* We have  $\mathbb{1}_{\text{Sets}} \cong \text{pt}$ .

More precisely, the full subcategory of the category  $\mathcal{M}_{\mathbb{E}_\infty}(\text{Sets})$  of ?? spanned by the symmetric monoidal categories  $(\text{Sets}, \otimes_{\text{Sets}}, \mathbb{1}_{\text{Sets}}, \lambda^{\text{Sets}}, \rho^{\text{Sets}}, \sigma^{\text{Sets}})$  satisfying [Items 1](#) and [2](#) is contractible.

#### PROOF 5.1.10.1.4 ► PROOF OF COROLLARY 5.1.10.1.3

Since  $\text{Sets}$  is locally presentable (??), it follows from ?? that [Item 1](#) is equivalent to the existence of an internal Hom as in [Item 1](#) of [Theorem 5.1.10.1.1](#). The result then follows from [Theorem 5.1.10.1.1](#). 

## 01PL 5.2 The Monoidal Category of Sets and Coproducts

## 01PM 5.2.1 Coproducts of Sets

See Section 4.2.3.

01PN 5.2.2 The Monoidal Unit

01PP

## DEFINITION 5.2.2.1.1 ► THE MONOIDAL UNIT OF $\sqcup$

The **monoidal unit** of the coproduct of sets is the functor

$$\emptyset^{\text{Sets}} : \text{pt} \rightarrow \text{Sets}$$

defined by

$$\emptyset_{\text{Sets}} \stackrel{\text{def}}{=} \emptyset,$$

where  $\emptyset$  is the empty set of Definition 4.3.1.1.1.

### 01PQ 5.2.3 The Associator


01PR

#### DEFINITION 5.2.3.1.1 ► THE ASSOCIATOR OF $\sqcup$

The associator of the coproduct of sets is the natural isomorphism

$$\alpha^{\text{Sets}, \coprod} : \coprod \circ (\coprod \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \coprod \circ (\text{id}_{\text{Sets}} \times \coprod) \circ \alpha^{\text{Cats}}_{\text{Sets}, \text{Sets}, \text{Sets}},$$

as in the diagram



whose component

$$\alpha_{X,Y,Z}^{\text{Sets}, \coprod}: (X \coprod Y) \coprod Z \xrightarrow{\sim} X \coprod (Y \coprod Z)$$

at  $(X, Y, Z)$  is given by

$$\alpha_{X,Y,Z}^{\text{Sets}, \coprod}(a) \stackrel{\text{def}}{=} \begin{cases} (0, x) & \text{if } a = (0, (0, x)), \\ (1, (0, y)) & \text{if } a = (0, (1, y)), \\ (1, (1, a)) & \text{if } a = (1, z) \end{cases}$$

for each  $a \in (X \coprod Y) \coprod Z$ .

#### PROOF 5.2.3.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.2.3.1.1

Unwinding the Definitions of  $(X \coprod Y) \coprod Z$  and  $X \coprod (Y \coprod Z)$

Firstly, we unwind the expressions for  $(X \coprod Y) \coprod Z$  and  $X \coprod (Y \coprod Z)$ . We have

$$\begin{aligned} (X \coprod Y) \coprod Z &\stackrel{\text{def}}{=} \{(0, a) \in S \mid a \in X \coprod Y\} \cup \{(1, z) \in S \mid z \in Z\} \\ &= \{(0, (0, x)) \in S \mid x \in X\} \cup \{(0, (1, y)) \in S \mid y \in Y\} \\ &\quad \cup \{(1, z) \in S \mid z \in Z\}, \end{aligned}$$

where  $S = \{0, 1\} \times ((X \coprod Y) \cup Z)$  and

$$\begin{aligned} X \coprod (Y \coprod Z) &\stackrel{\text{def}}{=} \{(0, x) \in S' \mid x \in X\} \cup \{(1, a) \in S' \mid a \in Y \coprod Z\} \\ &= \{(0, x) \in S' \mid x \in X\} \cup \{(1, (0, y)) \in S' \mid y \in Y\} \\ &\quad \cup \{(1, (1, z)) \in S' \mid z \in Z\}, \end{aligned}$$

where  $S' = \{0, 1\} \times (X \cup (Y \coprod Z))$ .

#### Invertibility

The inverse of  $\alpha_{X,Y,Z}^{\text{Sets}, \coprod}$  is the map

$$\alpha_{X,Y,Z}^{\text{Sets}, \coprod, -1} : X \coprod (Y \coprod Z) \rightarrow (X \coprod Y) \coprod Z$$

given by

$$\alpha_{X,Y,Z}^{\text{Sets}, \coprod, -1}(a) \stackrel{\text{def}}{=} \begin{cases} (0, (0, x)) & \text{if } a = (0, x), \\ (0, (1, y)) & \text{if } a = (1, (0, y)), \\ (1, z) & \text{if } a = (1, (1, z)) \end{cases}$$

for each  $a \in X \coprod Y \coprod Z$ . Indeed:

- *Invertibility I.* The map  $\alpha_{X,Y,Z}^{\text{Sets}, \coprod, -1} \circ \alpha_{X,Y,Z}^{\text{Sets}, \coprod}$  acts on elements as

$$(0, (0, x)) \mapsto (0, x) \mapsto (0, (0, x)),$$

$$\begin{aligned}(0, (0, y)) &\mapsto (1, (0, y)) \mapsto (0, (0, y)), \\ (1, z) &\mapsto (1, (1, z)) \mapsto (1, z)\end{aligned}$$

and hence is equal to the identity map of  $(X \coprod Y) \coprod Z$ .

- *Invertibility II.* The map  $\alpha_{X,Y,Z}^{\text{Sets}, \coprod} \circ \alpha_{X,Y,Z}^{\text{Sets}, \coprod, -1}$  acts on elements as

$$\begin{aligned}(0, x) &\mapsto (0, (0, x)) \mapsto (0, x), \\ (1, (0, y)) &\mapsto (0, (0, y)) \mapsto (1, (0, y)), \\ (1, (1, z)) &\mapsto (1, z) \mapsto (1, (1, z))\end{aligned}$$

and hence is equal to the identity map of  $X \coprod (Y \coprod Z)$ .

Therefore  $\alpha_{X,Y,Z}^{\text{Sets}, \coprod}$  is indeed an isomorphism.

### Naturality

We need to show that, given functions

$$\begin{aligned}f: X &\rightarrow X', \\ g: Y &\rightarrow Y', \\ h: Z &\rightarrow Z'\end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \coprod Y) \coprod Z & \xrightarrow{(f \coprod g) \coprod h} & (X' \coprod Y') \coprod Z' \\ \downarrow \alpha_{X,Y,Z}^{\text{Sets}, \coprod} & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}, \coprod} \\ X \coprod (Y \coprod Z) & \xrightarrow{f \coprod (g \coprod h)} & X' \coprod (Y' \coprod Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (0, (0, x)) & & (0, (0, x)) \mapsto (0, (0, f(x))) \\
 \downarrow & & \downarrow \\
 (0, x) \longmapsto (0, f(x)) & & (0, f(x)) \\
 (0, (1, y)) & & (0, (1, y)) \mapsto (0, (1, g(y))) \\
 \downarrow & & \downarrow \\
 (1, (0, y)) \mapsto (1, (0, g(y))) & & (1, (0, g(y))) \\
 (1, z) & & (1, z) \longmapsto (1, h(z)) \\
 \downarrow & & \downarrow \\
 (1, (1, z)) \mapsto (1, (1, h(z))) & & (1, (1, h(z)))
 \end{array}$$

and hence indeed commutes, showing  $\alpha^{\text{Sets}, \coprod}$  to be a natural transformation.

Being a Natural Isomorphism

Since  $\alpha^{\text{Sets}, \coprod}$  is natural and  $\alpha^{\text{Sets}, \coprod, -1}$  is a componentwise inverse to  $\alpha^{\text{Sets}, \coprod}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\lambda^{\text{Sets}, -1}$  is also natural. Thus  $\alpha^{\text{Sets}, \coprod}$  is a natural isomorphism. ■

#### 01PS 5.2.4 The Left Unitor

01PT

##### DEFINITION 5.2.4.1.1 ► THE LEFT UNITOR OF $\coprod$

The **left unitor of the coproduct of sets** is the natural isomorphism

$$\begin{array}{ccc}
 \text{pt} \times \text{Sets} & \xrightarrow{\emptyset^{\text{Sets}} \times \text{id}} & \text{Sets} \times \text{Sets} \\
 \lambda^{\text{Sets}, \coprod} : \coprod \circ (\emptyset^{\text{Sets}} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \lambda_{\text{Sets}}^{\text{Cats}_2} & \swarrow \quad \uparrow \quad \searrow & \downarrow \\
 & \lambda_{\text{Sets}}^{\text{Cats}_2} & \lambda^{\text{Sets}, \coprod} \\
 & \dashrightarrow & \dashrightarrow \\
 & & \text{Sets,}
 \end{array}$$

whose component

$$\lambda_X^{\text{Sets}, \coprod} : \emptyset \coprod X \xrightarrow{\sim} X$$

at  $X$  is given by

$$\lambda_X^{\text{Sets}, \coprod}((1, x)) \stackrel{\text{def}}{=} x$$

for each  $(1, x) \in \emptyset \coprod X$ .

#### PROOF 5.2.4.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.2.4.1.1

##### Unwinding the Definition of $\emptyset \coprod X$

Firstly, we unwind the expressions for  $\emptyset \coprod X$ . We have

$$\begin{aligned} \emptyset \coprod X &\stackrel{\text{def}}{=} \{(0, z) \in S \mid z \in \emptyset\} \cup \{(1, x) \in S \mid x \in X\} \\ &= \emptyset \cup \{(1, x) \in S \mid x \in X\} \\ &= \{(1, x) \in S \mid x \in X\}, \end{aligned}$$

where  $S = \{0, 1\} \times (\emptyset \cup X)$ .

##### Invertibility

The inverse of  $\lambda_X^{\text{Sets}, \coprod}$  is the map

$$\lambda_X^{\text{Sets}, \coprod, -1} : X \rightarrow \emptyset \coprod X$$

given by

$$\lambda_X^{\text{Sets}, \coprod, -1}(x) \stackrel{\text{def}}{=} (1, x)$$

for each  $x \in X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned} [\lambda_X^{\text{Sets}, \coprod, -1} \circ \lambda_X^{\text{Sets}, \coprod}]((1, x)) &= \lambda_X^{\text{Sets}, \coprod, -1}(\lambda_X^{\text{Sets}, \coprod}((1, x))) \\ &= \lambda_X^{\text{Sets}, \coprod, -1}(x) \\ &= (1, x) \\ &= [\text{id}_{\emptyset \coprod X}]((1, x)) \end{aligned}$$

for each  $(1, x) \in \emptyset \coprod X$ , and therefore we have

$$\lambda_X^{\text{Sets}, \coprod, -1} \circ \lambda_X^{\text{Sets}, \coprod} = \text{id}_{\emptyset \coprod X}.$$

- *Invertibility II.* We have

$$[\lambda_X^{\text{Sets}, \coprod} \circ \lambda_X^{\text{Sets}, \coprod, -1}](x) = \lambda_X^{\text{Sets}, \coprod}(\lambda_X^{\text{Sets}, \coprod, -1}(x))$$

$$\begin{aligned}
 &= \lambda_X^{\text{Sets}, \coprod, -1}(1, x) \\
 &= x \\
 &= [\text{id}_X](x)
 \end{aligned}$$

for each  $x \in X$ , and therefore we have

$$\lambda_X^{\text{Sets}, \coprod} \circ \lambda_X^{\text{Sets}, \coprod, -1} = \text{id}_X.$$

Therefore  $\lambda_X^{\text{Sets}, \coprod}$  is indeed an isomorphism.

### Naturality

We need to show that, given a function  $f: X \rightarrow Y$ , the diagram

$$\begin{array}{ccc}
 \emptyset \coprod X & \xrightarrow{\text{id}_{\emptyset} \coprod f} & \emptyset \coprod Y \\
 \downarrow \lambda_X^{\text{Sets}, \coprod} & & \downarrow \lambda_Y^{\text{Sets}, \coprod} \\
 X & \xrightarrow{f} & Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (1, x) & & (1, x) \mapsto (1, f(x)) \\
 \downarrow & & \downarrow \\
 x \mapsto f(x) & & f(x)
 \end{array}$$

and hence indeed commutes. Therefore  $\lambda^{\text{Sets}, \coprod}$  is a natural transformation.

### Being a Natural Isomorphism

Since  $\lambda^{\text{Sets}, \coprod}$  is natural and  $\lambda^{\text{Sets}, -1}$  is a componentwise inverse to  $\lambda^{\text{Sets}, \coprod}$ , it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\lambda^{\text{Sets}, -1}$  is also natural. Thus  $\lambda^{\text{Sets}, \coprod}$  is a natural isomorphism. □

**01PU 5.2.5 The Right Unitor**

**01PV**

**DEFINITION 5.2.5.1.1 ► THE RIGHT UNITOR OF  $\coprod$**

The **right unitor of the coproduct of sets** is the natural isomorphism

$$\begin{array}{ccc} \text{Sets} \times \text{pt} & \xrightarrow{\text{id} \times \emptyset^{\text{Sets}}} & \text{Sets} \times \text{Sets} \\ \rho^{\text{Sets}, \coprod} : \coprod \circ (\text{id} \times \emptyset^{\text{Sets}}) \xrightarrow{\sim} \rho_{\text{Sets}}^{\text{Cats}_2}, & \swarrow \quad \searrow & \downarrow \coprod \\ & \rho^{\text{Sets}, \coprod}_{\text{Sets}} & \\ & \rho_{\text{Sets}}^{\text{Cats}_2} & \end{array}$$

$\coprod$

$\text{Sets}$ ,

whose component

$$\rho_X^{\text{Sets}, \coprod} : X \coprod \emptyset \xrightarrow{\sim} X$$

at  $X$  is given by

$$\rho_X^{\text{Sets}, \coprod}((0, x)) \stackrel{\text{def}}{=} x$$

for each  $(0, x) \in X \coprod \emptyset$ .

**PROOF 5.2.5.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.2.5.1.1**

**Unwinding the Definition of  $X \coprod \emptyset$**

Firstly, we unwind the expression for  $X \coprod \emptyset$ . We have

$$\begin{aligned} X \coprod \emptyset &\stackrel{\text{def}}{=} \{(0, x) \in S \mid x \in X\} \cup \{(1, z) \in S \mid z \in \emptyset\} \\ &= \{(0, x) \in S \mid x \in X\} \cup \emptyset \\ &= \{(0, x) \in S \mid x \in X\}, \end{aligned}$$

where  $S = \{0, 1\} \times (X \cup \emptyset) = \{0, 1\} \times (\emptyset \cup X) = S$ .

**Invertibility**

The inverse of  $\rho_X^{\text{Sets}, \coprod}$  is the map

$$\rho_X^{\text{Sets}, \coprod, -1} : X \rightarrow X \coprod \emptyset$$

given by

$$\rho_X^{\text{Sets}, \coprod, -1}(x) \stackrel{\text{def}}{=} (0, x)$$

for each  $x \in X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned} [\rho_X^{\text{Sets}, \coprod, -1} \circ \rho_X^{\text{Sets}, \coprod}] (0, x) &= \rho_X^{\text{Sets}, \coprod, -1} (\rho_X^{\text{Sets}, \coprod} (0, x)) \\ &= \rho_X^{\text{Sets}, \coprod, -1} (x) \\ &= (0, x) \\ &= [\text{id}_X \coprod_{\emptyset}] (0, x) \end{aligned}$$

for each  $(0, x) \in \emptyset \coprod X$ , and therefore we have

$$\rho_X^{\text{Sets}, \coprod, -1} \circ \rho_X^{\text{Sets}, \coprod} = \text{id}_{\emptyset \coprod X}.$$

- *Invertibility II.* We have

$$\begin{aligned} [\rho_X^{\text{Sets}, \coprod} \circ \rho_X^{\text{Sets}, \coprod, -1}] (x) &= \rho_X^{\text{Sets}, \coprod} (\rho_X^{\text{Sets}, \coprod, -1} (x)) \\ &= \rho_X^{\text{Sets}, \coprod, -1} (0, x) \\ &= x \\ &= [\text{id}_X] (x) \end{aligned}$$

for each  $x \in X$ , and therefore we have

$$\rho_X^{\text{Sets}, \coprod} \circ \rho_X^{\text{Sets}, \coprod, -1} = \text{id}_X.$$

Therefore  $\rho_X^{\text{Sets}, \coprod}$  is indeed an isomorphism.

### Naturality

We need to show that, given a function  $f: X \rightarrow Y$ , the diagram

$$\begin{array}{ccc} X \coprod \emptyset & \xrightarrow{f \coprod \text{id}_{\emptyset}} & Y \coprod \emptyset \\ \rho_X^{\text{Sets}, \coprod} \downarrow & & \downarrow \rho_Y^{\text{Sets}, \coprod} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (0, x) & \mapsto & (1, f(x)) \\ \downarrow & & \downarrow \\ x & \longmapsto & f(x) \end{array}$$

and hence indeed commutes. Therefore  $\rho^{\text{Sets}, \coprod}$  is a natural transformation.

#### Being a Natural Isomorphism

Since  $\rho^{\text{Sets}, \coprod}$  is natural and  $\rho^{\text{Sets}, -1}$  is a componentwise inverse to  $\rho^{\text{Sets}, \coprod}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\rho^{\text{Sets}, -1}$  is also natural. Thus  $\rho^{\text{Sets}, \coprod}$  is a natural isomorphism. 

### 01PW 5.2.6 The Symmetry

01PX

#### DEFINITION 5.2.6.1.1 ► THE SYMMETRY OF $\coprod$

The **symmetry of the coproduct of sets** is the natural isomorphism

$$\sigma^{\text{Sets}, \coprod}: \coprod \xrightarrow{\sim} \coprod \circ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2}, \quad \begin{array}{ccc} \text{Sets} \times \text{Sets} & \xrightarrow{\coprod} & \text{Sets}, \\ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2} \swarrow & \downarrow \sigma^{\text{Sets}, \coprod} & \searrow \coprod \\ \text{Sets} \times \text{Sets} & & \end{array}$$

whose component

$$\sigma_{X,Y}^{\text{Sets}, \coprod}: X \coprod Y \xrightarrow{\sim} Y \coprod X$$

at  $X, Y \in \text{Obj}(\text{Sets})$  is defined by

$$\sigma_{X,Y}^{\text{Sets}, \coprod}(x, y) \stackrel{\text{def}}{=} (y, x)$$

for each  $(x, y) \in X \times Y$ .

#### PROOF 5.2.6.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.2.6.1.1

##### Unwinding the Definitions of $X \coprod Y$ and $Y \coprod X$

Firstly, we unwind the expressions for  $X \coprod Y$  and  $Y \coprod X$ . We have

$$X \coprod Y \stackrel{\text{def}}{=} \{(0, x) \in S \mid x \in X\} \cup \{(1, y) \in S \mid y \in Y\},$$

where  $S = \{0, 1\} \times (X \cup Y)$  and

$$Y \coprod X \stackrel{\text{def}}{=} \{(0, y) \in S' \mid y \in Y\} \cup \{(1, x) \in S' \mid x \in X\},$$

where  $S' = \{0, 1\} \times (Y \cup X) = \{0, 1\} \times (X \cup Y) = S$ .

### Invertibility

The inverse of  $\sigma_{X,Y}^{\text{Sets}, \coprod}$  is the map

$$\sigma_{X,Y}^{\text{Sets}, \coprod, -1} : Y \coprod X \rightarrow X \coprod Y$$

defined by

$$\sigma_{X,Y}^{\text{Sets}, \coprod, -1} \stackrel{\text{def}}{=} \sigma_{Y,X}^{\text{Sets}, \coprod}$$

and hence given by

$$\sigma_{X,Y}^{\text{Sets}, \coprod, -1}(z) \stackrel{\text{def}}{=} \begin{cases} (0, x) & \text{if } z = (1, x), \\ (1, y) & \text{if } z = (0, y) \end{cases}$$

for each  $z \in Y \coprod X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned} [\sigma_{X,Y}^{\text{Sets}, \coprod, -1} \circ \sigma_{X,Y}^{\text{Sets}, \coprod}] (0, x) &= \sigma_X^{\text{Sets}, \coprod, -1} (\sigma_X^{\text{Sets}, \coprod} (0, x)) \\ &= \sigma_X^{\text{Sets}, \coprod, -1} (1, x) \\ &= (0, x) \\ &= [\text{id}_{X \coprod Y}] (0, x) \end{aligned}$$

for each  $(0, x) \in X \coprod Y$  and

$$\begin{aligned} [\sigma_{X,Y}^{\text{Sets}, \coprod, -1} \circ \sigma_{X,Y}^{\text{Sets}, \coprod}] (1, y) &= \sigma_X^{\text{Sets}, \coprod, -1} (\sigma_X^{\text{Sets}, \coprod} (1, y)) \\ &= \sigma_X^{\text{Sets}, \coprod, -1} (0, y) \\ &= (1, y) \\ &= [\text{id}_{X \coprod Y}] (1, y) \end{aligned}$$

for each  $(1, y) \in X \coprod Y$ , and therefore we have

$$\sigma_{X,Y}^{\text{Sets}, \coprod, -1} \circ \sigma_{X,Y}^{\text{Sets}, \coprod} = \text{id}_{X \coprod Y}.$$

- *Invertibility II.* We have

$$\begin{aligned} [\sigma_{X,Y}^{\text{Sets}, \coprod} \circ \sigma_{X,Y}^{\text{Sets}, \coprod, -1}] (0, y) &= \sigma_X^{\text{Sets}, \coprod} (\sigma_X^{\text{Sets}, \coprod, -1} (0, y)) \\ &= \sigma_X^{\text{Sets}, \coprod, -1} (1, y) \\ &= (0, y) \end{aligned}$$

$$= [\text{id}_{Y \coprod X}](0, y)$$

for each  $(0, y) \in Y \coprod X$  and

$$\begin{aligned} [\sigma_{X,Y}^{\text{Sets}, \coprod} \circ \sigma_{X,Y}^{\text{Sets}, \coprod, -1}](1, x) &= \sigma_X^{\text{Sets}, \coprod}(\sigma_X^{\text{Sets}, \coprod, -1}(1, x)) \\ &= \sigma_X^{\text{Sets}, \coprod, -1}(0, x) \\ &= (1, x) \\ &= [\text{id}_{Y \coprod X}](1, x) \end{aligned}$$

for each  $(1, x) \in Y \coprod X$ , and therefore we have

$$\sigma_X^{\text{Sets}, \coprod} \circ \sigma_X^{\text{Sets}, \coprod, -1} = \text{id}_{Y \coprod X}.$$

Therefore  $\sigma_{X,Y}^{\text{Sets}, \coprod}$  is indeed an isomorphism.

### Naturality

We need to show that, given functions  $f: A \rightarrow X$  and  $g: B \rightarrow Y$ , the diagram

$$\begin{array}{ccc} A \coprod B & \xrightarrow{f \coprod g} & X \coprod Y \\ \downarrow \sigma_{A,B}^{\text{Sets}, \coprod} & & \downarrow \sigma_{X,Y}^{\text{Sets}, \coprod} \\ B \coprod A & \xrightarrow{g \coprod f} & Y \coprod X \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (0, a) & & (0, a) \mapsto (0, f(a)) \\ \downarrow & & \downarrow \\ (1, a) \mapsto (1, f(a)) & & (1, f(a)) \\ \\ (1, b) & & (1, b) \mapsto (1, g(b)) \\ \downarrow & & \downarrow \\ (0, b) \mapsto (0, g(b)) & & (0, g(b)) \end{array}$$

and hence indeed commutes. Therefore  $\sigma^{\text{Sets}, \coprod}$  is a natural transformation.

**Being a Natural Isomorphism**

Since  $\sigma^{\text{Sets}, \coprod}$  is natural and  $\sigma^{\text{Sets}, -1}$  is a componentwise inverse to  $\sigma^{\text{Sets}, \coprod}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\sigma^{\text{Sets}, -1}$  is also natural. Thus  $\sigma^{\text{Sets}, \coprod}$  is a natural isomorphism. 

**01PY 5.2.7 The Monoidal Category of Sets and Coproducts**

01PZ

**PROPOSITION 5.2.7.1.1 ► THE MONOIDAL STRUCTURE ON SETS ASSOCIATED TO  $\coprod$** 

The category  $\text{Sets}$  admits a closed symmetric monoidal category structure consisting of:

- *The Underlying Category.* The category  $\text{Sets}$  of pointed sets.
- *The Monoidal Product.* The coproduct functor

$$\coprod : \text{Sets} \times \text{Sets} \rightarrow \text{Sets}$$

of Item 1 of Proposition 4.2.3.1.4.

- *The Monoidal Unit.* The functor

$$\emptyset^{\text{Sets}} : \text{pt} \rightarrow \text{Sets}$$

of Definition 5.2.2.1.1.

- *The Associators.* The natural isomorphism

$$\alpha^{\text{Sets}, \coprod} : \coprod \circ (\coprod \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \coprod \circ (\text{id}_{\text{Sets}} \times \coprod) \circ \alpha_{\text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}}$$

of Definition 5.2.3.1.1.

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\text{Sets}, \coprod} : \coprod \circ (\emptyset^{\text{Sets}} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \lambda_{\text{Sets}}^{\text{Cats}_2}$$

of Definition 5.2.4.1.1.

- *The Right Unitors.* The natural isomorphism

$$\rho^{\text{Sets}, \coprod} : \coprod \circ (\text{id} \times \emptyset^{\text{Sets}}) \xrightarrow{\sim} \rho_{\text{Sets}}^{\text{Cats}_2}$$

of Definition 5.2.5.1.1.

- *The Symmetry.* The natural isomorphism

$$\sigma^{\text{Sets}, \coprod} : \times \xrightarrow{\sim} \times \circ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2}$$

of Definition 5.2.6.1.1.

**PROOF 5.2.7.1.2 ► PROOF OF PROPOSITION 5.2.7.1.1**


The Pentagon Identity

Let  $W, X, Y$  and  $Z$  be sets. We have to show that the diagram

$$\begin{array}{ccc}
 & (W \sqcup (X \sqcup Y)) \sqcup Z & \\
 \alpha_{W,X,Y}^{\text{Sets}, \sqcup} \sqcup \text{id}_Z \swarrow & & \searrow \alpha_{W,X \sqcup Y,Z}^{\text{Sets}, \sqcup} \\
 ((W \sqcup X) \sqcup Y) \sqcup Z & & W \sqcup ((X \sqcup Y) \sqcup Z) \\
 \downarrow \alpha_{W \sqcup X,Y,Z}^{\text{Sets}, \sqcup} & & \downarrow \text{id}_W \sqcup \alpha_{X,Y,Z}^{\text{Sets}, \sqcup} \\
 (W \sqcup X) \sqcup (Y \sqcup Z) & \xrightarrow{\alpha_{W,X,Y \sqcup Z}^{\text{Sets}, \sqcup}} & W \sqcup (X \sqcup (Y \sqcup Z))
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccccc}
 & & (0, (0, w)) & & \\
 & \nearrow & & \searrow & \\
 (0, (0, (0, w))) & & (0, (0, (0, w))) & & (0, w) \\
 \downarrow & & \downarrow & & \downarrow \\
 (0, (0, w)) & \longmapsto & (0, w) & & \\
 & \nearrow & & \searrow & \\
 & & (0, (1, (0, x))) & & \\
 & \nearrow & & \searrow & \\
 (0, (0, (1, x))) & & (0, (0, (1, x))) & & (1, (0, (0, x))) \\
 \downarrow & & \downarrow & & \downarrow \\
 (0, (1, x)) & \longmapsto & (1, (0, x)) & & 
 \end{array}$$



and therefore the pentagon identity is satisfied.

### The Triangle Identity

Let  $X$  and  $Y$  be sets. We have to show that the diagram


$$(X \coprod \emptyset) \coprod Y \xrightarrow{\alpha_{X, \emptyset, Y}^{\text{Sets}, \coprod}} X \coprod (\emptyset \coprod Y)$$

$$\rho_X^{\text{Sets}, \coprod} \coprod \text{id}_Y \quad \text{id}_X \coprod \lambda_Y^{\text{Sets}, \coprod}$$

$$\downarrow \qquad \qquad \downarrow$$

$$X \coprod Y$$

commutes. Indeed, this diagram acts on elements as



$$\begin{array}{ccc}
 (1, y) & & (1, y) \xrightarrow{\quad} (1, (1, y)) \\
 \swarrow & & \searrow \\
 (1, y) & & (1, y)
 \end{array}$$

and therefore the triangle identity is satisfied.


### The Left Hexagon Identity

Let  $X$ ,  $Y$ , and  $Z$  be sets. We have to show that the diagram

$$\begin{array}{ccc}
 & (X \sqcup Y) \sqcup Z & \\
 \alpha_{X,Y,Z}^{\text{Sets}} \swarrow & & \searrow \sigma_{X,Y}^{\text{Sets}} \sqcup \text{id}_Z \\
 X \sqcup (Y \sqcup Z) & & (Y \sqcup X) \sqcup Z \\
 \downarrow \sigma_{X,Y}^{\text{Sets}} \sqcup Z & & \downarrow \alpha_{Y,X,Z}^{\text{Sets}} \\
 (Y \sqcup Z) \sqcup X & & Y \sqcup (X \sqcup Z) \\
 \alpha_{Y,Z,X}^{\text{Sets}} \searrow & & \swarrow \text{id}_Y \sqcup \sigma_{X,Z}^{\text{Sets}}
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (0, (0, x)) & & (0, (0, x)) \\
 \swarrow & & \searrow \\
 (0, x) & & (0, (1, x)) \\
 \downarrow & & \downarrow \\
 (1, x) & & (1, (0, x)) \\
 \swarrow & & \searrow \\
 (1, (1, x)) & & (1, (1, x))
 \end{array}$$




and thus the left hexagon identity is satisfied.

### The Right Hexagon Identity

Let  $X$ ,  $Y$ , and  $Z$  be sets. We have to show that the diagram

$$\begin{array}{ccc}
 & X \coprod (Y \coprod Z) & \\
 (\alpha_{X,Y,Z}^{\text{Sets}})^{-1} \swarrow & & \searrow \text{id}_X \coprod \sigma_{Y,Z}^{\text{Sets}} \\
 (X \coprod Y) \coprod Z & & X \coprod (Z \coprod Y) \\
 \downarrow \sigma_{X \coprod Y, Z}^{\text{Sets}} & & \downarrow (\alpha_{X,Z,Y}^{\text{Sets}})^{-1} \\
 Z \coprod (X \coprod Y) & & (X \coprod Z) \coprod Y \\
 \downarrow (\alpha_{Z,X,Y}^{\text{Sets}})^{-1} & & \swarrow \sigma_{X,Z}^{\text{Sets}} \coprod \text{id}_Y \\
 (Z \coprod X) \coprod Y & &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as



and thus the right hexagon identity is satisfied. 

## 01Q0 5.3 The Bimonoidal Category of Sets, Products, and Coproducts

### 01Q1 5.3.1 The Left Distributor

01Q2

#### DEFINITION 5.3.1.1.1 ► THE LEFT DISTRIBUTOR OF $\times$ OVER $\coprod$

The **left distributor** of the product of sets over the coproduct of sets is the natural isomorphism

$$\delta_{\ell}^{\text{Sets}} : \times \circ (\text{id}_{\text{Sets}} \times \coprod) \xrightarrow{\sim} \coprod \circ ((\times) \times (\times)) \circ \mu_{4|(\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets})}^{\text{Cats}_2} \circ (\Delta_{\text{Sets}} \times (\text{id}_{\text{Sets}} \times \text{id}_{\text{Sets}}))$$

as in the diagram

$$\begin{array}{ccc}
 & (\text{Sets} \times \text{Sets}) \times (\text{Sets} \times \text{Sets}) & \\
 \Delta_{\text{Sets}} \times (\text{id}_{\text{Sets}} \times \text{id}_{\text{Sets}}) & \nearrow & \searrow \mu_{4|(\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets})}^{\text{Cats}_2} \\
 \text{Sets} \times (\text{Sets} \times \text{Sets}) & & (\text{Sets} \times \text{Sets}) \times (\text{Sets} \times \text{Sets}) \\
 \downarrow \text{id}_{\text{Sets}} \times \coprod & \swarrow \delta_{\ell}^{\text{Sets}} & \downarrow (\times) \times (\times) \\
 \text{Sets} \times \text{Sets} & & \text{Sets} \times \text{Sets}, \\
 & \searrow \times & \swarrow \coprod \\
 & \text{Sets} & 
 \end{array}$$

whose component

$$\delta_{\ell|X,Y,Z}^{\text{Sets}} : X \times (Y \coprod Z) \xrightarrow{\sim} (X \times Y) \coprod (X \times Z)$$

at  $(X, Y, Z)$  is defined by

$$\delta_{\ell|X,Y,Z}^{\text{Sets}}(x, a) \stackrel{\text{def}}{=} \begin{cases} (0, (x, y)) & \text{if } a = (0, y), \\ (1, (x, z)) & \text{if } a = (1, z) \end{cases}$$

for each  $(x, a) \in X \times (Y \coprod Z)$ .

**PROOF 5.3.1.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.3.1.1****Invertibility**

The inverse of  $\delta_{\ell|X,Y,Z}^{\text{Sets}}$  is the map

$$\delta_{\ell|X,Y,Z}^{\text{Sets},-1}: (X \times Y) \coprod (X \times Z) \xrightarrow{\sim} X \times (Y \coprod Z)$$

given by

$$\delta_{\ell|X,Y,Z}^{\text{Sets},-1}(a) \stackrel{\text{def}}{=} \begin{cases} (x, (0, y)) & \text{if } a = (0, (x, y)), \\ (x, (1, z)) & \text{if } a = (1, (x, z)) \end{cases}$$

for  $a \in (X \times Y) \coprod (X \times Z)$ . Indeed:

- *Invertibility I.* The map  $\delta_{\ell|X,Y,Z}^{\text{Sets}} \circ \delta_{\ell|X,Y,Z}^{\text{Sets},-1}$  acts on elements as

$$\begin{aligned} (x, (0, y)) &\mapsto (0, (x, y)) \mapsto (x, (0, y)), \\ (x, (1, z)) &\mapsto (1, (x, z)) \mapsto (x, (1, z)), \end{aligned}$$

but these are the two possible cases for elements of  $X \times (Y \coprod Z)$ . Hence the map is equal to the identity.

- *Invertibility II.* The map  $\delta_{\ell|X,Y,Z}^{\text{Sets}} \circ \delta_{\ell|X,Y,Z}^{\text{Sets},-1}$  acts on elements as

$$\begin{aligned} (0, (x, y)) &\mapsto (x, (0, y)) \mapsto (0, (x, y)), \\ (1, (x, z)) &\mapsto (x, (1, z)) \mapsto (1, (x, z)), \end{aligned}$$

but these are the two possible cases for elements of  $(X \times Y) \coprod (X \times Z)$ . Hence the map is equal to the identity.

Thus  $\delta_{\ell|X,Y,Z}^{\text{Sets}}$  is an isomorphism for all  $X, Y, Z$ .

**Naturality**

We need to show that, given functions

$$\begin{aligned} f: X &\rightarrow X', \\ g: Y &\rightarrow Y', \\ h: Z &\rightarrow Z' \end{aligned}$$

the diagram

$$\begin{array}{ccc} X \times (Y \coprod Z) & \xrightarrow{f \times (g \coprod h)} & X' \times (Y' \coprod Z') \\ \delta_{\ell|X,Y,Z}^{\text{Sets}} \downarrow & & \downarrow \delta_{\ell|X',Y',Z'}^{\text{Sets}} \\ (X \times Y) \coprod (X \times Z) & \xrightarrow{(f \times g) \coprod (f \times h)} & (X' \times Y') \coprod (X' \times Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (x, (0, y)) & & (x, (0, y)) \mapsto (f(x), (0, f(y))) \\
 \downarrow & & \downarrow \\
 (0, (x, y)) \mapsto (0, (f(x), g(y))) & & (0, (f(x), g(y)))
 \end{array}$$
  

$$\begin{array}{ccc}
 (x, (1, z)) & & (x, (1, z)) \mapsto (f(x), (1, h(z))) \\
 \downarrow & & \downarrow \\
 (1, (x, z)) \mapsto (1, (f(x), h(z))) & & (1, (f(x), h(z))),
 \end{array}$$

so it commutes, showing  $\delta_\ell^{\text{Sets}}$  to be a natural transformation.

#### Being a Natural Isomorphism

Since  $\delta_\ell^{\text{Sets}}$  is natural and  $\delta_\ell^{\text{Sets}, -1}$  is a componentwise inverse to  $\delta_\ell^{\text{Sets}}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\delta_\ell^{\text{Sets}, -1}$  is also natural. Thus  $\delta_\ell^{\text{Sets}}$  is a natural isomorphism. 

### 01Q3 5.3.2 The Right Distributor

#### 01Q4 DEFINITION 5.3.2.1.1 ► THE RIGHT DISTRIBUTOR OF $\times$ OVER $\coprod$

The **right distributor of the product of sets over the coproduct of sets** is the natural isomorphism

$$\delta_r^{\text{Sets}} : \times \circ (\coprod \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \coprod \circ ((\times) \times (\times)) \circ \mu_{4|(\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets})}^{\text{Cats}_2} \circ ((\text{id}_{\text{Sets}} \times \text{id}_{\text{Sets}}) \times \Delta_{\text{Sets}})$$

as in the diagram

$$\begin{array}{ccccc}
 & & (\text{Sets} \times \text{Sets}) \times (\text{Sets} \times \text{Sets}) & & \\
 & \swarrow & & \searrow & \\
 (\text{id}_{\text{Sets}} \times \text{id}_{\text{Sets}}) \times \Delta_{\text{Sets}} & & & \mu_{4|\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} & \\
 & & (\text{Sets} \times \text{Sets}) \times \text{Sets} & & (\text{Sets} \times \text{Sets}) \times (\text{Sets} \times \text{Sets}) \\
 & \downarrow & & & \downarrow (\times) \times (\times) \\
 \coprod \times \text{id}_{\text{Sets}} & & \delta_r^{\text{Sets}} & & \\
 & \searrow & & \nearrow & \\
 \text{Sets} \times \text{Sets} & & & & \text{Sets} \times \text{Sets}, \\
 & & \times & & \coprod \\
 & & \searrow & \nearrow & \\
 & & \text{Sets} & &
 \end{array}$$

whose component

$$\delta_{r|X,Y,Z}^{\text{Sets}} : (X \coprod Y) \times Z \xrightarrow{\sim} (X \times Z) \coprod (Y \times Z)$$

at  $(X, Y, Z)$  is defined by

$$\delta_{r|X,Y,Z}^{\text{Sets}}(a, z) \stackrel{\text{def}}{=} \begin{cases} (0, (x, z)) & \text{if } a = (0, x), \\ (1, (y, z)) & \text{if } a = (1, y) \end{cases}$$

for each  $(a, z) \in (X \coprod Y) \times Z$ .

#### PROOF 5.3.2.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.3.2.1.1

##### Invertibility

The inverse of  $\delta_{r|X,Y,Z}^{\text{Sets}}$  is the map

$$\delta_{r|X,Y,Z}^{\text{Sets}, -1} : (X \times Z) \coprod (Y \times Z) \xrightarrow{\sim} (X \coprod Y) \times Z$$

given by

$$\delta_{r|X,Y,Z}^{\text{Sets}, -1}(a) \stackrel{\text{def}}{=} \begin{cases} ((0, x), z) & \text{if } a = (0, (x, z)), \\ ((1, y), z) & \text{if } a = (1, (y, z)) \end{cases}$$

for  $a \in (X \times Z) \coprod (Y \times Z)$ . Indeed:

- *Invertibility I.* The map  $\delta_{r|X,Y,Z}^{\text{Sets}, -1} \circ \delta_{r|X,Y,Z}^{\text{Sets}}$  acts on elements as

$$\begin{aligned} ((0, x), z) &\mapsto (0, (x, z)) \mapsto (0, (x, z)), \\ ((1, y), z) &\mapsto (1, (y, z)) \mapsto (1, (y, z)), \end{aligned}$$

but these are the two possible cases for elements of  $(X \coprod Y) \times Z$ . Hence the map is equal to the identity.

- *Invertibility II.* The map  $\delta_{r|X,Y,Z}^{\text{Sets}} \circ \delta_{r|X,Y,Z}^{\text{Sets}, -1}$  acts on elements as

$$\begin{aligned} (0, (x, z)) &\mapsto ((0, x), z) \mapsto (0, (x, z)), \\ (1, (y, z)) &\mapsto ((1, y), z) \mapsto (1, (y, z)), \end{aligned}$$

but these are the two possible cases for elements of  $(X \times Z) \coprod (Y \times Z)$ . Hence the map is equal to the identity.

So  $\delta_{r|X,Y,Z}^{\text{Sets}}$  is an isomorphism for all  $X, Y, Z$ .

### Naturality

We need to show that, given functions

$$\begin{aligned} f: X &\rightarrow X', \\ g: Y &\rightarrow Y', \\ h: Z &\rightarrow Z' \end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \coprod Y) \times Z' & \xrightarrow{(f \coprod g) \times h} & (X' \coprod Y') \times Z' \\ \delta_{r|X,Y,Z}^{\text{Sets}} \downarrow & & \downarrow \delta_{r|X',Y',Z'}^{\text{Sets}} \\ (X \times Z) \coprod (Y \times Z) & \xrightarrow{(f \times h) \coprod (g \times h)} & (X' \times Z') \coprod (Y' \times Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} ((0, x), z) & & ((0, x), z) \mapsto ((0, f(x)), h(z)) \\ \downarrow & & \downarrow \\ (0, (x, z)) & \mapsto (0, (f(x), h(z))) & (0, (f(x), h(z))) \end{array}$$

$$\begin{array}{ccc}
 ((1, y), z) & & ((1, y), z) \mapsto ((1, g(y)), h(z)) \\
 \downarrow & & \downarrow \\
 (1, (y, z)) \mapsto (1, (g(y), h(z))) & & (1, (g(y), h(z)))
 \end{array}$$

so it commutes and  $\delta_r^{\text{Sets}}$  is a natural transformation.

Being a Natural Isomorphism

Since  $\delta_r^{\text{Sets}}$  is natural and  $\delta_r^{\text{Sets}, -1}$  is a componentwise inverse to  $\delta_r^{\text{Sets}}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\delta_r^{\text{Sets}, -1}$  is also natural. Thus  $\delta_r^{\text{Sets}}$  is a natural isomorphism. 

### 01Q5 5.3.3 The Left Annihilator

#### 01Q6 DEFINITION 5.3.3.1.1 ► THE LEFT ANNIHILATOR OF $\times$

The **left annihilator** of the product of sets is the natural isomorphism

$$\zeta_{\ell}^{\text{Sets}} : \emptyset^{\text{Sets}} \circ \mu_{4|(\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets})}^{\text{Cats}_2} \circ (\text{id}_{\text{pt}} \times \epsilon_{\text{Sets}}^{\text{Cats}_2}) \xrightarrow{\sim} \times \circ (\emptyset^{\text{Sets}} \times \text{id}_{\text{Sets}})$$

as in the diagram

$$\begin{array}{ccccc}
 & & \text{pt} \times \text{pt} & & \\
 & \nearrow \text{id}_{\text{pt}} \times \epsilon_{\text{Sets}}^{\text{Cats}_2} & & \searrow \mu_{4|(\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets})}^{\text{Cats}_2} & \\
 \text{pt} \times \text{Sets} & & & & \text{pt} \\
 & \swarrow \emptyset^{\text{Sets}} \times \text{id}_{\text{Sets}} & \Downarrow \zeta_{\ell}^{\text{Sets}} & \searrow \emptyset^{\text{Sets}} & \\
 \text{Sets} \times \text{Sets} & \xrightarrow{\times} & \text{Sets} & &
 \end{array}$$

with components

$$\zeta_{\ell|A}^{\text{Sets}} : \emptyset \times A \xrightarrow{\sim} \emptyset$$

given by  $\zeta_{\ell|A}^{\text{Sets}} \stackrel{\text{def}}{=} \text{pr}_1$ .

**PROOF 5.3.3.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.3.3.1.1****Invertibility**

The inverse of  $\zeta_{\ell|A}^{\text{Sets}}$  is the map

$$\zeta_{\ell|A}^{\text{Sets}, -1}: \emptyset \xrightarrow{\sim} \emptyset \times A$$

given by

$$\zeta_{\ell|A}^{\text{Sets}, -1} \stackrel{\text{def}}{=} \iota_A,$$

where  $\iota_A$  is as defined in [Construction 4.2.1.1.2](#):

- *Invertibility I.* The map  $\zeta_{\ell|A}^{\text{Sets}} \circ \iota_A: \emptyset \rightarrow \emptyset$  is equal to  $\text{id}_{\emptyset}$ , as  $\emptyset$  is the initial object of Sets.
- *Invertibility II.* The map  $\iota_A \circ \zeta_{\ell|A}^{\text{Sets}}$  is equal to the identity on every  $(x, a) \in \emptyset \times A$ , of which there are none.

Hence  $\zeta_{\ell|A}^{\text{Sets}}$  is an isomorphism.

**Naturality**

We need to show that given a function  $f: A \rightarrow B$ , the diagram

$$\begin{array}{ccc} \emptyset \times A & \xrightarrow{\text{id}_{\emptyset} \times f} & \emptyset \times B \\ \zeta_{\ell|A}^{\text{Sets}} \downarrow & & \downarrow \zeta_{\ell|B}^{\text{Sets}} \\ \emptyset & \xrightarrow{\text{id}_{\emptyset}} & \emptyset \end{array}$$

commutes. But since  $\emptyset \times A$  has no elements, this is trivially true.

**Being a Natural Isomorphism**

Since  $\zeta_{\ell}^{\text{Sets}}$  is natural and  $\zeta_{\ell}^{\text{Sets}, -1}$  is a componentwise inverse to  $\zeta_{\ell}^{\text{Sets}}$ , it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\zeta_{\ell}^{\text{Sets}, -1}$  is also natural. Thus  $\zeta_{\ell}^{\text{Sets}}$  is a natural isomorphism. 

**01Q7 5.3.4 The Right Annihilator**

01Q8

**DEFINITION 5.3.4.1.1 ► THE RIGHT ANNIHILATOR OF ×**

The **right annihilator of the product of sets** is the natural isomorphism

$$\zeta_r^{\text{Sets}}: \mathbb{O}^{\text{Sets}} \circ \mu_{4|\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} \circ (\epsilon_{\text{Sets}}^{\text{Cats}_2} \times \text{id}_{\text{pt}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Sets}} \times \mathbb{O}^{\text{Sets}})$$

as in the diagram

$$\begin{array}{ccccc}
 & & \text{pt} \times \text{pt} & & \\
 & \swarrow \epsilon_{\text{Sets}}^{\text{Cats}_2 \times \text{id}_{\text{pt}}} & & \searrow \mu_{4|\text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} & \\
 \text{Sets} \times \text{pt} & & & & \text{pt} \\
 & \searrow \text{id}_{\text{Sets}} \times \emptyset^{\text{Sets}} & \swarrow \zeta_r^{\text{Sets}} & & \downarrow \emptyset^{\text{Sets}} \\
 & & \text{Sets} \times \text{Sets} & \xrightarrow{x} & \text{Sets}
 \end{array}$$

with components

$$\zeta_{r|A}^{\text{Sets}} : A \times \emptyset \xrightarrow{\sim} \emptyset$$

given by  $\zeta_{r|A}^{\text{Sets}} \stackrel{\text{def}}{=} \text{pr}_2$ .

#### PROOF 5.3.4.1.2 ► PROOF OF THE CLAIMS MADE IN DEFINITION 5.3.4.1.1

##### Invertibility

The inverse of  $\zeta_{r|A}^{\text{Sets}}$  is the map

$$\zeta_{r|A}^{\text{Sets}, -1} : \emptyset \xrightarrow{\sim} A \times \emptyset$$

given by

$$\zeta_{r|A}^{\text{Sets}, -1} \stackrel{\text{def}}{=} \iota_A,$$

where  $\iota_A$  is as defined in [Construction 4.2.1.1.2](#):

- *Invertibility I.* The map  $\zeta_{r|A}^{\text{Sets}} \circ \iota_A : \emptyset \rightarrow \emptyset$  is equal to  $\text{id}_{\emptyset}$ , as  $\emptyset$  is the initial object of  $\text{Sets}$ .
- *Invertibility II.* The map  $\iota_A \circ \zeta_{r|A}^{\text{Sets}}$  is equal to the identity on every  $(a, x) \in A \times \emptyset$ , of which there are none.

Hence  $\zeta_{r|A}^{\text{Sets}}$  is an isomorphism.

##### Naturality

We need to show that given a function  $f: A \rightarrow B$ , the diagram

$$\begin{array}{ccc} A \times \emptyset & \xrightarrow{f \times \text{id}_\emptyset} & B \times \emptyset \\ \zeta_{r|A}^{\text{Sets}} \downarrow & & \downarrow \zeta_{r|B}^{\text{Sets}} \\ \emptyset & \xrightarrow{\text{id}_\emptyset} & \emptyset \end{array}$$

commutes. But since  $A \times \emptyset$  has no elements, this is trivially true.

Being a Natural Isomorphism

Since  $\zeta_r^{\text{Sets}}$  is natural and  $\zeta_r^{\text{Sets}, -1}$  is a componentwise inverse to  $\zeta_r^{\text{Sets}}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\zeta_r^{\text{Sets}, -1}$  is also natural. Thus  $\zeta_r^{\text{Sets}}$  is a natural isomorphism. □

### 01Q9 5.3.5 The Bimonoidal Category of Sets, Products, and Coproducts

01QA

#### PROPOSITION 5.3.5.1.1 ► THE BIMONOIDAL STRUCTURE ON SETS ASSOCIATED TO $\times$ AND $\coprod$

The category Sets admits a closed symmetric bimonoidal category structure consisting of:

- *The Underlying Category.* The category Sets of pointed sets.
- *The Additive Monoidal Product.* The coproduct functor

$$\coprod: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}$$

of Item 1 of Proposition 4.2.3.1.4.

- *The Multiplicative Monoidal Product.* The product functor

$$\times: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}$$

of Item 1 of Proposition 4.1.3.1.4.

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\text{Sets}}: \text{pt} \rightarrow \text{Sets}$$

of Definition 5.1.3.1.1.

- *The Monoidal Zero.* The functor

$$\emptyset^{\text{Sets}}: \text{pt} \rightarrow \text{Sets}$$

of Definition 5.1.3.1.1.

- *The Internal Hom.* The internal Hom functor

$$\text{Sets}: \text{Sets}^{\text{op}} \times \text{Sets} \rightarrow \text{Sets}$$

of ?? of ??.

- *The Additive Associators.* The natural isomorphism

$$\alpha^{\text{Sets}, \coprod}: \coprod \circ (\coprod \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \coprod \circ (\text{id}_{\text{Sets}} \times \coprod) \circ \alpha_{\text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}}$$

of [Definition 5.2.3.1.1](#).

- *The Additive Left Unitors.* The natural isomorphism

$$\lambda^{\text{Sets}, \coprod}: \coprod \circ (\emptyset^{\text{Sets}} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \lambda_{\text{Sets}}^{\text{Cats}_2}$$

of [Definition 5.2.4.1.1](#).

- *The Additive Right Unitors.* The natural isomorphism

$$\rho^{\text{Sets}, \coprod}: \coprod \circ (\text{id} \times \emptyset^{\text{Sets}}) \xrightarrow{\sim} \rho_{\text{Sets}}^{\text{Cats}_2}$$

of [Definition 5.2.5.1.1](#).

- *The Additive Symmetry.* The natural isomorphism

$$\sigma^{\text{Sets}, \coprod}: \coprod \xrightarrow{\sim} \coprod \circ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2}$$

of [Definition 5.2.6.1.1](#).

- *The Multiplicative Associators.* The natural isomorphism

$$\alpha^{\text{Sets}}: \times \circ (\times \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Sets}} \times \times) \circ \alpha_{\text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}}$$

of [Definition 5.1.4.1.1](#).

- *The Multiplicative Left Unitors.* The natural isomorphism

$$\lambda^{\text{Sets}}: \times \circ (\mathbb{1}^{\text{Sets}} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \lambda_{\text{Sets}}^{\text{Cats}_2}$$

of [Definition 5.1.5.1.1](#).

- *The Multiplicative Right Unitors.* The natural isomorphism

$$\rho^{\text{Sets}}: \times \circ (\text{id} \times \mathbb{1}^{\text{Sets}}) \xrightarrow{\sim} \rho_{\text{Sets}}^{\text{Cats}_2}$$

of [Definition 5.1.6.1.1](#).

- *The Multiplicative Symmetry.* The natural isomorphism

$$\sigma^{\text{Sets}} : \times \xrightarrow{\sim} \times \circ \sigma_{\text{Sets}, \text{Sets}}^{\text{Cats}_2}$$

of [Definition 5.1.7.1.1](#).

- *The Left Distributor.* The natural isomorphism

$$\delta_\ell^{\text{Sets}} : \times \circ (\text{id}_{\text{Sets}} \times \text{I}) \xrightarrow{\sim} \text{I} \circ ((\times) \times (\times)) \circ \mu_{4| \text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} \circ (\Delta_{\text{Sets}} \times (\text{id}_{\text{Sets}} \times \text{id}_{\text{Sets}}))$$

of [Definition 5.3.1.1.1](#).

- *The Right Distributor.* The natural isomorphism

$$\delta_r^{\text{Sets}} : \times \circ (\text{I} \times \text{id}_{\text{Sets}}) \xrightarrow{\sim} \text{I} \circ ((\times) \times (\times)) \circ \mu_{4| \text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} \circ ((\text{id}_{\text{Sets}} \times \text{id}_{\text{Sets}}) \times \Delta_{\text{Sets}})$$

of [Definition 5.3.2.1.1](#).

- *The Left Annihilator.* The natural isomorphism

$$\zeta_\ell^{\text{Sets}} : \emptyset^{\text{Sets}} \circ \mu_{4| \text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} \circ (\text{id}_{\text{pt}} \times \epsilon_{\text{Sets}}^{\text{Cats}_2}) \xrightarrow{\sim} \times \circ (\emptyset^{\text{Sets}} \times \text{id}_{\text{Sets}})$$

of [Definition 5.3.3.1.1](#).

- *The Right Annihilator.* The natural isomorphism

$$\zeta_r^{\text{Sets}} : \emptyset^{\text{Sets}} \circ \mu_{4| \text{Sets}, \text{Sets}, \text{Sets}, \text{Sets}}^{\text{Cats}_2} \circ (\epsilon_{\text{Sets}}^{\text{Cats}_2} \times \text{id}_{\text{pt}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Sets}} \times \emptyset^{\text{Sets}})$$

of [Definition 5.3.4.1.1](#).

**PROOF 5.3.5.1.2 ► PROOF OF PROPOSITION 5.3.5.1.1**

Omitted. 

# Appendices

## 5.A Other Chapters

### Preliminaries

1. Introduction
2. A Guide to the Literature

### Sets

3. Sets

### 4. Constructions With Sets

5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

### Relations

8. Relations

9. Constructions With Relations

10. Conditions on Relations

**Categories**

11. Categories

12. Presheaves and the Yoneda Lemma

**Monoidal Categories**

13. Constructions With Monoidal Categories

**Bicategories**

14. Types of Morphisms in Bicategories

**Extra Part**

15. Notes

# Chapter 6

## Pointed Sets

**02H3** This chapter contains some foundational material on pointed sets.

### Contents

---

<b>6.1</b>	<b>Pointed Sets</b>	.....	<b>331</b>
6.1.1	Foundations	.....	331
6.1.2	Morphisms of Pointed Sets	.....	333
6.1.3	The Category of Pointed Sets	.....	333
6.1.4	Elementary Properties of Pointed Sets	.....	334
6.1.5	Active and Inert Morphisms of Pointed Sets	.....	337
<b>6.2</b>	<b>Limits of Pointed Sets</b>	.....	<b>341</b>
6.2.1	The Terminal Pointed Set	.....	341
6.2.2	Products of Families of Pointed Sets	.....	341
6.2.3	Products	.....	343
6.2.4	Pullbacks	.....	346
6.2.5	Equalisers	.....	351
<b>6.3</b>	<b>Colimits of Pointed Sets</b>	.....	<b>354</b>
6.3.1	The Initial Pointed Set	.....	354
6.3.2	Coproducts of Families of Pointed Sets	.....	354
6.3.3	Coproducts	.....	357
6.3.4	Pushouts	.....	360
6.3.5	Coequalisers	.....	366
<b>6.4</b>	<b>Constructions With Pointed Sets</b>	.....	<b>368</b>
6.4.1	Free Pointed Sets	.....	368
6.4.2	Deleting Basepoints	.....	376
<b>6.A</b>	<b>Other Chapters</b>	.....	<b>383</b>

---

**0099 6.1 Pointed Sets**

**009A 6.1.1 Foundations**

009B

**DEFINITION 6.1.1.1 ► POINTED SETS**

A **pointed set**<sup>1</sup> is equivalently:

- An  $\mathbb{E}_0$ -monoid in  $(N_\bullet(\text{Sets}), \text{pt})$ .
- A pointed object in  $(\text{Sets}, \text{pt})$ .

<sup>1</sup>*Further Terminology:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, pointed sets are viewed as  $\mathbb{F}_1$ -**modules**.

009C

**REMARK 6.1.1.2 ► UNWINDING DEFINITION 6.1.1.1**

In detail, a **pointed set** is a pair  $(X, x_0)$  consisting of:

- *The Underlying Set.* A set  $X$ , called the **underlying set of**  $(X, x_0)$ .
- *The Basepoint.* A morphism

$$[x_0] : \text{pt} \rightarrow X$$

in  $\text{Sets}$ , determining an element  $x_0 \in X$ , called the **basepoint of**  $X$ .

009D

**EXAMPLE 6.1.1.3 ► THE ZERO SPHERE**

The **0-sphere**<sup>1</sup> is the pointed set  $(S^0, 0)$ <sup>2</sup> consisting of:

- *The Underlying Set.* The set  $S^0$  defined by

$$S^0 \stackrel{\text{def}}{=} \{0, 1\}.$$

- *The Basepoint.* The element 0 of  $S^0$ .

<sup>1</sup>*Further Terminology:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, the 0-sphere is viewed as the **underlying pointed set of the field with one element**.

<sup>2</sup>*Further Notation:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras,  $S^0$  is also denoted  $(\mathbb{F}_1, 0)$ .

009E

**EXAMPLE 6.1.1.4 ► THE TRIVIAL POINTED SET**

The **trivial pointed set** is the pointed set  $(\text{pt}, \star)$  consisting of:

- *The Underlying Set.* The punctual set  $\text{pt} \stackrel{\text{def}}{=} \{\star\}$ .
- *The Basepoint.* The element  $\star$  of  $\text{pt}$ .

01QB

The **standard pointed set with  $n + 1$  elements** is the pointed set  $\langle n \rangle$  consisting of

- *The Underlying Set.* The set  $\langle n \rangle$  defined by

$$\langle n \rangle \stackrel{\text{def}}{=} \{*\} \cup \{1, \dots, n\}.$$

- *The Basepoint.* The element  $*$  of  $\langle n \rangle$ .

### 009H 6.1.2 Morphisms of Pointed Sets

009J

#### DEFINITION 6.1.2.1.1 ► MORPHISMS OF POINTED SETS

A **morphism of pointed sets**<sup>1,2</sup> is equivalently:

- A morphism of  $\mathbb{E}_0$ -monoids in  $(N_\bullet(\text{Sets}), \text{pt})$ .
- A morphism of pointed objects in  $(\text{Sets}, \text{pt})$ .

<sup>1</sup>Further Terminology: Also called a **pointed function**.

<sup>2</sup>Further Terminology: In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, morphisms of pointed sets are also called **morphism of  $\mathbb{F}_1$ -modules**.

009K

#### REMARK 6.1.2.1.2 ► UNWINDING DEFINITION 6.1.2.1.1

In detail, a **morphism of pointed sets**  $f: (X, x_0) \rightarrow (Y, y_0)$  is a morphism of sets  $f: X \rightarrow Y$  such that the diagram

$$\begin{array}{ccc} & \text{pt} & \\ [x_0] \swarrow & & \searrow [y_0] \\ X & \xrightarrow{f} & Y \end{array}$$

commutes, i.e. such that

$$f(x_0) = y_0.$$

### 009L 6.1.3 The Category of Pointed Sets

009M

#### DEFINITION 6.1.3.1.1 ► THE CATEGORY OF POINTED SETS

The **category of pointed sets** is the category  $\text{Sets}_*$  defined equivalently as:

- The homotopy category of the  $\infty$ -category  $\text{Mon}_{\mathbb{E}_0}(N_\bullet(\text{Sets}), \text{pt})$  of ??.

- The category  $\text{Sets}_*$  of ??.

009N

**REMARK 6.1.3.1.2 ► UNWINDING DEFINITION 6.1.3.1.1**

In detail, the **category of pointed sets** is the category  $\text{Sets}_*$  where:

- *Objects.* The objects of  $\text{Sets}_*$  are pointed sets.
- *Morphisms.* The morphisms of  $\text{Sets}_*$  are morphisms of pointed sets.
- *Identities.* For each  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ , the unit map

$$\mathbb{1}_{(X, x_0)}^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*((X, x_0), (X, x_0))$$

of  $\text{Sets}_*$  at  $(X, x_0)$  is defined by<sup>1</sup>

$$\text{id}_{(X, x_0)}^{\text{Sets}_*} \stackrel{\text{def}}{=} \text{id}_X .$$

- *Composition.* For each  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ , the composition map

$$\circ_{(X, x_0), (Y, y_0), (Z, z_0)}^{\text{Sets}_*} : \text{Sets}_*((Y, y_0), (Z, z_0)) \times \text{Sets}_*((X, x_0), (Y, y_0)) \rightarrow \text{Sets}_*((X, x_0), (Z, z_0))$$

of  $\text{Sets}_*$  at  $((X, x_0), (Y, y_0), (Z, z_0))$  is defined by<sup>2</sup>

$$g \circ_{(X, x_0), (Y, y_0), (Z, z_0)}^{\text{Sets}_*} f \stackrel{\text{def}}{=} g \circ f .$$

<sup>1</sup>Note that  $\text{id}_X$  is indeed a morphism of pointed sets, as we have  $\text{id}_X(x_0) = x_0$ .

<sup>2</sup>Note that the composition of two morphisms of pointed sets is indeed a morphism of pointed sets, as we have

$$g(f(x_0)) = g(y_0) \\ = z_0, \quad \begin{array}{ccccc} & & \text{pt} & & \\ & [x_0] & \swarrow & \downarrow & \searrow [z_0] \\ X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \end{array}$$

009P 6.1.4 Elementary Properties of Pointed Sets

009Q

**PROPOSITION 6.1.4.1.1 ► ELEMENTARY PROPERTIES OF POINTED SETS**

Let  $(X, x_0)$  be a pointed set.

009R

1. *Completeness.* The category  $\text{Sets}_*$  of pointed sets and morphisms between them is complete, having in particular:

009S	(a) Products, described as in <a href="#">Definition 6.2.3.1.1</a> .
009T	(b) Pullbacks, described as in <a href="#">Definition 6.2.4.1.1</a> .
009U	(c) Equalisers, described as in <a href="#">Definition 6.2.5.1.1</a> .
009V	2. <i>Cocompleteness.</i> The category $\mathbf{Sets}_*$ of pointed sets and morphisms between them is cocomplete, having in particular:
009W	(a) Coproducts, described as in <a href="#">Definition 6.3.3.1.1</a> .
009X	(b) Pushouts, described as in <a href="#">Definition 6.3.4.1.1</a> ;
009Y	(c) Coequalisers, described as in <a href="#">Definition 6.3.5.1.1</a> .
009Z	3. <i>Failure To Be Cartesian Closed.</i> The category $\mathbf{Sets}_*$ is not Cartesian closed. <sup>1</sup>
00A0	4. <i>Morphisms From the Monoidal Unit.</i> We have a bijection of sets <sup>2</sup>
	$\mathbf{Sets}_*(S^0, X) \cong X,$
	natural in $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$ , internalising also to an isomorphism of pointed sets
	$\mathbf{Sets}_*(S^0, X) \cong (X, x_0),$
	again natural in $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$ .
00A1	5. <i>Relation to Partial Functions.</i> We have an equivalence of categories <sup>3</sup>
	$\mathbf{Sets}_* \xrightarrow{\text{ed.}} \mathbf{Sets}^{\text{part.}}$
	between the category of pointed sets and pointed functions between them and the category of sets and partial functions between them, where:
024U	(a) <i>From Pointed Sets to Sets With Partial Functions.</i> The equivalence
	$\xi: \mathbf{Sets}_* \xrightarrow{\cong} \mathbf{Sets}^{\text{part.}}$
	sends:
024V	i. A pointed set $(X, x_0)$ to $X$ .
024W	ii. A pointed function
	$f: (X, x_0) \rightarrow (Y, y_0)$
	to the partial function
	$\xi_f: X \rightarrow Y$
	defined on $f^{-1}(Y \setminus y_0)$ and given by
	$\xi_f(x) \stackrel{\text{def}}{=} f(x)$
	for each $x \in f^{-1}(Y \setminus y_0)$ .

024X

(b) *From Sets With Partial Functions to Pointed Sets.* The equivalence

$$\xi^{-1} : \text{Sets}^{\text{part.}} \xrightarrow{\cong} \text{Sets}_*$$

sends:

024Y

- i. A set  $X$  is to the pointed set  $(X, \star)$  with  $\star$  an element that is not in  $X$ .

024Z

- ii. A partial function

$$f : X \rightarrow Y$$

defined on  $U \subset X$  to the pointed function

$$\xi_f^{-1} : (X, x_0) \rightarrow (Y, y_0)$$

defined by

$$\xi_f(x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in U, \\ y_0 & \text{otherwise.} \end{cases}$$

for each  $x \in X$ .

<sup>1</sup>The category  $\text{Sets}_*$  does admit a natural monoidal closed structure, however; see Tensor Products of Pointed Sets.

<sup>2</sup>In other words, the forgetful functor

$$\text{忘} : \text{Sets}_* \rightarrow \text{Sets}$$

defined on objects by sending a pointed set to its underlying set is corepresentable by  $S^0$ .

<sup>3</sup>Warning: This is not an isomorphism of categories, only an equivalence.

#### PROOF 6.1.4.1.2 ► PROOF OF PROPOSITION 6.1.4.1.1

Item 1: Completeness

This follows from (the proofs) of Definitions 6.2.3.1.1, 6.2.4.1.1 and 6.2.5.1.1 and ??.

Item 2: Cocompleteness

This follows from (the proofs) of Definitions 6.3.3.1.1, 6.3.4.1.1 and 6.3.5.1.1 and ??.

Item 3: Failure To Be Cartesian Closed

See [MSE 2855868].

## Item 4: Morphisms From the Monoidal Unit

Since a morphism from  $S^0$  to a pointed set  $(X, x_0)$  sends  $0 \in S^0$  to  $x_0$  and then can send  $1 \in S^0$  to any element of  $X$ , we obtain a bijection between pointed maps  $S^0 \rightarrow X$  and the elements of  $X$ .

The isomorphism then

$$\mathbf{Sets}_*(S^0, X) \cong (X, x_0)$$

follows by noting that  $\Delta_{x_0} : S^0 \rightarrow X$ , the basepoint of  $\mathbf{Sets}_*(S^0, X)$ , corresponds to the pointed map  $S^0 \rightarrow X$  picking the element  $x_0$  of  $X$ , and thus we see that the bijection between pointed maps  $S^0 \rightarrow X$  and elements of  $X$  is compatible with basepoints, lifting to an isomorphism of pointed sets.

## Item 5: Relation to Partial Functions

See [MSE 884460].



## 01QC 6.1.5 Active and Inert Morphisms of Pointed Sets

01QD

## DEFINITION 6.1.5.1.1 ► ACTIVE AND INERT MORPHISMS OF POINTED SETS

Let  $f : (X, x_0) \rightarrow (Y, y_0)$  be a morphism of pointed sets.

01QE

1. The morphism  $f$  is **active** if  $f^{-1}(y_0) = x_0$ .

01QF

2. The morphism  $f$  is **inert** if, for each  $y \in Y$ , the set  $f^{-1}(y)$  has exactly one element.

01QG

## NOTATION 6.1.5.1.2 ► THE CATEGORY OF POINTED SETS AND ACTIVE MORPHISMS

We write  $\mathbf{Sets}_*^{\text{actv}}$  for the wide subcategory of  $\mathbf{Sets}_*$  spanned by pointed sets and the active maps between them.


01QH

## EXAMPLE 6.1.5.1.3 ► EXAMPLES OF ACTIVE AND INERT MAPS OF POINTED SETS

Here are some examples of active and inert maps of pointed sets.

01QJ

1. The map  $\mu : \langle 2 \rangle \rightarrow \langle 1 \rangle$  given by



is active but not inert.

01QK

2. The map  $f: \langle 2 \rangle \rightarrow \langle 2 \rangle$  given by

$$\begin{array}{ccc} 1 & \xlongleftarrow{\quad} & 1 \\ 2 & \swarrow & 2 \\ * & \xrightarrow{\quad} & * \end{array}$$

is inert but not active.

01QL

3. The map  $f: \langle 3 \rangle \rightarrow \langle 1 \rangle$  given by

$$\begin{array}{ccc} 1 & \xrightarrow{\quad} & 1 \\ 2 & \swarrow & \\ 3 & \swarrow & \\ * & \xrightarrow{\quad} & * \end{array}$$

is neither inert nor active. However, it factors as  $f = a \circ i$ , where

$$\begin{aligned} i: \langle 3 \rangle &\rightarrow \langle 2 \rangle, \\ a: \langle 2 \rangle &\rightarrow \langle 1 \rangle \end{aligned}$$

are the morphisms of pointed sets given by

$$\begin{array}{ccc} 1 & \xlongleftarrow{\quad} & 1 & 1 & \xrightarrow{\quad} & 1 \\ 2 & \xlongleftarrow{\quad} & 2 & 2 & \swarrow & \\ 3 & \swarrow & & * & \xlongleftarrow{\quad} & * \\ * & \xrightarrow{\quad} & * & & & \end{array}$$

with  $i$  being inert and  $a$  being active.

01QM

#### PROPOSITION 6.1.5.1.4 ► PROPERTIES OF ACTIVE AND INERT MAPS OF POINTED SETS

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

01QN

1. *Active-Inert Factorisation.* Every morphism of pointed sets  $f: (X, x_0) \rightarrow (Y, y_0)$  factors uniquely as

$$f = a \circ i,$$

where:

01QP

- (a) The map  $i: (X, x_0) \rightarrow (K, k_0)$  is an inert morphism of pointed sets

01QQ

- (b) The map  $a: (K, k_0) \rightarrow (Y, y_0)$  is an active morphism of pointed sets.

Moreover, this determines an orthogonal factorisation system in  $\text{Sets}_*$ .

#### PROOF 6.1.5.1.5 ► PROOF OF PROPOSITION 6.1.5.1.4

##### Item 1: Active-Inert Factorisation

Let  $f: X \rightarrow Y$  be a morphism of pointed sets. We can factor  $f$  as

$$X \xrightarrow{i} K \xrightarrow{a} Y,$$

where:

- $K$  is the pointed set given by

$$\begin{aligned} K &= \{x \in X \mid f(x) \neq y_0\} \cup \{x_0\} \\ &= (X \setminus f^{-1}(y_0)) \cup \{x_0\}; \end{aligned}$$

- $i: X \rightarrow K$  is the inert morphism of pointed sets given by

$$i(x) \stackrel{\text{def}}{=} \begin{cases} x & \text{if } x \in K, \\ x_0 & \text{otherwise} \end{cases}$$

for each  $x \in X$ ;

- $a: K \rightarrow Y$  is the active morphism of pointed sets given by

$$a(x) \stackrel{\text{def}}{=} f(x)$$

for each  $x \in K$ .

Next, let

$$\begin{array}{ccc} X & \xrightarrow{i} & Y \\ f \downarrow & & \downarrow g \\ A & \xrightarrow{a} & B \end{array}$$

be a commutative diagram in  $\text{Sets}_*$ . Consider the morphism  $\phi: Y \rightarrow A$  given by

$$\phi(y) = f(i^{-1}(y))$$

for each  $y \in Y$  (which is well-defined since, as  $i$  is inert,  $i^{-1}(y)$  is a singleton for all  $y \in Y$ ). We claim that  $\phi$  is the unique diagonal filler in the diagram

$$\begin{array}{ccc} X & \xrightarrow{i} & Y \\ f \downarrow & \nearrow \exists! \phi & \downarrow g \\ A & \xrightarrow{a} & B. \end{array}$$

Indeed, this diagram commutes, as we have

$$\begin{aligned} [\phi \circ i](x) &\stackrel{\text{def}}{=} \phi(i(x)) \\ &\stackrel{\text{def}}{=} f(i^{-1}(i(x))) \\ &= f(x) \end{aligned}$$

for each  $x \in X$  and

$$\begin{aligned} [a \circ \phi](y) &\stackrel{\text{def}}{=} a(\phi(y)) \\ &\stackrel{\text{def}}{=} a(f(i^{-1}(y))) \\ &\stackrel{\text{def}}{=} [a \circ f](i^{-1}(y)) \\ &= [g \circ i](i^{-1}(y)) \\ &\stackrel{\text{def}}{=} g(i(i^{-1}(y))) \\ &\stackrel{\text{def}}{=} g(y) \end{aligned}$$

for each  $y \in Y$ . Moreover, given another morphism  $\psi$  such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{i} & Y \\ f \downarrow & \nearrow \psi & \downarrow g \\ A & \xrightarrow{a} & B \end{array}$$

commutes, it follows that we must have  $\psi = \phi$ , since, given  $y \in Y$ , there exists a unique  $x \in X$  such that  $i(x) = y$ , so we have

$$\begin{aligned} \psi(y) &= \psi(i(x)) \\ &= f(x) \\ &= f(i^{-1}(y)) \\ &\stackrel{\text{def}}{=} \phi(y). \end{aligned}$$

This finishes the proof. □

**00A2 6.2 Limits of Pointed Sets**

**00A3 6.2.1 The Terminal Pointed Set**

**00A4 DEFINITION 6.2.1.1.1 ► THE TERMINAL POINTED SET**

The **terminal pointed set** is the terminal object of  $\text{Sets}_*$  as in ??.

**0250 CONSTRUCTION 6.2.1.1.2 ► CONSTRUCTION OF THE TERMINAL POINTED SET**

Concretely, the **terminal pointed set** is the pair  $((\text{pt}, \star), \{!_X\}_{(X, x_0) \in \text{Obj}(\text{Sets}_*)})$  consisting of:

- *The Limit.* The pointed set  $(\text{pt}, \star)$ .
- *The Cone.* The collection of morphisms of pointed sets

$$\{!_X : (X, x_0) \rightarrow (\text{pt}, \star)\}_{(X, x_0) \in \text{Obj}(\text{Sets}_*)}$$

defined by

$$!_X(x) \stackrel{\text{def}}{=} \star$$

for each  $x \in X$  and each  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

**PROOF 6.2.1.1.3 ► PROOF OF CONSTRUCTION 6.2.1.1.2**

We claim that  $(\text{pt}, \star)$  is the terminal object of  $\text{Sets}_*$ . Indeed, suppose we have a diagram of the form

$$(X, x_0) \quad (\text{pt}, \star)$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi : (X, x_0) \rightarrow (\text{pt}, \star)$$

making the diagram

$$(X, x_0) \xrightarrow[\exists !]{\phi} (\text{pt}, \star)$$

commute, namely  $!_X$ . 

**00A5 6.2.2 Products of Families of Pointed Sets**

Let  $\{(X_i, x_0^i)\}_{i \in I}$  be a family of pointed sets.

00A6

**DEFINITION 6.2.2.1.1 ► THE PRODUCT OF A FAMILY OF POINTED SETS**

The **product** of  $\{(X_i, x_0^i)\}_{i \in I}$  is the product of  $\{(X_i, x_0^i)\}_{i \in I}$  in  $\text{Sets}_*$  as in ??.

0251

**CONSTRUCTION 6.2.2.1.2 ► CONSTRUCTION OF THE PRODUCT OF A FAMILY OF POINTED SETS**

Concretely, the **product** of  $\{(X_i, x_0^i)\}_{i \in I}$  is the pair  $((\prod_{i \in I} X_i, (x_0^i)_{i \in I}), \{\text{pr}_i\}_{i \in I})$  consisting of:

- *The Limit.* The pointed set  $(\prod_{i \in I} X_i, (x_0^i)_{i \in I})$ .
- *The Cone.* The collection

$$\left\{ \text{pr}_i: \left( \prod_{i \in I} X_i, (x_0^i)_{i \in I} \right) \rightarrow (X_i, x_0^i) \right\}_{i \in I}$$

of maps given by

$$\text{pr}_i((x_j)_{j \in I}) \stackrel{\text{def}}{=} x_i$$

for each  $(x_j)_{j \in I} \in \prod_{i \in I} X_i$  and each  $i \in I$ .

**PROOF 6.2.2.1.3 ► PROOF OF CONSTRUCTION 6.2.2.1.2**

We claim that  $(\prod_{i \in I} X_i, (x_0^i)_{i \in I})$  is the categorical product of  $\{(X_i, x_0^i)\}_{i \in I}$  in  $\text{Sets}_*$ . Indeed, suppose we have, for each  $i \in I$ , a diagram of the form

$$\begin{array}{ccc} (P, *) & & \\ & \searrow p_i & \\ & (\prod_{i \in I} X_i, (x_0^i)_{i \in I}) & \xrightarrow{\text{pr}_i} (X_i, x_0^i) \end{array}$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi: (P, *) \rightarrow \left( \prod_{i \in I} X_i, (x_0^i)_{i \in I} \right)$$

making the diagram

$$\begin{array}{ccc} (P, *) & & \\ \downarrow \phi \quad \exists! & \searrow p_i & \\ (\prod_{i \in I} X_i, (x_0^i)_{i \in I}) & \xrightarrow{\text{pr}_i} & (X_i, x_0^i) \end{array}$$

commute, being uniquely determined by the condition  $\text{pr}_i \circ \phi = p_i$  for each  $i \in I$  via

$$\phi(x) = (p_i(x))_{i \in I}$$

for each  $x \in P$ . Note that this is indeed a morphism of pointed sets, as we have

$$\begin{aligned}\phi(*) &= (p_i(*))_{i \in I} \\ &= (x_0^i)_{i \in I},\end{aligned}$$

where we have used that  $p_i$  is a morphism of pointed sets for each  $i \in I$ .



00A7

**PROPOSITION 6.2.2.1.4 ► PROPERTIES OF PRODUCTS OF FAMILIES OF POINTED SETS**

Let  $\{(X_i, x_0^i)\}_{i \in I}$  be a family of pointed sets.

00A8

1. *Functoriality.* The assignment  $\{(X_i, x_0^i)\}_{i \in I} \mapsto (\prod_{i \in I} X_i, (x_0^i)_{i \in I})$  defines a functor

$$\prod_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}_*) \rightarrow \text{Sets}_*.$$

**PROOF 6.2.2.1.5 ► PROOF OF PROPOSITION 6.2.2.1.4**

Item 1: Functoriality

This follows from ?? of ??.



00A9

**6.2.3 Products**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00AA

**DEFINITION 6.2.3.1.1 ► PRODUCTS OF POINTED SETS**

The **product of**  $(X, x_0)$  **and**  $(Y, y_0)$  is the product of  $(X, x_0)$  and  $(Y, y_0)$  in  $\text{Sets}_*$  as in ??.

0252

**CONSTRUCTION 6.2.3.1.2 ► CONSTRUCTION OF PRODUCTS OF POINTED SETS**

Concretely, the **product of**  $(X, x_0)$  **and**  $(Y, y_0)$  is the pair consisting of:

- *The Limit.* The pointed set  $(X \times Y, (x_0, y_0))$ .

- *The Cone.* The morphisms of pointed sets

$$\begin{aligned} \text{pr}_1 &: (X \times Y, (x_0, y_0)) \rightarrow (X, x_0), \\ \text{pr}_2 &: (X \times Y, (x_0, y_0)) \rightarrow (Y, y_0) \end{aligned}$$

defined by

$$\begin{aligned} \text{pr}_1(x, y) &\stackrel{\text{def}}{=} x, \\ \text{pr}_2(x, y) &\stackrel{\text{def}}{=} y \end{aligned}$$

for each  $(x, y) \in X \times Y$ .

#### PROOF 6.2.3.1.3 ► PROOF OF CONSTRUCTION 6.2.3.1.2

We claim that  $(X \times Y, (x_0, y_0))$  is the categorical product of  $(X, x_0)$  and  $(Y, y_0)$  in  $\text{Sets}_*$ . Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & (P, *) & & \\ & p_1 \swarrow & & \searrow p_2 & \\ (X, x_0) & \xleftarrow[\text{pr}_1]{} & (X \times Y, (x_0, y_0)) & \xrightarrow[\text{pr}_2]{} & (Y, y_0) \end{array}$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi: (P, *) \rightarrow (X \times Y, (x_0, y_0))$$

making the diagram

$$\begin{array}{ccccc} & & (P, *) & & \\ & p_1 \swarrow & \downarrow \phi \exists! & \searrow p_2 & \\ (X, x_0) & \xleftarrow[\text{pr}_1]{} & (X \times Y, (x_0, y_0)) & \xrightarrow[\text{pr}_2]{} & (Y, y_0) \end{array}$$

commute, being uniquely determined by the conditions

$$\text{pr}_1 \circ \phi = p_1,$$

$$\text{pr}_2 \circ \phi = p_2$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each  $x \in P$ . Note that this is indeed a morphism of pointed sets, as we have

$$\begin{aligned}\phi(*) &= (p_1(*), p_2(*)) \\ &= (x_0, y_0),\end{aligned}$$

where we have used that  $p_1$  and  $p_2$  are morphisms of pointed sets.  $\blacksquare$

00AB

#### PROPOSITION 6.2.3.1.4 ► PROPERTIES OF PRODUCTS OF POINTED SETS

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

00AC

1. *Functionality.* The assignments

$$(X, x_0), (Y, y_0), ((X, x_0), (Y, y_0)) \mapsto (X \times Y, (x_0, y_0))$$

define functors

$$\begin{aligned}A \times -: \quad \text{Sets}_* &\rightarrow \text{Sets}_*, \\ - \times B: \quad \text{Sets}_* &\rightarrow \text{Sets}_*, \\ -_1 \times -_2: \quad \text{Sets}_* \times \text{Sets}_* &\rightarrow \text{Sets}_*,\end{aligned}$$

defined in the same way as the functors of Item 1 of Proposition 4.1.3.1.4.

01QR

2. *Lack of Adjointness.* The functors  $X \times -$  and  $- \times Y$  do not admit right adjoints.

00AD

3. *Associativity.* We have an isomorphism of pointed sets

$$((X \times Y) \times Z, ((x_0, y_0), z_0)) \cong (X \times (Y \times Z), (x_0, (y_0, z_0)))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ .

00AE

4. *Unitality.* We have isomorphisms of pointed sets

$$\begin{aligned}(\text{pt}, \star) \times (X, x_0) &\cong (X, x_0), \\ (X, x_0) \times (\text{pt}, \star) &\cong (X, x_0),\end{aligned}$$

natural in  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00AF

5. *Commutativity.* We have an isomorphism of pointed sets

$$(X \times Y, (x_0, y_0)) \cong (Y \times X, (y_0, x_0)),$$

natural in  $(X, x_0), (Y, y_0) \in \text{Obj}(\text{Sets}_*)$ .

00AG

6. *Symmetric Monoidality.* The triple  $(\text{Sets}_*, \times, (\text{pt}, \star))$  is a symmetric monoidal category.

#### PROOF 6.2.3.1.5 ► PROOF OF PROPOSITION 6.2.3.1.4

Item 1: Functoriality

This is a special case of functoriality of limits, ?? of ??.

Item 2: Lack of Adjointness

See [MSE 285586].

Item 3: Associativity

This follows from Item 4 of Proposition 4.1.3.1.4.

Item 4: Unitality

This follows from Item 5 of Proposition 4.1.3.1.4.

Item 5: Commutativity

This follows from Item 6 of Proposition 4.1.3.1.4.

Item 6: Symmetric Monoidality

This follows from Item 14 of Proposition 4.1.3.1.4. 

#### 00AH 6.2.4 Pullbacks

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets and let  $f: (X, x_0) \rightarrow (Z, z_0)$  and  $g: (Y, y_0) \rightarrow (Z, z_0)$  be morphisms of pointed sets.

00AJ

#### DEFINITION 6.2.4.1.1 ► PULLBACKS OF POINTED SETS

The **pullback of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  along  $(f, g)$**  is the pullback of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  along  $(f, g)$  in  $\text{Sets}_*$  as in ??.

0253

#### CONSTRUCTION 6.2.4.1.2 ► CONSTRUCTION OF PULLBACKS OF POINTED SETS

Concretely, the **pullback of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  along  $(f, g)$**  is the pair consisting of:

- *The Limit.* The pointed set  $(X \times_Z Y, (x_0, y_0))$ .
- *The Cone.* The morphisms of pointed sets

$$\begin{aligned} \text{pr}_1: (X \times_Z Y, (x_0, y_0)) &\rightarrow (X, x_0), \\ \text{pr}_2: (X \times_Z Y, (x_0, y_0)) &\rightarrow (Y, y_0) \end{aligned}$$

defined by

$$\begin{aligned}\text{pr}_1(x, y) &\stackrel{\text{def}}{=} x, \\ \text{pr}_2(x, y) &\stackrel{\text{def}}{=} y\end{aligned}$$

for each  $(x, y) \in X \times_Z Y$ .

**PROOF 6.2.4.1.3 ► PROOF OF CONSTRUCTION 6.2.4.1.2**

We claim that  $X \times_Z Y$  is the categorical pullback of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  with respect to  $(f, g)$  in  $\text{Sets}_*$ . First we need to check that the relevant pullback diagram commutes, i.e. that we have

$$\begin{array}{ccc}(X \times_Z Y, (x_0, y_0)) & \xrightarrow{\text{pr}_2} & (Y, y_0) \\ f \circ \text{pr}_1 = g \circ \text{pr}_2, & \text{pr}_1 \downarrow & \downarrow g \\ & & \\ (X, x_0) & \xrightarrow{f} & (Z, z_0).\end{array}$$

Indeed, given  $(x, y) \in X \times_Z Y$ , we have

$$\begin{aligned}[f \circ \text{pr}_1](x, y) &= f(\text{pr}_1(x, y)) \\ &= f(x) \\ &= g(y) \\ &= g(\text{pr}_2(x, y)) \\ &= [g \circ \text{pr}_2](x, y),\end{aligned}$$

where  $f(x) = g(y)$  since  $(x, y) \in X \times_Z Y$ . Next, we prove that  $X \times_Z Y$  satisfies the universal property of the pullback. Suppose we have a diagram of the form

$$\begin{array}{ccccc}(P, *) & \xrightarrow{p_2} & (X \times_Z Y, (x_0, y_0)) & \xrightarrow{-\text{pr}_2} & (Y, y_0) \\ & \swarrow p_1 & \downarrow \text{pr}_1 & \lrcorner & \downarrow g \\ & & (X, x_0) & \xrightarrow{f} & (Z, z_0)\end{array}$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi: (P, *) \rightarrow (X \times_Z Y, (x_0, y_0))$$

making the diagram

$$\begin{array}{ccccc}
 (P, *) & \xrightarrow{\quad p_1 \quad} & (X \times_Z Y, (x_0, y_0)) & \xrightarrow{\quad p_2 \quad} & (Y, y_0) \\
 & \exists! \phi \dashrightarrow & \downarrow \text{pr}_1 & \lrcorner & \downarrow g \\
 & & (X, x_0) & \xrightarrow{\quad f \quad} & (Z, z_0)
 \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned}
 \text{pr}_1 \circ \phi &= p_1, \\
 \text{pr}_2 \circ \phi &= p_2
 \end{aligned}$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each  $x \in P$ , where we note that  $(p_1(x), p_2(x)) \in X \times Y$  indeed lies in  $X \times_Z Y$  by the condition

$$f \circ p_1 = g \circ p_2,$$

which gives

$$f(p_1(x)) = g(p_2(x))$$

for each  $x \in P$ , so that  $(p_1(x), p_2(x)) \in X \times_Z Y$ . Lastly, we note that  $\phi$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned}
 \phi(*) &= (p_1(*), p_2(*)) \\
 &= (x_0, y_0),
 \end{aligned}$$

where we have used that  $p_1$  and  $p_2$  are morphisms of pointed sets. ■

00AK

**PROPOSITION 6.2.4.1.4 ► PROPERTIES OF PULLBACKS OF POINTED SETS**


Let  $(X, x_0)$ ,  $(Y, y_0)$ ,  $(Z, z_0)$ , and  $(A, a_0)$  be pointed sets.

00AL

1. *Functoriality.* The assignment  $(X, Y, Z, f, g) \mapsto X \times_{f, Z, g} Y$  defines a functor

$$-_1 \times_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}_*) \rightarrow \text{Sets}_*,$$

where  $\mathcal{P}$  is the category that looks like this:



In particular, the action on morphisms of  $-_1 \times_{-3} -_1$  is given by sending a morphism

$$\begin{array}{ccccc} X \times_Z Y & \xrightarrow{\quad} & Y & & \\ \downarrow & \lrcorner & \downarrow g & \searrow \psi & \\ X' \times_{Z'} Y' & \xrightarrow{\quad} & Y' & & \\ \downarrow & \lrcorner & \downarrow & & \\ X & \xrightarrow{f} & Z & & \\ \phi \searrow & & \downarrow & \swarrow \chi & \\ & & X' & \xrightarrow{f'} & Z' \end{array}$$

in  $\text{Fun}(\mathcal{P}, \text{Sets}_*)$  to the morphism of pointed sets

$$\xi: (X \times_Z Y, (x_0, y_0)) \xrightarrow{\exists!} (X' \times_{Z'} Y', (x'_0, y'_0))$$

given by

$$\xi(x, y) \stackrel{\text{def}}{=} (\phi(x), \psi(y))$$

for each  $(x, y) \in X \times_Z Y$ , which is the unique morphism of pointed sets making the diagram

$$\begin{array}{ccccc} X \times_Z Y & \xrightarrow{\quad} & Y & & \\ \downarrow & \lrcorner & \downarrow g & \searrow \psi & \\ X' \times_{Z'} Y' & \xrightarrow{\quad} & Y' & & \\ \downarrow & \lrcorner & \downarrow & & \\ X & \xrightarrow{f} & Z & & \\ \phi \searrow & & \downarrow & \swarrow \chi & \\ & & X' & \xrightarrow{f'} & Z' \end{array}$$

commute.

00AM

2. *Associativity.* Given a diagram

$$\begin{array}{ccccc} X & & Y & & Z \\ & \searrow f & \swarrow g & \searrow h & \swarrow k \\ & W & & V & \end{array}$$

in  $\text{Sets}_*$ , we have isomorphisms of pointed sets

$$(X \times_W Y) \times_V Z \cong (X \times_W Y) \times_Y (Y \times_V Z) \cong X \times_W (Y \times_V Z),$$

where these pullbacks are built as in the diagrams

$$\begin{array}{c} (X \times_W Y) \times_Y Z \\ \downarrow \quad \swarrow \quad \searrow \\ X \times_W Y \quad \quad \quad Y \times_V Z \\ \downarrow \quad \swarrow \quad \searrow \\ X \quad \quad \quad Y \quad \quad \quad Z \\ \downarrow f \quad \downarrow g \quad \downarrow h \quad \downarrow k \\ W \quad \quad V \quad \quad V \quad \quad Z \\ \text{and} \\ (X \times_W Y) \times_Y (Y \times_V Z) \\ \downarrow \quad \swarrow \quad \searrow \\ X \times_W Y \quad \quad \quad Y \times_V Z \\ \downarrow \quad \swarrow \quad \searrow \\ X \quad \quad \quad Y \quad \quad \quad Z \\ \downarrow f \quad \downarrow g \quad \downarrow h \quad \downarrow k \\ W \quad \quad V \quad \quad V \quad \quad Z \\ \text{and} \\ X \times_W (Y \times_V Z) \\ \downarrow \quad \swarrow \quad \searrow \\ X \quad \quad \quad Y \times_V Z \\ \downarrow \quad \swarrow \quad \searrow \\ X \quad \quad \quad Y \quad \quad \quad Z \\ \downarrow f \quad \downarrow g \quad \downarrow h \quad \downarrow k \\ W \quad \quad V \quad \quad V \quad \quad Z \end{array}$$

00AN

3. *Unitality.* We have isomorphisms of pointed sets

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ f \downarrow & \lrcorner & \downarrow f \\ X & \xlongequal{\quad} & X \end{array} \quad \begin{array}{c} X \times_X A \cong A, \\ A \times_X X \cong A, \end{array} \quad \begin{array}{ccc} A & \xrightarrow{f} & X \\ \parallel & \lrcorner & \parallel \\ X & \xrightarrow{f} & X. \end{array}$$

00AP

4. *Commutativity.* We have an isomorphism of pointed sets

$$\begin{array}{ccc} A \times_X B & \longrightarrow & B \\ \downarrow \lrcorner & & \downarrow g \\ A & \xrightarrow{f} & X, \end{array} \quad A \times_X B \cong B \times_X A \quad \begin{array}{ccc} B \times_X A & \longrightarrow & A \\ \downarrow \lrcorner & & \downarrow f \\ B & \xrightarrow{g} & X. \end{array}$$

00AQ

5. *Interaction With Products.* We have an isomorphism of pointed sets

$$\begin{array}{ccc} X \times_{\text{pt}} Y & \longrightarrow & Y \\ \downarrow \lrcorner & & \downarrow !_Y \\ X & \xrightarrow{!_X} & \text{pt}. \end{array}$$

00AR

6. *Symmetric Monoidality.* The triple  $(\text{Sets}_*, \times_X, X)$  is a symmetric monoidal category.

**PROOF 6.2.4.1.5 ▶ PROOF OF PROPOSITION 6.2.4.1.4****Item 1: Functoriality**

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for  $\xi$  following from the commutativity of the cube pullback diagram.

**Item 2: Associativity**

This follows from [Item 4 of Proposition 6.2.4.1.4](#).

**Item 3: Unitality**

This follows from [Item 6 of Proposition 4.1.4.1.7](#).

**Item 4: Commutativity**

This follows from [Item 7 of Proposition 4.1.4.1.7](#).

**Item 5: Interaction With Products**

This follows from [Item 10 of Proposition 4.1.4.1.7](#).

**Item 6: Symmetric Monoidality**

This follows from [Item 11 of Proposition 4.1.4.1.7](#). 

**00AS 6.2.5 Equalisers**

Let  $f, g: (X, x_0) \rightrightarrows (Y, y_0)$  be morphisms of pointed sets.

**00AT****DEFINITION 6.2.5.1.1 ▶ EQUALISERS OF POINTED SETS**

The **equaliser of**  $(f, g)$  is the equaliser of  $f$  and  $g$  in  $\text{Sets}_*$  as in ??.

**0254****CONSTRUCTION 6.2.5.1.2 ▶ CONSTRUCTION OF EQUALISERS OF POINTED SETS**

Concretely, the **equaliser of**  $(f, g)$  is the pair consisting of:

- *The Limit.* The pointed set  $(\text{Eq}(f, g), x_0)$ .
- *The Cone.* The morphism of pointed sets

$$\text{eq}(f, g): (\text{Eq}(f, g), x_0) \hookrightarrow (X, x_0)$$

given by the canonical inclusion  $\text{eq}(f, g) \hookrightarrow \text{Eq}(f, g) \hookrightarrow X$ .

**PROOF 6.2.5.1.3 ▶ PROOF OF CONSTRUCTION 6.2.5.1.2**

We claim that  $(\text{Eq}(f, g), x_0)$  is the categorical equaliser of  $f$  and  $g$  in  $\text{Sets}_*$ . First we need to check that the relevant equaliser diagram commutes, i.e. that we have

$$f \circ \text{eq}(f, g) = g \circ \text{eq}(f, g),$$

which indeed holds by the definition of the set  $\text{Eq}(f, g)$ . Next, we prove that  $\text{Eq}(f, g)$  satisfies the universal property of the equaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} (\text{Eq}(f, g), x_0) & \xrightarrow{\text{eq}(f, g)} & (X, x_0) \xrightarrow[g]{f} (Y, y_0) \\ & \searrow e & \\ & (E, *) & \end{array}$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi: (E, *) \rightarrow (\text{Eq}(f, g), x_0)$$

making the diagram

$$\begin{array}{ccc} (\text{Eq}(f, g), x_0) & \xrightarrow{\text{eq}(f, g)} & (X, x_0) \xrightarrow[g]{f} (Y, y_0) \\ \uparrow \phi \exists! & \nearrow e & \\ (E, *) & & \end{array}$$

commute, being uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

via

$$\phi(x) = e(x)$$

for each  $x \in E$ , where we note that  $e(x) \in A$  indeed lies in  $\text{Eq}(f, g)$  by the condition

$$f \circ e = g \circ e,$$

which gives

$$f(e(x)) = g(e(x))$$

for each  $x \in E$ , so that  $e(x) \in \text{Eq}(f, g)$ . Lastly, we note that  $\phi$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(*) &= e(*) \\ &= x_0, \end{aligned}$$

where we have used that  $e$  is a morphism of pointed sets. □

00AU

**PROPOSITION 6.2.5.1.4 ► PROPERTIES OF EQUALISERS OF POINTED SETS**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets and let  $f, g, h: (X, x_0) \rightarrow (Y, y_0)$  be morphisms of pointed sets.

00AV

1. *Associativity.* We have isomorphisms of pointed sets

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}$$

where  $\text{Eq}(f, g, h)$  is the limit of the diagram

$$(X, x_0) \xrightarrow[\substack{f \\ g \\ h}]{} (Y, y_0)$$

in  $\text{Sets}_*$ , being explicitly given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

00AW

2. *Unitality.* We have an isomorphism of pointed sets

$$\text{Eq}(f, f) \cong X.$$

00AX

3. *Commutativity.* We have an isomorphism of pointed sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

**PROOF 6.2.5.1.5 ► PROOF OF PROPOSITION 6.2.5.1.4**

Item 1: *Associativity*

This follows from Item 1 of Proposition 4.1.5.1.4.

Item 2: *Unitality*

This follows from Item 2 of Proposition 4.1.5.1.4.

Item 3: *Commutativity*

This follows from Item 3 of Proposition 4.1.5.1.4. 

## 00AY 6.3 Colimits of Pointed Sets

### 00AZ 6.3.1 The Initial Pointed Set

00B0

#### DEFINITION 6.3.1.1.1 ► THE INITIAL POINTED SET

The **initial pointed set** is the initial object of  $\text{Sets}_*$  as in ??.

0255

#### CONSTRUCTION 6.3.1.1.2 ► CONSTRUCTION OF THE INITIAL POINTED SET

Concretely, the **initial pointed set** is the pair  $((\text{pt}, \star), \{\iota_X\}_{(X, x_0) \in \text{Obj}(\text{Sets}_*)})$  consisting of:

- *The Limit.* The pointed set  $(\text{pt}, \star)$ .
- *The Cone.* The collection of morphisms of pointed sets

$$\{\iota_X: (\text{pt}, \star) \rightarrow (X, x_0)\}_{(X, x_0) \in \text{Obj}(\text{Sets}_*)}$$

defined by

$$\iota_X(\star) \stackrel{\text{def}}{=} x_0.$$

#### PROOF 6.3.1.1.3 ► PROOF OF CONSTRUCTION 6.3.1.1.2

We claim that  $(\text{pt}, \star)$  is the initial object of  $\text{Sets}_*$ . Indeed, suppose we have a diagram of the form

$$(\text{pt}, \star) \quad (X, x_0)$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi: (\text{pt}, \star) \rightarrow (X, x_0)$$

making the diagram

$$(\text{pt}, \star) \xrightarrow[\exists!]{\phi} (X, x_0)$$

commute, namely  $\iota_X$ . □

### 00B1 6.3.2 Coproducts of Families of Pointed Sets

Let  $\{(X_i, x_0^i)\}_{i \in I}$  be a family of pointed sets.

00B2

**DEFINITION 6.3.2.1.1 ► COPRODUCTS OF FAMILIES OF POINTED SETS**

The **coproduct of the family**  $\{(X_i, x_0^i)\}_{i \in I}$ <sup>1</sup> is the coproduct of  $\{(X_i, x_0^i)\}_{i \in I}$  in  $\text{Sets}_*$  as in ??.

<sup>1</sup>Further Terminology: Also called the **wedge sum of the family**  $\{(X_i, x_0^i)\}_{i \in I}$ .

0256

**CONSTRUCTION 6.3.2.1.2 ► CONSTRUCTION OF COPRODUCTS OF FAMILIES OF POINTED SETS**

Concretely, the **coproduct of the family**  $\{(X_i, x_0^i)\}_{i \in I}$  is the pair  $((\bigvee_{i \in I} X_i, p_0), \{\text{inj}_i\}_{i \in I})$  consisting of:

- *The Colimit.* The pointed set  $(\bigvee_{i \in I} X_i, p_0)$  consisting of:

- *The Underlying Set.* The set  $\bigvee_{i \in I} X_i$  defined by

$$\bigvee_{i \in I} X_i \stackrel{\text{def}}{=} (\coprod_{i \in I} X_i) / \sim,$$

where  $\sim$  is the equivalence relation on  $\coprod_{i \in I} X_i$  given by declaring

$$(i, x_0^i) \sim (j, x_0^j)$$

for each  $i, j \in I$ .

- *The Basepoint.* The element  $p_0$  of  $\bigvee_{i \in I} X_i$  defined by

$$\begin{aligned} p_0 &\stackrel{\text{def}}{=} [(i, x_0^i)] \\ &= [(j, x_0^j)] \end{aligned}$$

for any  $i, j \in I$ .

- *The Cocone.* The collection

$$\left\{ \text{inj}_i : (X_i, x_0^i) \rightarrow \left( \bigvee_{i \in I} X_i, p_0 \right) \right\}_{i \in I}$$

of morphism of pointed sets given by

$$\text{inj}_i(x) \stackrel{\text{def}}{=} (i, x)$$

for each  $x \in X_i$  and each  $i \in I$ .

**PROOF 6.3.2.1.3 ▶ PROOF OF CONSTRUCTION 6.3.2.1.2**

We claim that  $(\bigvee_{i \in I} X_i, p_0)$  is the categorical coproduct of  $\{(X_i, x_0^i)\}_{i \in I}$  in  $\text{Sets}_*$ . Indeed, suppose we have, for each  $i \in I$ , a diagram of the form

$$\begin{array}{ccc} & & (C, *) \\ & \nearrow \iota_i & \\ (X_i, x_0^i) & \xrightarrow{\text{inj}_i} & (\bigvee_{i \in I} X_i, p_0) \end{array}$$

in  $\text{Sets}_*$ . Then there exists a unique morphism of pointed sets

$$\phi: (\bigvee_{i \in I} X_i, p_0) \rightarrow (C, *)$$

making the diagram

$$\begin{array}{ccc} & & (C, *) \\ & \nearrow \iota_i & \uparrow \phi \quad \exists! \\ (X_i, x_0^i) & \xrightarrow{\text{inj}_i} & (\bigvee_{i \in I} X_i, p_0) \end{array}$$

commute, being uniquely determined by the condition  $\phi \circ \text{inj}_i = \iota_i$  for each  $i \in I$  via

$$\phi([(i, x)]) = \iota_i(x)$$

for each  $[(i, x)] \in \bigvee_{i \in I} X_i$ , where we note that  $\phi$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(p_0) &= \iota_i([(i, x_0^i)]) \\ &= *, \end{aligned}$$

as  $\iota_i$  is a morphism of pointed sets. □

00B3

**PROPOSITION 6.3.2.1.4 ▶ PROPERTIES OF COPRODUCTS OF FAMILIES OF POINTED SETS**

Let  $\{(X_i, x_0^i)\}_{i \in I}$  be a family of pointed sets.

00B4

- Functoriality.* The assignment  $\{(X_i, x_0^i)\}_{i \in I} \mapsto (\bigvee_{i \in I} X_i, p_0)$  defines a functor

$$\bigvee_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}_*) \rightarrow \text{Sets}_*.$$

**PROOF 6.3.2.1.5 ► PROOF OF PROPOSITION 6.3.2.1.4**

Item 1: Functoriality

This follows from ?? of ??.

**00B5 6.3.3 Coproducts**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00B6**

**DEFINITION 6.3.3.1.1 ► COPRODUCTS OF POINTED SETS**

The **coproduct of  $(X, x_0)$  and  $(Y, y_0)$** <sup>1</sup> is the coproduct of  $(X, x_0)$  and  $(Y, y_0)$  in Sets<sub>\*</sub> as in ??.

<sup>1</sup>Further Terminology: Also called the **wedge sum of  $(X, x_0)$  and  $(Y, y_0)$** .

**0257**

**CONSTRUCTION 6.3.3.1.2 ► CONSTRUCTION OF COPRODUCTS OF POINTED SETS**

Concretely, the **coproduct of  $(X, x_0)$  and  $(Y, y_0)$** , also called their **wedge sum**, is the pair consisting of:

- *The Colimit.* The pointed set  $(X \vee Y, p_0)$  consisting of:

- *The Underlying Set.* The set  $X \vee Y$  defined by

$$\begin{aligned} (X \vee Y, p_0) &\stackrel{\text{def}}{=} (X, x_0) \coprod (Y, y_0) \\ &\cong (X \coprod_{\text{pt}} Y, p_0) \\ &\cong (X \coprod Y / \sim, p_0), \end{aligned}$$

$$\begin{array}{ccc} X \vee Y & \xleftarrow{\quad} & Y \\ \uparrow \lrcorner & & \uparrow [y_0] \\ X & \xleftarrow{\quad [x_0] \quad} & \text{pt}, \end{array}$$

where  $\sim$  is the equivalence relation on  $X \coprod Y$  obtained by declaring  $(0, x_0) \sim (1, y_0)$ .

- *The Basepoint.* The element  $p_0$  of  $X \vee Y$  defined by

$$\begin{aligned} p_0 &\stackrel{\text{def}}{=} [(0, x_0)] \\ &= [(1, y_0)]. \end{aligned}$$

- *The Cocone.* The morphisms of pointed sets

$$\begin{aligned} \text{inj}_1: (X, x_0) &\rightarrow (X \vee Y, p_0), \\ \text{inj}_2: (Y, y_0) &\rightarrow (X \vee Y, p_0), \end{aligned}$$

given by

$$\text{inj}_1(x) \stackrel{\text{def}}{=} [(0, x)],$$

$$\text{inj}_2(y) \stackrel{\text{def}}{=} [(1, y)],$$

for each  $x \in X$  and each  $y \in Y$ .

**PROOF 6.3.3.1.3 ► PROOF OF CONSTRUCTION 6.3.3.1.2**

We claim that  $(X \vee Y, p_0)$  is the categorical coproduct of  $(X, x_0)$  and  $(Y, y_0)$  in  $\text{Sets}_*$ . Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & (C, *) & & \\ & \nearrow \iota_1 & & \searrow \iota_2 & \\ (X, x_0) & \xrightarrow[\text{inj}_1]{} & (X \vee Y, p_0) & \xleftarrow[\text{inj}_2]{} & (Y, y_0) \end{array}$$

in  $\text{Sets}$ . Then there exists a unique morphism of pointed sets

$$\phi: (X \vee Y, p_0) \rightarrow (C, *)$$

making the diagram

$$\begin{array}{ccccc} & & (C, *) & & \\ & \nearrow \iota_1 & \uparrow \phi \exists! & \searrow \iota_2 & \\ (X, x_0) & \xrightarrow[\text{inj}_1]{} & (X \vee Y, p_0) & \xleftarrow[\text{inj}_2]{} & (Y, y_0) \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned} \phi \circ \text{inj}_X &= \iota_X, \\ \phi \circ \text{inj}_Y &= \iota_Y \end{aligned}$$

via

$$\phi(z) = \begin{cases} \iota_X(x) & \text{if } z = [(0, x)] \text{ with } x \in X, \\ \iota_Y(y) & \text{if } z = [(1, y)] \text{ with } y \in Y \end{cases}$$

for each  $z \in X \vee Y$ , where we note that  $\phi$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(p_0) &= \iota_X([(0, x_0)]) \\ &= \iota_Y([(1, y_0)]) \\ &= *, \end{aligned}$$

as  $\iota_X$  and  $\iota_Y$  are morphisms of pointed sets.

00B7

**PROPOSITION 6.3.3.1.4 ► PROPERTIES OF WEDGE SUMS OF POINTED SETS**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00B8

1. *Functionality.* The assignments

$$(X, x_0), (Y, y_0), ((X, x_0), (Y, y_0)) \mapsto (X \vee Y, p_0)$$

define functors

$$\begin{aligned} X \vee - &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \vee Y &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ -_1 \vee -_2 &: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

00B9

2. *Associativity.* We have an isomorphism of pointed sets

$$(X \vee Y) \vee Z \cong X \vee (Y \vee Z),$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Sets}_*$ .

00BA

3. *Unitality.* We have isomorphisms of pointed sets

$$\begin{aligned} (\text{pt}, *) \vee (X, x_0) &\cong (X, x_0), \\ (X, x_0) \vee (\text{pt}, *) &\cong (X, x_0), \end{aligned}$$

natural in  $(X, x_0) \in \text{Sets}_*$ .

00BB

4. *Commutativity.* We have an isomorphism of pointed sets

$$X \vee Y \cong Y \vee X,$$

natural in  $(X, x_0), (Y, y_0) \in \text{Sets}_*$ .

00BC

5. *Symmetric Monoidality.* The triple  $(\text{Sets}_*, \vee, \text{pt})$  is a symmetric monoidal category.

00BD

6. *The Fold Map.* We have a natural transformation

$$\nabla: \vee \circ \Delta_{\text{Sets}_*}^{\text{Cats}} \Rightarrow \text{id}_{\text{Sets}_*},$$

called the **fold map**, whose component

$$\nabla_X: X \vee X \rightarrow X$$

at  $X$  is given by

$$\nabla_X(p) \stackrel{\text{def}}{=} \begin{cases} x & \text{if } p = [(0, x)], \\ x & \text{if } p = [(1, x)] \end{cases}$$

for each  $p \in X \vee X$ .

**PROOF 6.3.3.1.5 ▶ PROOF OF PROPOSITION 6.3.3.1.4****Item 1: Functoriality**

This follows from ?? of ??.

**Item 2: Associativity**

Omitted.

**Item 3: Unitality**

Omitted.

**Item 4: Commutativity**

Omitted.

**Item 5: Symmetric Monoidality**

Omitted.

**Item 6: The Fold Map**

Naturality for the transformation  $\nabla$  is the statement that, given a morphism of pointed sets  $f: (X, x_0) \rightarrow (Y, y_0)$ , we have

$$\begin{array}{ccc} X \vee X & \xrightarrow{\nabla_X} & X \\ \nabla_Y \circ (f \vee f) = f \circ \nabla_X, & \downarrow f \vee f & \downarrow f \\ Y \vee Y & \xrightarrow{\nabla_Y} & Y. \end{array}$$

Indeed, we have

$$\begin{aligned} [\nabla_Y \circ (f \vee f)]([(i, x)]) &= \nabla_Y([(i, f(x))]) \\ &= f(x) \\ &= f(\nabla_X([(i, x)])) \\ &= [f \circ \nabla_X]([(i, x)]) \end{aligned}$$

for each  $[(i, x)] \in X \vee X$ , and thus  $\nabla$  is indeed a natural transformation.

**00BE 6.3.4 Pushouts**

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets and let  $f: (Z, z_0) \rightarrow (X, x_0)$  and  $g: (Z, z_0) \rightarrow (Y, y_0)$  be morphisms of pointed sets.

**00BF****DEFINITION 6.3.4.1.1 ▶ PUSHOUTS OF POINTED SETS**

The **pushout** of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  along  $(f, g)$  is the pushout of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  along  $(f, g)$  in  $\text{Sets}_*$  as in

??.

0258

#### CONSTRUCTION 6.3.4.1.2 ► CONSTRUCTION OF PUSHOUTS OF POINTED SETS

Concretely, the **pushout of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  along  $(f, g)$**  is the pair consisting of:

- *The Colimit.* The pointed set  $(X \coprod_{f, Z, g} Y, p_0)$ , where:
  - The set  $X \coprod_{f, Z, g} Y$  is the pushout (of unpointed sets) of  $X$  and  $Y$  over  $Z$  with respect to  $f$  and  $g$ ;
  - We have  $p_0 = [x_0] = [y_0]$ .
- *The Cocone.* The morphisms of pointed sets

$$\begin{aligned}\text{inj}_1 &: (X, x_0) \rightarrow (X \coprod_Z Y, p_0), \\ \text{inj}_2 &: (Y, y_0) \rightarrow (X \coprod_Z Y, p_0)\end{aligned}$$

given by

$$\begin{aligned}\text{inj}_1(x) &\stackrel{\text{def}}{=} [(0, x)] \\ \text{inj}_2(y) &\stackrel{\text{def}}{=} [(1, y)]\end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ .

#### PROOF 6.3.4.1.3 ► PROOF OF ??

Firstly, we note that indeed  $[x_0] = [y_0]$ , as we have

$$\begin{aligned}x_0 &= f(z_0), \\ y_0 &= g(z_0)\end{aligned}$$

since  $f$  and  $g$  are morphisms of pointed sets, with the relation  $\sim$  on  $X \coprod_Z Y$  then identifying  $x_0 = f(z_0) \sim g(z_0) = y_0$ .

We now claim that  $(X \coprod_Z Y, p_0)$  is the categorical pushout of  $(X, x_0)$  and  $(Y, y_0)$  over  $(Z, z_0)$  with respect to  $(f, g)$  in  $\text{Sets}_*$ . First we need to check that the relevant pushout diagram commutes, i.e. that we have

$$\begin{array}{ccc} (X \coprod_Z Y, p_0) & \xleftarrow{\text{inj}_2} & (Y, y_0) \\ \text{inj}_1 \uparrow & & \uparrow g \\ (X, x_0) & \xleftarrow{f} & (Z, z_0). \end{array}$$

Indeed, given  $z \in Z$ , we have

$$\begin{aligned} [\text{inj}_1 \circ f](z) &= \text{inj}_1(f(z)) \\ &= [(0, f(z))] \\ &= [(1, g(z))] \\ &= \text{inj}_2(g(z)) \\ &= [\text{inj}_2 \circ g](z), \end{aligned}$$

where  $[(0, f(z))] = [(1, g(z))]$  by the definition of the relation  $\sim$  on  $X \coprod Y$  (the coproduct of unpointed sets of  $X$  and  $Y$ ). Next, we prove that  $X \coprod_Z Y$  satisfies the universal property of the pushout. Suppose we have a diagram of the form

$$\begin{array}{ccc} (P, *) & \xleftarrow{\quad \iota_2 \quad} & \\ \uparrow & & \\ (X \coprod_Z Y, p_0) & \xleftarrow{\quad \text{inj}_2 - \quad} & (Y, y_0) \\ \uparrow \text{inj}_1 & \lrcorner & \uparrow g \\ (X, x_0) & \xleftarrow{\quad f \quad} & (Z, z_0) \end{array}$$

in  $\text{Sets}_{*}$ . Then there exists a unique morphism of pointed sets

$$\phi: (X \coprod_Z Y, p_0) \rightarrow (P, *)$$

making the diagram

$$\begin{array}{ccc} (P, *) & \xleftarrow{\quad \iota_2 \quad} & \\ \uparrow \phi \exists! & & \\ (X \coprod_Z Y, p_0) & \xleftarrow{\quad \text{inj}_2 - \quad} & (Y, y_0) \\ \uparrow \text{inj}_1 & \lrcorner & \uparrow g \\ (X, x_0) & \xleftarrow{\quad f \quad} & (Z, z_0) \end{array}$$

commute, being uniquely determined by the conditions

$$\phi \circ \text{inj}_1 = \iota_1,$$

$$\phi \circ \text{inj}_2 = \iota_2$$

via

$$\phi(p) = \begin{cases} \iota_1(x) & \text{if } x = [(0, x)], \\ \iota_2(y) & \text{if } x = [(1, y)] \end{cases}$$

for each  $p \in X \coprod_Z Y$ , where the well-definedness of  $\phi$  is proven in the same way as in the proof of [Definition 4.2.4.1.1](#). Finally, we show that  $\phi$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(p_0) &= \phi([(0, x_0)]) \\ &= \iota_1(x_0) \\ &= *, \end{aligned}$$

or alternatively

$$\begin{aligned} \phi(p_0) &= \phi([(1, y_0)]) \\ &= \iota_2(y_0) \\ &= *, \end{aligned}$$

where we use that  $\iota_1$  (resp.  $\iota_2$ ) is a morphism of pointed sets. ■

00BG

#### PROPOSITION 6.3.4.1.4 ► PROPERTIES OF PUSHOUTS OF POINTED SETS


Let  $(X, x_0)$ ,  $(Y, y_0)$ ,  $(Z, z_0)$ , and  $(A, a_0)$  be pointed sets.

00BH

- Functionality.* The assignment  $(X, Y, Z, f, g) \mapsto X \coprod_{f, Z, g} Y$  defines a functor

$$-_1 \coprod_{-3} -_1: \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets}_*,$$

where  $\mathcal{P}$  is the category that looks like this:



In particular, the action on morphisms of  $-_1 \amalg_{-3} -_1$  is given by sending a morphism

$$\begin{array}{ccccc}
 X \amalg_Z Y & \xleftarrow{\quad \lrcorner \quad} & Y & \xrightarrow{\psi} & \\
 \uparrow & & \uparrow & & \\
 X' \amalg_{Z'} Y' & \xleftarrow{\quad \lrcorner \quad} & Y' & \xleftarrow{g} & \\
 \uparrow & & \downarrow g' & & \\
 X & \xleftarrow{f} & Z & \xrightarrow{\chi} & Z' \\
 \downarrow \phi & & \downarrow & & \downarrow g' \\
 X' & \xleftarrow{f'} & Z' & & 
 \end{array}$$

in  $\text{Fun}(\mathcal{P}, \text{Sets}_*)$  to the morphism of pointed sets

$$\xi: (X \amalg_Z Y, p_0) \xrightarrow{\exists!} (X' \amalg_{Z'} Y', p'_0)$$

given by

$$\xi(p) \stackrel{\text{def}}{=} \begin{cases} \phi(x) & \text{if } p = [(0, x)], \\ \psi(y) & \text{if } p = [(1, y)] \end{cases}$$

for each  $p \in X \amalg_Z Y$ , which is the unique morphism of pointed sets making the diagram

$$\begin{array}{ccccc}
 X \amalg_Z Y & \xleftarrow{\quad \lrcorner \quad} & Y & \xrightarrow{\psi} & \\
 \uparrow & \searrow & \uparrow & & \\
 X' \amalg_{Z'} Y' & \xleftarrow{\quad \lrcorner \quad} & Y' & \xleftarrow{g} & \\
 \uparrow & & \downarrow g' & & \\
 X & \xleftarrow{f} & Z & \xrightarrow{\chi} & Z' \\
 \downarrow \phi & & \downarrow & & \downarrow g' \\
 X' & \xleftarrow{f'} & Z' & & 
 \end{array}$$

commute.

00BJ

2. *Associativity.* Given a diagram

$$\begin{array}{ccccc}
 X & & Y & & Z \\
 & \swarrow f & \nearrow g & \swarrow h & \nearrow k \\
 W & & V & & 
 \end{array}$$

in Sets, we have isomorphisms of pointed sets

$$(X \coprod_W Y) \coprod_V Z \cong (X \coprod_W Y) \coprod_Y (Y \coprod_V Z) \cong X \coprod_W (Y \coprod_V Z),$$

where these pullbacks are built as in the diagrams

$$\begin{array}{c} (X \coprod_W Y) \coprod_V Z \\ \nearrow \quad \searrow \\ X \coprod_W Y & & Z, \\ \nearrow f \quad \nearrow g \quad \nearrow h \quad \nearrow k \\ W \quad Y \quad V \quad Z \end{array} \quad \begin{array}{c} (X \coprod_W Y) \coprod_Y (Y \coprod_V Z) \\ \nearrow \quad \searrow \\ X \coprod_W Y & Y \coprod_V Z & Z, \\ \nearrow f \quad \nearrow g \quad \nearrow h \quad \nearrow k \\ W \quad Y \quad V \quad Z \end{array} \quad \begin{array}{c} X \coprod_W (Y \coprod_V Z) \\ \nearrow \quad \searrow \\ X & Y \coprod_V Z & Z, \\ \nearrow f \quad \nearrow g \quad \nearrow h \quad \nearrow k \\ W \quad Y \quad V \quad Z \end{array}$$

00BK

3. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ \uparrow f & & \uparrow f \\ X & \xlongequal{\quad} & X \end{array} \quad \begin{array}{c} X \coprod_X A \cong A, \\ A \coprod_X X \cong A, \end{array} \quad \begin{array}{ccc} A & \xleftarrow{f} & X \\ \parallel & & \parallel \\ X & \xleftarrow{f} & X. \end{array}$$

00BL

4. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc} X \coprod_Z Y & \xleftarrow{\quad} & Y \\ \uparrow f & & \uparrow g \\ X & \xleftarrow{f} & Z, \end{array} \quad X \coprod_Z Y \cong Y \coprod_Z X \quad \begin{array}{ccc} Y \coprod_Z X & \xleftarrow{\quad} & X \\ \uparrow f & & \uparrow g \\ Y & \xleftarrow{g} & Z. \end{array}$$

00BM

5. *Interaction With Coproducts.* We have

$$\begin{array}{c} X \vee Y \xleftarrow{\quad} Y \\ \uparrow f \qquad \uparrow [y_0] \\ X \xleftarrow{[x_0]} \text{pt.} \end{array} \quad X \coprod_{\text{pt}} Y \cong X \vee Y,$$

00BN

6. *Symmetric Monoidality.* The triple  $(\text{Sets}_*, \coprod_X, (X, x_0))$  is a symmetric monoidal category.

**PROOF 6.3.4.1.5 ▶ PROOF OF PROPOSITION 6.3.4.1.4****Item 1: Functoriality**

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for  $\xi$  following from the commutativity of the cube pushout diagram.

**Item 2: Associativity**

This follows from [Item 3 of Proposition 4.2.4.1.8](#).

**Item 3: Unitality**

This follows from [Item 5 of Proposition 4.2.4.1.8](#).

**Item 4: Commutativity**

This follows from [Item 6 of Proposition 4.2.4.1.8](#).

**Item 5: Interaction With Coproducts**

Omitted.

**Item 6: Symmetric Monoidality**

Omitted. 

**00BP 6.3.5 Coequalisers**

Let  $f, g: (X, x_0) \rightrightarrows (Y, y_0)$  be morphisms of pointed sets.

**00BQ****DEFINITION 6.3.5.1.1 ▶ COEQUALISERS OF POINTED SETS**

The **coequaliser of**  $(f, g)$  is the pointed set  $(\text{CoEq}(f, g), [y_0])$ .

**0259****CONSTRUCTION 6.3.5.1.2 ▶ CONSTRUCTION OF COEQUALISERS OF POINTED SETS**

The **coequaliser of**  $(f, g)$  is the pair  $((\text{CoEq}(f, g), [y_0]), \text{coeq}(f, g))$  consisting of:

- *The Colimit.* The pointed set  $(\text{CoEq}(f, g), [y_0])$ , where  $\text{CoEq}(f, g)$  is the coequaliser of  $f$  and  $g$  as in [Definition 4.2.5.1.1](#).
- *The Cocone.* The map

$$\text{coeq}(f, g): Y \twoheadrightarrow (\text{CoEq}(f, g), [y_0])$$

given by the quotient map, as in [Item 2 of Construction 4.2.5.1.2](#).

## PROOF 6.3.5.1.3 ▶ PROOF OF CONSTRUCTION 6.3.5.1.2

We claim that  $(\text{CoEq}(f, g), [y_0])$  is the categorical coequaliser of  $f$  and  $g$  in  $\text{Sets}_*$ . First we need to check that the relevant coequaliser diagram commutes, i.e. that we have

$$\text{coeq}(f, g) \circ f = \text{coeq}(f, g) \circ g.$$

Indeed, we have

$$\begin{aligned} [\text{coeq}(f, g) \circ f](x) &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](f(x)) \\ &\stackrel{\text{def}}{=} [f(x)] \\ &= [g(x)] \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](g(x)) \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g) \circ g](x) \end{aligned}$$

for each  $x \in X$ . Next, we prove that  $\text{CoEq}(f, g)$  satisfies the universal property of the coequaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} (X, x_0) & \xrightarrow[\substack{f \\ g}]{} & (Y, y_0) & \xrightarrow{\text{coeq}(f, g)} & (\text{CoEq}(f, g), [y_0]) \\ & & & \searrow c & \\ & & & & (C, *) \end{array}$$

in  $\text{Sets}$ . Then, since  $c(f(a)) = c(g(a))$  for each  $a \in A$ , it follows from [Items 4 and 5 of Proposition 10.6.2.1.3](#) that there exists a unique map  $\phi: \text{CoEq}(f, g) \xrightarrow{\exists!} C$  making the diagram

$$\begin{array}{ccc} (X, x_0) & \xrightarrow[\substack{f \\ g}]{} & (Y, y_0) & \xrightarrow{\text{coeq}(f, g)} & (\text{CoEq}(f, g), [y_0]) \\ & & & \searrow c & \downarrow \phi \exists! \\ & & & & (C, *) \end{array}$$

commute, where we note that  $\phi$  is indeed a morphism of pointed sets since

$$\begin{aligned} \phi([y_0]) &= [\phi \circ \text{coeq}(f, g)]([y_0]) \\ &= c([y_0]) \\ &= *, \end{aligned}$$

where we have used that  $c$  is a morphism of pointed sets. □

00BR

**PROPOSITION 6.3.5.1.4 ► PROPERTIES OF COEQUALISERS OF POINTED SETS**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets and let  $f, g, h: (X, x_0) \rightarrow (Y, y_0)$  be morphisms of pointed sets.

00BS

1. *Associativity.* We have isomorphisms of pointed sets

$$\begin{aligned} \text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h) &\cong \text{CoEq}(f, g, h) \cong \text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g), \\ &= \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h) \end{aligned}$$

where  $\text{CoEq}(f, g, h)$  is the colimit of the diagram

$$(X, x_0) \begin{array}{c} \xrightarrow{f} \\[-1ex] \xrightarrow[g]{\quad} \\[-1ex] \xrightarrow{h} \end{array} (Y, y_0)$$

in  $\text{Sets}_*$ .

00BT

2. *Unitality.* We have an isomorphism of pointed sets

$$\text{CoEq}(f, f) \cong B.$$

00BU

3. *Commutativity.* We have an isomorphism of pointed sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

**PROOF 6.3.5.1.5 ► PROOF OF PROPOSITION 6.3.5.1.4**

**Item 1: Associativity**

This follows from [Item 1 of Proposition 4.2.5.1.7](#).

**Item 2: Unitality**

This follows from [Item 2 of Proposition 4.2.5.1.7](#).

**Item 3: Commutativity**

This follows from [Item 3 of Proposition 4.2.5.1.7](#). 

**00BV 6.4 Constructions With Pointed Sets****00BW 6.4.1 Free Pointed Sets**

Let  $X$  be a set.

00BX

## DEFINITION 6.4.1.1.1 ► FREE POINTED SETS

The **free pointed set on  $X$**  is the pointed set  $X^+$  consisting of:

- *The Underlying Set.* The set  $X^+$  defined by<sup>1</sup>

$$\begin{aligned} X^+ &\stackrel{\text{def}}{=} X \coprod \text{pt} \\ &\stackrel{\text{def}}{=} X \coprod \{\star\}. \end{aligned}$$

- *The Basepoint.* The element  $\star$  of  $X^+$ .

<sup>1</sup>*Further Notation:* We sometimes write  $\star_X$  for the basepoint of  $X^+$  for clarity, especially when there are multiple free pointed sets involved in the current discussion.

00BY

## PROPOSITION 6.4.1.1.2 ► PROPERTIES OF FREE POINTED SETS

Let  $X$  be a set.

00BZ

1. *Functoriality.* The assignment  $X \mapsto X^+$  defines a functor

$$(-)^+: \text{Sets} \rightarrow \text{Sets}_*,$$

where:

- *Action on Objects.* For each  $X \in \text{Obj}(\text{Sets})$ , we have

$$[(-)^+](X) \stackrel{\text{def}}{=} X^+,$$

where  $X^+$  is the pointed set of [Definition 6.4.1.1.1](#).

- *Action on Morphisms.* For each morphism  $f: X \rightarrow Y$  of  $\text{Sets}$ , the image

$$f^+: X^+ \rightarrow Y^+$$

of  $f$  by  $(-)^+$  is the map of pointed sets defined by

$$f^+(x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in X, \\ \star_Y & \text{if } x = \star_X. \end{cases}$$

00C0

2. *Adjointness.* We have an adjunction

$$((-)^+ \dashv \overline{\text{pt}}): \text{Sets} \begin{array}{c} \xrightarrow{(-)^+} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Sets}_*((X^+, \star_X), (Y, y_0)) \cong \text{Sets}(X, Y),$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $(Y, y_0) \in \text{Obj}(\text{Sets}_*)$ .

00C1

3. *Symmetric Strong Monoidality With Respect to Wedge Sums.* The free pointed set functor of [Item 1](#) has a symmetric strong monoidal structure

$$((-)^+, (-)^+, \coprod, (-)_{\mathbb{1}}^+): (\text{Sets}, \coprod, \emptyset) \rightarrow (\text{Sets}_*, \vee, \text{pt}),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^{+, \coprod}: X^+ \vee Y^+ &\xrightarrow{\sim} (X \coprod Y)^+, \\ (-)_{\mathbb{1}}^+: \text{pt} &\xrightarrow{\sim} \emptyset^+, \end{aligned}$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

00C2

4. *Symmetric Strong Monoidality With Respect to Smash Products.* The free pointed set functor of [Item 1](#) has a symmetric strong monoidal structure

$$((-)^+, (-)^+, (-)_{\mathbb{1}}^+): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}_*, \wedge, S^0),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^+: X^+ \wedge Y^+ &\xrightarrow{\sim} (X \times Y)^+, \\ (-)_{\mathbb{1}}^+: S^0 &\xrightarrow{\sim} \text{pt}^+, \end{aligned}$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

#### PROOF 6.4.1.1.3 ► PROOF OF PROPOSITION 6.4.1.1.2

##### Item 1: Functoriality

We claim that  $(-)^+$  is indeed a functor:

- *Preservation of Identities.* Let  $X \in \text{Obj}(\text{Sets})$ . We have

$$\text{id}_X^+(x) \stackrel{\text{def}}{=} \begin{cases} x & \text{if } x \in X, \\ \star_X & \text{if } x = \star_X, \end{cases}$$

for each  $x \in X^+$ , so  $\text{id}_X^+ = \text{id}_{X^+}$ .

- *Preservation of Composition.* Given morphisms of sets

$$\begin{aligned} f: X &\rightarrow Y, \\ g: Y &\rightarrow Z, \end{aligned}$$

we have

$$\begin{aligned}[g^+ \circ f^+](x) &\stackrel{\text{def}}{=} g^+(f^+(x)) \\ &\stackrel{\text{def}}{=} g^+(f(x)) \\ &\stackrel{\text{def}}{=} g(f(x)) \\ &\stackrel{\text{def}}{=} [g \circ f]^+(x)\end{aligned}$$

for each  $x \in X$  and

$$\begin{aligned}[g^+ \circ f^+](\star_X) &\stackrel{\text{def}}{=} g^+(f^+(\star_X)) \\ &\stackrel{\text{def}}{=} g^+(\star_Y) \\ &\stackrel{\text{def}}{=} \star_Z \\ &\stackrel{\text{def}}{=} [g \circ f]^+(\star_X),\end{aligned}$$

so  $(g \circ f)^+ = g^+ \circ f^+$ .

This finishes the proof.

### Item 2: Adjointness

We proceed in a few steps:

025A

- *Map I.* We define a map

$$\Phi_{X,Y}: \text{Sets}_*(X^+, Y) \rightarrow \text{Sets}(X, Y)$$

by sending a morphism of pointed sets

$$\xi: (X^+, \star_X) \rightarrow (Y, y_0)$$

to the function

$$\xi^\dagger: X \rightarrow Y$$

given by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \xi(x)$$

for each  $x \in X$ .

025B

- *Map II.* We define a map

$$\Psi_{X,Y}: \text{Sets}(X, Y) \rightarrow \text{Sets}_*(X^+, Y)$$

given by sending a function  $\xi: X \rightarrow Y$  to the morphism of pointed sets

$$\xi^\dagger: (X^+, \star_X) \rightarrow (Y, y_0)$$

defined by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \begin{cases} \xi(x) & \text{if } x \in X, \\ y_0 & \text{if } x = \star_X \end{cases}$$

for each  $x \in X^+$ .

025C

- *Invertibility I.* Given a morphism of pointed sets

$$\xi: (X^+, \star_X) \rightarrow (Y, y_0),$$

we have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(\xi)) \\ &= \Psi_{X,Y}(\xi^\dagger) \\ &\stackrel{\text{def}}{=} [\![x \mapsto \begin{cases} \xi^\dagger(x) & \text{if } x \in X \\ y_0 & \text{if } x = \star_X \end{cases}]\!] \\ &= [\![x \mapsto \begin{cases} \xi(x) & \text{if } x \in X \\ y_0 & \text{if } x = \star_X \end{cases}]\!] \\ &= \xi \\ &\stackrel{\text{def}}{=} [\text{id}_{\text{Sets}_*(X^+, Y)}](\xi). \end{aligned}$$

Therefore we have

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Sets}_*(X^+, Y)}.$$

025D

- *Invertibility II.* Given a map of sets  $\xi: X \rightarrow Y$ , we have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(\xi)) \\ &= \Phi_{X,Y}(\xi^\dagger) \\ &= \Phi_{X,Y}([\![x \mapsto \begin{cases} \xi(x) & \text{if } x \in X \\ y_0 & \text{if } x = \star_X \end{cases}]\!]) \\ &= [\![x \mapsto \xi(x)]\!] \\ &= \xi \\ &\stackrel{\text{def}}{=} [\text{id}_{\text{Sets}(X, Y)}](\xi). \end{aligned}$$

Therefore we have

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X, Y)}.$$

025E

- *Naturality for  $\Phi$ , Part I.* We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (X', x'_0),$$

the diagram

$$\begin{array}{ccc} \text{Sets}_*(X'^+, Y) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}(X', Y) \\ f^* \downarrow & & \downarrow f^* \\ \text{Sets}_*(X^+, Y) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \end{array}$$

commutes. Indeed, given a morphism of pointed sets  $\xi: X'^+ \rightarrow Y$ , we have

$$\begin{aligned} [\Phi_{X,Y} \circ f^*](\xi) &= \Phi_{X,Y}(f^*(\xi)) \\ &= \Phi_{X,Y}(\xi \circ f) \\ &= \xi \circ f \\ &= \Phi_{X',Y}(\xi) \circ f \\ &= f^*(\Phi_{X',Y}(\xi)) \\ &= f^*(\Phi_{X',Y}(\xi)) \\ &= [f^* \circ \Phi_{X',Y}](\xi). \end{aligned}$$

Therefore we have

$$\Phi_{X,Y} \circ f^* = f^* \circ \Phi_{X',Y}$$

and the naturality diagram for  $\Phi$  above indeed commutes.

025F

- *Naturality for  $\Phi$ , Part II.* We need to show that, given a morphism of pointed sets

$$g: (Y, y_0) \rightarrow (Y', y'_0),$$

the diagram

$$\begin{array}{ccc} \text{Sets}_*(X^+, Y) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\ g_* \downarrow & & \downarrow g_* \\ \text{Sets}_*(X^+, Y'), & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y') \end{array}$$

commutes. Indeed, given a morphism of pointed sets

$$\xi^\dagger: X^+ \rightarrow Y,$$

we have

$$\begin{aligned}
 [\Phi_{X,Y'} \circ g_*](\xi) &= \Phi_{X,Y'}(g_*(\xi)) \\
 &= \Phi_{X,Y'}(g \circ \xi) \\
 &= g \circ \xi \\
 &= g \circ \Phi_{X,Y'}(\xi) \\
 &= g_*(\Phi_{X,Y'}(\xi)) \\
 &= [g_* \circ \Phi_{X,Y'}](\xi).
 \end{aligned}$$

Therefore we have

$$\Phi_{X,Y'} \circ g_* = g_* \circ \Phi_{X,Y'}$$

and the naturality diagram for  $\Phi$  above indeed commutes.

025G

- *Naturality for  $\Psi$ .* Since  $\Phi$  is natural in each argument and  $\Phi$  is a componentwise inverse to  $\Psi$  in each argument, it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\Psi$  is also natural in each argument.

This finishes the proof.

### Item 3: Symmetric Strong Monoidality With Respect to Wedge Sums

We construct the strong monoidal structure on  $(-)^+$  with respect to  $\coprod$  and  $\vee$  as follows:

- *The Strong Monoidality Constraints.* The isomorphism

$$(-)_{X,Y}^{+, \coprod} : X^+ \vee Y^+ \xrightarrow{\sim} (X \coprod Y)^+$$

is given by

$$(-)_{X,Y}^{+, \coprod}(z) = \begin{cases} x & \text{if } z = [(0, x)] \text{ with } x \in X, \\ y & \text{if } z = [(1, y)] \text{ with } y \in Y, \\ \star_X \coprod_Y & \text{if } z = [(0, \star_X)], \\ \star_X \coprod_Y & \text{if } z = [(1, \star_Y)] \end{cases}$$

for each  $z \in X^+ \vee Y^+$ , with inverse

$$(-)_{X,Y}^{+, \coprod, -1} : (X \coprod Y)^+ \xrightarrow{\sim} X^+ \vee Y^+$$

given by

$$(-)_{X,Y}^{+, \coprod, -1}(z) \stackrel{\text{def}}{=} \begin{cases} [(0, x)] & \text{if } z = [(0, x)], \\ [(1, y)] & \text{if } z = [(1, y)], \\ p_0 & \text{if } z = \star_X \coprod_Y \end{cases}$$

for each  $z \in (X \coprod Y)^+$ .

- *The Strong Monoidal Unity Constraint.* The isomorphism

$$(-)_{X,Y}^{+, \coprod, \mathbb{1}} : \text{pt} \xrightarrow{\sim} \emptyset^+$$

is given by sending  $\star_X$  to  $\star_\emptyset$ .

The verification that these isomorphisms satisfy the coherence conditions making the functor  $(-)^+$  into a symmetric strong monoidal functor is omitted.

#### Item 4: Symmetric Strong Monoidality With Respect to Smash Product

We construct the strong monoidal structure on  $(-)^+$  with respect to  $\times$  and  $\wedge$  as follows:

- *The Strong Monoidality Constraints.* The isomorphism

$$(-)_{X,Y}^+ : X^+ \wedge Y^+ \xrightarrow{\sim} (X \times Y)^+$$

is given by

$$(-)_{X,Y}^+(x \wedge y) = \begin{cases} (x, y) & \text{if } x \neq \star_X \text{ and } y \neq \star_Y \\ \star_{X \times Y} & \text{otherwise} \end{cases}$$

for each  $x \wedge y \in X^+ \wedge Y^+$ , with inverse

$$(-)_{X,Y}^{+,-1} : (X \times Y)^+ \xrightarrow{\sim} X^+ \wedge Y^+$$

given by

$$(-)_{X,Y}^{+,-1}(z) \stackrel{\text{def}}{=} \begin{cases} x \wedge y & \text{if } z = (x, y) \text{ with } (x, y) \in X \times Y, \\ \star_X \wedge \star_Y & \text{if } z = \star_{X \times Y}, \end{cases}$$

for each  $z \in (X \times Y)^+$ .

- *The Strong Monoidal Unity Constraint.* The isomorphism

$$(-)_{X,Y}^{+,\mathbb{1}} : S^0 \xrightarrow{\sim} \text{pt}^+$$

is given by sending 0 to  $\star_{\text{pt}}$  and 1 to  $\star$ , where  $\text{pt}^+ = \{\star, \star_{\text{pt}}\}$ .

The verification that these isomorphisms satisfy the coherence conditions making the functor  $(-)^+$  into a symmetric strong monoidal functor is omitted.

**01QS 6.4.2 Deleting Basepoints**

Let  $(X, x_0)$  be a pointed set.

01QT

**DEFINITION 6.4.2.1.1 ► SETS WITH DELETED BASEPOINTS**

The **set with deleted basepoint associated to  $X$**  is the set  $X^-$  defined by

$$X^- \stackrel{\text{def}}{=} X \setminus \{x_0\}.$$

01QU

**PROPOSITION 6.4.2.1.2 ► PROPERTIES OF SETS WITH DELETED BASEPOINTS**

Let  $(X, x_0)$  be a pointed set.

01QV

1. *Functoriality.* The assignment  $(X, x_0) \mapsto X^-$  defines a functor

$$X^- : \text{Sets}_*^{\text{actv}} \rightarrow \text{Sets},$$

where:

- *Action on Objects.* For each  $X \in \text{Obj}(\text{Sets}_*^{\text{actv}})$ , we have

$$[(-)^-](X) \stackrel{\text{def}}{=} X^-,$$

where  $X^-$  is the set of **Definition 6.4.2.1.1**.

- *Action on Morphisms.* For each morphism  $f: X \rightarrow Y$  of  $\text{Sets}_*^{\text{actv}}$ , the image

$$f^-: X^- \rightarrow Y^-$$

of  $f$  by  $(-)^-$  is the map defined by

$$f^-(x) \stackrel{\text{def}}{=} f(x)$$

for each  $x \in X^-$ .

01QW

2. *Adjoint Equivalence.* We have an adjoint equivalence of categories

$$((-)^- \dashv (-)^+): \quad \text{Sets}_*^{\text{actv}} \begin{array}{c} \xrightarrow{(-)^-} \\[-1ex] \xleftarrow{(-)^+} \end{array}_{\perp_{\text{eq}}} \text{Sets},$$

witnessed by a bijection of sets

$$\text{Sets}(X^-, Y) \cong \text{Sets}_*(X, Y^+),$$

natural in  $X \in \text{Obj}(\text{Sets}_*)$  and  $Y \in \text{Obj}(\text{Sets})$ , and by isomorphisms

$$\begin{aligned} (X^-)^+ &\cong X, \\ (Y^+)^- &\cong Y, \end{aligned}$$

once again natural in  $X \in \text{Obj}(\text{Sets}_*)$  and  $Y \in \text{Obj}(\text{Sets})$ .

01QX

3. *Symmetric Strong Monoidality With Respect to Wedge Sums.* The functor of [Item 1](#) has a symmetric strong monoidal structure

$$((-)^-, (-)^{-,\vee}, (-)_{\perp}^{-,\vee}): (\text{Sets}_*^{\text{actv}}, \vee, \text{pt}), \rightarrow (\text{Sets}, \coprod, \emptyset),$$

being equipped with isomorphisms of pointed sets

$$(-)_{X,Y}^{-,\vee}: X^- \coprod Y^- \xrightarrow{\sim} (X \vee Y)^-,$$

$$(-)_{\perp}^{-,\vee}: \emptyset \xrightarrow{\sim} \text{pt}^-,$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

01QY

4. *Symmetric Strong Monoidality With Respect to Smash Products.* The free pointed set functor of [Item 1](#) has a symmetric strong monoidal structure

$$((-)^-, (-)^{-,\times}, (-)_{\perp}^{-,\times}): (\text{Sets}_*^{\text{actv}}, \wedge, S^0), \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with isomorphisms of pointed sets

$$(-)_{X,Y}^-: X^- \times Y^- \xrightarrow{\sim} (X \wedge Y)^-,$$

$$(-)_{\perp}^-: \text{pt} \xrightarrow{\sim} (S^0)^-,$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

#### PROOF 6.4.2.1.3 ► PROOF OF PROPOSITION 6.4.2.1.2

##### Item 1: Functoriality

We claim that  $(-)^-$  is indeed a functor:

- *Preservation of Identities.* Let  $X \in \text{Obj}(\text{Sets})$ . We have

$$\text{id}_X^-(x) \stackrel{\text{def}}{=} x$$

for each  $x \in X^-$ , so  $\text{id}_X^- = \text{id}_{X^-}$ .

- *Preservation of Composition.* Given morphisms of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

$$g: (Y, y_0) \rightarrow (Z, z_0),$$

we have

$$[g^- \circ f^-](x) \stackrel{\text{def}}{=} g^-(f^-(x))$$

$$\begin{aligned} &\stackrel{\text{def}}{=} g^-(f(x)) \\ &\stackrel{\text{def}}{=} g(f(x)) \\ &\stackrel{\text{def}}{=} [g \circ f]^{-}(x) \end{aligned}$$

for each  $x \in X$ , so  $(g \circ f)^- = g^- \circ f^-$ .

This finishes the proof.

**Item 2: Adjoint Equivalence**

We proceed in a few steps:

025H

1. *Map I.* We define a map

$$\Phi_{X,Y}: \text{Sets}(X^-, Y) \rightarrow \text{Sets}_*^{\text{actv}}(X, Y^+)$$

by sending a map  $\xi: X^- \rightarrow Y$  to the active morphism of pointed sets

$$\xi^\dagger: X \rightarrow Y^+$$

given by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \begin{cases} \xi(x) & \text{if } x \in X^-, \\ \star_Y & \text{if } x = x_0, \end{cases}$$

for each  $x \in X$ , where this morphism is indeed active since  $\xi(x) \in Y = Y^+ \setminus \{\star_Y\}$  for all  $x \in X^-$ .

025J

2. *Map II.* We define a map

$$\Psi_{X,Y}: \text{Sets}_*^{\text{actv}}(X, Y^+) \rightarrow \text{Sets}(X^-, Y)$$

given by sending an active morphism of pointed sets  $\xi: X \rightarrow Y^+$  to the map

$$\xi^\dagger: X^- \rightarrow Y$$

defined by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \xi(x)$$

for each  $x \in X^-$ , which is indeed well-defined (in that  $\xi(x) \in Y$  for all  $x \in X^-$ ) since  $\xi$  is active.

025K

3. *Invertibility I.* Given a map of sets  $\xi: X^- \rightarrow Y$ , we have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(\xi)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}([\![x \mapsto \begin{cases} \xi(x) & \text{if } x \in X^-, \\ \star_Y & \text{if } x = x_0 \end{cases}]\!]) \end{aligned}$$

$$\begin{aligned}
&= \llbracket x \mapsto \xi(x) \rrbracket \\
&= \xi \\
&= [\text{id}_{\text{Sets}(X^-, Y)}](\xi).
\end{aligned}$$

Therefore we have

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Sets}(X^-, Y)}.$$

**025L** 4. *Invertibility II.* Given a morphism of pointed sets

$$\xi: (X, x_0) \rightarrow (Y^+, \star_Y),$$

we have

$$\begin{aligned}
[\Phi_{X,Y} \circ \Psi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(\xi)) \\
&= \Phi_{X,Y}(\llbracket x \mapsto \xi(x) \rrbracket) \\
&= \llbracket x \mapsto \begin{cases} \xi(x) & \text{if } x \in X^- \\ \star_Y & \text{if } x = x_0 \end{cases} \rrbracket \\
&= \xi \\
&= [\text{id}_{\text{Sets}_*^{\text{actv}}(X, Y^+)}](\xi).
\end{aligned}$$

Therefore we have

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}_*^{\text{actv}}(X, Y^+)}.$$

**025M** 5. *Naturality for  $\Phi$ , Part I.* We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (X', x'_0),$$

the diagram

$$\begin{array}{ccc}
\text{Sets}(X'^-, Y) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}_*^{\text{actv}}(X', Y^+) \\
f^* \downarrow & & \downarrow f^* \\
\text{Sets}_*(X^-, Y) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}_*^{\text{actv}}(X, Y^+)
\end{array}$$

commutes. Indeed, given a map of sets  $\xi: X' \rightarrow Y$ , we have

$$[\Phi_{X,Y} \circ f^*](\xi) = \Phi_{X,Y}(f^*(\xi))$$

$$\begin{aligned}
&= \Phi_{X,Y}(\xi \circ f) \\
&= [\![x \mapsto \begin{cases} \xi(f(x)) & \text{if } f(x) \in X'^{-} \\ \star_Y & \text{if } f(x) = x'_0 \end{cases}]\!] \\
&= f^*([\![x' \mapsto \begin{cases} \xi(x') & \text{if } x' \in X'^{-} \\ \star_Y & \text{if } x' = x'_0 \end{cases}]\!]) \\
&= f^*(\Phi_{X',Y}(\xi)) \\
&= [f^* \circ \Phi_{X',Y}](\xi).
\end{aligned}$$

Therefore we have

$$\Phi_{X,Y} \circ f^* = f^* \circ \Phi_{X',Y},$$

and the naturality diagram for  $\Phi$  above indeed commutes.

025N

6. *Naturality for  $\Phi$ , Part II.* We need to show that, given a morphism of pointed sets

$$g: (Y, y_0) \rightarrow (Y', y'_0),$$

the diagram

$$\begin{array}{ccc}
\text{Sets}(X^-, Y) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}_*^{\text{actv}}(X, Y^+) \\
g_* \downarrow & & \downarrow g_* \\
\text{Sets}(X^-, Y') & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}_*^{\text{actv}}(X, Y'^+)
\end{array}$$

commutes. Indeed, given a map of sets  $\xi: X^- \rightarrow Y$ , we have

$$\begin{aligned}
[\Phi_{X,Y'} \circ g_*](\xi) &= \Phi_{X,Y'}(g_*(\xi)) \\
&= \Phi_{X,Y'}(g \circ \xi) \\
&= [\![x \mapsto \begin{cases} g(\xi(x)) & \text{if } x \in X^- \\ \star_{Y'} & \text{if } x = x_0 \end{cases}]\!] \\
&= g_*([\![x \mapsto \begin{cases} \xi(x) & \text{if } x \in X^- \\ \star_Y & \text{if } x = x_0 \end{cases}]\!]) \\
&= g_*(\Phi_{X,Y}(\xi)) \\
&= [g_* \circ \Phi_{X,Y}](\xi).
\end{aligned}$$

Therefore we have

$$\Phi_{X,Y'} \circ g_* = g_* \circ \Phi_{X,Y},$$

and the naturality diagram for  $\Phi$  above indeed commutes.

025P

7. *Naturality for  $\Psi$ .* Since  $\Phi$  is natural in each argument and  $\Phi$  is a componentwise inverse to  $\Psi$  in each argument, it follows from [Item 2 of Proposition 11.9.7.1.2](#) that  $\Psi$  is also natural in each argument.

025Q

8. *Fully Faithfulness of  $(-)^-$ .* We aim to show that the assignment  $f \mapsto f^-$  sets up a bijection

$$(-)_{X,Y}^-: \text{Sets}_*^{\text{actv}}(X, Y) \xrightarrow{\sim} \text{Sets}(X^-, Y^-).$$

Indeed, the inverse map

$$(-)_{X,Y}^{-,-1}: \text{Sets}(X^-, Y^-) \xrightarrow{\sim} \text{Sets}_*^{\text{actv}}(X, Y)$$

is given by sending a map of sets  $f: X^- \rightarrow Y^-$  to the active morphism of pointed sets  $f^\dagger: X \rightarrow Y$  defined by

$$f^\dagger(x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in X^-, \\ y_0 & \text{if } x = x_0 \end{cases}$$

for each  $x \in X$ .

025R

9. *Essential Surjectivity of  $(-)^-$ .* We need to show that, given an object  $X \in \text{Obj}(\text{Sets})$ , there exists some  $X' \in \text{Obj}(\text{Sets}_*^{\text{actv}})$  such that  $(X')^- \cong X$ . Indeed, taking  $X' = X^+$ , we have

$$\begin{aligned} (X^+)^- &\stackrel{\text{def}}{=} (X \cup \{\star_X\})^- \\ &\stackrel{\text{def}}{=} (X \cup \{\star_X\}) \setminus \{\star_X\} \\ &= X, \end{aligned}$$

and thus we have in fact an equality  $(X^+)^- = X$ , showing  $(-)^-$  to be essentially surjective.

025S

10. *The Functor  $(-)^-$  Is an Equivalence.* Since  $(-)^-$  is fully faithful and essentially surjective, it is an equivalence by [Item 1 of Proposition 11.6.7.1.2](#).

This finishes the proof.

**Item 3: Symmetric Strong Monoidality With Respect to Wedge Sums**

We construct the strong monoidal structure on  $(-)^-$  with respect to  $\vee$  and  $\coprod$  as follows:

- *The Strong Monoidality Constraints.* The isomorphism

$$(-)_{X,Y}^{-,\vee}: X^- \coprod Y^- \xrightarrow{\sim} (X \vee Y)^-$$

is given by

$$(-)_{X,Y}^{-,\vee}(z) = \begin{cases} [(0,x)] & \text{if } z = (0,x) \text{ with } x \in X, \\ [(1,y)] & \text{if } z = (1,y) \text{ with } y \in Y \end{cases}$$

for each  $z \in X^- \coprod Y^-$ , with inverse

$$(-)_{X,Y}^{-,\vee,-1}: (X \vee Y)^- \xrightarrow{\sim} X^- \coprod Y^-$$

given by

$$(-)_{X,Y}^{-,\vee,-1}(z) \stackrel{\text{def}}{=} \begin{cases} (0,x) & \text{if } z = [(0,x)], \\ (1,y) & \text{if } z = [(1,y)], \end{cases}$$

for each  $z \in (X \vee Y)^-$ .

- *The Strong Monoidal Unity Constraint.* The isomorphism

$$(-)_{X,Y}^{+,\vee,1}: \emptyset \xrightarrow{\sim} \text{pt}^-$$

is an equality.

The verification that these isomorphisms satisfy the coherence conditions making the functor  $(-)^-$  into a symmetric strong monoidal functor is omitted.

#### Item 4: Symmetric Strong Monoidality With Respect to Smash Product

We construct the strong monoidal structure on  $(-)^+$  with respect to  $\wedge$  and  $\times$  as follows:

- *The Strong Monoidality Constraints.* The isomorphism

$$(-)_{X,Y}^-: X^- \times Y^- \xrightarrow{\sim} (X \wedge Y)^-$$

is given by

$$(-)_{X,Y}^-(x, y) = x \wedge y$$

for each  $(x, y) \in X^- \times Y^-$ , with inverse

$$(-)_{X,Y}^{-,-1}: (X \wedge Y)^- \xrightarrow{\sim} X^- \times Y^-$$

given by

$$(-)_{X,Y}^{-,-1}(x \wedge y) \stackrel{\text{def}}{=} (x, y)$$

for each  $x \wedge y \in (X \wedge Y)^-$ .

- *The Strong Monoidal Unity Constraint.* The isomorphism

$$(-)_{X,Y}^{-,1} : \text{pt} \xrightarrow{\sim} (S^0)^-$$

is given by sending  $\star$  to 1.

The verification that these isomorphisms satisfy the coherence conditions making the functor  $(-)^+$  into a symmetric strong monoidal functor is omitted. 

# Appendices

## 6.A Other Chapters

### Preliminaries

1. Introduction
2. A Guide to the Literature

### Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

### Relations

8. Relations
9. Constructions With Relations

### 10. Conditions on Relations

### Categories

### 11. Categories

### 12. Presheaves and the Yoneda Lemma

### Monoidal Categories

### 13. Constructions With Monoidal Categories

### Bicategories

### 14. Types of Morphisms in Bicategories

### Extra Part

### 15. Notes

## Chapter 7

# Tensor Products of Pointed Sets

**02H3** In this chapter we introduce, construct, and study tensor products of pointed sets. The most well-known among these is the *smash product of pointed sets*

$$\wedge : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*,$$

introduced in [Section 7.5.1](#), defined via a universal property as inducing a bijection between the following data:

- Pointed maps  $f: X \wedge Y \rightarrow Z$ .
- Maps of sets  $f: X \times Y \rightarrow Z$  satisfying

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ .

As it turns out, however, dropping either of the *bilinearity* conditions

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

while retaining the other leads to two other tensor products of pointed sets,

$$\begin{aligned} \lhd : \text{Sets}_* \times \text{Sets}_* &\rightarrow \text{Sets}_*, \\ \rhd : \text{Sets}_* \times \text{Sets}_* &\rightarrow \text{Sets}_*, \end{aligned}$$

called the *left* and *right tensor products of pointed sets*. In contrast to  $\wedge$ , which turns out to endow  $\text{Sets}_*$  with a monoidal category structure ([Proposition 7.5.9.1.1](#)), these do not admit invertible associators and unitors, but do endow  $\text{Sets}_*$  with the structure of a skew monoidal category, however ([Propositions 7.3.8.1.1](#) and [7.4.8.1.1](#)).

Finally, in addition to the tensor products  $\lhd$ ,  $\rhd$ , and  $\wedge$ , we also have a

“tensor product” of the form

$$\odot : \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*,$$

called the *tensor* of sets with pointed sets. All in all, these tensor products assemble into a family of functors of the form

$$\begin{aligned}\otimes_{k,\ell} &: \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_\ell}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_{k+\ell}}(\text{Sets}), \\ \triangleleft_{i,k} &: \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_k}(\text{Sets}), \\ \triangleright_{i,k} &: \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_k}(\text{Sets}),\end{aligned}$$

where  $k, \ell, i \in \mathbb{N}$  with  $i \leq k - 1$ . Together with the Cartesian product  $\times$  of Sets, the tensor products studied in this chapter form the cases:

- $(k, \ell) = (-1, -1)$  for the Cartesian product of Sets;
- $(k, \ell) = (0, -1)$  and  $(-1, 0)$  for the tensor of sets with pointed sets of [Definition 7.2.1.1.1](#);
- $(i, k) = (-1, 0)$  for the left and right tensor products of pointed sets of [Sections 7.3 and 7.4](#);
- $(k, \ell) = (-1, -1)$  for the smash product of pointed sets of [Section 7.5](#).

In this chapter, we will carefully define and study bilinearity for pointed sets, as well as all the tensor products described above. Then, in ??, we will extend these to tensor products involving also monoids and commutative monoids, which will end up covering all cases up to  $k, \ell \leq 2$ , and hence *all* cases since  $\mathbb{E}_k$ -monoids on Sets are the same as  $\mathbb{E}_2$ -monoids on Sets when  $k \geq 2$ .

## Contents

---

<b>7.1</b>	<b>Bilinear Morphisms of Pointed Sets .....</b>	<b>386</b>
7.1.1	Left Bilinear Morphisms of Pointed Sets .....	386
7.1.2	Right Bilinear Morphisms of Pointed Sets .....	387
7.1.3	Bilinear Morphisms of Pointed Sets .....	388
<b>7.2</b>	<b>Tensors and Cotensors of Pointed Sets by Sets .....</b>	<b>390</b>
7.2.1	Tensors of Pointed Sets by Sets.....	390
7.2.2	Cotensors of Pointed Sets by Sets.....	398
<b>7.3</b>	<b>The Left Tensor Product of Pointed Sets.....</b>	<b>405</b>
7.3.1	Foundations .....	405
7.3.2	The Left Internal Hom of Pointed Sets .....	411
7.3.3	The Left Skew Unit.....	414
7.3.4	The Left Skew Associator .....	414
7.3.5	The Left Skew Left Unitor .....	417
7.3.6	The Left Skew Right Unitor .....	420

7.3.7	The Diagonal.....	422
7.3.8	The Left Skew Monoidal Structure on Pointed Sets Associated to $\triangleleft$ .....	424
7.3.9	Monoids With Respect to the Left Tensor Product of Pointed Sets.....	428
7.4	<b>The Right Tensor Product of Pointed Sets.....</b>	433
7.4.1	Foundations .....	433
7.4.2	The Right Internal Hom of Pointed Sets .....	438
7.4.3	The Right Skew Unit .....	441
7.4.4	The Right Skew Associator .....	442
7.4.5	The Right Skew Left Unitor .....	445
7.4.6	The Right Skew Right Unitor .....	447
7.4.7	The Diagonal.....	450
7.4.8	The Right Skew Monoidal Structure on Pointed Sets Associated to $\triangleright$ .....	451
7.4.9	Monoids With Respect to the Right Tensor Prod- uct of Pointed Sets .....	455
7.5	<b>The Smash Product of Pointed Sets.....</b>	460
7.5.1	Foundations .....	460
7.5.2	The Internal Hom of Pointed Sets .....	470
7.5.3	The Monoidal Unit.....	473
7.5.4	The Associator .....	473
7.5.5	The Left Unitor .....	475
7.5.6	The Right Unitor .....	478
7.5.7	The Symmetry .....	481
7.5.8	The Diagonal.....	483
7.5.9	The Monoidal Structure on Pointed Sets Associ- ated to $\wedge$ .....	486
7.5.10	The Universal Property of $(\text{Sets}_*, \wedge, S^0)$ .....	491
7.5.11	Monoids With Respect to the Smash Product of Pointed Sets .....	516
7.5.12	Comonoids With Respect to the Smash Product of Pointed Sets.....	517
7.6	<b>Miscellany .....</b>	519
7.6.1	The Smash Product of a Family of Pointed Sets... 519	
7.A	<b>Other Chapters .....</b>	520

## 00C4 7.1 Bilinear Morphisms of Pointed Sets

### 00C5 7.1.1 Left Bilinear Morphisms of Pointed Sets

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

00C6

## DEFINITION 7.1.1.1.1 ► LEFT BILINEAR MORPHISMS OF POINTED SETS

A **left bilinear morphism of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is a map of sets

$$f: X \times Y \rightarrow Z$$

satisfying the following condition:<sup>1,2</sup>

(★) *Left Unital Bilinearity.* The diagram

$$\begin{array}{ccc} & \text{pt} \times \text{pt} & \\ \text{id}_{\text{pt}} \times \epsilon_Y \nearrow & \searrow \curvearrowright & \\ \text{pt} \times Y & & \text{pt} \\ \downarrow [x_0] \times \text{id}_Y & & \downarrow [z_0] \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commutes, i.e. for each  $y \in Y$ , we have

$$f(x_0, y) = z_0.$$

<sup>1</sup>*Slogan:* The map  $f$  is left bilinear if it preserves basepoints in its first argument.

<sup>2</sup>Succinctly,  $f$  is bilinear if we have

$$f(x_0, y) = z_0$$

for each  $y \in Y$ .

00C7

## DEFINITION 7.1.1.1.2 ► THE SET OF LEFT BILINEAR MORPHISMS OF POINTED SETS

The **set of left bilinear morphisms of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is the set  $\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z)$  defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is left bilinear}\}.$$

00C8 7.1.2 Right Bilinear Morphisms of Pointed Sets

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

00C9

## DEFINITION 7.1.2.1.1 ► RIGHT BILINEAR MORPHISMS OF POINTED SETS

A **right bilinear morphism of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is a map of sets

$$f: X \times Y \rightarrow Z$$

satisfying the following condition:<sup>1,2</sup>

(★) *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \epsilon_X \times \text{id}_{\text{pt}} \nearrow & \searrow \curvearrowright & \\
 X \times \text{pt} & & \text{pt} \\
 \text{id}_X \times [y_0] \searrow & & \swarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each  $x \in X$ , we have

$$f(x, y_0) = z_0.$$

<sup>1</sup>Slogan: The map  $f$  is right bilinear if it preserves basepoints in its second argument.

<sup>2</sup>Succinctly,  $f$  is bilinear if we have

$$f(x, y_0) = z_0$$

for each  $x \in X$ .

00CA

#### DEFINITION 7.1.2.1.2 ► THE SET OF RIGHT BILINEAR MORPHISMS OF POINTED SETS

The **set of right bilinear morphisms of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is the set  $\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z)$  defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is right bilinear}\}.$$

00CB

#### 7.1.3 BILINEAR MORPHISMS OF POINTED SETS

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

00CC

#### DEFINITION 7.1.3.1.1 ► BILINEAR MORPHISMS OF POINTED SETS

A **bilinear morphism of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is a map of sets

$$f: X \times Y \rightarrow Z$$

that is both left bilinear and right bilinear.

00CD

#### REMARK 7.1.3.1.2 ► UNWINDING DEFINITION 7.1.3.1.1

In detail, a **bilinear morphism of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

satisfying the following conditions:<sup>1,2</sup>

025T

1. *Left Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \text{id}_{\text{pt}} \times \epsilon_Y \nearrow & \searrow \text{dashed} \curvearrowright & \\
 \text{pt} \times Y & & \text{pt} \\
 \downarrow [x_0] \times \text{id}_Y & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each  $y \in Y$ , we have

$$f(x_0, y) = z_0.$$

025U

2. *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \epsilon_X \times \text{id}_{\text{pt}} \nearrow & \searrow \text{dashed} \curvearrowright & \\
 X \times \text{pt} & & \text{pt} \\
 \downarrow \text{id}_X \times [y_0] & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each  $x \in X$ , we have

$$f(x, y_0) = z_0.$$

<sup>1</sup>*Slogan:* The map  $f$  is bilinear if it preserves basepoints in each argument.

<sup>2</sup>Succinctly,  $f$  is bilinear if we have

$$\begin{aligned}
 f(x_0, y) &= z_0, \\
 f(x, y_0) &= z_0
 \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ .

00CE

#### DEFINITION 7.1.3.1.3 ► THE SET OF BILINEAR MORPHISMS OF POINTED SETS

The **set of bilinear morphisms of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is the set  $\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$  defined by

$$\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is bilinear}\}.$$

## 00CF 7.2 Tensors and Cotensors of Pointed Sets by Sets

### 00CG 7.2.1 Tensors of Pointed Sets by Sets

Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

00CH

#### DEFINITION 7.2.1.1.1 ► TENSORS OF POINTED SETS BY SETS

The **tensor of**  $(X, x_0)$  **by**  $A$ <sup>1</sup> is the tensor  $A \odot (X, x_0)$ <sup>2</sup> of  $(X, x_0)$  by  $A$  as in ??.

<sup>1</sup>Further Terminology: Also called the **copower of**  $(X, x_0)$  **by**  $A$ .

<sup>2</sup>Further Notation: Often written  $A \odot X$  for simplicity.

00CJ

#### REMARK 7.2.1.2 ► UNWINDING DEFINITION 7.2.1.1.1

In detail, the **tensor of**  $(X, x_0)$  **by**  $A$  is the pointed set  $A \odot (X, x_0)$  satisfying the following universal property:

(★) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K)),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ .

This universal property is in turn equivalent to the following one:

(★) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}_{\mathbb{E}_0}^\otimes(A \times X, K),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ , where  $\text{Sets}_{\mathbb{E}_0}^\otimes(A \times X, K)$  is the set defined by

$$\text{Sets}_{\mathbb{E}_0}^\otimes(A \times X, K) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times X, K) \mid \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } f(a, x_0) = k_0 \end{array} \right\}.$$

#### PROOF 7.2.1.1.3 ► PROOF OF THE EQUIVALENCE IN REMARK 7.2.1.1.2

We claim that we have a bijection

$$\text{Sets}(A, \text{Sets}_*(X, K)) \cong \text{Sets}_{\mathbb{E}_0}^\otimes(A \times X, K)$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ . Indeed, this bijection is a restriction of the bijection

$$\text{Sets}(A, \text{Sets}(X, K)) \cong \text{Sets}(A \times X, K)$$

of Item 2 of Proposition 4.1.3.1.4:

- A map

$$\xi: A \longrightarrow \text{Sets}_*(X, K),$$

$$a \mapsto (\xi_a: X \rightarrow K),$$

in  $\text{Sets}(A, \text{Sets}_*(X, K))$  gets sent to the map

$$\xi^\dagger: A \times X \rightarrow K$$

defined by

$$\xi^\dagger(a, x) \stackrel{\text{def}}{=} \xi_a(x)$$

for each  $(a, x) \in A \times X$ , which indeed lies in  $\text{Sets}_{\mathbb{E}_0}^\otimes(A \times X, K)$ , as we have

$$\begin{aligned} \xi^\dagger(a, x_0) &\stackrel{\text{def}}{=} \xi_a(x_0) \\ &\stackrel{\text{def}}{=} k_0 \end{aligned}$$

for each  $a \in A$ , where we have used that  $\xi_a \in \text{Sets}_*(X, K)$  is a morphism of pointed sets.

- Conversely, a map

$$\xi: A \times X \rightarrow K$$

in  $\text{Sets}_{\mathbb{E}_0}^\otimes(A \times X, K)$  gets sent to the map

$$\xi^\dagger: A \longrightarrow \text{Sets}_*(X, K),$$

$$a \mapsto (\xi_a^\dagger: X \rightarrow K),$$

where

$$\xi_a^\dagger: X \rightarrow K$$

is the map defined by

$$\xi_a^\dagger(x) \stackrel{\text{def}}{=} \xi(a, x)$$

for each  $x \in X$ , and indeed lies in  $\text{Sets}_*(X, K)$ , as we have

$$\begin{aligned} \xi_a^\dagger(x_0) &\stackrel{\text{def}}{=} \xi(a, x_0) \\ &\stackrel{\text{def}}{=} k_0. \end{aligned}$$

This finishes the proof. □

00CK

## CONSTRUCTION 7.2.1.1.4 ► CONSTRUCTION OF TENSORS OF POINTED SETS BY SETS

Concretely, the **tensor of**  $(X, x_0)$  **by**  $A$  is the pointed set  $A \odot (X, x_0)$  consisting of:

- *The Underlying Set.* The set  $A \odot X$  given by

$$A \odot X \cong \bigvee_{a \in A} (X, x_0),$$

where  $\bigvee_{a \in A} (X, x_0)$  is the wedge product of the  $A$ -indexed family  $((X, x_0))_{a \in A}$  of Definition 6.3.2.1.1.

- *The Basepoint.* The point  $[(a, x_0)] = [(a', x_0)]$  of  $\bigvee_{a \in A} (X, x_0)$ .

## PROOF 7.2.1.1.5 ► PROOF OF CONSTRUCTION 7.2.1.1.4

(Proven below in a bit.)



00CL

## NOTATION 7.2.1.1.6 ► ELEMENTS OF TENSORS OF POINTED SETS BY SETS

We write  $a \odot x$  for the element  $[(a, x)]$  of

$$\begin{aligned} A \odot X &\cong \bigvee_{a \in A} (X, x_0) \\ &\stackrel{\text{def}}{=} (\coprod_{i \in I} X_i) / \sim. \end{aligned}$$

00CM

## REMARK 7.2.1.1.7 ► BASEPOINTS OF TENSORS OF POINTED SETS BY SETS

Taking the tensor of any element of  $A$  with the basepoint  $x_0$  of  $X$  leads to the same element in  $A \odot X$ , i.e. we have

$$a \odot x_0 = a' \odot x_0,$$

for each  $a, a' \in A$ . This is due to the equivalence relation  $\sim$  on

$$\bigvee_{a \in A} (X, x_0) \stackrel{\text{def}}{=} \coprod_{a \in A} X / \sim$$

identifying  $(a, x_0)$  with  $(a', x_0)$ , so that the equivalence class  $a \odot x_0$  is independent from the choice of  $a \in A$ .

**PROOF 7.2.1.8 ► PROOF OF CONSTRUCTION 7.2.1.4**

We claim we have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K))$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ .

**025V**

1. *Map I.* We define a map

$$\Phi_K : \text{Sets}_*(A \odot X, K) \rightarrow \text{Sets}(A, \text{Sets}_*(X, K))$$

by sending a morphism of pointed sets

$$\xi : (A \odot X, a \odot x_0) \rightarrow (K, k_0)$$

to the map of sets

$$\begin{aligned} \xi^\dagger : A &\longrightarrow \text{Sets}_*(X, K), \\ a &\mapsto (\xi_a : X \rightarrow K), \end{aligned}$$

where

$$\xi_a : (X, x_0) \rightarrow (K, k_0)$$

is the morphism of pointed sets defined by

$$\xi_a(x) \stackrel{\text{def}}{=} \xi(a \odot x)$$

for each  $x \in X$ . Note that we have

$$\begin{aligned} \xi_a(x_0) &\stackrel{\text{def}}{=} \xi(a \odot x_0) \\ &= k_0, \end{aligned}$$

so that  $\xi_a$  is indeed a morphism of pointed sets, where we have used that  $\xi$  is a morphism of pointed sets.

**025W**

2. *Map II.* We define a map

$$\Psi_K : \text{Sets}(A, \text{Sets}_*(X, K)) \rightarrow \text{Sets}_*(A \odot X, K)$$

given by sending a map

$$\begin{aligned} \xi : A &\longrightarrow \text{Sets}_*(X, K), \\ a &\mapsto (\xi_a : X \rightarrow K), \end{aligned}$$

to the morphism of pointed sets

$$\xi^\dagger: (A \odot X, a \odot x_0) \rightarrow (K, k_0)$$

defined by

$$\xi^\dagger(a \odot x) \stackrel{\text{def}}{=} \xi_a(x)$$

for each  $a \odot x \in A \odot X$ . Note that  $\xi^\dagger$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \xi^\dagger(a \odot x_0) &\stackrel{\text{def}}{=} \xi_a(x_0) \\ &= k_0, \end{aligned}$$

where we have used that  $\xi(a) \in \text{Sets}_*(X, K)$  is a morphism of pointed sets.

025X

3. *Invertibility I.* We claim that

$$\Psi_K \circ \Phi_K = \text{id}_{\text{Sets}_*(A \odot X, K)}.$$

Indeed, given a morphism of pointed sets

$$\xi: (A \odot X, a \odot x_0) \rightarrow (K, k_0),$$

we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= \Psi_K(\Phi_K(\xi)) \\ &= \Psi_K(\llbracket a \mapsto \llbracket x \mapsto \xi(a \odot x) \rrbracket \rrbracket) \\ &= \Psi_K(\llbracket a' \mapsto \llbracket x' \mapsto \xi(a' \odot x') \rrbracket \rrbracket) \\ &= \llbracket a \odot x \mapsto \text{ev}_x(\text{ev}_a(\llbracket a' \mapsto \llbracket x' \mapsto \xi(a' \odot x') \rrbracket \rrbracket)) \rrbracket \\ &= \llbracket a \odot x \mapsto \text{ev}_x(\llbracket x' \mapsto \xi(a \odot x') \rrbracket) \rrbracket \\ &= \llbracket a \odot x \mapsto \xi(a \odot x) \rrbracket \\ &= \xi. \end{aligned}$$

025Y

4. *Invertibility II.* We claim that

$$\Phi_K \circ \Psi_K = \text{id}_{\text{Sets}(A, \text{Sets}_*(X, K))}.$$

Indeed, given a morphism  $\xi: A \rightarrow \text{Sets}_*(X, K)$ , we have

$$\begin{aligned} [\Phi_K \circ \Psi_K](\xi) &= \Phi_K(\Psi_K(\xi)) \\ &= \Phi_K(\llbracket a \odot x \mapsto \xi_a(x) \rrbracket) \\ &= \llbracket a \mapsto \llbracket x \mapsto \xi_a(x) \rrbracket \rrbracket \\ &= \llbracket a \mapsto \xi(a) \rrbracket \\ &= \xi. \end{aligned}$$

025Z

5. *Naturality of  $\Phi$ .* We need to show that, given a morphism of pointed sets

$$\phi: (K, k_0) \rightarrow (K', k'_0),$$

the diagram

$$\begin{array}{ccc} \text{Sets}_*(A \odot X, K) & \xrightarrow{\Phi_K} & \text{Sets}(A, \text{Sets}_*(X, K)) \\ \phi_* \downarrow & & \downarrow (\phi_*)_* \\ \text{Sets}_*(A \odot X, K') & \xrightarrow{\Phi_{K'}} & \text{Sets}(A, \text{Sets}_*(X, K')) \end{array}$$

commutes. Indeed, given a morphism of pointed sets

$$\xi: (A \odot X, a \odot x_0) \rightarrow (K, k_0),$$

we have

$$\begin{aligned} [\Phi_{K'} \circ \phi_*](\xi) &= \Phi_{K'}(\phi_*(\xi)) \\ &= \Phi_{K'}(\phi \circ \xi) \\ &= (\phi \circ \xi)^\dagger \\ &= \llbracket a \mapsto \phi \circ \xi(a \odot -) \rrbracket \\ &= \llbracket a \mapsto \phi_*(\xi(a \odot -)) \rrbracket \\ &= (\phi_*)_*(\llbracket a \mapsto \xi(a \odot -) \rrbracket) \\ &= (\phi_*)_*(\Phi_K(\xi)) \\ &= [(\phi_*)_* \circ \Phi_K](\xi). \end{aligned}$$

0260

6. *Naturality of  $\Psi$ .* Since  $\Phi$  is natural and  $\Phi$  is a componentwise inverse to  $\Psi$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\Psi$  is also natural.

This finishes the proof. □

00CN

#### PROPOSITION 7.2.1.9 ▶ PROPERTIES OF TENSORS OF POINTED SETS BY SETS

Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

00CP

1. *Functionality.* The assignments  $A, (X, x_0), (A, (X, x_0))$  define functors

$$\begin{aligned} A \odot -: &\quad \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \odot X: &\quad \text{Sets} \rightarrow \text{Sets}_*, \\ -_1 \odot -_2: &\quad \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

In particular, given:

- A map of sets  $f: A \rightarrow B$ ;
- A pointed map  $\phi: (X, x_0) \rightarrow (Y, y_0)$ ;

the induced map

$$f \odot \phi: A \odot X \rightarrow B \odot Y$$

is given by

$$[f \odot \phi](a \odot x) \stackrel{\text{def}}{=} f(a) \odot \phi(x)$$

for each  $a \odot x \in A \odot X$ .

**00CQ** 2. *Adjointness I.* We have an adjunction

$$(- \odot X \dashv \text{Sets}_*(X, -)): \text{Sets} \begin{array}{c} \xrightarrow{- \odot X} \\ \perp \\ \xleftarrow{\text{Sets}_*(X, -)} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K)),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

**00CR** 3. *Adjointness II.* We have an adjunctions

$$(A \odot - \dashv A \pitchfork -): \text{Sets}_* \begin{array}{c} \xrightarrow{A \odot -} \\ \perp \\ \xleftarrow{A \pitchfork -} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Hom}_{\text{Sets}_*}(A \odot X, Y) \cong \text{Hom}_{\text{Sets}_*}(X, A \pitchfork Y),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

**00CS** 4. *As a Weighted Colimit.* We have

$$A \odot X \cong \text{colim}^{[A]}(X),$$

where in the right hand side we write:

- $A$  for the functor  $A: \text{pt} \rightarrow \text{Sets}$  picking  $A \in \text{Obj}(\text{Sets})$ ;
- $X$  for the functor  $X: \text{pt} \rightarrow \text{Sets}_*$  picking  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00CT

5. *Iterated Tensors.* We have an isomorphism of pointed sets

$$A \odot (B \odot X) \cong (A \times B) \odot X,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$  and  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00CU

6. *Interaction With Hom.* We have a natural isomorphism

$$\text{Sets}_*(A \odot X, -) \cong A \pitchfork \text{Sets}_*(X, -).$$

00CV

7. *The Tensor Evaluation Map.* For each  $X, Y \in \text{Obj}(\text{Sets}_*)$ , we have a map

$$\text{ev}_{X,Y}^\odot : \text{Sets}_*(X, Y) \odot X \rightarrow Y,$$

natural in  $X, Y \in \text{Obj}(\text{Sets}_*)$ , and given by

$$\text{ev}_{X,Y}^\odot(f \odot x) \stackrel{\text{def}}{=} f(x)$$

for each  $f \odot x \in \text{Sets}_*(X, Y) \odot X$ .

00CW

8. *The Tensor Coevaluation Map.* For each  $A \in \text{Obj}(\text{Sets})$  and each  $X \in \text{Obj}(\text{Sets}_*)$ , we have a map

$$\text{coev}_{A,X}^\odot : A \rightarrow \text{Sets}_*(X, A \odot X),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X \in \text{Obj}(\text{Sets}_*)$ , and given by

$$\text{coev}_{A,X}^\odot(a) \stackrel{\text{def}}{=} \llbracket x \mapsto a \odot x \rrbracket$$

for each  $a \in A$ .

#### PROOF 7.2.1.1.10 ▶ PROOF OF PROPOSITION 7.2.1.1.9

**Item 1: Functoriality**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 2: Adjointness I**

This is simply a rephrasing of [Definition 7.2.1.1.1](#).

**Item 3: Adjointness II**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 4: As a Weighted Colimit**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

## Item 5: Iterated Tensors

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

## Item 6: Interaction With Hom

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

## Item 7: The Tensor Evaluation Map

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

## Item 8: The Tensor Coevaluation Map

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ . 

**00CX 7.2.2 Cotensors of Pointed Sets by Sets**

Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

00CY

**DEFINITION 7.2.2.1.1 ► COTENSORS OF POINTED SETS BY SETS**

The **cotensor of  $(X, x_0)$  by  $A$** <sup>1</sup> is the cotensor  $A \pitchfork (X, x_0)$ <sup>2</sup> of  $(X, x_0)$  by  $A$  as in ??.

<sup>1</sup>Further Terminology: Also called the **power of  $(X, x_0)$  by  $A$** .

<sup>2</sup>Further Notation: Often written  $A \pitchfork X$  for simplicity.

00CZ

**REMARK 7.2.2.1.2 ► UNWINDING DEFINITION 7.2.2.1.1**

In detail, the **cotensor of  $(X, x_0)$  by  $A$**  is the pointed set  $A \pitchfork (X, x_0)$  satisfying the following universal property:

(★) We have a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}(A, \text{Sets}_*(K, X)),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ .

This universal property is in turn equivalent to the following one:

(★) We have a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}_{\mathbb{E}_0}^\otimes(A \times K, X),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ , where  $\text{Sets}_{\mathbb{E}_0}^\otimes(A \times K, X)$  is the set defined by

$$\text{Sets}_{\mathbb{E}_0}^\otimes(A \times K, X) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times K, X) \mid \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } f(a, k_0) = x_0 \end{array} \right\}.$$

**PROOF 7.2.2.1.3 ► PROOF OF THE EQUIVALENCE IN REMARK 7.2.2.1.2**

This follows from the bijection

$$\text{Sets}(A, \text{Sets}_*(K, X)) \cong \text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$  constructed in the proof of [Remark 7.2.1.1.2](#).



00D0

**CONSTRUCTION 7.2.2.1.4 ► CONSTRUCTION OF COTENSORS OF POINTED SETS BY SETS**

Concretely, the **cotensor of**  $(X, x_0)$  **by**  $A$  is the pointed set  $A \pitchfork (X, x_0)$  consisting of:

- *The Underlying Set.* The set  $A \pitchfork X$  given by

$$A \pitchfork X \cong \bigwedge_{a \in A} (X, x_0),$$

where  $\bigwedge_{a \in A} (X, x_0)$  is the smash product of the  $A$ -indexed family  $((X, x_0))_{a \in A}$  of [Definition 7.6.1.1.1](#).

- *The Basepoint.* The point  $[(x_0)_{a \in A}] = [(x_0, x_0, x_0, \dots)]$  of  $\bigwedge_{a \in A} (X, x_0)$ .

**PROOF 7.2.2.1.5 ► PROOF OF CONSTRUCTION 7.2.2.1.4**

We claim we have a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}(A, \text{Sets}_*(K, X)),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ .

0261

1. *Map I.* We define a map

$$\Phi_K : \text{Sets}_*(K, A \pitchfork X) \rightarrow \text{Sets}(A, \text{Sets}_*(K, X)),$$

by sending a morphism of pointed sets

$$\xi : (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

to the map of sets

$$\begin{aligned} \xi^\dagger : A &\longrightarrow \text{Sets}_*(K, X), \\ a &\mapsto (\xi_a : K \rightarrow X), \end{aligned}$$

where

$$\xi_a: (K, k_0) \rightarrow (X, x_0)$$

is the morphism of pointed sets defined by

$$\xi_a(k) = \begin{cases} x_a^k & \text{if } \xi(k) \neq [(x_0)_{a \in A}], \\ x_0 & \text{if } \xi(k) = [(x_0)_{a \in A}] \end{cases}$$

for each  $k \in K$ , where  $x_a^k$  is the  $a$ th component of  $\xi(k) = [(x_a^k)_{a \in A}]$ . Note that:

0262

- (a) The definition of  $\xi_a(k)$  is independent of the choice of equivalence class. Indeed, suppose we have

$$\begin{aligned} \xi(k) &= [(x_a^k)_{a \in A}] \\ &= [(y_a^k)_{a \in A}] \end{aligned}$$

with  $x_a^k \neq y_a^k$  for some  $a \in A$ . Then there exist  $a_x, a_y \in A$  such that  $x_{a_x}^k = y_{a_y}^k = x_0$ . The equivalence relation  $\sim$  on  $\prod_{a \in A} X$  then forces

$$\begin{aligned} [(x_a^k)_{a \in A}] &= [(x_0)_{a \in A}], \\ [(y_a^k)_{a \in A}] &= [(x_0)_{a \in A}], \end{aligned}$$

however, and  $\xi_a(k)$  is defined to be  $x_0$  in this case.

0263

- (b) The map  $\xi_a$  is indeed a morphism of pointed sets, as we have

$$\xi_a(k_0) = x_0$$

since  $\xi(k_0) = [(x_0)_{a \in A}]$  as  $\xi$  is a morphism of pointed sets and  $\xi_a(k_0)$ , defined to be the  $a$ th component of  $[(x_0)_{a \in A}]$ , is equal to  $x_0$ .

0264

2. *Map II.* We define a map

$$\Psi_K: \text{Sets}(A, \text{Sets}_*(K, X)) \rightarrow \text{Sets}_*(K, A \pitchfork X),$$

given by sending a map

$$\begin{aligned} \xi: A &\longrightarrow \text{Sets}_*(K, X), \\ a &\mapsto (\xi_a: K \rightarrow X), \end{aligned}$$

to the morphism of pointed sets

$$\xi^\dagger: (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

defined by

$$\xi^\dagger(k) \stackrel{\text{def}}{=} [(\xi_a(k))_{a \in A}]$$

for each  $k \in K$ . Note that  $\xi^\dagger$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \xi^\dagger(k_0) &\stackrel{\text{def}}{=} [(\xi_a(k_0))_{a \in A}] \\ &= x_0, \end{aligned}$$

where we have used that  $\xi_a \in \text{Sets}_*(K, X)$  is a morphism of pointed sets for each  $a \in A$ .

0265

3. *Invertibility I.* We claim that

$$\Psi_K \circ \Phi_K = \text{id}_{\text{Sets}_*(K, A \pitchfork X)}.$$

Indeed, given a morphism of pointed sets

$$\xi: (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= \Psi_K(\Phi_K(\xi)) \\ &= \Psi_K([\![a \mapsto \xi_a]\!]) \\ &= \Psi_K([\![a' \mapsto \xi_{a'}]\!]) \\ &= [\![k \mapsto ([\text{ev}_a([\![a' \mapsto \xi_{a'}(k)]\!])]_{a \in A})]\!] \\ &= [\![k \mapsto [(\xi_a(k))_{a \in A}]]\!]. \end{aligned}$$

Now, we have two cases:

0266

(a) If  $\xi(k) = [(x_0)_{a \in A}]$ , we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= [\![k \mapsto [(\xi_a(k))_{a \in A}]]\!] \\ &= [\![k \mapsto [(x_0)_{a \in A}]]\!] \\ &= [\![k \mapsto \xi(k)]\!] \\ &= \xi. \end{aligned}$$

0267

(b) If  $\xi(k) \neq [(x_0)_{a \in A}]$  and  $\xi(k) = [(x_a^k)_{a \in A}]$  instead, we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= [\![k \mapsto [(\xi_a(k))_{a \in A}]]\!] \\ &= [\![k \mapsto [(x_a^k)_{a \in A}]]\!] \\ &= [\![k \mapsto \xi(k)]\!] \\ &= \xi. \end{aligned}$$

In both cases, we have  $[\Psi_K \circ \Phi_K](\xi) = \xi$ , and thus we are done.

0268

4. *Invertibility II.* We claim that

$$\Phi_K \circ \Psi_K = \text{id}_{\text{Sets}(A, \text{Sets}_*(K, X))}.$$

Indeed, given a morphism  $\xi: A \rightarrow \text{Sets}_*(K, X)$ , we have

$$\begin{aligned} [\Phi_K \circ \Psi_K](\xi) &= \Phi_K(\Psi_K(\xi)) \\ &= \Phi_K([\![k \mapsto [(\xi_a(k))_{a \in A}]\!]) \\ &= [\![a \mapsto [\![k \mapsto \xi_a(k)]\!]]\!] \\ &= \xi \end{aligned}$$

0269

5. *Naturality of  $\Psi$ .* We need to show that, given a morphism of pointed sets

$$\phi: (K, k_0) \rightarrow (K', k'_0),$$

the diagram

$$\begin{array}{ccc} \text{Sets}(A, \text{Sets}_*(K', X)) & \xrightarrow{\Psi_{K'}} & \text{Sets}_*(K', A \pitchfork X) \\ (\phi^*)_* \downarrow & & \downarrow \phi^* \\ \text{Sets}(A, \text{Sets}_*(K, X)) & \xrightarrow{\Psi_K} & \text{Sets}_*(K, A \pitchfork X) \end{array}$$

commutes. Indeed, given a map of sets

$$\begin{aligned} \xi: A &\longrightarrow \text{Sets}_*(K', X), \\ a &\mapsto (\xi_a: K' \rightarrow X), \end{aligned}$$

we have

$$\begin{aligned} [\Psi_K \circ (\phi^*)_*](\xi) &= \Psi_K((\phi^*)_*(\xi)) \\ &= \Psi_K((\phi^*)_*([\![a \mapsto \xi_a]\!])) \\ &= \Psi_K([\![a \mapsto \phi^*(\xi_a)]\!]) \\ &= \Psi_K([\![a \mapsto [\![k \mapsto \xi_a(\phi(k))]\!]]\!]) \\ &= [\![k \mapsto [(\xi_a(\phi(k)))_{a \in A}]\!]] \\ &= \phi^*([\![k' \mapsto [(\xi_a(k'))_{a \in A}]\!]]\!) \\ &= \phi^*(\Psi_{K'}(\xi)) \\ &= [\phi^* \circ \Psi_{K'}](\xi). \end{aligned}$$

026A

6. *Naturality of  $\Phi$ .* Since  $\Psi$  is natural and  $\Psi$  is a componentwise inverse to  $\Phi$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\Phi$  is also natural.

This finishes the proof. 

00D1

#### PROPOSITION 7.2.2.1.6 ► PROPERTIES OF COTENSORS OF POINTED SETS BY SETS

Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

00D2

1. *Functionality.* The assignments  $A, (X, x_0), (A, (X, x_0))$  define functors

$$\begin{aligned} A \pitchfork - & : \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \pitchfork X & : \text{Sets}^{\text{op}} \rightarrow \text{Sets}_*, \\ -_1 \pitchfork -_2 & : \text{Sets}^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

In particular, given:

- A map of sets  $f: A \rightarrow B$ ;
- A pointed map  $\phi: (X, x_0) \rightarrow (Y, y_0)$ ;

the induced map

$$f \odot \phi: A \pitchfork X \rightarrow B \pitchfork Y$$

is given by

$$[f \odot \phi]([(x_a)_{a \in A}]) \stackrel{\text{def}}{=} [(\phi(x_{f(a)}))_{a \in A}]$$

for each  $[(x_a)_{a \in A}] \in A \pitchfork X$ .

00D3

2. *Adjointness I.* We have an adjunction

$$(- \pitchfork X \dashv \text{Sets}_*(-, X)): \text{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{- \pitchfork X} \\ \perp \\ \xleftarrow{\text{Sets}_*(-, X)} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Sets}_*^{\text{op}}(A \pitchfork X, K) \cong \text{Sets}(A, \text{Sets}_*(K, X)),$$

i.e. by a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}(A, \text{Sets}_*(K, X)),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

00D4

3. *Adjointness II.* We have an adjunctions

$$(A \odot - \dashv A \pitchfork -): \text{Sets}_* \begin{array}{c} \xrightarrow{A \odot -} \\ \perp \\ \xleftarrow{A \pitchfork -} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Hom}_{\text{Sets}_*}(A \odot X, Y) \cong \text{Hom}_{\text{Sets}_*}(X, A \pitchfork Y),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

00D5

4. *As a Weighted Limit.* We have

$$A \pitchfork X \cong \lim^{[A]}(X),$$

where in the right hand side we write:

- $A$  for the functor  $A: \text{pt} \rightarrow \text{Sets}$  picking  $A \in \text{Obj}(\text{Sets})$ ;
- $X$  for the functor  $X: \text{pt} \rightarrow \text{Sets}_*$  picking  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00D6

5. *Iterated Cotensors.* We have an isomorphism of pointed sets

$$A \pitchfork (B \pitchfork X) \cong (A \times B) \pitchfork X,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$  and  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00D7

6. *Commutativity With Homs.* We have natural isomorphisms

$$A \pitchfork \text{Sets}_*(X, -) \cong \text{Sets}_*(A \odot X, -),$$

$$A \pitchfork \text{Sets}_*(-, Y) \cong \text{Sets}_*(-, A \pitchfork Y).$$

00D8

7. *The Cotensor Evaluation Map.* For each  $X, Y \in \text{Obj}(\text{Sets}_*)$ , we have a map

$$\text{ev}_{X,Y}^\pitchfork: X \rightarrow \text{Sets}_*(X, Y) \pitchfork Y,$$

natural in  $X, Y \in \text{Obj}(\text{Sets}_*)$ , and given by

$$\text{ev}_{X,Y}^\pitchfork(x) \stackrel{\text{def}}{=} [(f(x))_{f \in \text{Sets}_*(X,Y)}]$$

for each  $x \in X$ .

00D9

8. *The Cotensor Coevaluation Map.* For each  $X \in \text{Obj}(\text{Sets}_*)$  and each  $A \in \text{Obj}(\text{Sets})$ , we have a map

$$\text{coev}_{A,X}^\pitchfork: A \rightarrow \text{Sets}_*(A \pitchfork X, X),$$

natural in  $X \in \text{Obj}(\text{Sets}_*)$  and  $A \in \text{Obj}(\text{Sets})$ , and given by

$$\text{coev}_{A,X}^\pitchfork(a) \stackrel{\text{def}}{=} [[(x_b)_{b \in A}] \mapsto x_a]]$$

for each  $a \in A$ .

**PROOF 7.2.2.1.7 ► PROOF OF PROPOSITION 7.2.2.1.6****Item 1: Functoriality**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 2: Adjointness I**

This is simply a rephrasing of [Definition 7.2.2.1.1](#).

**Item 3: Adjointness II**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 4: As a Weighted Limit**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 5: Iterated Cotensors**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 6: Commutativity With Hom**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 7: The Cotensor Evaluation Map**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ .

**Item 8: The Cotensor Coevaluation Map**

This is the special case of ?? of ?? for  $C = \text{Sets}_*$ . 

**00DA 7.3 The Left Tensor Product of Pointed Sets****00DB 7.3.1 Foundations**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00DC DEFINITION 7.3.1.1.1 ► THE LEFT TENSOR PRODUCT OF POINTED SETS**

The **left tensor product of pointed sets** is the functor<sup>1</sup>

$$\triangleleft : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{id} \times \tilde{\otimes}} \text{Sets}_* \times \text{Sets} \xrightarrow{\beta_{\text{Sets}_*, \text{Sets}}^{\text{Cats}_2}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*,$$

where:

- $\tilde{\otimes} : \text{Sets}_* \times \text{Sets} \rightarrow \text{Sets}$  is the forgetful functor from pointed sets to sets.
- $\beta_{\text{Sets}_*, \text{Sets}}^{\text{Cats}_2} : \text{Sets}_* \times \text{Sets} \xrightarrow{\sim} \text{Sets} \times \text{Sets}_*$  is the braiding of  $\text{Cats}_2$ , i.e. the functor witnessing the isomorphism

$$\text{Sets}_* \times \text{Sets} \cong \text{Sets} \times \text{Sets}_*.$$

- $\odot: \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*$  is the tensor functor of Item 1 of Proposition 7.2.1.1.9.

<sup>1</sup>Further Notation: Also written  $\triangleleft_{\text{Sets}_*}$ .

00DD

#### REMARK 7.3.1.1.2 ► UNWINDING DEFINITION 7.3.1.1.1: UNIVERSAL PROPERTY I

The left tensor product of pointed sets satisfies the following natural bijection:

$$\text{Sets}_*(X \triangleleft Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z).$$

That is to say, the following data are in natural bijection:

026B

1. Pointed maps  $f: X \triangleleft Y \rightarrow Z$ .

026C

2. Maps of sets  $f: X \times Y \rightarrow Z$  satisfying  $f(x_0, y) = z_0$  for each  $y \in Y$ .

00DE

#### REMARK 7.3.1.1.3 ► UNWINDING DEFINITION 7.3.1.1.1: UNIVERSAL PROPERTY II

The left tensor product of pointed sets may be described as follows:

- The left tensor product of  $(X, x_0)$  and  $(Y, y_0)$  is the pair  $((X \triangleleft Y, x_0 \triangleleft y_0), \iota)$  consisting of
  - A pointed set  $(X \triangleleft Y, x_0 \triangleleft y_0)$ ;
  - A left bilinear morphism of pointed sets  $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \triangleleft Y$ ;

satisfying the following universal property:

(★) Given another such pair  $((Z, z_0), f)$  consisting of

- \* A pointed set  $(Z, z_0)$ ;
- \* A left bilinear morphism of pointed sets  $f: (X \times Y, (x_0, y_0)) \rightarrow Z$ ;

there exists a unique morphism of pointed sets  $X \triangleleft Y \xrightarrow{\exists!} Z$  making the diagram

$$\begin{array}{ccc} & X \triangleleft Y & \\ \iota \nearrow & \downarrow & \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

00DF

## CONSTRUCTION 7.3.1.1.4 ► THE LEFT TENSOR PRODUCT OF POINTED SETS

In detail, the **left tensor product of**  $(X, x_0)$  **and**  $(Y, y_0)$  is the pointed set  $(X \triangleleft Y, [x_0])$  consisting of:

- *The Underlying Set.* The set  $X \triangleleft Y$  defined by

$$\begin{aligned} X \triangleleft Y &\stackrel{\text{def}}{=} |Y| \odot X \\ &\cong \bigvee_{y \in Y} (X, x_0), \end{aligned}$$

where  $|Y|$  denotes the underlying set of  $(Y, y_0)$ .

- *The Underlying Basepoint.* The point  $[(y_0, x_0)]$  of  $\bigvee_{y \in Y} (X, x_0)$ , which is equal to  $[(y, x_0)]$  for any  $y \in Y$ .

## PROOF 7.3.1.1.5 ► PROOF OF CONSTRUCTION 7.3.1.1.4

Since  $\bigvee_{y \in Y} (X, x_0)$  is defined as the quotient of  $\coprod_{y \in Y} X$  by the equivalence relation  $R$  generated by declaring  $(y, x) \sim (y', x')$  if  $x = x' = x_0$ , we have, by ??, a natural bijection

$$\text{Sets}_*(X \triangleleft Y, Z) \cong \text{Hom}_{\text{Sets}}^R(\coprod_{y \in Y} X, Z),$$

where  $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$  is the set

$$\text{Hom}_{\text{Sets}}^R(\coprod_{y \in Y} X, Z) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(\coprod_{y \in Y} X, Z) \mid \begin{array}{l} \text{for each } x, y \in X, \text{ if} \\ (y, x) \sim_R (y', x'), \text{ then} \\ f(y, x) = f(y', x') \end{array} \right\}.$$

However, the condition  $(y, x) \sim_R (y', x')$  only holds when:

026D

1. We have  $x = x'$  and  $y = y'$ .

026E

2. We have  $x = x' = x_0$ .

So, given  $f \in \text{Hom}_{\text{Sets}}(\coprod_{y \in Y} X, Z)$  with a corresponding  $\bar{f}: X \triangleleft Y \rightarrow Z$ , the latter case above implies

$$\begin{aligned} f([(y, x_0)]) &= f([(y', x_0)]) \\ &= f([(y_0, x_0)]), \end{aligned}$$

and since  $\bar{f}: X \triangleleft Y \rightarrow Z$  is a pointed map, we have

$$f([(y_0, x_0)]) = \bar{f}([(y_0, x_0)])$$

$$= z_0.$$

Thus the elements  $f$  in  $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$  are precisely those functions  $f: X \times Y \rightarrow Z$  satisfying the equality

$$f(x_0, y) = z_0$$

for each  $y \in Y$ , giving an equality

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) = \text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z)$$

of sets, which when composed with our earlier isomorphism

$$\text{Sets}_*(X \triangleleft Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z),$$

gives our desired natural bijection, finishing the proof. 

00DG

#### NOTATION 7.3.1.1.6 ► ELEMENTS OF LEFT TENSOR PRODUCTS OF POINTED SETS

We write<sup>1</sup>  $x \triangleleft y$  for the element  $[(y, x)]$  of

$$X \triangleleft Y \cong |Y| \odot X.$$

<sup>1</sup>Further Notation: Also written  $x \triangleleft_{\text{Sets}_*} y$ .

00DH

#### REMARK 7.3.1.1.7 ► BASEPOINTS OF LEFT TENSOR PRODUCTS OF POINTED SETS

Employing the notation introduced in [Notation 7.3.1.1.6](#), we have

$$x_0 \triangleleft y_0 = x_0 \triangleleft y$$

for each  $y \in Y$ , and

$$x_0 \triangleleft y = x_0 \triangleleft y'$$

for each  $y, y' \in Y$ .

00DJ

#### PROPOSITION 7.3.1.1.8 ► PROPERTIES OF LEFT TENSOR PRODUCTS OF POINTED SETS

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00DK

1. *Functionality.* The assignments  $X, Y, (X, Y) \mapsto X \triangleleft Y$  define functors

$$\begin{aligned} X \triangleleft -: \quad \text{Sets}_* &\longrightarrow \text{Sets}_*, \\ - \triangleleft Y: \quad \text{Sets}_* &\longrightarrow \text{Sets}_*, \\ -_1 \triangleleft -_2: \text{Sets}_* \times \text{Sets}_* &\longrightarrow \text{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f: (X, x_0) &\rightarrow (A, a_0), \\ g: (Y, y_0) &\rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \triangleleft g: X \triangleleft Y \rightarrow A \triangleleft B$$

is given by

$$[f \triangleleft g](x \triangleleft y) \stackrel{\text{def}}{=} f(x) \triangleleft g(y)$$

for each  $x \triangleleft y \in X \triangleleft Y$ .

00DL

2. *Adjointness I.* We have an adjunction

$$\left( - \triangleleft Y \dashv [Y, -]_{\mathbf{Sets}_*}^\triangleleft \right): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \triangleleft Y} \\ \perp \\ \xleftarrow{[Y, -]_{\mathbf{Sets}_*}^\triangleleft} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, [Y, Z]_{\mathbf{Sets}_*}^\triangleleft)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ , where  $[X, Y]_{\mathbf{Sets}_*}^\triangleleft$  is the pointed set of [Definition 7.3.2.1.1](#).

00DM

3. *Adjointness II.* The functor

$$X \triangleleft -: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

00DN

4. *Adjointness III.* We have a  $\overline{\mathbb{E}}$ -relative adjunction

$$(X \triangleleft - \dashv \mathbf{Sets}_*(X, -)): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \triangleleft -} \\ \perp_{\overline{\mathbb{E}}} \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}}(|Y|, \mathbf{Sets}_*(X, Z))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ .

**PROOF 7.3.1.1.9 ► PROOF OF PROPOSITION 7.3.1.1.8****Item 1: Functoriality**

This follows from the definition of  $\triangleleft$  as a composition of functors ([Definition 7.3.1.1.1](#)).

**Item 2: Adjointness I**

This follows from [Item 3 of Proposition 7.2.1.1.9](#).

**Item 3: Adjointness II**

For  $X \triangleleft -$  to admit a right adjoint would require it to preserve colimits by ?? of ?. However, we have

$$\begin{aligned} X \triangleleft \text{pt} &\stackrel{\text{def}}{=} |\text{pt}| \odot X \\ &\cong X \\ &\not\cong \text{pt}, \end{aligned}$$

and thus we see that  $X \triangleleft -$  does not have a right adjoint.

**Item 4: Adjointness III**

This follows from [Item 2 of Proposition 7.2.1.1.9](#). 

00DP

**REMARK 7.3.1.1.10 ► ON THE FAILURE OF  $X \triangleleft -$  TO BE A LEFT ADJOINT**

Here is some intuition on why  $X \triangleleft -$  fails to be a left adjoint. [Item 4 of Proposition 7.3.1.1.8](#) states that we have a natural bijection

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}}(|Y|, \mathbf{Sets}_*(X, Z)),$$

so it would be reasonable to wonder whether a natural bijection of the form

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(Y, \mathbf{Sets}_*(X, Z)),$$

also holds, which would give  $X \triangleleft - \dashv \mathbf{Sets}_*(X, -)$ . However, such a bijection would require every map

$$f: X \triangleleft Y \rightarrow Z$$

to satisfy

$$f(x \triangleleft y_0) = z_0$$

for each  $x \in X$ , whereas we are imposing such a basepoint preservation condition only for elements of the form  $x_0 \triangleleft y$ . Thus  $\mathbf{Sets}_*(X, -)$  can't be a right adjoint for  $X \triangleleft -$ , and as shown by [Item 3 of Proposition 7.3.1.1.8](#), no functor can.<sup>1</sup>

<sup>1</sup>The functor  $\mathbf{Sets}_*(X, -)$  is instead right adjoint to  $X \wedge -$ , the smash product of pointed sets of [Definition 7.5.1.1.1](#). See [Item 2 of Proposition 7.5.1.1.12](#).

**00DQ 7.3.2 The Left Internal Hom of Pointed Sets**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00DR**

**DEFINITION 7.3.2.1.1 ► THE LEFT INTERNAL HOM OF POINTED SETS**

The **left internal Hom<sup>1</sup>** of pointed sets is the functor

$$[-, -]_{\text{Sets}_*}^\triangleleft : \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_*^{\text{op}} \times \text{Sets}_* \xrightarrow{\text{Forget} \times \text{id}} \text{Sets}_*^{\text{op}} \times \text{Sets}_* \xrightarrow{\pitchfork} \text{Sets}_*,$$

where:

- $\text{Forget} : \text{Sets}_* \rightarrow \text{Sets}$  is the forgetful functor from pointed sets to sets.
- $\pitchfork : \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$  is the cotensor functor of **Item 1** of **Proposition 7.2.2.1.6**.

<sup>1</sup>For a proof that  $[-, -]_{\text{Sets}_*}^\triangleleft$  is indeed the left internal Hom of  $\text{Sets}_*$  with respect to the left tensor product of pointed sets, see **Item 2** of **Proposition 7.3.1.1.8**.

**00DS**

**REMARK 7.3.2.1.2 ► UNWINDING DEFINITION 7.3.2.1.1, I: UNIVERSAL PROPERTY**

The left internal Hom of pointed sets satisfies the following universal property:

$$\text{Sets}_*(X \triangleleft Y, Z) \cong \text{Sets}_*(X, [Y, Z]_{\text{Sets}_*}^\triangleleft)$$

That is to say, the following data are in bijection:

**026F**

1. Pointed maps  $f : X \triangleleft Y \rightarrow Z$ .

**026G**

2. Pointed maps  $f : X \rightarrow [Y, Z]_{\text{Sets}_*}^\triangleleft$ .

**00DT**

**REMARK 7.3.2.1.3 ► UNWINDING DEFINITION 7.3.2.1.1, II: EXPLICIT DESCRIPTION**

In detail, the **left internal Hom** of  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $([X, Y]_{\text{Sets}_*}^\triangleleft, [(y_0)_{x \in X}])$  consisting of:

- *The Underlying Set.* The set  $[X, Y]_{\text{Sets}_*}^\triangleleft$  defined by

$$\begin{aligned} [X, Y]_{\text{Sets}_*}^\triangleleft &\stackrel{\text{def}}{=} |X| \pitchfork Y \\ &\cong \bigwedge_{x \in X} (Y, y_0), \end{aligned}$$

where  $|X|$  denotes the underlying set of  $(X, x_0)$ .

- *The Underlying Basepoint.* The point  $[(y_0)_{x \in X}]$  of  $\bigwedge_{x \in X} (Y, y_0)$ .

00DU

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

- 00DV** 1. *Functionality.* The assignments  $X, Y, (X, Y) \mapsto [X, Y]_{\text{Sets}_*}^\triangleleft$  define functors

$$\begin{aligned}[X, -]_{\text{Sets}_*}^\triangleleft : \text{Sets}_* &\rightarrow \text{Sets}_*, \\ [-, Y]_{\text{Sets}_*}^\triangleleft : \text{Sets}_*^{\text{op}} &\rightarrow \text{Sets}_*, \\ [-_1, -_2]_{\text{Sets}_*}^\triangleleft : \text{Sets}_*^{\text{op}} \times \text{Sets}_* &\rightarrow \text{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned}f &: (X, x_0) \rightarrow (A, a_0), \\g &: (Y, y_0) \rightarrow (B, b_0),\end{aligned}$$

the induced map

$$[f, g]_{\text{Sets}_*}^\triangleleft : [A, Y]_{\text{Sets}_*}^\triangleleft \rightarrow [X, B]_{\text{Sets}_*}^\triangleleft$$

is given by

$$[f, g]_{\text{Sets}_*}^\triangleleft([(y_a)_{a \in A}]) \stackrel{\text{def}}{=} [(g(y_{f(x)}))_{x \in X}]$$

for each  $[(y_a)_{a \in A}] \in [A, Y]_{\text{Sets}_*}^\triangleleft$ .

- 00DW** 2. *Adjointness I.* We have an adjunction

$$\left( - \triangleleft Y \dashv [Y, -]_{\text{Sets}_*}^\triangleleft \right) : \text{Sets}_* \begin{array}{c} \xrightarrow{- \triangleleft Y} \\ \perp \\ \xleftarrow{[Y, -]_{\text{Sets}_*}^\triangleleft} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\text{Sets}_*}(X, [Y, Z]_{\text{Sets}_*}^\triangleleft)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$

- 00DX** 3. *Adjointness II.* The functor

$$X \triangleleft - : \text{Sets}_* \rightarrow \text{Sets}_*$$

does not admit a right adjoint.

**PROOF 7.3.2.1.5 ► PROOF OF PROPOSITION 7.3.2.1.4****Item 1: Functoriality**

This follows from the definition of  $[-, -]_{\text{Sets}_*}^\triangleleft$  as a composition of functors ([Definition 7.3.2.1.1](#)).

**Item 2: Adjointness I**

This is a repetition of [Item 2](#) of [Proposition 7.3.1.1.8](#), and is proved there.

**Item 3: Adjointness II**

This is a repetition of [Item 3](#) of [Proposition 7.3.1.1.8](#), and is proved there. 

**00DY 7.3.3 The Left Skew Unit****00DZ DEFINITION 7.3.3.1.1 ► THE LEFT SKEW UNIT OF  $\triangleleft$** 

The **left skew unit** of the left tensor product of pointed sets is the functor

$$\mathbb{1}_{\text{Sets}_*}^{\triangleleft} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*}^{\triangleleft} \stackrel{\text{def}}{=} S^0.$$

**00E0 7.3.4 The Left Skew Associator****00E1 DEFINITION 7.3.4.1.1 ► THE LEFT SKEW ASSOCIATOR OF  $\triangleleft$** 

The **skew associator** of the left tensor product of pointed sets is the natural transformation

$$\alpha^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\triangleleft \times \text{id}_{\text{Sets}_*}) \Longrightarrow \triangleleft \circ (\text{id}_{\text{Sets}_*} \times \triangleleft) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}}$$

as in the diagram

$$\begin{array}{ccc}
 & \text{Sets}_* \times (\text{Sets}_* \times \text{Sets}_*) & \\
 & \swarrow \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}} & \searrow \text{id} \times \triangleleft \\
 (\text{Sets}_* \times \text{Sets}_*) \times \text{Sets}_* & \xrightarrow{\quad \quad} & \text{Sets}_* \times \text{Sets}_* \\
 \downarrow \triangleleft \times \text{id} & \xrightarrow{\quad \quad} & \downarrow \triangleleft \\
 \text{Sets}_* \times \text{Sets}_* & \xrightarrow{\alpha^{\text{Sets}_*, \triangleleft}} & \text{Sets}_*,
 \end{array}$$

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} : (X \triangleleft Y) \triangleleft Z \rightarrow X \triangleleft (Y \triangleleft Z)$$

at  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\begin{aligned} (X \triangleleft Y) \triangleleft Z &\stackrel{\text{def}}{=} |Z| \odot (X \triangleleft Y) \\ &\stackrel{\text{def}}{=} |Z| \odot (|Y| \odot X) \\ &\cong \bigvee_{z \in Z} |Y| \odot X \\ &\cong \bigvee_{z \in Z} (\bigvee_{y \in Y} X) \\ &\rightarrow \bigvee_{[(z,y)] \in \bigvee_{z \in Z} Y} X \\ &\cong \bigvee_{[(z,y)] \in |Z| \odot Y} X \\ &\cong ||Z| \odot Y| \odot X \\ &\stackrel{\text{def}}{=} |Y \triangleleft Z| \odot X \\ &\stackrel{\text{def}}{=} X \triangleleft (Y \triangleleft Z), \end{aligned}$$

where the map

$$\bigvee_{z \in Z} (\bigvee_{y \in Y} X) \rightarrow \bigvee_{(z,y) \in \bigvee_{z \in Z} Y} X$$

is given by  $[(z, [(y, x)])] \mapsto [([(z, y)], x)]$ .

#### PROOF 7.3.4.1.2 ► PROOF OF DEFINITION 7.3.4.1.1

(Proven below in a bit.)



00E2

#### REMARK 7.3.4.1.3 ► UNWINDING DEFINITION 7.3.4.1.1

Unwinding the notation for elements, we have

$$\begin{aligned} [(z, [(y, x)])] &\stackrel{\text{def}}{=} [(z, x \triangleleft y)] \\ &\stackrel{\text{def}}{=} (x \triangleleft y) \triangleleft z \end{aligned}$$

and

$$\begin{aligned} [([(z, y)], x)] &\stackrel{\text{def}}{=} [(y \triangleleft z, x)] \\ &\stackrel{\text{def}}{=} x \triangleleft (y \triangleleft z). \end{aligned}$$

So, in other words,  $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$  acts on elements via

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} ((x \triangleleft y) \triangleleft z) \stackrel{\text{def}}{=} x \triangleleft (y \triangleleft z)$$

for each  $(x \triangleleft y) \triangleleft z \in (X \triangleleft Y) \triangleleft Z$ .

00E3

**REMARK 7.3.4.1.4 ► NON-INVERTIBILITY OF THE SKEW ASSOCIATOR OF  $\triangleleft$** 

Taking  $y = y_0$ , we see that the morphism  $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$  acts on elements as

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}((x \triangleleft y_0) \triangleleft z) \stackrel{\text{def}}{=} x \triangleleft (y_0 \triangleleft z).$$

However, by the definition of  $\triangleleft$ , we have  $y_0 \triangleleft z = y_0 \triangleleft z'$  for all  $z, z' \in Z$ , preventing  $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$  from being non-invertible.

**PROOF 7.3.4.1.5 ► PROOF OF DEFINITION 7.3.4.1.1**

Firstly, note that, given  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}: (X \triangleleft Y) \triangleleft Z \rightarrow X \triangleleft (Y \triangleleft Z)$$

is indeed a morphism of pointed sets, as we have

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}((x_0 \triangleleft y_0) \triangleleft z_0) = x_0 \triangleleft (y_0 \triangleleft z_0).$$

Next, we claim that  $\alpha^{\text{Sets}_*, \triangleleft}$  is a natural transformation. We need to show that, given morphisms of pointed sets

$$\begin{aligned} f: (X, x_0) &\rightarrow (X', x'_0), \\ g: (Y, y_0) &\rightarrow (Y', y'_0), \\ h: (Z, z_0) &\rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \triangleleft Y) \triangleleft Z & \xrightarrow{(f \triangleleft g) \triangleleft h} & (X' \triangleleft Y') \triangleleft Z' \\ \downarrow \alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*, \triangleleft} \\ X \triangleleft (Y \triangleleft Z) & \xrightarrow{f \triangleleft (g \triangleleft h)} & X' \triangleleft (Y' \triangleleft Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \triangleleft y) \triangleleft z & \longmapsto & (f(x) \triangleleft g(y)) \triangleleft h(z) \\ \downarrow & & \downarrow \\ x \triangleleft (y \triangleleft z) & \longmapsto & f(x) \triangleleft (g(y) \triangleleft h(z)) \end{array}$$

and hence indeed commutes, showing  $\alpha^{\text{Sets}_*, \triangleleft}$  to be a natural transformation. This finishes the proof. ■

**00E4 7.3.5 The Left Skew Left Unit**

00E5

**DEFINITION 7.3.5.1.1 ► THE LEFT SKEW LEFT UNIT OF  $\triangleleft$**

The **skew left unit** of the left tensor product of pointed sets is the natural transformation

$$\begin{array}{ccc} \text{pt} \times \text{Sets}_* & \xrightarrow{\mathbb{1}_{\text{Sets}_*} \times \text{id}} & \text{Sets}_* \times \text{Sets}_* \\ \downarrow & \nearrow \lambda_{\text{Sets}_*, \triangleleft} & \downarrow \triangleleft \\ \lambda^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\mathbb{1}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) & \xrightarrow{\sim} & \lambda^{\text{Cats}_2}_{\text{Sets}_*} \\ \downarrow & \nearrow \lambda^{\text{Cats}_2}_{\text{Sets}_*} & \downarrow \\ & & \text{Sets}_*, \end{array}$$

whose component

$$\lambda_X^{\text{Sets}_*, \triangleleft} : S^0 \triangleleft X \rightarrow X$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by the composition

$$\begin{aligned} S^0 \triangleleft X &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X, \end{aligned}$$

where  $\bigvee_{x \in X} S^0 \rightarrow X$  is the map given by

$$\begin{aligned} [(x, 0)] &\mapsto x_0, \\ [(x, 1)] &\mapsto x \end{aligned}$$

for each  $x \in X$ .

**PROOF 7.3.5.1.2 ► PROOF OF DEFINITION 7.3.5.1.1**

(Proven below in a bit.)



00E6

**REMARK 7.3.5.1.3 ► UNWINDING DEFINITION 7.3.5.1.1**

In other words,  $\lambda_X^{\text{Sets}_*, \triangleleft}$  acts on elements as

$$\begin{aligned} \lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x) &\stackrel{\text{def}}{=} x_0, \\ \lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x) &\stackrel{\text{def}}{=} x \end{aligned}$$

for each  $1 \triangleleft x \in S^0 \triangleleft X$ .

00E7

The morphism  $\lambda_X^{\text{Sets}_*, \triangleleft}$  is almost invertible, with its would-be-inverse

$$\phi_X: X \rightarrow S^0 \triangleleft X$$

given by

$$\phi_X(x) \stackrel{\text{def}}{=} 1 \triangleleft x$$

for each  $x \in X$ . Indeed, we have

$$\begin{aligned} [\lambda_X^{\text{Sets}_*, \triangleleft} \circ \phi](x) &= \lambda_X^{\text{Sets}_*, \triangleleft}(\phi(x)) \\ &= \lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

so that

$$\lambda_X^{\text{Sets}_*, \triangleleft} \circ \phi = \text{id}_X$$

and

$$\begin{aligned} [\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft}](1 \triangleleft x) &= \phi(\lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x)) \\ &= \phi(x) \\ &= 1 \triangleleft x \\ &= [\text{id}_{S^0 \triangleleft X}](1 \triangleleft x), \end{aligned}$$

but

$$\begin{aligned} [\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft}](0 \triangleleft x) &= \phi(\lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x)) \\ &= \phi(x_0) \\ &= 1 \triangleleft x_0, \end{aligned}$$

where  $0 \triangleleft x \neq 1 \triangleleft x_0$ . Thus

$$\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft} \stackrel{?}{=} \text{id}_{S^0 \triangleleft X}$$

holds for all elements in  $S^0 \triangleleft X$  except one.

#### PROOF 7.3.5.1.5 ▶ PROOF OF DEFINITION 7.3.5.1.1

Firstly, note that, given  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\lambda_X^{\text{Sets}_*, \triangleleft}: S^0 \triangleleft X \rightarrow X$$

is indeed a morphism of pointed sets, as we have

$$\lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x_0) = x_0.$$

Next, we claim that  $\lambda^{\text{Sets}_*, \triangleleft}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} S^0 \triangleleft X & \xrightarrow{\text{id}_{S^0} \triangleleft f} & S^0 \triangleleft Y \\ \lambda_X^{\text{Sets}_*, \triangleleft} \downarrow & & \downarrow \lambda_Y^{\text{Sets}_*, \triangleleft} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 \triangleleft x & & 0 \triangleleft x \longmapsto 0 \triangleleft f(x) \\ \downarrow & & \downarrow \\ x_0 \longmapsto f(x_0) & & y_0 \end{array}$$

and

$$\begin{array}{ccc} 1 \triangleleft x \longmapsto 1 \triangleleft f(x) & & \\ \downarrow & & \downarrow \\ x \longmapsto f(x) & & \end{array}$$

and hence indeed commutes, showing  $\lambda^{\text{Sets}_*, \triangleleft}$  to be a natural transformation. This finishes the proof. ■

### 00E8 7.3.6 The Left Skew Right Unit

00E9

DEFINITION 7.3.6.1.1 ► THE LEFT SKEW RIGHT UNIT OF  $\triangleleft$

The **skew right unit of the left tensor product of pointed sets** is the natural transformation

$$\begin{array}{ccc}
 \text{Sets}_* \times \text{pt} & \xrightarrow{\text{id} \times \mathbb{1}^{\text{Sets}_*}} & \text{Sets}_* \times \text{Sets}_* \\
 \rho^{\text{Sets}_*, \triangleleft} : \rho_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleleft \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}), & \swarrow \quad \searrow & \downarrow \triangleleft \\
 & \rho^{\text{Sets}_*, \triangleleft} & \\
 & \rho_{\text{Sets}_*}^{\text{Cats}_2} & \\
 & \searrow & \downarrow \triangleleft \\
 & & \text{Sets}_*,
 \end{array}$$

whose component

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft S^0$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by the composition

$$\begin{aligned}
 X &\rightarrow X \vee X \\
 &\cong |S^0| \odot X \\
 &\cong X \triangleleft S^0,
 \end{aligned}$$

where  $X \rightarrow X \vee X$  is the map sending  $X$  to the second factor of  $X$  in  $X \vee X$ .

#### PROOF 7.3.6.1.2 ▶ PROOF OF DEFINITION 7.3.6.1.1

(Proven below in a bit.)



00EA

#### REMARK 7.3.6.1.3 ▶ UNWINDING DEFINITION 7.3.6.1.1

In other words,  $\rho_X^{\text{Sets}_*, \triangleleft}$  acts on elements as

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} [(1, x)]$$

i.e. by

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} x \triangleleft 1$$

for each  $x \in X$ .

00EB

#### REMARK 7.3.6.1.4 ▶ NON-INVERTIBILITY OF THE SKEW RIGHT UNIT OF $\triangleleft$

The morphism  $\rho_X^{\text{Sets}_*, \triangleleft}$  is non-invertible, as it is non-surjective when viewed as a map of sets, since the elements  $x \triangleleft 0$  of  $X \triangleleft S^0$  with  $x \neq x_0$  are outside the image of  $\rho_X^{\text{Sets}_*, \triangleleft}$ , which sends  $x$  to  $x \triangleleft 1$ .

**PROOF 7.3.6.1.5 ▶ PROOF OF DEFINITION 7.3.6.1.1**

Firstly, note that, given  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft S^0$$

is indeed a morphism of pointed sets as we have

$$\begin{aligned}\rho_X^{\text{Sets}_*, \triangleleft}(x_0) &= x_0 \triangleleft 1 \\ &= x_0 \triangleleft 0.\end{aligned}$$

Next, we claim that  $\rho^{\text{Sets}_*, \triangleleft}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \rho_X^{\text{Sets}_*, \triangleleft} \downarrow & & \downarrow \rho_Y^{\text{Sets}_*, \triangleleft} \\ X \triangleleft S^0 & \xrightarrow[f \triangleleft \text{id}_{S^0}]{} & Y \triangleleft S^0 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \mapsto & f(x) \\ \downarrow & & \downarrow \\ x \triangleleft 0 & \mapsto & f(x) \triangleleft 0 \end{array}$$

and hence indeed commutes, showing  $\rho^{\text{Sets}_*, \triangleleft}$  to be a natural transformation. This finishes the proof. ■

**00EC 7.3.7 The Diagonal**

00ED

**DEFINITION 7.3.7.1.1 ▶ THE DIAGONAL OF  $\triangleleft$**

The **diagonal of the left tensor product of pointed sets** is the natural transformation

$$\Delta^\triangleleft : \text{id}_{\text{Sets}_*} \Rightarrow \triangleleft \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

whose component

$$\Delta_X^\triangleleft : (X, x_0) \rightarrow (X \triangleleft X, x_0 \triangleleft x_0)$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\Delta_X^\triangleleft(x) \stackrel{\text{def}}{=} x \triangleleft x$$

for each  $x \in X$ .

#### PROOF 7.3.7.1.2 ► PROOF OF DEFINITION 7.3.7.1.1

Being a Morphism of Pointed Sets

We have

$$\Delta_X^\triangleleft(x_0) \stackrel{\text{def}}{=} x_0 \triangleleft x_0,$$

and thus  $\Delta_X^\triangleleft$  is a morphism of pointed sets.

#### Naturality

We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\triangleleft \downarrow & & \downarrow \Delta_Y^\triangleleft \\ X \triangleleft X & \xrightarrow{f \triangleleft f} & Y \triangleleft Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \xrightarrow{\quad} & f(x) \\ \downarrow & & \downarrow \\ x \triangleleft x & \xrightarrow{\quad} & f(x) \triangleleft f(x) \end{array}$$

and hence indeed commutes, showing  $\Delta^\triangleleft$  to be natural. ■

### 00EE 7.3.8 The Left Skew Monoidal Structure on Pointed Sets Associated to $\triangleleft$

00EF PROPOSITION 7.3.8.1.1 ► THE LEFT SKEW MONOIDAL STRUCTURE ON POINTED SETS ASSOCIATED TO  $\triangleleft$

The category  $\text{Sets}_*$  admits a left-closed left skew monoidal category structure consisting of:

- *The Underlying Category.* The category  $\text{Sets}_*$  of pointed sets.
- *The Left Skew Monoidal Product.* The left tensor product functor

$$\triangleleft : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Definition 7.3.1.1.1.

- *The Left Internal Skew Hom.* The left internal Hom functor

$$[-, -]_{\text{Sets}_*}^\triangleleft : \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Definition 7.3.2.1.1.

- *The Left Skew Monoidal Unit.* The functor

$$\mathbb{1}_{\text{Sets}_*, \triangleleft} : \text{pt} \rightarrow \text{Sets}_*$$

of Definition 7.3.3.1.1.

- *The Left Skew Associators.* The natural transformation

$$\alpha^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\triangleleft \times \text{id}_{\text{Sets}_*}) \Longrightarrow \triangleleft \circ (\text{id}_{\text{Sets}_*} \times \triangleleft) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}}$$

of Definition 7.3.4.1.1.

- *The Left Skew Left Unitors.* The natural transformation

$$\lambda^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\mathbb{1}^{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \lambda_{\text{Sets}_*}^{\text{Cats}_2}$$

of Definition 7.3.5.1.1.

- *The Left Skew Right Unitors.* The natural transformation

$$\rho^{\text{Sets}_*, \triangleleft} : \rho_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleleft \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*})$$

of Definition 7.3.6.1.1.

### PROOF 7.3.8.1.2 ► PROOF OF PROPOSITION 7.3.8.1.1

#### The Pentagon Identity

Let  $(W, w_0)$ ,  $(X, x_0)$ ,  $(Y, y_0)$  and  $(Z, z_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccccc}
 & & (W \triangleleft (X \triangleleft Y)) \triangleleft Z & & \\
 & \swarrow \alpha_{W,X,Y}^{\text{Sets}_*, \triangleleft} \triangleleft \text{id}_Z & & \searrow \alpha_{W,X \triangleleft Y,Z}^{\text{Sets}_*, \triangleleft} & \\
 ((W \triangleleft X) \triangleleft Y) \triangleleft Z & & & & W \triangleleft ((X \triangleleft Y) \triangleleft Z) \\
 & \downarrow \alpha_{W \triangleleft X, Y, Z}^{\text{Sets}_*, \triangleleft} & & \downarrow \text{id}_W \triangleleft \alpha_{X, Y, Z}^{\text{Sets}_*, \triangleleft} & \\
 & (W \triangleleft X) \triangleleft (Y \triangleleft Z) & \xrightarrow{\alpha_{W, X, Y \triangleleft Z}^{\text{Sets}_*, \triangleleft}} & W \triangleleft (X \triangleleft (Y \triangleleft Z)) &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccccc}
 & & (w \triangleleft (x \triangleleft y)) \triangleleft z & & \\
 & \swarrow & & \searrow & \\
 ((w \triangleleft x) \triangleleft y) \triangleleft z & & & & w \triangleleft ((x \triangleleft y) \triangleleft z) \\
 \downarrow & & & & \downarrow \\
 (w \triangleleft x) \triangleleft (y \triangleleft z) & \longmapsto & w \triangleleft (x \triangleleft (y \triangleleft z))
 \end{array}$$

and thus we see that the pentagon identity is satisfied.

### The Left Skew Left Triangle Identity

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 (S^0 \triangleleft X) \triangleleft Y & \xrightarrow{\alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleleft}} & S^0 \triangleleft (X \triangleleft Y) \\
 \lambda_X^{\text{Sets}_*, \triangleleft} \text{id}_Y \searrow & & \downarrow \lambda_{X \triangleleft Y}^{\text{Sets}_*, \triangleleft} \\
 & & X \triangleleft Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (0 \triangleleft x) \triangleleft y & \longmapsto & 0 \triangleleft (x \triangleleft y) \\
 \downarrow & & \downarrow \\
 x_0 \triangleleft y = x_0 \triangleleft y_0
 \end{array}$$

and

$$\begin{array}{ccc}
 (1 \triangleleft x) \triangleleft y & \longmapsto & 1 \triangleleft (x \triangleleft y) \\
 \downarrow & & \downarrow \\
 x \triangleleft y
 \end{array}$$

and hence indeed commutes. Thus the left skew triangle identity is satisfied.

### The Left Skew Right Triangle Identity

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc} X \triangleleft Y & & \\ \downarrow \rho_{X \triangleleft Y}^{\text{Sets}_*, \triangleleft} & \searrow \text{id}_X \triangleleft \rho_Y^{\text{Sets}_*, \triangleleft} & \\ (X \triangleleft Y) \triangleleft S^0 & \xrightarrow{\alpha_{X,Y,S^0}^{\text{Sets}_*, \triangleleft}} & X \triangleleft (Y \triangleleft S^0) \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleleft y & & \\ \downarrow & \nearrow & \\ (x \triangleleft y) \triangleleft 1 & \mapsto & x \triangleleft (y \triangleleft 1) \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

### The Left Skew Middle Triangle Identity

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc} X \triangleleft Y & \xlongequal{\quad} & X \triangleleft Y \\ \downarrow \rho_X^{\text{Sets}_*, \triangleleft} \triangleleft \text{id}_Y & & \uparrow \text{id}_A \triangleleft \lambda_Y^{\text{Sets}_*, \triangleleft} \\ (X \triangleleft S^0) \triangleleft Y & \xrightarrow{\alpha_{X,S^0,Y}^{\text{Sets}_*, \triangleleft}} & X \triangleleft (S^0 \triangleleft Y) \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleleft y & \xlongequal{\quad} & x \triangleleft y \\ \downarrow & & \uparrow \\ (x \triangleleft 1) \triangleleft y & \mapsto & x \triangleleft (1 \triangleleft y) \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

### The Zig-Zag Identity

We have to show that the diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{\rho_{S^0}^{\text{Sets}_*, \triangleleft}} & S^0 \triangleleft S^0 \\ & \searrow & \downarrow \lambda_{S^0}^{\text{Sets}_*, \triangleleft} \\ & & S^0 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 & \xrightarrow{\quad} & 0 \triangleleft 1 \\ & \swarrow & \downarrow \\ & & 0 \end{array}$$

and

$$\begin{array}{ccc} 1 & \xrightarrow{\quad} & 1 \triangleleft 1 \\ & \swarrow & \downarrow \\ & & 1 \end{array}$$

and hence indeed commutes. Thus the zig-zag identity is satisfied.

#### Left Skew Monoidal Left-Closedness

This follows from [Item 2 of Proposition 7.3.1.1.8](#). 

### 00EG 7.3.9 Monoids With Respect to the Left Tensor Product of Pointed Sets

00EH

#### PROPOSITION 7.3.9.1.1 ► MONOIDS WITH RESPECT TO $\triangleleft$

The category of monoids on  $(\text{Sets}_*, \triangleleft, S^0)$  is isomorphic to the category of “monoids with left zero”<sup>1</sup> and morphisms between them.

<sup>1</sup>A monoid with left zero is defined similarly as the monoids with zero of [??](#). Succinctly, they are monoids  $(A, \mu_A, \eta_A)$  with a special element  $0_A$  satisfying

$$0_A a = 0_A$$

for each  $a \in A$ .

## PROOF 7.3.9.1.2 ► PROOF OF PROPOSITION 7.3.9.1.1

Monoids on  $(\text{Sets}_*, \triangleleft, S^0)$

A monoid on  $(\text{Sets}_*, \triangleleft, S^0)$  consists of:

- *The Underlying Object.* A pointed set  $(A, 0_A)$ .
- *The Multiplication Morphism.* A morphism of pointed sets

$$\mu_A: A \triangleleft A \rightarrow A,$$

determining a left bilinear morphism of pointed sets

$$\begin{aligned} A \times A &\longrightarrow A \\ (a, b) &\longmapsto ab. \end{aligned}$$

- *The Unit Morphism.* A morphism of pointed sets

$$\eta_A: S^0 \rightarrow A$$

picking an element  $1_A$  of  $A$ .

satisfying the following conditions:

026H

1. *Associativity.* The diagram

$$\begin{array}{ccc} & A \triangleleft (A \triangleleft A) & \\ \alpha_{A,A,A}^{\text{Sets}_*, \triangleleft} \nearrow & & \searrow \text{id}_A \triangleleft \mu_A \\ (A \triangleleft A) \triangleleft A & & A \triangleleft A \\ \downarrow \mu_A \triangleleft \text{id}_A & & \downarrow \mu_A \\ A \triangleleft A & \xrightarrow{\mu_A} & A \end{array}$$

026J

2. *Left Unitality.* The diagram

$$\begin{array}{ccc} S^0 \triangleleft A & \xrightarrow{\eta_A \times \text{id}_A} & A \triangleleft A \\ & \searrow \lambda_A^{\text{Sets}_*, \triangleleft} & \downarrow \mu_A \\ & & A \end{array}$$

commutes.

026K

## 3. Right Unitality. The diagram

$$\begin{array}{ccc}
 A & \xrightarrow{\rho_A^{\text{Sets}_{*}, \triangleleft}} & A \triangleleft S^0 \\
 \parallel & & \downarrow \text{id}_A \times \eta_A \\
 A & \xleftarrow{\mu_A} & A \triangleleft A
 \end{array}$$

commutes.

Being a left-bilinear morphism of pointed sets, the multiplication map satisfies

$$0_A a = 0_A$$

for each  $a \in A$ . Now, the associativity, left unitality, and right unitality conditions act on elements as follows:

026L

## 1. Associativity. The associativity condition acts as

$$\begin{array}{ccc}
 & a \triangleleft (b \triangleleft c) & \\
 & \swarrow \quad \searrow & \\
 (a \triangleleft b) \triangleleft c & & a \triangleleft bc \\
 \swarrow & & \downarrow \\
 ab \triangleleft c & \longmapsto & a(bc)
 \end{array}$$

This gives

$$(ab)c = a(bc)$$

for each  $a, b, c \in A$ .

026M

## 2. Left Unitality. The left unitality condition acts:

026N

(a) On  $0 \triangleleft a$  as

$$\begin{array}{ccc}
 0 \triangleleft a & & 0 \triangleleft a \xrightarrow{\quad} 0_A \triangleleft a \\
 \swarrow & & \downarrow \\
 0_A & & 0_A a.
 \end{array}$$

026P

(b) On  $1 \triangleleft a$  as

$$\begin{array}{ccc} 1 \triangleleft a & \xrightarrow{\quad} & 1 \triangleleft a \xrightarrow{\quad} 1_A \triangleleft a \\ \downarrow & \nearrow a & \downarrow \\ a & & 1_A a. \end{array}$$

This gives

$$\begin{aligned} 1_A a &= a, \\ 0_A a &= 0_A \end{aligned}$$

for each  $a \in A$ .

026Q

3. Right Unitality. The right unitality condition acts as

$$\begin{array}{ccc} a & \xrightarrow{\quad} & a \triangleright 1 \\ \downarrow & & \downarrow \\ a & & a 1_A \xleftarrow{\quad} a \triangleleft 1_A \end{array}$$

This gives

$$a 1_A = a$$

for each  $a \in A$ .Thus we see that monoids with respect to  $\triangleleft$  are exactly monoids with left zero.**Morphisms of Monoids on  $(\text{Sets}_*, \triangleleft, S^0)$** A morphism of monoids on  $(\text{Sets}_*, \triangleleft, S^0)$  from  $(A, \mu_A, \eta_A, 0_A)$  to  $(B, \mu_B, \eta_B, 0_B)$  is a morphism of pointed sets

$$f: (A, 0_A) \rightarrow (B, 0_B)$$

satisfying the following conditions:

026R

1. *Compatibility With the Multiplication Morphisms.* The diagram

$$\begin{array}{ccc} A \triangleleft A & \xrightarrow{f \triangleleft f} & B \triangleleft B \\ \mu_A \downarrow & & \downarrow \mu_B \\ A & \xrightarrow{f} & B \end{array}$$

commutes.

026S

2. *Compatibility With the Unit Morphisms.* The diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{\eta_A} & A \\ & \searrow \eta_B & \downarrow f \\ & & B \end{array}$$

commutes.

These act on elements as

$$\begin{array}{ccc} a \triangleleft b & & a \triangleleft b \mapsto f(a) \triangleleft f(b) \\ \downarrow & & \downarrow \\ ab & \longmapsto & f(ab) \\ & & f(a)f(b) \end{array}$$

and

$$\begin{array}{ccc} 0 & & 0 \mapsto 0_A \\ & \searrow & \downarrow \\ & 0_B & f(0_A) \end{array}$$

and

$$\begin{array}{ccc} 1 & & 1 \mapsto 1_A \\ \downarrow & & \downarrow \\ 1_B & & f(1_A) \end{array}$$

giving

$$f(ab) = f(a)f(b),$$

$$f(0_A) = 0_B,$$

$$f(1_A) = 1_B,$$

for each  $a, b \in A$ , which is exactly a morphism of monoids with left zero.

### Identities and Composition

Similarly, the identities and composition of  $\text{Mon}(\text{Sets}_*, \triangleleft, S^0)$  can be easily seen to agree with those of monoids with left zero, which finishes the proof. 

## 00EJ 7.4 The Right Tensor Product of Pointed Sets

### 00EK 7.4.1 Foundations

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

#### 00EL DEFINITION 7.4.1.1.1 ► THE RIGHT TENSOR PRODUCT OF POINTED SETS

The **right tensor product of pointed sets** is the functor<sup>1</sup>

$$\triangleright : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{Forget} \times \text{id}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*,$$

where:

- $\text{Forget} : \text{Sets}_* \rightarrow \text{Sets}$  is the forgetful functor from pointed sets to sets.
- $\odot : \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*$  is the tensor functor of **Item 1** of [Proposition 7.2.1.1.9](#).

<sup>1</sup>Further Notation: Also written  $\triangleright_{\text{Sets}_*}$ .

#### 00EM REMARK 7.4.1.1.2 ► UNWINDING DEFINITION 7.4.1.1.1: UNIVERSAL PROPERTY I

The right tensor product of pointed sets satisfies the following natural bijection:

$$\text{Sets}_*(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z).$$

That is to say, the following data are in natural bijection:

#### 026T

1. Pointed maps  $f : X \triangleright Y \rightarrow Z$ .

#### 026U

2. Maps of sets  $f : X \times Y \rightarrow Z$  satisfying  $f(x, y_0) = z_0$  for each  $x \in X$ .

#### 00EN

#### REMARK 7.4.1.1.3 ► UNWINDING DEFINITION 7.4.1.1.1: UNIVERSAL PROPERTY II

The right tensor product of pointed sets may be described as follows:

- The right tensor product of  $(X, x_0)$  and  $(Y, y_0)$  is the pair  $((X \triangleright Y, x_0 \triangleright y_0), \iota)$  consisting of
  - A pointed set  $(X \triangleright Y, x_0 \triangleright y_0)$ ;
  - A right bilinear morphism of pointed sets  $\iota : (X \times Y, (x_0, y_0)) \rightarrow X \triangleright Y$ ;

satisfying the following universal property:

- (★) Given another such pair  $((Z, z_0), f)$  consisting of
- \* A pointed set  $(Z, z_0)$ ;
  - \* A right bilinear morphism of pointed sets  $f: (X \times Y, (x_0, y_0)) \rightarrow X \triangleright Y$ ;

there exists a unique morphism of pointed sets  $X \triangleright Y \xrightarrow{\exists!} Z$  making the diagram

$$\begin{array}{ccc} & X \triangleright Y & \\ l \nearrow & \downarrow & \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

00EP

#### CONSTRUCTION 7.4.1.1.4 ► THE RIGHT TENSOR PRODUCT OF POINTED SETS

In detail, the **right tensor product** of  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $(X \triangleright Y, [y_0])$  consisting of:

- *The Underlying Set.* The set  $X \triangleright Y$  defined by

$$\begin{aligned} X \triangleright Y &\stackrel{\text{def}}{=} |X| \odot Y \\ &\cong \bigvee_{x \in X} (Y, y_0), \end{aligned}$$

where  $|X|$  denotes the underlying set of  $(X, x_0)$ .

- *The Underlying Basepoint.* The point  $[(x_0, y_0)]$  of  $\bigvee_{x \in X} (Y, y_0)$ , which is equal to  $[(x, y_0)]$  for any  $x \in X$ .

#### PROOF 7.4.1.1.5 ► PROOF OF CONSTRUCTION 7.4.1.1.4

Since  $\bigvee_{y \in Y} (X, x_0)$  is defined as the quotient of  $\coprod_{x \in X} Y$  by the equivalence relation  $R$  generated by declaring  $(x, y) \sim (x', y')$  if  $y = y' = y_0$ , we have, by ??, a natural bijection

$$\text{Sets}_*(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}}^R(\coprod_{X \in X} Y, Z),$$

where  $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$  is the set

$$\text{Hom}_{\text{Sets}}^R(\coprod_{x \in X} Y, Z) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(\coprod_{x \in X} Y, Z) \mid \begin{array}{l} \text{for each } x, y \in X, \text{ if} \\ (x, y) \sim_R (x', y'), \text{ then} \\ f(x, y) = f(x', y') \end{array} \right\}.$$

However, the condition  $(x, y) \sim_R (x', y')$  only holds when:

- 026V** 1. We have  $x = x'$  and  $y = y'$ .
- 026W** 2. We have  $y = y' = y_0$ .

So, given  $f \in \text{Hom}_{\text{Sets}}(\coprod_{x \in X} Y, Z)$  with a corresponding  $\bar{f}: X \triangleright Y \rightarrow Z$ , the latter case above implies

$$\begin{aligned} f([(x, y_0)]) &= f([(x', y_0)]) \\ &= f([(x_0, y_0)]), \end{aligned}$$

and since  $\bar{f}: X \triangleright Y \rightarrow Z$  is a pointed map, we have

$$\begin{aligned} f([(x_0, y_0)]) &= \bar{f}([(x_0, y_0)]) \\ &= z_0. \end{aligned}$$

Thus the elements  $f$  in  $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$  are precisely those functions  $f: X \times Y \rightarrow Z$  satisfying the equality

$$f(x, y_0) = z_0$$

for each  $y \in Y$ , giving an equality

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) = \text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z)$$

of sets, which when composed with our earlier isomorphism

$$\text{Sets}_*(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z),$$

gives our desired natural bijection, finishing the proof. 

**00EQ**

#### NOTATION 7.4.1.1.6 ► ELEMENTS OF RIGHT TENSOR PRODUCTS OF POINTED SETS

We write<sup>1</sup>  $x \triangleright y$  for the element  $[(x, y)]$  of

$$X \triangleright Y \cong |X| \odot Y.$$

<sup>1</sup>Further Notation: Also written  $x \triangleright_{\text{Sets}_*} y$ .

00ER

**REMARK 7.4.1.1.7 ► BASEPOINTS OF RIGHT TENSOR PRODUCTS OF POINTED SETS**

Employing the notation introduced in [Notation 7.4.1.1.6](#), we have

$$x_0 \triangleright y_0 = x \triangleright y_0$$

for each  $x \in X$ , and

$$x \triangleright y_0 = x' \triangleright y_0$$

for each  $x, x' \in X$ .

00ES

**PROPOSITION 7.4.1.1.8 ► PROPERTIES OF RIGHT TENSOR PRODUCTS OF POINTED SETS**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00ET

- Functoriality.* The assignments  $X, Y, (X, Y) \mapsto X \triangleright Y$  define functors

$$\begin{aligned} X \triangleright - : \quad \text{Sets}_* &\rightarrow \text{Sets}_*, \\ - \triangleright Y : \quad \text{Sets}_* &\rightarrow \text{Sets}_*, \\ -_1 \triangleright -_2 : \text{Sets}_* \times \text{Sets}_* &\rightarrow \text{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f : (X, x_0) &\rightarrow (A, a_0), \\ g : (Y, y_0) &\rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \triangleright g : X \triangleright Y \rightarrow A \triangleright B$$

is given by

$$[f \triangleright g](x \triangleright y) \stackrel{\text{def}}{=} f(x) \triangleright g(y)$$

for each  $x \triangleright y \in X \triangleright Y$ .

00EU

- Adjointness I.* We have an adjunction

$$\left( X \triangleright - \dashv [X, -]_{\text{Sets}_*}^\triangleright \right) : \quad \text{Sets}_* \begin{array}{c} \xrightarrow{X \triangleright -} \\ \perp \\ \xleftarrow{[X, -]_{\text{Sets}_*}^\triangleright} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}_*}(Y, [X, Z]_{\text{Sets}_*}^\triangleright)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ , where  $[X, Y]_{\text{Sets}_*}^\triangleright$  is the pointed set of [Definition 7.4.2.1.1](#).

00EV

3. *Adjointness II.* The functor

$$- \triangleright Y : \text{Sets}_* \rightarrow \text{Sets}_*$$

does not admit a right adjoint.

00EW

4. *Adjointness III.* We have a  $\perp_{\text{忘}}$ -relative adjunction

$$(- \triangleright Y \dashv \text{Sets}_*(Y, -)) : \text{Sets}_* \begin{array}{c} \xrightarrow{- \triangleright Y} \\ \perp_{\text{忘}} \\ \xleftarrow{\text{Sets}_*(Y, -)} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}}(|X|, \text{Sets}_*(Y, Z))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ .

## PROOF 7.4.1.1.9 ► PROOF OF PROPOSITION 7.4.1.1.8

## Item 1: Functoriality

This follows from the definition of  $\triangleright$  as a composition of functors ([Definition 7.4.1.1.1](#)).

## Item 2: Adjointness I

This follows from [Item 3 of Proposition 7.2.1.1.9](#).

## Item 3: Adjointness II

For  $- \triangleright Y$  to admit a right adjoint would require it to preserve colimits by ?? of ???. However, we have

$$\begin{aligned} \text{pt} \triangleright X &\stackrel{\text{def}}{=} |\text{pt}| \odot X \\ &\cong X \\ &\not\cong \text{pt}, \end{aligned}$$

and thus we see that  $- \triangleright Y$  does not have a right adjoint.

## Item 4: Adjointness III

This follows from [Item 2 of Proposition 7.2.1.1.9](#). 

00EX

REMARK 7.4.1.1.10 ► ON THE FAILURE OF  $- \triangleright Y$  TO BE A LEFT ADJOINT

Here is some intuition on why  $- \triangleright Y$  fails to be a left adjoint. [Item 4 of Proposition 7.3.1.1.8](#) states that we have a natural bijection

$$\text{Hom}_{\text{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}}(|X|, \text{Sets}_*(Y, Z)),$$

so it would be reasonable to wonder whether a natural bijection of the form

$$\text{Hom}_{\text{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}_*}(X, \text{Sets}_*(Y, Z)),$$

also holds, which would give  $- \triangleright Y \dashv \text{Sets}_*(Y, -)$ . However, such a bijection would require every map

$$f: X \triangleright Y \rightarrow Z$$

to satisfy

$$f(x_0 \triangleright y) = z_0$$

for each  $x \in X$ , whereas we are imposing such a basepoint preservation condition only for elements of the form  $x \triangleright y_0$ . Thus  $\text{Sets}_*(Y, -)$  can't be a right adjoint for  $- \triangleright Y$ , and as shown by Item 3 of Proposition 7.4.1.1.8, no functor can.<sup>1</sup>

<sup>1</sup>The functor  $\text{Sets}_*(Y, -)$  is instead right adjoint to  $- \wedge Y$ , the smash product of pointed sets of Definition 7.5.1.1.1. See Item 2 of Proposition 7.5.1.1.12.

## 00EY 7.4.2 The Right Internal Hom of Pointed Sets

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00EZ

### DEFINITION 7.4.2.1.1 ► THE RIGHT INTERNAL HOM OF POINTED SETS

The **right internal Hom**<sup>1</sup> of pointed sets is the functor

$$[-, -]_{\text{Sets}_*}^\triangleright : \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_*^{\text{op}} \times \text{Sets}_* \xrightarrow{\text{忘} \times \text{id}} \text{Sets}_*^{\text{op}} \times \text{Sets}_* \xrightarrow{\pitchfork} \text{Sets}_*,$$

where:

- 忘:  $\text{Sets}_* \rightarrow \text{Sets}$  is the forgetful functor from pointed sets to sets.
- $\pitchfork: \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$  is the cotensor functor of Item 1 of Proposition 7.2.2.1.6.

<sup>1</sup>For a proof that  $[-, -]_{\text{Sets}_*}^\triangleright$  is indeed the right internal Hom of  $\text{Sets}_*$  with respect to the right tensor product of pointed sets, see Item 2 of Proposition 7.4.1.1.8.

00F0

### REMARK 7.4.2.1.2 ► UNWINDING DEFINITION 7.4.2.1.1, I: COMPARISON WITH $[-, -]_{\text{Sets}_*}^\triangleleft$

We have

$$[-, -]_{\text{Sets}_*}^\triangleleft = [-, -]_{\text{Sets}_*}^\triangleright.$$

00F1

The right internal Hom of pointed sets satisfies the following universal property:

$$\text{Sets}_*(X \triangleright Y, Z) \cong \text{Sets}_*(Y, [X, Z]_{\text{Sets}_*}^\triangleright)$$

That is to say, the following data are in bijection:

026X

1. Pointed maps  $f: X \triangleright Y \rightarrow Z$ .

026Y

2. Pointed maps  $f: Y \rightarrow [X, Z]_{\text{Sets}_*}^\triangleright$ .

00F2

#### REMARK 7.4.2.1.4 ► UNWINDING DEFINITION 7.4.2.1.1, III: EXPLICIT DESCRIPTION

In detail, the **right internal Hom** of  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $([X, Y]_{\text{Sets}_*}^\triangleright, [(y_0)_{x \in X}])$  consisting of:

- *The Underlying Set.* The set  $[X, Y]_{\text{Sets}_*}^\triangleright$  defined by

$$\begin{aligned} [X, Y]_{\text{Sets}_*}^\triangleright &\stackrel{\text{def}}{=} |X| \pitchfork Y \\ &\cong \bigwedge_{x \in X} (Y, y_0), \end{aligned}$$

where  $|X|$  denotes the underlying set of  $(X, x_0)$ .

- *The Underlying Basepoint.* The point  $[(y_0)_{x \in X}]$  of  $\wedge_{x \in X} (Y, y_0)$ .

00F3

#### PROPOSITION 7.4.2.1.5 ► PROPERTIES OF RIGHT INTERNAL HOMS OF POINTED SETS

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00F4

1. *Functionality.* The assignments  $X, Y, (X, Y) \mapsto [X, Y]_{\text{Sets}_*}^\triangleright$  define functors

$$\begin{aligned} [X, -]_{\text{Sets}_*}^\triangleright : \text{Sets}_* &\rightarrow \text{Sets}_*, \\ [-, Y]_{\text{Sets}_*}^\triangleright : \text{Sets}_*^{\text{op}} &\rightarrow \text{Sets}_*, \\ [-_1, -_2]_{\text{Sets}_*}^\triangleright : \text{Sets}_*^{\text{op}} \times \text{Sets}_* &\rightarrow \text{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f : (X, x_0) &\rightarrow (A, a_0), \\ g : (Y, y_0) &\rightarrow (B, b_0), \end{aligned}$$

the induced map

$$[f, g]_{\text{Sets}_*}^\triangleright : [A, Y]_{\text{Sets}_*}^\triangleright \rightarrow [X, B]_{\text{Sets}_*}^\triangleright$$

is given by

$$[f, g]_{\text{Sets}_*}^\triangleright([(y_a)_{a \in A}]) \stackrel{\text{def}}{=} [(g(y_{f(x)}))_{x \in X}]$$

for each  $[(y_a)_{a \in A}] \in [A, Y]_{\text{Sets}_*}^\triangleright$ .

00F5 2. *Adjointness I.* We have an adjunction

$$\left( X \triangleright - \dashv [X, -]_{\text{Sets}_*}^\triangleright \right): \text{Sets}_* \begin{array}{c} \xrightarrow{X \triangleright -} \\ \perp \\ \xleftarrow{[X, -]_{\text{Sets}_*}^\triangleright} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}_*}(Y, [X, Z]_{\text{Sets}_*}^\triangleright)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ , where  $[X, Y]_{\text{Sets}_*}^\triangleright$  is the pointed set of [Definition 7.4.2.1.1](#).

00F6 3. *Adjointness II.* The functor

$$- \triangleright Y: \text{Sets}_* \rightarrow \text{Sets}_*$$

does not admit a right adjoint.

#### PROOF 7.4.2.1.6 ▶ PROOF OF PROPOSITION 7.4.2.1.5

##### Item 1: Functoriality

This follows from the definition of  $[-, -]_{\text{Sets}_*}^\triangleright$  as a composition of functors ([Definition 7.4.2.1.1](#)).

##### Item 2: Adjointness I

This is a repetition of [Item 2](#) of [Proposition 7.4.1.1.8](#), and is proved there.

##### Item 3: Adjointness II

This is a repetition of [Item 3](#) of [Proposition 7.4.1.1.8](#), and is proved there. 

### 00F7 7.4.3 The Right Skew Unit

00F8

#### DEFINITION 7.4.3.1.1 ▶ THE RIGHT SKEW UNIT OF $\triangleright$

The **right skew unit of the right tensor product of pointed sets** is the functor

$$\mathbb{1}_{\text{Sets}_*}^{\triangleright}: \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*}^{\triangleright} \stackrel{\text{def}}{=} S^0.$$

**00F9 7.4.4 The Right Skew Associator**

00FA

DEFINITION 7.4.4.1.1 ► THE RIGHT SKEW ASSOCIATOR OF  $\triangleright$ 

The **skew associator of the right tensor product of pointed sets** is the natural transformation

$$\alpha^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id}_{\text{Sets}_*} \times \triangleright) \Rightarrow \triangleright \circ (\triangleright \times \text{id}_{\text{Sets}_*}) \circ \alpha^{\text{Cats}, -1}_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}$$

as in the diagram

$$\begin{array}{ccc}
 & (\text{Sets}_* \times \text{Sets}_*) \times \text{Sets}_* & \\
 \alpha^{\text{Cats}, -1}_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*} \nearrow & \swarrow \triangleright \times \text{id} & \\
 \text{Sets}_* \times (\text{Sets}_* \times \text{Sets}_*) & & \text{Sets}_* \times \text{Sets}_* \\
 \downarrow \text{id} \times \triangleright & \quad \quad \quad \alpha^{\text{Sets}_*, \triangleright} \quad \quad \quad \downarrow \triangleright & \\
 \text{Sets}_* \times \text{Sets}_* & \xrightarrow{\triangleright} & \text{Sets}_*,
 \end{array}$$

whose component

$$\alpha_{X, Y, Z}^{\text{Sets}_*, \triangleright} : X \triangleright (Y \triangleright Z) \rightarrow (X \triangleright Y) \triangleright Z$$

at  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\begin{aligned}
 X \triangleright (Y \triangleright Z) &\stackrel{\text{def}}{=} |X| \odot (Y \triangleright Z) \\
 &\stackrel{\text{def}}{=} |X| \odot (|Y| \odot Z) \\
 &\cong \bigvee_{x \in X} (|Y| \odot Z) \\
 &\cong \bigvee_{x \in X} (\bigvee_{y \in Y} Z) \\
 &\rightarrow \bigvee_{[(x, y)] \in \bigvee_{x \in X} Y} Z \\
 &\cong \bigvee_{[(x, y)] \in |X| \odot Y} Z \\
 &\cong ||X| \odot Y| \odot Z \\
 &\stackrel{\text{def}}{=} |X \triangleright Y| \odot Z \\
 &\stackrel{\text{def}}{=} (X \triangleright Y) \triangleright Z,
 \end{aligned}$$

where the map

$$\bigvee_{x \in X} (\bigvee_{y \in Y} Z) \rightarrow \bigvee_{[(x, y)] \in \bigvee_{x \in X} Y} Z$$

is given by  $[(x, [(y, z)])] \mapsto [([(x, y)], z)]$ .

**PROOF 7.4.4.1.2 ▶ PROOF OF DEFINITION 7.4.4.1.1**

(Proven below in a bit.)

**00FB****REMARK 7.4.4.1.3 ▶ UNWINDING DEFINITION 7.4.4.1.1**

Unwinding the notation for elements, we have

$$\begin{aligned} [(x, [(y, z)])] &\stackrel{\text{def}}{=} [(x, y \triangleright z)] \\ &\stackrel{\text{def}}{=} x \triangleright (y \triangleright z) \end{aligned}$$

and

$$\begin{aligned} [([(x, y)], z)] &\stackrel{\text{def}}{=} [(x \triangleright y, z)] \\ &\stackrel{\text{def}}{=} (x \triangleright y) \triangleright z. \end{aligned}$$

So, in other words,  $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}$  acts on elements via

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}(x \triangleright (y \triangleright z)) \stackrel{\text{def}}{=} (x \triangleright y) \triangleright z$$

for each  $x \triangleright (y \triangleright z) \in X \triangleright (Y \triangleright Z)$ .

**00FC****REMARK 7.4.4.1.4 ▶ NON-INVERTIBILITY OF THE SKew ASSOCIATOR OF  $\triangleright$** 

Taking  $y = y_0$ , we see that the morphism  $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}$  acts on elements as

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}(x \triangleright (y_0 \triangleright z)) \stackrel{\text{def}}{=} (x \triangleright y_0) \triangleright z.$$

However, by the definition of  $\triangleright$ , we have  $x \triangleright y_0 = x' \triangleright y_0$  for all  $x, x' \in X$ , preventing  $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}$  from being non-invertible.

**PROOF 7.4.4.1.5 ▶ PROOF OF DEFINITION 7.4.4.1.1**

Firstly, note that, given  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}: X \triangleright (Y \triangleright Z) \rightarrow (X \triangleright Y) \triangleright Z$$

is indeed a morphism of pointed sets, as we have

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}(x_0 \triangleright (y_0 \triangleright z_0)) = (x_0 \triangleright y_0) \triangleright z_0.$$

Next, we claim that  $\alpha^{\text{Sets}_*, \triangleright}$  is a natural transformation. We need to show that, given morphisms of pointed sets

$$f: (X, x_0) \rightarrow (X', x'_0),$$

$$\begin{aligned} g: (Y, y_0) &\rightarrow (Y', y'_0), \\ h: (Z, z_0) &\rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} X \triangleright (Y \triangleright Z) & \xrightarrow{f \triangleright (g \triangleright h)} & X' \triangleright (Y' \triangleright Z') \\ \alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*, \triangleright} \\ (X \triangleright Y) \triangleright Z & \xrightarrow{(f \triangleright g) \triangleright h} & (X' \triangleright Y') \triangleright Z' \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleright (y \triangleright z) & \longmapsto & f(x) \triangleright (g(y) \triangleright h(z)) \\ \downarrow & & \downarrow \\ (x \triangleright y) \triangleright z & \longmapsto & (f(x) \triangleright g(y)) \triangleright h(z) \end{array}$$

and hence indeed commutes, showing  $\alpha^{\text{Sets}_*, \triangleright}$  to be a natural transformation. This finishes the proof. □

#### 00FD 7.4.5 The Right Skew Left Unit

00FE

##### DEFINITION 7.4.5.1.1 ► THE RIGHT SKEW LEFT UNIT OF $\triangleright$

The **skew left unit** of the right tensor product of pointed sets is the natural transformation

$$\begin{array}{ccc} \text{pt} \times \text{Sets}_* & \xrightarrow{\mathbb{1}_{\text{Sets}_*} \times \text{id}} & \text{Sets}_* \times \text{Sets}_* \\ \lambda^{\text{Sets}_*, \triangleright} : \lambda_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleright \circ (\mathbb{1}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) & \swarrow \quad \searrow & \downarrow \triangleright \\ \lambda_{\text{Sets}_*}^{\text{Cats}_2} & & \text{Sets}_*, \end{array}$$

whose component

$$\lambda_X^{\text{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright X$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by the composition

$$X \rightarrow X \vee X$$

$$\begin{aligned} &\cong |S^0| \odot X \\ &\cong S^0 \triangleright X, \end{aligned}$$

where  $X \rightarrow X \vee X$  is the map sending  $X$  to the second factor of  $X$  in  $X \vee X$ .

**PROOF 7.4.5.1.2 ▶ PROOF OF DEFINITION 7.4.5.1.1**

(Proven below in a bit.)



00FF

**REMARK 7.4.5.1.3 ▶ UNWINDING DEFINITION 7.4.5.1.1**

In other words,  $\lambda_X^{\text{Sets}_*, \triangleright}$  acts on elements as

$$\lambda_X^{\text{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} [(1, x)]$$

i.e. by

$$\lambda_X^{\text{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} 1 \triangleright x$$

for each  $x \in X$ .

00FG

**REMARK 7.4.5.1.4 ▶ NON-INVERTIBILITY OF THE SKEW LEFT UNIT OF  $\triangleright$**

The morphism  $\lambda_X^{\text{Sets}_*, \triangleright}$  is non-invertible, as it is non-surjective when viewed as a map of sets, since the elements  $0 \triangleright x$  of  $S^0 \triangleright X$  with  $x \neq x_0$  are outside the image of  $\lambda_X^{\text{Sets}_*, \triangleright}$ , which sends  $x$  to  $1 \triangleright x$ .

**PROOF 7.4.5.1.5 ▶ PROOF OF DEFINITION 7.4.5.1.1**

Firstly, note that, given  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\lambda_X^{\text{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright X$$

is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \lambda_X^{\text{Sets}_*, \triangleright}(x_0) &= 1 \triangleright x_0 \\ &= 0 \triangleright x_0. \end{aligned}$$

Next, we claim that  $\lambda^{\text{Sets}_*, \triangleright}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \lambda_X^{\text{Sets}_*, \triangleright} \downarrow & & \downarrow \lambda_Y^{\text{Sets}_*, \triangleright} \\ S^0 \triangleright X & \xrightarrow{\text{id}_{S^0 \triangleright f}} & S^0 \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \mapsto & f(x) \\ \downarrow & & \downarrow \\ 1 \triangleright x & \mapsto & 1 \triangleright f(x) \end{array}$$

and hence indeed commutes, showing  $\lambda^{\text{Sets}_*, \triangleright}$  to be a natural transformation. This finishes the proof. □

#### 00FH 7.4.6 The Right Skew Right Unitor

00FJ

##### DEFINITION 7.4.6.1.1 ► THE RIGHT SKEW RIGHT UNITOR OF $\triangleright$

The **skew right unitor of the right tensor product of pointed sets** is the natural transformation

$$\begin{array}{ccc} \text{Sets}_* \times \text{pt} & \xrightarrow{\text{id} \times 1^{\text{Sets}_*}} & \text{Sets}_* \times \text{Sets}_* \\ \rho^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id} \times 1^{\text{Sets}_*}) \xrightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}, & \swarrow \rho_{\text{Sets}_*}^{\text{Cats}_2} & \downarrow \triangleright \\ & & \text{Sets}_*, \end{array}$$

whose component

$$\rho_X^{\text{Sets}_*, \triangleright} : X \triangleright S^0 \rightarrow X$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by the composition

$$\begin{aligned} X \triangleright S^0 &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X, \end{aligned}$$

where  $\vee_{x \in X} S^0 \rightarrow X$  is the map given by

$$\begin{aligned} [(x, 0)] &\mapsto x_0, \\ [(x, 1)] &\mapsto x \end{aligned}$$

for each  $x \in X$ .

#### PROOF 7.4.6.1.2 ▶ PROOF OF DEFINITION 7.4.6.1.1

(Proven below in a bit.)



00FK

#### REMARK 7.4.6.1.3 ▶ UNWINDING DEFINITION 7.4.6.1.1

In other words,  $\rho_X^{\text{Sets}_*, \triangleright}$  acts on elements as

$$\begin{aligned} \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 0) &\stackrel{\text{def}}{=} x_0, \\ \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 1) &\stackrel{\text{def}}{=} x \end{aligned}$$

for each  $x \triangleright 1 \in X \triangleright S^0$ .

00FL

#### REMARK 7.4.6.1.4 ▶ NON-INVERTIBILITY OF THE SKEW RIGHT UNIT OF $\triangleright$

The morphism  $\rho_X^{\text{Sets}_*, \triangleright}$  is almost invertible, with its would-be-inverse

$$\phi_X : X \rightarrow X \triangleright S^0$$

given by

$$\phi_X(x) \stackrel{\text{def}}{=} x \triangleright 1$$

for each  $x \in X$ . Indeed, we have

$$\begin{aligned} [\rho_X^{\text{Sets}_*, \triangleright} \circ \phi](x) &= \rho_X^{\text{Sets}_*, \triangleright}(\phi(x)) \\ &= \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 1) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

so that

$$\rho_X^{\text{Sets}_*, \triangleright} \circ \phi = \text{id}_X$$

and

$$\begin{aligned} [\phi \circ \rho_X^{\text{Sets}_*, \triangleright}](x \triangleright 1) &= \phi(\rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 1)) \\ &= \phi(x) \\ &= x \triangleright 1 \end{aligned}$$

$$= [\text{id}_{X \triangleright S^0}](x \triangleright 1),$$

but

$$\begin{aligned} [\phi \circ \rho_X^{\text{Sets}_*, \triangleright}](x \triangleright 0) &= \phi(\rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 0)) \\ &= \phi(x_0) \\ &= 1 \triangleright x_0, \end{aligned}$$

where  $x \triangleright 0 \neq 1 \triangleright x_0$ . Thus

$$\phi \circ \rho_X^{\text{Sets}_*, \triangleright} \stackrel{?}{=} \text{id}_{X \triangleright S^0}$$

holds for all elements in  $X \triangleright S^0$  except one.

#### PROOF 7.4.6.1.5 ► PROOF OF DEFINITION 7.4.6.1.1

Firstly, note that, given  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\rho_X^{\text{Sets}_*, \triangleright}: X \triangleright S^0 \rightarrow X$$

is indeed a morphism of pointed sets as we have

$$\rho_X^{\text{Sets}_*, \triangleright}(x_0 \triangleright 0) = x_0.$$

Next, we claim that  $\rho^{\text{Sets}_*, \triangleright}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X \triangleright S^0 & \xrightarrow{f \triangleright \text{id}_{S^0}} & Y \triangleright S^0 \\ \rho_X^{\text{Sets}_*, \triangleright} \downarrow & & \downarrow \rho_Y^{\text{Sets}_*, \triangleright} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleright 0 & & x \triangleright 0 \longmapsto f(x) \triangleright 0 \\ \downarrow & & \downarrow \\ x_0 \longmapsto f(x_0) & & y_0 \end{array}$$

and

$$\begin{array}{ccc} x \triangleright 1 & \longmapsto & f(x) \triangleright 1 \\ \downarrow & & \downarrow \\ x & \longmapsto & f(x) \end{array}$$

and hence indeed commutes, showing  $\rho^{\text{Sets}_*, \triangleright}$  to be a natural transformation. This finishes the proof. 

### 00FM 7.4.7 The Diagonal

00FN

#### DEFINITION 7.4.7.1.1 ► THE DIAGONAL OF $\triangleright$

The **diagonal of the right tensor product of pointed sets** is the natural transformation

$$\Delta^\triangleright : \text{id}_{\text{Sets}_*} \Longrightarrow \triangleright \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2}, \quad \begin{array}{ccc} \text{Sets}_* & \xrightarrow{\text{id}_{\text{Sets}_*}} & \text{Sets}_* \\ \Delta_{\text{Sets}_*}^{\text{Cats}_2} \swarrow & \Downarrow \Delta^\triangleright & \searrow \triangleright \\ \text{Sets}_* \times \text{Sets}_*, & & \end{array}$$

whose component

$$\Delta_X^\triangleright : (X, x_0) \rightarrow (X \triangleright X, x_0 \triangleright x_0)$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\Delta_X^\triangleright(x) \stackrel{\text{def}}{=} x \triangleright x$$

for each  $x \in X$ .

#### PROOF 7.4.7.1.2 ► PROOF OF DEFINITION 7.4.7.1.1

##### Being a Morphism of Pointed Sets

We have

$$\Delta_X^\triangleright(x_0) \stackrel{\text{def}}{=} x_0 \triangleright x_0,$$

and thus  $\Delta_X^\triangleright$  is a morphism of pointed sets.

##### Naturality

We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\triangleright \downarrow & & \downarrow \Delta_Y^\triangleright \\ X \triangleright X & \xrightarrow{f \triangleright f} & Y \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ x \triangleright x & \longmapsto & f(x) \triangleright f(x) \end{array}$$

and hence indeed commutes, showing  $\Delta^\triangleright$  to be natural. 

### 00FP 7.4.8 The Right Skew Monoidal Structure on Pointed Sets Associated to $\triangleright$

00FQ

#### PROPOSITION 7.4.8.1.1 ► THE RIGHT SKEW MONOIDAL STRUCTURE ON POINTED SETS ASSOCIATED TO $\triangleright$

The category  $\text{Sets}_*$  admits a right-closed right skew monoidal category structure consisting of:

- *The Underlying Category.* The category  $\text{Sets}_*$  of pointed sets.
- *The Right Skew Monoidal Product.* The right tensor product functor

$$\triangleright: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Definition 7.4.1.1.1.

- *The Right Internal Skew Hom.* The right internal Hom functor

$$[-, -]_{\text{Sets}_*}^\triangleright: \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Definition 7.4.2.1.1.

- *The Right Skew Monoidal Unit.* The functor

$$\mathbb{1}_{\text{Sets}_*, \triangleright}: \text{pt} \rightarrow \text{Sets}_*$$

of Definition 7.4.3.1.1.

- *The Right Skew Associators.* The natural transformation

$$\alpha^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id}_{\text{Sets}_*} \times \triangleright) \Rightarrow \triangleright \circ (\triangleright \times \text{id}_{\text{Sets}_*}) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}, -1}$$

of [Definition 7.4.4.1.1](#).

- *The Right Skew Left Unitors.* The natural transformation

$$\lambda^{\text{Sets}_*, \triangleright} : \lambda_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleright \circ (\mathbb{1}^{\text{Sets}_*} \times \text{id}_{\text{Sets}_*})$$

of [Definition 7.4.5.1.1](#).

- *The Right Skew Right Unitors.* The natural transformation

$$\rho^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}) \xrightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}$$

of [Definition 7.4.6.1.1](#).

#### PROOF 7.4.8.1.2 ► PROOF OF PROPOSITION 7.4.8.1.1

##### The Pentagon Identity

Let  $(W, w_0)$ ,  $(X, x_0)$ ,  $(Y, y_0)$  and  $(Z, z_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 & W \triangleright ((X \triangleright Y) \triangleright Z) & \\
 & \swarrow \alpha_{W,X,Y}^{\text{Sets}_*, \triangleright} \circ \text{id}_Z & \searrow \alpha_{W,X \triangleright Y,Z}^{\text{Sets}_*, \triangleright} \\
 W \triangleright (X \triangleright (Y \triangleright Z)) & & (W \triangleright (X \triangleright Y)) \triangleright Z \\
 & \swarrow \alpha_{W \triangleright X,Y,Z}^{\text{Sets}_*, \triangleright} & \searrow \text{id}_W \triangleright \alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright} \\
 & (W \triangleright X) \triangleright (Y \triangleright Z) & \xrightarrow{\alpha_{W,X,Y \triangleright Z}^{\text{Sets}_*, \triangleright}} ((W \triangleright X) \triangleright Y) \triangleright Z
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccccc}
 & & w \triangleright ((x \triangleright y) \triangleright z) & & \\
 & \nearrow & & \searrow & \\
 w \triangleright (x \triangleright (y \triangleright z)) & & & & (w \triangleright (x \triangleright y)) \triangleright z \\
 \downarrow & & & & \downarrow \\
 (w \triangleright x) \triangleright (y \triangleright z) & \longleftarrow & & & ((w \triangleright x) \triangleright y) \triangleright z
 \end{array}$$

and thus we see that the pentagon identity is satisfied.

#### The Right Skew Left Triangle Identity

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright Y & & \\
 \downarrow \lambda_{X \triangleright Y}^{\text{Sets}_*, \triangleright} & \searrow \lambda_X^{\text{Sets}_*, \triangleright} \triangleright \text{id}_Y & \\
 S^0 \triangleright (X \triangleright Y) & \xrightarrow[\alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleright}]{} & (S^0 \triangleright X) \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright y & & \\
 \downarrow & \swarrow & \\
 1 \triangleright (x \triangleright y) & \longleftarrow & (1 \triangleright x) \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the left skew triangle identity is satisfied.

#### The Right Skew Right Triangle Identity

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc} X \triangleright (Y \triangleright S^0) & \xrightarrow{\text{id}_X \triangleright \rho_Y^{\text{Sets}_*, \triangleright}} & (X \triangleright Y) \triangleright S^0 \\ & \searrow \alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleright} & \downarrow \rho_{X \triangleright Y}^{\text{Sets}_*, \triangleright} \\ & & X \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleright (y \triangleright 0) & \longmapsto & (x \triangleright y) \triangleright 0 \\ \swarrow & & \downarrow \\ & x \triangleright y_0 = x_0 \triangleright y_0 & \end{array}$$

and

$$\begin{array}{ccc} x \triangleright (y \triangleright 1) & \longmapsto & (x \triangleright y) \triangleright 1 \\ \swarrow & & \downarrow \\ & x \triangleright y & \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

### The Right Skew Middle Triangle Identity

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc} X \triangleright Y & \xlongequal{\quad} & X \triangleright Y \\ \text{id}_X \triangleright \lambda_Y^{\text{Sets}_*, \triangleright} \downarrow & & \uparrow \rho_X^{\text{Sets}_*, \triangleright \triangleright \text{id}_Y} \\ X \triangleright (S^0 \triangleright Y) & \xrightarrow{\alpha_{X, S^0, Y}^{\text{Sets}_*, \triangleright}} & (X \triangleright S^0) \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleright y & \longleftrightarrow & x \triangleright y \\ \downarrow & & \uparrow \\ x \triangleright (1 \triangleright y) & \longmapsto & (x \triangleright 1) \triangleright y \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

The Zig-Zag Identity

We have to show that the diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{\lambda_{S^0}^{\text{Sets}_*, \triangleright}} & S^0 \triangleright S^0 \\ & \searrow & \downarrow \rho_{S^0}^{\text{Sets}_*, \triangleright} \\ & & S^0 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 & \xrightarrow{\quad} & 1 \triangleright 0 \\ \swarrow & & \downarrow \\ & 0 & \end{array}$$

and

$$\begin{array}{ccc} 1 & \xrightarrow{\quad} & 1 \triangleright 1 \\ \swarrow & & \downarrow \\ & 1 & \end{array}$$

and hence indeed commutes. Thus the zig-zag identity is satisfied.

Right Skew Monoidal Right-Closedness

This follows from Item 2 of Proposition 7.4.1.1.8. 

**00FR 7.4.9 Monoids With Respect to the Right Tensor Product of Pointed Sets**

00FS

**PROPOSITION 7.4.9.1.1 ▶ MONOIDS WITH RESPECT TO  $\triangleright$**

The category of monoids on  $(\text{Sets}_*, \triangleright, S^0)$  is isomorphic to the category of “monoids with right zero”<sup>1</sup> and morphisms between them.

<sup>1</sup>A monoid with right zero is defined similarly as the monoids with zero of ??.  
Succinctly, they are monoids  $(A, \mu_A, \eta_A)$  with a special element  $0_A$  satisfying

$$0_A a = 0_A$$

for each  $a \in A$ .

## PROOF 7.4.9.1.2 ► PROOF OF PROPOSITION 7.4.9.1.1

Monoids on  $(\text{Sets}_*, \triangleright, S^0)$

A monoid on  $(\text{Sets}_*, \triangleright, S^0)$  consists of:

- *The Underlying Object.* A pointed set  $(A, 0_A)$ .
- *The Multiplication Morphism.* A morphism of pointed sets

$$\mu_A: A \triangleright A \rightarrow A,$$

determining a right bilinear morphism of pointed sets

$$\begin{aligned} A \times A &\longrightarrow A \\ (a, b) &\longmapsto ab. \end{aligned}$$

- *The Unit Morphism.* A morphism of pointed sets

$$\eta_A: S^0 \rightarrow A$$

picking an element  $1_A$  of  $A$ .

satisfying the following conditions:

026Z

1. *Associativity.* The diagram

$$\begin{array}{ccc} & A \triangleright (A \triangleright A) & \\ \alpha_{A,A,A}^{\text{Sets}_*, \triangleright} \nearrow & & \searrow \text{id}_A \triangleright \mu_A \\ (A \triangleright A) \triangleright A & & A \triangleright A \\ \downarrow \mu_A \triangleright \text{id}_A & & \downarrow \mu_A \\ A \triangleright A & \xrightarrow{\mu_A} & A \end{array}$$

0270

2. *Left Unitality.* The diagram

$$\begin{array}{ccc} A & \xrightarrow{\lambda_A^{\text{Sets}_*, \triangleright}} & S^0 \triangleright A \\ \parallel & & \downarrow \eta_A \times \text{id}_A \\ A & \xleftarrow{\mu_A} & A \triangleright A \end{array}$$

commutes.

0271

## 3. Right Unitality. The diagram

$$\begin{array}{ccc} A \triangleright S^0 & \xrightarrow{\text{id}_A \times \eta_A} & A \triangleright A \\ & \searrow \rho_A^{\text{Sets}_*, \triangleright} & \downarrow \mu_A \\ & & A \end{array}$$

commutes.

Being a right-bilinear morphism of pointed sets, the multiplication map satisfies

$$0_A a = 0_A$$

for each  $a \in A$ . Now, the associativity, left unitality, and right unitality conditions act on elements as follows:

0272

## 1. Associativity. The associativity condition acts as

$$\begin{array}{ccccc} & & a \triangleright (b \triangleright c) & & \\ & \nearrow & & \swarrow & \\ (a \triangleright b) \triangleright c & & (a \triangleright b) \triangleright c & & a \triangleright bc \\ \searrow & & \nearrow & & \swarrow \\ ab \triangleright c & \longmapsto & (ab)c & & a(bc) \end{array}$$

This gives

$$(ab)c = a(bc)$$

for each  $a, b, c \in A$ .

0273

## 2. Left Unitality. The left unitality condition acts as

$$\begin{array}{ccc} a & & a \longmapsto 1 \triangleright a \\ \downarrow & & \downarrow \\ a & & 1_A a \longleftarrow 1_A \triangleright a \end{array}$$

This gives

$$1_A a = a$$

for each  $a \in A$ .

0274

3. *Right Unitality.* The right unitality condition acts:

0275

(a) On  $1 \triangleright 0$  as

$$\begin{array}{ccc} 1 \triangleright 0 & & a \triangleright 0 \mapsto a \triangleright 0_A \\ \swarrow & & \downarrow \\ 0_A & & a0_A. \end{array}$$

0276

(b) On  $a \triangleright 1$  as

$$\begin{array}{ccc} a \triangleright 1 & & a \triangleright 1 \mapsto a \triangleright 1_A \\ \swarrow & & \downarrow \\ a & & a1_A. \end{array}$$

This gives

$$\begin{aligned} a1_A &= a, \\ a0_A &= 0_A \end{aligned}$$

for each  $a \in A$ .

Thus we see that monoids with respect to  $\triangleright$  are exactly monoids with right zero.

#### Morphisms of Monoids on $(\text{Sets}_*, \triangleright, S^0)$

A morphism of monoids on  $(\text{Sets}_*, \triangleright, S^0)$  from  $(A, \mu_A, \eta_A, 0_A)$  to  $(B, \mu_B, \eta_B, 0_B)$  is a morphism of pointed sets

$$f: (A, 0_A) \rightarrow (B, 0_B)$$

satisfying the following conditions:

0277

1. *Compatibility With the Multiplication Morphisms.* The diagram

$$\begin{array}{ccc} A \triangleright A & \xrightarrow{f \triangleright f} & B \triangleright B \\ \mu_A \downarrow & & \downarrow \mu_B \\ A & \xrightarrow{f} & B \end{array}$$

commutes.

0278

2. *Compatibility With the Unit Morphisms.* The diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{\eta_A} & A \\ & \searrow \eta_B & \downarrow f \\ & & B \end{array}$$

commutes.

These act on elements as

$$\begin{array}{ccc} a \triangleright b & & a \triangleright b \mapsto f(a) \triangleright f(b) \\ \downarrow & & \downarrow \\ ab & \longmapsto & f(ab) \\ & & f(a)f(b) \end{array}$$

and

$$\begin{array}{ccc} 0 & \nearrow & 0 \mapsto 0_A \\ & \searrow & \downarrow \\ & 0_B & f(0_A) \end{array}$$

and

$$\begin{array}{ccc} 1 & \nearrow & 1 \mapsto 1_A \\ & \searrow & \downarrow \\ & 1_B & f(1_A) \end{array}$$

giving

$$f(ab) = f(a)f(b),$$

$$f(0_A) = 0_B,$$

$$f(1_A) = 1_B,$$

for each  $a, b \in A$ , which is exactly a morphism of monoids with right zero.

**Identities and Composition**

Similarly, the identities and composition of  $\text{Mon}(\text{Sets}_*, \triangleright, S^0)$  can be easily seen to agree with those of monoids with right zero, which finishes the proof. 

## 00FT 7.5 The Smash Product of Pointed Sets

### 00FU 7.5.1 Foundations

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00FV

#### DEFINITION 7.5.1.1.1 ► SMASH PRODUCTS OF POINTED SETS

The **smash product of**  $(X, x_0)$  **and**  $(Y, y_0)$ <sup>1</sup> is the pointed set  $X \wedge Y$ <sup>2</sup> satisfying the bijection

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z),$$

naturally in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ .

<sup>1</sup>*Further Terminology:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, the smash product  $X \wedge Y$  is also called the **tensor product of  $\mathbb{F}_1$ -modules of**  $(X, x_0)$  **and**  $(Y, y_0)$  or the **tensor product of**  $(X, x_0)$  **and**  $(Y, y_0)$  **over**  $\mathbb{F}_1$ .

<sup>2</sup>*Further Notation:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, the smash product  $X \wedge Y$  is also denoted  $X \otimes_{\mathbb{F}_1} Y$ .

00FW

#### REMARK 7.5.1.2 ► UNWINDING DEFINITION 7.5.1.1: THE UNIVERSAL PROPERTY I

That is to say, the smash product of pointed sets is defined so as to induce a bijection between the following data:

- Pointed maps  $f: X \wedge Y \rightarrow Z$ .
- Maps of sets  $f: X \times Y \rightarrow Z$  satisfying

$$f(x_0, y) = z_0,$$

$$f(x, y_0) = z_0$$

for each  $x \in X$  and each  $y \in Y$ .

00FX

#### REMARK 7.5.1.3 ► UNWINDING DEFINITION 7.5.1.1: THE UNIVERSAL PROPERTY II

The smash product of pointed sets may be described as follows:

- The smash product of  $(X, x_0)$  and  $(Y, y_0)$  is the pair  $((X \wedge Y, x_0 \wedge y_0), \iota)$  consisting of
  - A pointed set  $(X \wedge Y, x_0 \wedge y_0)$ ;
  - A bilinear morphism of pointed sets  $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \wedge Y$ ;

satisfying the following universal property:

- (★) Given another such pair  $((Z, z_0), f)$  consisting of

\* A pointed set  $(Z, z_0)$ ;

\* A bilinear morphism of pointed sets  $f: (X \times Y, (x_0, y_0)) \rightarrow X \wedge Y$ ;

there exists a unique morphism of pointed sets  $X \wedge Y \xrightarrow{\exists!} Z$  making the diagram

$$\begin{array}{ccc} & X \wedge Y & \\ l \nearrow & \downarrow & \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

00FY

#### CONSTRUCTION 7.5.1.1.4 ► SMASH PRODUCTS OF POINTED SETS

Concretely, the smash product of  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $(X \wedge Y, x_0 \wedge y_0)$  consisting of:

- *The Underlying Set.* The set  $X \wedge Y$  defined by

$$X \wedge Y \cong (X \times Y) / \sim_R,$$

where  $\sim_R$  is the equivalence relation on  $X \times Y$  obtained by declaring

$$\begin{aligned} (x_0, y) &\sim_R (x_0, y'), \\ (x, y_0) &\sim_R (x', y_0) \end{aligned}$$

for each  $x, x' \in X$  and each  $y, y' \in Y$ .

- *The Basepoint.* The element  $[(x_0, y_0)]$  of  $X \wedge Y$  given by the equivalence class of  $(x_0, y_0)$  under the equivalence relation  $\sim$  on  $X \times Y$ .

#### PROOF 7.5.1.1.5 ► PROOF OF CONSTRUCTION 7.5.1.1.4

By ??, we have a natural bijection

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z),$$

where  $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$  is the set

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid \begin{array}{l} \text{for each } x, y \in X, \text{ if} \\ (x, y) \sim_R (x', y'), \text{ then} \\ f(x, y) = f(x', y') \end{array} \right\}.$$

However, the condition  $(x, y) \sim_R (x', y')$  only holds when:

0279 1. We have  $x = x'$  and  $y = y'$ .

027A 2. The following conditions are satisfied:

027B (a) We have  $x = x_0$  or  $y = y_0$ .

027C (b) We have  $x' = x_0$  or  $y' = y_0$ .

So, given  $f \in \text{Hom}_{\text{Sets}}(X \times Y, Z)$  with a corresponding  $\bar{f}: X \wedge Y \rightarrow Z$ , the latter case above implies

$$\begin{aligned} f(x_0, y) &= f(x, y_0) \\ &= f(x_0, y_0), \end{aligned}$$

and since  $\bar{f}: X \wedge Y \rightarrow Z$  is a pointed map, we have

$$\begin{aligned} f(x_0, y_0) &= \bar{f}(x_0, y_0) \\ &= z_0. \end{aligned}$$

Thus the elements  $f$  in  $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$  are precisely those functions  $f: X \times Y \rightarrow Z$  satisfying the equalities

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ , giving an equality

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) = \text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$$

of sets, which when composed with our earlier isomorphism

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z),$$

gives our desired natural bijection, finishing the proof. □

00FZ

**REMARK 7.5.1.1.6 ► ON THE CONSTRUCTION OF THE SMASH PRODUCT OF POINTED SETS**

It is also somewhat common to write

$$X \wedge Y \stackrel{\text{def}}{=} \frac{X \times Y}{X \vee Y},$$

identifying  $X \vee Y$  with the subspace  $(\{x_0\} \times Y) \cup (X \times \{y_0\})$  of  $X \times Y$ , and having the quotient be defined by declaring  $(x, y) \sim (x', y')$  iff we have  $(x, y), (x', y') \in X \vee Y$ .

01VX

**CONSTRUCTION 7.5.1.1.7 ► A SECOND CONSTRUCTION OF THE SMASH PRODUCT OF POINTED SETS**

Alternatively, the smash product of  $(X, x_0)$  and  $(Y, y_0)$  may be constructed as the pointed set  $X \wedge Y$  given by

$$\begin{aligned} X \wedge Y &\cong \bigvee_{x \in X^-} Y \\ &\cong \bigvee_{y \in Y^-} X. \end{aligned}$$

**PROOF 7.5.1.1.8 ► PROOF OF CONSTRUCTION 7.5.1.1.7**

Indeed, since  $X \cong \bigvee_{x \in X^-} S^0$ , we have

$$\begin{aligned} X \wedge Y &\cong (\bigvee_{x \in X^-} S^0) \wedge Y \\ &\cong \bigvee_{x \in X^-} S^0 \wedge Y \\ &\cong \bigvee_{x \in X^-} Y, \end{aligned}$$

where we have used that  $\wedge$  preserves colimits in both variables via ?? for the second isomorphism above, since it has right adjoints in both variables by [Item 2](#).

A similar proof applies to the isomorphism  $X \wedge Y \cong \bigvee_{y \in Y^-} X$ . 

00G0

**NOTATION 7.5.1.1.9 ► ELEMENTS OF SMASH PRODUCTS OF POINTED SETS**

We write  $x \wedge y$  for the element  $[(x, y)]$  of  $X \wedge Y \cong X \times Y / \sim$ .

00G1

**REMARK 7.5.1.1.10 ► BASEPOINTS OF SMASH PRODUCTS OF POINTED SETS**

Employing the notation introduced in [Notation 7.5.1.1.9](#), we have

$$\begin{aligned} x_0 \wedge y_0 &= x \wedge y_0, \\ &= x_0 \wedge y \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ , and

$$\begin{aligned} x \wedge y_0 &= x' \wedge y_0, \\ x_0 \wedge y &= x_0 \wedge y' \end{aligned}$$

for each  $x, x' \in X$  and each  $y, y' \in Y$ .

00G2

**EXAMPLE 7.5.1.1.11 ► EXAMPLES OF SMASH PRODUCTS OF POINTED SETS**

Here are some examples of smash products of pointed sets.

00G3

- Smashing With pt.* For any pointed set  $X$ , we have isomorphisms of pointed sets

$$\begin{aligned} \text{pt} \wedge X &\cong \text{pt}, \\ X \wedge \text{pt} &\cong \text{pt}. \end{aligned}$$

00G4

- Smashing With  $S^0$ .* For any pointed set  $X$ , we have isomorphisms of pointed sets

$$\begin{aligned} S^0 \wedge X &\cong X, \\ X \wedge S^0 &\cong X. \end{aligned}$$

00G5

**PROPOSITION 7.5.1.1.12 ► PROPERTIES OF SMASH PRODUCTS OF POINTED SETS**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00G6

- Functionality.* The assignments  $X, Y, (X, Y) \mapsto X \wedge Y$  define functors

$$\begin{aligned} X \wedge -: \quad \text{Sets}_* &\rightarrow \text{Sets}_*, \\ - \wedge Y: \quad \text{Sets}_* &\rightarrow \text{Sets}_*, \\ -_1 \wedge -_2: \text{Sets}_* \times \text{Sets}_* &\rightarrow \text{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f: (X, x_0) &\rightarrow (A, a_0), \\ g: (Y, y_0) &\rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \wedge g: X \wedge Y \rightarrow A \wedge B$$

is given by

$$[f \wedge g](x \wedge y) \stackrel{\text{def}}{=} f(x) \wedge g(y)$$

for each  $x \wedge y \in X \wedge Y$ .

**00G7** 2. *Adjointness.* We have adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*,$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*,$$

witnessed by bijections

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(A, Z)),$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ .

**00G8** 3. *Enriched Adjointness.* We have  $\mathbf{Sets}_*$ -enriched adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*,$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*,$$

witnessed by isomorphisms of pointed sets

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)),$$

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)),$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ .

**00G9** 4. *As a Pushout.* We have an isomorphism

$$X \wedge Y \cong \mathrm{pt} \coprod_{X \vee Y} (X \times Y),$$

$$\begin{array}{ccc} X \wedge Y & \xhookleftarrow{\quad} & X \times Y \\ \uparrow \lrcorner & & \uparrow \wr \\ \mathrm{pt} & \xleftarrow{!} & X \vee Y, \end{array}$$

natural in  $X, Y \in \text{Obj}(\text{Sets}_*)$ , where the pushout is taken in  $\text{Sets}$ , and the embedding  $\iota: X \vee Y \hookrightarrow X \times Y$  is defined following [Remark 7.5.1.1.6](#).

00GA

5. *Distributivity Over Wedge Sums.* We have isomorphisms of pointed sets

$$\begin{aligned} X \wedge (Y \vee Z) &\cong (X \wedge Y) \vee (X \wedge Z), \\ (X \vee Y) \wedge Z &\cong (X \wedge Z) \vee (Y \wedge Z), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ .

**PROOF 7.5.1.1.13 ▶ PROOF OF PROPOSITION 7.5.1.1.12**

**Item 1: Functoriality**

The map  $f \wedge g$  comes from [Item 4 of Proposition 10.6.2.1.3](#) via the map

$$f \wedge g: X \times Y \rightarrow A \wedge B$$

sending  $(x, y)$  to  $f(x) \wedge g(y)$ , which we need to show satisfies

$$[f \wedge g](x, y) = [f \wedge g](x', y')$$

for each  $(x, y), (x', y') \in X \times Y$  with  $(x, y) \sim_R (x', y')$ , where  $\sim_R$  is the relation constructing  $X \wedge Y$  as

$$X \wedge Y \cong (X \times Y)/\sim_R$$

in [Construction 7.5.1.1.4](#). The condition defining  $\sim$  is that at least one of the following conditions is satisfied:

027D

1. We have  $x = x'$  and  $y = y'$ ;

027E

2. Both of the following conditions are satisfied:

027F

- (a) We have  $x = x_0$  or  $y = y_0$ .

027G

- (b) We have  $x' = x_0$  or  $y' = y_0$ .

We have five cases:

027H

1. In the first case, we clearly have

$$[f \wedge g](x, y) = [f \wedge g](x', y')$$

since  $x = x'$  and  $y = y'$ .

027J

2. If  $x = x_0$  and  $x' = x_0$ , we have

$$\begin{aligned}[f \wedge g](x_0, y) &\stackrel{\text{def}}{=} f(x_0) \wedge g(y) \\ &= a_0 \wedge g(y) \\ &= a_0 \wedge g(y') \\ &= f(x_0) \wedge g(y') \\ &\stackrel{\text{def}}{=} [f \wedge g](x_0, y').\end{aligned}$$

027K

3. If  $x = x_0$  and  $y' = y_0$ , we have

$$\begin{aligned}[f \wedge g](x_0, y) &\stackrel{\text{def}}{=} f(x_0) \wedge g(y) \\ &= a_0 \wedge g(y) \\ &= a_0 \wedge b_0 \\ &= f(x') \wedge b_0 \\ &= f(x') \wedge g(y_0) \\ &\stackrel{\text{def}}{=} [f \wedge g](x', y_0).\end{aligned}$$

027L

4. If  $y = y_0$  and  $x' = x_0$ , we have

$$\begin{aligned}[f \wedge g](x, y_0) &\stackrel{\text{def}}{=} f(x) \wedge g(y_0) \\ &= f(x) \wedge b_0 \\ &= a_0 \wedge b_0 \\ &= a_0 \wedge g(y') \\ &= f(x_0) \wedge g(y') \\ &\stackrel{\text{def}}{=} [f \wedge g](x_0, y').\end{aligned}$$

027M

5. If  $y = y_0$  and  $y' = y_0$ , we have

$$\begin{aligned}[f \wedge g](x, y_0) &\stackrel{\text{def}}{=} f(x) \wedge g(y_0) \\ &= f(x) \wedge b_0 \\ &= f(x') \wedge b_0 \\ &= f(x) \wedge g(y_0) \\ &\stackrel{\text{def}}{=} [f \wedge g](x', y_0).\end{aligned}$$

Thus  $f \wedge g$  is well-defined. Next, we claim that  $\wedge$  preserves identities and composition:

- *Preservation of Identities.* We have

$$\begin{aligned} [\text{id}_X \wedge \text{id}_Y](x \wedge y) &\stackrel{\text{def}}{=} \text{id}_X(x) \wedge \text{id}_Y(y) \\ &= x \wedge y \\ &= [\text{id}_{X \wedge Y}](x \wedge y) \end{aligned}$$

for each  $x \wedge y \in X \wedge Y$ , and thus

$$\text{id}_X \wedge \text{id}_Y = \text{id}_{X \wedge Y}.$$

- *Preservation of Composition.* Given pointed maps

$$\begin{aligned} f: (X, x_0) &\rightarrow (X', x'_0), \\ h: (X', x'_0) &\rightarrow (X'', x''_0), \\ g: (Y, y_0) &\rightarrow (Y', y'_0), \\ k: (Y', y'_0) &\rightarrow (Y'', y''_0), \end{aligned}$$

we have

$$\begin{aligned} [(h \circ f) \wedge (k \circ g)](x \wedge y) &\stackrel{\text{def}}{=} h(f(x)) \wedge k(g(y)) \\ &\stackrel{\text{def}}{=} [h \wedge k](f(x) \wedge g(y)) \\ &\stackrel{\text{def}}{=} [h \wedge k]([f \wedge g](x \wedge y)) \\ &\stackrel{\text{def}}{=} [(h \wedge k) \circ (f \wedge g)](x \wedge y) \end{aligned}$$

for each  $x \wedge y \in X \wedge Y$ , and thus

$$(h \circ f) \wedge (k \circ g) = (h \wedge k) \circ (f \wedge g).$$

This finishes the proof.

### Item 2: Adjointness

We prove only the adjunction  $- \wedge Y \dashv \mathbf{Sets}_*(Y, -)$ , witnessed by a natural bijection

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

as the proof of the adjunction  $X \wedge - \dashv \mathbf{Sets}_*(X, -)$  is similar. We claim we have a bijection

$$\text{Hom}_{\mathbf{Sets}_*}^\otimes(X \times Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , implying the desired adjunction. Indeed, this bijection is a restriction of the bijection

$$\text{Sets}(X \times Y, Z) \cong \text{Sets}(X, \text{Sets}(Y, Z))$$

of Item 2 of Proposition 4.1.3.1.4:

- A map

$$\xi: X \times Y \rightarrow Z$$

in  $\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$  gets sent to the pointed map

$$\begin{aligned}\xi^\dagger: (X, x_0) &\rightarrow (\text{Sets}_*(Y, Z), \Delta_{z_0}), \\ x &\longmapsto (\xi_x^\dagger: Y \rightarrow Z),\end{aligned}$$

where  $\xi_x^\dagger: Y \rightarrow Z$  is the map defined by

$$\xi_x^\dagger(y) \stackrel{\text{def}}{=} \xi(x, y)$$

for each  $y \in Y$ , where:

- The map  $\xi^\dagger$  is indeed pointed, as we have

$$\begin{aligned}\xi_{x_0}^\dagger(y) &\stackrel{\text{def}}{=} \xi(x_0, y) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each  $y \in Y$ . Thus  $\xi_{x_0}^\dagger = \Delta_{z_0}$  and  $\xi^\dagger$  is pointed.

- The map  $\xi_x^\dagger$  indeed lies in  $\text{Sets}_*(Y, Z)$ , as we have

$$\begin{aligned}\xi_x^\dagger(y_0) &\stackrel{\text{def}}{=} \xi(x, y_0) \\ &\stackrel{\text{def}}{=} z_0.\end{aligned}$$

- Conversely, a map

$$\xi: (X, x_0) \rightarrow (\text{Sets}_*(Y, Z), \Delta_{z_0}),$$

$$x \longmapsto (\xi_x: Y \rightarrow Z),$$

in  $\text{Hom}_{\text{Sets}_*}(X, \text{Sets}_*(Y, Z))$  gets sent to the map

$$\xi^\dagger: X \times Y \rightarrow Z$$

defined by

$$\xi^\dagger(x, y) \stackrel{\text{def}}{=} \xi_x(y)$$

for each  $(x, y) \in X \times Y$ , which indeed lies in  $\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$ , as:

– *Left Bilinearity.* We have

$$\begin{aligned}\xi^\dagger(x_0, y) &\stackrel{\text{def}}{=} \xi_{x_0}(y) \\ &\stackrel{\text{def}}{=} \Delta_{z_0}(y) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each  $y \in Y$ , since  $\xi_{x_0} = \Delta_{z_0}$  as  $\xi$  is assumed to be a pointed map.

– *Right Bilinearity.* We have

$$\begin{aligned}\xi^\dagger(x, y_0) &\stackrel{\text{def}}{=} \xi_x(y_0) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each  $x \in X$ , since  $\xi_x \in \mathbf{Sets}_*(Y, Z)$  is a morphism of pointed sets.

This finishes the proof.

**Item 3: Enriched Adjointness**

This follows from **Item 2** and ?? of ??.

**Item 4: As a Pushout**

Following the description of **Remark 4.2.4.1.4**, we have

$$\text{pt} \coprod_{X \vee Y} (X \times Y) \cong (\text{pt} \times (X \times Y)) / \sim,$$

where  $\sim$  identifies the element  $\star$  in  $\text{pt}$  with all elements of the form  $(x_0, y)$  and  $(x, y_0)$  in  $X \times Y$ . Thus **Item 4** of **Proposition 10.6.2.1.3** coupled with **Remark 7.5.1.1.10** then gives us a well-defined map

$$\text{pt} \coprod_{X \vee Y} (X \times Y) \rightarrow X \wedge Y$$

via  $[(\star, (x, y))] \mapsto x \wedge y$ , with inverse

$$X \wedge Y \rightarrow \text{pt} \coprod_{X \vee Y} (X \times Y)$$

given by  $x \wedge y \mapsto [(\star, (x, y))]$ .

**Item 5: Distributivity Over Wedge Sums**

This follows from **Proposition 7.5.9.1.1**, ?? of ??, and the fact that  $\vee$  is the coproduct in  $\mathbf{Sets}_*$  (**Definition 6.3.3.1.1**). 

## 00GB 7.5.2 The Internal Hom of Pointed Sets

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00GC

## DEFINITION 7.5.2.1.1 ► THE INTERNAL HOM OF POINTED SETS

The **internal Hom**<sup>1</sup> of pointed sets from  $(X, x_0)$  to  $(Y, y_0)$  is the pointed set  $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ <sup>2</sup> consisting of:

- *The Underlying Set.* The set  $\mathbf{Sets}_*((X, x_0), (Y, y_0))$  of morphisms of pointed sets from  $(X, x_0)$  to  $(Y, y_0)$ .
- *The Basepoint.* The element

$$\Delta_{y_0}: (X, x_0) \rightarrow (Y, y_0)$$

of  $\mathbf{Sets}_*((X, x_0), (Y, y_0))$  given by

$$\Delta_{y_0}(x) \stackrel{\text{def}}{=} y_0$$

for each  $x \in X$ .

<sup>1</sup>For a proof that  $\mathbf{Sets}_*$  is indeed the internal Hom of  $\mathbf{Sets}_*$  with respect to the smash product of pointed sets, see Item 2 of Proposition 7.5.1.12.

<sup>2</sup>Further Notation: Also written  $\mathbf{Hom}_{\mathbf{Sets}_*}(X, Y)$ .

00GD

## PROPOSITION 7.5.2.1.2 ► PROPERTIES OF THE INTERNAL HOM OF POINTED SETS

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00GE

1. *Functionality.* The assignments  $X, Y, (X, Y) \mapsto \mathbf{Sets}_*(X, Y)$  define functors

$$\begin{aligned} \mathbf{Sets}_*(X, -): \quad & \mathbf{Sets}_* & \rightarrow \mathbf{Sets}_*, \\ \mathbf{Sets}_*(-, Y): \quad & \mathbf{Sets}_*^{\text{op}} & \rightarrow \mathbf{Sets}_*, \\ \mathbf{Sets}_*(-, -): \quad & \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* & \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f: (X, x_0) & \rightarrow (A, a_0), \\ g: (Y, y_0) & \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$\mathbf{Sets}_*(f, g): \mathbf{Sets}_*(A, Y) \rightarrow \mathbf{Sets}_*(X, B)$$

is given by

$$[\mathbf{Sets}_*(f, g)](\phi) \stackrel{\text{def}}{=} g \circ \phi \circ f$$

for each  $\phi \in \mathbf{Sets}_*(A, Y)$ .

00GF

2. *Adjointness.* We have adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)): \quad \begin{array}{c} \mathbf{Sets}_* \xrightarrow{\quad X \wedge - \quad} \mathbf{Sets}_*, \\ \perp \\ \mathbf{Sets}_*(X, -) \end{array}$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)): \quad \begin{array}{c} \mathbf{Sets}_* \xrightarrow{\quad - \wedge Y \quad} \mathbf{Sets}_*, \\ \perp \\ \mathbf{Sets}_*(Y, -) \end{array}$$

witnessed by bijections

$$\begin{aligned} \text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) &\cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)), \\ \text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) &\cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(A, Z)), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ .

00GG

3. *Enriched Adjointness.* We have  $\mathbf{Sets}_*$ -enriched adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)): \quad \begin{array}{c} \mathbf{Sets}_* \xrightarrow{\quad X \wedge - \quad} \mathbf{Sets}_*, \\ \perp \\ \mathbf{Sets}_*(X, -) \end{array}$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)): \quad \begin{array}{c} \mathbf{Sets}_* \xrightarrow{\quad - \wedge Y \quad} \mathbf{Sets}_*, \\ \perp \\ \mathbf{Sets}_*(Y, -) \end{array}$$

witnessed by isomorphisms of pointed sets

$$\begin{aligned} \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)), \\ \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ .

#### PROOF 7.5.2.1.3 ▶ PROOF OF PROPOSITION 7.5.2.1.2

**Item 1:** Functoriality

This follows from **Item 1** of [Proposition 4.3.5.1.2](#) and from the equalities

$$\begin{aligned} g \circ \Delta_{y_0} &= \Delta_{z_0}, \\ \Delta_{y_0} \circ f &= \Delta_{y_0} \end{aligned}$$

for morphisms  $f: (K, k_0) \rightarrow (X, x_0)$  and  $g: (Y, y_0) \rightarrow (Z, z_0)$ , which guarantee pre- and postcomposition by morphisms of pointed sets to

also be morphisms of pointed sets.

## Item 2: Adjointness

This is a repetition of Item 2 of Proposition 7.5.1.1.12, and is proved there.

### Item 3: Enriched Adjointness

This is a repetition of Item 3 of Proposition 7.5.1.1.12, and is proved there. 

00GH 7.5.3 The Monoidal Unit

00GJ

### DEFINITION 7.5.3.1.1 ► THE MONOIDAL UNIT OF $\wedge$

The **monoidal unit of the smash product of pointed sets** is the functor

$$1^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*} \stackrel{\text{def}}{=} S^0.$$

00GK 7.5.4 The Associator


00GL

### DEFINITION 7.5.4.1.1 ► THE ASSOCIATOR OF $\wedge$

The **associator** of the smash product of pointed sets is the natural isomorphism

$$\alpha^{\text{Sets}_*} : \wedge \circ (\wedge \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \wedge \circ (\text{id}_{\text{Sets}_*} \times \wedge) \circ \alpha^{\text{Cats}}_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*},$$

as in the diagram



whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*}: (X \wedge Y) \wedge Z \xrightarrow{\sim} X \wedge (Y \wedge Z)$$

at  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\alpha_{X,Y,Z}^{\text{Sets}_*}((x \wedge y) \wedge z) \stackrel{\text{def}}{=} x \wedge (y \wedge z)$$

for each  $(x \wedge y) \wedge z \in (X \wedge Y) \wedge Z$ .

#### PROOF 7.5.4.1.2 ► PROOF OF DEFINITION 7.5.4.1.1

##### Well-Definedness

Let  $[(x, y), z] = [(x', y'), z']$  be an element in  $(X \wedge Y) \wedge Z$ . Then either:

027N 1. We have  $x = x'$ ,  $y = y'$ , and  $z = z'$ .

027P 2. Both of the following conditions are satisfied:

027Q (a) We have  $x = x_0$  or  $y = y_0$  or  $z = z_0$ .

027R (b) We have  $x' = x_0$  or  $y' = y_0$  or  $z' = z_0$ .

In the first case,  $\alpha_{X,Y,Z}^{\text{Sets}_*}$  clearly sends both elements to the same element in  $X \wedge (Y \wedge Z)$ . Meanwhile, in the latter case both elements are equal to the basepoint  $(x_0 \wedge y_0) \wedge z_0$  of  $(X \wedge Y) \wedge Z$ , which gets sent to the basepoint  $x_0 \wedge (y_0 \wedge z_0)$  of  $X \wedge (Y \wedge Z)$ .

##### Being a Morphism of Pointed Sets

As just mentioned, we have

$$\alpha_{X,Y,Z}^{\text{Sets}_*}((x_0 \wedge y_0) \wedge z_0) \stackrel{\text{def}}{=} x_0 \wedge (y_0 \wedge z_0),$$

and thus  $\alpha_{X,Y,Z}^{\text{Sets}_*}$  is a morphism of pointed sets.

##### Invertibility

The inverse of  $\alpha_{X,Y,Z}^{\text{Sets}_*}$  is given by the morphism

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1}: X \wedge (Y \wedge Z) \xrightarrow{\sim} (X \wedge Y) \wedge Z$$

defined by

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1}(x \wedge (y \wedge z)) \stackrel{\text{def}}{=} (x \wedge y) \wedge z$$

for each  $x \wedge (y \wedge z) \in X \wedge (Y \wedge Z)$ .

##### Naturality

We need to show that, given morphisms of pointed sets

$$f: (X, x_0) \rightarrow (X', x'_0),$$

$$\begin{aligned} g: (Y, y_0) &\rightarrow (Y', y'_0), \\ h: (Z, z_0) &\rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \wedge Y) \wedge Z & \xrightarrow{(f \wedge g) \wedge h} & (X' \wedge Y') \wedge Z' \\ \alpha_{X,Y,Z}^{\text{Sets}_*} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*} \\ X \wedge (Y \wedge Z) & \xrightarrow{f \wedge (g \wedge h)} & X' \wedge (Y' \wedge Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \wedge y) \wedge z & \longmapsto & (f(x) \wedge g(y)) \wedge h(z) \\ \downarrow & & \downarrow \\ x \wedge (y \wedge z) & \longmapsto & f(x) \wedge (g(y) \wedge h(z)) \end{array}$$

and hence indeed commutes, showing  $\alpha^{\text{Sets}_*}$  to be a natural transformation.

#### Being a Natural Isomorphism

Since  $\alpha^{\text{Sets}_*}$  is natural and  $\alpha^{\text{Sets}_*, -1}$  is a componentwise inverse to  $\alpha^{\text{Sets}_*}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\alpha^{\text{Sets}_*, -1}$  is also natural. Thus  $\alpha^{\text{Sets}_*}$  is a natural isomorphism. ■

### 00GM 7.5.5 The Left Unitor

00GN

#### DEFINITION 7.5.5.1.1 ► THE LEFT UNITOR OF $\wedge$

The **left unitor of the smash product of pointed sets** is the natural isomorphism

$$\begin{array}{ccc} \text{pt} \times \text{Sets}_* & \xrightarrow{\mathbb{1}_{\text{Sets}_*} \times \text{id}} & \text{Sets}_* \times \text{Sets}_* \\ \lambda^{\text{Sets}_*}: \wedge \circ (\mathbb{1}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \lambda_{\text{Sets}_*}^{\text{Cats}_2} & \swarrow \quad \searrow & \downarrow \wedge \\ & \lambda_{\text{Sets}_*}^{\text{Cats}_2} & \text{Sets}_*, \end{array}$$

whose component

$$\lambda_X^{\text{Sets}_*} : S^0 \wedge X \xrightarrow{\sim} X$$

at  $X \in \text{Obj}(\text{Sets}_*)$  is given by

$$\begin{aligned} 0 \wedge x &\mapsto x_0, \\ 1 \wedge x &\mapsto x \end{aligned}$$

for each  $x \in X$ .

#### PROOF 7.5.5.1.2 ▶ PROOF OF DEFINITION 7.5.5.1.1

##### Well-Definedness

Let  $[(x, y)] = [(x', y')]$  be an element in  $S^0 \wedge X$ . Then either:

**027S** 1. We have  $x = x'$  and  $y = y'$ .

**027T** 2. Both of the following conditions are satisfied:

**027U** (a) We have  $x = 0$  or  $y = x_0$ .

**027V** (b) We have  $x' = 0$  or  $y' = x_0$ .

In the first case,  $\lambda_X^{\text{Sets}_*}$  clearly sends both elements to the same element in  $X$ . Meanwhile, in the latter case both elements are equal to the basepoint  $0 \wedge x_0$  of  $S^0 \wedge X$ , which gets sent to the basepoint  $x_0$  of  $X$ .

##### Being a Morphism of Pointed Sets

As just mentioned, we have

$$\lambda_X^{\text{Sets}_*}(0 \wedge x_0) \stackrel{\text{def}}{=} x_0,$$

and thus  $\lambda_X^{\text{Sets}_*}$  is a morphism of pointed sets.

##### Invertibility

The inverse of  $\lambda_X^{\text{Sets}_*}$  is the morphism

$$\lambda_X^{\text{Sets}_*, -1} : X \xrightarrow{\sim} S^0 \wedge X$$

defined by

$$\lambda_X^{\text{Sets}_*, -1}(x) \stackrel{\text{def}}{=} 1 \wedge x$$

for each  $x \in X$ . Indeed:

027W

1. *Invertibility I.* We have

$$\begin{aligned} [\lambda_X^{\text{Sets}_*, -1} \circ \lambda_X^{\text{Sets}_*}](0 \wedge x) &= \lambda_X^{\text{Sets}_*, -1}(\lambda_X^{\text{Sets}_*}(0 \wedge x)) \\ &= \lambda_X^{\text{Sets}_*, -1}(x_0) \\ &= 1 \wedge x_0 \\ &= 0 \wedge x, \end{aligned}$$

and

$$\begin{aligned} [\lambda_X^{\text{Sets}_*, -1} \circ \lambda_X^{\text{Sets}_*}](1 \wedge x) &= \lambda_X^{\text{Sets}_*, -1}(\lambda_X^{\text{Sets}_*}(1 \wedge x)) \\ &= \lambda_X^{\text{Sets}_*, -1}(x) \\ &= 1 \wedge x \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\lambda_X^{\text{Sets}_*, -1} \circ \lambda_X^{\text{Sets}_*} = \text{id}_{S^0 \wedge X}.$$

027X

2. *Invertibility II.* We have

$$\begin{aligned} [\lambda_X^{\text{Sets}_*} \circ \lambda_X^{\text{Sets}_*, -1}](x) &= \lambda_X^{\text{Sets}_*}(\lambda_X^{\text{Sets}_*, -1}(x)) \\ &= \lambda_X^{\text{Sets}_*, -1}(1 \wedge x) \\ &= x \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\lambda_X^{\text{Sets}_*} \circ \lambda_X^{\text{Sets}_*, -1} = \text{id}_X.$$

This shows  $\lambda_X^{\text{Sets}_*}$  to be invertible.

### Naturality

We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} S^0 \wedge X & \xrightarrow{\text{id}_{S^0 \wedge f}} & S^0 \wedge Y \\ \lambda_X^{\text{Sets}_*} \downarrow & & \downarrow \lambda_Y^{\text{Sets}_*} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 \wedge x & & 0 \wedge x \xrightarrow{\quad} 0 \wedge f(x) \\ \downarrow & & \downarrow \\ x_0 \xrightarrow{\quad} f(x_0) & & y_0 \end{array}$$

and

$$\begin{array}{ccc} 1 \wedge x \xrightarrow{\quad} 1 \wedge f(x) & & \\ \downarrow & & \downarrow \\ x \xrightarrow{\quad} f(x) & & \end{array}$$

and hence indeed commutes, showing  $\lambda^{\text{Sets}_*}$  to be a natural transformation.

#### Being a Natural Isomorphism

Since  $\lambda^{\text{Sets}_*}$  is natural and  $\lambda^{\text{Sets}_*, -1}$  is a componentwise inverse to  $\lambda^{\text{Sets}_*}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\lambda^{\text{Sets}_*, -1}$  is also natural. Thus  $\lambda^{\text{Sets}_*}$  is a natural isomorphism. □

### 00GP 7.5.6 The Right Unitor

00GQ

#### DEFINITION 7.5.6.1.1 ▶ THE RIGHT UNITOR OF $\wedge$

The **right unitor of the smash product of pointed sets** is the natural isomorphism

$$\begin{array}{ccc} \text{Sets}_* \times \text{pt} & \xrightarrow{\quad \text{id} \times 1^{\text{Sets}_*} \quad} & \text{Sets}_* \times \text{Sets}_* \\ \rho^{\text{Sets}_*} : \wedge \circ (\text{id} \times 1^{\text{Sets}_*}) \xrightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}, & \swarrow \text{dashed} & \searrow \text{double} \\ & \rho_{\text{Sets}_*}^{\text{Cats}_2} & \end{array}$$

$\wedge$

whose component

$$\rho_X^{\text{Sets}_*} : X \wedge S^0 \xrightarrow{\sim} X$$

at  $X \in \text{Obj}(\text{Sets}_*)$  is given by

$$x \wedge 0 \mapsto x_0,$$

$$x \wedge 1 \mapsto x$$

for each  $x \in X$ .

#### PROOF 7.5.6.1.2 ► PROOF OF DEFINITION 7.5.6.1.1

##### Well-Definedness

Let  $[(x, y)] = [(x', y')]$  be an element in  $X \wedge S^0$ . Then either:

027Y

1. We have  $x = x'$  and  $y = y'$ .

027Z

2. Both of the following conditions are satisfied:

0280

(a) We have  $x = x_0$  or  $y = 0$ .

0281

(b) We have  $x' = x_0$  or  $y' = 0$ .

In the first case,  $\rho_X^{\text{Sets}_*}$  clearly sends both elements to the same element in  $X$ . Meanwhile, in the latter case both elements are equal to the basepoint  $x_0 \wedge 0$  of  $X \wedge S^0$ , which gets sent to the basepoint  $x_0$  of  $X$ .

##### Being a Morphism of Pointed Sets

As just mentioned, we have

$$\rho_X^{\text{Sets}_*}(x_0 \wedge 0) \stackrel{\text{def}}{=} x_0,$$

and thus  $\rho_X^{\text{Sets}_*}$  is a morphism of pointed sets.

##### Invertibility

The inverse of  $\rho_X^{\text{Sets}_*}$  is the morphism

$$\rho_X^{\text{Sets}_*, -1} : X \xrightarrow{\sim} X \wedge S^0$$

defined by

$$\rho_X^{\text{Sets}_*, -1}(x) \stackrel{\text{def}}{=} x \wedge 1$$

for each  $x \in X$ . Indeed:

0282

1. *Invertibility I.* We have

$$\begin{aligned} [\rho_X^{\text{Sets}_*, -1} \circ \rho_X^{\text{Sets}_*}](x \wedge 0) &= \rho_X^{\text{Sets}_*, -1}(\rho_X^{\text{Sets}_*}(x \wedge 0)) \\ &= \rho_X^{\text{Sets}_*, -1}(x_0) \\ &= x_0 \wedge 1 \\ &= x \wedge 0, \end{aligned}$$

and

$$\begin{aligned} [\rho_X^{\text{Sets}_*, -1} \circ \rho_X^{\text{Sets}_*}](x \wedge 1) &= \rho_X^{\text{Sets}_*, -1}(\rho_X^{\text{Sets}_*}(x \wedge 1)) \\ &= \rho_X^{\text{Sets}_*, -1}(x) \\ &= x \wedge 1 \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\rho_X^{\text{Sets}_*, -1} \circ \rho_X^{\text{Sets}_*} = \text{id}_{X \wedge S^0}.$$

0283

2. *Invertibility II.* We have

$$\begin{aligned} [\rho_X^{\text{Sets}_*} \circ \rho_X^{\text{Sets}_*, -1}](x) &= \rho_X^{\text{Sets}_*}(\rho_X^{\text{Sets}_*, -1}(x)) \\ &= \rho_X^{\text{Sets}_*, -1}(x \wedge 1) \\ &= x \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\rho_X^{\text{Sets}_*} \circ \rho_X^{\text{Sets}_*, -1} = \text{id}_X.$$

This shows  $\rho_X^{\text{Sets}_*}$  to be invertible.

### Naturality

We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X \wedge S^0 & \xrightarrow{f \wedge \text{id}_{S^0}} & Y \wedge S^0 \\ \rho_X^{\text{Sets}_*} \downarrow & & \downarrow \rho_Y^{\text{Sets}_*} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge 0 & \longmapsto & x \wedge 0 \longmapsto f(x) \wedge 0 \\ \downarrow & & \downarrow \\ x_0 \longmapsto f(x_0) & & y_0 \end{array}$$

and

$$\begin{array}{ccc} x \wedge 1 & \longmapsto & f(x) \wedge 1 \\ \downarrow & & \downarrow \\ x & \longmapsto & f(x) \end{array}$$

and hence indeed commutes, showing  $\rho^{\text{Sets}_*}$  to be a natural transformation.

#### Being a Natural Isomorphism

Since  $\rho^{\text{Sets}_*}$  is natural and  $\rho^{\text{Sets}_*, -1}$  is a componentwise inverse to  $\rho^{\text{Sets}_*}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\rho^{\text{Sets}_*, -1}$  is also natural. Thus  $\rho^{\text{Sets}_*}$  is a natural isomorphism. 

### 00GR 7.5.7 The Symmetry

00GS

#### DEFINITION 7.5.7.1.1 ► THE SYMMETRY OF $\wedge$

The **symmetry of the smash product of pointed sets** is the natural isomorphism

$$\sigma^{\text{Sets}_*} : \wedge \xrightarrow{\sim} \wedge \circ \sigma_{\text{Sets}_*, \text{Sets}_*}^{\text{Cats}_2}, \quad \begin{array}{ccc} \text{Sets}_* \times \text{Sets}_* & \xrightarrow{\wedge} & \text{Sets}_*, \\ \sigma_{\text{Sets}_*, \text{Sets}_*}^{\text{Cats}_2} \swarrow & \Downarrow \sigma^{\text{Sets}_*} & \searrow \wedge \\ \text{Sets}_* \times \text{Sets}_* & & \end{array}$$

whose component

$$\sigma_{X,Y}^{\text{Sets}_*} : X \wedge Y \xrightarrow{\sim} Y \wedge X$$

at  $X, Y \in \text{Obj}(\text{Sets}_*)$  is defined by

$$\sigma_{X,Y}^{\text{Sets}_*}(x \wedge y) \stackrel{\text{def}}{=} y \wedge x$$

for each  $x \wedge y \in X \wedge Y$ .

#### PROOF 7.5.7.1.2 ► PROOF OF DEFINITION 7.5.7.1.1

##### Well-Definedness

Let  $[(x, y)] = [(x', y')]$  be an element in  $X \wedge Y$ . Then either:

0284

1. We have  $x = x'$  and  $y = y'$ .

0285

2. Both of the following conditions are satisfied:

0286

(a) We have  $x = x_0$  or  $y = y_0$ .

0287

(b) We have  $x' = x_0$  or  $y' = y_0$ .

In the first case,  $\sigma_X^{\text{Sets}_*}$  clearly sends both elements to the same element in  $X$ . Meanwhile, in the latter case both elements are equal to the basepoint  $x_0 \wedge y_0$  of  $X \wedge Y$ , which gets sent to the basepoint  $y_0 \wedge x_0$  of  $Y \wedge X$ .

### Being a Morphism of Pointed Sets

As just mentioned, we have

$$\sigma_X^{\text{Sets}_*}(x_0 \wedge y_0) \stackrel{\text{def}}{=} y_0 \wedge x_0,$$

and thus  $\sigma_X^{\text{Sets}_*}$  is a morphism of pointed sets.

### Invertibility

The inverse of  $\sigma_{X,Y}^{\text{Sets}_*}$  is given by the morphism

$$\sigma_{X,Y}^{\text{Sets}_*, -1}: Y \wedge X \xrightarrow{\sim} X \wedge Y$$

defined by

$$\sigma_{X,Y}^{\text{Sets}_*, -1}(y \wedge x) \stackrel{\text{def}}{=} x \wedge y$$

for each  $y \wedge x \in Y \wedge X$ .

### Naturality

We need to show that, given morphisms of pointed sets

$$\begin{aligned} f: (X, x_0) &\rightarrow (A, a_0), \\ g: (Y, y_0) &\rightarrow (B, b_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{f \wedge g} & A \wedge B \\ \sigma_{X,Y}^{\text{Sets}_*} \downarrow & & \downarrow \sigma_{A,B}^{\text{Sets}_*} \\ Y \wedge X & \xrightarrow{g \wedge f} & B \wedge A \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge y & \longmapsto & f(x) \wedge g(y) \\ \downarrow & & \downarrow \\ y \wedge x & \longmapsto & g(y) \wedge f(x) \end{array}$$

and hence indeed commutes, showing  $\sigma^{\text{Sets}_*}$  to be a natural transformation.

#### Being a Natural Isomorphism

Since  $\sigma^{\text{Sets}_*}$  is natural and  $\sigma^{\text{Sets}_*, -1}$  is a componentwise inverse to  $\sigma^{\text{Sets}_*}$ , it follows from Item 2 of Proposition 11.9.7.1.2 that  $\sigma^{\text{Sets}_*, -1}$  is also natural. Thus  $\sigma^{\text{Sets}_*}$  is a natural isomorphism. □

### 00GT 7.5.8 The Diagonal

00GU

#### DEFINITION 7.5.8.1.1 ► THE DIAGONAL OF $\wedge$

The **diagonal of the smash product of pointed sets** is the natural transformation

$$\Delta^\wedge: \text{id}_{\text{Sets}_*} \Rightarrow \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

whose component

$$\Delta_X^\wedge: (X, x_0) \rightarrow (X \wedge X, x_0 \wedge x_0)$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by the composition

$$\begin{aligned} (X, x_0) &\xrightarrow{\Delta_X^\wedge} (X \times X, (x_0, x_0)) \\ &\longrightarrow ((X \times X)/\sim, [(x_0, x_0)]) \\ &\stackrel{\text{def}}{=} (X \wedge X, x_0 \wedge x_0) \end{aligned}$$

in  $\text{Sets}_*$ , and thus by

$$\Delta_X^\wedge(x) \stackrel{\text{def}}{=} x \wedge x$$

for each  $x \in X$ .

#### PROOF 7.5.8.1.2 ► PROOF OF DEFINITION 7.5.8.1.1

#### Being a Morphism of Pointed Sets

We have

$$\Delta_X^\wedge(x_0) \stackrel{\text{def}}{=} x_0 \wedge x_0,$$

and thus  $\Delta_X^\wedge$  is a morphism of pointed sets.

### Naturality

We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\wedge \downarrow & & \downarrow \Delta_Y^\wedge \\ X \wedge X & \xrightarrow{f \wedge f} & Y \wedge Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ x \wedge x & \longmapsto & f(x) \wedge f(x) \end{array}$$

and hence indeed commutes, showing  $\Delta^\wedge$  to be natural. □

00GV

### PROPOSITION 7.5.8.1.3 ▶ PROPERTIES OF THE DIAGONAL OF $\wedge$

Let  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00GW

1. *Monoidality.* The diagonal

$$\Delta^\wedge: \text{id}_{\text{Sets}_*} \Longrightarrow \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

of the smash product of pointed sets is a monoidal natural transformation:

00GX

(a) *Compatibility With Strong Monoidality Constraints.* For each  $(X, x_0), (Y, y_0) \in \text{Obj}(\text{Sets}_*)$ , the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{\Delta_X^\wedge \wedge \Delta_Y^\wedge} & (X \wedge X) \wedge (Y \wedge Y) \\ & \searrow \Delta_{X \wedge Y}^\wedge & \downarrow \lrcorner \\ & & (X \wedge Y) \wedge (X \wedge Y) \end{array}$$

commutes.

00GY

(b) *Compatibility With Strong Unitality Constraints.* The diagrams

$$\begin{array}{ccc} S^0 & \xrightarrow{\Delta_{S^0}^\wedge} & S^0 \wedge S^0 \\ \searrow & \downarrow \lambda_{S^0}^{\text{Sets}_*, -1} & \searrow \\ & S^0 & \end{array} \quad \begin{array}{ccc} S^0 & \xrightarrow{\Delta_{S^0}^\wedge} & S^0 \wedge S^0 \\ \searrow & \downarrow \rho_{S^0}^{\text{Sets}_*, -1} & \searrow \\ & S^0 & \end{array}$$

commute, i.e. we have

$$\begin{aligned} \Delta_{S^0}^\wedge &= \lambda_{S^0}^{\text{Sets}_*, -1} \\ &= \rho_{S^0}^{\text{Sets}_*, -1}, \end{aligned}$$

where we recall that the equalities

$$\begin{aligned} \lambda_{S^0}^{\text{Sets}_*} &= \rho_{S^0}^{\text{Sets}_*}, \\ \lambda_{S^0}^{\text{Sets}_*, -1} &= \rho_{S^0}^{\text{Sets}_*, -1} \end{aligned}$$

are always true in any monoidal category by ?? of ??.

00GZ

2. *The Diagonal of the Unit.* The component

$$\Delta_{S^0}^\wedge : S^0 \xrightarrow{\sim} S^0 \wedge S^0$$

of  $\Delta^\wedge$  at  $S^0$  is an isomorphism.

#### PROOF 7.5.8.1.4 ► PROOF OF PROPOSITION 7.5.8.1.3

**Item 1: Monoidality**

We claim that  $\Delta^\wedge$  is indeed monoidal:

0288

1. **Item 1a: Compatibility With Strong Monoidality Constraints:** We need to show that the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{\Delta_X^\wedge \wedge \Delta_Y^\wedge} & (X \wedge X) \wedge (Y \wedge Y) \\ & \searrow \Delta_{X \wedge Y}^\wedge & \downarrow \text{?} \\ & & (X \wedge Y) \wedge (X \wedge Y) \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge y & \xrightarrow{\quad} & (x \wedge x) \wedge (y \wedge y) \\ & \swarrow & \downarrow \\ & & (x \wedge y) \wedge (x \wedge y) \end{array}$$

and hence indeed commutes.

0289

2. *Item 1b: Compatibility With Strong Unitality Constraints:* As shown in the proof of [Definition 7.5.5.1.1](#), the inverse of the left unit of  $\text{Sets}_*$  with respect to the smash product of pointed sets at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\lambda_X^{\text{Sets}_*, -1}(x) \stackrel{\text{def}}{=} 1 \wedge x$$

for each  $x \in X$ , so when  $X = S^0$ , we have

$$\begin{aligned} \lambda_{S^0}^{\text{Sets}_*, -1}(0) &\stackrel{\text{def}}{=} 1 \wedge 0, \\ \lambda_{S^0}^{\text{Sets}_*, -1}(1) &\stackrel{\text{def}}{=} 1 \wedge 1. \end{aligned}$$

But since  $1 \wedge 0 = 0 \wedge 0$  and

$$\begin{aligned} \Delta_{S^0}^\wedge(0) &\stackrel{\text{def}}{=} 0 \wedge 0, \\ \Delta_{S^0}^\wedge(1) &\stackrel{\text{def}}{=} 1 \wedge 1, \end{aligned}$$

it follows that we indeed have  $\Delta_{S^0}^\wedge = \lambda_{S^0}^{\text{Sets}_*, -1}$ .

This finishes the proof.

**Item 2: The Diagonal of the Unit**

This follows from [Item 1](#) and the invertibility of the left/right unit of  $\text{Sets}_*$  with respect to  $\wedge$ , proved in the proof of [Definition 7.5.5.1.1](#) for the left unit or the proof of [Definition 7.5.6.1.1](#) for the right unit. ☐

00H0 7.5.9 The Monoidal Structure on Pointed Sets Associated to  $\wedge$ 

00H1 PROPOSITION 7.5.9.1.1 ▶ THE MONOIDAL STRUCTURE ON POINTED SETS ASSOCIATED TO

$\wedge$

The category  $\text{Sets}_*$  admits a closed monoidal category with diagonals structure consisting of:

- *The Underlying Category.* The category  $\text{Sets}_*$  of pointed sets.

- *The Monoidal Product.* The smash product functor

$$\wedge : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Item 1 of [Proposition 7.5.1.1.12](#).

- *The Internal Hom.* The internal Hom functor

$$\mathbf{Sets}_* : \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Item 1 of [Proposition 7.5.2.1.2](#).

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*$$

of [Definition 7.5.3.1.1](#).

- *The Associators.* The natural isomorphism

$$\alpha^{\text{Sets}_*} : \wedge \circ (\wedge \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \wedge \circ (\text{id}_{\text{Sets}_*} \times \wedge) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}}$$

of [Definition 7.5.4.1.1](#).

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\text{Sets}_*} : \wedge \circ (\mathbb{1}^{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \lambda_{\text{Sets}_*}^{\text{Cats}_2}$$

of [Definition 7.5.5.1.1](#).

- *The Right Unitors.* The natural isomorphism

$$\rho^{\text{Sets}_*} : \wedge \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}) \xrightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}$$

of [Definition 7.5.6.1.1](#).

- *The Symmetry.* The natural isomorphism

$$\sigma^{\text{Sets}_*} : \wedge \xrightarrow{\sim} \wedge \circ \sigma_{\text{Sets}_*, \text{Sets}_*}^{\text{Cats}_2}$$

of [Definition 7.5.7.1.1](#).

- *The Diagonals.* The monoidal natural transformation

$$\Delta^\wedge : \text{id}_{\text{Sets}_*} \Longrightarrow \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2}$$

of [Definition 7.5.8.1.1](#).

**PROOF 7.5.9.1.2 ► PROOF OF PROPOSITION 7.5.9.1.1****The Pentagon Identity**

Let  $(W, w_0)$ ,  $(X, x_0)$ ,  $(Y, y_0)$  and  $(Z, z_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccccc}
 & & (W \wedge (X \wedge Y)) \wedge Z & & \\
 & \swarrow \alpha_{W,X,Y}^{\text{Sets}_*} \wedge \text{id}_Z & & \searrow \alpha_{W,X \wedge Y,Z}^{\text{Sets}_*} & \\
 ((W \wedge X) \wedge Y) \wedge Z & & & & W \wedge ((X \wedge Y) \wedge Z) \\
 & \downarrow \alpha_{W \wedge X,Y,Z}^{\text{Sets}_*} & & \downarrow \text{id}_W \wedge \alpha_{X,Y,Z}^{\text{Sets}_*} & \\
 & (W \wedge X) \wedge (Y \wedge Z) & \xrightarrow{\alpha_{W,X,Y \wedge Z}^{\text{Sets}_*}} & & W \wedge (X \wedge (Y \wedge Z))
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccccc}
 & & (w \wedge (x \wedge y)) \wedge z & & \\
 & \swarrow & & \searrow & \\
 ((w \wedge x) \wedge y) \wedge z & & & & w \wedge ((x \wedge y) \wedge z) \\
 & \downarrow & & \downarrow & \\
 & (w \wedge x) \wedge (y \wedge z) & \longmapsto & & w \wedge (x \wedge (y \wedge z))
 \end{array}$$

and thus we see that the pentagon identity is satisfied.

**The Triangle Identity**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc} (X \wedge S^0) \wedge Y & \xrightarrow{\alpha_{X,S^0,Y}^{\text{Sets}_*}} & X \wedge (S^0 \wedge Y) \\ & \searrow \rho_X^{\text{Sets}_*} \wedge \text{id}_Y & \swarrow \text{id}_X \wedge \lambda_Y^{\text{Sets}_*} \\ & X \wedge Y & \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \wedge 0) \wedge y & \xleftarrow{\quad} & (x \wedge 0) \wedge y \xrightarrow{\quad} x \wedge (0 \wedge y) \\ \downarrow & & \downarrow \\ x_0 \wedge y & & x \wedge y_0 \end{array}$$

and

$$\begin{array}{ccc} (x \wedge 1) \wedge y & \xrightarrow{\quad} & x \wedge (1 \wedge y) \\ \downarrow & & \downarrow \\ x \wedge y & & x \wedge y \end{array}$$

and thus we see that the triangle identity is satisfied.

### The Left Hexagon Identity

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccccc} & & (X \wedge Y) \wedge Z & & \\ & \swarrow \alpha_{X,Y,Z}^{\text{Sets}_*} & & \searrow \beta_{X,Y}^{\text{Sets}_*} \wedge \text{id}_Z & \\ X \wedge (Y \wedge Z) & & & & (Y \wedge X) \wedge Z \\ \downarrow \beta_{X,Y \wedge Z}^{\text{Sets}_*} & & & & \downarrow \alpha_{Y,X,Z}^{\text{Sets}_*} \\ (Y \wedge Z) \wedge X & & & & Y \wedge (X \wedge Z) \\ & \searrow \alpha_{Y,Z,X}^{\text{Sets}_*} & & \swarrow \text{id}_Y \wedge \beta_{X,Z}^{\text{Sets}_*} & \\ & & Y \wedge (Z \wedge X) & & \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccccc}
 & & (x \wedge y) \wedge z & & \\
 & \swarrow & & \searrow & \\
 x \wedge (y \wedge z) & & & & (y \wedge x) \wedge z \\
 \downarrow & & & & \downarrow \\
 (y \wedge z) \wedge x & & & & y \wedge (x \wedge z) \\
 \swarrow & & & & \searrow \\
 & & y \wedge (z \wedge x) & &
 \end{array}$$

and thus we see that the left hexagon identity is satisfied.

### The Right Hexagon Identity

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccccc}
 & X \wedge (Y \wedge Z) & & & \\
 (\alpha_{X,Y,Z}^{\text{Sets}_*})^{-1} \swarrow & & \searrow \text{id}_X \wedge \beta_{Y,Z}^{\text{Sets}_*} & & \\
 (X \wedge Y) \wedge Z & & & & X \wedge (Z \wedge Y) \\
 \downarrow \beta_{X \wedge Y, Z}^{\text{Sets}_*} & & & & \downarrow (\alpha_{X,Z,Y}^{\text{Sets}_*})^{-1} \\
 Z \wedge (X \wedge Y) & & & & (X \wedge Z) \wedge Y \\
 \swarrow (\alpha_{Z,X,Y}^{\text{Sets}_*})^{-1} & & \searrow \beta_{X,Z}^{\text{Sets}_*} \wedge \text{id}_Y & & \\
 (Z \wedge X) \wedge Y & & & &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccccc}
 & & x \wedge (y \wedge z) & & \\
 & \swarrow & & \searrow & \\
 (x \wedge y) \wedge z & & & & x \wedge (z \wedge y) \\
 \downarrow & & & & \downarrow \\
 z \wedge (x \wedge y) & & & & (x \wedge z) \wedge y \\
 \swarrow & & & \searrow & \\
 (z \wedge x) \wedge y & & & &
 \end{array}$$

and thus we see that the right hexagon identity is satisfied.

Monoidal Closedness

This follows from [Item 2 of Proposition 7.5.1.1.12](#).

Existence of Monoidal Diagonals

This follows from [Items 1 and 2 of Proposition 7.5.8.1.3](#). 

### 01VY 7.5.10 The Universal Property of $(\text{Sets}_*, \wedge, S^0)$

01VZ

#### THEOREM 7.5.10.1.1 ► THE UNIVERSAL PROPERTY OF $(\text{Sets}_*, \wedge, S^0)$

The symmetric monoidal structure on the category  $\text{Sets}_*$  of [Proposition 7.5.9.1.1](#) is uniquely determined by the following requirements:

01W0

1. *Existence of an Internal Hom.* The tensor product

$$\otimes_{\text{Sets}_*} : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of  $\text{Sets}_*$  admits an internal Hom  $[-_1, -_2]_{\text{Sets}_*}$ .

01W1

2. *The Unit Object Is  $S^0$ .* We have  $1_{\text{Sets}_*} \cong S^0$ .

More precisely, the full subcategory of the category  $\mathcal{M}_{\mathbb{E}_\infty}^{\text{cl}}(\text{Sets}_*)$  of ?? spanned by the closed symmetric monoidal categories  $(\text{Sets}_*, \otimes_{\text{Sets}_*}, [-_1, -_2]_{\text{Sets}_*}, 1_{\text{Sets}_*}, \lambda^{\text{Sets}_*}, \rho^{\text{Sets}_*}, \sigma^{\text{Sets}_*})$  satisfying [Items 1 and 2](#) is contractible (i.e. equivalent to the punctual category).

## PROOF 7.5.10.1.2 ► PROOF OF THEOREM 7.5.10.1.1

## Unwinding the Statement

Let  $(\text{Sets}_*, \otimes_{\text{Sets}_*}, [-_1, -_2]_{\text{Sets}_*}, \mathbb{1}_{\text{Sets}_*}, \lambda', \rho', \sigma')$  be a closed symmetric monoidal category satisfying **Items 1** and **2**. We need to show that the identity functor

$$\text{id}_{\text{Sets}_*} : \text{Sets}_* \rightarrow \text{Sets}_*$$

admits a *unique* closed symmetric monoidal functor structure

$$\begin{aligned}\text{id}_{\text{Sets}_*}^{\otimes} &: X \otimes_{\text{Sets}_*} Y \xrightarrow{\sim} X \wedge Y, \\ \text{id}_{\text{Sets}_*}^{\text{Hom}} &: [X, Y]_{\text{Sets}_*} \xrightarrow{\sim} \text{Sets}_*(X, Y), \\ \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\otimes} &: \mathbb{1}_{\text{Sets}_*} \xrightarrow{\sim} S^0,\end{aligned}$$

making it into a symmetric monoidal strongly closed isomorphism of categories from  $(\text{Sets}_*, \otimes_{\text{Sets}_*}, [-_1, -_2]_{\text{Sets}_*}, \mathbb{1}_{\text{Sets}_*}, \lambda', \rho', \sigma')$  to the closed symmetric monoidal category  $(\text{Sets}_*, \times, \text{Sets}_*(-_1, -_2), \mathbb{1}_{\text{Sets}_*}, \lambda^{\text{Sets}_*}, \rho^{\text{Sets}_*}, \sigma^{\text{Sets}_*})$  of **Proposition 7.5.9.1.1**.

Constructing an Isomorphism  $[-_1, -_2]_{\text{Sets}_*} \cong \text{Sets}_*(-_1, -_2)$ 

By ??, we have a natural isomorphism

$$\text{Sets}_*(S^0, [-_1, -_2]_{\text{Sets}_*}) \cong \text{Sets}_*(-_1, -_2).$$

By **Item 4** of **Proposition 6.1.4.1.1**, we also have a natural isomorphism

$$\text{Sets}_*(S^0, [-_1, -_2]_{\text{Sets}_*}) \cong [-_1, -_2]_{\text{Sets}_*}.$$

Composing both natural isomorphisms, we obtain a natural isomorphism

$$\text{Sets}_*(-_1, -_2) \cong [-_1, -_2]_{\text{Sets}_*}.$$

Given  $X, Y \in \text{Obj}(\text{Sets}_*)$ , we will write

$$\text{id}_{X,Y}^{\text{Hom}} : \text{Sets}_*(X, Y) \xrightarrow{\sim} [X, Y]_{\text{Sets}_*}$$

for the component of this isomorphism at  $(X, Y)$ .

Constructing an Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$ 

Since  $\otimes_{\text{Sets}_*}$  is adjoint in each variable to  $[-_1, -_2]_{\text{Sets}_*}$  by assumption and  $\wedge$  is adjoint in each variable to  $\text{Sets}_*(-_1, -_2)$  by **Item 2** of **Proposition 4.3.5.1.2**, uniqueness of adjoints ?? gives us natural isomorphisms

$$X \otimes_{\text{Sets}_*} - \cong X \wedge -,$$

$$- \otimes_{\text{Sets}_*} Y \cong Y \wedge -.$$

By ??, we then have  $\otimes_{\text{Sets}_*} \cong \wedge$ . We will write

$$\text{id}_{\text{Sets}_*|X,Y}^{\otimes}: X \otimes_{\text{Sets}_*} Y \xrightarrow{\sim} X \wedge Y$$

for the component of this isomorphism at  $(X, Y)$ .

**Alternative Construction of an Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$**

Alternatively, we may construct a natural isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$  as follows:

01W2 1. Let  $X \in \text{Obj}(\text{Sets}_*)$ .

01W3 2. Since  $\otimes_{\text{Sets}_*}$  is part of a closed monoidal structure, it preserves colimits in each variable by ??.

01W4 3. Since  $X \cong \bigvee_{x \in X^-} S^0$  and  $\otimes_{\text{Sets}_*}$  preserves colimits in each variable, we have

$$\begin{aligned} X \otimes_{\text{Sets}_*} Y &\cong \left( \bigvee_{x \in X^-} S^0 \right) \otimes_{\text{Sets}_*} Y \\ &\cong \bigvee_{x \in X^-} (S^0 \otimes_{\text{Sets}_*} Y) \\ &\cong \bigvee_{x \in X^-} Y \\ &\cong \bigvee_{x \in X^-} S^0 \wedge Y \\ &\cong \left( \bigvee_{x \in X^-} S^0 \right) \wedge Y \\ &\cong X \wedge Y, \end{aligned}$$

naturally in  $Y \in \text{Obj}(\text{Sets}_*)$ , where we have used that  $S^0$  is the monoidal unit for  $\otimes_{\text{Sets}_*}$ . Thus  $X \otimes_{\text{Sets}_*} - \cong X \wedge -$  for each  $X \in \text{Obj}(\text{Sets}_*)$ .

01W5 4. Similarly,  $- \otimes_{\text{Sets}_*} Y \cong - \wedge Y$  for each  $Y \in \text{Obj}(\text{Sets}_*)$ .

01W6 5. By ??, we then have  $\otimes_{\text{Sets}_*} \cong \wedge$ .

Below, we'll show that if a natural isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$  exists, then it must be unique. This will show that the isomorphism constructed above is equal to the isomorphism  $\text{id}_{\text{Sets}_*|X,Y}^{\otimes}: X \otimes_{\text{Sets}_*} Y \rightarrow X \wedge Y$  from before.

Constructing an Isomorphism  $\text{id}_{\mathbb{1}}^{\otimes} : \mathbb{1}_{\text{Sets}_*} \rightarrow S^0$

We define an isomorphism  $\text{id}_{\mathbb{1}}^{\otimes} : \mathbb{1}_{\text{Sets}_*} \rightarrow S^0$  as the composition

$$\mathbb{1}_{\text{Sets}_*} \xrightarrow[\sim]{\rho_{\mathbb{1}\text{Sets}_*}^{\text{Sets}_*, -1}} \mathbb{1}_{\text{Sets}_*} \wedge S^0 \xrightarrow[\sim]{\text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_*}^{\otimes, -1}} \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 \xrightarrow[\sim]{\lambda'_{S^0}} S^0$$

in  $\text{Sets}_*$ .

Monoidal Left Unity of the Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$

We have to show that the diagram

$$\begin{array}{ccc} S^0 \otimes_{\text{Sets}_*} X & \xrightarrow{\text{id}_{S^0 \otimes_{\text{Sets}_*} X}^{\otimes}} & S^0 \wedge X \\ \text{id}_{\mathbb{1}\text{Sets}_*}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_X & \nearrow & \searrow \lambda_X^{\text{Sets}_*} \\ \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} X & \xrightarrow{\lambda'_X} & X \end{array}$$

commutes. To this end, we will first show that the diagram

$$\begin{array}{ccc} S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{S^0 \wedge S^0}^{\otimes}} & S^0 \wedge S^0 \\ \text{id}_{\mathbb{1}\text{Sets}_*}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_{S^0} & \nearrow & \searrow \lambda_{S^0}^{\text{Sets}_*} \\ \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\lambda'_{S^0}} & S^0, \end{array} \quad (\dagger)$$

corresponding to the case  $X = S^0$ , commutes. Indeed, consider the diagram

$$\begin{array}{ccccccc} \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\rho_{\mathbb{1}\text{Sets}_*, S^0}^{\text{Sets}_*, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0}} & (\mathbb{1}_{\text{Sets}_*} \wedge S^0) \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{(\mathbb{1}_{\text{Sets}_*} \wedge S^0) \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0}} & (\mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\lambda'_{S^0} \otimes_{\text{Sets}_*} \text{id}_{S^0}} & S^0 \otimes_{\text{Sets}_*} S^0 \\ \downarrow \text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_* S^0}^{\otimes} & (1) & \downarrow \text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_* S^0}^{\otimes} & (2) & \downarrow \text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_* S^0}^{\otimes} & (3) & \downarrow \text{id}_{\text{Sets}_*|S^0 S^0}^{\otimes} \\ \mathbb{1}_{\text{Sets}_*} \wedge S^0 & \xrightarrow{\rho_{\mathbb{1}\text{Sets}_* \wedge S^0}^{\text{Sets}_*, -1} \wedge \text{id}_{S^0}} & (\mathbb{1}_{\text{Sets}_*} \wedge S^0) \wedge S^0 & \xrightarrow{-\text{id}_{\mathbb{1}\text{Sets}_* \wedge S^0}^{\otimes, -1} \wedge \text{id}_{S^0}} & (\mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0) \wedge S^0 & \xrightarrow{\lambda'_{S^0} \wedge \text{id}_{S^0}} & S^0 \wedge S^0 \\ \downarrow \text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_* S^0}^{\otimes} & (4) & \downarrow \text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_* S^0}^{\otimes} & (5) & \downarrow \text{id}_{\text{Sets}_*|\mathbb{1}\text{Sets}_* S^0}^{\otimes} & (6) & \downarrow \text{id}_{\text{Sets}_*|S^0 S^0}^{\otimes} \\ \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & & S^0 \end{array}$$

whose boundary diagram corresponds to the diagram (†) above. In this diagram:

- Subdiagrams (1), (2), and (3) commute by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes}$ .
- Subdiagram (4) commutes by ??.
- Subdiagram (5) commutes by the naturality of  $\rho^{\text{Sets}_*, -1}$ .
- Subdiagram (6) commutes trivially.
- Subdiagram (7) commutes by the naturality of  $\rho^{\text{Sets}_*}$ , where the equality  $\rho_{S^0}^{\text{Sets}_*} = \lambda_{S^0}^{\text{Sets}_*}$  comes from ??.

Since all subdiagrams commute, so does the boundary diagram, i.e. the diagram (†) above. As a result, the diagram

$$\begin{array}{ccc}
 S^0 \wedge S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0, S^0}^{\otimes, -1}} & S^0 \otimes_{\text{Sets}_*} S^0 \\
 \lambda_{S^0}^{\text{Sets}_*, -1} \nearrow & & \searrow \text{id}_{\mathbb{1}|S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0} \\
 S^0 & \xrightarrow{\lambda'^{-1}_{S^0}} & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0
 \end{array}$$

also commutes. Now, let  $X \in \text{Obj}(\text{Sets}_*)$ , let  $x \in X$ , and consider the diagram

$$\begin{array}{ccccc}
 S^0 \wedge S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0, S^0}^{\otimes, -1}} & S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\lambda'^{-1}_{S^0} \wedge \text{id}_{S^0}} & S^0 \otimes_{\text{Sets}_*} S^0 \\
 \lambda_{S^0}^{\text{Sets}_*, -1} \nearrow & & & \searrow & \\
 S^0 & \xrightarrow{\lambda'^{-1}_{S^0}} & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & & \\
 \downarrow \text{id}_{S^0} \wedge [x] & & \downarrow \text{id}_{S^0} \otimes_{\text{Sets}_*} [x] & & \downarrow \text{id}_{\mathbb{1}_{\text{Sets}_*}} \wedge [x] \\
 X & \xrightarrow{\lambda_X'^{-1}} & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} X & & 
 \end{array}$$

(1)      (2)      (3)      (4)

(5)      (6)      (7)

Since:

- Subdiagram (5) commutes by the naturality of  $\lambda'^{-1}$ .
- Subdiagram (‡) commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $\text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1}$ .
- Subdiagram (1) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes, -1}$ .
- Subdiagram (3) commutes by the naturality of  $\lambda^{\text{Sets}_*, -1}$ .

it follows that the diagram

$$\begin{array}{ccccc}
 & S^0 \wedge X & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,X}^{\otimes, -1}} & S^0 \otimes_{\text{Sets}_*} X & \\
 \lambda_X^{\text{Sets}_*, -1} \nearrow & & & & \searrow \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_X \\
 S^0 \xrightarrow{[x]} X & \xrightarrow{\lambda'_X^{-1}} & & & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} X
 \end{array}$$

Here's a step-by-step showcase of this argument: [\[Link\]](#). We then have

$$\begin{aligned}
 \lambda'_X^{-1}(x) &= [\lambda'_X^{-1} \circ [x]](1) \\
 &= [( \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1} \wedge \text{id}_X ) \circ \text{id}_{\text{Sets}_*|S^0,X}^{\otimes, -1} \circ \lambda_X^{\text{Sets}_*, -1} \circ [x])(1) \\
 &= [( \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1} \wedge \text{id}_X ) \circ \text{id}_{\text{Sets}_*|S^0,X}^{\otimes, -1} \circ \lambda_X^{\text{Sets}_*, -1}](x)
 \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\lambda'_X^{-1} = (\text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1} \wedge \text{id}_X) \circ \text{id}_{\text{Sets}_*|S^0,X}^{\otimes, -1} \circ \lambda_X^{\text{Sets}_*, -1}.$$

Taking inverses then gives

$$\lambda'_X = \lambda_X^{\text{Sets}_*} \circ \text{id}_{\text{Sets}_*|S^0,X}^{\otimes} \circ (\text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes} \wedge \text{id}_X),$$

showing that the diagram

$$\begin{array}{ccccc}
 & S^0 \otimes_{\text{Sets}_*} X & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,X}^{\otimes}} & S^0 \wedge X & \\
 \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_X \nearrow & & & & \searrow \lambda_X^{\text{Sets}_*} \\
 \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} X & \xrightarrow{\lambda'_X} & & & X
 \end{array}$$

indeed commutes.

Braidedness of the Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$

We have to show that the diagram

$$\begin{array}{ccc} X \otimes_{\text{Sets}_*} Y & \xrightarrow{\text{id}_{\text{Sets}_*|X,Y}^\otimes} & X \wedge Y \\ \sigma'_{X,Y} \downarrow & & \downarrow \sigma_{X,Y}^{\text{Sets}_*} \\ Y \otimes_{\text{Sets}_*} X & \xrightarrow{\text{id}_{\text{Sets}_*|Y,X}^\otimes} & Y \wedge X \end{array}$$

commutes. To this end, we will first show that the diagram

$$\begin{array}{ccc} S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,S^0}^\otimes} & S^0 \wedge S^0 \\ \sigma'_{S^0,S^0} \downarrow & (\dagger) & \downarrow \sigma_{S^0,S^0}^{\text{Sets}_*} \\ S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,S^0}^\otimes} & S^0 \wedge S^0 \end{array}$$

commutes. To that end, we will first show that the diagram

$$\begin{array}{ccc} S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,\mathbb{1}_{\text{Sets}_*}}^\otimes} & S^0 \wedge \mathbb{1}_{\text{Sets}_*} \\ \sigma'_{S^0,\mathbb{1}_{\text{Sets}_*}} \downarrow & (\ddagger) & \downarrow \sigma_{S^0,\mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*} \\ \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|\mathbb{1}_{\text{Sets}_*},S^0}^\otimes} & \mathbb{1}_{\text{Sets}_*} \wedge S^0 \end{array}$$

commutes, and, to this end, we will first show that the diagram

$$\begin{array}{ccc} S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,S^0}^\otimes} & S^0 \wedge S^0 \\ \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\text{Sets}_*|\mathbb{1}}^\otimes \uparrow & (\S) & \downarrow \lambda_{S^0}^{\text{Sets}_*} \\ S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\rho'_{S^0}} & S^0 \end{array}$$

commutes. Indeed, consider the diagram

The diagram consists of several nodes and arrows. At the top right is  $S^0 \wedge S^0$ . At the top left is  $S^0 \otimes_{Sets_*} S^0$ . At the bottom right is  $S^0$ . At the bottom left is  $S^0 \otimes_{Sets_*} \mathbb{1}_{Sets_*}$ . On the right side, there is a vertical arrow  $\lambda_{S^0}^{Sets_*}$  pointing down from  $S^0 \wedge S^0$  to  $S^0$ . On the left side, there is a vertical arrow  $\text{id}_{S^0} \otimes_{Sets_*} \text{id}_{\mathbb{1}|Sets_*}^\otimes$  pointing down from  $S^0 \otimes_{Sets_*} S^0$  to  $S^0 \otimes_{Sets_*} \mathbb{1}_{Sets_*}$ . In the center, there is a node  $\mathbb{1}_{Sets_*} \otimes_{Sets_*} S^0$  with an upward arrow  $\text{id}_{Sets_*|1_{Sets_*},S^0}^\otimes$  from below and a downward arrow  $\text{id}_{Sets_*|1_{Sets_*},S^0}^\otimes$  to the bottom center node. The bottom center node has a rightward arrow  $\lambda'_{S^0}$  to the bottom right  $S^0$ . The bottom left node has a rightward arrow  $\text{id}_{\mathbb{1}|Sets_*}^\otimes$  to the bottom right  $S^0$ . The top left node has two diagonal arrows: one to the top right  $S^0 \wedge S^0$  and one to the bottom left  $S^0 \otimes_{Sets_*} \mathbb{1}_{Sets_*}$ . The top right node has two diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$  and one to the bottom right  $S^0$ . The center node has three diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$ , one to the top right  $S^0 \wedge S^0$ , and one to the bottom center node. The bottom center node has three diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$ , one to the top right  $S^0 \wedge S^0$ , and one to the bottom right  $S^0$ . The bottom left node has two diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$  and one to the top right  $S^0 \wedge S^0$ . The top left node has two diagonal arrows: one to the top right  $S^0 \wedge S^0$  and one to the bottom left  $S^0 \otimes_{Sets_*} \mathbb{1}_{Sets_*}$ . The top right node has two diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$  and one to the bottom right  $S^0$ . The center node has two diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$  and one to the top right  $S^0 \wedge S^0$ . The bottom center node has two diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$  and one to the top right  $S^0 \wedge S^0$ . The bottom left node has two diagonal arrows: one to the top left  $S^0 \otimes_{Sets_*} S^0$  and one to the top right  $S^0 \wedge S^0$ .

whose boundary diagram corresponds to diagram (§) above. Since:

- Subdiagram (1) commutes by the naturality of  $\text{id}_{Sets_*}^\otimes$ ;
- Subdiagrams (2) and (3) commute by the functoriality of  $\otimes$ ;
- Subdiagram (4) commutes by the left monoidal unity of  $(\text{id}^\otimes, \text{id}_\mathbb{1}^\otimes)$ , which we proved above;
- Subdiagram (5) commutes by the naturality of  $\lambda'$ ;
- Subdiagram (6) commutes by the naturality of  $\rho'$ , where the equality  $\rho'_{\mathbb{1}_{Sets_*}} = \lambda'_{\mathbb{1}_{Sets_*}}$  comes from ??;

it follows that the boundary diagram, i.e. diagram (§), also commutes.

Next, consider the diagram

$$\begin{array}{ccccc}
 & S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\quad \text{id}_{\text{Sets}_*|S^0, \mathbb{1}_{\text{Sets}_*}}^{\otimes} \quad} & S^0 \wedge \mathbb{1}_{\text{Sets}_*} & \\
 & \downarrow & & \downarrow & \\
 & \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\otimes} & \searrow & \swarrow \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\otimes-1} & \\
 & & (2) & & \\
 & S^0 \otimes_{\text{Sets}_*} S^0 - \text{id}_{\text{Sets}_*|S^0, S^0}^{\otimes} & \rightarrow & S^0 \wedge S^0 & \\
 & \downarrow & & \downarrow \lambda_{S^0}^{\text{Sets}_*} & \downarrow \lambda_{\mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*} \\
 & (§) & & (3) & (6) \\
 & \downarrow \sigma'_{S^0, \mathbb{1}_{\text{Sets}_*}} & & \downarrow \sigma_{S^0, \mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*} & \downarrow \rho_{\mathbb{1}_{\text{Sets}_*}, -1}^{\text{Sets}_*} \\
 & S^0 & \xrightarrow{\quad \text{id}_{S^0}^{\otimes-1} \quad} & \mathbb{1}_{\text{Sets}_*} & \\
 & \downarrow \lambda'^{-1}_{S^0} & & \downarrow \text{id}_{\mathbb{1}_{\text{Sets}_*}|1}^{\otimes-1} & \downarrow \rho_{\mathbb{1}_{\text{Sets}_*}, S^0}^{\text{Sets}_*} \\
 & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\quad \text{id}_{\text{Sets}_*|S^0}^{\otimes} \quad} & \mathbb{1}_{\text{Sets}_*} \wedge S^0 &
 \end{array}$$

whose boundary diagram corresponds to the diagram (‡) above. Since:

- Subdiagrams (1) and (6) commute by ??;
- Subdiagram (2) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes}$ ;
- Subdiagram (§) commutes, as was shown above;
- Subdiagram (3) commutes by the naturality of  $\lambda^{\text{Sets}_*}$ ;
- Subdiagram (4) commutes trivially;
- Subdiagram (5) commutes by Item 2c of Item 2 of Proposition 13.1.1.4, whose proof uses only the left monoidal unity of  $(\text{id}^{\otimes}, \text{id}_{\mathbb{1}})$ , which has been proven above;

it follows that the boundary diagram, i.e. diagram (‡), also commutes.

Next, consider the diagram

$$\begin{array}{ccccc}
 S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\quad \text{id}_{S^0, S^0}^\otimes \quad} & S^0 \wedge S^0 \\
 \downarrow \sigma'_{S^0, S^0} & \searrow \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\mathbb{1}}^{\otimes, -1} & (1) & \swarrow \text{id}_{S^0} \wedge \text{id}_{\mathbb{1}}^{\otimes, -1} & \downarrow \sigma_{S^0, S^0}^{\text{Sets}_*} \\
 & & S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\quad \text{id}_{S^0, \mathbb{1}_{\text{Sets}_*}}^\otimes \quad} & S^0 \wedge \mathbb{1}_{\text{Sets}_*} \\
 & (2) & \downarrow \sigma'_{S^0, \mathbb{1}_{\text{Sets}_*}} & & (3) \\
 & & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\quad \text{id}_{\mathbb{1}_{\text{Sets}_*}, S^0}^\otimes \quad} & \mathbb{1}_{\text{Sets}_*} \wedge S^0 \\
 & & \downarrow \text{id}_{\mathbb{1}}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_{S^0} & & \downarrow \text{id}_{\mathbb{1}}^{\otimes} \wedge \text{id}_{S^0} \\
 & & (4) & & 
 \end{array}$$

whose boundary diagram corresponds to the diagram  $(\dagger)$ . Since:

- Subdiagram (1) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^\otimes$ ;
- Subdiagram (2) commutes by the naturality of  $\sigma'$  and the fact that  $\text{id}_{\mathbb{1}}^\otimes$  is invertible;
- Subdiagram  $(\ddagger)$  commutes as proved above;
- Subdiagram (3) commutes by the naturality of  $\sigma_{S^0, \mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*}$  and the fact that  $\text{id}_{\mathbb{1}}^\otimes$  is invertible;
- Subdiagram (4) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^\otimes$ ;

it follows that the boundary diagram, i.e. diagram  $(\dagger)$  also commutes.

Taking inverses for the diagram ( $\dagger$ ), we see that the diagram

$$\begin{array}{ccc} S^0 \wedge S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 \\ \sigma_{S^0,S^0}^{\text{Sets}_*, -1} \downarrow & (\ddagger) & \downarrow \sigma'_{S^0,S^0}^{-1} \\ S^0 \wedge S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 \end{array}$$

commutes as well. Now, let  $X, Y \in \text{Obj}(\text{Sets}_*)$ , let  $x \in X$ , let  $y \in Y$ , and consider the diagram

$$\begin{array}{ccccc} S^0 \wedge S^0 & \xrightarrow{\text{id}_{S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 & & \\ \sigma_{S^0,S^0}^{\text{Sets}_*, -1} \downarrow & \searrow [y] \wedge [x] & \downarrow \sigma'_{S^0,S^0}^{-1} & \searrow [y] \otimes_{\text{Sets}_*} [x] & \\ S^0 \wedge X & \xrightarrow{\text{id}_{\text{Sets}_*|Y,A}^{\otimes,-1}} & Y \wedge X & \xrightarrow{\text{id}_{\text{Sets}_*|Y,A}^{\otimes,-1}} & Y \otimes_{\text{Sets}_*} X \\ \sigma_{A,Y}^{\text{Sets}_*, -1} \downarrow & & \sigma_{A,Y}^{\text{Sets}_*, -1} \downarrow & & \downarrow \sigma'_{A,Y}^{-1} \\ S^0 \wedge S^0 & \xrightarrow{\text{id}_{S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 & & \\ \downarrow [x] \wedge [y] & & \downarrow [x] \otimes_{\text{Sets}_*} [y] & & \\ X \wedge Y & \xrightarrow{\text{id}_{\text{Sets}_*|A,Y}^{\otimes,-1}} & X \otimes_{\text{Sets}_*} Y & & \end{array}$$

which we partition into subdiagrams as follows:

$$\begin{array}{ccc} S^0 \wedge S^0 & \xrightarrow{\text{id}_{S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 \\ \sigma_{S^0,S^0}^{\text{Sets}_*, -1} \downarrow & \searrow [y] \wedge [x] & \searrow [y] \otimes_{\text{Sets}_*} [x] \\ S^0 \wedge X & \xrightarrow{\text{id}_{\text{Sets}_*|Y,X}^{\otimes,-1}} & Y \wedge X \\ \sigma_{X,Y}^{\text{Sets}_*, -1} \downarrow & (2) & \downarrow \sigma'_{X,Y}^{-1} \\ X \wedge Y & \xrightarrow{\text{id}_{\text{Sets}_*|X,Y}^{\otimes,-1}} & X \otimes_{\text{Sets}_*} Y \end{array} \quad \begin{array}{ccc} S^0 \wedge S^0 & \xrightarrow{\text{id}_{S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 \\ \sigma_{S^0,S^0}^{\text{Sets}_*, -1} \downarrow & (\ddagger) & \downarrow \sigma'_{S^0,S^0}^{-1} \\ S^0 \wedge S^0 & \xrightarrow{\text{id}_{S^0,S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 \\ \downarrow [x] \wedge [y] & \searrow [x] \otimes_{\text{Sets}_*} [y] & \downarrow \sigma'_{X,Y}^{-1} \\ X \wedge Y & \xrightarrow{\text{id}_{\text{Sets}_*|X,Y}^{\otimes,-1}} & X \otimes_{\text{Sets}_*} Y \end{array}$$

(3)

Since:

- Subdiagram (2) commutes by the naturality of  $\sigma^{\text{Sets}_*, -1}$ .
- Subdiagram (5) commutes by the naturality of  $\text{id}^{\otimes, -1}$ .
- Subdiagram (¶) commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $\sigma'^{-1}$ .
- Subdiagram (1) commutes by the naturality of  $\text{id}^{\otimes, -1}$ .

it follows that the diagram

$$\begin{array}{ccccc}
 S^0 \wedge S^0 & \xrightarrow{\quad /y \wedge /x \quad} & Y \wedge X & \xrightarrow{\text{id}_{\text{Sets}_*|Y,X}^\otimes} & Y \otimes_{\text{Sets}_*} X \\
 & \downarrow & \downarrow \sigma_{X,Y}^{\text{Sets}_*} & & \downarrow \sigma'_{X,Y} \\
 & & X \wedge Y & \xrightarrow{\text{id}_{\text{Sets}_*|X,Y}^\otimes} & X \otimes_{\text{Sets}_*} Y
 \end{array}$$

commutes. We then have

$$\begin{aligned}
 [\text{id}_{\text{Sets}_*|X,Y}^{\otimes, -1} \circ \sigma_{X,Y}^{\text{Sets}_*, -1}](y, x) &= [\text{id}_{\text{Sets}_*|X,Y}^{\otimes, -1} \circ \sigma_{X,Y}^{\text{Sets}_*, -1} \circ ([y] \wedge [x])](1, 1) \\
 &= [\sigma'^{-1}_{X,Y} \circ \text{id}_{\text{Sets}_*|Y,X}^{\otimes, -1} \circ ([y] \wedge [x])](1, 1) \\
 &= [\sigma'^{-1}_{X,Y} \circ \text{id}_{\text{Sets}_*|Y,X}^{\otimes, -1}](y, x)
 \end{aligned}$$

for each  $(y, x) \in Y \wedge X$ , and thus we have

$$\text{id}_{\text{Sets}_*|X,Y}^{\otimes, -1} \circ \sigma_{X,Y}^{\text{Sets}_*, -1} = \sigma'^{-1}_{X,Y} \circ \text{id}_{\text{Sets}_*|Y,X}^{\otimes, -1}.$$

Taking inverses then gives

$$\sigma_{X,Y}^{\text{Sets}_*} \circ \text{id}_{\text{Sets}_*|X,Y}^\otimes = \text{id}_{\text{Sets}_*|Y,X}^\otimes \circ \sigma'_{X,Y},$$

showing that the diagram

$$\begin{array}{ccc}
 A \otimes_{\text{Sets}_*} B & \xrightarrow{\text{id}_{\text{Sets}_*|A,B}^\otimes} & A \wedge B \\
 \downarrow \sigma'_{A,B} & & \downarrow \sigma_{A,B}^{\text{Sets}_*} \\
 B \otimes_{\text{Sets}_*} A & \xrightarrow{\text{id}_{\text{Sets}_*|B,A}^\otimes} & B \wedge A
 \end{array}$$

indeed commutes.

Monoidal Right Unity of the Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$

We have to show that the diagram

$$\begin{array}{ccc} X \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{S^0}^{\otimes_{\text{Sets}_*}|X,S^0}} & X \wedge S^0 \\ \text{id}_X \otimes_{\text{Sets}_*} \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\otimes} & \nearrow & \searrow \rho_X^{\text{Sets}_*} \\ X \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\rho'_X} & X \end{array}$$

commutes. To this end, we will first show that the diagram

$$\begin{array}{ccc} S^0 \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{S^0}^{\otimes_{\text{Sets}_*}|S^0,S^0}} & S^0 \wedge S^0 \\ \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_{S^0} & \nearrow & \searrow \rho_{S^0}^{\text{Sets}_*} \\ S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\rho'_{S^0}} & S^0, \end{array} \quad (\dagger)$$

corresponding to the case  $X = S^0$ , commutes. First, notice that we may write

$$\sigma'_{S^0, \mathbb{1}_{\text{Sets}_*}} : S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} \rightarrow \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0$$

as the composition

$$\begin{array}{c} S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} \xrightarrow{\text{id}_{S^0, \mathbb{1}_{\text{Sets}_*}}^{\otimes}} S^0 \wedge \mathbb{1}_{\text{Sets}_*} \\ \xrightarrow{\lambda_{\mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*}} \mathbb{1}_{\text{Sets}_*} \\ \xrightarrow{\rho_{\mathbb{1}_{\text{Sets}_*}, -1}^{\text{Sets}_*}} \mathbb{1}_{\text{Sets}_*} \wedge S^0 \\ \xrightarrow{\text{id}_{\mathbb{1}_{\text{Sets}_*}, S^0}^{\otimes, -1}} \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0. \end{array}$$

Indeed, we may write this composition as part of the diagram

$$\begin{array}{ccccc} S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\text{id}_{S^0, \mathbb{1}_{\text{Sets}_*}}^{\otimes}} & S^0 \wedge \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\lambda_{\mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*}} & \mathbb{1}_{\text{Sets}_*} \\ \downarrow \sigma'_{S^0, \mathbb{1}_{\text{Sets}_*}} & (1) & \downarrow \sigma_{S^0, \mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*} & (2) & \downarrow \rho_{\mathbb{1}_{\text{Sets}_*}, -1}^{\text{Sets}_*} \\ \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{\mathbb{1}_{\text{Sets}_*}, S^0}^{\otimes}} & \mathbb{1}_{\text{Sets}_*} \wedge S^0 & \xrightarrow{\text{id}_{\mathbb{1}_{\text{Sets}_*}, S^0}^{\otimes, -1}} & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0, \end{array}$$

which commutes since:

- Subdiagram (1) commutes by the braidedness of  $\text{id}^\otimes$ , as proved above.
- Subdiagram (2) commutes by ??.

Next, consider the diagram

$$\begin{array}{ccccccc}
 S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\text{id}_{S^0} \otimes_{\text{Sets}_*} \rho_{S^0}^{\text{Sets}_*, -1}} & S^0 \otimes_{\text{Sets}_*} (\mathbb{1}_{\text{Sets}_*} \wedge S^0) & \xrightarrow{\text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\text{Sets}_*|S^0}^{\otimes, -1}} & S^0 \otimes_{\text{Sets}_*} (\mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0) & \xrightarrow{\text{id}_{S^0} \otimes_{\text{Sets}_*} \lambda'_{S^0}} & S^0 \otimes_{\text{Sets}_*} S^0 \\
 \downarrow \text{id}_{\text{Sets}_*|S^0, \mathbb{1}_{\text{Sets}_*}}^\otimes & & \downarrow & & \downarrow & & \downarrow \text{id}_{\text{Sets}_*|S^0, S^0}^\otimes \\
 S^0 \wedge \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\text{id}_{S^0} \wedge \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*, -1}} & S^0 \wedge (\mathbb{1}_{\text{Sets}_*} \wedge S^0) & \xrightarrow{\text{id}_{S^0} \wedge \text{id}_{\text{Sets}_*|S^0}^{\otimes, -1}} & S^0 \wedge (\mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0) & \xrightarrow{\text{id}_{S^0} \wedge \lambda'_{S^0}} & S^0 \wedge S^0 \\
 \downarrow \lambda_{\mathbb{1}_{\text{Sets}_*}}^{\text{Sets}_*} & & \downarrow \lambda_{\mathbb{1}_{\text{Sets}_*} \wedge S^0}^{\text{Sets}_*} & & \downarrow \lambda_{\mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0}^{\text{Sets}_*} & & \downarrow \lambda_{S^0}^{\text{Sets}_*} = \rho_{S^0}^{\text{Sets}_*} \\
 \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\rho_{\mathbb{1}_{\text{Sets}_*}, -1}^{\text{Sets}_*}} & \mathbb{1}_{\text{Sets}_*} \wedge S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0}^{\otimes, -1}} & \mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\lambda'_{S^0}} & S^0,
 \end{array}$$

whose boundary diagram corresponds to the diagram  $(\dagger)$  above, since the composition in red is equal to  $\sigma'_{S^0, \mathbb{1}_{\text{Sets}_*}}$  as proved above, and then the composition in red composed with  $\lambda'_{S^0}$  is equal to  $\rho'_{S^0}$  by ???. In this diagram:

- Subdiagrams (1), (2), and (3) commute by the naturality of  $\text{id}_{\text{Sets}_*}^\otimes$ .
- Subdiagrams (4), (5), and (6) commute by the naturality of  $\lambda^{\text{Sets}_*}$ , where the equality  $\lambda_{S^0}^{\text{Sets}_*} = \rho_{S^0}^{\text{Sets}_*}$  comes from ??.

Since all subdiagrams commute, so does the boundary diagram, i.e. the diagram  $(\dagger)$  above. As a result, the diagram

$$\begin{array}{ccc}
 S^0 \wedge S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|S^0, S^0}^{\otimes, -1}} & S^0 \otimes_{\text{Sets}_*} S^0 \\
 \rho_{S^0}^{\text{Sets}_*, -1} \nearrow & & \searrow \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\mathbb{1}_{\text{Sets}_*}}^{\otimes, -1} \\
 S^0 & \xrightarrow{\rho'_{S^0}^{-1}} & S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*}.
 \end{array}$$

also commutes. Now, let  $X \in \text{Obj}(\text{Sets}_*)$ , let  $x \in X$ , and consider the diagram

$$\begin{array}{ccccc}
& S^0 \wedge S^0 & \xrightarrow{\text{id}_{S^0 \wedge S^0}^{\otimes,-1}} & S^0 \otimes_{\text{Sets}_*} S^0 & \\
\begin{matrix} \nearrow \rho_{S^0}^{\text{Sets}_*, -1} \\ \downarrow \end{matrix} & \downarrow & \text{iff} & \downarrow & \begin{matrix} \searrow id_{S^0 \wedge S^0}^{\otimes,-1} \wedge id_{S^0 \otimes_{\text{Sets}_*} S^0}^{\otimes,-1} \\ \downarrow \end{matrix} \\
S^0 & \xrightarrow{\rho'^{-1}_{S^0}} & S^0 \otimes_{\text{Sets}_*} 1_{\text{Sets}_*} & & \\
\downarrow id_{S^0} \wedge [x] & \downarrow (1) & \downarrow id_{S^0 \otimes_{\text{Sets}_*} S^0} [x] & & \downarrow id_{1_{\text{Sets}_*}} \wedge [x] \\
& X \wedge S^0 & \xrightarrow{\text{id}_{X \wedge S^0}^{\otimes,-1}} & X \otimes_{\text{Sets}_*} S^0 & \\
\downarrow \rho_X^{\text{Sets}_*, -1} & \downarrow (5) & \text{iff} & \downarrow id_X \wedge id_{X \otimes_{\text{Sets}_*} S^0}^{\otimes,-1} & \downarrow \\
X & \xrightarrow{\rho'^{-1}_X} & X \otimes_{\text{Sets}_*} 1_{\text{Sets}_*} & &
\end{array}$$

Since:

- Subdiagram (5) commutes by the naturality of  $\rho'^{-1}$ .
- Subdiagram ( $\dagger$ ) commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $\text{id}_{1_{\text{Sets}_*}}^{\otimes,-1}$ .
- Subdiagram (1) commutes by the naturality of  $\text{id}_{S^0}^{\otimes,-1}$ .
- Subdiagram (3) commutes by the naturality of  $\rho^{\text{Sets}_*, -1}$ .

it follows that the diagram

$$\begin{array}{ccc}
X \wedge S^0 & \xrightarrow{\text{id}_{X \wedge S^0}^{\otimes,-1}} & X \otimes_{\text{Sets}_*} S^0 \\
\rho_X^{\text{Sets}_*, -1} \nearrow & & \searrow id_X \otimes_{\text{Sets}_*} \text{id}_{1_{\text{Sets}_*}}^{\otimes,-1} \\
S^0 \xrightarrow{[x]} X & \xrightarrow{\rho'^{-1}_X} & X \otimes_{\text{Sets}_*} 1_{\text{Sets}_*}
\end{array}$$

Here's a step-by-step showcase of this argument: [\[Link\]](#). We then have

$$\rho'^{-1}_X(a) = [\rho'^{-1}_X \circ [x]](1)$$

$$\begin{aligned}
&= [(\text{id}_X \wedge \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1}) \circ \text{id}_{\text{Sets}_*|S^0, X}^{\otimes, -1} \circ \rho_X^{\text{Sets}_*, -1} \circ [x]](1) \\
&= [(\text{id}_X \wedge \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1}) \circ \text{id}_{\text{Sets}_*|S^0, X}^{\otimes, -1} \circ \rho_X^{\text{Sets}_*, -1}](a)
\end{aligned}$$

for each  $a \in X$ , and thus we have

$$\rho'_X = (\text{id}_X \wedge \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes, -1}) \circ \text{id}_{\text{Sets}_*|S^0, X}^{\otimes, -1} \circ \rho_X^{\text{Sets}_*, -1}.$$

Taking inverses then gives

$$\rho'_X = \rho_X^{\text{Sets}_*} \circ \text{id}_{\text{Sets}_*|S^0, X}^{\otimes} \circ (\text{id}_X \wedge \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes}),$$

showing that the diagram

$$\begin{array}{ccc}
X \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\text{id}_{\text{Sets}_*|X, S^0}^{\otimes}} & X \wedge S^0 \\
\text{id}_X \otimes_{\text{Sets}_*} \text{id}_{\mathbb{1}|\text{Sets}_*}^{\otimes} \swarrow & & \searrow \rho_X^{\text{Sets}_*} \\
X \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*} & \xrightarrow{\rho'_X} & X
\end{array}$$

indeed commutes.

Monoidality of the Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$

We have to show that the diagram

$$\begin{array}{ccc}
& (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z & \\
\text{id}_{\text{Sets}_*|X, Y}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_Z \swarrow & & \searrow \alpha'_{X, Y, Z} \\
(X \wedge Y) \otimes_{\text{Sets}_*} Z & & X \otimes_{\text{Sets}_*} (Y \otimes_{\text{Sets}_*} Z) \\
\downarrow \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes} & & \downarrow \text{id}_X \otimes_{\text{Sets}_*} \text{id}_{\text{Sets}_*|Y, Z}^{\otimes} \\
(X \wedge Y) \wedge Z & & X \otimes_{\text{Sets}_*} (Y \wedge Z) \\
\searrow \alpha_{X, Y, Z}^{\text{Sets}_*} & & \swarrow \text{id}_{\text{Sets}_*|X, Y \wedge Z}^{\otimes} \\
& X \wedge (Y \wedge Z) &
\end{array}$$

commutes. To this end, we will first prove that the diagram

$$\begin{array}{ccc}
 & (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & \\
 \text{id}_{\text{Sets}_*|S^0,S^0}^\otimes \otimes_{\text{Sets}_*} \text{id}_{S^0} & \swarrow & \searrow \alpha'_{S^0,S^0,S^0} \\
 (S^0 \wedge S^0) \otimes_{\text{Sets}_*} S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \otimes_{\text{Sets}_*} S^0) \\
 \downarrow \text{id}_{\text{Sets}_*|S^0 \wedge S^0,S^0}^\otimes & (\dagger) & \downarrow \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\text{Sets}_*|S^0,S^0}^\otimes \\
 (S^0 \wedge S^0) \wedge S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \wedge S^0) \\
 \downarrow \alpha_{S^0,S^0,S^0}^{\text{Sets}_*} & & \downarrow \text{id}_{\text{Sets}_*|S^0 \wedge S^0,S^0}^\otimes \\
 S^0 \wedge (S^0 \wedge S^0) & &
 \end{array}$$

commutes, and, to that end, we will first show that the diagram

$$\begin{array}{ccc}
 & (S^0 \otimes_{\text{Sets}_*} \mathbb{1}_{\text{Sets}_*}) \otimes_{\text{Sets}_*} S^0 & \\
 \text{id}_{\text{Sets}_*|S^0,\mathbb{1}_{\text{Sets}_*}}^\otimes \otimes_{\text{Sets}_*} \text{id}_{S^0} & \swarrow & \searrow \alpha'_{S^0,\mathbb{1}_{\text{Sets}_*},S^0} \\
 (S^0 \wedge \mathbb{1}_{\text{Sets}_*}) \otimes_{\text{Sets}_*} S^0 & & S^0 \otimes_{\text{Sets}_*} (\mathbb{1}_{\text{Sets}_*} \otimes_{\text{Sets}_*} S^0) \\
 \downarrow \text{id}_{\text{Sets}_*|S^0 \wedge \mathbb{1}_{\text{Sets}_*},S^0}^\otimes & (\ddagger) & \downarrow \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{\text{Sets}_*|\mathbb{1}_{\text{Sets}_*},S^0}^\otimes \\
 (S^0 \wedge \mathbb{1}_{\text{Sets}_*}) \wedge S^0 & & S^0 \otimes_{\text{Sets}_*} (\mathbb{1}_{\text{Sets}_*} \wedge S^0) \\
 \downarrow \alpha_{S^0,\mathbb{1}_{\text{Sets}_*},S^0}^{\text{Sets}_*} & & \downarrow \text{id}_{\text{Sets}_*|S^0 \wedge \mathbb{1}_{\text{Sets}_*},S^0}^\otimes \\
 S^0 \wedge (\mathbb{1}_{\text{Sets}_*} \wedge S^0) & &
 \end{array}$$

commutes. Indeed, consider the diagram

whose boundary diagram corresponds to diagram (‡) above. Since:

- Subdiagrams (1), (4), (5), (8), and (11) commute by the naturality of  $\text{id}_{\text{Sets}_*}^\otimes$ ;
- Subdiagram (2) commutes by the right monoidal unity of  $(\text{id}_{\text{Sets}_*}^\otimes, \text{id}_{\mathbb{1}|\text{Sets}_*}^\otimes)$ ;
- Subdiagram (3) commutes by the triangle identity for  $(\alpha', \lambda', \rho')$ ;
- Subdiagram (6) commutes by ??;
- Subdiagram (7) commutes by the naturality of  $\rho^{\text{Sets}_*}$ ;
- Subdiagram (9) commutes by ??;
- Subdiagram (10) commutes by ??;

it follows that the boundary diagram, i.e. diagram  $(\ddagger)$ , also commutes. Consider now the diagram

whose boundary corresponds to diagram  $(\dagger)$  above. Since:

- Subdiagrams (1), (3), (4), and (6) commute by the naturality of  $\text{id}_{Sets_*}^{⊗}$ ;
- Subdiagram  $(\ddagger)$  commutes, as proved above;
- Subdiagram (2) commutes by the naturality of  $\alpha'$ ;
- Subdiagram (5) commutes by the naturality of  $\alpha_{Sets_*}^{Sets_*}$ ;

it follows that the boundary diagram, i.e. diagram  $(\dagger)$ , also commutes.

Taking inverses on the diagram  $(\dagger)$ , we see that the diagram

$$\begin{array}{ccc}
 & S^0 \wedge (S^0 \wedge S^0) & \\
 \alpha_{S^0, S^0, S^0}^{\text{Sets}_*, -1} \swarrow & & \searrow \text{id}_{\text{Sets}_* | S^0, S^0 \wedge S^0}^{\otimes, -1} \\
 (S^0 \wedge S^0) \wedge S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \wedge S^0) \\
 \downarrow \text{id}_{\text{Sets}_* | S^0 \wedge S^0, S^0}^{\otimes, -1} & (\dagger) & \downarrow \text{id}_{S^0 \otimes_{\text{Sets}_*} S^0} \circ \text{id}_{\text{Sets}_* | S^0, S^0}^{\otimes, -1} \\
 (S^0 \wedge S^0) \otimes_{\text{Sets}_*} S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \otimes_{\text{Sets}_*} S^0) \\
 \downarrow \text{id}_{\text{Sets}_* | S^0, S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0} & & \downarrow \alpha'_{S^0, S^0, S^0}^{\otimes, -1} \\
 (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \otimes_{\text{Sets}_*} S^0)
 \end{array}$$

commutes as well. Now, let  $X, Y, Z \in \text{Obj}(\text{Sets}_*)$ , let  $x \in X$ , let  $y \in Y$ , let  $z \in Z$ , and consider the diagram

$$\begin{array}{ccccc}
 & S^0 \wedge (S^0 \wedge S^0) & & X \wedge (Y \wedge Z) & \\
 \alpha_{S^0, S^0, S^0}^{\text{Sets}_*, -1} \swarrow & \nearrow \text{id}_{S^0, S^0, S^0}^{\otimes, -1} & & \downarrow \text{id}_{\text{Sets}_* | X, Y \wedge Z}^{\otimes, -1} & \\
 (S^0 \wedge S^0) \wedge S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \wedge S^0) & & X \otimes_{\text{Sets}_*} (Y \wedge Z) \\
 \downarrow \text{id}_{S^0 \wedge S^0, S^0}^{\otimes, -1} & \downarrow \text{id}_{S^0 \otimes_{\text{Sets}_*} S^0} \circ \text{id}_{S^0, S^0}^{\otimes, -1} & \downarrow \text{id}_{S^0 \otimes_{\text{Sets}_*} S^0} \circ \text{id}_{S^0, S^0}^{\otimes, -1} & & \downarrow \text{id}_{X \otimes_{\text{Sets}_*} (Y \wedge Z)} \circ \text{id}_{S^0, S^0}^{\otimes, -1} \\
 (S^0 \wedge S^0) \otimes_{\text{Sets}_*} S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \otimes_{\text{Sets}_*} S^0) & & X \otimes_{\text{Sets}_*} (Y \wedge Z) \\
 \downarrow \text{id}_{S^0, S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0} & \downarrow \text{id}_{S^0, S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0} & \downarrow \text{id}_{S^0, S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0} & & \downarrow \text{id}_{X \otimes_{\text{Sets}_*} (Y \wedge Z)} \circ \text{id}_{S^0, S^0}^{\otimes, -1} \\
 (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & & (X \wedge Y) \wedge Z & & X \otimes_{\text{Sets}_*} (Y \wedge Z) \\
 \downarrow \text{id}_{(S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} & \downarrow \text{id}_{(X \wedge Y) \wedge Z}^{\otimes, -1} & \downarrow \text{id}_{(X \wedge Y) \otimes_{\text{Sets}_*} Z}^{\otimes, -1} & & \downarrow \text{id}_{X \otimes_{\text{Sets}_*} (Y \wedge Z)} \circ \text{id}_{S^0, S^0}^{\otimes, -1} \\
 (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & & (X \wedge Y) \otimes_{\text{Sets}_*} Z & & X \otimes_{\text{Sets}_*} (Y \otimes_{\text{Sets}_*} Z) \\
 \downarrow \text{id}_{S^0, S^0}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_{S^0} & \downarrow \text{id}_{(X \wedge Y) \otimes_{\text{Sets}_*} Z}^{\otimes, -1} & \downarrow \text{id}_{(X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z}^{\otimes, -1} & & \downarrow \text{id}_{X \otimes_{\text{Sets}_*} (Y \otimes_{\text{Sets}_*} Z)} \circ \text{id}_{S^0, S^0}^{\otimes, -1} \\
 (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & & (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z & & (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z
 \end{array}$$

which we partition into subdiagrams as follows:

$$\begin{array}{ccc}
 & S^0 \wedge (S^0 \wedge S^0) & \\
 \swarrow_{\alpha_{S^0 \wedge S^0 \wedge S^0}^{-1}} & & \searrow i_{S^0 \wedge S^0 \wedge S^0}^{-1} \\
 (S^0 \wedge S^0) \wedge S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \wedge S^0) \\
 \downarrow \text{id}_{S^0 \wedge S^0 \wedge S^0}^{\otimes -1} & (\dagger) & \downarrow \text{id}_{S^0} \otimes_{\text{Sets}_*} \text{id}_{S^0, S^0}^{\otimes -1} \\
 (S^0 \wedge S^0) \otimes_{\text{Sets}_*} S^0 & & S^0 \otimes_{\text{Sets}_*} (S^0 \otimes_{\text{Sets}_*} S^0) \\
 \downarrow i_{S^0 \wedge S^0 \otimes_{\text{Sets}_*} S^0}^{-1} & \searrow d_{S^0, S^0}^{-1} & \\
 (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & &
 \end{array}$$

$$\begin{array}{ccccc}
 & S^0 \wedge (S^0 \wedge S^0) & & X \wedge (Y \wedge Z) & \\
 \swarrow_{\alpha_{S^0 \wedge S^0 \wedge S^0}^{-1}} & & \searrow^{[x] \wedge ([y] \wedge [z])} & \swarrow_{\alpha_{X \wedge Y, Z}^{-1}} & \\
 (S^0 \wedge S^0) \wedge S^0 & & (X \wedge Y) \wedge Z & & \\
 \downarrow \text{id}_{S^0 \wedge S^0 \wedge S^0}^{\otimes -1} & \searrow^{([x] \wedge [y]) \wedge [z]} & \downarrow \text{id}_{\text{Sets}_*[X \wedge Y, Z]}^{\otimes -1} & & \\
 (S^0 \wedge S^0) \otimes_{\text{Sets}_*} S^0 & & (X \wedge Y) \otimes_{\text{Sets}_*} Z & & \\
 \downarrow i_{S^0 \wedge S^0 \otimes_{\text{Sets}_*} S^0}^{-1} & \searrow^{([x] \wedge [y]) \otimes_{\text{Sets}_*} [z]} & \downarrow i_{\text{Sets}_*[X \wedge Y, Z]}^{\otimes -1} & & \\
 (S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & & (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z & & \\
 \downarrow i_{S^0 \otimes_{\text{Sets}_*} S^0 \otimes_{\text{Sets}_*} S^0}^{-1} & \searrow^{([x] \otimes_{\text{Sets}_*} [y]) \otimes_{\text{Sets}_*} [z]} & & & \\
 (3) & & & &
 \end{array}$$

$$\begin{array}{ccccc}
S^0 \wedge (S^0 \wedge S^0) & \xrightarrow{\quad id_{S^0 \wedge S^0}^{\otimes, -1} \quad} & S^0 \otimes_{\text{Sets}_*} (S^0 \wedge S^0) & \xrightarrow{\quad id_{\text{Sets}_*}^{\otimes, -1} \quad} & X \wedge (Y \wedge Z) \\
& \searrow [x] \wedge ([y] \wedge [z]) & \nearrow i' & \nearrow (4) & \downarrow id_{\text{Sets}_* | X \wedge Y \wedge Z}^{\otimes, -1} \\
& & S^0 \otimes_{\text{Sets}_*} (S^0 \wedge S^0) & \xrightarrow{\quad id_{\text{Sets}_*}^{\otimes, -1} \quad} & X \otimes_{\text{Sets}_*} (Y \wedge Z) \\
& & \downarrow id_{S^0 \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} & & \downarrow id_X \otimes_{\text{Sets}_*} id_{\text{Sets}_* | Y, Z}^{\otimes, -1} \\
& & S^0 \otimes_{\text{Sets}_*} (S^0 \otimes_{\text{Sets}_*} S^0) & \xrightarrow{\quad id_{S^0 \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} \quad} & X \otimes_{\text{Sets}_*} (Y \otimes_{\text{Sets}_*} Z) \\
& \swarrow id_{S^0 \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} & \searrow [x] \otimes_{\text{Sets}_*} ([y] \otimes_{\text{Sets}_*} [z]) & \nearrow id_{(X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z}^{\otimes, -1} & \\
(S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0 & \xrightarrow{\quad id_{(S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} \quad} & & & (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z \\
& \searrow id_{(S^0 \otimes_{\text{Sets}_*} S^0) \otimes_{\text{Sets}_*} S^0}^{\otimes, -1} & & & \swarrow id_{(X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z}^{\otimes, -1} \\
& & (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z & &
\end{array}$$

(5)

Since:

- Subdiagram (1) commutes by the naturality of  $\alpha^{\text{Sets}_*, -1}$ .
- Subdiagram (2) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes, -1}$ .
- Subdiagram (3) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes, -1}$ .
- Subdiagram ( $\dagger$ ) commutes, as proved above.
- Subdiagram (4) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes, -1}$ .
- Subdiagram (5) commutes by the naturality of  $\text{id}_{\text{Sets}_*}^{\otimes, -1}$ .
- Subdiagram (6) commutes by the naturality of  $\alpha'^{-1}$ .

it follows that the diagram

$$\begin{array}{ccccc}
 & S^0 \wedge (S^0 \wedge S^0) & & & \\
 & \downarrow & & & \\
 & [x] \wedge ([y] \wedge [z]) & & & \\
 & \downarrow & & & \\
 X \wedge (Y \wedge Z) & & & & \\
 \alpha_{X,Y,Z}^{\text{Sets}_*, -1} \swarrow & & \searrow \text{id}_{\text{Sets}_*|X,Y \wedge Z}^{\otimes, -1} \\
 (X \wedge Y) \wedge Z & & & X \otimes_{\text{Sets}_*} (Y \wedge Z) & \\
 \downarrow \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes, -1} & & & \downarrow \text{id}_X \wedge \text{id}_{\text{Sets}_*|Y, Z}^{\otimes, -1} & \\
 (X \wedge Y) \otimes_{\text{Sets}_*} Z & & & X \otimes_{\text{Sets}_*} (Y \otimes_{\text{Sets}_*} Z) & \\
 \downarrow \text{id}_{\text{Sets}_*|X, Y}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_Z & & & \downarrow \alpha'_{X,Y,Z}^{\text{Sets}_*, -1} & \\
 (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z & & & &
 \end{array}$$

also commutes. We then have

$$\begin{aligned}
 & \left[ (\text{id}_{\text{Sets}_*|X, Y}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_Z) \circ \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes, -1} \right. \\
 & \quad \left. \circ \alpha_{X,Y,Z}^{\text{Sets}_*, -1} \right] (x, (y, z)) = \left[ (\text{id}_{\text{Sets}_*|X, Y}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_Z) \circ \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes, -1} \right. \\
 & \quad \left. \circ \alpha_{X,Y,Z}^{\text{Sets}_*, -1} \circ ([x] \wedge ([y] \wedge [z])) \right] (1, (1, 1)) \\
 & = \left[ \alpha'_{X,Y,Z}^{\text{Sets}_*, -1} \circ (\text{id}_X \wedge \text{id}_{\text{Sets}_*|Y, Z}^{\otimes, -1}) \right. \\
 & \quad \left. \circ \text{id}_{\text{Sets}_*|X, Y \wedge Z}^{\otimes, -1} \circ ([x] \wedge ([y] \wedge [z])) \right] (1, (1, 1)) \\
 & = [\alpha'_{X,Y,Z}^{\text{Sets}_*, -1} \circ (\text{id}_X \wedge \text{id}_{\text{Sets}_*|Y, Z}^{\otimes, -1}) \circ \text{id}_{\text{Sets}_*|X, Y \wedge Z}^{\otimes, -1}] (x, (y, z))
 \end{aligned}$$

for each  $(x, (y, z)) \in X \wedge (Y \wedge Z)$ , and thus we have

$$(\text{id}_{\text{Sets}_*|X, Y}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_Z) \circ \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes, -1} \circ \alpha_{X,Y,Z}^{\text{Sets}_*, -1} = \alpha'_{X,Y,Z}^{\text{Sets}_*, -1} \circ (\text{id}_X \wedge \text{id}_{\text{Sets}_*|Y, Z}^{\otimes, -1}) \circ \text{id}_{\text{Sets}_*|X, Y \wedge Z}^{\otimes, -1}.$$

Taking inverses then gives

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1} \circ \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes, -1} \circ (\text{id}_{\text{Sets}_*|X, Y}^{\otimes, -1} \otimes_{\text{Sets}_*} \text{id}_Z) = \text{id}_{\text{Sets}_*|X, Y \wedge Z}^{\otimes, -1} \circ (\text{id}_X \wedge \text{id}_{\text{Sets}_*|Y, Z}^{\otimes, -1}) \circ \alpha'_{X,Y,Z}^{\text{Sets}_*, -1},$$

showing that the diagram

$$\begin{array}{ccc}
 & (X \otimes_{\text{Sets}_*} Y) \otimes_{\text{Sets}_*} Z & \\
 \text{id}_{\text{Sets}_*|X,Y}^{\otimes} \otimes_{\text{Sets}_*} \text{id}_Z & \swarrow & \searrow d'_{X,Y,Z} \\
 (X \wedge Y) \otimes_{\text{Sets}_*} Z & & X \otimes_{\text{Sets}_*} (Y \otimes_{\text{Sets}_*} Z) \\
 \downarrow \text{id}_{\text{Sets}_*|X \wedge Y, Z}^{\otimes} & & \downarrow \text{id}_X \otimes_{\text{Sets}_*} \text{id}_{\text{Sets}_*|Y, Z}^{\otimes} \\
 (X \wedge Y) \wedge Z & & X \otimes_{\text{Sets}_*} (Y \wedge Z) \\
 \downarrow \alpha_{X,Y,Z}^{\text{Sets}_*} & & \downarrow \text{id}_{\text{Sets}_*|X, Y \wedge Z}^{\otimes} \\
 X \wedge (Y \wedge Z) & &
 \end{array}$$

indeed commutes.

Uniqueness of the Isomorphism  $\otimes_{\text{Sets}_*} \cong \wedge$

Let  $\phi, \psi: -_1 \otimes_{\text{Sets}_*} -_2 \Rightarrow -_1 \wedge -_2$  be natural isomorphisms. Since these isomorphisms are compatible with the unitors of  $\text{Sets}_*$  with respect to  $\wedge$  and  $\otimes$  (as shown above), we have

$$\begin{aligned}
 \lambda'_Y &= \lambda_Y^{\text{Sets}_*} \circ \phi_{S^0, Y} \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \otimes_{\text{Sets}} \text{id}_Y), \\
 \lambda'_Y &= \lambda_Y^{\text{Sets}_*} \circ \psi_{S^0, Y} \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes} \otimes_{\text{Sets}} \text{id}_Y).
 \end{aligned}$$

Postcomposing both sides with  $\lambda_Y^{\text{Sets}_*, -1}$  and then precomposing both sides with  $\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_Y$  gives

$$\begin{aligned}
 \lambda_Y^{\text{Sets}_*, -1} \circ \lambda'_Y \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_Y) &= \phi_{S^0, Y}, \\
 \lambda_Y^{\text{Sets}_*, -1} \circ \lambda'_Y \circ (\text{id}_{\mathbb{1}|\text{Sets}}^{\otimes, -1} \otimes_{\text{Sets}} \text{id}_Y) &= \psi_{S^0, Y},
 \end{aligned}$$

and thus we have

$$\phi_{S^0, Y} = \psi_{S^0, Y}$$

for each  $Y \in \text{Obj}(\text{Sets}_*)$ . Now, let  $x \in X$  and consider the naturality diagrams

$$\begin{array}{ccc}
 S^0 \wedge Y & \xrightarrow{[x] \wedge \text{id}_Y} & X \wedge Y \\
 \phi_{S^0, Y} \downarrow & & \downarrow \phi_{X, Y} \\
 S^0 \otimes_{\text{Sets}_*} Y & \xrightarrow{[x] \otimes_{\text{Sets}_*} \text{id}_Y} & X \otimes_{\text{Sets}_*} Y
 \end{array}
 \quad
 \begin{array}{ccc}
 S^0 \wedge Y & \xrightarrow{[x] \wedge \text{id}_Y} & X \wedge Y \\
 \psi_{S^0, Y} \downarrow & & \downarrow \psi_{X, Y} \\
 S^0 \otimes_{\text{Sets}_*} Y & \xrightarrow{[x] \otimes_{\text{Sets}_*} \text{id}_Y} & X \otimes_{\text{Sets}_*} Y
 \end{array}$$

for  $\phi$  and  $\psi$  with respect to the morphisms  $[x]$  and  $\text{id}_Y$ . Having shown that  $\phi_{S^0,Y} = \psi_{S^0,Y}$ , we have

$$\begin{aligned}\phi_{X,Y}(x,y) &= [\phi_{X,Y} \circ ([x] \wedge \text{id}_Y)](1,y) \\ &= [([x] \otimes_{\text{Sets}_*} \text{id}_Y) \circ \phi_{S^0,Y}](1,y) \\ &= [([x] \otimes_{\text{Sets}_*} \text{id}_Y) \circ \psi_{S^0,Y}](1,y) \\ &= [\psi_{X,Y} \circ ([x] \wedge \text{id}_Y)](1,y) \\ &= \psi_{X,Y}(x,y)\end{aligned}$$

for each  $(x,y) \in X \wedge Y$ . Therefore we have

$$\phi_{X,Y} = \psi_{X,Y}$$

for each  $X, Y \in \text{Obj}(\text{Sets}_*)$  and thus  $\phi = \psi$ , showing the isomorphism  $\otimes_{\text{Sets}_*} \cong \times$  to be unique. 

01W7

#### COROLLARY 7.5.10.1.3 ► A SECOND UNIVERSAL PROPERTY FOR $(\text{Sets}_*, \wedge, S^0)$

The symmetric monoidal structure on the category  $\text{Sets}_*$  of [Proposition 7.5.9.1.1](#) is uniquely determined by the following requirements:

01W8

1. *Two-Sided Preservation of Colimits.* The tensor product

$$\otimes_{\text{Sets}_*} : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of  $\text{Sets}_*$  preserves colimits separately in each variable.

01W9

2. *The Unit Object Is  $S^0$ .* We have  $1_{\text{Sets}_*} \cong S^0$ .

More precisely, the full subcategory of the category  $\mathcal{M}_{\mathbb{E}_\infty}(\text{Sets}_*)$  of ?? spanned by the symmetric monoidal categories  $(\text{Sets}_*, \otimes_{\text{Sets}_*}, 1_{\text{Sets}_*}, \lambda^{\text{Sets}_*}, \rho^{\text{Sets}_*}, \sigma^{\text{Sets}_*})$  satisfying [Items 1](#) and [2](#) is contractible.

#### PROOF 7.5.10.1.4 ► PROOF OF COROLLARY 7.5.10.1.3

Since  $\text{Sets}_*$  is locally presentable (??), it follows from ?? that [Corollary 7.5.10.1.3](#) is equivalent to the existence of an internal Hom as in [Item 1](#) of [Theorem 7.5.10.1.1](#). The result then follows from [Theorem 7.5.10.1.1](#). 

01WA

**COROLLARY 7.5.10.1.5 ► A THIRD UNIVERSAL PROPERTY OF THE SMASH PRODUCT OF POINTED SETS**

The symmetric monoidal structure on the category  $\text{Sets}_*$  is the unique symmetric monoidal structure on  $\text{Sets}_*$  such that the free pointed set functor

$$(-)^+ : \text{Sets} \rightarrow \text{Sets}_*$$

admits a symmetric monoidal structure, i.e. the full subcategory of the category  $\mathcal{M}_{\mathbb{E}_\infty}(\text{Sets}_*)$  of ?? spanned by the symmetric monoidal categories  $(\text{Sets}_*, \otimes_{\text{Sets}_*}, \mathbb{1}_{\text{Sets}_*}, \lambda^{\text{Sets}_*}, \rho^{\text{Sets}_*}, \sigma^{\text{Sets}_*})$  with respect to which  $(-)^+$  admits a symmetric monoidal structure is contractible.

**PROOF 7.5.10.1.6 ► PROOF OF COROLLARY 7.5.10.1.5**

Let  $(\otimes_{\text{Sets}_*}, \mathbb{1}_{\text{Sets}_*}, \lambda^{\text{Sets}_*}, \rho^{\text{Sets}_*}, \sigma^{\text{Sets}_*})$  be a symmetric monoidal structure on  $\text{Sets}_*$  such that  $(-)^+$  admits a symmetric monoidal structure with respect to  $\otimes_{\text{Sets}_*}$  and  $\wedge$ . We have isomorphisms

$$\begin{aligned} X \otimes_{\text{Sets}_*} Y &\cong (X^-)^+ \otimes_{\text{Sets}_*} (Y^-)^+ \\ &\cong (X^- \times Y^-)^+ \\ &\cong (X^-)^+ \wedge (Y^-)^+ \\ &\cong X \wedge Y, \end{aligned}$$

all natural in  $X$  and  $Y$ . Now, since  $\wedge$  preserves colimits in both variables and  $\otimes_{\text{Sets}_*} \cong \wedge$ , it follows that  $\otimes_{\text{Sets}_*}$  also preserves colimits in both variables, so the result then follows from [Corollary 7.5.10.1.3](#). 

00H6 **7.5.11 Monoids With Respect to the Smash Product of Pointed Sets**

00H7

**PROPOSITION 7.5.11.1.1 ► MONOIDS WITH RESPECT TO  $\wedge$** 

The category of monoids on  $(\text{Sets}_*, \wedge, S^0)$  is isomorphic to the category of monoids with zero and morphisms between them.

**PROOF 7.5.11.1.2 ► PROOF OF PROPOSITION 7.5.11.1.1**

See ??, in particular ??, ??, and ??.



**00H8 7.5.12 Comonoids With Respect to the Smash Product of Pointed Sets**

00H9

**PROPOSITION 7.5.12.1.1 ► COMONOIDS WITH RESPECT TO  $\wedge$**

The symmetric monoidal functor

$$((-)^+, (-)^{+,\times}, (-)_{\mathbb{1}}^{+,\times}): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}_*, \wedge, S^0),$$

of [Item 4 of Proposition 6.4.1.1.2](#) lifts to an equivalence of categories

$$\begin{aligned} \text{CoMon}(\text{Sets}_*, \wedge, S^0) &\xrightarrow{\text{eq.}} \text{CoMon}(\text{Sets}, \times, \text{pt}) \\ &\cong \text{Sets}. \end{aligned}$$

**PROOF 7.5.12.1.2 ► PROOF OF PROPOSITION 7.5.12.1.1**

We follow [[PS19](#), Lemma 2.4].

**Faithfulness**

Given morphisms  $f, g: X \rightarrow Y$ , if  $f^+ = g^+$ , then we have

$$\begin{aligned} f(x) &\stackrel{\text{def}}{=} f^+(x) \\ &= g^+(x) \\ &\stackrel{\text{def}}{=} g(x) \end{aligned}$$

for each  $x \in X^+$ , and thus  $f = g$ , showing  $(-)^+$  to be faithful.

**Fullness**

Let  $f: X^+ \rightarrow Y^+$  be a morphism of comonoids in  $\text{Sets}_*$ . By counitality, the diagram

$$\begin{array}{ccc} X^+ & \xrightarrow{f} & Y^+ \\ \epsilon_X^+ \searrow & & \swarrow \epsilon_Y^+ \\ & S^0 & \end{array}$$

commutes. If  $f(x) = \star_Y$  for  $x \neq \star_X$ , the commutativity of this diagram then gives

$$\begin{aligned} 1 &= \epsilon_X^+(x) \\ &= \epsilon_Y^+(f(x)) \\ &= \epsilon_Y^+(\star_Y) \\ &= 0, \end{aligned}$$

which is a contradiction. Thus  $f$  is an active morphism of pointed sets, so there exists a map  $f^-$  such that  $(f^-)^+ = f$  (Item 1 of Proposition 6.4.2.1.2).

### Essential Surjectivity

Let  $(X, \Delta_X, \epsilon_X)$  be a comonoid in  $\text{Sets}_*$ . We claim that

$$\Delta_X(x) = x \wedge x$$

for each  $x \in X$  with  $x \neq \star_X$ . Indeed:

- Suppose that  $x \neq \star_X$  and write  $\Delta_X(x) = x_1 \wedge x_2$ .
- Since  $\text{id}_X \wedge \epsilon_X$  is pointed, we have

$$[\text{id}_X \wedge \epsilon_X](x_1 \wedge x_2) = \star_{X \wedge S^0}.$$

- The counitality condition for  $\Delta_X$ , corresponding to the commutativity of the diagram

$$\begin{array}{ccc} X & \xrightarrow{\Delta_X} & X \wedge X \\ & \searrow \rho_X^{\text{Sets}_*, -1} & \downarrow \text{id}_X \wedge \epsilon_X \\ & & X \wedge S^0 \end{array}$$

gives

$$\begin{aligned} x \wedge 1 &= \rho_X^{\text{Sets}_*, -1}(x) \\ &= [\text{id}_X \wedge \epsilon_X \circ \Delta_X](x) \\ &= [\text{id}_X \wedge \epsilon_X](\Delta_X(x)) \\ &= [\text{id}_X \wedge \epsilon_X](x_1 \wedge x_2) \\ &= \star_{X \wedge S^0}, \end{aligned}$$

which is a contradiction. Thus  $x_1 \neq \star_X$ .

- Similarly, if  $x \neq \star_X$ , then  $x_2 \neq \star_X$ .
- Next, we claim that  $\epsilon_X(x_2) = 1$ , as otherwise we would have

$$\begin{aligned} \star_{X \wedge S^0} &= x_1 \wedge 0 \\ &= [\text{id}_X \wedge \epsilon_X](x_1 \wedge x_2) \\ &= [\text{id}_X \wedge \epsilon_X](\Delta_X(x)) \end{aligned}$$

$$\begin{aligned}
&= [\text{id}_X \wedge \epsilon_X \circ \Delta_X](x) \\
&= \rho_X^{\text{Sets}_*, -1}(x) \\
&= x \wedge 1,
\end{aligned}$$

a contradiction. Thus  $\epsilon_X(x_2) = 1$ .

- Similarly, if  $x \neq \star_X$ , then  $\epsilon_X(x_1) = 1$ .
- Now, since  $\Delta_X$  is counital, we have

$$\begin{aligned}
x \wedge 1 &= \rho_X^{\text{Sets}_*, -1}(x) \\
&= [\text{id}_X \wedge \epsilon_X \circ \Delta_X](x) \\
&= [\text{id}_X \wedge \epsilon_X](\Delta_X(x)) \\
&= [\text{id}_X \wedge \epsilon_X](x_1 \wedge x_2) \\
&= x_1 \wedge 1,
\end{aligned}$$

so  $x = x_1$ .

- Similarly,  $x = x_2$ , and we are done.

Next, notice that  $X \cong \epsilon_X^{-1}(0) \coprod \epsilon_X^{-1}(1)$ , and let  $x \in \epsilon_X^{-1}(0)$ . We then have

$$\begin{aligned}
[(\text{id}_X \wedge \epsilon_X) \circ \Delta_X](x) &= [\text{id}_X \wedge \epsilon_X](x \wedge x) \\
&= x \wedge 0 \\
&= \star_{X \wedge S^0}.
\end{aligned}$$

The counity condition for  $\Delta_X$  then gives  $x = \star_X$ , so  $\epsilon_X^{-1}(0) = \{\star_X\}$ . Thus we have  $(\epsilon_X^{-1}(1))^+ \cong X$ , and this isomorphism is compatible with the comonoid structures when equipping  $\epsilon_X^{-1}(1)$  with its unique comonoid structure. This shows that  $(-)^+$  is essentially surjective.

### Equivalence

Since  $(-)^+$  is fully faithful and essentially surjective, it is an equivalence by Item 1b of Item 1 of Proposition 11.6.7.1.2. 

## 00HA 7.6 Miscellany

### 00HB 7.6.1 The Smash Product of a Family of Pointed Sets

Let  $\{(X_i, x_0^i)\}_{i \in I}$  be a family of pointed sets.

00HC

**DEFINITION 7.6.1.1.1 ► THE SMASH PRODUCT OF A FAMILY OF POINTED SETS**

The **smash product of the family**  $\{(X_i, x_0^i)\}_{i \in I}$  is the pointed set  $\bigwedge_{i \in I} X_i$  consisting of:

- *The Underlying Set.* The set  $\bigwedge_{i \in I} X_i$  defined by

$$\bigwedge_{i \in I} X_i \stackrel{\text{def}}{=} \left( \prod_{i \in I} X_i \right) / \sim,$$

where  $\sim$  is the equivalence relation on  $\prod_{i \in I} X_i$  obtained by declaring

$$(x_i)_{i \in I} \sim (y_i)_{i \in I}$$

if there exist  $i_0 \in I$  such that  $x_{i_0} = x_0$  and  $y_{i_0} = y_0$ , for each  $(x_i)_{i \in I}, (y_i)_{i \in I} \in \prod_{i \in I} X_i$ .

- *The Basepoint.* The element  $[(x_0)_{i \in I}]$  of  $\bigwedge_{i \in I} X_i$ .

# Appendices

## 7.A Other Chapters

<b>Preliminaries</b>	10. Conditions on Relations
1. Introduction	
2. A Guide to the Literature	
<b>Sets</b>	<b>Categories</b>
3. Sets	11. Categories
4. Constructions With Sets	12. Presheaves and the Yoneda Lemma
5. Monoidal Structures on the Category of Sets	<b>Monoidal Categories</b>
6. Pointed Sets	13. Constructions With Monoidal Categories
7. Tensor Products of Pointed Sets	<b>Bicategories</b>
<b>Relations</b>	14. Types of Morphisms in Bicategories
8. Relations	<b>Extra Part</b>
9. Constructions With Relations	15. Notes

## **Part III**

# **Relations**

# Chapter 8

## Relations

**02H3** This chapter contains some material about relations. Notably, we discuss and explore:

- 028A** 1. The definition of relations ([Section 8.1.1](#)).
- 028B** 2. How relations may be viewed as decategorification of profunctors ([Section 8.1.2](#)).
- 02KS** 3. The various kinds of categories that relations form, namely:
  - 02KT** (a) A category ([Section 8.3.2](#)).
  - 02KU** (b) A monoidal category ([Section 8.3.3](#)).
  - 02KV** (c) A 2-category ([Section 8.3.4](#)).
  - 02KW** (d) A double category ([Section 8.3.5](#)).
- 028H** 4. The various categorical properties of the 2-category of relations, including:
  - 028J** (a) The self-duality of **Rel** and **Rel** ([Proposition 8.5.1.1.1](#)).
  - 028K** (b) Identifications of equivalences and isomorphisms in **Rel** with bijections ([Proposition 8.5.2.1.3](#)).
  - 028L** (c) Identifications of adjunctions in **Rel** with functions ([Proposition 8.5.3.1.1](#)).
  - 028M** (d) Identifications of monads in **Rel** with preorders (??).
  - 028N** (e) Identifications of comonads in **Rel** with subsets (??).
  - 028P** (f) A description of the monoids and comonoids in **Rel** with respect to the Cartesian product ([Remark 8.5.9.1.1](#)).
  - 028Q** (g) Characterisations of monomorphisms in **Rel** ([Old Tag 15.2.1.1.11](#)).
  - 028R** (h) Characterisations of 2-categorical notions of monomorphisms in **Rel** ([Old Tag 15.2.1.1.13](#)).
  - 028S** (i) Characterisations of epimorphisms in **Rel** ([Old Tag 15.2.1.1.23](#)).

- 028T (j) Characterisations of 2-categorical notions of epimorphisms in **Rel** ([Old Tag 15.2.1.1.26](#)).
- 028U (k) The partial co/completeness of **Rel** ([Proposition 8.5.12.1.1](#)).
- 028V (l) The existence or non-existence of Kan extensions and Kan lifts in **Rel** ([Old Tag 15.2.1.1.7](#)).
- 028W (m) The closedness of **Rel** ([Proposition 8.5.17.1.1](#)).
- 028X (n) The identification of **Rel** with the category of free algebras of the powerset monad on Sets ([Proposition 8.5.18.1.1](#)).

**02KX** 5. The adjoint pairs

$$\begin{aligned} R_! \dashv R_{-1} : \mathcal{P}(A) &\rightleftarrows \mathcal{P}(B), \\ R^{-1} \dashv R_* : \mathcal{P}(B) &\rightleftarrows \mathcal{P}(A) \end{aligned}$$

of functors (morphisms of posets) between  $\mathcal{P}(A)$  and  $\mathcal{P}(B)$  induced by a relation  $R: A \rightarrow B$ , as well as the properties of  $R_!$ ,  $R_{-1}$ ,  $R^{-1}$ , and  $R_*$  ([Section 8.7](#)).

Of particular note are the following points:

- 02KY (a) These two pairs of adjoint functors are the counterpart for relations of the adjoint triple  $f_! \dashv f^{-1} \dashv f_*$  induced by a function  $f: A \rightarrow B$  studied in [Section 4.6](#).
- 02KZ (b) We have  $R_{-1} = R^{-1}$  iff  $R$  is total and functional ([Item 8 of Proposition 8.7.2.1.4](#)).
- 02L0 (c) As a consequence of the previous item, when  $R$  comes from a function  $f$ , the pair of adjunctions

$$R_! \dashv R_{-1} = R^{-1} \dashv R_*$$

reduces to the triple adjunction

$$f_! \dashv f^{-1} \dashv f_*$$

from [Section 4.6](#).

- 02L1 (d) The pairs  $R_! \dashv R_{-1}$  and  $R^{-1} \dashv R_*$  turn out to be rather important later on, as they appear in the definition and study of continuous, open, and closed relations between topological spaces ([??](#)).
- 028Y** 6. A description of two notions of “skew composition” on  $\text{Rel}(A, B)$ , giving rise to left and right skew monoidal structures analogous to the left skew monoidal structure on  $\text{Fun}(C, \mathcal{D})$  appearing in the definition of a relative monad ([Sections 8.8 and 8.9](#)).

This chapter is under revision. TODO:

1. Replicate [Section 8.5](#) for apartness composition
2. Revise [Section 8.7](#)
3. Add subsection “A Six Functor Formalism for Sets, Part 2”, now with relations, building upon [Section 8.7](#).
4. Replicate [Section 8.7](#) for apartness composition
5. Revise sections on skew monoidal structures on  $\mathbf{Rel}(A, B)$
6. Replicate the sections on skew monoidal structures on  $\mathbf{Rel}(A, B)$  for apartness composition.
7. Explore relative co/monads in  $\mathbf{Rel}$ , defined to be co/monoids in  $\mathbf{Rel}(A, B)$  with its left/right skew monoidal structures of [Sections 8.8](#) and [8.9](#)
8. functional total relations defined with “satisfying the following equivalent conditions.”

## Contents

---

<b>8.1 Relations .....</b>	<b>526</b>
8.1.1 Foundations .....	526
8.1.2 Relations as Decategorifications of Profunctors ...	530
8.1.3 Composition of Relations.....	532
8.1.4 Apartness Composition of Relations.....	540
8.1.5 The Converse of a Relation .....	547
<b>8.2 Examples of Relations.....</b>	<b>551</b>
8.2.1 Elementary Examples of Relations.....	551
8.2.2 The Graph of a Function .....	554
8.2.3 The Inverse of a Function .....	558
8.2.4 Representable Relations.....	560
<b>8.3 Categories of Relations .....</b>	<b>561</b>
8.3.1 The Category of Relations Between Two Sets ....	561
8.3.2 The Category of Relations .....	561
8.3.3 The Closed Symmetric Monoidal Category of Relations .....	562
8.3.4 The 2-Category of Relations.....	568
8.3.5 The Double Category of Relations .....	569
<b>8.4 Categories of Relations With Apartness Composition .</b>	<b>577</b>
8.4.1 The Category of Relations With Apartness Composition .....	577
8.4.2 The 2-Category of Relations With Apartness Composition .....	578
8.4.3 The Linear Bicategory of Relations .....	580

---

8.4.4 Other Categorical Structures With Apartness Composition.....	580
<b>8.5 Properties of the 2-Category of Relations.....</b>	<b>581</b>
8.5.1 Self-Duality.....	581
8.5.2 Isomorphisms and Equivalences.....	582
8.5.3 Internal Adjunctions.....	584
8.5.4 Internal Monads.....	586
8.5.5 Internal Comonads .....	588
8.5.6 Modules Over Internal Monads.....	590
8.5.7 Comodules Over Internal Comonads.....	591
8.5.8 Eilenberg–Moore and Kleisli Objects.....	593
8.5.9 Co/Monoids .....	593
8.5.10 Monomorphisms and 2-Categorical Monomorphisms .....	594
8.5.11 Epimorphisms and 2-Categorical Epimorphisms..	612
8.5.12 Co/Limits .....	632
8.5.13 Internal Left Kan Extensions .....	632
8.5.14 Internal Left Kan Lifts.....	635
8.5.15 Internal Right Kan Extensions .....	636
8.5.16 Internal Right Kan Lifts.....	642
8.5.17 Closedness .....	646
8.5.18 Rel as a Category of Free Algebras.....	647
<b>8.6 Properties of the 2-Category of Relations With Apartness Composition.....</b>	<b>648</b>
8.6.1 Self-Duality.....	648
8.6.2 Isomorphisms and Equivalences.....	649
8.6.3 Internal Adjunctions.....	651
8.6.4 Internal Monads.....	651
8.6.5 Internal Comonads .....	652
8.6.6 Modules Over Internal Monads.....	654
8.6.7 Comodules Over Internal Comonads.....	654
8.6.8 Eilenberg–Moore and Kleisli Objects.....	654
8.6.9 Monomorphisms .....	654
8.6.10 2-Categorical Monomorphisms .....	654
8.6.11 Epimorphisms.....	654
8.6.12 2-Categorical Epimorphisms .....	654
8.6.13 Co/Limits .....	654
8.6.14 Internal Left Kan Extensions .....	654
8.6.15 Internal Left Kan Lifts.....	654
8.6.16 Internal Right Kan Extensions .....	654
8.6.17 Internal Right Kan Lifts.....	654

8.6.18	Coclosedness.....	654
<b>8.7</b>	<b>The Adjoint Pairs <math>R_! \dashv R_{-1}</math> and <math>R^{-1} \dashv R_*</math>.....</b>	<b>654</b>
8.7.1	Direct Images .....	654
8.7.2	Coinverse Images .....	660
8.7.3	Inverse Images.....	666
8.7.4	Codirect Images.....	671
8.7.5	Functoriality of Powersets .....	678
<b>8.8</b>	<b>The Left Skew Monoidal Structure on <math>\text{Rel}(A, B)</math> .....</b>	<b>680</b>
8.8.1	The Left Skew Monoidal Product .....	680
8.8.2	The Left Skew Monoidal Unit.....	681
8.8.3	The Left Skew Associators.....	681
8.8.4	The Left Skew Left Unitors .....	682
8.8.5	The Left Skew Right Unitors .....	683
8.8.6	The Left Skew Monoidal Structure on $\text{Rel}(A, B)$ ..	684
<b>8.9</b>	<b>The Right Skew Monoidal Structure on <math>\text{Rel}(A, B)</math> .....</b>	<b>685</b>
8.9.1	The Right Skew Monoidal Product .....	685
8.9.2	The Right Skew Monoidal Unit.....	686
8.9.3	The Right Skew Associators.....	686
8.9.4	The Right Skew Left Unitors .....	687
8.9.5	The Right Skew Right Unitors .....	688
8.9.6	The Right Skew Monoidal Structure on $\text{Rel}(A, B)$	
689		
<b>8.A</b>	<b>Other Chapters.....</b>	<b>690</b>

## 00HE 8.1 Relations

### 00HF 8.1.1 Foundations

Let  $A$  and  $B$  be sets.

01D0

#### DEFINITION 8.1.1.1 ► RELATIONS

A **relation**  $R: A \rightarrow B$  from  $A$  to  $B^{1,2}$  is equivalently:

00HR

1. A subset  $R$  of  $A \times B$ .

00HS

2. A function from  $A \times B$  to  $\{\text{true}, \text{false}\}$ .

00HT

3. A function from  $A$  to  $\mathcal{P}(B)$ .

00HU

4. A function from  $B$  to  $\mathcal{P}(A)$ .

00HV

5. A cocontinuous morphism of posets from  $(\mathcal{P}(A), \subset)$  to  $(\mathcal{P}(B), \subset)$ .

02C6

6. A continuous morphism of posets from  $(\mathcal{P}(B), \supset)$  to  $(\mathcal{P}(A), \supset)$ .

<sup>1</sup>Further Terminology: Also called a **multivalued function from A to B**.

<sup>2</sup>Further Terminology: When  $A = B$ , we also call  $R \subset A \times A$  a **relation on A**.

#### PROOF 8.1.1.1.2 ► PROOF OF THE EQUIVALENCES IN DEFINITION 8.1.1.1

(We will prove that **Items 1** to **6** are indeed equivalent in a bit.)



01QZ

#### REMARK 8.1.1.3 ► UNWINDING ITEM 1, I

We may think of a relation  $R: A \rightarrow B$  as a function from  $A$  to  $B$  that is *multivalued*, assigning to each element  $a$  in  $A$  a set  $R(a)$  of elements of  $B$ , thought of as the *set of values of R at a*.

Note that this includes also the possibility of  $R$  having no value at all on a given  $a \in A$  when  $R(a) = \emptyset$ .

01R0

#### REMARK 8.1.1.4 ► UNWINDING ITEM 2, II

Another way of stating the equivalence between **Items 1** to **5** of **Definition 8.1.1.1** is by saying that we have bijections of sets

$$\begin{aligned} \{\text{relations from } A \text{ to } B\} &\cong \mathcal{P}(A \times B) \\ &\cong \text{Sets}(A \times B, \{\text{true, false}\}) \\ &\cong \text{Sets}(A, \mathcal{P}(B)) \\ &\cong \text{Sets}(B, \mathcal{P}(A)) \\ &\cong \text{Pos}^{\mathcal{D}}(\mathcal{P}(A), \mathcal{P}(B)) \\ &\cong \text{Pos}^{\mathcal{C}}(\mathcal{P}(B), \mathcal{P}(A)) \end{aligned}$$

natural in  $A, B \in \text{Obj}(\text{Sets})$ , where  $\mathcal{P}(A)$  and  $\mathcal{P}(B)$  are endowed with the poset structure given by inclusion.

#### PROOF 8.1.1.1.5 ► PROOF OF THE EQUIVALENCES IN DEFINITION 8.1.1.1

We claim that **Items 1** to **5** are indeed equivalent:

- **Item 1**  $\iff$  **Item 2**: This is a special case of **Items 2** and **3** of **Proposition 4.5.1.1.4**.
- **Item 2**  $\iff$  **Item 3**: This follows from the bijections

$$\text{Sets}(A \times B, \{\text{true, false}\}) \cong \text{Sets}(A, \text{Sets}(B, \{\text{true, false}\}))$$

$$\cong \text{Sets}(A, \mathcal{P}(B)),$$

where the last bijection is from [Items 2 and 3](#) of [Proposition 4.5.1.1.4](#).

- [Item 2  \$\iff\$  Item 4](#): This follows from the bijections

$$\begin{aligned} \text{Sets}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Sets}(B, \text{Sets}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Sets}(B, \mathcal{P}(A)), \end{aligned}$$

where again the last bijection is from [Items 2 and 3](#) of [Proposition 4.5.1.1.4](#).

- [Item 2  \$\iff\$  Item 5](#): This follows from the universal property of the powerset  $\mathcal{P}(X)$  of a set  $X$  as the free cocompletion of  $X$  via the characteristic embedding

$$\chi_X: X \hookrightarrow \mathcal{P}(X)$$

of  $X$  into  $\mathcal{P}(X)$ , as in [Proposition 4.4.5.1.1](#). In particular, the bijection

$$\text{Sets}(A, \mathcal{P}(B)) \cong \text{Pos}^{\mathcal{D}}(\mathcal{P}(A), \mathcal{P}(B))$$

is given by extending each  $f: A \rightarrow \mathcal{P}(B)$  in  $\text{Sets}(A, \mathcal{P}(B))$  from  $A$  to all of  $\mathcal{P}(A)$  by taking its left Kan extension along  $\chi_X$ , recovering the direct image function  $f_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$  of  $f$  of [Definition 4.6.1.1.1](#).

- [Item 5  \$\iff\$  Item 6](#): Omitted.

This finishes the proof. 

00HH

#### NOTATION 8.1.1.6 ► FURTHER NOTATION FOR RELATIONS

Let  $A$  and  $B$  be sets and let  $R: A \rightarrow B$  be a relation from  $A$  to  $B$ .

00HM

1. We write  $\text{Rel}(A, B)$  for the set of relations from  $A$  to  $B$ .

00HN

2. We write  $\text{Rel}(A, B)$  for the sub-poset of  $(\mathcal{P}(A \times B), \subset)$  spanned by the relations from  $A$  to  $B$ .

00HJ

3. Given  $a \in A$  and  $b \in B$ , we write  $a \sim_R b$  to mean  $(a, b) \in R$ .

00HK

4. When viewing  $R$  as a function

$$R: A \times B \rightarrow \{\text{t, f}\},$$

we write  $R_a^b$  for the value of  $R$  at  $(a, b)$ .<sup>1</sup>

<sup>1</sup>The choice to write  $R_a^b$  in place of  $R_b^a$  is to keep the notation consistent with the notation we will later employ for profunctors in ??.

00HW

**PROPOSITION 8.1.1.7 ► PROPERTIES OF RELATIONS**

Let  $A$  and  $B$  be sets and let  $R, S: A \rightarrow B$  be relations.

00HX

1. *End Formula for the Set of Inclusions of Relations.* We have

$$\text{Hom}_{\text{Rel}(A,B)}(R, S) \cong \int_{a \in A} \int_{b \in B} \text{Hom}_{\{\text{t,f}\}}(R_a^b, S_a^b).$$

**PROOF 8.1.1.8 ► PROOF OF PROPOSITION 8.1.1.7****Item 1: End Formula for the Set of Inclusions of Relations**

Unwinding the expression inside the end on the right hand side, we have

$$\int_{a \in A} \int_{b \in B} \text{Hom}_{\{\text{t,f}\}}(R_a^b, S_a^b) \cong \begin{cases} \text{pt} & \text{if, for each } a \in A \text{ and each } b \in B, \\ & \text{we have } \text{Hom}_{\{\text{t,f}\}}(R_a^b, S_a^b) \cong \text{pt} \\ \emptyset & \text{otherwise.} \end{cases}$$

Since we have  $\text{Hom}_{\{\text{t,f}\}}(R_a^b, S_a^b) = \{\text{true}\} \cong \text{pt}$  exactly when  $R_a^b = \text{false}$  or  $R_a^b = S_a^b = \text{true}$ , we get

$$\int_{a \in A} \int_{b \in B} \text{Hom}_{\{\text{t,f}\}}(R_a^b, S_a^b) \cong \begin{cases} \text{pt} & \text{if, for each } a \in A \text{ and each } b \in B, \\ & \text{if } a \sim_R b, \text{ then } a \sim_S b, \\ \emptyset & \text{otherwise.} \end{cases}$$

On the left hand-side, we have

$$\text{Hom}_{\text{Rel}(A,B)}(R, S) \cong \begin{cases} \text{pt} & \text{if } R \subset S, \\ \emptyset & \text{otherwise.} \end{cases}$$

Since  $(a \sim_R b \implies a \sim_S b)$  iff  $R \subset S$ , the two sets above are isomorphic. This finishes the proof. 

**00HY 8.1.2 Relations as Decategorifications of Profunctors**

00HZ

**REMARK 8.1.2.1.1 ► RELATIONS AS DECATERGORIFICATIONS OF PROFUNCTORS I**

The notion of a relation is a decategorification of that of a profunctor:

028Z

1. A profunctor from a category  $C$  to a category  $\mathcal{D}$  is a functor

$$\mathbf{p}: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}.$$

0290

2. A relation on sets  $A$  and  $B$  is a function

$$R: A \times B \rightarrow \{\text{true, false}\}.$$

Here we notice that:

- The opposite  $X^{\text{op}}$  of a set  $X$  is itself, as  $(-)^{\text{op}}: \text{Cats} \rightarrow \text{Cats}$  restricts to the identity endofunctor on Sets.
- The values that profunctors and relations take are analogous:
  - A category is enriched over the category

$$\text{Sets} \stackrel{\text{def}}{=} \text{Cats}_0$$

of sets, with profunctors taking values on it.

- A set is enriched over the set

$$\{\text{true, false}\} \stackrel{\text{def}}{=} \text{Cats}_{-1}$$

of classical truth values, with relations taking values on it.

00J0

**REMARK 8.1.2.1.2 ► RELATIONS AS DECATERGORIFICATIONS OF PROFUNCTORS II**

Extending Remark 8.1.2.1.1, the equivalent definitions of relations in Definition 8.1.1.1.1 are also related to the corresponding ones for profunctors (??), which state that a profunctor  $\mathbf{p}: C \nrightarrow \mathcal{D}$  is equivalently:

00J1

1. A functor  $\mathbf{p}: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$ .

00J2

2. A functor  $\mathbf{p}: C \rightarrow \text{PSh}(\mathcal{D})$ .

00J3

3. A functor  $\mathbf{p}: \mathcal{D}^{\text{op}} \rightarrow \text{CoPSh}(C)$ .

00J4

4. A colimit-preserving functor  $\mathbf{p}: \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$ .

02C7

5. A limit-preserving functor  $\mathbf{p}: \text{CoPSh}(\mathcal{D})^{\text{op}} \rightarrow \text{CoPSh}(C)^{\text{op}}$ .

Indeed:

- The equivalence between [Items 1](#) and [2](#) (and also that between [Items 1](#) and [3](#), which is proved analogously) is an instance of currying, both for profunctors as well as for relations, using the isomorphisms

$$\begin{aligned}\text{Sets}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Sets}(A, \text{Sets}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Sets}(A, \mathcal{P}(B)),\end{aligned}$$

and

$$\begin{aligned}\text{Fun}(\mathcal{D}^{\text{op}} \times \mathcal{D}, \text{Sets}) &\cong \text{Fun}(C, \text{Fun}(\mathcal{D}^{\text{op}}, \text{Sets})) \\ &\cong \text{Fun}(C, \text{PSh}(\mathcal{D})).\end{aligned}$$

- The equivalence between [Items 2](#) and [4](#) follows from the universal properties of:

- The powerset  $\mathcal{P}(X)$  of a set  $X$  as the free cocompletion of  $X$  via the characteristic embedding

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

of  $X$  into  $\mathcal{P}(X)$ , as stated and proved in [Proposition 4.4.5.1.1](#).

- The category  $\text{PSh}(C)$  of presheaves on a category  $C$  as the free cocompletion of  $C$  via the Yoneda embedding

$$\mathfrak{y} : C \hookrightarrow \text{PSh}(C)$$

of  $C$  into  $\text{PSh}(C)$ , as stated and proved in ?? of [Proposition 12.1.4.1.3](#).

- The equivalence between [Items 3](#) and [5](#) follows from the universal properties of:

- The powerset  $\mathcal{P}(X)$  of a set  $X$  as the free completion of  $X$  via the characteristic embedding

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

of  $X$  into  $\mathcal{P}(X)$ , as stated and proved in [Proposition 4.4.6.1.1](#).

- The category  $\text{CoPSh}(\mathcal{D})^{\text{op}}$  of copresheaves on a category  $\mathcal{D}$  as the free completion of  $\mathcal{D}$  via the dual Yoneda embedding

$$\mathfrak{F} : \mathcal{D} \hookrightarrow \text{CoPSh}(\mathcal{D})^{\text{op}}$$

of  $\mathcal{D}$  into  $\text{CoPSh}(\mathcal{D})^{\text{op}}$ , as stated and proved in ?? of [Proposition 12.1.4.1.3](#).

**00QT 8.1.3 Composition of Relations**

Let  $A$ ,  $B$ , and  $C$  be sets and let  $R: A \rightarrow B$  and  $S: B \rightarrow C$  be relations.

**00QU**

**DEFINITION 8.1.3.1.1 ► COMPOSITION OF RELATIONS**

The **composition of  $R$  and  $S$**  is the relation  $S \diamond R$  defined as follows:

**02D1**

- Viewing relations from  $A$  to  $C$  as subsets of  $A \times C$ , we define

$$S \diamond R \stackrel{\text{def}}{=} \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } a \sim_R b \text{ and } b \sim_S c \end{array} \right\}.$$

**02D2**

- Viewing relations as functions  $A \times B \rightarrow \{\text{true}, \text{false}\}$ , we define

$$\begin{aligned} (S \diamond R)^{-1}_{-2} &\stackrel{\text{def}}{=} \int^{b \in B} S_b^{-1} \times R_{-2}^b \\ &= \bigvee_{b \in B} S_b^{-1} \times R_{-2}^b, \end{aligned}$$

where the join  $\vee$  is taken in the poset  $(\{\text{true}, \text{false}\}, \preceq)$  of [Definition 3.2.2.1.3](#).

**02D3**

- Viewing relations as functions  $A \rightarrow \mathcal{P}(B)$ , we define

$$\begin{array}{ccc} B & \xrightarrow{S} & \mathcal{P}(C), \\ S \diamond R \stackrel{\text{def}}{=} \text{Lan}_{\chi_B}(S) \circ R, & \chi_B \downarrow & \nearrow \text{Lan}_{\chi_B}(S) \\ A & \xrightarrow[R]{} & \mathcal{P}(B) \end{array}$$

where  $\text{Lan}_{\chi_B}(S)$  is computed by the formula

$$\begin{aligned} [\text{Lan}_{\chi_B}(S)](V) &\cong \int^{b \in B} \chi_{\mathcal{P}(B)}(\chi_b, V) \odot S(b) \\ &\cong \int^{b \in B} \chi_V(b) \odot S(b) \\ &\cong \bigcup_{b \in B} \chi_V(b) \odot S(b) \\ &\cong \bigcup_{b \in V} S(b) \end{aligned}$$

for each  $V \in \mathcal{P}(B)$ , so we have<sup>1</sup>

$$\begin{aligned} [S \diamond R](a) &\stackrel{\text{def}}{=} S(R(a)) \\ &\stackrel{\text{def}}{=} \bigcup_{b \in R(a)} S(b). \end{aligned}$$

for each  $a \in A$ .

<sup>1</sup>That is: the relation  $R$  may send  $a \in A$  to a number of elements  $\{b_i\}_{i \in I}$  in  $B$ , and then the relation  $S$  may send the image of each of the  $b_i$ 's to a number of elements  $\{S(b_i)\}_{i \in I} = \left\{\{c_{j_i}\}_{j_i \in J_i}\right\}_{i \in I}$  in  $C$ .

02D4

## REMARK 8.1.3.1.2 ► COMPOSING RELATIONS WITH RIGHT KAN EXTENSIONS

You might wonder what happens if we instead define an alternative composition of relations  $\diamond'$  via right Kan extensions. In this case, we would take the right Kan extension of  $S$  along the dual characteristic embedding  $B \hookrightarrow \mathcal{P}(B)^{\text{op}}$ :

$$\begin{array}{ccc} B & \xrightarrow{S} & \mathcal{P}(C). \\ S \diamond' R \stackrel{\text{def}}{=} \text{Ran}_{\chi_B}(S) \circ R, & \downarrow \chi_B & \nearrow \text{Ran}_{\chi'_B}(S) \\ A & \xrightarrow[R]{} & \mathcal{P}(B)^{\text{op}} \end{array}$$

In this case, we would have<sup>1</sup>

$$[S \diamond' R](a) \stackrel{\text{def}}{=} \bigcap_{b \in R(a)} S(b).$$

This alternative composition turns out to actually be a different kind of structure: it's an internal right Kan extension in **Rel**, namely  $\text{Ran}_{R^\dagger}(S)$  — see [Section 8.5.15](#).

<sup>1</sup>If we replace  $R(a)$  with  $B \setminus R(a)$ , defining

$$S \square R \stackrel{\text{def}}{=} \bigcap_{b \in B \setminus R(a)} S(b),$$

we instead obtain the apartness composition of relations; see [Section 8.1.4](#).

00QV

## EXAMPLE 8.1.3.1.3 ► EXAMPLES OF COMPOSITION OF RELATIONS

Here are some examples of composition of relations.

02AX

1. Composing Less/Greater Than Equal With Greater/Less Than Equal Signs. Let  $A = B = C = \mathbb{R}$ . We have

$$\begin{aligned} \leq \diamond \geq &= \sim_{\text{triv}}, \\ \geq \diamond \leq &= \sim_{\text{triv}}. \end{aligned}$$

02AY

2. Composing Less/Greater Than Equal Signs With Less/Greater Than Equal Signs. Let  $A = B = C = \mathbb{R}$ . We have

$$\begin{aligned} \leq \diamond \leq &= \leq, \\ \geq \diamond \geq &= \geq. \end{aligned}$$

00QW

Let  $R: A \rightarrow B$ ,  $S: B \rightarrow C$ , and  $T: C \rightarrow D$  be relations.

02D5

1. *Functionality.* The assignments  $R, S, (R, S) \mapsto S \diamond R$  define functors

$$\begin{aligned} S \diamond -: & \quad \text{Rel}(A, B) & \rightarrow \text{Rel}(A, C), \\ - \diamond R: & \quad \text{Rel}(B, C) & \rightarrow \text{Rel}(A, C), \\ -_1 \diamond -_2: & \quad \text{Rel}(B, C) \times \text{Rel}(A, B) & \rightarrow \text{Rel}(A, C). \end{aligned}$$

In particular, given relations

$$A \xrightarrow[R_1]{\quad} B \xrightarrow[S_1]{\quad} C, \\ A \xrightarrow[R_2]{\quad} B \xrightarrow[S_2]{\quad} C,$$

if  $R_1 \subset R_2$  and  $S_1 \subset S_2$ , then  $S_1 \diamond R_1 \subset S_2 \diamond R_2$ .

00QY

2. *Associativity.* We have

$$(T \diamond S) \diamond R = T \diamond (S \diamond R).$$

That is, we have

$$\bigcup_{b \in R(a)} \bigcup_{c \in S(b)} T(c) = \bigcup_{c \in \bigcup_{b \in R(a)} S(b)} T(c)$$

for each  $a \in A$ .

00QZ

3. *Unitality.* We have

$$\begin{aligned} \Delta_B \diamond R &= R, \\ R \diamond \Delta_A &= R. \end{aligned}$$

That is, we have

$$\begin{aligned} \bigcup_{b \in R(a)} \{b\} &= R(a), \\ \bigcup_{a \in \{a\}} R(a) &= R(a) \end{aligned}$$

for each  $a \in A$ .

02D6

4. *Relation to Apartness Composition of Relations.* We have

$$\begin{aligned} (S \diamond R)^c &= S^c \square R^c, \\ (S \square R)^c &= S^c \diamond R^c, \end{aligned}$$

where  $(-)^c$  is the complement functor of [Section 4.3.11](#). In particular,  $\diamond$  is a special case of apartness composition of relations, as we have

$$S \diamond R = (S^c \square R^c)^c.$$

This is also compatible with units, as we have  $\Delta_A^c = \nabla_A$ .

02D7

5. *Linear Distributivity.* We have inclusions of relations

$$\begin{aligned} T \diamond (S \square R) &\subset (T \diamond S) \square R, \\ (T \square S) \diamond R &\subset T \square (S \diamond R). \end{aligned}$$

That is, we have

$$\begin{aligned} T\left(\bigcap_{b \in B \setminus R(a)} S(b)\right) &\subset \bigcap_{b \in B \setminus R(a)} T(S(b)) \\ \bigcup_{b \in R(a)} \bigcap_{c \in C \setminus S(b)} T(c) &\subset \bigcap_{c \in C \setminus S(R(a))} T(c) \end{aligned}$$

or, unwinding the expression for  $S(R(a))$ , we have

$$\begin{aligned} \bigcup_{c \in \bigcap_{b \in B \setminus R(a)} S(b)} T(c) &\subset \bigcap_{b \in B \setminus R(a)} \bigcup_{c \in S(b)} T(c) \\ \bigcup_{b \in R(a)} \bigcap_{c \in C \setminus S(b)} T(c) &\subset \bigcap_{c \in C \setminus \bigcup_{b \in R(a)} S(b)} T(c) \end{aligned}$$

for each  $a \in A$ .

00R0

6. *Interaction With Converses.* We have

$$(S \diamond R)^\dagger = R^\dagger \diamond S^\dagger.$$

00QX

7. *Interaction With Ranges and Domains.* We have

$$\begin{aligned} \text{dom}(S \diamond R) &\subset \text{dom}(R), \\ \text{range}(S \diamond R) &\subset \text{range}(S). \end{aligned}$$

#### PROOF 8.1.3.1.5 ► PROOF OF PROPOSITION 8.1.3.1.4

##### Item 1: Functoriality

We have

$$\begin{aligned} S_1 \diamond R_1 &\stackrel{\text{def}}{=} \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{there exists some } b \in B, \text{ such} \\ \text{that } a \sim_{R_1} b \text{ or } b \sim_{S_1} c \end{array} \right\} \\ &\subset \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{there exists some } b \in B, \text{ such} \\ \text{that } a \sim_{R_2} b \text{ or } b \sim_{S_2} c \end{array} \right\} \\ &\stackrel{\text{def}}{=} S_2 \diamond R_2. \end{aligned}$$

This finishes the proof.

**Item 2: Associativity, Proof I**

Indeed, we have

$$\begin{aligned}
 (T \diamond S) \diamond R &\stackrel{\text{def}}{=} \left( \int^{c \in C} T_c^{-1} \times S_{-2}^c \right) \diamond R \\
 &\stackrel{\text{def}}{=} \int^{b \in B} \left( \int^{c \in C} T_c^{-1} \times S_b^c \right) \times R_{-2}^b \\
 &= \int^{b \in B} \int^{c \in C} (T_c^{-1} \times S_b^c) \times R_{-2}^b \\
 &= \int^{c \in C} \int^{b \in B} (T_c^{-1} \times S_b^c) \times R_{-2}^b \\
 &= \int^{c \in C} \int^{b \in B} T_c^{-1} \times (S_b^c \times R_{-2}^b) \\
 &= \int^{c \in C} T_c^{-1} \times \left( \int^{b \in B} S_b^c \times R_{-2}^b \right) \\
 &\stackrel{\text{def}}{=} \int^{c \in C} T_c^{-1} \times (S \diamond R)_{-2}^c \\
 &\stackrel{\text{def}}{=} T \diamond (S \diamond R).
 \end{aligned}$$

In the language of relations, given  $a \in A$  and  $d \in D$ , the stated equality witnesses the equivalence of the following two statements:

02D8

1. We have  $a \sim_{(T \diamond S) \diamond R} d$ , i.e. there exists some  $b \in B$  such that:

- We have  $a \sim_R b$ ;
- We have  $b \sim_{T \diamond S} d$ , i.e. there exists some  $c \in C$  such that:
  - We have  $b \sim_S c$ ;
  - We have  $c \sim_T d$ ;

02D9

2. We have  $a \sim_{T \diamond (S \diamond R)} d$ , i.e. there exists some  $c \in C$  such that:

- We have  $a \sim_{S \diamond R} c$ , i.e. there exists some  $b \in B$  such that:
  - We have  $a \sim_R b$ ;
  - We have  $b \sim_S c$ ;
- We have  $c \sim_T d$ ;

both of which are equivalent to the statement

- (★) There exist  $b \in B$  and  $c \in C$  such that  $a \sim_R b \sim_S c \sim_T d$ .

## Item 2: Associativity, Proof II

Using **Item 3 of Definition 8.1.3.1.1**, we have

$$\begin{aligned} [(T \diamond S) \diamond R](a) &\stackrel{\text{def}}{=} \bigcup_{b \in R(a)} (T \diamond S)(b) \\ &\stackrel{\text{def}}{=} \bigcup_{b \in R(a)} \bigcup_{c \in S(b)} T(c) \end{aligned}$$

on the one hand and

$$\begin{aligned} [T \diamond (S \diamond R)](a) &\stackrel{\text{def}}{=} \bigcup_{c \in [S \diamond R](a)} T(c) \\ &\stackrel{\text{def}}{=} \bigcup_{c \in \bigcup_{b \in R(a)} S(b)} T(c) \end{aligned}$$

on the other, so we want to prove an equality of the form

$$\bigcup_{b \in R(a)} \bigcup_{c \in S(b)} T(c) = \bigcup_{c \in \bigcup_{b \in R(a)} S(b)} T(c).$$

This then follows from an application of **Item 2 of Proposition 4.3.6.1.2** in which we consider  $X = D$ , consider  $\mathcal{P}(\mathcal{P}(\mathcal{P}(D)))$ , take  $U = U_c = T(c)$ , take  $A$  to be

$$A_b \stackrel{\text{def}}{=} \{T(c) \in \mathcal{P}(D) \mid c \in S(b)\},$$

and then finally take

$$\begin{aligned} \mathcal{A} &\stackrel{\text{def}}{=} \{A_b \in \mathcal{P}(\mathcal{P}(D)) \mid b \in R(a)\} \\ &\stackrel{\text{def}}{=} \{\{T(c) \in \mathcal{P}(D) \mid c \in S(b)\} \mid b \in R(a)\}. \end{aligned}$$

Indeed, we have

$$\begin{aligned} \bigcup_{A \in \mathcal{A}} (\bigcup_{U \in A} U) &= \bigcup_{A_b \in \mathcal{A}} (\bigcup_{c \in S(b)} T(c)) \\ &= \bigcup_{b \in R(a)} (\bigcup_{c \in S(b)} T(c)) \end{aligned}$$

and

$$\begin{aligned} \bigcup_{U \in \bigcup_{A \in \mathcal{A}} A} U &= \bigcup_{U_c \in \bigcup_{b \in R(a)} A_b} U_c \\ &= \bigcup_{T(c) \in \bigcup_{b \in R(a)} A_b} T(c) \end{aligned}$$

$$= \bigcup_{c \in \bigcup_{b \in R(a)} S(b)} T(c).$$

This finishes the proof.

**Item 3: Unitality**

Indeed, we have

$$\begin{aligned} \Delta_B \diamond R &\stackrel{\text{def}}{=} \int^{b \in B} (\Delta_B)_b^{-1} \times R_{-2}^b \\ &= \bigvee_{b \in B} (\Delta_B)_b^{-1} \times R_{-2}^b \\ &= \bigvee_{\substack{b \in B \\ b=-1}} R_{-2}^b \\ &= R_{-2}^{-1}, \end{aligned}$$

and

$$\begin{aligned} R \diamond \Delta_A &\stackrel{\text{def}}{=} \int^{a \in A} R_a^{-1} \times (\Delta_A)_{-2}^a \\ &= \bigvee_{a \in B} R_a^{-1} \times (\Delta_A)_{-2}^a \\ &= \bigvee_{\substack{a \in B \\ a=-2}} R_a^{-1} \\ &= R_{-2}^{-1}. \end{aligned}$$

In the language of relations, given  $a \in A$  and  $b \in B$ :

- The equality

$$\Delta_B \diamond R = R$$

witnesses the equivalence of the following two statements:

- We have  $a \sim_b B$ .
- There exists some  $b' \in B$  such that:
  - \* We have  $a \sim_R b'$
  - \* We have  $b' \sim_{\Delta_B} b$ , i.e.  $b' = b$ .

- The equality

$$R \diamond \Delta_A = R$$

witnesses the equivalence of the following two statements:

- There exists some  $a' \in A$  such that:
  - \* We have  $a \sim_{\Delta_B} a'$ , i.e.  $a = a'$ .
  - \* We have  $a' \sim_R b$
- We have  $a \sim_b B$ .

**Item 4: Relation to Apartness Composition of Relations**

This is a repetition of **Item 4** of [Proposition 8.1.4.1.3](#) and is proved there.

**Item 5: Linear Distributivity**

This is a repetition of **Item 5** of [Proposition 8.1.4.1.3](#) and is proved there.

**Item 6: Interaction With Converses**

This is a repetition of **Item 3** of [Proposition 8.1.5.1.3](#) and is proved there.

**Item 7: Interaction With Ranges and Domains**

We have

$$\begin{aligned}\text{dom}(S \diamond R) &\stackrel{\text{def}}{=} \{a \in A \mid a \sim_{S \diamond R} c \text{ for some } c \in C\}, \\ &= \left\{ a \in A \mid \begin{array}{l} \text{there exists some } b \in B \text{ and } c \in C \\ \text{such that } a \sim_R b \text{ and } b \sim_R c \end{array} \right\}, \\ &\subset \left\{ a \in A \mid \begin{array}{l} \text{there exists some } b \in B \\ \text{such that } a \sim_R b \end{array} \right\}, \\ &\stackrel{\text{def}}{=} \text{dom}(R)\end{aligned}$$

and

$$\begin{aligned}\text{range}(S \diamond R) &\stackrel{\text{def}}{=} \{c \in C \mid a \sim_{S \diamond R} c \text{ for some } a \in A\}, \\ &= \left\{ c \in C \mid \begin{array}{l} \text{there exists some } a \in A \text{ and } b \in B \\ \text{such that } a \sim_R b \text{ and } b \sim_R c \end{array} \right\}, \\ &\subset \left\{ c \in C \mid \begin{array}{l} \text{there exists some } b \in B \\ \text{such that } b \sim_S c \end{array} \right\}, \\ &\stackrel{\text{def}}{=} \text{range}(S).\end{aligned}$$

This finishes the proof. □

## 02CY 8.1.4 Apartness Composition of Relations

Let  $A$ ,  $B$ , and  $C$  be sets and let  $R: A \nrightarrow B$  and  $S: B \nrightarrow C$  be relations.

02DA

**DEFINITION 8.1.4.1.1 ► APARTNESS COMPOSITION OF RELATIONS**

The **apartness composition of  $R$  and  $S$**  is the relation  $S \square R$  defined as follows:

- Viewing relations as subsets of  $A \times C$ , we define

$$S \square R \stackrel{\text{def}}{=} \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ a \sim_R b \text{ or } b \sim_S c \end{array} \right\}.$$

- Viewing relations as functions  $A \times C \rightarrow \{\text{true, false}\}$ , we define

$$\begin{aligned} (S \square R)^{-1}_{-2} &\stackrel{\text{def}}{=} \int_{b \in B} S_b^{-1} \sqcup R_{-2}^b \\ &= \bigwedge_{b \in B} S_b^{-1} \sqcup R_{-2}^b, \end{aligned}$$

where the meet  $\wedge$  is taken in the poset  $(\{\text{true, false}\}, \preceq)$  of [Definition 3.2.2.1.3](#).

- Viewing relations as functions  $A \rightarrow \mathcal{P}(C)$ , we define

$$[S \square R](a) \stackrel{\text{def}}{=} \bigcap_{b \in B \setminus R(a)} S(b)$$

for each  $a \in A$ .

02DB

**EXAMPLE 8.1.4.1.2 ► EXAMPLES OF APARTNESS COMPOSITION OF RELATIONS**

Here are some examples of apartness composition of relations.

02DC

1. *Composing Less/Greater Than Equal With Greater/Less Than Equal Signs.* Let  $A = B = C = \mathbb{R}$ . We have

$$\begin{aligned} \leq \square \geq &= \emptyset, \\ \geq \square \leq &= \emptyset. \end{aligned}$$

02DD

2. *Composing Less/Greater Than Equal Signs With Less/Greater Than Equal Signs.* Let  $A = B = C = \mathbb{R}$ . We have

$$\begin{aligned} \leq \square \leq &= \emptyset, \\ \geq \square \geq &= \emptyset. \end{aligned}$$

02DE

3. *Equality and Inequality.* Let  $A = B = C = \mathbb{Z}$ . We have

$$\begin{aligned} = \square \neq &= =, \\ \neq \square = &= =. \end{aligned}$$

02DF

4. *Subset Inclusion.* Let  $X$  be a set with at least three elements and consider the relations  $\subset$  and  $\supset$  in  $\mathcal{P}(X)$ . We have

$$\supset \square \subset = \{(U, V) \in \mathcal{P}(X) \mid U = \emptyset \text{ or } V = \emptyset\}.$$

02DG

#### PROPOSITION 8.1.4.1.3 ► PROPERTIES OF APARTNESS COMPOSITION OF RELATIONS

Let  $R: A \rightarrow B$ ,  $S: B \rightarrow C$ , and  $T: C \rightarrow D$  be relations.

02DH

1. *Functionality.* The assignments  $R, S, (R, S) \mapsto S \square R$  define functors

$$\begin{aligned} S \square -: \quad \mathbf{Rel}(A, B) &\rightarrow \mathbf{Rel}(A, C), \\ - \square R: \quad \mathbf{Rel}(B, C) &\rightarrow \mathbf{Rel}(A, C), \\ -_1 \square -_2: \quad \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B) &\rightarrow \mathbf{Rel}(A, C). \end{aligned}$$

In particular, given relations

$$A \xrightarrow[R_1]{\quad} B \xrightarrow[S_1]{\quad} C, \quad A \xrightarrow[R_2]{\quad} B \xrightarrow[S_2]{\quad} C,$$

if  $R_1 \subset R_2$  and  $S_1 \subset S_2$ , then  $S_1 \square R_1 \subset S_2 \square R_2$ .

02DJ

2. *Associativity.* We have

$$(T \square S) \square R = T \square (S \square R).$$

02DK

3. *Unitality.* We have

$$\begin{aligned} \nabla_B \square R &= R, \\ R \square \nabla_A &= R. \end{aligned}$$

02DL

4. *Relation to Composition of Relations.* We have

$$\begin{aligned} (S \square R)^c &= S^c \diamond R^c, \\ (S \diamond R)^c &= S^c \square R^c, \end{aligned}$$

where  $(-)^c$  is the complement functor of [Section 4.3.11](#). In particular,  $\square$  is a special case of composition of relations, as we have

$$S \square R = (S^c \diamond R^c)^c.$$

This is also compatible with units, as we have  $\nabla_A^c = \Delta_A$ .

02DM

5. *Linear Distributivity.* We have inclusions of relations

$$\begin{aligned} T \diamond (S \square R) &\subset (T \diamond S) \square R, \\ (T \square S) \diamond R &\subset T \square (S \diamond R). \end{aligned}$$

02DN

6. *Interaction With Convereses.* We have

$$(S \square R)^\dagger = R^\dagger \square S^\dagger.$$



## Item 1: Functoriality

We have

$$\begin{aligned} S_1 \square R_1 &\stackrel{\text{def}}{=} \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ a \sim_{R_1} b \text{ or } b \sim_{S_1} c \end{array} \right\} \\ &\subset \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ a \sim_{R_2} b \text{ or } b \sim_{S_2} c \end{array} \right\} \\ &\stackrel{\text{def}}{=} S_2 \square R_2. \end{aligned}$$

This finishes the proof.

## Item 2: Associativity

Indeed, we have

$$\begin{aligned} (T \square S) \square R &\stackrel{\text{def}}{=} \left( \int_{c \in C} T_c^{-1} \sqcup S_{-2}^c \right) \square R \\ &\stackrel{\text{def}}{=} \int_{b \in B} \left( \int_{c \in C} T_c^{-1} \sqcup S_b^c \right) \sqcup R_{-2}^b \\ &= \int_{b \in B} \int_{c \in C} (T_c^{-1} \sqcup S_b^c) \sqcup R_{-2}^b \\ &= \int_{c \in C} \int_{b \in B} (T_c^{-1} \sqcup S_b^c) \sqcup R_{-2}^b \\ &= \int_{c \in C} \int_{b \in B} T_c^{-1} \sqcup (S_b^c \sqcup R_{-2}^b) \\ &= \int_{c \in C} T_c^{-1} \sqcup \left( \int_{b \in B} S_b^c \sqcup R_{-2}^b \right) \\ &\stackrel{\text{def}}{=} \int_{c \in C} T_c^{-1} \sqcup (S \square R)_{-2}^c \\ &\stackrel{\text{def}}{=} T \square (S \square R). \end{aligned}$$

In the language of relations, given  $a \in A$  and  $d \in D$ , the stated equality witnesses the equivalence of the following two statements:

- We have  $a \sim_{(T \square S) \square R} d$ , i.e. there exists some  $b \in B$  such that:
  - We have  $a \sim_R b$ ;
  - We have  $b \sim_{T \square S} d$ , i.e. there exists some  $c \in C$  such that:
    - \* We have  $b \sim_S c$ ;
    - \* We have  $c \sim_T d$ ;
- We have  $a \sim_{T \square (S \square R)} d$ , i.e. there exists some  $c \in C$  such that:

- We have  $a \sim_{S \square R} c$ , i.e. there exists some  $b \in B$  such that:
  - \* We have  $a \sim_R b$ ;
  - \* We have  $b \sim_S c$ ;
- We have  $c \sim_T d$ ;

both of which are equivalent to the statement

- There exist  $b \in B$  and  $c \in C$  such that  $a \sim_R b \sim_S c \sim_T d$ .

### Item 3: Unitality

Indeed, we have

$$\begin{aligned}
 \nabla_B \square R &\stackrel{\text{def}}{=} \int_{b \in B} (\nabla_B)_b^{-1} \sqcup R_{-2}^b \\
 &= \bigwedge_{b \in B} (\nabla_B)_b^{-1} \sqcup R_{-2}^b \\
 &= \left( \bigwedge_{\substack{b \in B \\ b=-1}} (\nabla_B)_b^{-1} \sqcup R_{-2}^b \right) \wedge \left( \bigwedge_{\substack{b \in B \\ b \neq -1}} (\nabla_B)_b^{-1} \sqcup R_{-2}^b \right) \\
 &= ((\nabla_B)^{-1}_{-1} \sqcup R_{-2}^{-1}) \wedge \left( \bigwedge_{\substack{b \in B \\ b \neq -1}} t \sqcup R_{-2}^b \right) \\
 &= (f \sqcup R_{-2}^{-1}) \wedge \left( \bigwedge_{\substack{b \in B \\ b \neq -1}} t \right) \\
 &= R_{-2}^{-1} \wedge t \\
 &= R_{-2}^{-1},
 \end{aligned}$$

and

$$\begin{aligned}
 R \square \nabla_A &\stackrel{\text{def}}{=} \int_{a \in A} R_a^{-1} \sqcup (\nabla_A)_{-2}^a \\
 &= \bigwedge_{a \in A} R_a^{-1} \sqcup (\nabla_A)_{-2}^a \\
 &= \left( \bigwedge_{\substack{a \in A \\ a=-2}} R_a^{-1} \sqcup (\nabla_A)_{-2}^a \right) \wedge \left( \bigwedge_{\substack{a \in A \\ a \neq -2}} R_a^{-1} \sqcup (\nabla_A)_{-2}^a \right) \\
 &= (R_{-2}^{-1} \sqcup (\nabla_A)_{-2}^{-2}) \wedge \left( \bigwedge_{\substack{a \in A \\ a \neq -2}} R_a^{-1} \sqcup t \right) \\
 &= (R_{-2}^{-1} \sqcup f) \wedge \left( \bigwedge_{\substack{a \in A \\ a \neq -2}} t \right)
 \end{aligned}$$

$$\begin{aligned} &= R_{-2}^{-1} \wedge t \\ &= R_{-2}^{-1}, \end{aligned}$$

This finishes the proof.

#### Item 4: Relation to Composition of Relations

We proceed in a few steps.

- We have  $a \sim_{(S \square R)^c} b$  iff  $a \not\sim_{S \square R} b$ .
- We have  $a \not\sim_{S \square R} b$  iff the assertion “for each  $b \in B$ , we have  $a \sim_R b$  or  $b \sim_S c$ ” is false.
- That happens iff there exists some  $b \in B$  such that  $a \not\sim_R b$  and  $b \not\sim_S c$ .
- That happens iff there exists some  $b \in B$  such that  $a \sim_{R^c} b$  and  $b \sim_{S^c} c$ .

The second equality then follows from the first one by [Item 3 of Proposition 4.3.11.1.2](#).

#### Item 5: Linear Distributivity

We have

$$\begin{aligned} T \diamond (S \square R) &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{there exists some } c \in C \text{ such that} \\ \text{that } a \sim_{S \square R} c \text{ and } c \sim_T d \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{there exists some } c \in C \text{ such that} \\ c \sim_T d \text{ and, for each } b \in B, \\ \text{we have } a \sim_R b \text{ or } b \sim_S c \end{array} \right\} \\ &= \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{the following conditions are satisfied:} \\ \begin{array}{l} 1. \text{ For each } b \in B, \text{ we have } a \sim_R b \text{ or } b \sim_S c. \\ 2. \text{ There exists } c \in C \text{ such that } c \sim_T d. \end{array} \end{array} \right\} \\ &\subset \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ at least one of the} \\ \text{following conditions is satisfied:} \\ \begin{array}{l} 1. \text{ We have } a \sim_R b. \\ 2. \text{ There exists } c \in C \text{ such that } b \sim_S c \\ \text{and } c \sim_T d. \end{array} \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ a \sim_R b \text{ or there exists some } c \in C \\ \text{such that } b \sim_S c \text{ and } c \sim_T d \end{array} \right\} \end{aligned}$$

$$\begin{aligned} &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ a \sim_R b \text{ or } b \sim_{T \diamond S} d \end{array} \right\} \\ &\stackrel{\text{def}}{=} (T \diamond S) \square R \end{aligned}$$

and

$$\begin{aligned} (T \square S) \diamond R &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } a \sim_R b \text{ and } b \sim_{T \square S} d \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } a \sim_R b \text{ and, for each } c \in C, \\ \text{we have } b \sim_S c \text{ or } c \sim_T d \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{there exists some } b \in B \text{ satisfying} \\ \text{the following conditions:} \\ \begin{array}{l} 1. \text{ We have } a \sim_R b. \\ 2. \text{ For each } c \in C, \text{ we have } b \sim_S c \\ \text{or } c \sim_T d. \end{array} \end{array} \right\} \\ &\subset \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{for each } c \in C, \text{ at least one of the} \\ \text{following conditions is satisfied:} \\ \begin{array}{l} 1. \text{ We have } c \sim_T d. \\ 2. \text{ There exists some } b \in B \text{ such that} \\ \text{we have } a \sim_R b \text{ and } b \sim_S c \end{array} \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{for each } c \in C, \text{ we have } c \sim_T d \\ \text{or there exists some } b \in B, \text{ such that} \\ \text{we have } a \sim_R b \text{ and } b \sim_S c \end{array} \right\} \\ &\stackrel{\text{def}}{=} \left\{ (d, a) \in D \times A \mid \begin{array}{l} \text{for each } c \in C, \text{ we have} \\ a \sim_{S \circ R} c \text{ or } c \sim_T d \end{array} \right\} \\ &\subset T \square (S \diamond R). \end{aligned}$$

This finishes the proof.

**Item 6: Interaction With Converses**

This is a repetition of **Item 4** of **Proposition 8.1.5.1.3** and is proved there.



### 00QE 8.1.5 The Converse of a Relation

Let  $A$ ,  $B$ , and  $C$  be sets and let  $R \subset A \times B$  be a relation.

00QF

**DEFINITION 8.1.5.1.1 ► THE CONVERSE OF A RELATION**

The **converse** of  $R^1$  is the relation  $R^\dagger$  defined as follows:

- Viewing relations as subsets, we define

$$R^\dagger \stackrel{\text{def}}{=} \{(b, a) \in B \times A \mid \text{we have } a \sim_R b\}.$$

- Viewing relations as functions  $A \times B \rightarrow \{\text{true, false}\}$ , we define

$$[R^\dagger]_b^a \stackrel{\text{def}}{=} R_a^b$$

for each  $(b, a) \in B \times A$ .

- Viewing relations as functions  $A \rightarrow \mathcal{P}(B)$ , we define<sup>2</sup>

$$R^\dagger(b) \stackrel{\text{def}}{=} \{a \in A \mid b \in R(a)\}$$

for each  $b \in B$ .

<sup>1</sup>Further Terminology: Also called the **opposite** of  $R$  or the **transpose** of  $R$ .

<sup>2</sup>Note that  $R^\dagger(b) = R^{-1}(\{b\})$ .

00QG

**EXAMPLE 8.1.5.1.2 ► EXAMPLES OF CONVERSES OF RELATIONS**

Here are some examples of converses of relations.

00QH

1. *Less Than Equal Signs.* We have  $(\leq)^\dagger = \geq$ .

00QJ

2. *Greater Than Equal Signs.* Dually to Item 1, we have  $(\geq)^\dagger = \leq$ .

00QK

3. *Functions.* Let  $f: A \rightarrow B$  be a function. We have

$$\begin{aligned} \text{Gr}(f)^\dagger &= f^{-1}, \\ (f^{-1})^\dagger &= \text{Gr}(f), \end{aligned}$$

where  $\text{Gr}(f)$  and  $f^{-1}$  are the relations of Sections 8.2.2 and 8.2.3.

00QL

**PROPOSITION 8.1.5.1.3 ► PROPERTIES OF CONVERSES OF RELATIONS**

Let  $R: A \rightarrow B$  and  $S: B \rightarrow C$  be relations.

00QM

1. *Functionality.* The assignment  $R \mapsto R^\dagger$  defines a functor (i.e. morphism of posets)

$$(-)^\dagger: \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(B, A).$$

In other words, given relations  $R, S: A \rightarrow B$ , we have:

00QN

(★) If  $R \subset S$ , then  $R^\dagger \subset S^\dagger$ .

2. *Interaction With Ranges and Domains.* We have

$$\begin{aligned}\text{dom}(R^\dagger) &= \text{range}(R), \\ \text{range}(R^\dagger) &= \text{dom}(R).\end{aligned}$$

00QP

3. *Interaction With Composition.* We have

$$(S \diamond R)^\dagger = R^\dagger \diamond S^\dagger.$$

02DP

4. *Interaction With Apartness Composition.* We have

$$(S \square R)^\dagger = R^\dagger \square S^\dagger.$$

00QR

5. *Invertibility.* We have

$$(R^\dagger)^\dagger = R.$$

00QS

6. *Identity I.* We have

$$\Delta_A^\dagger = \Delta_A.$$

02DQ

7. *Identity II.* We have

$$\nabla_A^\dagger = \nabla_A.$$

#### PROOF 8.1.5.1.4 ► PROOF OF PROPOSITION 8.1.5.1.3

##### Item 1: Functoriality

We have

$$\begin{aligned}R^\dagger &\stackrel{\text{def}}{=} \{a \in A \mid b \in R(a)\} \\ &\subset \{a \in A \mid b \in S(a)\} \\ &\stackrel{\text{def}}{=} S^\dagger.\end{aligned}$$

This finishes the proof.

##### Item 2: Interaction With Ranges and Domains

We have

$$\begin{aligned}\text{dom}(R^\dagger) &\stackrel{\text{def}}{=} \{b \in B \mid b \sim_{R^\dagger} a \text{ for some } a \in A\} \\ &= \{b \in B \mid a \sim_R b \text{ for some } a \in A\}\end{aligned}$$

$$\stackrel{\text{def}}{=} \text{range}(R)$$

and

$$\begin{aligned} \text{range}(R^\dagger) &\stackrel{\text{def}}{=} \{a \in A \mid b \sim_{R^\dagger} a \text{ for some } b \in B\} \\ &= \{a \in A \mid a \sim_R b \text{ for some } b \in B\} \\ &\stackrel{\text{def}}{=} \text{dom}(R). \end{aligned}$$

This finishes the proof.

**Item 3: Interaction With Composition**

We have

$$\begin{aligned} (S \diamond R)^\dagger &\stackrel{\text{def}}{=} \{(c, a) \in C \times A \mid c \sim_{(S \diamond R)^\dagger} a\} \\ &= \{(c, a) \in C \times A \mid a \sim_{S \diamond R} c\} \\ &= \left\{ (c, a) \in C \times A \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } a \sim_R b \text{ and } b \sim_S c \end{array} \right\} \\ &= \left\{ (c, a) \in C \times A \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } b \sim_{R^\dagger} a \text{ and } c \sim_S b \end{array} \right\} \\ &= \left\{ (c, a) \in C \times A \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } c \sim_{S^\dagger} b \text{ and } b \sim_{R^\dagger} a \end{array} \right\} \\ &\stackrel{\text{def}}{=} R^\dagger \diamond S^\dagger. \end{aligned}$$

This finishes the proof.

**Item 4: Interaction With Apartness Composition**

We have

$$\begin{aligned} (S \square R)^\dagger &\stackrel{\text{def}}{=} \{(c, a) \in C \times A \mid c \sim_{(S \square R)^\dagger} a\} \\ &= \{(c, a) \in C \times A \mid a \sim_{S \square R} c\} \\ &= \left\{ (c, a) \in C \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ a \sim_R b \text{ or } b \sim_S c \end{array} \right\} \\ &= \left\{ (c, a) \in C \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ b \sim_{R^\dagger} a \text{ or } c \sim_{S^\dagger} b \end{array} \right\} \\ &= \left\{ (c, a) \in C \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ we have} \\ c \sim_{S^\dagger} b \text{ or } b \sim_{R^\dagger} a \end{array} \right\} \\ &\stackrel{\text{def}}{=} R^\dagger \square S^\dagger. \end{aligned}$$

This finishes the proof.

**Item 5: Invertibility**

We have

$$\begin{aligned}(R^\dagger)^\dagger &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid b \sim_{R^\dagger} a\} \\ &= \{(a, b) \in A \times B \mid a \sim_R b\} \\ &\stackrel{\text{def}}{=} R.\end{aligned}$$

This finishes the proof.

**Item 6: Identity I**

We have

$$\begin{aligned}\Delta_A^\dagger &\stackrel{\text{def}}{=} \{(a, b) \in A \times A \mid a \sim_{\Delta_A} b\} \\ &= \{(a, b) \in A \times A \mid a = b\} \\ &= \Delta_A.\end{aligned}$$

This finishes the proof.

**Item 7: Identity II**

We have

$$\begin{aligned}\nabla_A^\dagger &\stackrel{\text{def}}{=} \{(a, b) \in A \times A \mid a \sim_{\nabla_A} b\} \\ &= \{(a, b) \in A \times A \mid a \neq b\} \\ &= \nabla_A.\end{aligned}$$

This finishes the proof. 

## 02CZ 8.2 Examples of Relations

### 00J5 8.2.1 Elementary Examples of Relations

00J6

#### EXAMPLE 8.2.1.1.1 ► THE TRIVIAL RELATION

The **trivial relation on  $A$  and  $B$**  is the relation  $\sim_{\text{triv}}$  defined equivalently as follows:

0291

- As a subset of  $A \times B$ , we have

$$\sim_{\text{triv}} \stackrel{\text{def}}{=} A \times B.$$

0292

- As a function from  $A \times B$  to {true, false}, the relation  $\sim_{\text{triv}}$  is the constant function

$$\Delta_{\text{true}}: A \times B \rightarrow \{\text{true, false}\}$$

from  $A \times B$  to {true, false} taking the value true.

- 0293 3. As a function from  $A$  to  $\mathcal{P}(B)$ , the relation  $\sim_{\text{triv}}$  is the function

$$\Delta_{\text{true}} : A \rightarrow \mathcal{P}(B)$$

defined by

$$\Delta_{\text{true}}(a) \stackrel{\text{def}}{=} B$$

for each  $a \in A$ .

- 0294 4. Lastly, it is the unique relation  $R$  on  $A$  and  $B$  such that we have  $a \sim_R b$  for each  $a \in A$  and each  $b \in B$ .

00J7 **EXAMPLE 8.2.1.1.2 ► THE COTRIVIAL RELATION**

The **cotrivial relation on  $A$  and  $B$**  is the relation  $\sim_{\text{cotriv}}$  defined equivalently as follows:

- 0295 1. As a subset of  $A \times B$ , we have

$$\sim_{\text{cotriv}} \stackrel{\text{def}}{=} \emptyset.$$

- 0296 2. As a function from  $A \times B$  to {true, false}, the relation  $\sim_{\text{cotriv}}$  is the constant function

$$\Delta_{\text{false}} : A \times B \rightarrow \{\text{true, false}\}$$

from  $A \times B$  to {true, false} taking the value false.

- 0297 3. As a function from  $A$  to  $\mathcal{P}(B)$ , the relation  $\sim_{\text{cotriv}}$  is the function

$$\Delta_{\text{false}} : A \rightarrow \mathcal{P}(B)$$

defined by

$$\Delta_{\text{false}}(a) \stackrel{\text{def}}{=} \emptyset$$

for each  $a \in A$ .

- 0298 4. Lastly, it is the unique relation  $R$  on  $A$  and  $B$  such that we have  $a \not\sim_R b$  for each  $a \in A$  and each  $b \in B$ .

00J8 **EXAMPLE 8.2.1.1.3 ► THE CHARACTERISTIC RELATION OF A SET**

The characteristic relation  $\chi_X$  on  $X$  of [Definition 4.5.3.1.1](#):

02DR

- As a subset of  $X \times X$ , we have

$$\begin{aligned}\sim_{\chi_X} &\stackrel{\text{def}}{=} \Delta_X \\ &\stackrel{\text{def}}{=} \{(x, x) \in X \times X\}.\end{aligned}$$

02DS

- As a function from  $X \times X$  to {true, false}, we have

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each  $x, y \in X$ .

02DT

- As a function from  $X$  to  $\mathcal{P}(X)$ , we have

$$\chi_X(x) \stackrel{\text{def}}{=} \{x\}$$

for each  $x \in X$ .

02DU

#### EXAMPLE 8.2.1.4 ► THE ANTIAGONAL RELATION ON A SET

The **antidiagonal relation on  $X$**  is the relation  $\nabla_X$  defined equivalently as follows:

02DV

- As a subset of  $X \times X$ , we have

$$\begin{aligned}\sim_{\nabla_X} &\stackrel{\text{def}}{=} \nabla_X \\ &\stackrel{\text{def}}{=} X \setminus \Delta_X \\ &= \{(x, y) \in X \times X \mid x \neq y\}.\end{aligned}$$

02DW

- As a function from  $X \times X$  to {true, false}, we have

$$\nabla_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } a \neq b, \\ \text{false} & \text{if } a = b \end{cases}$$

for each  $x, y \in X$ .

02DX

- As a function from  $X$  to  $\mathcal{P}(X)$ , we have

$$\nabla_X(x) \stackrel{\text{def}}{=} X \setminus \{x\}$$

for each  $x \in X$ .

02DY

**EXAMPLE 8.2.1.5 ► PARTIAL FUNCTIONS**

Partial functions may be viewed (or defined) as being exactly those relations which are functional; see [Section 10.1.1](#).

00J9

**EXAMPLE 8.2.1.6 ► SQUARE ROOTS**

Square roots are examples of relations:

0299

1. *Square Roots in  $\mathbb{R}$* . The assignment  $x \mapsto \sqrt{x}$  defines a relation

$$\sqrt{-} : \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$$

from  $\mathbb{R}$  to itself, being explicitly given by

$$\sqrt{x} \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x = 0, \\ \{-\sqrt{|x|}, \sqrt{|x|}\} & \text{if } x \neq 0. \end{cases}$$

029A

2. *Square Roots in  $\mathbb{Q}$* . Square roots in  $\mathbb{Q}$  are similar to square roots in  $\mathbb{R}$ , though now additionally it may also occur that  $\sqrt{-} : \mathbb{Q} \rightarrow \mathcal{P}(\mathbb{Q})$  sends a rational number  $x$  (e.g. 2) to the empty set (since  $\sqrt{2} \notin \mathbb{Q}$ ).

00JA

**EXAMPLE 8.2.1.7 ► COMPLEX LOGARITHMS**

The complex logarithm defines a relation

$$\log : \mathbb{C} \rightarrow \mathcal{P}(\mathbb{C})$$

from  $\mathbb{C}$  to itself, where we have

$$\log(a + bi) \stackrel{\text{def}}{=} \left\{ \log(\sqrt{a^2 + b^2}) + i \arg(a + bi) + (2\pi i)k \mid k \in \mathbb{Z} \right\}$$

for each  $a + bi \in \mathbb{C}$ .

00JB

**EXAMPLE 8.2.1.8 ► MORE EXAMPLES OF RELATIONS**

See [[Wik25](#)] for more examples of relations, such as antiderivation, inverse trigonometric functions, and inverse hyperbolic functions.

**00P0 8.2.2 The Graph of a Function**

Let  $f : A \rightarrow B$  be a function.

00P1

**DEFINITION 8.2.2.1.1 ► THE GRAPH OF A FUNCTION**

The **graph of  $f$**  is the relation  $\text{Gr}(f) : A \rightarrow B$  defined as follows:<sup>1</sup>

- Viewing relations from  $A$  to  $B$  as subsets of  $A \times B$ , we define

$$\text{Gr}(f) \stackrel{\text{def}}{=} \{(a, f(a)) \in A \times B \mid a \in A\}.$$

- Viewing relations from  $A$  to  $B$  as functions  $A \times B \rightarrow \{\text{true}, \text{false}\}$ , we define

$$\text{Gr}(f)_a^b \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } b = f(a), \\ \text{false} & \text{otherwise} \end{cases}$$

for each  $(a, b) \in A \times B$ .

- Viewing relations from  $A$  to  $B$  as functions  $A \rightarrow \mathcal{P}(B)$ , we define

$$[\text{Gr}(f)](a) \stackrel{\text{def}}{=} \{f(a)\}$$

for each  $a \in A$ , i.e. we define  $\text{Gr}(f)$  as the composition

$$A \xrightarrow{f} B \xhookrightarrow{\chi_B} \mathcal{P}(B).$$

---

<sup>1</sup>*Further Terminology and Notation:* When  $f = \text{id}_A$ , we write  $\text{Gr}(A)$  for  $\text{Gr}(\text{id}_A)$ , calling it the **graph of  $A$** .

00P2

**PROPOSITION 8.2.2.1.2 ► PROPERTIES OF GRAPHS OF FUNCTIONS**

Let  $f : A \rightarrow B$  be a function.

00P3

1. *Functionality.* The assignment  $A \mapsto \text{Gr}(A)$  defines a functor

$$\text{Gr} : \text{Sets} \rightarrow \text{Rel}$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$\text{Gr}(A) \stackrel{\text{def}}{=} A.$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(\text{Sets})$ , the action on Hom-sets

$$\text{Gr}_{A,B} : \text{Sets}(A, B) \rightarrow \underbrace{\text{Rel}(\text{Gr}(A), \text{Gr}(B))}_{\stackrel{\text{def}}{=} \text{Rel}(A, B)}$$

of  $\text{Gr}$  at  $(A, B)$  is defined by

$$\text{Gr}_{A,B}(f) \stackrel{\text{def}}{=} \text{Gr}(f),$$

where  $\text{Gr}(f)$  is the graph of  $f$  as in [Definition 8.2.2.1.1](#).

In particular, the following statements are true:

- *Preservation of Identities.* We have

$$\text{Gr}(\text{id}_A) = \chi_A$$

for each  $A \in \text{Obj}(\text{Sets})$ .

- *Preservation of Composition.* We have

$$\text{Gr}(g \circ f) = \text{Gr}(g) \diamond \text{Gr}(f)$$

for each pair of functions  $f: A \rightarrow B$  and  $g: B \rightarrow C$ .

00P5

2. *Adjointness.* We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_!): \text{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_!} \end{array} \text{Rel},$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(A), B) \cong \text{Sets}(A, \mathcal{P}(B))$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $B \in \text{Obj}(\text{Rel})$ .

00P7

3. *Cocontinuity.* The functor  $\text{Gr}: \text{Sets} \rightarrow \text{Rel}$  of [Item 1](#) preserves colimits.

00P4

4. *Adjointness Inside **Rel**.* We have an internal adjunction

$$(\text{Gr}(f) \dashv f^{-1}): A \begin{array}{c} \xrightarrow{\text{Gr}(f)} \\ \perp \\ \xleftarrow{f^{-1}} \end{array} B$$

in **Rel**, where  $f^{-1}$  is the inverse of  $f$  of [Definition 8.2.3.1.1](#).

00P6

5. *Interaction With Converses.* We have

$$\begin{aligned} \text{Gr}(f)^\dagger &= f^{-1}, \\ (f^{-1})^\dagger &= \text{Gr}(f). \end{aligned}$$

00P8

6. *Characterisations.* Let  $R: A \rightarrow B$  be a relation. The following conditions are equivalent:

00P9

- (a) There exists a function  $f: A \rightarrow B$  such that  $R = \text{Gr}(f)$ .

00PA

- (b) The relation  $R$  is total and functional.

00PB

- (c) The inverse and coinverse images of  $R$  agree, i.e. we have  $R^{-1} = R_{-1}$ .

00PC

- (d) The relation  $R$  has a right adjoint  $R^\dagger$  in **Rel**.

**PROOF 8.2.2.1.3 ► PROOF OF PROPOSITION 8.2.2.1.2****Item 1: Functoriality**

Omitted.

**Item 2: Adjointness**

This is a repetition of [Proposition 4.4.4.1.1](#), and is proved there.

**Item 3: Cocontinuity**

This follows from [Item 2](#) and ??.

**Item 4: Adjointness Inside Rel**

We need to check that there are inclusions

$$\begin{aligned}\chi_A &\subset f^{-1} \diamond \text{Gr}(f), \\ \text{Gr}(f) \diamond f^{-1} &\subset \chi_B.\end{aligned}$$

These correspond respectively to the following conditions:

**02AR**

1. For each  $a \in A$ , there exists some  $b \in B$  such that  $a \sim_{\text{Gr}(f)} b$  and  $b \sim_{f^{-1}} a$ .
2. For each  $a, b \in A$ , if  $a \sim_{\text{Gr}(f)} b$  and  $b \sim_{f^{-1}} a$ , then  $a = b$ .

**02AS**

In other words, the first condition states that the image of any  $a \in A$  by  $f$  is nonempty, whereas the second condition states that  $f$  is not multivalued. As  $f$  is a function, both of these statements are true, and we are done.

**Item 5: Interaction With Converses**

Omitted.

**Item 6: Characterisations**

We claim that [Items 6a](#) to [6d](#) are indeed equivalent:

- [Item 6a](#)  $\iff$  [Item 6b](#). This is shown in the proof of [Proposition 8.5.2.1.3](#).
- [Item 6b](#)  $\implies$  [Item 6c](#). If  $R$  is total and functional, then, for each  $a \in A$ , the set  $R(a)$  is a singleton. Since the conditions

- $R(a) \cap V \neq \emptyset$ ;
- $R(a) \subset V$ ;

are equivalent when  $R(a)$  is a singleton, it follows that the sets

$$R^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \cap V \neq \emptyset\},$$

$$R_{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \subset V\}$$

are equal for all  $V \in \mathcal{P}(B)$ .

- *Item 6c*  $\implies$  *Item 6b*. We claim that  $R$  is indeed total and functional:

– *Totality*. We proceed in a few steps:

- \* If we had  $R(a) = \emptyset$  for some  $a \in A$ , then we would have  $a \in R_{-1}(\emptyset)$ , so that  $R_{-1}(\emptyset) \neq \emptyset$ .
- \* But since  $R^{-1}(\emptyset) = \emptyset$ , this would imply  $R_{-1}(\emptyset) \neq R^{-1}(\emptyset)$ , a contradiction.
- \* Thus  $R(a) \neq \emptyset$  for all  $a \in A$  and  $R$  is total.

– *Functionality*. If  $R^{-1} = R_{-1}$ , then we have

$$\begin{aligned}\{a\} &= R^{-1}(\{b\}) \\ &= R_{-1}(\{b\})\end{aligned}$$

for each  $b \in R(a)$  and each  $a \in A$ , and thus  $R(a) \subset \{b\}$ . But since  $R$  is total, we must have  $R(a) = \{b\}$ , so  $R$  is functional.

- *Item 6a*  $\iff$  *Item 6d*. This follows from [Proposition 8.5.3.1.1](#).

This finishes the proof. 

### 00PD 8.2.3 The Inverse of a Function

Let  $f: A \rightarrow B$  be a function.

00PE

#### DEFINITION 8.2.3.1.1 ► THE INVERSE OF A FUNCTION

The **inverse of  $f$**  is the relation  $f^{-1}: B \nrightarrow A$  defined as follows:

- Viewing relations from  $B$  to  $A$  as subsets of  $B \times A$ , we define

$$f^{-1} \stackrel{\text{def}}{=} \{(b, f^{-1}(b)) \in B \times A \mid a \in A\},$$

where

$$f^{-1}(b) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = b\}$$

for each  $b \in B$ .

- Viewing relations from  $B$  to  $A$  as functions  $B \times A \rightarrow \{\text{true, false}\}$ , we define

$$[f^{-1}]_a^b \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if there exists } a \in A \text{ with } f(a) = b, \\ \text{false} & \text{otherwise} \end{cases}$$

for each  $(b, a) \in B \times A$ .

- Viewing relations from  $B$  to  $A$  as functions  $B \rightarrow \mathcal{P}(A)$ , we define

$$f^{-1}(b) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = b\}$$

for each  $b \in B$ .

00PF

### PROPOSITION 8.2.3.1.2 ► PROPERTIES OF INVERSES OF FUNCTIONS

Let  $f: A \rightarrow B$  be a function.

00PG

1. *Functionality.* The assignment  $A \mapsto A$ ,  $f \mapsto f^{-1}$  defines a functor

$$(-)^{-1}: \mathbf{Sets} \rightarrow \mathbf{Rel}$$

where

- *Action on Objects.* For each  $A \in \mathbf{Obj}(\mathbf{Sets})$ , we have

$$[(-)^{-1}](A) \stackrel{\text{def}}{=} A.$$

- *Action on Morphisms.* For each  $A, B \in \mathbf{Obj}(\mathbf{Sets})$ , the action on Hom-sets

$$(-)_{A,B}^{-1}: \mathbf{Sets}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of  $(-)^{-1}$  at  $(A, B)$  is defined by

$$(-)_{A,B}^{-1}(f) \stackrel{\text{def}}{=} [(-)^{-1}](f),$$

where  $f^{-1}$  is the inverse of  $f$  as in [Definition 8.2.3.1.1](#).

In particular, the following statements are true:

- *Preservation of Identities.* We have

$$\text{id}_A^{-1} = \chi_A$$

for each  $A \in \mathbf{Obj}(\mathbf{Sets})$ .

- *Preservation of Composition.* We have

$$(g \circ f)^{-1} = g^{-1} \diamond f^{-1}$$

for pair of functions  $f: A \rightarrow B$  and  $g: B \rightarrow C$ .

00PH

2. *Adjointness Inside Rel.* We have an adjunction

$$\left( \text{Gr}(f) \dashv f^{-1} \right): A \begin{array}{c} \xrightarrow{\quad \text{Gr}(f) \quad} \\[-1ex] \perp \\[-1ex] \xleftarrow{\quad f^{-1} \quad} \end{array} B$$

in **Rel**.

00PJ

3. *Interaction With Converses of Relations.* We have

$$(f^{-1})^\dagger = \text{Gr}(f), \\ \text{Gr}(f)^\dagger = f^{-1}.$$

#### PROOF 8.2.3.1.3 ► PROOF OF PROPOSITION 8.2.3.1.2

Item 1: Functoriality

Omitted.

Item 2: Adjointness Inside Rel

This is a repetition of [Item 4 of Proposition 8.2.2.1.2](#) and is proved there.

Item 3: Interaction With Converses of Relations

This is a repetition of [Item 5 of Proposition 8.2.2.1.2](#) and is proved there.



## 00PK 8.2.4 Representable Relations

Let  $A$  and  $B$  be sets.

00PL

#### DEFINITION 8.2.4.1.1 ► REPRESENTABLE RELATIONS

Let  $f: A \rightarrow B$  and  $g: B \rightarrow A$  be functions.<sup>1</sup>

02AT

1. The **representable relation associated to  $f$**  is the relation  $\chi_f: A \rightarrow B$  defined as the composition

$$A \times B \xrightarrow{f \times \text{id}_B} B \times B \xrightarrow{\chi_B} \{\text{true}, \text{false}\},$$

i.e. given by declaring  $a \sim_{\chi_f} b$  iff  $f(a) = b$ .

02AU

2. The **corepresentable relation associated to  $g$**  is the relation  $\chi^g: B \rightarrow A$  defined as the composition

$$B \times A \xrightarrow{g \times \text{id}_A} A \times A \xrightarrow{\chi_A} \{\text{true}, \text{false}\},$$

i.e. given by declaring  $b \sim_{\chi^g} a$  iff  $g(b) = a$ .

<sup>1</sup>More generally, given functions

$$\begin{aligned} f: A &\rightarrow C, \\ g: B &\rightarrow D \end{aligned}$$

and a relation  $B \nrightarrow D$ , we may consider the composite relation

$$A \times B \xrightarrow{f \times g} C \times D \xrightarrow{R} \{\text{true, false}\},$$

for which we have  $a \sim_{R \circ (f \times g)} b$  iff  $f(a) \sim_R g(b)$ .

## 00JQ 8.3 Categories of Relations

### 01R1 8.3.1 The Category of Relations Between Two Sets

#### 01R2 DEFINITION 8.3.1.1 ► THE CATEGORY OF RELATIONS BETWEEN TWO SETS

The **category of relations from  $A$  to  $B$**  is the category  $\text{Rel}(A, B)$  defined by<sup>1</sup>

$$\text{Rel}(A, B) \stackrel{\text{def}}{=} \text{Rel}(A, B)_{\text{pos}},$$

where  $\text{Rel}(A, B)_{\text{pos}}$  is the posetal category associated to the poset  $\text{Rel}(A, B)$  of Item 2 of Notation 8.1.1.6 and Definition 11.2.7.1.1.

<sup>1</sup>Here we choose to abuse notation by writing  $\text{Rel}(A, B)$  instead of  $\text{Rel}(A, B)_{\text{pos}}$  for the posetal category of relations from  $A$  to  $B$ , even though the same notation is used for the poset of relations from  $A$  to  $B$ .

### 00JR 8.3.2 The Category of Relations

#### 00JS DEFINITION 8.3.2.1.1 ► THE CATEGORY OF RELATIONS

The **category of relations** is the category  $\text{Rel}$  where

- *Objects.* The objects of  $\text{Rel}$  are sets.
- *Morphisms.* For each  $A, B \in \text{Obj}(\text{Sets})$ , we have

$$\text{Rel}(A, B) \stackrel{\text{def}}{=} \text{Rel}(A, B).$$

- *Identities.* For each  $A \in \text{Obj}(\text{Rel})$ , the unit map

$$\mathbb{1}_A^{\text{Rel}}: \text{pt} \rightarrow \text{Rel}(A, A)$$

of  $\text{Rel}$  at  $A$  is defined by

$$\text{id}_A^{\text{Rel}} \stackrel{\text{def}}{=} \chi_A(-_1, -_2),$$

where  $\chi_A(-_1, -_2)$  is the characteristic relation of  $A$  of Example 8.2.1.1.3.

- *Composition.* For each  $A, B, C \in \text{Obj}(\text{Rel})$ , the composition map

$$\circ_{A,B,C}^{\text{Rel}} : \text{Rel}(B, C) \times \text{Rel}(A, B) \rightarrow \text{Rel}(A, C)$$

of Rel at  $(A, B, C)$  is defined by

$$S \circ_{A,B,C}^{\text{Rel}} R \stackrel{\text{def}}{=} S \diamond R$$

for each  $(S, R) \in \text{Rel}(B, C) \times \text{Rel}(A, B)$ , where  $S \diamond R$  is the composition of  $S$  and  $R$  of [Definition 8.1.3.1.1](#).

### **00JT 8.3.3 The Closed Symmetric Monoidal Category of Relations**

#### **00JU 8.3.3.1 The Monoidal Product**

**00JV**

##### **DEFINITION 8.3.3.1.1 ► THE MONOIDAL PRODUCT OF Rel**

The **monoidal product of Rel** is the functor

$$\times : \text{Rel} \times \text{Rel} \rightarrow \text{Rel}$$

where

- *Action on Objects.* For each  $A, B \in \text{Obj}(\text{Rel})$ , we have

$$\times(A, B) \stackrel{\text{def}}{=} A \times B,$$

where  $A \times B$  is the Cartesian product of sets of [Definition 4.1.3.1.1](#).

- *Action on Morphisms.* For each  $(A, C), (B, D) \in \text{Obj}(\text{Rel} \times \text{Rel})$ , the action on morphisms

$$\times_{(A,C),(B,D)} : \text{Rel}(A, B) \times \text{Rel}(C, D) \rightarrow \text{Rel}(A \times C, B \times D)$$

of  $\times$  is given by sending a pair of morphisms  $(R, S)$  of the form

$$\begin{aligned} R &: A \rightarrow B, \\ S &: C \rightarrow D \end{aligned}$$

to the relation

$$R \times S : A \times C \rightarrow B \times D$$

of [Definition 9.2.6.1.1](#).

### 00JW 8.3.3.2 The Monoidal Unit

00JX

## DEFINITION 8.3.3.2.1 ► THE MONOIDAL UNIT OF Rel

The **monoidal unit** of Rel is the functor

$$1^{\text{Rel}} : \text{pt} \rightarrow \text{Rel}$$

picking the set

$$\mathbb{1}_{\text{Rel}} \stackrel{\text{def}}{=} \text{pt}$$

of Rel.

### 00JY 8.3.3.3 The Associator


00JZ

### **DEFINITION 8.3.3.3.1 ► THE ASSOCIATOR OF Rel**

The **associator** of Rel is the natural isomorphism

$$\alpha^{\text{Rel}} : \times \circ ((\times) \times \text{id}) \xrightarrow{\sim} \times \circ (\text{id} \times (\times)) \circ \alpha_{\text{Rel}, \text{Rel}, \text{Rel}}^{\text{Cats}},$$

as in the diagram



whose component

$$\alpha_{A,B,C}^{\text{Rel}}: (A \times B) \times C \rightarrow A \times (B \times C)$$

at  $A, B, C \in \text{Obj}(\text{Rel})$  is the relation defined by declaring

$$((a,b),c) \sim_{\alpha^{\text{Rel}}_{A,B,C}} (a',(b',c'))$$

iff  $a = a'$ ,  $b = b'$ , and  $c = c'$ .

## 00K0 8.3.3.4 The Left Unit

00K1

## DEFINITION 8.3.3.4.1 ► THE LEFT UNIT OF Rel

The **left unit** of Rel is the natural isomorphism

$$\begin{array}{ccc} \text{pt} \times \text{Rel} & \xrightarrow{\text{id}_{\text{Rel}} \times \text{id}} & \text{Rel} \times \text{Rel}, \\ \lambda^{\text{Rel}} : \times \circ (\text{id}_{\text{Rel}} \times \text{id}) \xrightarrow{\sim} \lambda_{\text{Rel}}^{\text{Cats}_2}, & \swarrow \quad \searrow & \downarrow \times \\ \lambda_{\text{Rel}}^{\text{Cats}_2} & & \text{Rel} \end{array}$$

whose component

$$\lambda_A^{\text{Rel}} : \text{id}_{\text{Rel}} \times A \dashrightarrow A$$

at  $A$  is defined by declaring

$$(\star, a) \sim_{\lambda_A^{\text{Rel}}} b$$

iff  $a = b$ .

## 00K2 8.3.3.5 The Right Unit

00K3

## DEFINITION 8.3.3.5.1 ► THE RIGHT UNIT OF Rel

The **right unit** of Rel is the natural isomorphism

$$\rho^{\text{Rel}} : \times \circ (\text{id} \times \mathbb{1}_{\text{Rel}}) \xrightarrow{\sim} \rho_{\text{Rel}}^{\text{Cats}_2},$$

$$\begin{array}{ccc} \text{Rel} \times \text{pt} & \xrightarrow{\text{id} \times \mathbb{1}_{\text{Rel}}} & \text{Rel} \times \text{Rel}, \\ & \searrow \rho_{\text{Rel}}^{\text{Cats}_2} & \downarrow \times \\ & \nearrow \rho^{\text{Rel}} & \\ & \text{Rel} & \end{array}$$

whose component

$$\rho_A^{\text{Rel}} : A \times \mathbb{1}_{\text{Rel}} \rightarrow A$$

at  $A$  is defined by declaring

$$(a, \star) \sim_{\rho_A^{\text{Rel}}} b$$

iff  $a = b$ .

00K4 8.3.3.6 The Symmetry

00K5

## DEFINITION 8.3.3.6.1 ► THE SYMMETRY OF Rel

The **symmetry** of Rel is the natural isomorphism

$$\sigma^{\text{Rel}} : \times \Longrightarrow \times \circ \sigma_{\text{Rel}, \text{Rel}}^{\text{Cats}_2},$$

$$\begin{array}{ccc} \text{Rel} \times \text{Rel} & \xrightarrow{\times} & \text{Rel}, \\ & \searrow \sigma_{\text{Rel}, \text{Rel}}^{\text{Cats}_2} & \downarrow \sigma^{\text{Rel}} \\ & \nearrow & \swarrow \times \\ & \text{Rel} \times \text{Rel} & \end{array}$$

whose component

$$\sigma_{A,B}^{\text{Rel}} : A \times B \rightarrow B \times A$$

at  $(A, B)$  is defined by declaring

$$(a, b) \sim_{\sigma_{A,B}^{\text{Rel}}} (b', a')$$

iff  $a = a'$  and  $b = b'$ .

## 00K6 8.3.3.7 The Internal Hom

00K7

## DEFINITION 8.3.3.7.1 ► THE INTERNAL HOM OF Rel

The **internal Hom of Rel** is the functor

$$\text{Rel}: \text{Rel}^{\text{op}} \times \text{Rel} \rightarrow \text{Rel}$$

defined

- On objects by sending  $A, B \in \text{Obj}(\text{Rel})$  to the set  $\text{Rel}(A, B)$  of ?? of ??.
- On morphisms by pre/post-composition defined as in [Definition 8.1.3.1.1](#).

00K8

## PROPOSITION 8.3.3.7.2 ► PROPERTIES OF THE INTERNAL HOM OF Rel

Let  $A, B, C \in \text{Obj}(\text{Rel})$ .

00K9

1. *Adjointness.* We have adjunctions

$$(A \times - \dashv \text{Rel}(A, -)): \begin{array}{c} \text{Rel} \xrightarrow{\quad A \times - \quad} \text{Rel}, \\ \perp \\ \text{Rel}(A, -) \end{array}$$

$$(- \times B \dashv \text{Rel}(B, -)): \begin{array}{c} \text{Rel} \xrightarrow{\quad - \times B \quad} \text{Rel}, \\ \perp \\ \text{Rel}(B, -) \end{array}$$

witnessed by bijections

$$\begin{aligned} \text{Rel}(A \times B, C) &\cong \text{Rel}(A, \text{Rel}(B, C)), \\ \text{Rel}(A \times B, C) &\cong \text{Rel}(B, \text{Rel}(A, C)), \end{aligned}$$

natural in  $A, B, C \in \text{Obj}(\text{Rel})$ .

## PROOF 8.3.3.7.3 ► PROOF OF PROPOSITION 8.3.3.7.2

## Item 1: Adjointness

Indeed, we have

$$\begin{aligned} \text{Rel}(A \times B, C) &\stackrel{\text{def}}{=} \text{Sets}(A \times B \times C, \{\text{true}, \text{false}\}) \\ &\stackrel{\text{def}}{=} \text{Rel}(A, B \times C) \end{aligned}$$

$$\stackrel{\text{def}}{=} \text{Rel}(A, \text{Rel}(B, C)),$$

and similarly for the bijection  $\text{Rel}(A \times B, C) \cong \text{Rel}(B, \text{Rel}(A, C))$ . □

### 00KA 8.3.3.8 The Closed Symmetric Monoidal Category of Relations

00KB

#### PROPOSITION 8.3.3.8.1 ► THE CLOSED SYMMETRIC MONOIDAL CATEGORY OF RELATIONS

The category  $\text{Rel}$  admits a closed symmetric monoidal category structure consisting of<sup>1</sup>

- *The Underlying Category.* The category  $\text{Rel}$  of sets and relations of [Definition 8.3.2.1.1](#).

- *The Monoidal Product.* The functor

$$\times: \text{Rel} \times \text{Rel} \rightarrow \text{Rel}$$

of [Definition 8.3.3.1.1](#).

- *The Internal Hom.* The internal Hom functor

$$\text{Rel}: \text{Rel}^{\text{op}} \times \text{Rel} \rightarrow \text{Rel}$$

of [Definition 8.3.3.7.1](#).

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\text{Rel}}: \text{pt} \rightarrow \text{Rel}$$

of [Definition 8.3.3.2.1](#).

- *The Associators.* The natural isomorphism

$$\alpha^{\text{Rel}}: \times \circ (\times \times \text{id}_{\text{Rel}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Rel}} \times \times) \circ \alpha_{\text{Rel}, \text{Rel}, \text{Rel}}^{\text{Cats}_2}$$

of [Definition 8.3.3.3.1](#).

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\text{Rel}}: \times \circ (\mathbb{1}^{\text{Rel}} \times \text{id}_{\text{Rel}}) \xrightarrow{\sim} \lambda_{\text{Rel}}^{\text{Cats}_2}$$

of [Definition 8.3.3.4.1](#).

- *The Right Unitors.* The natural isomorphism

$$\rho^{\text{Rel}}: \times \circ (\text{id} \times \mathbb{1}^{\text{Rel}}) \xrightarrow{\sim} \rho_{\text{Rel}}^{\text{Cats}_2}$$

of [Definition 8.3.3.5.1](#).

- *The Symmetry.* The natural isomorphism

$$\sigma^{\text{Rel}} : \times \xrightarrow{\sim} \times \circ \sigma_{\text{Rel}, \text{Rel}}^{\text{Cats}_2}$$

of [Definition 8.3.3.6.1](#).

 **Warning:** This is not a Cartesian monoidal structure, as the product on Rel is in fact given by the disjoint union of sets; see ??.

**PROOF 8.3.3.8.2 ▶ PROOF OF PROPOSITION 8.3.3.8.1**

Omitted. 

#### 00KC 8.3.4 The 2-Category of Relations

00KD

**DEFINITION 8.3.4.1.1 ► THE 2-CATEGORY OF RELATIONS**

The **2-category of relations** is the locally posetal 2-category **Rel** where

- **Objects.** The objects of **Rel** are sets.
- **Hom-Objects.** For each  $A, B \in \text{Obj}(\text{Sets})$ , we have

$$\begin{aligned}\text{Hom}_{\text{Rel}}(A, B) &\stackrel{\text{def}}{=} \text{Rel}(A, B) \\ &\stackrel{\text{def}}{=} (\text{Rel}(A, B), \subset).\end{aligned}$$

- **Identities.** For each  $A \in \text{Obj}(\text{Rel})$ , the unit map

$$1_A^{\text{Rel}} : \text{pt} \rightarrow \text{Rel}(A, A)$$

of **Rel** at  $A$  is defined by

$$\text{id}_A^{\text{Rel}} \stackrel{\text{def}}{=} \chi_A(-_1, -_2),$$

where  $\chi_A(-_1, -_2)$  is the characteristic relation of  $A$  of [Example 8.2.1.1.3](#).

- **Composition.** For each  $A, B, C \in \text{Obj}(\text{Rel})$ , the composition map<sup>1</sup>

$$\circ_{A,B,C}^{\text{Rel}} : \text{Rel}(B, C) \times \text{Rel}(A, B) \rightarrow \text{Rel}(A, C)$$

of **Rel** at  $(A, B, C)$  is defined by

$$S \circ_{A,B,C}^{\text{Rel}} R \stackrel{\text{def}}{=} S \diamond R$$

for each  $(S, R) \in \text{Rel}(B, C) \times \text{Rel}(A, B)$ , where  $S \diamond R$  is the composition of  $S$  and  $R$  of [Definition 8.1.3.1.1](#).

---

<sup>1</sup>That this is indeed a morphism of posets is proven in ?? of [Proposition 8.1.3.1.4](#).

00KE 8.3.5 The Double Category of Relations

00KF 8.3.5.1 The Double Category of Relations

00KG

**DEFINITION 8.3.5.1.1 ► THE DOUBLE CATEGORY OF RELATIONS**

The **double category of relations** is the locally posetal double category  $\text{Rel}^{\text{dbl}}$  where

- **Objects.** The objects of  $\text{Rel}^{\text{dbl}}$  are sets.
- **Vertical Morphisms.** The vertical morphisms of  $\text{Rel}^{\text{dbl}}$  are maps of sets  $f : A \rightarrow B$ .
- **Horizontal Morphisms.** The horizontal morphisms of  $\text{Rel}^{\text{dbl}}$  are

relations  $R: A \rightarrow X$ .

- **2-Morphisms.** A 2-cell

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ X & \xrightarrow{S} & Y \end{array}$$

of  $\text{Rel}^{\text{dbl}}$  is either non-existent or an inclusion of relations of the form

$$\begin{array}{ccc} A \times B & \xrightarrow{R} & \{\text{true, false}\} \\ R \subset S \circ (f \times g), \quad f \times g \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ X \times Y & \xrightarrow{S} & \{\text{true, false}\}. \end{array}$$

- **Horizontal Identities.** The horizontal unit functor of  $\text{Rel}^{\text{dbl}}$  is the functor of [Definition 8.3.5.2.1](#).
- **Vertical Identities.** For each  $A \in \text{Obj}(\text{Rel}^{\text{dbl}})$ , we have

$$\text{id}_A^{\text{Rel}^{\text{dbl}}} \stackrel{\text{def}}{=} \text{id}_A.$$

- **Identity 2-Morphisms.** For each horizontal morphism  $R: A \rightarrow B$  of  $\text{Rel}^{\text{dbl}}$ , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ \text{id}_A \downarrow & \Downarrow \text{id}_R & \downarrow \text{id}_B \\ A & \xrightarrow{R} & B \end{array}$$

of  $R$  is the identity inclusion

$$\begin{array}{ccc} B \times A & \xrightarrow{R} & \{\text{true, false}\} \\ R \subset R, \quad \text{id}_B \times \text{id}_A \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times A & \xrightarrow{R} & \{\text{true, false}\}. \end{array}$$

- **Horizontal Composition.** The horizontal composition functor of  $\text{Rel}^{\text{dbl}}$  is the functor of [Definition 8.3.5.3.1](#).

- *Vertical Composition of 1-Morphisms.* For each composable pair  $A \xrightarrow{F} B \xrightarrow{G} C$  of vertical morphisms of  $\text{Rel}^{\text{dbl}}$ , i.e. maps of sets, we have
$$g \circ^{\text{Rel}^{\text{dbl}}} f \stackrel{\text{def}}{=} g \circ f.$$
- *Vertical Composition of 2-Morphisms.* The vertical composition of 2-morphisms in  $\text{Rel}^{\text{dbl}}$  is defined as in [Definition 8.3.5.4.1](#).
- *Associators.* The associators of  $\text{Rel}^{\text{dbl}}$  are defined as in [Definition 8.3.5.1](#).
- *Left Unitors.* The left unitors of  $\text{Rel}^{\text{dbl}}$  are defined as in [Definition 8.3.5.6.1](#).
- *Right Unitors.* The right unitors of  $\text{Rel}^{\text{dbl}}$  are defined as in [Definition 8.3.5.7.1](#).

### 00KH 8.3.5.2 Horizontal Identities

00KJ

#### DEFINITION 8.3.5.2.1 ► THE HORIZONTAL IDENTITIES OF $\text{Rel}^{\text{dbl}}$

The **horizontal unit functor** of  $\text{Rel}^{\text{dbl}}$  is the functor

$$\mathbb{1}^{\text{Rel}^{\text{dbl}}} : \text{Rel}_0^{\text{dbl}} \rightarrow \text{Rel}_1^{\text{dbl}}$$

of  $\text{Rel}^{\text{dbl}}$  is the functor where

- *Action on Objects.* For each  $A \in \text{Obj}(\text{Rel}_0^{\text{dbl}})$ , we have

$$\mathbb{1}_A \stackrel{\text{def}}{=} \chi_A(-_1, -_2).$$

- *Action on Morphisms.* For each vertical morphism  $f: A \rightarrow B$  of  $\text{Rel}^{\text{dbl}}$ , i.e. each map of sets  $f$  from  $A$  to  $B$ , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{\mathbb{1}_A} & A \\ f \downarrow & \parallel & \downarrow f \\ B & \xrightarrow{\mathbb{1}_B} & B \end{array}$$

of  $f$  is the inclusion

$$\begin{array}{ccc} A \times A & \xrightarrow{\chi_A(-_1, -_2)} & \{\text{true}, \text{false}\} \\ \chi_B \circ (f \times f) \subset \chi_A, & f \times f \downarrow & \subset \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ B \times B & \xrightarrow{\chi_B(-_1, -_2)} & \{\text{true}, \text{false}\} \end{array}$$

of Item 1 of [Proposition 4.5.3.1.3](#).

### 00KK 8.3.5.3 Horizontal Composition

00KL

#### DEFINITION 8.3.5.3.1 ► THE HORIZONTAL COMPOSITION OF $\text{Rel}^{\text{dbl}}$

The **horizontal composition functor** of  $\text{Rel}^{\text{dbl}}$  is the functor

$$\odot^{\text{Rel}^{\text{dbl}}} : \text{Rel}_1^{\text{dbl}} \times_{\text{Rel}_0^{\text{dbl}}} \text{Rel}_1^{\text{dbl}} \rightarrow \text{Rel}_1^{\text{dbl}}$$

of  $\text{Rel}^{\text{dbl}}$  is the functor where

- *Action on Objects.* For each composable pair  $A \xrightarrow{R} B \xrightarrow{S} C$  of horizontal morphisms of  $\text{Rel}^{\text{dbl}}$ , we have

$$S \odot R \stackrel{\text{def}}{=} S \diamond R,$$

where  $S \diamond R$  is the composition of  $R$  and  $S$  of [Definition 8.1.3.1.1](#).

- *Action on Morphisms.* For each horizontally composable pair

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ X & \xrightarrow{T} & Y \end{array} \quad \begin{array}{ccc} B & \xrightarrow{S} & C \\ g \downarrow & \Downarrow \beta & \downarrow h \\ Y & \xrightarrow{U} & Z \end{array}$$

of 2-morphisms of  $\text{Rel}^{\text{dbl}}$ , i.e. for each pair

$$\begin{array}{ccc} A \times B & \xrightarrow{R} & \{\text{true, false}\} \\ f \times g \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ X \times Y & \xrightarrow{T} & \{\text{true, false}\} \end{array} \quad \begin{array}{ccc} B \times C & \xrightarrow{S} & \{\text{true, false}\} \\ g \times h \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ Y \times Z & \xrightarrow{U} & \{\text{true, false}\} \end{array}$$

of inclusions of relations, the horizontal composition

$$\begin{array}{ccc} A & \xrightarrow{S \odot R} & C \\ f \downarrow & \Downarrow \beta \odot \alpha & \downarrow h \\ X & \xrightarrow{U \odot T} & Z \end{array}$$

of  $\alpha$  and  $\beta$  is the inclusion of relations

$$\begin{array}{ccc} A \times C & \xrightarrow{S \diamond R} & \{\text{true, false}\} \\ (U \diamond T) \circ (f \times h) \subset (S \diamond R) & f \times h \downarrow & \subset \qquad \downarrow \text{id}_{\{\text{true, false}\}} \\ X \times Z & \xrightarrow{U \diamond T} & \{\text{true, false}\}. \end{array}$$

#### PROOF 8.3.5.3.2 ► PROOF OF THE INCLUSION IN DEFINITION 8.3.5.3.1

The inclusion of relations

$$(U \diamond T) \circ (f \times h) \subset (S \diamond R)$$

follows from the fact that the statement

- We have  $a \sim_{(U \diamond T) \circ (f \times h)} c$ , i.e.  $f(a) \sim_{U \diamond T} h(c)$ , i.e. there exists some  $y \in Y$  such that:
  - We have  $f(a) \sim_T y$ .
  - We have  $y \sim_U h(c)$ .

is implied by the statement

- We have  $a \sim_{S \diamond R} c$ , i.e. there exists some  $b \in B$  such that:
  - We have  $a \sim_R b$ .
  - We have  $b \sim_S c$ .

since:

- If  $a \sim_R b$ , then  $f(a) \sim_T g(b)$ , as  $T \circ (f \times g) \subset R$ ;
- If  $b \sim_S c$ , then  $g(b) \sim_U h(c)$ , as  $U \circ (g \times h) \subset S$ .

This finishes the proof. □

#### 00KM 8.3.5.4 Vertical Composition of 2-Morphisms

00KN

#### DEFINITION 8.3.5.4.1 ► THE VERTICAL COMPOSITION OF 2-MORPHISMS IN $\text{Rel}^{\text{dbl}}$

The **vertical composition** in  $\text{Rel}^{\text{dbl}}$  is defined as follows: for each vertically composable pair

$$\begin{array}{ccc} A & \xrightarrow{R} & X \\ f \downarrow & \parallel \alpha & \downarrow g \\ B & \xrightarrow{S} & Y \end{array} \quad \begin{array}{ccc} B & \xrightarrow{S} & Y \\ h \downarrow & \parallel \beta & \downarrow k \\ C & \xrightarrow{T} & Z \end{array}$$

of 2-morphisms of  $\text{Rel}^{\text{dbl}}$ , i.e. for each each pair

$$\begin{array}{ccc} A \times X & \xrightarrow{R} & \{\text{true, false}\} \\ f \times g \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times Y & \xrightarrow{S} & \{\text{true, false}\} \end{array} \quad \begin{array}{ccc} B \times Y & \xrightarrow{S} & \{\text{true, false}\} \\ h \times k \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ C \times Z & \xrightarrow{T} & \{\text{true, false}\} \end{array}$$

of inclusions of relations, we define the vertical composition

$$\begin{array}{ccc} A & \xrightarrow{R} & X \\ h \circ f \downarrow & \parallel \beta \circ \alpha & \downarrow k \circ g \\ C & \xrightarrow{T} & Z \end{array}$$

of  $\alpha$  and  $\beta$  as the inclusion of relations

$$T \circ [(h \circ f) \times (k \circ g)] \subset R, \quad \begin{array}{ccc} A \times X & \xrightarrow{R} & \{\text{true, false}\} \\ (h \circ f) \times (k \circ g) \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ C \times Z & \xrightarrow{T} & \{\text{true, false}\} \end{array}$$

given by the pasting of inclusions

$$\begin{array}{ccc} A \times X & \xrightarrow{R} & \{\text{true, false}\} \\ f \times g \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times Y & \xrightarrow{S} & \{\text{true, false}\} \\ h \times k \downarrow & \curvearrowright & \downarrow \text{id}_{\{\text{true, false}\}} \\ C \times Z & \xrightarrow{T} & \{\text{true, false}\}. \end{array}$$

**PROOF 8.3.5.4.2 ► PROOF OF THE INCLUSION IN DEFINITION 8.3.5.4.1**

The inclusion

$$T \circ [(h \circ f) \times (k \circ g)] \subset R$$

follows from the fact that, given  $(a, x) \in A \times X$ , the statement

- We have  $h(f(a)) \sim_T k(g(x))$ ;

is implied by the statement

- We have  $a \sim_R x$ ;

since

- If  $a \sim_R x$ , then  $f(a) \sim_S g(x)$ , as  $S \circ (f \times g) \subset R$ ;
- If  $b \sim_S y$ , then  $h(b) \sim_T k(y)$ , as  $T \circ (h \times k) \subset S$ , and thus, in particular:
  - If  $f(a) \sim_S g(x)$ , then  $h(f(a)) \sim_T k(g(x))$ .

This finishes the proof. 

**00KP 8.3.5.5 The Associators**

00KQ

DEFINITION 8.3.5.1 ► THE ASSOCIATORS OF  $\text{Rel}^{\text{dbl}}$ 

For each composable triple

$$A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D$$

of horizontal morphisms of  $\text{Rel}^{\text{dbl}}$ , the component

$$\alpha_{T,S,R}^{\text{Rel}^{\text{dbl}}} : (T \odot S) \odot R \xrightarrow{\sim} T \odot (S \odot R), \quad \begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow{S} & C & \xrightarrow{T} & D \\ \downarrow \text{id}_A & & \downarrow \alpha_{T,S,R}^{\text{Rel}^{\text{dbl}}} & & & & \downarrow \text{id}_D \\ A & \xrightarrow[R]{\quad} & B & \xrightarrow[S]{\quad} & C & \xrightarrow[T]{\quad} & D \end{array}$$

of the associator of  $\text{Rel}^{\text{dbl}}$  at  $(R, S, T)$  is the identity inclusion<sup>1</sup>

$$\begin{array}{ccc} A \times B & \xrightarrow{(T \diamond S) \diamond R} & \{\text{true, false}\} \\ (T \diamond S) \diamond R = T \diamond (S \diamond R) & \parallel & \equiv & \downarrow \text{id}_{\{\text{true, false}\}} \\ A \times B & \xrightarrow{T \diamond (S \diamond R)} & \{\text{true, false}\}. \end{array}$$

<sup>1</sup>As proved in Item 2 of Proposition 8.1.3.1.4.

00KR 8.3.5.6 The Left Unitors

00KS

DEFINITION 8.3.6.1 ► THE LEFT UNITORS OF  $\text{Rel}^{\text{dbl}}$ 

For each horizontal morphism  $R: A \rightarrow B$  of  $\text{Rel}^{\text{dbl}}$ , the component

$$\lambda_R^{\text{Rel}^{\text{dbl}}} : \mathbb{1}_B \odot R \xrightarrow{\sim} R, \quad \begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow{\mathbb{1}_B} & B \\ \downarrow \text{id}_A & & \downarrow \lambda_R^{\text{Rel}^{\text{dbl}}} & & \downarrow \text{id}_B \\ A & \xrightarrow[R]{\quad} & B & & \end{array}$$

of the left unit of  $\text{Rel}^{\text{dbl}}$  at  $R$  is the identity inclusion<sup>1</sup>

$$\begin{array}{ccc} A \times B & \xrightarrow{\chi_B \diamond R} & \{\text{true, false}\} \\ R = \chi_B \diamond R, & \parallel & \equiv & \downarrow \text{id}_{\{\text{true, false}\}} \\ A \times B & \xrightarrow[R]{\quad} & \{\text{true, false}\}. \end{array}$$

<sup>1</sup>As proved in Item 3 of Proposition 8.1.3.1.4.

**00KT 8.3.5.7 The Right Unitors**

**00KU**

**DEFINITION 8.3.5.7.1 ► THE RIGHT UNITORS OF  $\text{Rel}^{\text{dbl}}$**

For each horizontal morphism  $R: A \rightarrow B$  of  $\text{Rel}^{\text{dbl}}$ , the component

$$\rho_R^{\text{Rel}^{\text{dbl}}}: R \odot \mathbb{1}_A \xrightarrow{\sim} R,$$

$$\begin{array}{ccccc} & & \mathbb{1}_A & & \\ A & \xrightarrow{\quad} & A & \xrightarrow{\quad R \quad} & B \\ id_A \downarrow & & \rho_R^{\text{Rel}^{\text{dbl}}} \parallel & & id_B \downarrow \\ A & \xrightarrow{\quad R \quad} & B & & \end{array}$$

of the right unitor of  $\text{Rel}^{\text{dbl}}$  at  $R$  is the identity inclusion<sup>1</sup>

$$\begin{array}{ccc} A \times B & \xrightarrow{R \diamond \chi_A} & \{\text{true, false}\} \\ \parallel & \not\equiv & \downarrow id_{\{\text{true, false}\}} \\ A \times B & \xrightarrow{R} & \{\text{true, false}\}. \end{array}$$

<sup>1</sup>As proved in Item 3 of Proposition 8.1.3.1.4.

**02DZ 8.4 Categories of Relations With Apartness Composition**

**02E0 8.4.1 The Category of Relations With Apartness Composition**

**02E1**

**DEFINITION 8.4.1.1.1 ► THE CATEGORY OF RELATIONS WITH APARTNESS COMPOSITION**

The **category of relations with apartness composition** is the category  $\text{Rel}^\square$  where

- *Objects.* The objects of  $\text{Rel}^\square$  are sets.
- *Morphisms.* For each  $A, B \in \text{Obj}(\text{Sets})$ , we have

$$\text{Rel}^\square(A, B) \stackrel{\text{def}}{=} \text{Rel}(A, B).$$

- *Identities.* For each  $A \in \text{Obj}(\text{Rel}^\square)$ , the unit map

$$\mathbb{1}_A^{\text{Rel}^\square}: \text{pt} \rightarrow \text{Rel}(A, A)$$

of  $\text{Rel}^\square$  at  $A$  is defined by

$$id_A^{\text{Rel}^\square} \stackrel{\text{def}}{=} \nabla_A(-1, -2),$$

where  $\nabla_A(-_1, -_2)$  is the antidiagonal relation of  $A$  of [Example 8.2.1.1.4](#).

- *Composition.* For each  $A, B, C \in \text{Obj}(\text{Rel}^\square)$ , the composition map

$$\circ_{A,B,C}^{\text{Rel}^\square}: \text{Rel}(B, C) \times \text{Rel}(A, B) \rightarrow \text{Rel}(A, C)$$

of  $\text{Rel}^\square$  at  $(A, B, C)$  is defined by

$$S \circ_{A,B,C}^{\text{Rel}^\square} R \stackrel{\text{def}}{=} S \square R$$

for each  $(S, R) \in \text{Rel}(B, C) \times \text{Rel}(A, B)$ , where  $S \diamond R$  is the composition of  $S$  and  $R$  of [Definition 8.1.4.1.1](#).

02E2

#### PROPOSITION 8.4.1.1.2 ► ISOMORPHISM BETWEEN $\text{Rel}$ AND $\text{Rel}^\square$

The functor

$$(-)^c: \text{Rel} \rightarrow \text{Rel}^\square$$

given by the identity on objects and by  $R \mapsto R^c$  on morphisms is an isomorphism of categories.

#### PROOF 8.4.1.1.3 ► PROOF OF PROPOSITION 8.4.1.1.2

By [Item 4 of Proposition 8.1.4.1.3](#), we see that  $(-)^c$  is indeed a functor. By [Item 1 of Proposition 11.6.8.1.3](#), it suffices to show that  $(-)^c$  is bijective on objects (which follows by definition) and fully faithful. Indeed, the map

$$(-)^c: \text{Rel}(A, B) \rightarrow \text{Rel}(A, B)$$

defined by the assignment  $R \mapsto R^c$  is a bijection by [Item 3 of Proposition 4.3.11.1.2](#). Thus  $(-)^c$  is an isomorphism of categories. 

02E3 8.4.2 The 2-Category of Relations With Apartness Composition

02E4

#### DEFINITION 8.4.2.1.1 ► THE 2-CATEGORY OF RELATIONS WITH APARTNESS COMPOSITION

The **2-category of relations with apartness composition** is the locally posetal 2-category **Rel** where

- **Objects.** The objects of **Rel** are sets.
- **Hom-Objects.** For each  $A, B \in \text{Obj}(\text{Sets})$ , we have

$$\begin{aligned} \text{Hom}_{\text{Rel}}(A, B) &\stackrel{\text{def}}{=} \text{Rel}(A, B) \\ &\stackrel{\text{def}}{=} (\text{Rel}(A, B), \subset). \end{aligned}$$

- *Identities.* For each  $A \in \text{Obj}(\mathbf{Rel})$ , the unit map

$$\mathbb{1}_A^{\mathbf{Rel}} : \text{pt} \rightarrow \mathbf{Rel}(A, A)$$

of **Rel** at  $A$  is defined by

$$\text{id}_A^{\mathbf{Rel}} \stackrel{\text{def}}{=} \chi_A(-_1, -_2),$$

where  $\chi_A(-_1, -_2)$  is the characteristic relation of  $A$  of [Example 8.2.1.1.3](#).

- *Composition.* For each  $A, B, C \in \text{Obj}(\mathbf{Rel})$ , the composition map<sup>1</sup>

$$\circ_{A,B,C}^{\mathbf{Rel}} : \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, C)$$

of **Rel** at  $(A, B, C)$  is defined by

$$S \circ_{A,B,C}^{\mathbf{Rel}} R \stackrel{\text{def}}{=} S \diamond R$$

for each  $(S, R) \in \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B)$ , where  $S \diamond R$  is the composition of  $S$  and  $R$  of [Definition 8.1.3.1.1](#).

<sup>1</sup>That this is indeed a morphism of posets is proven in ?? of [Proposition 8.1.4.1.3](#).

02E5

### PROPOSITION 8.4.2.1.2 ► 2-ISOMORPHISM BETWEEN **Rel** AND **Rel**<sup>□,co</sup>

The functor

$$(-)^c : \mathbf{Rel} \rightarrow \mathbf{Rel}^{\square, \text{co}}$$

given by the identity on objects and by  $R \mapsto R^c$  on 1-morphisms is a 2-isomorphism of 2-categories.

### PROOF 8.4.2.1.3 ► PROOF OF PROPOSITION 8.4.2.1.2

By [Item 4 of Proposition 8.1.4.1.3](#), we see that  $(-)^c$  is indeed a functor.

By [Item 1 of Proposition 4.3.11.1.2](#), it is also a 2-functor.

By ??, it suffices to show that  $(-)^c$  is:

- Bijective on objects, which follows by definition.
- Bijective on 1-morphisms, which was shown in [Definition 8.4.1.1.1](#).
- Bijective on 2-morphisms, which follows from [Item 1 of Proposition 4.3.11.1.2](#).

Thus  $(-)^c$  is indeed a 2-isomorphism of categories. 

**02E6 8.4.3 The Linear Bicategory of Relations**

02E7

**DEFINITION 8.4.3.1.1 ► THE LINEAR BICATEGORY OF RELATIONS**

The **linear bicategory of relations** is the linear bicategory consisting of:

- *The Underlying Bicategory I.* The bicategory Rel of [Definition 8.3.4.1.1](#).
- *The Underlying Bicategory II.* The bicategory Rel of [Definition 8.4.2.1.1](#).
- *Linear Distributors.* The inclusions

$$\delta_{R,S,T}^\ell : T \diamond (S \square R) \hookrightarrow (T \diamond S) \square R,$$

$$\delta_{R,S,T}^r : (T \square S) \diamond R \hookrightarrow T \square (S \diamond R)$$

of Item 5 of [Proposition 8.1.4.1.3](#).

**PROOF 8.4.3.1.2 ► PROOF OF THE CLAIMS IN DEFINITION 8.4.3.1.1**

Since Rel and Rel<sup>□</sup> are locally posetal, the commutativity of the coherence conditions for linear bicategories follows automatically (?? of ??).



**02E8 8.4.4 Other Categorical Structures With Apartness Composition**

02E9

**REMARK 8.4.4.1.1 ► OTHER CATEGORICAL STRUCTURES WITH APARTNESS COMPOSITION**

It seems apartness composition fails to form the following categorical structures:

02EA

- *Monoidal Category With Products.* Products don't seem to endow Rel<sup>□</sup> with a monoidal structure.

02EB

- *Monoidal Category With Coproducts.* Coproducts also don't seem to endow Rel<sup>□</sup> with a monoidal structure.

02EC

- *Double Categorical Structure.* It seems the apartness composition of relations doesn't form a double category in a natural<sup>1</sup> way.

<sup>1</sup>I.e. such that the composition of vertical morphisms is the usual composition of functions, as in Sets.

## 00KV 8.5 Properties of the 2-Category of Relations

### 00KW 8.5.1 Self-Duality

00KX

#### PROPOSITION 8.5.1.1.1 ► SELF-DUALITY FOR THE (2-)CATEGORY OF RELATIONS

The 2-/category of relations is self-dual:

00KY

1. *Self-Duality I.* We have an isomorphism

$$\mathbf{Rel}^{\text{op}} \cong \mathbf{Rel}$$

of categories.

00KZ

2. *Self-Duality II.* We have a 2-isomorphism

$$\mathbf{Rel}^{\text{op}} \cong \mathbf{Rel}$$

of 2-categories.

#### PROOF 8.5.1.1.2 ► PROOF OF PROPOSITION 8.5.1.1.1

##### Item 1: Self-Duality I

We claim that the functor

$$(-)^\dagger : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$$

given by the identity on objects and by  $R \mapsto R^\dagger$  on morphisms is an isomorphism of categories. Note that this is indeed a functor by [Items 3 and 6 of Proposition 8.1.5.1.3](#).

By [Item 1 of Proposition 11.6.8.1.3](#), it suffices to show that  $(-)^{\dagger}$  is bijective on objects (which follows by definition) and fully faithful. Indeed, the map

$$(-)^\dagger : \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(B, A)$$

defined by the assignment  $R \mapsto R^\dagger$  is a bijection by [Item 5 of Proposition 8.1.5.1.3](#), showing  $(-)^{\dagger}$  to be fully faithful.

##### Item 2: Self-Duality II

We claim that the 2-functor

$$(-)^\dagger : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$$

given by the identity on objects, by  $R \mapsto R^\dagger$  on morphisms, and by preserving inclusions on 2-morphisms via [Item 1 of Proposition 8.1.5.1.3](#), is an isomorphism of categories.

By ??, it suffices to show that  $(-)^{\dagger}$  is:

- Bijective on objects, which follows by definition.
- Bijective on 1-morphisms, which was shown in [Item 1](#).
- Bijective on 2-morphisms, which follows from [Item 1](#) of [Proposition 8.1.5.1.3](#).

Thus  $(-)^{\dagger}$  is indeed a 2-isomorphism of categories. 

### 00L0 8.5.2 Isomorphisms and Equivalences

Let  $R: A \rightarrow B$  be a relation from  $A$  to  $B$ .

02ED

#### LEMMA 8.5.2.1.1 ► CONDITIONS INVOLVING A RELATION AND ITS CONVERSE I

The conditions below are row-wise equivalent:

CONDITION	INCLUSION
$R$ is functional	$R \diamond R^{\dagger} \subset \Delta_B$
$R$ is total	$\Delta_A \subset R^{\dagger} \diamond R$
$R$ is injective	$R^{\dagger} \diamond R \subset \Delta_A$
$R$ is surjective	$\Delta_B \subset R \diamond R^{\dagger}$

#### PROOF 8.5.2.1.2 ► PROOF OF LEMMA 8.5.2.1.1

Functionality Is Equivalent to  $R \diamond R^{\dagger} \subset \Delta_B$

The condition  $R \diamond R^{\dagger} \subset \Delta_B$  unwinds to

- (★) For each  $b, b' \in B$ , if there exists some  $a \in A$  such that  $b \sim_{R^{\dagger}} a$  and  $a \sim_R b'$ , then  $b = b'$ .

Since  $b \sim_{R^{\dagger}} a$  is the same as  $a \sim_R b$ , the condition says that  $a \sim_R b$  and  $a \sim_R b'$  imply  $b = b'$ . This is precisely the condition for  $R$  to be functional.

Totality Is Equivalent to  $\Delta_A \subset R^{\dagger} \diamond R$

The condition  $\Delta_A \subset R^{\dagger} \diamond R$  unwinds to

- (★) For each  $a, a' \in A$ , if  $a = a'$ , then there exists some  $b \in B$  such that  $a \sim_R b$  and  $b \sim_{R^{\dagger}} a'$ .

Since  $b \sim_{R^\dagger} a'$  is the same as  $a' \sim_R b$ , the condition says that for each  $a \in A$ , there is some  $b \in B$  with  $b \in R(a)$ , so  $R(a) \neq \emptyset$ . This is precisely the condition for  $R$  to be total.

**Injectivity Is Equivalent to  $R^\dagger \diamond R \subset \Delta_A$**

The condition  $R^\dagger \diamond R \subset \Delta_A$  unwinds to

- (★) For each  $a, a' \in A$ , if there exists some  $b \in B$  such that  $a \sim_R b$  and  $b \sim_{R^\dagger} a'$ , then  $a = a'$ .

Since  $b \sim_{R^\dagger} a'$  is the same as  $a' \sim_R b$ , the condition says that for each  $b \in B$ , if  $a \sim_R b$  and  $a' \sim_R b$ , then  $a = a'$ . This is precisely the condition for  $R$  to be injective.

**Surjectivity Is Equivalent to  $\Delta_B \subset R \diamond R^\dagger$**

The condition  $\Delta_B \subset R \diamond R^\dagger$  unwinds to

- (★) For each  $b, b' \in B$ , if  $b = b'$ , then there exists some  $a \in A$  such that  $b \sim_{R^\dagger} a$  and  $a \sim_R b'$ .

Since  $b \sim_{R^\dagger} a$  is the same as  $a \sim_R b$ , the condition says that for each  $b \in B$ , there is some  $a \in A$  with  $b \in R(a)$ , so  $R^{-1}(b) \neq \emptyset$ . This is precisely the condition for  $R$  to be surjective. 

00L1

**PROPOSITION 8.5.2.1.3 ► ISOMORPHISMS AND EQUIVALENCES IN Rel**

The following conditions are equivalent:

00L2

1. The relation  $R: A \rightarrow B$  is an equivalence in **Rel**, i.e.:

- (★) There exists a relation  $R^{-1}: B \rightarrow A$  from  $B$  to  $A$  together with isomorphisms

$$\begin{aligned} R^{-1} \diamond R &\cong \Delta_A, \\ R \diamond R^{-1} &\cong \Delta_B. \end{aligned}$$

00L3

2. The relation  $R: A \rightarrow B$  is an isomorphism in **Rel**, i.e.:

- (★) There exists a relation  $R^{-1}: B \rightarrow A$  from  $B$  to  $A$  such that we have

$$\begin{aligned} R^{-1} \diamond R &= \Delta_A, \\ R \diamond R^{-1} &= \Delta_B. \end{aligned}$$

00L4

3. There exists a bijection  $f: A \xrightarrow{\sim} B$  with  $R = \text{Gr}(f)$ .

## PROOF 8.5.2.1.4 ► PROOF OF PROPOSITION 8.5.2.1.3

We claim that **Items 1** to **3** are indeed equivalent:

- **Item 1**  $\iff$  **Item 2**: This follows from the fact that **Rel** is locally posetal, so that natural isomorphisms and equalities of 1-morphisms in **Rel** coincide.
- **Item 2**  $\implies$  **Item 3**: We proceed in a few steps:
  - First, note that the equalities in **Item 2** imply  $R \dashv R^{-1}$  and thus, by **Proposition 8.5.3.1.1**, there exists a function  $f_R : A \rightarrow B$  associated to  $R$ .
  - By **Lemma 8.5.2.1.1**,  $f_R$  is a bijection.
- **Item 3**  $\implies$  **Item 2**: By **Item 4** of **Proposition 8.2.2.1.2**, we have an adjunction  $\text{Gr}(f) \dashv f^{-1}$ , giving inclusions
 
$$\Delta_A \subset f^{-1} \diamond \text{Gr}(f),$$

$$\text{Gr}(f) \diamond f^{-1} \subset \Delta_B.$$

If  $f$  is bijective, then the reverse inclusions are also true by **Lemma 8.5.2.1.1**.

This finishes the proof. 

## 00L5 8.5.3 Internal Adjunctions

Let  $A$  and  $B$  be sets.

00L6

## PROPOSITION 8.5.3.1.1 ► ADJUNCTIONS IN Rel

We have a natural bijection

$$\left\{ \begin{array}{c} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\} \cong \left\{ \begin{array}{c} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\},$$

with every adjunction in **Rel** being of the form  $\text{Gr}(f) \dashv f^{-1}$  for some function  $f$ .

## PROOF 8.5.3.1.2 ▶ PROOF OF PROPOSITION 8.5.3.1.1

We proceed step by step:

029B

1. *From Adjunctions in **Rel** to Functions.* An adjunction in **Rel** from  $A$  to  $B$  consists of a pair of relations

$$\begin{aligned} R: A &\rightarrow B, \\ S: B &\rightarrow A, \end{aligned}$$

together with inclusions

$$\begin{aligned} \Delta_A &\subset S \diamond R, \\ R \diamond S &\subset \Delta_B. \end{aligned}$$

By Lemma 8.5.2.1.1,  $R$  is total and functional. In particular,  $R(a)$  is a singleton for all  $a \in A$ . Defining  $f_R$  such that  $f_R(a)$  is the unique element of  $R(a)$  then gives us our desired function, forming a map

$$\left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\}.$$

Moreover, by uniqueness of adjoints (??), this implies also that  $S = f^{-1}$ .

029K

2. *From Functions to Adjunctions in **Rel**.* By Item 4 of Proposition 8.2.2.1.2, every function  $f: A \rightarrow B$  gives rise to an adjunction  $\text{Gr}(f) \dashv f^{-1}$  in **Rel**, giving a map

$$\left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\}.$$

029L

3. *Invertibility: From Functions to Adjunctions Back to Functions.* We need to show that starting with a function  $f: A \rightarrow B$ , passing to  $\text{Gr}(f) \dashv f^{-1}$ , and then passing again to a function gives  $f$  again. This follows from the fact that we have  $a \sim_{\text{Gr}(f)} b$  iff  $f(a) = b$ .

029M

4. *Invertibility: From Adjunctions to Functions Back to Adjunctions.* We need to show that, given an adjunction  $R \dashv S$  in **Rel** giving rise to a function  $f_{R,S}: A \rightarrow B$ , we have

$$\begin{aligned} \text{Gr}(f_{R,S}) &= R, \\ f_{R,S}^{-1} &= S. \end{aligned}$$

We check these explicitly:

- $\text{Gr}(f_{R,S}) = R$ . We have

$$\begin{aligned}\text{Gr}(f_{R,S}) &\stackrel{\text{def}}{=} \{(a, f_{R,S}(a)) \in A \times B \mid a \in A\} \\ &\stackrel{\text{def}}{=} \{(a, R(a)) \in A \times B \mid a \in A\} \\ &= R.\end{aligned}$$

- $f_{R,S}^{-1} = S$ . We first claim that, given  $a \in A$  and  $b \in B$ , the following conditions are equivalent:

- We have  $a \sim_R b$ .
- We have  $b \sim_S a$ .

Indeed:

- *If  $a \sim_R b$ , then  $b \sim_S a$ :* We proceed in a few steps.
  - \* Since  $\Delta_A \subset S \diamond R$ , there exists  $k \in B$  such that  $a \sim_R k$  and  $k \sim_S a$ .
  - \* Since  $a \sim_R b$  and  $R$  is functional, we have  $k = b$ .
  - \* Thus  $b \sim_S a$ .
- *If  $b \sim_S a$ , then  $a \sim_R b$ :* We proceed in a few steps.
  - \* First note that, since  $R$  is total, we have  $a \sim_R b'$  for some  $b' \in B$ .
  - \* Since  $R \diamond S \subset \Delta_B$ ,  $b \sim_S a$ , and  $a \sim_R b'$ , we have  $b = b'$ .
  - \* Thus  $a \sim_R b$ .

Having show this, we now have

$$\begin{aligned}f_{R,S}^{-1}(b) &\stackrel{\text{def}}{=} \{a \in A \mid f_{R,S}(a) = b\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid a \sim_R b\} \\ &= \{a \in A \mid b \sim_S a\} \\ &\stackrel{\text{def}}{=} S(b).\end{aligned}$$

for each  $b \in B$ , and thus  $f_{R,S}^{-1} = S$ .

This finishes the proof. 

#### 00L7 8.5.4 Internal Monads

Let  $X$  be a set.

#### 00L8 PROPOSITION 8.5.4.1.1 ► INTERNAL MONADS IN $\mathbf{Rel}$

We have a natural identification<sup>1</sup>

$$\left\{ \begin{array}{l} \text{Monads in} \\ \mathbf{Rel} \text{ on } X \end{array} \right\} \cong \{\text{Preorders on } X\}.$$

<sup>1</sup>See also ?? for an extension of this correspondence to “relative monads in  $\mathbf{Rel}$ ”.

#### PROOF 8.5.4.1.2 ► PROOF OF PROPOSITION 8.5.4.1.1

A monad in  $\mathbf{Rel}$  on  $X$  consists of a relation  $R: X \rightarrow X$  together with maps

$$\begin{aligned} \mu_R: R \diamond R &\subset R, \\ \eta_R: \Delta_X &\subset R \end{aligned}$$

making the diagrams

The diagram consists of four separate commutative squares. Top-left:  $\Delta_X \diamond R \xrightarrow{\eta_R \diamond \text{id}_R} R \diamond R \xrightarrow{\mu_R} R$ . Top-right:  $R \diamond (R \diamond R) \xrightarrow{\alpha_{R,R,R}^{\mathbf{Rel}}} R \diamond R \xrightarrow{\text{id}_R \diamond \mu_R} R$ . Bottom-left:  $R \diamond R \xrightarrow{\mu_R} R$ . Bottom-right:  $R \diamond \Delta_X \xrightarrow{\text{id}_R \diamond \eta_R} R \diamond R \xrightarrow{\mu_R} R$ .

commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic (?? of ??), and hence all that is left is the data of the two maps  $\mu_R$  and  $\eta_R$ , which correspond respectively to the following conditions:

029N

1. For each  $x, z \in X$ , if there exists some  $y \in Y$  such that  $x \sim_R y$  and  $y \sim_R z$ , then  $x \sim_R z$ .
2. For each  $x \in X$ , we have  $x \sim_R x$ .

029P

These are exactly the requirements for  $R$  to be a preorder (??). Conversely, any preorder  $\preceq$  gives rise to a pair of maps  $\mu_\preceq$  and  $\eta_\preceq$ , forming a monad on  $X$ . ■

02EE

#### EXAMPLE 8.5.4.1.3 ► CODENSITY MONADS IN $\mathbf{Rel}$

Let  $R: A \rightarrow B$  be a relation.

02EF

1. The codensity monad  $\text{Ran}_R(R) : B \dashv B$  is given by

$$[\text{Ran}_R(R)](b) = \bigcap_{a \in R^{-1}(b)} R(b)$$

for each  $b \in B$ . Thus, it corresponds to the preorder

$$\preceq_{\text{Ran}_R(R)} : B \times B \rightarrow \{\text{t}, \text{f}\}$$

on  $B$  obtained by declaring  $b \preceq_{\text{Ran}_R(R)} b'$  iff the following equivalent conditions are satisfied:

02EG

(a) For each  $a \in A$ , if  $a \sim_R b$ , then  $a \sim_R b'$ .

02EH

(b) We have  $R^{-1}(b) \subset R^{-1}(b')$ .

02EJ

2. The dual codensity monad  $\text{Rift}_R(R) : A \dashv A$  is given by

$$[\text{Rift}_R(R)](a) = \{a' \in A \mid R(a') \subset R(a)\}$$

for each  $a \in A$ . Thus, it corresponds to the preorder

$$\preceq_{\text{Rift}_R(R)} : A \times A \rightarrow \{\text{t}, \text{f}\}$$

on  $A$  obtained by declaring  $a \preceq_{\text{Rift}_R(R)} a'$  iff the following equivalent conditions are satisfied:

02EK

(a) For each  $a \in A$ , if  $a \sim_R b$ , then  $a' \sim_R b$ .

02EL

(b) We have  $R(a') \subset R(a)$ .

### 00L9 8.5.5 Internal Comonads

Let  $X$  be a set.

00LA

**PROPOSITION 8.5.5.1.1 ► INTERNAL COMONADS IN Rel**

We have a natural identification

$$\left\{ \begin{array}{l} \text{Comonads in} \\ \text{\textbf{Rel} on } X \end{array} \right\} \cong \{\text{Subsets of } X\}.$$

#### PROOF 8.5.5.1.2 ▶ PROOF OF PROPOSITION 8.5.5.1.1

A comonad in **Rel** on  $X$  consists of a relation  $R: X \nrightarrow X$  together with maps

$$\begin{aligned} \Delta_R: R &\subset R \diamond R, \\ \epsilon_R: R &\subset \Delta_X \end{aligned}$$

making the diagrams

commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic (?? of ??), and hence all that is left is the data of the two maps  $\Delta_R$  and  $\epsilon_R$ , which correspond respectively to the following conditions:

029Q

1. For each  $x, y \in X$ , if  $x \sim_R y$ , then there exists some  $k \in X$  such that  $x \sim_R k$  and  $k \sim_R y$ .
2. For each  $x, y \in X$ , if  $x \sim_R y$ , then  $x = y$ .

029R

The second condition implies that  $R \subset \Delta_X$ , so  $R$  must be a subset of  $X$ . Taking  $k = y$  in the first condition above then shows it to be trivially satisfied. Conversely, any subset  $U$  of  $X$  satisfies  $U \subset \Delta_X$ , defining a comonad as above. ■

02EM

#### EXAMPLE 8.5.5.1.3 ▶ DENSITY COMONADS IN **Rel**

Let  $f: A \rightarrow B$  be a function.

02EN

1. The density comonad  $\text{Lan}_f(f) : B \rightarrow B$  is given by

$$[\text{Lan}_f(f)](b) = \bigcup_{a \in f^{-1}(b)} f(a)$$

for each  $b \in B$ . Thus, it corresponds to the image  $\text{Im}(f)$  of  $f$  as a subset of  $B$ .

02EP

2. The dual density comonad  $\text{Lift}_{f^\dagger}(f^\dagger) : A \rightarrow A$  is given by

$$[\text{Lift}_{f^\dagger}(f^\dagger)](b) = \bigcup_{a \in f^{-1}(b)} f(a)$$

for each  $b \in B$ . Thus, it also corresponds to the image  $\text{Im}(f)$  of  $f$  as a subset of  $B$ .

### 02EQ 8.5.6 Modules Over Internal Monads

Let  $A$  be a set.

02ER

#### PROPOSITION 8.5.6.1.1 ► MODULES OVER INTERNAL MONADS IN Rel

Let  $\preceq_A$  be a preorder on  $A$ , viewed also as an internal monad on  $A$  via Proposition 8.5.4.1.1.

02ES

1. *Left Modules.* We have a natural identification

$$\{\text{Left modules over } \preceq_A\} \cong \left\{ \begin{array}{l} \text{Relations } R : B \rightarrow A \text{ such that,} \\ \text{for each } b \in B, \text{ the set } R(b) \text{ is} \\ \text{upward-closed in } A \end{array} \right\}.$$

02ET

2. *Right Modules.* We have a natural identification

$$\{\text{Right modules over } \preceq_A\} \cong \left\{ \begin{array}{l} \text{Relations } R : A \rightarrow B \text{ such that,} \\ \text{for each } b \in B, \text{ the set } R^{-1}(b) \text{ is} \\ \text{downward-closed in } A \end{array} \right\}.$$

02EU

3. *Bimodules.* We have a natural identification

$$\{\text{Bimodules over } \preceq_A\} \cong \left\{ \begin{array}{l} \text{Quadruples } (B, C, R, S) \text{ such that:} \\ \left. \begin{array}{l} 1. \text{ For each } b \in B, \text{ the set } R(b) \text{ is} \\ \text{upward-closed in } A. \\ 2. \text{ For each } c \in C, \text{ the set } S^{-1}(c) \text{ is} \\ \text{downward-closed in } A. \end{array} \right\} \end{array} \right\}.$$

## PROOF 8.5.6.1.2 ► PROOF OF PROPOSITION 8.5.6.1.1

## Item 1: Left Modules

A left module over  $\preceq_A$  in **Rel** consists of a relation  $R: B \rightarrow A$  together with an inclusion

$$\alpha_B: \preceq_A \diamond R \subset R$$

making appropriate diagrams commute. Since **Rel** is locally posetal, however, the commutativity of the diagrams in question is automatic (?? of ??), and hence all that is left is the data of the inclusion  $\alpha_B$ . This corresponds to the following condition:

- (★) For each  $a, a' \in A$ , if there exists some  $b \in B$  such that  $b \sim_R a$  and  $a \preceq_a a'$ , then  $b \sim_R a'$ .

This condition is equivalent to  $R(b)$  being downward-closed for all  $b \in B$ .

## Item 2: Right Modules

The proof is dual to Item 1, and is therefore omitted.

## Item 3: Bimodules

Since **Rel** is locally posetal, the diagram encoding the compatibility conditions for a bimodule commutes automatically (?? of ??), and hence a bimodule is just a left module along with a right module. 

02EV 8.5.7 Comodules Over Internal Comonads

Let  $A$  be a set.

02EW

## PROPOSITION 8.5.7.1.1 ► COMODULES OVER INTERNAL COMONADS IN Rel

Let  $U$  be a subset of  $A$ , viewed also as an internal comonad on  $A$  via Proposition 8.5.5.1.1.

02EX

1. *Left Comodules.* We have a natural identification

$$\{\text{Left comodules over } U\} \cong \left\{ \begin{array}{l} \text{Relations } R: B \rightarrow A \text{ such that,} \\ \text{for each } b \in B, \text{ we have } R(b) \subset U \end{array} \right\}.$$

02EY

2. *Right Comodules.* We have a natural identification

$$\{\text{Right comodules over } U\} \cong \left\{ \begin{array}{l} \text{Relations } R: A \rightarrow B \text{ such that,} \\ \text{for each } b \in B, \text{ we have } R^{-1}(b) \subset U \end{array} \right\}.$$

02EZ

3. *Bicomodules.* We have a natural identification

$$\{\text{Bicomodules over } U\} \cong \left\{ \begin{array}{l} \text{Quadruples } (B, C, R, S) \text{ such that:} \\ \text{1. For each } b \in B, \text{ we have } R(b) \subset U \\ \text{2. For each } c \in C, \text{ we have } S^{-1}(c) \subset U \end{array} \right\}.$$

#### PROOF 8.5.7.1.2 ► PROOF OF PROPOSITION 8.5.7.1.1

##### Item 1: Left Comodules

A left comodule over  $U$  in **Rel** consists of a relation  $R: B \rightarrow A$  together with an inclusion

$$R \subset U \diamond R$$

making appropriate diagrams commute. Since **Rel** is locally posetal, however, the commutativity of the diagrams in question is automatic (?? of ??), and hence all that is left is the data of the inclusion. This corresponds to the following condition:

- (★) For each  $b \in B$ , if  $b \sim_R a$ , then there exists some  $a' \in A$  such that  $b \sim_R a'$  and  $a' \sim_U a$ .

Since  $a' \sim_U a$  is true if  $a = a'$  and  $a \in U$ , this condition ends up being equivalent to  $R(b) \subset U$ .

##### Item 2: Right Comodules

A right comodule over  $U$  in **Rel** consists of a relation  $R: A \rightarrow B$  together with an inclusion

$$R \subset R \diamond U$$

making appropriate diagrams commute. Since **Rel** is locally posetal, however, the commutativity of the diagrams in question is automatic

(?? of ??), and hence all that is left is the data of the inclusion. This corresponds to the following condition:

- (★) For each  $a \in A$ , if  $a \sim_R b$ , then there exists some  $x \in A$  such that  $a \sim_U x$  and  $x \sim_R b$ .

Since  $a \sim_U x$  is true if  $a = x$  and  $a \in U$ , this condition ends up being equivalent to  $R^{-1}(b) \subset U$ .

**Item 3: Bicomodules**

Since **Rel** is locally posetal, the diagram encoding the compatibility conditions for a bimodule commutes automatically (?? of ??), and hence a bicomodule is just a left comodule along with a right comodule. 

### 02F0 8.5.8 Eilenberg–Moore and Kleisli Objects

Let  $X$  be a set.

02F1

**PROPOSITION 8.5.8.1.1 ► EILENBERG–MOORE AND KLEISLI OBJECTS IN Rel**

Let  $R$  be a preorder on  $X$ , viewed as an internal monad on  $X$  via [Proposition 8.5.4.1.1](#).

02F2

1. *Eilenberg–Moore Objects in Rel.* The Eilenberg–Moore object for  $R$  exists iff it is an equivalence relation, in which case it is the quotient  $X/\sim_R$  of  $X$  by  $R$ .

02F3

2. *Kleisli Objects in Rel.* [...]

**PROOF 8.5.8.1.2 ► PROOF OF PROPOSITION 8.5.8.1.1**

Omitted. 

### 00LB 8.5.9 Co/Monoids

00LC

**REMARK 8.5.9.1.1 ► Co/MONOIDS IN Rel**

The monoids in **Rel** with respect to the Cartesian monoidal structure of [Proposition 8.3.3.8.1](#) are called *hypermonoids*, and their theory is explored in ?. Similarly, the comonoids in **Rel** are called *hypercomonoids*, and they are defined and studied in ?.

**00LD 8.5.10 Monomorphisms and 2-Categorical Monomorphisms**

02PS

## EXPLANATION 8.5.10.1.1 ► MONOMORPHISMS IN Rel


In this section, we characterise:

- The 1-categorical monomorphisms in Rel, following ??.
- The 2-categorical monomorphisms in **Rel**, following [Section 14.1](#).

More specifically:

- [Proposition 8.5.10.1.2](#) gives *conceptual* characterisations of the monomorphisms in Rel.
- [Proposition 8.5.10.1.4](#) gives *point-set* characterisations of the monomorphisms in Rel.
- [Propositions 8.5.10.1.13](#) and [8.5.10.1.15](#) characterise the 2-categorical monomorphisms in **Rel**.<sup>1</sup>

Essentially, a monomorphism  $R: A \rightarrow B$  in Rel is a relation that is total and injective. Therefore, it looks like this:



In particular:

02PT

1.  $R$  should contain an injection  $f: A \hookrightarrow B$  embedding a copy of  $A$  into  $B$ .

02PU

2.  $R$  can be non-functional, mapping elements of  $A$  to multiple elements of  $B$  (but not to more than one in  $\text{Im}(f)$ ).

02PV

3.  $R$  doesn't need to be surjective, so  $B$  can have elements that aren't in the image of  $R$ .

<sup>1</sup>*Summary:* As it turns out, every 1-morphism in **Rel** is representably faithful and most other notions of 2-categorical monomorphism agree with the usual (1-categorical) notion of monomorphism.

00LF

**PROPOSITION 8.5.10.1.2 ► CHARACTERISATIONS OF MONOMORPHISMS IN Rel I**

Let  $R: A \rightarrow B$  be a relation. The following conditions are equivalent:<sup>1</sup>

02PW

1. The relation  $R$  is a monomorphism in Rel.

02PX

2. The relation  $R$  is total and injective.

02PY

3. The direct image function

$$R_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to  $R$  is injective.

02PZ

4. The codirect image function

$$R_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to  $R$  is injective.

02Q0

5. The direct image functor

$$R_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

associated to  $R$  is full.

02Q1

6. The codirect image functor

$$R_*: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

associated to  $R$  is full.

02Q2

7. For each pair of relations  $S, T: X \rightrightarrows A$ , the following condition is satisfied:

(★) If  $R \diamond S \subset R \diamond T$ , then  $S \subset T$ .

02Q3

8. There exists an injective function  $f: A \hookrightarrow B$  satisfying the following conditions:<sup>2</sup>

02Q4

(a) We have  $\text{Gr}(f) \subset R$ .<sup>3</sup>

02Q5

(b) The diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & \mathcal{P}(B) \\ & \searrow \chi_A & \downarrow R^{-1} \\ & \mathcal{P}(A) & \end{array}$$

commutes.<sup>4</sup>

02Q6

9. The diagram

$$\begin{array}{ccc} A & \xrightarrow{R} & \mathcal{P}(B) \\ & \searrow \chi_A & \downarrow R^{-1} \\ & & \mathcal{P}(A) \end{array}$$

commutes. In other words, we have

$$R^{-1}(R(a)) = \{a\}$$

for each  $a \in A$ .

02S7

10. We have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_!} & \mathcal{P}(B) \\ R_{-1} \circ R_! = \text{id}_{\mathcal{P}(A)} & \swarrow & \downarrow R_{-1} \\ & & \mathcal{P}(A). \end{array}$$

In other words, we have

$$U = \underbrace{\{a \in A \mid R(a) \subset R(U)\}}_{=R_{-1}(R_!(U))}$$

for each  $U \in \mathcal{P}(A)$ .

02S8

11. We have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_!} & \mathcal{P}(B) \\ R^{-1} \circ R_! = \text{id}_{\mathcal{P}(A)} & \swarrow & \downarrow R^{-1} \\ & & \mathcal{P}(A). \end{array}$$

In other words, we have

$$U = \underbrace{\{a \in A \mid R(a) \cap R(U) \neq \emptyset\}}_{=R^{-1}(R_!(U))}$$

for each  $U \in \mathcal{P}(A)$ .

02S9

12. We have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_*} & \mathcal{P}(B) \\ R^{-1} \circ R_* = \text{id}_{\mathcal{P}(A)} & \swarrow & \downarrow R^{-1} \\ & & \mathcal{P}(A). \end{array}$$

In other words, we have

$$U = \underbrace{\left\{ a \in A \mid \begin{array}{l} \text{there exists some } b \in R(a) \\ \text{such that we have } R^{-1}(b) \subset U \end{array} \right\}}_{=R^{-1}(R_*(U))}$$

for each  $U \in \mathcal{P}(A)$ .

02SA

13. We have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_*} & \mathcal{P}(B) \\ R_{-1} \circ R_* = \text{id}_{\mathcal{P}(A)} & \swarrow & \downarrow R_{-1} \\ & & \mathcal{P}(A). \end{array}$$

In other words, we have

$$U = \underbrace{\left\{ a \in A \mid R^{-1}(R(a)) \subset U \right\}}_{=R_{-1}(R_*(U))}$$

for each  $U \in \mathcal{P}(A)$ .<sup>1</sup>Items 3 to 6 unwind respectively to the following statements:

- For each  $U, V \in \mathcal{P}(A)$ , if  $R_!(U) = R_!(V)$ , then  $U = V$ .
- For each  $U, V \in \mathcal{P}(A)$ , if  $R_*(U) = R_*(V)$ , then  $U = V$ .
- For each  $U, V \in \mathcal{P}(A)$ , if  $R_!(U) \subset R_!(V)$ , then  $U \subset V$ .
- For each  $U, V \in \mathcal{P}(A)$ , if  $R_*(U) \subset R_*(V)$ , then  $U \subset V$ .

<sup>2</sup>We are assuming the axiom of choice for this item (Item 8).<sup>3</sup>In other words, for each  $a \in A$ , we have  $f(a) \in R(a)$ .<sup>4</sup>In other words, for each  $a \in A$ , we have  $R^{-1}(f(a)) = \{a\}$ .

**PROOF 8.5.10.1.3 ► PROOF OF PROPOSITION 8.5.10.1.2**

We will prove this by showing:

- Step 1: Item 1  $\iff$  Item 2.
- Step 2: Item 3  $\iff$  Item 4 and Item 4  $\iff$  Item 6.
- Step 3: Item 1  $\iff$  Item 3.
- Step 4: Item 3  $\iff$  Item 5.
- Step 5: Item 5  $\iff$  Item 7.
- Step 6: Item 1  $\iff$  Item 8.
- Step 7: Item 1  $\iff$  Item 9.
- Step 8: Item 1  $\iff$  Item 10.
- Step 9: Item 1  $\iff$  Item 11.
- Step 10: Item 1  $\iff$  Item 12.
- Step 11: Item 1  $\iff$  Item 13.

**Step 1: Item 1  $\iff$  Item 2**

We defer this proof to [Corollary 8.5.10.1.8](#).

**Step 2: Item 3  $\iff$  Item 4 and Item 4  $\iff$  Item 6**

This follows from [Item 7 of Proposition 8.7.1.1.4](#).

**Step 3: First Proof of Item 1  $\iff$  Item 3**

We claim that [Items 1](#) and [3](#) are equivalent:

- [Item 1  \$\implies\$  Item 3](#): Let  $U, V \in \mathcal{P}(A)$  and consider the diagram

$$\begin{array}{ccc} \text{pt} & \xrightarrow[U]{\quad} & A \xrightarrow[R]{\quad} B. \\ & \parallel & \\ & V & \end{array}$$

By [Remark 8.7.1.1.3](#), we have

$$\begin{aligned} R_!(U) &= R \diamond U, \\ R_!(V) &= R \diamond V. \end{aligned}$$

Now, if  $R \diamond U = R \diamond V$ , i.e.  $R_!(U) = R_!(V)$ , then  $U = V$  since  $R$  is assumed to be a monomorphism, showing  $R_!$  to be injective.

- *Item 3  $\implies$  Item 1*: Conversely, suppose that  $R_!$  is injective, consider the diagram

$$X \xrightarrow[S]{\quad\quad} A \xrightarrow[R]{\quad\quad} B,$$

$\begin{array}{c} S \\ \parallel \\ T \end{array}$

and suppose that  $R \diamond S = R \diamond T$ . Note that, since  $R_!$  is injective, given a diagram of the form

$$\text{pt} \xrightarrow[U]{\quad\quad} A \xrightarrow[R]{\quad\quad} B,$$

$\begin{array}{c} U \\ \parallel \\ V \end{array}$

if  $R_!(U) = R \diamond U = R \diamond V = R_!(V)$ , then  $U = V$ . In particular, for each  $x \in X$ , we may consider the diagram

$$\text{pt} \xrightarrow[x]{\quad\quad} X \xrightarrow[S]{\quad\quad} A \xrightarrow[R]{\quad\quad} B,$$

$\begin{array}{c} [x] \\ \parallel \\ T \end{array}$

where we have  $R \diamond S \diamond [x] = R \diamond T \diamond [x]$ , implying that we have

$$S(x) = S \diamond [x] = T \diamond [x] = T(x)$$

for each  $x \in X$ . Thus  $S = T$  and  $R$  is a monomorphism.

#### Step 3.5: Second Proof of Item 1 $\iff$ Item 3

A more abstract proof can also be given, following [MSE 350788]:

- *Proposition 8.5.10.1.2  $\implies$  Proposition 8.5.10.1.4*: Assume that  $R$  is a monomorphism.

- We first notice that the functor  $\text{Rel}(\text{pt}, -) : \text{Rel} \rightarrow \text{Sets}$  maps  $R$  to  $R_!$  by Remark 8.7.1.1.3.
- Since  $\text{Rel}(\text{pt}, -)$  preserves all limits by ?? of ??, it follows by ?? of ?? that  $\text{Rel}(\text{pt}, -)$  also preserves monomorphisms.
- Since  $R$  is a monomorphism and  $\text{Rel}(\text{pt}, -)$  maps  $R$  to  $R_!$ , it follows that  $R_!$  is also a monomorphism.
- Since the monomorphisms in  $\text{Sets}$  are precisely the injections (?? of ??), it follows that  $R_!$  is injective.

- *Proposition 8.5.10.1.4  $\implies$  Proposition 8.5.10.1.2*: Assume that  $R_!$  is injective.

- We first notice that the functor  $\text{Rel}(\text{pt}, -) : \text{Rel} \rightarrow \text{Sets}$  maps  $R$  to  $R_!$  by Remark 8.7.1.1.3.

- Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that  $R_!$  is a monomorphism.
- Since  $\text{Rel}(\text{pt}, -)$  is faithful, it follows by ?? of ?? that  $\text{Rel}(\text{pt}, -)$  reflects monomorphisms.
- Since  $R_!$  is a monomorphism and  $\text{Rel}(\text{pt}, -)$  maps  $R$  to  $R_!$ , it follows that  $R$  is also a monomorphism.

**Step 4: Item 3  $\iff$  Item 5**

We claim that **Items 3** and **5** are equivalent:

- **Item 3  $\implies$  Item 5:** We proceed in a few steps:

- Let  $U, V \in \mathcal{P}(A)$  such that  $R_!(U) \subset R_!(V)$ , assume  $R_!$  to be injective, and consider the set  $U \cup V$ .
- Since  $R_!(U) \subset R_!(V)$ , we have

$$\begin{aligned} R_!(U \cup V) &= R_!(U) \cup R_!(V) \\ &= R_!(V), \end{aligned}$$

where we have used **Item 5** of **Proposition 8.7.1.1.4** for the first equality.

- Since  $R_!$  is injective, this implies  $U \cup V = V$ .
- Thus  $U \subset V$ , as we wished to show.

- **Item 3  $\implies$  Item 5:** We proceed in a few steps:

- Suppose **Item 5** holds, and let  $U, V \in \mathcal{P}(A)$  such that  $R_!(U) = R_!(V)$ .
- Since  $R_!(U) = R_!(V)$ , we have  $R_!(U) \subset R_!(V)$  and  $R_!(V) \subset R_!(U)$ .
- By assumption, this implies  $U \subset V$  and  $V \subset U$ .
- Thus  $U = V$ , showing  $R_!$  to be injective.

**Step 5: Item 5  $\iff$  Item 7**

We claim that **Items 5** and **7** are equivalent:

- **Item 5  $\implies$  Item 7:** Let  $U, V \in \mathcal{P}(A)$  and consider the diagram

$$\text{pt} \quad \begin{array}{c} \xrightarrow{\hspace{1cm} U \hspace{1cm}} \\ \xrightleftharpoons[\hspace{1cm} V \hspace{1cm}]{} \\ \xrightarrow{\hspace{1cm} R \hspace{1cm}} \end{array} A \longrightarrow B.$$

By Remark 8.7.1.1.3, we have

$$R_!(U) = R \diamond U,$$

$$R_!(V) = R \diamond V.$$

Now, if  $R \diamond U \subset R \diamond V$ , then  $R_!(U) \subset R_!(V)$ . By assumption, we then have  $U \subset V$ .

- *Item 7  $\implies$  Item 5:* Consider the diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad S \quad} & A & \xrightarrow{\quad R \quad} & B, \\ & \downarrow T & & & \end{array}$$

and suppose that  $R \diamond S \subset R \diamond T$ . Note that, by assumption, given a diagram of the form

$$\begin{array}{ccc} \text{pt} & \xrightarrow{\quad U \quad} & A & \xrightarrow{\quad R \quad} & B, \\ & \downarrow V & & & \end{array}$$

if  $R_!(U) = R \diamond U \subset R \diamond V = R_!(V)$ , then  $U \subset V$ . In particular, for each  $x \in X$ , we may consider the diagram

$$\begin{array}{ccccc} & [x] & & & \\ \text{pt} & \xrightarrow{\quad} & X & \xrightarrow{\quad S \quad} & A & \xrightarrow{\quad R \quad} & B, \\ & & & \downarrow T & & & \end{array}$$

for which we have  $R \diamond S \diamond [x] \subset R \diamond T \diamond [x]$ , implying that we have

$$S(x) = S \diamond [x] \subset T \diamond [x] = T(x)$$

for each  $x \in X$ , implying  $S \subset T$ .

This finishes the proof.

**Step 6: Item 1  $\iff$  Item 8**

We defer this proof to Corollary 8.5.10.1.6.

**Step 7: Item 1  $\iff$  Item 9**

We defer this proof to Corollary 8.5.10.1.10.

**Step 8: Item 1  $\iff$  Item 10**

We defer this proof to Corollary 8.5.10.1.6.

**Step 9: Item 1  $\iff$  Item 11**

We defer this proof to Corollary 8.5.10.1.10.

**Step 10: Item 1  $\iff$  Item 12**

We defer this proof to Corollary 8.5.10.1.10.

**Step 11: Item 1  $\iff$  Item 13**

We defer this proof to Corollary 8.5.10.1.10. 

00LG

**PROPOSITION 8.5.10.1.4 ► CHARACTERISATIONS OF MONOMORPHISMS IN Rel II**

02Q7

Let  $R: A \rightarrow B$  be a relation. The following conditions are equivalent:

02Q8

1. The relation  $R$  is a monomorphism in Rel.
2. For each  $a \in A$  and each  $U \in \mathcal{P}(A)$ , if  $R(a) \subset R(U)$ , then  $a \in U$ .

02Q9

3. For each  $a \in A$ , there exists some  $b \in B$  such that  $R^{-1}(b) = \{a\}$ .

**PROOF 8.5.10.1.5 ► PROOF OF PROPOSITION 8.5.10.1.4**

We will prove this by showing:

- Step 1: Item 1  $\implies$  Item 2.
- Step 2: Item 2  $\implies$  Item 3.
- Step 3: Item 3  $\implies$  Item 1.

**Step 1: Item 1  $\implies$  Item 2**

We proceed in a few steps:

- If  $R$  is a monomorphism, then, by Item 3 of Proposition 8.5.10.1.2, the functor

$$R_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

is full.

- As a result, given  $a \in A$  and  $U \in \mathcal{P}(A)$  such that  $R(a) \subset R(U)$ , it follows that  $\{a\} \subset U$ .
- Thus, we have  $a \in U$ .

**Step 2: Item 2  $\implies$  Item 3**

We proceed in a few steps:

- Let  $a \in A$  and consider the subset  $U = A \setminus \{a\}$ .
- Since  $a \notin U$ , we have  $R(a) \not\subset R(U)$  by the contrapositive of Item 2.
- As a result, there must exist some  $b \in R(a)$  with  $b \notin R(U)$ .
- In particular, we have  $a \in R^{-1}(b)$ .
- Moreover, the condition  $b \notin R(U) = R(A \setminus \{a\})$  means that, if  $a' \in A \setminus \{a\}$ , then  $a' \notin R^{-1}(b)$ .

- Thus  $R^{-1}(b) = \{a\}$ .

**Step 3: Item 3  $\implies$  Item 1**

We proceed in a few steps:

- By the equivalence between **Items 1** and **5** of **Proposition 8.5.10.1.2**, to show that  $R$  is a monomorphism it suffices to prove that, for each  $U, V \in \mathcal{P}(A)$ , if  $R(U) \subset R(V)$ , then  $U \subset V$ .
- So let  $u \in U$  and assume  $R(U) \subset R(V)$ .
- By assumption, there exists some  $b \in B$  with  $R^{-1}(b) = \{u\}$ .
- In particular,  $b \in R(U)$ .
- Since  $R(U) \subset R(V)$ , we also have  $b \in R(V)$ .
- Thus, there exists some  $v \in V$  with  $b \in R(v)$ .
- However,  $R^{-1}(b) = \{u\}$ , so we must in fact have  $v = u$ .
- Therefore  $u \in V$ , showing that  $U \subset V$ .

This finishes the proof. 

00LH

**COROLLARY 8.5.10.1.6 ▶ CHARACTERISATIONS OF MONOMORPHISMS IN Rel III**

**Items 1, 8 and 10 of Proposition 8.5.10.1.2** are indeed equivalent.

**PROOF 8.5.10.1.7 ▶ PROOF OF COROLLARY 8.5.10.1.6**

**Item 1  $\iff$  Item 8**

We claim that **Item 3** of **Proposition 8.5.10.1.4** is equivalent to **Item 8** of **Proposition 8.5.10.1.2**:

- **Item 3 of Proposition 8.5.10.1.4  $\implies$  Item 8:** By assumption, given  $a \in A$ , there exists some  $b \in B$  such that  $R^{-1}(b) = \{a\}$ . Invoking the axiom of choice, we may pick one such  $b$  for each  $a \in A$ , giving us our desired function  $f: A \rightarrow B$ . All the requirements listed in **Item 8 of Proposition 8.5.10.1.2** then follow by construction.
- **Item 8  $\implies$  Item 3 of Proposition 8.5.10.1.4:** Given  $a \in A$ , we may pick  $b = f(a)$ , in which case  $R^{-1}(f(a))$  will be equal to  $\{a\}$  by assumption.

By [Proposition 8.5.10.1.4](#), Item 3 of [Proposition 8.5.10.1.4](#) is equivalent to Item 1 of [Proposition 8.5.10.1.4](#). Since Item 1 of [Proposition 8.5.10.1.4](#) is exactly the same condition as Item 1 of [Proposition 8.5.10.1.2](#), the result follows.

Item 1  $\iff$  Item 10

Indeed, we have

$$\begin{aligned} [R_{-1} \circ R_!](U) &\stackrel{\text{def}}{=} R_{-1}(R_!(U)) \\ &\stackrel{\text{def}}{=} \{a \in A \mid R(a) \subset R(U)\} \end{aligned}$$

for each  $U \in \mathcal{P}(A)$ . As a result, the condition  $R_{-1} \circ R_! = \text{id}_{\mathcal{P}(A)}$  becomes

$$\{a \in A \mid R(a) \subset R(U)\} = U,$$

which holds precisely when Item 2 of [Proposition 8.5.10.1.4](#) does. By [Proposition 8.5.10.1.4](#), that in turn holds precisely if Item 1 of [Proposition 8.5.10.1.4](#) holds. Since Item 1 of [Proposition 8.5.10.1.4](#) is exactly the same condition as Item 1 of [Proposition 8.5.10.1.2](#), the result follows. □

02QA

#### COROLLARY 8.5.10.1.8 ▶ CHARACTERISATIONS OF MONOMORPHISMS IN Rel IV

[Items 1 and 2 of Proposition 8.5.10.1.2](#) are indeed equivalent.<sup>1</sup>

<sup>1</sup>I.e. a relation is a monomorphism in Rel iff it is total and injective.

#### PROOF 8.5.10.1.9 ▶ PROOF OF COROLLARY 8.5.10.1.8

We claim that [Items 1 and 2 of Proposition 8.5.10.1.2](#) are indeed equivalent:

- [Item 1  \$\implies\$  Item 2](#): First, note that  $R$  is total by [Item 3 of Proposition 8.5.10.1.4](#). Next, let  $a, a' \in A$  such that  $a \sim_R b$  and  $a' \sim_R b$  for some  $b \in B$  and consider the diagram

$$\begin{array}{ccc} \text{pt} & \xrightarrow{\quad [a] \quad} & A \xrightarrow{\quad R \quad} B \\ & \parallel & \\ & \xrightarrow{\quad [a'] \quad} & \end{array}$$

Then:

- Since  $\star \sim_{[a]} a$  and  $a \sim_R b$ , we have  $\star \sim_{R \circ [a]} b$ .
- Similarly,  $\star \sim_{R \circ [a']} b$ .

- Thus  $R \diamond [a] = R \diamond [a']$ .
- Since  $R$  is a monomorphism, we have  $[a] = [a']$ , so  $a = a'$ .
- *Item 2  $\implies$  Item 1:* Consider the diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad S \quad} & A & \xrightarrow{\quad R \quad} & B, \\ & \downarrow T & & & \end{array}$$

where  $R \diamond S = R \diamond T$ , and let  $x \in X$  and  $a \in A$  such that  $x \sim_S a$ .

- Since  $R$  is total and  $a \in A$ , there exists some  $b \in B$  such that  $a \sim_R b$ .
- In this case, we have  $x \sim_{R \diamond S} b$ , and since  $R \diamond S = R \diamond T$ , we have also  $x \sim_{R \diamond T} b$ .
- Thus there must exist some  $a' \in A$  such that  $x \sim_T a'$  and  $a' \sim_R b$ .
- However, since  $a \sim_R b$  and  $a' \sim_R b$ , we must have  $a = a'$  by condition  $(\star)$ .
- Thus  $x \sim_T a$  as well.
- A similar argument shows that if  $x \sim_T a$ , then  $x \sim_S a$ .
- Thus  $S = T$ , showing  $R$  to be a monomorphism.

This finishes the proof. □

02SB

#### COROLLARY 8.5.10.1.10 ▶ CHARACTERISATIONS OF MONOMORPHISMS IN Rel V

Items 1, 9 and 11 to 13 of Proposition 8.5.10.1.2 are indeed equivalent.

#### PROOF 8.5.10.1.11 ▶ PROOF OF COROLLARY 8.5.10.1.10

We will prove this by showing:

- Step 1: *Item 1  $\implies$  Item 11.*
- Step 2: *Item 11  $\implies$  Item 1.*
- Step 3: *Item 1  $\implies$  Item 12.*
- Step 4: *Item 12  $\implies$  Item 1.*
- Step 5: *Item 9  $\iff$  Item 13.*

- Step 6: Item 9  $\iff$  Item 2.

**Step 1: Item 1  $\implies$  Item 11**

Assume that  $R$  is a monomorphism, which is equivalent to  $R$  being total and injective by [Proposition 8.5.10.1.2](#). Let  $S(U) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \cap R(U) \neq \emptyset\}$ . We need to show that  $U = S(U)$  for any  $U \in \mathcal{P}(A)$  by proving double inclusion.

- $U \subset S(U)$ : Let  $u \in U$ .
  - Since  $R$  is total, we have  $R(u) \neq \emptyset$ .
  - By definition,  $R(U) = \bigcup_{x \in U} R(x)$ , so  $R(u) \subset R(U)$ .
  - Therefore,  $R(u) \cap R(U) = R(u)$ .
  - Since  $R(u) \neq \emptyset$ , we have  $R(u) \cap R(U) \neq \emptyset$ .
  - By the definition of  $S(U)$ , it follows that  $u \in S(U)$ .
- $S(U) \subset U$ : Let  $a \in S(U)$ .
  - By assumption,  $R(a) \cap R(U) \neq \emptyset$ .
  - This means  $R(a) \cap \bigcup_{u \in U} R(u) \neq \emptyset$ .
  - Using the distributivity of intersection over union, this is equivalent to  $\bigcup_{u \in U} (R(a) \cap R(u)) \neq \emptyset$ .
  - For this union of sets to be non-empty, at least one of the sets in the union must be non-empty. Thus, there must exist some  $u \in U$  such that  $R(a) \cap R(u) \neq \emptyset$ .
  - Since  $R$  is injective, the images of distinct elements are disjoint. For the intersection  $R(a) \cap R(u)$  to be non-empty, we must therefore have  $a = u$ .
  - Since  $u \in U$ , it follows that  $a \in U$ .

As both inclusions hold, we conclude that  $U = S(U)$ .

**Step 2: Item 11  $\implies$  Item 1**

Assume that for every  $U \in \mathcal{P}(A)$ , we have  $U = \{a \in A \mid R(a) \cap R(U) \neq \emptyset\}$ . We must show that  $R$  is both total and injective.

- *Totality*: Let  $a \in A$ . We must show that  $R(a) \neq \emptyset$ .
  - Consider the singleton set  $U = \{a\}$ .
  - By assumption,  $U = \{x \in A \mid R(x) \cap R(U) \neq \emptyset\}$ .
  - Since  $a \in U$ , we must have  $R(a) \cap R(U) \neq \emptyset$ .

- Substituting  $U = \{a\}$ , we get  $R(a) \cap R(\{a\}) \neq \emptyset$ .
- Since  $R(\{a\}) = R(a)$ , this simplifies to  $R(a) \cap R(a) = R(a) \neq \emptyset$ .
- Thus  $R(a) \neq \emptyset$  for all  $a \in A$ , showing  $R$  to be total.
- *Injectivity:* Let  $a, a' \in A$  such that  $a \neq a'$ . We must show that  $R(a) \cap R(a') = \emptyset$ .
  - Consider the singleton set  $U = \{a\}$ .
  - By assumption,  $U = \{x \in A \mid R(x) \cap R(U) \neq \emptyset\}$ .
  - Since  $a \neq a'$ , we have  $a' \notin U$ .
  - Therefore,  $a'$  cannot satisfy the membership condition for  $U$ . This means  $R(a') \cap R(U) = \emptyset$ .
  - Substituting  $U = \{a\}$ , we get  $R(a') \cap R(\{a\}) = \emptyset$ , which simplifies to  $R(a') \cap R(a) = \emptyset$ .
  - As this holds for any pair of distinct elements, the relation  $R$  is injective.

This completes the proof.

**Step 3: Item 1  $\implies$  Item 12**

We proceed by taking a specific choice of subset  $U$ :

- Let  $a$  be an arbitrary element of  $A$ . By our assumption, the condition  $R^{-1}(R_*(U)) = U$  must hold for the singleton set  $U = \{a\}$ .
- From  $R^{-1}(R_*(\{a\})) = \{a\}$ , it follows that  $a \in R^{-1}(R_*(\{a\}))$ .
- This means there must exist some  $b \in R(a)$  such that  $R^{-1}(b) \subset \{a\}$ .
- The condition  $b \in R(a)$  implies that  $a \in R^{-1}(b)$ . Therefore,  $R^{-1}(b)$  is a non-empty subset of  $\{a\}$ .
- The only non-empty subset of  $\{a\}$  is  $\{a\}$  itself.
- Thus, we must have  $R^{-1}(b) = \{a\}$ .

**Step 4: Item 12  $\implies$  Item 1**

By **Item 2 of Proposition 8.5.10.1.4**, for each  $a \in A$ , there exists some  $b \in B$  such that  $R^{-1}(b) = \{a\}$ . We need to show that  $R^{-1}(R_*(U)) = U$  for any  $U \in \mathcal{P}(A)$ , which requires proving two set inclusions.

- $R^{-1}(R_*(U)) \subset U$ : We proceed in a few steps:

- Let  $a \in R^{-1}(R_*(U))$ .
- By definition, there exists some  $b \in R(a)$  such that  $R^{-1}(b) \subset U$ .
- Since  $b \in R(a)$  implies  $a \in R^{-1}(b)$ , it follows immediately that  $a \in U$ .
- Thus,  $R^{-1}(R_*(U)) \subset U$ .
- $U \subset R^{-1}(R_*(U))$ : Let  $a \in U$ . By assumption, there exists an element  $b \in B$  such that  $R^{-1}(b) = \{a\}$ . Thus  $R^{-1}(b) \subset U$ , so  $a \in R^{-1}(R_*(U))$ .

Combining both inclusions gives  $R^{-1}(R_*(U)) = U$ .

Step 5: Item 9  $\iff$  Item 13

We claim that **Items 9** and **13** are equivalent:

- **Item 13  $\implies$  Item 9**: Let  $a \in A$ .
  - First, let  $U = \{a\}$ . By assumption, we have
 
$$\{a\} = \{a' \in A \mid R^{-1}(R(a')) \subset \{a\}\}.$$
 Since  $a$  is in the set on the left-hand side, it must also be in the set on the right-hand side. Thus  $R^{-1}(R(a')) \subset \{a\}$  must be true.
  - Next, consider the complement  $U = A \setminus \{a\}$ . By assumption, we have
 
$$A \setminus \{a\} = \{a' \in A \mid R^{-1}(R(a')) \subset A \setminus \{a\}\}$$
 Since  $a$  is not in the set on the left-hand side, it cannot be in the set on the right-hand side. Thus  $R^{-1}(R(a)) \not\subset A \setminus \{a\}$ .

- The statement  $R^{-1}(R(a)) \not\subset A \setminus \{a\}$  implies that there exists an element  $x \in R^{-1}(R(a))$  such that  $x \notin A \setminus \{a\}$ . The only such element is  $a$ , so we must have  $a \in R^{-1}(R(a))$ .
- Combining these two results, namely  $R^{-1}(R(a)) \subset \{a\}$  and  $a \in R^{-1}(R(a))$ , we conclude that  $R^{-1}(R(a)) = \{a\}$ , as we wished to show.

- **Item 9  $\implies$  Item 13**: We have

$$\begin{aligned} R^{-1}(R_*(U)) &= \{a \in A \mid R^{-1}(R(a)) \subset U\} \\ &= \{a \in A \mid \{a\} \subset U\} \\ &= U. \end{aligned}$$

Step 6: Item 9  $\iff$  Item 2

We claim that **Items 2** and **9** are equivalent:

- **Item 9  $\implies$  Item 2:** By definition,

$$R^{-1}(R(a)) = \{x \in A \mid R(x) \cap R(a) \neq \emptyset\}.$$

The condition  $R^{-1}(R(a)) = \{a\}$  implies two facts:

- The element  $a$  must belong to the set  $\{x \in A \mid R(x) \cap R(a) \neq \emptyset\}$ . For this to be true, the condition must hold for  $x = a$ , so  $R(a) \cap R(a) \neq \emptyset$ . This is equivalent to  $R(a) \neq \emptyset$ . Since this must hold for all  $a \in A$ , the relation  $R$  is total.
- Any element  $x \in A$  such that  $x \neq a$  must not belong to the set. This means that for any  $x \neq a$ , we must have  $R(x) \cap R(a) = \emptyset$ . This means the image sets of distinct elements of  $A$  are pairwise disjoint.

Thus,  $R$  is total and injective.

- **Item 2  $\implies$  Item 9:** Let  $a \in A$ . We wish to show  $R^{-1}(R(a)) = \{a\}$ .

- Let  $x \in R^{-1}(R(a))$ . By definition, this means  $R(x) \cap R(a) \neq \emptyset$ .
- Since the image sets are pairwise disjoint, this can only be true if  $x = a$ .
- Therefore,  $R^{-1}(R(a)) \subset \{a\}$ .
- Since  $R$  is total,  $R(a)$  is non-empty.
- Thus  $R(a) \cap R(a) \neq \emptyset$ , which implies  $a \in R^{-1}(R(a))$ .
- Therefore,  $\{a\} \subset R^{-1}(R(a))$ .

Combining both inclusions, we have  $R^{-1}(R(a)) = \{a\}$ .

This finishes the proof. 

02QB

## REMARK 8.5.10.1.12 ► MONOMORPHISMS IN Rel GIVE RISE TO ANTICHAINS

Taking the contrapositive of **Item 2** of **Proposition 8.5.10.1.4** and letting  $U = \{a'\}$  shows that the subset

$$\{R(a) \in \mathcal{P}(B) \mid a \in A\}$$

of  $\mathcal{P}(B)$  forms an antichain in  $\mathcal{P}(B)$ . The converse however, fails.

02QC

**PROPOSITION 8.5.10.1.13 ► CHARACTERISATIONS OF 2-CATEGORICAL MONOMORPHISMS  
IN **Rel I****

Every 1-morphism in **Rel** is representably faithful.

**PROOF 8.5.10.1.14 ► PROOF OF PROPOSITION 8.5.10.1.13**

A relation  $R: A \rightarrow B$  will be representably faithful in **Rel** iff, for each  $X \in \text{Obj}(\mathbf{Rel})$ , the functor

$$R_*: \mathbf{Rel}(X, A) \rightarrow \mathbf{Rel}(X, B)$$

given by postcomposition by  $R$  is faithful. This happens iff the morphism

$$R_{*|S,T}: \text{Hom}_{\mathbf{Rel}(X,A)}(S, T) \rightarrow \text{Hom}_{\mathbf{Rel}(X,B)}(R \diamond S, R \diamond T)$$

is injective for each  $S, T \in \text{Obj}(\mathbf{Rel}(X, A))$ .

However, since **Rel** is locally posetal, the Hom-set  $\text{Hom}_{\mathbf{Rel}(X,A)}(S, T)$  is either empty or a singleton. As a result, the map  $R_{*|S,T}$  will necessarily be injective in either of these cases. 

02QD

**PROPOSITION 8.5.10.1.15 ► CHARACTERISATIONS OF 2-CATEGORICAL MONOMORPHISMS  
IN **Rel II****

Let  $R: A \rightarrow B$  be a relation. The following conditions are equivalent:

02QE

1. The morphism  $R: A \rightarrow B$  is a monomorphism in **Rel**.

02QF

2. The 1-morphism  $R: A \rightarrow B$  is representably full in **Rel**.

02QG

3. The 1-morphism  $R: A \rightarrow B$  is representably fully faithful in **Rel**.

02QH

4. The 1-morphism  $R: A \rightarrow B$  is pseudomonic in **Rel**.

02QJ

5. The 1-morphism  $R: A \rightarrow B$  is representably essentially injective in **Rel**.

02QK

6. The 1-morphism  $R: A \rightarrow B$  is representably conservative in **Rel**.

**PROOF 8.5.10.1.16 ► PROOF OF PROPOSITION 8.5.10.1.15**

We will prove this by showing:

- Step 1: Item 1  $\iff$  Item 2.
- Step 2: Item 2  $\iff$  Item 3.

- Step 3: Item 3  $\iff$  Item 4  $\iff$  Item 5  $\iff$  Item 6.

Step 1: Item 1  $\iff$  Item 2

The condition that  $R$  is representably full corresponds precisely to [Item 7](#) of [Proposition 8.5.10.1.2](#), so this follows by [Proposition 8.5.10.1.2](#).

Step 2: Item 2  $\iff$  Item 3

This follows from Step 1 and [Proposition 8.5.10.1.13](#).

Step 3: Item 3  $\iff$  Item 4  $\iff$  Item 5  $\iff$  Item 6

Since **Rel** is locally posetal, the conditions in [Items 4](#) to [6](#) all collapse to the one of [Item 3](#). 

### 00LW 8.5.11 Epimorphisms and 2-Categorical Epimorphisms

02QL

#### EXPLANATION 8.5.11.1.1 ▶ EPIMORPHISMS IN Rel


In this section, we characterise:

- The 1-categorical epimorphisms in **Rel**, following ??.
- The 2-categorical epimorphisms in **Rel**, following [Section 14.2](#).

More specifically:

- [Proposition 8.5.11.1.2](#) gives *conceptual* characterisations of the epimorphisms in **Rel**.
- [Proposition 8.5.11.1.4](#) gives *point-set* characterisations of the epimorphisms in **Rel**.
- [Warning 8.5.11.1.10](#) lists a few conditions that look natural but fail to characterise epimorphisms in **Rel**.
- [Propositions 8.5.11.1.13](#) and [8.5.11.1.15](#) characterise the 2-categorical epimorphisms in **Rel**.<sup>1</sup>

Essentially, an epimorphism  $R: A \rightarrow B$  in **Rel** looks like this:



In particular:

**02QM**

1.  $R$  should contain a surjection  $f: A \twoheadrightarrow B$ .

**02QN**

2.  $R$  doesn't need to be injective, so  $R$  can map different elements of  $A$  to the same element of  $B$ .

**02QP**

3.  $R$  can be non-functional, mapping elements of  $A$  to multiple elements of  $B$ .

**02QQ**

4.  $R$  can be non-total, so  $R$  doesn't need to be defined on all of  $A$ .

**02QR**

5. For each  $b \in B$ , there must exist some  $a \in A$  with  $R(a) = \{b\}$ .

Moreover, if  $R$  is functional, then being an epimorphism is equivalent to being surjective.

<sup>1</sup>*Summary:* As it turns out, every 1-morphism in **Rel** is representably faithful and most other notions of 2-categorical epimorphism agree with the usual (1-categorical) notion of epimorphism.

**00LY**

#### PROPOSITION 8.5.11.1.2 ► CHARACTERISATIONS OF EPIMORPHISMS IN **Rel** I

Let  $R: A \rightarrow B$  be a relation. The following conditions are equivalent:<sup>1</sup>

**02QS**

1. The relation  $R$  is an epimorphism in **Rel**.

**02QT**

2. The inverse image function

$$R^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to  $R$  is injective.

02QU

3. The coinverse image function

$$R_{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to  $R$  is injective.

02QV

4. The inverse image functor

$$R^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

associated to  $R$  is full.

02QW

5. The coinverse image functor

$$R_{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

associated to  $R$  is full.

02QX

6. For each pair of relations  $S, T: B \nrightarrow X$ , the following condition is satisfied:

(★) If  $S \diamond R \subset T \diamond R$ , then  $S \subset T$ .

02QY

7. There exists an injective function  $f: B \hookrightarrow A$  satisfying the following conditions:<sup>2</sup>

(a) We have  $\text{Gr}(f) \subset R^\dagger$ .<sup>3</sup>

(b) The diagram

$$\begin{array}{ccc} B & \xrightarrow{f} & A \\ & \searrow \chi_B & \downarrow R \\ & & \mathcal{P}(B) \end{array}$$

commutes.<sup>4</sup>

02R1

8. We have

$$\begin{array}{ccc} \mathcal{P}(B) & \xrightarrow{R^{-1}} & \mathcal{P}(A) \\ R_* \circ R^{-1} = \text{id}_{\mathcal{P}(B)} & \cong & \downarrow R_* \\ & & \mathcal{P}(B). \end{array}$$

In other words, we have

$$U = \underbrace{\{b \in B \mid R^{-1}(b) \subset R^{-1}(U)\}}_{=R_*(R^{-1}(U))}$$

for each  $U \in \mathcal{P}(B)$ .

02SC

9. We have

$$\begin{array}{ccc} \mathcal{P}(B) & \xrightarrow{R_{-1}} & \mathcal{P}(A) \\ R_! \circ R_{-1} = \text{id}_{\mathcal{P}(B)} & \searrow & \downarrow R_! \\ & & \mathcal{P}(B). \end{array}$$

In other words, we have

$$U = \underbrace{\left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in R^{-1}(b) \\ \text{such that we have } R(a) \subset U \end{array} \right\}}_{=R_!(R_{-1}(U))}$$

for each  $U \in \mathcal{P}(B)$ .

<sup>1</sup>Items 2 to 5 unwind respectively to the following statements:

- For each  $U, V \in \mathcal{P}(B)$ , if  $R^{-1}(U) = R^{-1}(V)$ , then  $U = V$ .
- For each  $U, V \in \mathcal{P}(B)$ , if  $R_{-1}(U) = R_{-1}(V)$ , then  $U = V$ .
- For each  $U, V \in \mathcal{P}(B)$ , if  $R^{-1}(U) \subset R^{-1}(V)$ , then  $U \subset V$ .

<sup>2</sup>We are assuming the axiom of choice for this item (Item 7).

<sup>3</sup>In other words, for each  $b \in B$ , we have  $f(b) \in R^{-1}(b)$ .

<sup>4</sup>In other words, for each  $b \in B$ , we have  $R(f(b)) = \{b\}$ .

#### PROOF 8.5.11.1.3 ► PROOF OF PROPOSITION 8.5.11.1.2

We will prove this by showing:

- Step 1: Item 2  $\iff$  Item 3 and Item 3  $\iff$  Item 5.
- Step 2: Item 1  $\iff$  Item 2.
- Step 3: Item 2  $\iff$  Item 4.
- Step 4: Item 4  $\iff$  Item 6.
- Step 5: Item 1  $\iff$  Item 7.
- Step 6: Item 1  $\iff$  Item 8.
- Step 7: Item 1  $\iff$  Item 9.

Step 1: Item 2  $\iff$  Item 3 and Item 3  $\iff$  Item 5

This follows from [Item 7 of Proposition 8.7.3.1.4](#).

**Step 2: First Proof of Item 1  $\iff$  Item 2**

We claim that [Items 1](#) and [2](#) are equivalent:

- [Item 1  \$\implies\$  Item 2](#): Let  $U, V \in \mathcal{P}(B)$  and consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ & \dashrightarrow & \dashrightarrow \\ & & V \end{array} \xrightarrow{U} \text{pt.}$$

By [Remark 8.7.1.1.3](#), we have

$$\begin{aligned} R^{-1}(U) &= U \diamond R, \\ R^{-1}(V) &= V \diamond R. \end{aligned}$$

Now, if  $U \diamond R = V \diamond R$ , i.e.  $R^{-1}(U) = R^{-1}(V)$ , then  $U = V$  since  $R$  is assumed to be an epimorphism, showing  $R^{-1}$  to be injective.

- [Item 2  \$\implies\$  Item 1](#): Conversely, suppose that  $R^{-1}$  is injective, consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ & \dashrightarrow & \dashrightarrow \\ & & T \end{array} \xrightarrow{S} X,$$

and suppose that  $S \diamond R = T \diamond R$ . Note that, since  $R^{-1}$  is injective, given a diagram of the form

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ & \dashrightarrow & \dashrightarrow \\ & & V \end{array} \xrightarrow{U} \text{pt},$$

if  $R^{-1}(U) = U \diamond R = V \diamond R = R^{-1}(V)$ , then  $U = V$ . In particular, for each  $x \in X$ , we may consider the diagram

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow{S} & X \\ & \dashrightarrow & \dashrightarrow & \dashrightarrow & \dashrightarrow \\ & & T & & [x] \\ & & & & \dashrightarrow \end{array} \xrightarrow{\quad} \text{pt},$$

for which we have  $[x] \diamond S \diamond R = [x] \diamond T \diamond R$ , implying that we have

$$S^{-1}(x) = [x] \diamond S = [x] \diamond T = T^{-1}(x)$$

for each  $x \in X$ . Thus  $S = T$  and  $R$  is an epimorphism.

**Step 2.5: Second Proof of Item 1  $\iff$  Item 2**

A more abstract proof can also be given, following [\[MSE 350788\]](#):

- *Item 1  $\implies$  Item 2:* Assume that  $R$  is an epimorphism.
  - We first notice that the functor  $\text{Rel}(-, \text{pt}) : \text{Rel}^{\text{op}} \rightarrow \text{Sets}$  maps  $R$  to  $R^{-1}$  by Remark 8.7.3.1.2.
  - Since  $\text{Rel}(-, \text{pt})$  preserves limits by ?? of ??, it follows by ?? of ?? that  $\text{Rel}(-, \text{pt})$  also preserves epimorphisms.
  - That is:  $\text{Rel}(-, \text{pt})$  sends epimorphisms in  $\text{Rel}^{\text{op}}$  to epimorphisms in  $\text{Sets}$ .
  - The epimorphisms  $\text{Rel}^{\text{op}}$  are precisely the epimorphisms in  $\text{Rel}$  by ?? of ??.
  - Since  $R$  is an epimorphism and  $\text{Rel}(-, \text{pt})$  maps  $R$  to  $R^{-1}$ , it follows that  $R^{-1}$  is an epimorphism.
  - Since the epimorphisms in  $\text{Sets}$  are precisely the injections (?? of ??), it follows that  $R^{-1}$  is injective.
- *Item 2  $\implies$  Item 1:* Assume that  $R^{-1}$  is injective.
  - We first notice that the functor  $\text{Rel}(-, \text{pt}) : \text{Rel}^{\text{op}} \rightarrow \text{Sets}$  maps  $R$  to  $R^{-1}$  by Remark 8.7.3.1.2.
  - Since the epimorphisms in  $\text{Sets}$  are precisely the injections (?? of ??), it follows that  $R^{-1}$  is an epimorphism.
  - Since  $\text{Rel}(-, \text{pt})$  is faithful, it follows by ?? of ?? that  $\text{Rel}(-, \text{pt})$  reflects epimorphisms.
  - That is:  $\text{Rel}(-, \text{pt})$  reflects epimorphisms in  $\text{Sets}$  to epimorphisms in  $\text{Rel}^{\text{op}}$ .
  - The epimorphisms  $\text{Rel}^{\text{op}}$  are precisely the epimorphisms in  $\text{Rel}$  by ?? of ??.
  - Since  $R^{-1}$  is an epimorphism and  $\text{Rel}(-, \text{pt})$  maps  $R$  to  $R^{-1}$ , it follows that  $R$  is an epimorphism.

**Step 3: Item 2  $\iff$  Item 4**

We claim that Items 2 and 4 are equivalent:

- *Item 2  $\implies$  Item 4:* We proceed in a few steps:

- Let  $U, V \in \mathcal{P}(B)$  such that  $R^{-1}(U) \subset R^{-1}(V)$ , assume  $R^{-1}$  to be injective, and consider the set  $U \cup V$ .
- Since  $R^{-1}(U) \subset R^{-1}(V)$ , we have

$$R^{-1}(U \cup V) = R^{-1}(U) \cup R^{-1}(V)$$

$$= R^{-1}(V),$$

where we have used [Item 5 of Proposition 8.7.3.1.4](#) for the first equality.

- Since  $R^{-1}$  is injective, this implies  $U \cup V = V$ .
- Thus  $U \subset V$ , as we wished to show.

- [Item 2  \$\implies\$  Item 4](#): We proceed in a few steps:

- Suppose [Item 4](#) holds, and let  $U, V \in \mathcal{P}(B)$  such that  $R^{-1}(U) = R^{-1}(V)$ .
- Since  $R^{-1}(U) = R^{-1}(V)$ , we have  $R^{-1}(U) \subset R^{-1}(V)$  and  $R^{-1}(V) \subset R^{-1}(U)$ .
- By assumption, this implies  $U \subset V$  and  $V \subset U$ .
- Thus  $U = V$ , showing  $R^{-1}$  to be injective.

#### Step 4: Item 4 $\iff$ Item 6

We claim that [Items 4](#) and [6](#) are equivalent:

- [Item 4  \$\implies\$  Item 6](#): Let  $U, V \in \mathcal{P}(B)$  and consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{\quad R \quad} & B \\ & \dashrightarrow & \end{array} \begin{array}{c} U \\ \parallel \\ V \end{array} \xrightarrow{\quad \text{pt.} \quad}$$

By [Remark 8.7.3.1.2](#), we have

$$\begin{aligned} R^{-1}(U) &= U \diamond R, \\ R^{-1}(V) &= V \diamond R. \end{aligned}$$

Now, if  $U \diamond R \subset V \diamond R$ , then  $R^{-1}(U) \subset R^{-1}(V)$ . By assumption, we then have  $U \subset V$ .

- [Item 6  \$\implies\$  Item 4](#): Consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{\quad R \quad} & B \\ & \dashrightarrow & \end{array} \begin{array}{c} S \\ \parallel \\ T \end{array} \xrightarrow{\quad X \quad}$$

and suppose that  $S \diamond R \subset T \diamond R$ . Note that, by assumption, given a diagram of the form

$$\begin{array}{ccc} A & \xrightarrow{\quad R \quad} & B \\ & \dashrightarrow & \end{array} \begin{array}{c} U \\ \parallel \\ V \end{array} \xrightarrow{\quad \text{pt.} \quad}$$

if  $R^{-1}(U) = R \diamond U \subset R \diamond V = R^{-1}(V)$ , then  $U \subset V$ . In particular, for each  $x \in X$ , we may consider the diagram

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow[S]{T} & X \xrightarrow{[x]} \text{pt}, \end{array}$$

for which we have  $[x] \diamond S \diamond R \subset [x] \diamond T \diamond R$ , implying that we have

$$S^{-1}(x) = [x] \diamond S \subset [x] \diamond T = T^{-1}(x)$$

for each  $x \in X$ , implying  $S \subset T$ .

This finishes the proof.

**Step 5:** Item 1  $\iff$  Item 7

We defer this proof to [Corollary 8.5.11.1.6](#).

**Step 6:** Item 1  $\iff$  Item 8

We defer this proof to [Corollary 8.5.11.1.6](#).

**Step 6:** Item 1  $\iff$  Item 9

We defer this proof to [Old Tag 15.2.1.1.24](#).



00LZ

#### PROPOSITION 8.5.11.1.4 ► CHARACTERISATIONS OF EPIMORPHISMS IN Rel II

Let  $R: A \nrightarrow B$  be a relation. The following conditions are equivalent:

02R2

1. The relation  $R$  is an epimorphism in Rel.

02R3

2. For each  $b \in B$  and each  $U \in \mathcal{P}(B)$ , if  $R^{-1}(b) \subset R^{-1}(U)$ , then  $b \in U$ .

02R4

3. For each  $b \in B$ , there exists some  $a \in A$  such that  $R(a) = \{b\}$ .

Moreover, if  $R$  is an epimorphism, then it is surjective, and the converse holds if  $R$  is functional.

#### PROOF 8.5.11.1.5 ► PROOF OF PROPOSITION 8.5.11.1.4

We will prove this by showing:

- Step 1: [Item 1  \$\implies\$  Item 2](#).
- Step 2: [Item 2  \$\implies\$  Item 3](#).
- Step 3: [Item 3  \$\implies\$  Item 1](#).

- Step 4: The second half of the statement of [Proposition 8.5.11.1.2](#).

**Step 1: Item 1  $\implies$  Item 2**

We proceed in a few steps:

- If  $R$  is an epimorphism, then, by [Item 2 of Proposition 8.5.11.1.2](#), the functor

$$R^{-1}: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

is full.

- As a result, given  $b \in B$  and  $U \in \mathcal{P}(B)$  such that  $R^{-1}(b) \subset R^{-1}(U)$ , it follows that  $\{b\} \subset U$ .
- Thus, we have  $b \in U$ .

**Step 2: Item 2  $\implies$  Item 3**

We proceed in a few steps:

- Let  $b \in B$  and consider the subset  $U = B \setminus \{b\}$ .
- Since  $b \notin U$ , we have  $R^{-1}(b) \not\subset R^{-1}(U)$  by the contrapositive of [Item 2](#).
- As a result, there must exist some  $a \in R^{-1}(b)$  with  $a \notin R^{-1}(U)$ .
- In particular, we have  $b \in R(a)$ .
- Moreover, the condition  $a \notin R^{-1}(U) = R^{-1}(B \setminus \{b\})$  means that, if  $b' \in B \setminus \{b\}$ , then  $b' \notin R(a)$ .
- Thus  $R(a) = \{b\}$ .

**Step 3: Item 3  $\implies$  Item 1**

We proceed in a few steps:

- By the equivalence between [Items 1 and 4 of Proposition 8.5.11.1.2](#), to show that  $R$  is an epimorphism it suffices to prove that, for each  $U, V \in \mathcal{P}(B)$ , if  $R^{-1}(U) \subset R^{-1}(V)$ , then  $U \subset V$ .
- So let  $u \in U$  and assume  $R^{-1}(U) \subset R^{-1}(V)$ .
- By assumption, there exists some  $a \in A$  with  $R(a) = \{u\}$ .
- In particular,  $a \in R^{-1}(U)$ .

- Since  $R^{-1}(U) \subset R^{-1}(V)$ , we also have  $a \in R^{-1}(V)$ .
- Thus, there exists some  $v \in V$  with  $a \in R(v)$ .
- However,  $R(a) = \{u\}$ , so we must in fact have  $v = u$ .
- Therefore  $u \in V$ , showing that  $U \subset V$ .

**Step 4: Proof of the Second Half of Proposition 8.5.11.1.4**

We claim that  $R$  being an epimorphism implies surjectivity, and the converse holds if  $R$  is functional:

- *If  $R$  Is an Epimorphism, Then  $R$  Is Surjective:* Consider the diagram

$$A \xrightarrow{\quad R \quad} B \xrightarrow{\quad S \quad} \{0, 1\}, \quad T$$

where  $b \sim_S 0$  for each  $b \in B$  and where we have

$$b \sim_T \begin{cases} 0 & \text{if } b \in \text{Im}(R), \\ 1 & \text{otherwise} \end{cases}$$

for each  $b \in B$ .

– We claim that  $S \diamond R = T \diamond R$ :

- \* If  $R(a) = \emptyset$ , then

$$\begin{aligned} [S \diamond R](a) &= \emptyset \\ [T \diamond R](a) &= \emptyset \end{aligned}$$

by the definition of relational composition, so  $[S \diamond R](a) = [T \diamond R](a)$ .

- \* If  $R(a) \neq \emptyset$ , then we have  $a \sim_{S \diamond R} 0$  and  $a \sim_{T \diamond R} 0$  by the definition of  $S$  and  $T$ , with no element of  $A$  being related to 1 by  $S \diamond R$  or  $T \diamond R$ .

– Now, since  $R$  is an epimorphism, we have  $S = T$ .

– However, by the definition of  $T$ , this implies  $\text{Im}(R) = B$ .

– Thus  $R$  is surjective.

- *If  $R$  Is Functional and Surjective, Then  $R$  Is an Epimorphism:* Let  $U, V \in \mathcal{P}(B)$ , consider the diagram

$$A \xrightarrow{\quad R \quad} B \xrightarrow{\quad U \quad} \text{pt}, \quad V$$

where  $R^{-1}(U) = R^{-1}(V)$ , and let  $b \in U$ .

- By surjectivity, there exists some  $a \in A$  such that  $a \in R^{-1}(b)$ .
- Since  $R_{-1}(U) = R_{-1}(V)$ , if  $R(a) \subset U$ , then  $R(a) \subset V$ .
- Since  $R$  is functional, we have  $R(a) = \{b\}$ , so  $R(a) \subset U$ .
- Thus,  $R(a) \subset V$ , and  $b \in V$ .
- A similar argument shows that if  $b \in V$ , then  $b \in U$ .
- Thus  $U = V$ , showing  $R_{-1}$  to be injective.
- By the equivalence between [Items 1](#) and [3](#) of [Proposition 8.5.11.1.2](#), this shows  $R$  to be an epimorphism.

This finishes the proof. 

00M0

#### COROLLARY 8.5.11.1.6 ► CHARACTERISATIONS OF EPIMORPHISMS IN Rel III

[Items 1, 7](#) and [8](#) of [Proposition 8.5.11.1.2](#) are indeed equivalent.

#### PROOF 8.5.11.1.7 ► PROOF OF COROLLARY 8.5.11.1.6

Item 1  $\iff$  Item 7

We claim that [Item 3](#) of [Proposition 8.5.11.1.4](#) is equivalent to [Item 7](#) of [Proposition 8.5.11.1.2](#):

- [Item 3 of Proposition 8.5.11.1.4](#)  $\implies$  [Item 7](#): By assumption, given  $b \in B$ , there exists some  $a \in A$  such that  $R(a) = \{b\}$ . Invoking the axiom of choice, we may pick one such  $a$  for each  $b \in B$ , giving us our desired function  $f: B \rightarrow A$ . All the requirements listed in [Item 7](#) then follow by construction.
- [Item 7](#)  $\implies$  [Item 3 of Proposition 8.5.11.1.4](#): Given  $b \in B$ , we may pick  $a = f(b)$ , in which case  $R(f(b))$  will be equal to  $\{b\}$  by assumption.

By [Proposition 8.5.11.1.4](#), [Item 3](#) of [Proposition 8.5.11.1.4](#) is equivalent to [Item 1](#) of [Proposition 8.5.11.1.4](#). Since [Item 1](#) of [Proposition 8.5.11.1.4](#) is exactly the same condition as [Item 1](#) of [Proposition 8.5.11.1.2](#), the result follows.

Item 1  $\iff$  Item 8

Indeed, we have

$$\begin{aligned} [R_* \circ R^{-1}](U) &\stackrel{\text{def}}{=} R_*(R^{-1}(U)) \\ &\stackrel{\text{def}}{=} \{b \in B \mid R^{-1}(b) \subset R^{-1}(U)\} \end{aligned}$$

for each  $U \in \mathcal{P}(B)$ . As a result, the condition  $R_* \circ R^{-1} = \text{id}_{\mathcal{P}(B)}$  becomes

$$\{b \in B \mid R^{-1}(b) \subset R^{-1}(U)\} = U,$$

which holds precisely when [Item 2 of Proposition 8.5.11.1.4](#) does. By [Proposition 8.5.11.1.4](#), that in turn holds precisely if [Item 1 of Proposition 8.5.11.1.4](#) holds. Since [Item 1 of Proposition 8.5.11.1.4](#) is exactly the same condition as [Item 1 of Proposition 8.5.11.1.2](#), the result follows.



02SD

**COROLLARY 8.5.11.1.8 ► CHARACTERISATIONS OF EPIMORPHISMS IN Rel IV**

[Items 1 and 9 of Proposition 8.5.11.1.2](#) are indeed equivalent.

**PROOF 8.5.11.1.9 ► PROOF OF OLD TAG 15.2.1.1.24****Item 1  $\implies$  Item 9**

To show that  $R$  is an epimorphism, we will prove that for each  $b \in B$ , there exists some  $a \in A$  such that  $R(a) = \{b\}$ . This will then follow from [Item 3 of Proposition 8.5.11.1.4](#).

- Let  $b \in B$  and consider  $U = \{b\}$ .
- By assumption, we have  $U = R_!(R_{-1}(U))$ .
- In particular, this means that  $b \in R_!(R_{-1}(\{b\}))$ .
- Unwinding the definition, this means there exists some  $a \in R^{-1}(b)$  such that  $R(a) \subset \{b\}$ .
  - The condition  $a \in R^{-1}(b)$  implies that  $b \in R(a)$ .
  - The condition  $R(a) \subset \{b\}$  implies that every element of  $R(a)$  must be  $b$ .
- For  $R(a)$  to be a non-empty subset of  $\{b\}$ , it must be the case that  $R(a) = \{b\}$ .

This completes the proof.

**Item 9  $\implies$  Item 1**

We wish to show that for any  $U \in \mathcal{P}(B)$ , we have  $U = R_!(R_{-1}(U))$ . This requires proving two set inclusions.

- $R_!(R_{-1}(U)) \subset U$ : Let  $b \in R_!(R_{-1}(U))$ .

- By definition, there exists an  $a \in A$  such that  $a \in R^{-1}(b)$  and  $R(a) \subset U$ .
- The condition  $a \in R^{-1}(b)$  means that  $b \in R(a)$ .
- Since  $b \in R(a)$  and  $R(a) \subset U$ , it follows directly that  $b \in U$ .
- Therefore,  $R_!(R_{-1}(U)) \subset U$ .
- $U \subset R_!(R_{-1}(U))$ : Let  $b \in U$ .
  - By Item 3 of Proposition 8.5.11.1.4, there exists an element  $a \in A$  such that  $R(a) = \{b\}$ .
  - We must verify that this choice of  $a$  places  $b$  into the set  $R_!(R_{-1}(U))$ . This requires checking two conditions:
    - \*  $a \in R^{-1}(b)$ : Since  $R(a) = \{b\}$ , we have  $b \in R(a)$ , which is equivalent to  $a \in R^{-1}(b)$ .
    - \*  $R(a) \subset U$ : Since  $R(a) = \{b\}$  and we assumed  $b \in U$ , we have  $\{b\} \subset U$ , so the condition holds.
  - As both conditions are met, it follows that  $b \in R_!(R_{-1}(U))$ .
  - Therefore,  $U \subset R_!(R_{-1}(U))$ .

As both inclusions hold, we conclude that  $U = R_!(R_{-1}(U))$ , which is precisely the statement of Item 9. 

**WARNING 8.5.11.1.10 ► NATURAL CONDITIONS THAT FAIL TO CHARACTERISE EPIMORPHISMS IN Rel**

The following conditions are equivalent and imply  $R$  is an epimorphism, but the converse may fail. Thus they are *not* equivalent to  $R$  being an epimorphism:

02SF

1. The relation  $R$  is a surjective partial function.

02SG

2. The diagram

$$\begin{array}{ccc} B & \xrightarrow{R^{-1}} & \mathcal{P}(A) \\ & \searrow \chi_B & \downarrow R_! \\ & & \mathcal{P}(B) \end{array}$$

commutes. In other words, we have

$$R_!(R^{-1}(b)) = \{b\}$$

for each  $b \in B$ .

02SH

3. We have

$$\begin{array}{ccc} \mathcal{P}(B) & \xrightarrow{R^{-1}} & \mathcal{P}(A) \\ R_! \circ R^{-1} = \text{id}_{\mathcal{P}(B)} & \swarrow & \downarrow R_! \\ & & \mathcal{P}(B). \end{array}$$

In other words, we have

$$U = \underbrace{\{b \in B \mid R^{-1}(b) \cap R^{-1}(U) \neq \emptyset\}}_{=R_!(R^{-1}(U))}$$

for each  $U \in \mathcal{P}(B)$ .

02SJ

4. We have

$$\begin{array}{ccc} \mathcal{P}(B) & \xrightarrow{R_{-1}} & \mathcal{P}(A) \\ R_* \circ R_{-1} = \text{id}_{\mathcal{P}(B)} & \swarrow & \downarrow R_* \\ & & \mathcal{P}(B). \end{array}$$

In other words, we have

$$U = \underbrace{\{b \in B \mid R(R^{-1}(b)) \subset U\}}_{=R_*(R_{-1}(U))}$$

for each  $U \in \mathcal{P}(B)$ .

## PROOF 8.5.11.1.11 ► PROOF OF WARNING 8.5.11.1.10

First, note that the relation depicted in [Explanation 8.5.11.1.1](#) is not a surjective partial function, but it is an epimorphism in Rel by [Proposition 8.5.11.1.4](#), the next proposition. Moreover, partial surjective functions are epimorphisms by [Proposition 8.5.11.1.4](#). For the rest of the proposition, we proceed by showing:

- Step 1: [Item 1](#)  $\iff$  [Item 2](#).
- Step 2: [Item 2](#)  $\implies$  [Item 3](#).
- Step 3: [Item 3](#)  $\implies$  [Item 2](#).
- Step 4: [Item 2](#)  $\implies$  [Item 4](#).
- Step 5: [Item 4](#)  $\implies$  [Item 1](#).

Step 1: Item 1  $\iff$  Item 2

Note that we have

$$R_!(R^{-1}(b)) \stackrel{\text{def}}{=} \{b' \in B \mid R^{-1}(b') \cap R^{-1}(b) \neq \emptyset\}.$$

We now claim [Items 1](#) and [2](#) are equivalent:

- [Item 1](#)  $\implies$  [Item 2](#): We proceed in a few steps:
  - Since  $R$  is functional,  $R^{-1}(b)$  has at most one element.
  - Since  $R$  is surjective,  $R^{-1}(b)$  has at least one element.
  - Thus,  $R^{-1}(b)$  is a singleton.
  - The set  $R(R^{-1}(b))$  will then be precisely  $\{b\}$ .
- [Item 2](#)  $\implies$  [Item 1](#): We claim  $R$  is functional and surjective.
  - *Functionality*. The inclusion

$$R_!(R^{-1}(b)) \subset \{b\}$$

implies that if  $a \in R(b')$  and  $a \in R(b)$ , then  $b = b'$ . Thus  $R$  must be functional.

- *Surjectivity*. The inclusion

$$\{b\} \subset R_!(R^{-1}(b))$$

implies  $R^{-1}(b) \neq \emptyset$ , so  $R$  must be surjective.

Since  $R$  is functional and surjective, it is a surjective partial function.

**Step 2: Item 2  $\implies$  Item 3**

We have

$$\begin{aligned} [R_! \circ R^{-1}](U) &\stackrel{\text{def}}{=} R_!(R^{-1}(U)) \\ &= R_!\left(R^{-1}\left(\bigcup_{u \in U} \{u\}\right)\right) \\ &= R_!\left(\bigcup_{u \in U} R^{-1}(\{u\})\right) \\ &= \bigcup_{u \in U} R_!(R^{-1}(\{u\})) \\ &= \bigcup_{u \in U} \{u\} \\ &= U \end{aligned}$$

for each  $U \in \mathcal{P}(B)$ , where we have used:

- ?? of [Proposition 8.7.3.1.4](#) for the third equality.
- ?? of [Proposition 8.7.1.1.4](#) for the fourth equality.
- [Item 2](#) of this proposition for the fifth equality.

**Step 3: Item 3  $\implies$  Item 2**

Taking  $U = \{b\}$  gives  $R_!(R^{-1}(b)) = \{b\}$ .

**Step 4: Item 2  $\implies$  Item 4**

We have

$$\begin{aligned} R_*(R_{-1}(U)) &= \{b \in B \mid R(R^{-1}(b)) \subset U\} \\ &= \{b \in B \mid \{b\} \subset U\} \\ &= U. \end{aligned}$$

**Step 5: Item 4  $\implies$  Item 1**

Suppose that for each  $U \in \mathcal{P}(B)$ , we have  $R_*(R_{-1}(U)) = U$ . We must show that  $R$  is functional and surjective.

- *Functionality:* We show that if  $b, b' \in R(a)$ , then  $b = b'$ .

- Consider the singleton set  $U = \{b\}$ . By the assumed identity, we have

$$\{b\} = \{b \in B \mid R(R^{-1}(b)) \subset \{b\}\}.$$

- Since  $b$  is an element of the set on the left-hand side, it must satisfy the membership condition on the right-hand side. Thus, we have  $R(R^{-1}(b)) \subset \{b\}$ .
- By assumption,  $b \in R(a)$ , which implies  $a \in R^{-1}(b)$ .
- By assumption, we also have  $b' \in R(a)$ .
- Since  $a \in R^{-1}(b)$ , it follows that the image of  $a$  is contained in the image of the set  $R^{-1}(b)$ , i.e.,  $R(a) \subset R(R^{-1}(b))$ .
- Combining these steps, we have  $b' \in R(a) \subset R(R^{-1}(b))$ .
- As we established that  $R(R^{-1}(b)) \subset \{b\}$ , we must have  $b' \in \{b\}$ .
- Therefore,  $b' = b$ , which shows  $R$  to be functional.
- *Surjectivity:* We show that for each  $b \in B$ , the preimage set  $R^{-1}(b)$  is non-empty.
  - Consider the empty set  $U = \emptyset$ . By the assumed identity, we have

$$\emptyset = \{b \in B \mid R(R^{-1}(b)) \subset \emptyset\}.$$

- The identity thus states that there is no element  $b \in B$  for which  $R(R^{-1}(b))$  is the empty set.
- In other words, for each  $b \in B$ , we must have  $R(R^{-1}(b)) \neq \emptyset$ .
- The image of a set  $R(S)$  is empty iff the set  $S$  is empty.
- Therefore, the condition  $R(R^{-1}(b)) \neq \emptyset$  is equivalent to the condition  $R^{-1}(b) \neq \emptyset$ .
- Thus,  $R$  is surjective.

Since  $R$  is both functional and surjective, it is a surjective partial function. This finishes the proof. □

02R7

**REMARK 8.5.11.1.12 ► EPIMORPHISMS IN Rel GIVE RISE TO ANTICHAINS**

Taking the contrapositive of Item 2 of Proposition 8.5.11.1.4 and letting  $U = \{b'\}$  shows that the subset

$$\{R^{-1}(b) \in \mathcal{P}(A) \mid b \in B\}$$

of  $\mathcal{P}(A)$  forms an antichain in  $\mathcal{P}(A)$ . The converse however, fails.

02R8

Every 1-morphism in **Rel** is corepresentably faithful.

**PROOF 8.5.11.1.14 ▶ PROOF OF PROPOSITION 8.5.11.1.13**

A relation  $R: A \rightarrow B$  will be corepresentably faithful in **Rel** iff, for each  $X \in \text{Obj}(\mathbf{Rel})$ , the functor

$$R^*: \mathbf{Rel}(B, X) \rightarrow \mathbf{Rel}(A, X)$$

given by precomposition by  $R$  is faithful. This happens iff the morphism

$$R_{S,T}^*: \text{Hom}_{\mathbf{Rel}(B, X)}(S, T) \rightarrow \text{Hom}_{\mathbf{Rel}(A, X)}(S \diamond R, T \diamond R)$$

is injective for each  $S, T \in \text{Obj}(\mathbf{Rel}(B, X))$ .

However, since **Rel** is locally posetal, the Hom-set  $\text{Hom}_{\mathbf{Rel}(B, X)}(S, T)$  is either empty or a singleton. As a result, the map  $R_{S,T}^*$  will necessarily be injective in either of these cases. 

**PROPOSITION 8.5.11.1.15 ▶ CHARACTERISATIONS OF 2-CATEGORICAL EPIMORPHISMS IN **Rel II****

02R9

Let  $R: A \rightarrow B$  be a relation. The following conditions are equivalent:

02RA

1. The morphism  $R: A \rightarrow B$  is an epimorphism in **Rel**.

02RB

2. The 1-morphism  $R: A \rightarrow B$  is corepresentably full in **Rel**.

02RC

3. The 1-morphism  $R: A \rightarrow B$  is corepresentably fully faithful in **Rel**.

02RD

4. The 1-morphism  $R: A \rightarrow B$  is pseudoepic in **Rel**.

02RE

5. The 1-morphism  $R: A \rightarrow B$  is corepresentably essentially injective in **Rel**.

02RF

6. The 1-morphism  $R: A \rightarrow B$  is corepresentably conservative in **Rel**.

**PROOF 8.5.11.1.16 ▶ PROOF OF PROPOSITION 8.5.11.1.15**

We will prove this by showing:

- Step 1: Item 1  $\iff$  Item 2.
- Step 2: Item 2  $\iff$  Item 3.
- Step 3: Item 3  $\iff$  Item 4  $\iff$  Item 5  $\iff$  Item 6.

Step 1: Item 1  $\iff$  Item 2

The condition that  $R$  is representably full corresponds precisely to **Item 6** of [Proposition 8.5.11.1.2](#), so this follows by [Proposition 8.5.11.1.2](#).

Step 2: Item 2  $\iff$  Item 3

This follows from Step 1 and [Proposition 8.5.11.1.13](#).

Step 3: Item 3  $\iff$  Item 4  $\iff$  Item 5  $\iff$  Item 6

Since **Rel** is locally posetal, the conditions in **Items 4** to **6** all collapse to the one of **Item 3**. 

### 00ME 8.5.12 Co/Limits

00MF

#### PROPOSITION 8.5.12.1.1 ► CO/LIMITS IN Rel

This will be properly written later on.

#### PROOF 8.5.12.1.2 ► PROOF OF PROPOSITION 8.5.12.1.1

Omitted. 

### 00NH 8.5.13 Internal Left Kan Extensions

00NJ

#### PROPOSITION 8.5.13.1.1 ► INTERNAL LEFT KAN EXTENSIONS IN Rel

Let  $R: A \rightarrow B$  be a relation.

00NK

1. *Non-Existence of All Internal Left Kan Extensions in Rel.* Not all relations in **Rel** admit left Kan extensions.

00NL

2. *Characterisation of Relations Admitting Internal Left Kan Extensions Along Them.* The following conditions are equivalent:

02A1

(a) The left Kan extension

$$\text{Lan}_R: \text{Rel}(A, X) \rightarrow \text{Rel}(B, X)$$

along  $R$  exists.

02A2

(b) The relation  $R$  admits a left adjoint in **Rel**.

02A3

(c) The relation  $R$  is of the form  $\text{Gr}(f)$  (as in [Definition 8.2.2.1.1](#)) for some function  $f$ .

**PROOF 8.5.13.1.2 ► PROOF OF PROPOSITION 8.5.13.1.1**

**Item 1: Non-Existence of All Internal Left Kan Extensions in **Rel****

By **Item 2**, it suffices to take a relation that doesn't have a left adjoint.

**Item 2: Characterisation of Relations Admitting Left Kan Extensions**

This proof is mostly due to Tim Campion, via [MO 460693].

- We may view precomposition

$$-\diamond R: \mathbf{Rel}(B, C) \rightarrow \mathbf{Rel}(A, C)$$

with  $R: A \nrightarrow B$  as a cocontinuous functor from  $\mathcal{P}(B \times C)$  to  $\mathcal{P}(A \times C)$  (via **Item 5** of **Definition 8.1.1.1.1**).

- By the adjoint functor theorem (??), this map has a left adjoint iff it preserves limits.
- If  $C = \emptyset$ , this holds trivially.
- Otherwise,  $C$  admits pt as a retract, and we reduce to the case  $C = \text{pt}$  via ??.
- For the case  $C = \text{pt}$ , a relation  $T: B \nrightarrow \text{pt}$  is the same as a subset of  $B$ , and  $-\diamond R$  becomes the inverse image functor  $R^{-1}$  of **Section 8.7.3**.
- Now, again by the adjoint functor theorem,  $R^{-1}$  preserves limits exactly when it has a left adjoint.
- Finally  $R^{-1}$  has a left adjoint precisely when  $R = \text{Gr}(f)$  for  $f$  a function (**Item 8** of **Proposition 8.7.3.1.4**).

This finishes the proof. □

02F7

**EXAMPLE 8.5.13.1.3 ► INTERNAL LEFT KAN EXTENSIONS ALONG FUNCTIONS**

Given a function  $f: A \rightarrow B$ , the left Kan extension

$$\text{Lan}_f: \mathbf{Rel}(A, X) \rightarrow \mathbf{Rel}(B, X)$$

along  $f$  exists by **Item 2** of **Proposition 8.5.13.1.1**. Explicitly, given a relation  $R: A \nrightarrow X$ , the left Kan extension

$$\text{Lan}_f(R): B \nrightarrow X,$$

may be described as follows:

02F8 1. We declare  $b \sim_{\text{Lan}_f(R)} x$  iff there exists some  $a \in R$  such that  $b = f(a)$  and  $a \sim_R x$ .

02F9 2. We have<sup>1</sup>

$$[\text{Lan}_f(R)](b) = \bigcup_{a \in f^{-1}(b)} R(a)$$

for each  $b \in B$ .

<sup>1</sup>Cf. Item 3 of Proposition 8.5.15.1.2.

#### REMARK 8.5.13.1.4 ► ILLUSTRATING THE FAILURE OF INTERNAL LEFT KAN EXTENSIONS IN Rel TO EXIST

02FA Following Example 8.5.13.1.3, given a relation  $R: A \rightarrow B$  and a relation  $F: A \rightarrow X$ , we could perhaps try to define an “honorary” left Kan extension

$$\text{Lan}'_R(F): B \rightarrow X$$

by

$$[\text{Lan}'_R(F)](b) \stackrel{\text{def}}{=} \bigcup_{a \in R^{-1}(b)} F(a)$$

for each  $b \in B$ .

The failure of  $\text{Lan}'_R(F)$  to be a Kan extension can then be seen as follows. Let  $G: B \rightarrow X$  be a relation. If  $\text{Lan}'_R(F)$  were a left Kan extension, then the following conditions **would be** equivalent:

02FB 1. For each  $b \in B$ , we have  $\bigcup_{a \in R^{-1}(b)} F(a) \subset G(b)$ .

02FC 2. For each  $a \in A$ , we have  $F(a) \subset \bigcup_{b \in R(a)} G(b)$ .

The issue is two-fold:

- *Totality*. If  $R$  isn’t total, then the implication Item 1  $\Rightarrow$  Item 2 fails.
- *Functionality*. If  $R$  isn’t functional, then the implication Item 2  $\Rightarrow$  Item 1 fails.

#### QUESTION 8.5.13.1.5 ► EXISTENCE OF SPECIFIC INTERNAL LEFT KAN EXTENSIONS OF RELATIONS

00NM Given relations  $S: A \rightarrow X$  and  $R: A \rightarrow B$ , is there a characterisation of when the left Kan extension<sup>1</sup>

$$\text{Lan}_S(R): B \rightarrow X$$

exists in terms of properties of  $R$  and  $S$ ?  
 This question also appears as [MO 461592].

<sup>1</sup>Specifically for  $R$  and  $S$ , not  $\text{Lan}_S$  the functor.

### 00NP 8.5.14 Internal Left Kan Lifts

00NQ

#### PROPOSITION 8.5.14.1.1 ► INTERNAL LEFT KAN LIFTS IN **Rel**

Let  $R: A \rightarrow B$  be a relation.

00NR

1. *Non-Existence of All Internal Left Kan Lifts in **Rel**.* Not all relations in **Rel** admit left Kan lifts.

00NS

2. *Characterisation of Relations Admitting Internal Left Kan Lifts Along Them.* The following conditions are equivalent:

02A4

- (a) The left Kan lift

$$\text{Lift}_R: \mathbf{Rel}(X, B) \rightarrow \mathbf{Rel}(X, A)$$

along  $R$  exists.

02A5

- (b) The relation  $R$  admits a right adjoint in **Rel**.

02A6

- (c) The relation  $R$  is of the form  $f^{-1}$  (as in Definition 8.2.3.1.1) for some function  $f$ .

#### PROOF 8.5.14.1.2 ► PROOF OF PROPOSITION 8.5.14.1.1

Item 1: Non-Existence of All Internal Left Kan Lifts in **Rel**

By Item 2, it suffices to take a relation that doesn't have a right adjoint.

Item 2: Characterisation of Relations Admitting Left Kan Lifts Along  $R^\dagger$

This proof is dual to that of Item 2 of Proposition 8.5.13.1.1, and is therefore omitted. 

02FG

#### EXAMPLE 8.5.14.1.3 ► INTERNAL LEFT KAN LIFTS ALONG FUNCTIONS

Given a function  $f: A \rightarrow B$ , the left Kan lift

$$\text{Lift}_{f^\dagger}: \mathbf{Rel}(X, A) \rightarrow \mathbf{Rel}(X, B)$$

along  $f^\dagger$  exists by [Item 2 of Proposition 8.5.14.1.1](#). Explicitly, given a relation  $R: X \nrightarrow A$ , the left Kan lift

$$\text{Lift}_{f^\dagger}(R): X \nrightarrow B,$$

is given by

$$\begin{aligned} [\text{Lift}_f(R)](x) &= [\text{Gr}(f) \diamond R](a) \\ &= \bigcup_{a \in R(x)} f(a) \end{aligned}$$

for each  $x \in X$ .

00NT

**QUESTION 8.5.14.1.4 ► EXISTENCE OF SPECIFIC INTERNAL LEFT KAN LIFTS OF RELATIONS**

Given relations  $S: A \nrightarrow X$  and  $R: A \nrightarrow B$ , is there a characterisation of when the left Kan lift<sup>1</sup>

$$\text{Lift}_S(R): X \nrightarrow A$$

exists in terms of properties of  $R$  and  $S$ ?

This question also appears as [[MO 461592](#)].

<sup>1</sup>Specifically for  $R$  and  $S$ , not  $\text{Lift}_S$  the functor.

**00NV 8.5.15 Internal Right Kan Extensions**

Let  $A$ ,  $B$ , and  $X$  be sets and let  $R: A \nrightarrow B$  and  $F: A \nrightarrow X$  be relations.

02FH

**MOTIVATION 8.5.15.1.1 ► SETTING FOR INTERNAL RIGHT KAN EXTENSIONS IN Rel**

We want to understand internal right Kan extensions in **Rel**, which look like this:

Note in particular here that  $F: A \nrightarrow X$  is a relation from  $A$  to  $X$ . These will form a functor

$$\text{Ran}_R: \mathbf{Rel}(A, X) \rightarrow \mathbf{Rel}(B, X)$$

that is right adjoint to the precomposition by  $R$  functor

$$R^*: \mathbf{Rel}(B, X) \rightarrow \mathbf{Rel}(A, X).$$

00NW

**PROPOSITION 8.5.15.1.2 ► INTERNAL RIGHT KAN EXTENSIONS IN  $\mathbf{Rel}$** 

The internal right Kan extension of  $F$  along  $R$  is the relation  $\text{Ran}_R(F)$  described as follows:

02A7

- Viewing relations from  $B$  to  $X$  as subsets of  $B \times X$ , we have

$$\text{Ran}_R(F) = \left\{ (b, x) \in B \times X \mid \begin{array}{l} \text{for each } a \in A, \text{ if } a \sim_R b, \\ \text{then we have } a \sim_F x \end{array} \right\}.$$

02A8

- Viewing relations as functions  $B \times X \rightarrow \{\text{true}, \text{false}\}$ , we have

$$\begin{aligned} (\text{Ran}_R(F))^{-1}_{-2} &= \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_a^{-2}, F_a^{-1}) \\ &= \bigwedge_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_a^{-2}, F_a^{-1}), \end{aligned}$$

where the meet  $\wedge$  is taken in the poset  $(\{\text{true}, \text{false}\}, \preceq)$  of Definition 3.2.2.1.3.

02FL

- Viewing relations as functions  $B \rightarrow \mathcal{P}(X)$ , we have

$$\text{Ran}_R(F) = \text{Ran}_{\chi'_A}(F) \circ R^{-1}, \quad \begin{array}{ccc} A & \xrightarrow{F} & \mathcal{P}(X), \\ \chi_A \downarrow & \swarrow & \nearrow \text{Ran}_{\chi_A}(F) \\ B & \xrightarrow[R^{-1}]{} & \mathcal{P}(A)^{\text{op}} \end{array}$$

where  $\text{Ran}_{\chi'_B}(F)$  is computed by the formula

$$\begin{aligned} [\text{Ran}_{\chi'_A}(F)](V) &\cong \int_{a \in A} \chi_{\mathcal{P}(A)^{\text{op}}}(V, \chi_a) \pitchfork F(a) \\ &\cong \int_{a \in A} \chi_{\mathcal{P}(A)}(\chi_a, V) \pitchfork F(a) \\ &\cong \int_{a \in A} \chi_V(a) \pitchfork F(a) \\ &\cong \bigcap_{a \in A} \chi_V(a) \pitchfork F(a) \\ &\cong \bigcap_{a \in V} F(a) \end{aligned}$$

for each  $V \in \mathcal{P}(B)$ , so we have

$$[\text{Ran}_R(F)](b) = \bigcap_{a \in R^{-1}(b)} F(a)$$

for each  $b \in B$ .

#### PROOF 8.5.15.1.3 ► PROOF OF PROPOSITION 8.5.15.1.2

We have

$$\begin{aligned} \text{Hom}_{\mathbf{Rel}(A,X)}(F \diamond R, T) &\cong \int_{a \in A} \int_{x \in X} \mathbf{Hom}_{\{\text{t},\text{f}\}}((F \diamond R)_a^x, T_a^x) \\ &\cong \int_{a \in A} \int_{x \in X} \mathbf{Hom}_{\{\text{t},\text{f}\}}\left(\left(\int^{b \in B} F_b^x \times R_a^b\right), T_a^x\right) \\ &\cong \int_{a \in A} \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t},\text{f}\}}(F_b^x \times R_a^b, T_a^x) \\ &\cong \int_{a \in A} \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t},\text{f}\}}(F_b^x, \mathbf{Hom}_{\{\text{t},\text{f}\}}(R_a^b, T_a^x)) \\ &\cong \int_{b \in B} \int_{x \in X} \int_{a \in A} \mathbf{Hom}_{\{\text{t},\text{f}\}}(F_b^x, \mathbf{Hom}_{\{\text{t},\text{f}\}}(R_a^b, T_a^x)) \\ &\cong \int_{b \in B} \int_{x \in X} \mathbf{Hom}_{\{\text{t},\text{f}\}}(F_b^x, \int_{a \in A} \mathbf{Hom}_{\{\text{t},\text{f}\}}(R_a^b, T_a^x)) \\ &\cong \text{Hom}_{\mathbf{Rel}(B,X)}(F, \int_{a \in A} \mathbf{Hom}_{\{\text{t},\text{f}\}}(R_a^{-2}, T_a^{-1})) \end{aligned}$$

naturally in each  $F \in \mathbf{Rel}(B, X)$  and each  $T \in \mathbf{Rel}(A, X)$ , showing that

$$\int_{a \in A} \mathbf{Hom}_{\{\text{t},\text{f}\}}(R_a^{-2}, T_a^{-1})$$

is right adjoint to the precomposition functor  $- \diamond R$ , being thus the right Kan extension along  $R$ . Here we have used the following results, respectively (i.e. for each  $\cong$  sign):

- 02A9 1. Item 1 of Proposition 8.1.1.1.7.
- 02AA 2. Definition 8.1.3.1.1.
- 02AB 3. ?? of ??.
- 02AC 4. Proposition 3.2.2.1.5.
- 02AD 5. ?? of ??.
- 02AE 6. ?? of ??.

02AF

7. Item 1 of Proposition 8.1.1.7.

This finishes the proof. 

02FU

**EXAMPLE 8.5.15.1.4 ► EXAMPLES OF INTERNAL RIGHT KAN EXTENSIONS OF RELATIONS**

Here are some examples of internal right Kan extensions of relations.

02FV

1. *Orthogonal Complements.* Let  $A = B = X = \mathcal{V}$  be an inner product space, and let  $R = F = \perp$  be the orthogonality relation, so that we have

$$\begin{aligned} R(v) &= v^\perp \\ F(u) &= u^\perp, \end{aligned}$$

for each  $u, v \in \mathcal{V}$ , where

$$v^\perp \stackrel{\text{def}}{=} \{u \in V \mid v \perp u\}$$

is the orthogonal complement of  $v$ . The right Kan extension  $\text{Ran}_R(F)$  is then given by

$$\begin{aligned} [\text{Ran}_R(F)](v) &= \bigcap_{u \in R^{-1}(v)} F(u) \\ &= \bigcap_{\substack{u \in V \\ u \perp v}} u^\perp \\ &= (v^\perp)^\perp, \end{aligned}$$

the double orthogonal complement. In particular:

- If  $\mathcal{V}$  is finite-dimensional, then  $[\text{Ran}_R(F)](v) = \text{Span}(v)$ .
- If  $\mathcal{V}$  is a Hilbert space, then  $[\text{Ran}_R(F)](v) = \overline{\text{Span}(v)}$ .

02FW

2. *Galois Connections and Closure Operators.* Let:

- $B = X = (P, \preceq_P)$  and  $A = (Q, \preceq_Q)$  be posets;
- $(f, g)$  be a Galois connection (adjunction) between  $P$  and  $Q$ ;
- $R, F: Q \nrightarrow P$  be the relations defined by

$$\begin{aligned} R(q) &\stackrel{\text{def}}{=} \{p \in P \mid q \preceq_Q f(p)\}, \\ F(q) &\stackrel{\text{def}}{=} \{p \in P \mid p \preceq_P g(q)\} \end{aligned}$$

for each  $q \in Q$ .

We have

$$\begin{aligned} [\text{Ran}_R(F)](p) &= \bigcap_{q \in R^{-1}(p)} F(q) \\ &= \bigcap_{\substack{q \in Q \\ q \preceq_Q f(p)}} \{p \in P \mid p \preceq_P g(q)\} \\ &= \{p \in P \mid p \preceq_P g(f(q))\} \\ &= \downarrow g(f(p)), \end{aligned}$$

the down set of  $g(f(p))$ . In other words,  $\text{Ran}_R(F)$  is the closure operator on  $P$  associated with the Galois connection  $(f, g)$ .

02FX

### PROPOSITION 8.5.15.1.5 ► PROPERTIES OF INTERNAL RIGHT KAN EXTENSIONS IN **Rel**

Let  $A, B, C$  and  $X$  be sets and let  $R: A \rightarrow B$ ,  $S: B \rightarrow C$ , and  $F: A \rightarrow X$  be relations.

02FY

1. *Functionality.* The assignments  $R, F, (R, F) \mapsto \text{Ran}_R(F)$  define functors

$$\begin{aligned} \text{Ran}_{(-)}(F) : \quad \text{Rel}(A, B)^{\text{op}} &\rightarrow \text{Rel}(B, X), \\ \text{Ran}_R : \quad \text{Rel}(A, X) &\rightarrow \text{Rel}(B, X), \\ \text{Ran}_{(-1)}(-_2) : \text{Rel}(A, X) \times \text{Rel}(A, B)^{\text{op}} &\rightarrow \text{Rel}(B, X). \end{aligned}$$

In other words, given relations

$$A \xrightarrow[R_1]{\quad\quad} B \quad A \xrightarrow[F_1]{\quad\quad} X,$$

if  $R_1 \subset R_2$  and  $F_1 \subset F_2$ , then  $\text{Ran}_{R_2}(F_1) \subset \text{Ran}_{R_1}(F_2)$ .

02FZ

2. *Interaction With Composition.* We have

$$\text{Ran}_{S \circ R}(F) = \text{Ran}_S(\text{Ran}_R(F))$$

and an equality

of pasting diagrams in **Rel**.

02G0

3. *Interaction With Converses.* We have

$$\text{Ran}_R(F)^\dagger = \text{Rift}_{R^\dagger}(F^\dagger).$$

02G1

4. *Interaction With Inverse Images.* We have

$$[\text{Ran}_R(F)]^{-1}(x) = \{b \in B \mid R^{-1}(b) \subset F^{-1}(x)\}$$

for each  $x \in X$ .

#### PROOF 8.5.15.1.6 ► PROOF OF PROPOSITION 8.5.15.1.5

##### Item 1: Functoriality

We have

$$\begin{aligned} [\text{Ran}_{R_2}(F_1)](b) &= \bigcap_{a \in R_2^{-1}(b)} F_1(a) \\ &\subset \bigcap_{a \in R_1^{-1}(b)} F_1(a) \\ &\subset \bigcap_{a \in R_1^{-1}(b)} F_2(a) \\ &= [\text{Ran}_{R_1}(F_2)](b) \end{aligned}$$

for each  $b \in B$ , so we therefore have  $\text{Ran}_{R_2}(F_1) \subset \text{Ran}_{R_1}(F_2)$ .

##### Item 2: Interaction With Composition

This holds in a general bicategory with the necessary right Kan extensions, being therefore a special case of ??.

##### Item 3: Interaction With Converses

We have

$$\begin{aligned} [\text{Rift}_{R^\dagger}(F^\dagger)](x) &= \{b \in B \mid R^\dagger(b) \subset F^\dagger(x)\} \\ &= \{b \in B \mid R^{-1}(b) \subset F^{-1}(x)\} \\ &= \text{Ran}_R(F)^{-1}(x) \\ &= \text{Ran}_R(F)^\dagger(x) \end{aligned}$$

where we have used [Proposition 8.5.16.1.2](#) and [Item 4](#).

##### Item 4: Interaction With Inverse Images

We proceed in a few steps.

- We have  $b \in [\text{Ran}_R(F)]^{-1}(x)$  iff, for each  $a \in R^{-1}(b)$ , we have  $b \in F(a)$ .
- This holds iff, for each  $a \in R^{-1}(b)$ , we have  $a \in F^{-1}(b)$ .
- This holds iff  $R^{-1}(b) \subset F^{-1}(b)$ .

This finishes the proof. 

### 00NX 8.5.16 Internal Right Kan Lifts

Let  $A$ ,  $B$ , and  $X$  be sets and let  $R: A \nrightarrow B$  and  $F: X \nrightarrow B$  be relations.

02G2

#### MOTIVATION 8.5.16.1.1 ► SETTING FOR INTERNAL RIGHT KAN LIFTS IN Rel

We want to understand internal right Kan lifts in **Rel**, which look like this:

$$\begin{array}{ccc} & A & \\ \text{Rift}_R(F) & \nearrow \swarrow & \downarrow R \\ X & \xrightarrow[F]{\quad} & B. \end{array}$$

Note in particular here that  $F: B \nrightarrow X$  is a relation from  $B$  to  $X$ . These will form a functor

$$\text{Rift}_R: \mathbf{Rel}(X, B) \rightarrow \mathbf{Rel}(X, A)$$

that is right adjoint to the postcomposition by  $R$  functor

$$R_*: \mathbf{Rel}(X, A) \rightarrow \mathbf{Rel}(X, B).$$

00NY

#### PROPOSITION 8.5.16.1.2 ► INTERNAL RIGHT KAN LIFTS IN Rel

The internal right Kan lift of  $F$  along  $R$  is the relation  $\text{Rift}_R(F)$  described as follows:

02AG

- Viewing relations from  $X$  to  $A$  as subsets of  $X \times A$ , we have

$$\text{Rift}_R(F) = \left\{ (x, a) \in X \times A \mid \begin{array}{l} \text{for each } b \in B, \text{ if } a \sim_R b, \\ \text{then we have } x \sim_F b \end{array} \right\}.$$

02AH

- Viewing relations as functions  $X \times A \rightarrow \{\text{true, false}\}$ , we have

$$(\text{Rift}_R(F))_{-2}^{-1} = \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_{-1}^b, F_{-2}^b)$$

$$= \bigwedge_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_{-1}^b, F_{-2}^b),$$

where the meet  $\wedge$  is taken in the poset  $(\{\text{true, false}\}, \preceq)$  of [Definition 3.2.2.1.3](#).

**02G3** 3. Viewing relations as functions  $X \rightarrow \mathcal{P}(A)$ , we have

$$[\text{Rift}_R(F)](x) = \{a \in A \mid R(a) \subset F(x)\}$$

for each  $a \in A$ .

#### PROOF 8.5.16.1.3 ► PROOF OF PROPOSITION 8.5.16.1.2

We have

$$\begin{aligned} \mathbf{Hom}_{\mathbf{Rel}(X,B)}(R \diamond F, T) &\cong \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}((R \diamond F)_x^b, T_x^b) \\ &\cong \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}((\int^{a \in A} R_a^b \times F_x^a), T_x^b) \\ &\cong \int_{x \in X} \int_{b \in B} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_a^b \times F_x^a, T_x^b) \\ &\cong \int_{x \in X} \int_{b \in B} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(F_x^a, \mathbf{Hom}_{\{\text{t,f}\}}(R_a^b, T_x^b)) \\ &\cong \int_{x \in X} \int_{a \in A} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(F_x^a, \mathbf{Hom}_{\{\text{t,f}\}}(R_a^b, T_x^b)) \\ &\cong \int_{x \in X} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(F_x^a, \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_a^b, T_x^b)) \\ &\cong \mathbf{Hom}_{\mathbf{Rel}(X,A)}(F, \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_{-1}^b, T_{-2}^b)) \end{aligned}$$

naturally in each  $F \in \mathbf{Rel}(X, A)$  and each  $T \in \mathbf{Rel}(X, B)$ , showing that

$$\int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_{-1}^b, F_{-2}^b)$$

is right adjoint to the postcomposition functor  $R \diamond -$ , being thus the right Kan lift along  $R$ . Here we have used the following results, respectively (i.e. for each  $\cong$  sign):

**02AJ** 1. [Item 1 of Proposition 8.1.1.7](#).

**02AK** 2. [Definition 8.1.3.1.1](#).

**02AL** 3. ?? of ??.

02AM 4. [Proposition 3.2.2.1.5.](#)

02AN 5. ?? of ??.

02AP 6. ?? of ??.

02AQ 7. [Item 1 of Proposition 8.1.1.1.7.](#)

This finishes the proof. 

02G4 **EXAMPLE 8.5.16.1.4 ► EXAMPLES OF INTERNAL RIGHT KAN EXTENSIONS OF RELATIONS**

Here are some examples of internal right Kan lifts of relations.

02G5 1. *Pullbacks.* Let  $p: A \rightarrow B$  and  $f: X \rightarrow B$  be functions. We have

$$\begin{aligned} [\text{Rift}_{\text{Gr}(p)}(\text{Gr}(f))](x) &= \{a \in A \mid [\text{Gr}(p)](a) \subset [\text{Gr}(f)](x)\} \\ &= \{a \in A \mid p(a) = f(x)\}. \end{aligned}$$

Thus, as a subset of  $X \times A$ , the right Kan lift  $\text{Rift}_{\text{Gr}(p)}(\text{Gr}(f))$  corresponds precisely to the pullback  $X \times_B A$  of  $X$  and  $A$  along  $p$  and  $f$  of [Section 4.1.4](#).

02G6 **PROPOSITION 8.5.16.1.5 ► PROPERTIES OF INTERNAL RIGHT KAN LIFTS IN Rel**

Let  $A, B, C$  and  $X$  be sets and let  $R: A \rightarrow B$ ,  $S: B \rightarrow C$ , and  $F: X \rightarrow B$  be relations.

02G7 1. *Functionality.* The assignments  $R, F, (R, F) \mapsto \text{Rift}_R(F)$  define functors

$$\begin{aligned} \text{Rift}_{(-)}(F) : \quad \text{Rel}(A, B)^{\text{op}} &\rightarrow \text{Rel}(B, X), \\ \text{Rift}_R : \quad \text{Rel}(A, X) &\rightarrow \text{Rel}(B, X), \\ \text{Rift}_{(-1)}(-_2) : \text{Rel}(A, X) \times \text{Rel}(A, B)^{\text{op}} &\rightarrow \text{Rel}(B, X). \end{aligned}$$

In other words, given relations

$$A \begin{array}{c} R_1 \\[-1ex] \bot\!\!\!/\end{array} B \qquad A \begin{array}{c} F_1 \\[-1ex] \bot\!\!\!/\end{array} X,$$

if  $R_1 \subset R_2$  and  $F_1 \subset F_2$ , then  $\text{Rift}_{R_2}(F_1) \subset \text{Rift}_{R_1}(F_2)$ .

02G8 2. *Interaction With Composition.* We have

$$\text{Rift}_{S \circ R}(F) = \text{Rift}_R(\text{Ran}_S(F))$$

and an equality

$$\begin{array}{ccc}
 \begin{array}{c}
 \text{Rift}_R(\text{Rift}_S(F)) \\
 \text{Rift}_S(F) \\
 X \xrightarrow[F]{\quad} C
 \end{array}
 & = &
 \begin{array}{c}
 \text{Rift}_{S \circ R}(F) \\
 X \xrightarrow[F]{\quad} C
 \end{array}
 \end{array}$$

of pasting diagrams in **Rel**.

02G9

3. *Interaction With Converses.* We have

$$\text{Rift}_R(F)^\dagger = \text{Ran}_{R^\dagger}(F^\dagger).$$

02GA

4. *Interaction With Inverse Images.* We have

$$\begin{array}{c}
 \text{Rift}_R(F)^\dagger = \text{Ran}_{\chi'_B}(F^\dagger) \circ R, \\
 A \xrightarrow[R]{\quad} \mathcal{P}(B)^{\text{op}}
 \end{array}
 \quad
 \begin{array}{c}
 B \xrightarrow[F^\dagger]{\quad} \mathcal{P}(X), \\
 \chi_B \downarrow \quad \swarrow \quad \text{Ran}_{\chi_A}(F^{-1})
 \end{array}$$

where  $\text{Ran}_{\chi_A}(F^\dagger)$  is computed by the formula

$$\begin{aligned}
 [\text{Ran}_{\chi_A}(F^\dagger)](U) &\cong \int_{a \in A} \chi_{\mathcal{P}(B)^{\text{op}}}(U, \chi_a) \pitchfork F^\dagger(a) \\
 &\cong \int_{a \in A} \chi_{\mathcal{P}(B)}(\chi_a, U) \pitchfork F^{-1}(a) \\
 &\cong \int_{a \in A} \chi_U(a) \pitchfork F(a) \\
 &\cong \bigcap_{a \in A} \chi_U(a) \pitchfork F(a) \\
 &\cong \bigcap_{a \in U} F(a)
 \end{aligned}$$

for each  $U \in \mathcal{P}(A)$ , so we have

$$[\text{Rift}_R(F)]^{-1}(a) = \bigcap_{b \in R(a)} F^{-1}(b)$$

for each  $a \in A$ .

**PROOF 8.5.16.1.6 ► PROOF OF PROPOSITION 8.5.16.1.5****Item 1: Functoriality**

We have

$$\begin{aligned} [\text{Rift}_{R_2}(F_1)](x) &= \{a \in A \mid R_2(a) \subset F_1(x)\} \\ &\subset \{a \in A \mid R_1(a) \subset F_1(x)\} \\ &\subset \{a \in A \mid R_1(a) \subset F_2(x)\} \\ &= \text{Rift}_{R_1}(F_2) \end{aligned}$$

for each  $x \in X$ , so we therefore have  $\text{Rift}_{R_2}(F_1) \subset \text{Rift}_{R_1}(F_2)$ .

**Item 2: Interaction With Composition**

This holds in a general bicategory with the necessary right Kan lifts, being therefore a special case of ??.

**Item 3: Interaction With Converses**

This follows from Item 3 of Proposition 8.5.15.1.5 by duality.

**Item 4: Interaction With Inverse Images**

We proceed in a few steps.

- We have  $x \in \text{Rift}_R(F)^\dagger(a)$  iff  $a \in \text{Rift}_R(F)(x)$ .
- This holds iff  $R(a) \subset F(x)$ .
- This holds iff, for each  $b \in R(a)$ , we have  $b \in F(x)$ .
- This holds iff, for each  $b \in R(a)$ , we have  $x \in F^{-1}(b)$ .
- This holds iff  $x \in \bigcap_{b \in R(a)} F^{-1}(b)$ .

This finishes the proof. 

**00MJ 8.5.17 Closedness**

00MK

**PROPOSITION 8.5.17.1.1 ► CLOSEDNESS OF Rel**

The 2-category **Rel** is a closed bicategory, there being, for each  $R: A \rightarrow B$  and set  $X$ , a pair of adjunctions

$$(R^* \dashv \text{Ran}_R): \quad \text{Rel}(B, X) \begin{array}{c} \xrightarrow{R^*} \\ \perp \\ \xleftarrow{\text{Ran}_R} \end{array} \text{Rel}(A, X),$$

$$(R_! \dashv \text{Rift}_R): \quad \text{Rel}(X, A) \begin{array}{c} \xrightarrow{R_!} \\ \perp \\ \xleftarrow{\text{Rift}_R} \end{array} \text{Rel}(X, B),$$

witnessed by bijections

$$\begin{aligned} \text{Rel}(S \diamond R, T) &\cong \text{Rel}(S, \text{Ran}_R(T)), \\ \text{Rel}(R \diamond U, V) &\cong \text{Rel}(U, \text{Rift}_R(V)), \end{aligned}$$

natural in  $S \in \text{Rel}(B, X)$ ,  $T \in \text{Rel}(A, X)$ ,  $U \in \text{Rel}(X, A)$ , and  $V \in \text{Rel}(X, B)$ .

#### PROOF 8.5.17.1.2 ► PROOF OF PROPOSITION 8.5.17.1.1

This follows from ????.



### 00ML 8.5.18 Rel as a Category of Free Algebras

#### 00MM PROPOSITION 8.5.18.1.1 ► Rel AS A CATEGORY OF FREE ALGEBRAS

We have an isomorphism of categories

$$\text{Rel} \cong \text{FreeAlg}_{\mathcal{P}_!}(\text{Sets}),$$

where  $\mathcal{P}_!$  is the powerset monad of ??.

#### PROOF 8.5.18.1.2 ► PROOF OF PROPOSITION 8.5.18.1.1

Omitted.



**02GB 8.6 Properties of the 2-Category of Relations With Apartness Composition**

**02GC 8.6.1 Self-Duality**

**02GD PROPOSITION 8.6.1.1.1 ► SELF-DUALITY FOR THE (2-)CATEGORY OF RELATIONS WITH APARTNESS COMPOSITION**

The 2-/category of relations with apartness-composition-is self-dual:

- 02GE** 1. *Self-Duality I.* We have an isomorphism

$$(\mathbf{Rel}^{\square})^{\text{op}} \cong \mathbf{Rel}^{\square}$$

of categories.

- 02GF** 2. *Self-Duality II.* We have a 2-isomorphism

$$(\mathbf{Rel}^{\square})^{\text{op}} \cong \mathbf{Rel}^{\square}$$

of 2-categories.

**PROOF 8.6.1.2 ► PROOF OF PROPOSITION 8.6.1.1.1**

**Item 1: Self-Duality I**

We claim that the functor

$$(-)^{\dagger}: (\mathbf{Rel}^{\square})^{\text{op}} \rightarrow \mathbf{Rel}^{\square}$$

given by the identity on objects and by  $R \mapsto R^{\dagger}$  on morphisms is an isomorphism of categories. Note that this is indeed a functor by [Items 4](#) and [7 of Proposition 8.1.5.1.3](#).

By [Item 1 of Proposition 11.6.8.1.3](#), it suffices to show that  $(-)^{\dagger}$  is bijective on objects (which follows by definition) and fully faithful. Indeed, the map

$$(-)^{\dagger}: \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(B, A)$$

defined by the assignment  $R \mapsto R^{\dagger}$  is a bijection by [Item 5 of Proposition 8.1.5.1.3](#), showing  $(-)^{\dagger}$  to be fully faithful.

**Item 2: Self-Duality II**

We claim that the 2-functor

$$(-)^{\dagger}: \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$$

given by the identity on objects, by  $R \mapsto R^\dagger$  on morphisms, and by preserving inclusions on 2-morphisms via [Item 1 of Proposition 8.1.5.1.3](#), is an isomorphism of categories.

By ??, it suffices to show that  $(-)^{\dagger}$  is:

- Bijective on objects, which follows by definition.
- Bijective on 1-morphisms, which was shown in [Item 1](#).
- Bijective on 2-morphisms, which follows from [Item 1 of Proposition 8.1.5.1.3](#).

Thus  $(-)^{\dagger}$  is indeed a 2-isomorphism of categories. 

### 02GG 8.6.2 Isomorphisms and Equivalences

Let  $R: A \rightarrow B$  be a relation from  $A$  to  $B$ , and recall that  $R^c \stackrel{\text{def}}{=} B \times A \setminus R$ .

#### 02GH LEMMA 8.6.2.1.1 ► CONDITIONS INVOLVING A RELATION AND ITS CONVERSE II

The conditions below are row-wise equivalent:

CONDITION	INCLUSION
$R^c$ is functional	$\nabla_B \subset R \square R^\dagger$
$R^c$ is total	$R \square R^\dagger \subset \nabla_A$
$R^c$ is injective	$\nabla_A \subset R^\dagger \square R$
$R^c$ is surjective	$R^\dagger \square R \subset \nabla_B$

#### PROOF 8.6.2.1.2 ► PROOF OF LEMMA 8.6.2.1.1

This follows from [Lemma 8.5.2.1.1](#) and [Item 4 of Proposition 8.1.4.1.3](#). For instance:

- Suppose we have  $R \square R^\dagger \subset \nabla_B$ .
- Taking complements, we obtain  $\nabla_B^c \subset (R \square R^\dagger)^c$ .
- Applying [Item 4 of Proposition 8.1.4.1.3](#), this becomes  $\Delta_B \subset R^c \diamond (R^\dagger)^c$ .
- Then, by [Lemma 8.5.2.1.1](#), this is equivalent to  $R^c$  being total.

The proof of the other equivalences is similar, and thus omitted. 

## 8.6. Properties of the 2-Category of Relations With Apartness Composition

650

02SK

### REMARK 8.6.2.1.3 ► UNWINDING LEMMA 8.6.2.1.1

The statements in [Lemma 8.6.2.1.1](#) unwind to the following:

INCLUSION	QUANTIFIER	CONDITION
$\nabla_B \subset R \square R^\dagger$	For each $b_1, b_2 \in B$	If $b_1 \neq b_2$ , then, for each $a \in A$ , we have $a \sim_R b_1$ or $a \sim_R b_2$ .
$R \square R^\dagger \subset \nabla_B$	For each $b_1, b_2 \in B$	If, for each $a \in A$ , $a \sim_R b_1$ or $a \sim_R b_2$ , then $b_1 \neq b_2$ .
$\nabla_A \subset R^\dagger \square R$	For each $a_1, a_2 \in A$	If $a_1 \neq a_2$ , then, for each $b \in B$ , we have $a_1 \sim_R b$ or $a_2 \sim_R b$ .
$R^\dagger \square R \subset \nabla_A$	For each $a_1, a_2 \in A$	If, for each $b \in B$ , $a_1 \sim_R b$ or $a_2 \sim_R b$ , then $a_1 \neq a_2$ .

Equivalently:

INCLUSION	QUANTIFIER	IF	THEN
$\nabla_B \subset R \square R^\dagger$	For each $b_1, b_2 \in B$	$b_1 \neq b_2$	$R^{-1}(b_1) \cup R^{-1}(b_2) = A$
$R \square R^\dagger \subset \nabla_B$	For each $b_1, b_2 \in B$	$R^{-1}(b_1) \cup R^{-1}(b_2) = A$	$b_1 \neq b_2$
$\nabla_A \subset R^\dagger \square R$	For each $a_1, a_2 \in A$	$a_1 \neq a_2$	$R(a_1) \cup R(a_2) = B$
$R^\dagger \square R \subset \nabla_A$	For each $a_1, a_2 \in A$	$R(a_1) \cup R(a_2) = B$	$a_1 \neq a_2$

02GJ

### PROPOSITION 8.6.2.1.4 ► ISOMORPHISMS AND EQUIVALENCES IN $\mathbf{Rel}^\square$

The following conditions are equivalent:

02GK

1. The relation  $R: A \rightarrow B$  is an equivalence in  $\mathbf{Rel}^\square$ , i.e.:

(★) There exists a relation  $R^{-1}: B \rightarrow A$  from  $B$  to  $A$  together with isomorphisms

$$R^{-1} \square R \cong \nabla_A,$$

$$R \square R^{-1} \cong \nabla_B.$$

02GL

2. The relation  $R: A \rightarrow B$  is an isomorphism in  $\mathbf{Rel}$ , i.e.:

(★) There exists a relation  $R^{-1}: B \rightarrow A$  from  $B$  to  $A$  such that we have

$$R^{-1} \square R = \nabla_A,$$

$$R \square R^{-1} = \nabla_B.$$

02GM 3. There exists a bijection  $f: B \xrightarrow{\sim} A$  with  $R^c = f^{-1}$ .

**PROOF 8.6.2.1.5 ▶ PROOF OF PROPOSITION 8.6.2.1.4**

This follows from [Proposition 8.5.2.1.3](#) and [Item 4 of Proposition 8.1.4.1.3](#). 

02GN **8.6.3 Internal Adjunctions**

Let  $A$  and  $B$  be sets.

02L3 **PROPOSITION 8.6.3.1.1 ▶ ADJUNCTIONS IN  $\mathbf{Rel}^\square$**

We have a natural bijection

$$\left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel}^\square \\ \text{from } A \text{ to } B \end{array} \right\} \cong \left\{ \begin{array}{l} \text{Functions} \\ \text{from } B \text{ to } A \end{array} \right\},$$

with every adjunction in  $\mathbf{Rel}^\square$  being of the form  $(f^{-1})^c \dashv \text{Gr}(f)^c$  for some function  $f: B \rightarrow A$ .

**PROOF 8.6.3.1.2 ▶ PROOF OF PROPOSITION 8.6.3.1.1**

This follows from [Proposition 8.5.3.1.1](#) and [Item 4 of Proposition 8.1.4.1.3](#). 

02GP **8.6.4 Internal Monads**

Let  $X$  be a set.

02L4 **PROPOSITION 8.6.4.1.1 ▶ INTERNAL MONADS IN  $\mathbf{Rel}^\square$**

We have a natural identification

$$\left\{ \begin{array}{l} \text{Monads in} \\ \mathbf{Rel}^\square \text{ on } X \end{array} \right\} \cong \{\text{Subsets of } X\}.$$

**PROOF 8.6.4.1.2 ► PROOF OF PROPOSITION 8.6.4.1.1**

This follows from [Proposition 8.6.4.1.1](#) and [Item 4 of Proposition 8.1.4.1.3](#).

**02GQ 8.6.5 Internal Comonads**

Let  $X$  be a set.

**02L5**

**PROPOSITION 8.6.5.1.1 ► INTERNAL COMONADS IN  $\mathbf{Rel}^\square$**

We have a natural identification

$$\left\{ \begin{array}{l} \text{Comonads in} \\ \mathbf{Rel}^\square \text{ on } X \end{array} \right\} \cong \{\text{Strict total orders on } X\}.$$


**PROOF 8.6.5.1.2 ► PROOF OF PROPOSITION 8.6.5.1.1**

A comonad in  $\mathbf{Rel}^\square$  on  $X$  consists of a relation  $R: X \rightarrow X$  together with maps

$$\Delta_R: R \subset R \square R,$$

$$\epsilon_R: R \subset \nabla_X$$

making the diagrams



commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic (?? of ??), and hence all that is left is the data of the two maps  $\mu_R$  and  $\eta_R$ , which correspond respectively to the following conditions:

**02L6**

1. For each  $x, z \in X$ , if  $x \sim_R z$ , then, for each  $y \in X$ , we have  $x \sim_R y$  or  $y \sim_R z$ .
2. For each  $x, y \in X$ , if  $x \sim_R y$ , then  $x \neq y$ .

**02L7**

Replacing  $\sim_R$  with  $<_R$  and taking the contrapositive of each condition, we obtain:

- 02L8 1. For each  $x, z \in X$ , if there exists some  $y \in X$  such that  $x <_R y$  and  $y <_R z$ , then  $x <_R z$ .
- 02L9 2. For each  $x \in X$ , we have  $x \not<_R x$ .

These are exactly the requirements for  $R$  to be a strict linear order (??). Conversely, any strict linear order  $<_R$  gives rise to a pair of maps  $\Delta_{<_R}$  and  $\epsilon_{<_R}$ , forming a comonad on  $X$ . 


02LA

#### EXAMPLE 8.6.5.1.3 ► DENSITY COMONADS IN Rel

Let  $R: A \nrightarrow B$  be a relation.

02LB

1. The codensity monad  $\text{Ran}_R(R): B \nrightarrow B$  is given by

$$[\text{Ran}_R(R)](b) = \bigcap_{a \in R^{-1}(b)} R(b)$$


for each  $b \in B$ . Thus, it corresponds to the preorder

$$\preceq_{\text{Ran}_R(R)}: B \times B \rightarrow \{\text{t}, \text{f}\}$$

on  $B$  obtained by declaring  $b \preceq_{\text{Ran}_R(R)} b'$  iff the following equivalent conditions are satisfied:

02LC


- (a) For each  $a \in A$ , if  $a \sim_R b$ , then  $a \sim_R b'$ .

02LD

- (b) We have  $R^{-1}(b) \subset R^{-1}(b')$ .

02LE

2. The dual codensity monad  $\text{Rift}_R(R): A \nrightarrow A$  is given by

$$[\text{Rift}_R(R)](a) = \{a' \in A \mid R(a') \subset R(a)\}$$


for each  $a \in A$ . Thus, it corresponds to the preorder

$$\preceq_{\text{Rift}_R(R)}: A \times A \rightarrow \{\text{t}, \text{f}\}$$

on  $A$  obtained by declaring  $a \preceq_{\text{Rift}_R(R)} a'$  iff the following equivalent conditions are satisfied:

02LF

(a) For each  $a \in A$ , if  $a \sim_R b$ , then  $a' \sim_R b$ .

02LG

(b) We have  $R(a') \subset R(a)$ .

02LH 8.6.6 Modules Over Internal Monads

02LJ 8.6.7 Comodules Over Internal Comonads

02LK 8.6.8 Eilenberg–Moore and Kleisli Objects

02GS 8.6.9 Monomorphisms

02GT 8.6.10 2-Categorical Monomorphisms

02GU 8.6.11 Epimorphisms

02GV 8.6.12 2-Categorical Epimorphisms

02GW 8.6.13 Co/Limits

This will be expanded later on.

02LL 8.6.14 Internal Left Kan Extensions

02LM 8.6.15 Internal Left Kan Lifts

02LN 8.6.16 Internal Right Kan Extensions

02LP 8.6.17 Internal Right Kan Lifts

02LQ 8.6.18 Coclosedness

**00R7 8.7 The Adjoint Pairs  $R_! \dashv R_{-1}$  and  $R^{-1} \dashv R_*$** **00R8 8.7.1 Direct Images**

Let  $X$  and  $Y$  be sets and let  $R: X \nrightarrow Y$  be a relation.

00R9

**DEFINITION 8.7.1.1.1 ► DIRECT IMAGES**

The **direct image function associated to  $R$**  is the function<sup>1</sup>

$$R_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by<sup>2</sup>

$$R_!(U) \stackrel{\text{def}}{=} \bigcup_{a \in U} R(a)$$

$$= \left\{ b \in Y \mid \begin{array}{l} \text{there exists some } a \in U \\ \text{such that } b \in R(a) \end{array} \right\}$$

for each  $U \in \mathcal{P}(X)$ .

<sup>1</sup>Further Notation: Also written simply  $R: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ .

<sup>2</sup>Further Terminology: The set  $R(U)$  is called the **direct image of  $U$  by  $R$** .

#### WARNING 8.7.1.2 ► NOTATION FOR DIRECT IMAGES IS CONFUSING



Notation for direct images between powersets is tricky; see [Warning 4.6.1.3](#). Here we'll try to align our notation for relations with that for functions.

00RA

#### REMARK 8.7.1.3 ► UNWINDING DEFINITION 8.7.1.1

Identifying subsets of  $X$  with relations from pt to  $X$  via [Item 3 of Proposition 4.4.1.1.4](#), we see that the direct image function associated to  $R$  is equivalently the function

$$R_!: \underbrace{\mathcal{P}(X)}_{\cong \text{Rel(pt, } X)} \rightarrow \underbrace{\mathcal{P}(Y)}_{\cong \text{Rel(pt, } Y)}$$

defined by

$$R_!(U) \stackrel{\text{def}}{=} R \diamond U$$

for each  $U \in \mathcal{P}(X)$ , where  $R \diamond U$  is the composition

$$\text{pt} \xrightarrow{U} X \xrightarrow{R} Y.$$

00RB

#### PROPOSITION 8.7.1.4 ► PROPERTIES OF DIRECT IMAGE FUNCTIONS

Let  $R: X \rightarrow Y$  be a relation.

00RC

1. *Functionality.* The assignment  $U \mapsto R_!(U)$  defines a functor

$$R_!: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset)$$

where

- *Action on Objects.* For each  $U \in \mathcal{P}(X)$ , we have

$$[R_!] (U) \stackrel{\text{def}}{=} R_!(U).$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(X)$ :

00RD

- If  $U \subset V$ , then  $R_!(U) \subset R_!(V)$ .

2. *Adjointness.* We have an adjunction

$$(R_! \dashv R_{-1}): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{R_!} \\ \perp \\ \xleftarrow{R_{-1}} \end{array} \mathcal{P}(Y),$$

witnessed by:

02H1

(a) Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\hookrightarrow R_{-1} \circ R_!, \\ R_! \circ R_{-1} &\hookrightarrow \text{id}_{\mathcal{P}(Y)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset R_{-1}(R_!(U)), \\ R_!(R_{-1}(V)) &\subset V \end{aligned}$$

indexed by  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$

02H2

(b) A bijections of sets

$$\text{Hom}_{\mathcal{P}(X)}(R_!(U), V) \cong \text{Hom}_{\mathcal{P}(X)}(U, R_{-1}(V)),$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ . In particular:

(★) The following conditions are equivalent:

- We have  $R_!(U) \subset V$ .
- We have  $U \subset R_{-1}(V)$ .

00RE

3. *Preservation of Colimits.* We have an equality of sets

$$R_!(\bigcup_{i \in I} U_i) = \bigcup_{i \in I} R_!(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} R_!(U) \cup R_!(V) &= R_!(U \cup V), \\ R_!(\emptyset) &= \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00RF

4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$R_!(\bigcap_{i \in I} U_i) \subset \bigcap_{i \in I} R_!(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} R_!(U \cap V) &\subset R_!(U) \cap R_!(V), \\ R_!(X) &\subset Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00RG

5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of [Item 1](#) has a symmetric strict monoidal structure

$$(R_!, R_!^\otimes, R_{*|1}^\otimes): (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} R_{*|U,V}^\otimes: R_!(U) \cup R_!(V) &\xrightarrow{=} R_!(U \cup V), \\ R_{*|1}^\otimes: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00RH

6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of [Item 1](#) has a symmetric oplax monoidal structure

$$(R_!, R_!^\otimes, R_{*|1}^\otimes): (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with inclusions

$$\begin{aligned} R_{*|U,V}^\otimes: R_!(U \cap V) &\subset R_!(U) \cap R_!(V), \\ R_{*|1}^\otimes: R_!(X) &\subset Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00RJ

7. *Relation to Codirect Images.* We have

$$R_!(U) = Y \setminus R_*(X \setminus U)$$

for each  $U \in \mathcal{P}(X)$ .

**PROOF 8.7.1.1.5 ▶ PROOF OF PROPOSITION 8.7.1.1.4**

Item 1: Functoriality

Clear.

**Item 2: Adjointness**

This follows from ?? of ??.

**Item 3: Preservation of Colimits**

This follows from **Item 2** and ?? of ??.

**Item 4: Oplax Preservation of Limits**

Omitted.

**Item 5: Symmetric Strict Monoidality With Respect to Unions**

This follows from **Item 3**.

**Item 6: Symmetric Oplax Monoidality With Respect to Intersections**

This follows from **Item 4**.

**Item 7: Relation to Codirect Images**

The proof proceeds in the same way as in the case of functions (?? of [Proposition 4.6.1.1.5](#)): applying **Item 7** of [Proposition 8.7.4.1.4](#) to  $A \setminus U$ , we have

$$\begin{aligned} R_*(X \setminus U) &= Y \setminus R_!(X \setminus (X \setminus U)) \\ &= Y \setminus R_!(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} R_!(U) &= Y \setminus (Y \setminus R_!(U)), \\ &= Y \setminus R_*(X \setminus U), \end{aligned}$$

which finishes the proof. 

00RK

**PROPOSITION 8.7.1.1.6 ► PROPERTIES OF THE DIRECT IMAGE FUNCTION OPERATION**

Let  $R: X \rightarrow Y$  be a relation.

00RL

- Functionality I.* The assignment  $R \mapsto R_!$  defines a function

$$(-)_!: \text{Rel}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

00RM

- Functionality II.* The assignment  $R \mapsto R_!$  defines a function

$$(-)_!: \text{Rel}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

00RN

- Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have<sup>1</sup>

$$(\chi_X)_! = \text{id}_{\mathcal{P}(X)}.$$

00RP

4. *Interaction With Composition.* For each pair of composable relations  $R: X \rightarrow Y$  and  $S: Y \rightarrow C$ , we have<sup>2</sup>

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{R_!} & \mathcal{P}(Y) \\ (S \diamond R)_! = S_! \circ R_!, & \searrow & \downarrow S_! \\ & (S \diamond R)_! & \mathcal{P}(C). \end{array}$$

<sup>1</sup>That is, the postcomposition function

$$(\chi_X)_!: \text{Rel(pt, } X) \rightarrow \text{Rel(pt, } X)$$

is equal to  $\text{id}_{\text{Rel(pt, } X)}$ .

<sup>2</sup>That is, we have

$$\begin{array}{ccc} \text{Rel(pt, } X) & \xrightarrow{R_!} & \text{Rel(pt, } Y) \\ (S \diamond R)_! = S_! \circ R_!, & \searrow & \downarrow S_! \\ & (S \diamond R)_! & \text{Rel(pt, } C). \end{array}$$

#### PROOF 8.7.1.1.7 ► PROOF OF PROPOSITION 8.7.1.1.6

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

Indeed, we have

$$\begin{aligned} (\chi_X)_!(U) &\stackrel{\text{def}}{=} \bigcup_{a \in U} \chi_X(a) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} \{a\} \\ &= U \\ &\stackrel{\text{def}}{=} \text{id}_{\mathcal{P}(X)}(U) \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ . Thus  $(\chi_X)_! = \text{id}_{\mathcal{P}(X)}$ .

Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned}
 (S \diamond R)_!(U) &\stackrel{\text{def}}{=} \bigcup_{a \in U} [S \diamond R](a) \\
 &\stackrel{\text{def}}{=} \bigcup_{a \in U} S(R(a)) \\
 &\stackrel{\text{def}}{=} \bigcup_{a \in U} S_!(R(a)) \\
 &= S_!(\bigcup_{a \in U} R(a)) \\
 &\stackrel{\text{def}}{=} S_!(R_!(U)) \\
 &\stackrel{\text{def}}{=} [S_! \circ R_!](U)
 \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , where we used Item 3 of Proposition 8.7.1.1.4. Thus  $(S \diamond R)_! = S_! \circ R_!$ . □

### 00RQ 8.7.2 Coinverse Images

Let  $X$  and  $Y$  be sets and let  $R: X \rightarrow Y$  be a relation.

00RR

#### DEFINITION 8.7.2.1.1 ► COINVERSE IMAGES

The **coinverse image function associated to  $R$**  is the function

$$R_{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$$

defined by<sup>1</sup>

$$R_{-1}(V) \stackrel{\text{def}}{=} \{a \in X \mid R(a) \subset V\}$$

for each  $V \in \mathcal{P}(Y)$ .

---

<sup>1</sup>Further Terminology: The set  $R_{-1}(V)$  is called the **coinverse image of  $V$  by  $R$** .

00RS

#### REMARK 8.7.2.1.2 ► UNWINDING DEFINITION 8.7.2.1.1

Identifying subsets of  $Y$  with relations from pt to  $Y$  via Item 3 of Proposition 4.4.1.1.4, we see that the inverse image function associated to  $R$  is equivalently the function

$$R_{-1}: \underbrace{\mathcal{P}(Y)}_{\cong \text{Rel(pt, } Y)} \rightarrow \underbrace{\mathcal{P}(X)}_{\cong \text{Rel(pt, } X)}$$

defined by

$$R_{-1}(V) \stackrel{\text{def}}{=} \text{Rift}_R(V), \quad \begin{array}{ccc} X & & \\ \text{Rift}_R(V) & \nearrow \swarrow & \downarrow R \\ \text{pt} & \xrightarrow[V]{} & Y, \end{array}$$

and being explicitly computed by

$$\begin{aligned} R_{-1}(V) &\stackrel{\text{def}}{=} \text{Rift}_R(V) \\ &\cong \int_{b \in Y} \text{Hom}_{\{\text{t,f}\}}(R_{-1}^b, V_{-2}^b), \end{aligned}$$

where we have used ??.

#### PROOF 8.7.2.1.3 ► PROOF OF REMARK 8.7.2.1.2

We have

$$\begin{aligned} \text{Rift}_R(V) &\cong \int_{b \in Y} \text{Hom}_{\{\text{t,f}\}}(R_{-1}^b, V_{-2}^b) \\ &= \left\{ a \in X \mid \int_{b \in Y} \text{Hom}_{\{\text{t,f}\}}(R_a^b, V_\star^b) = \text{true} \right\} \\ &= \left\{ a \in X \mid \begin{array}{l} \text{for each } b \in Y, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\ &\quad \left. \begin{array}{l} 1. \text{ We have } R_a^b = \text{false} \\ 2. \text{ The following conditions hold:} \end{array} \right. \\ &\quad \left. \begin{array}{l} (a) \text{ We have } R_a^b = \text{true} \\ (b) \text{ We have } V_\star^b = \text{true} \end{array} \right\} \\ &= \left\{ a \in X \mid \begin{array}{l} \text{for each } b \in Y, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\ &\quad \left. \begin{array}{l} 1. \text{ We have } b \notin R(a) \\ 2. \text{ The following conditions hold:} \end{array} \right. \\ &\quad \left. \begin{array}{l} (a) \text{ We have } b \in R(a) \\ (b) \text{ We have } b \in V \end{array} \right\} \\ &= \{a \in X \mid \text{for each } b \in R(a), \text{ we have } b \in V\} \\ &= \{a \in X \mid R(a) \subset V\} \end{aligned}$$

$$\stackrel{\text{def}}{=} R_{-1}(V).$$

This finishes the proof. 

00RT

#### PROPOSITION 8.7.2.1.4 ► PROPERTIES OF COINVERSE IMAGES

00RU

Let  $R: X \rightarrow Y$  be a relation.

1. *Functoriality.* The assignment  $V \mapsto R_{-1}(V)$  defines a functor

$$R_{-1}: (\mathcal{P}(Y), \subset) \rightarrow (\mathcal{P}(X), \subset)$$

where

- *Action on Objects.* For each  $V \in \mathcal{P}(Y)$ , we have

$$[R_{-1}](V) \stackrel{\text{def}}{=} R_{-1}(V).$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(Y)$ :

- If  $U \subset V$ , then  $R_{-1}(U) \subset R_{-1}(V)$ .

00RV

2. *Adjointness.* We have an adjunction

$$(R_! \dashv R_{-1}): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{R_!} \\ \perp \\ \xleftarrow{R_{-1}} \end{array} \mathcal{P}(Y),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(X)}(R_!(U), V) \cong \text{Hom}_{\mathcal{P}(X)}(U, R_{-1}(V)),$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ , i.e. such that:

- (★) The following conditions are equivalent:

- We have  $R_!(U) \subset V$ .
- We have  $U \subset R_{-1}(V)$ .

00RW

3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} R_{-1}(U_i) \subset R_{-1}\left(\bigcup_{i \in I} U_i\right),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} R_{-1}(U) \cup R_{-1}(V) &\subset R_{-1}(U \cup V), \\ \emptyset &\subset R_{-1}(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00RX

4. *Preservation of Limits.* We have an equality of sets

$$R_{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} R_{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} R_{-1}(U \cap V) &= R_{-1}(U) \cap R_{-1}(V), \\ R_{-1}(Y) &= Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00RY

5. *Symmetric Lax Monoidality With Respect to Unions.* The codirect image function of [Item 1](#) has a symmetric lax monoidal structure

$$(R_{-1}, R_{-1}^\otimes, R_{-1|1\!\!1}^\otimes): (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} R_{-1|U,V}^\otimes: R_{-1}(U) \cup R_{-1}(V) &\subset R_{-1}(U \cup V), \\ R_{-1|1\!\!1}^\otimes: \emptyset &\subset R_{-1}(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00RZ

6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of [Item 1](#) has a symmetric strict monoidal structure

$$(R_{-1}, R_{-1}^\otimes, R_{-1|1\!\!1}^\otimes): (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with equalities

$$\begin{aligned} R_{-1|U,V}^\otimes: R_{-1}(U \cap V) &\xrightarrow{=} R_{-1}(U) \cap R_{-1}(V), \\ R_{-1|1\!\!1}^\otimes: R_{-1}(X) &\xrightarrow{=} Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00S0

7. *Interaction With Inverse Images I.* We have

$$R_{-1}(V) = X \setminus R^{-1}(Y \setminus V)$$

for each  $V \in \mathcal{P}(Y)$ .

00S1

8. *Interaction With Inverse Images II.* Let  $R: X \nrightarrow Y$  be a relation from  $X$  to  $Y$ .

00S2

(a) If  $R$  is a total relation, then we have an inclusion of sets

$$R_{-1}(V) \subset R^{-1}(V)$$

natural in  $V \in \mathcal{P}(Y)$ .

00S3

(b) If  $R$  is total and functional, then the above inclusion is in fact an equality.

00S4

(c) Conversely, if we have  $R_{-1} = R^{-1}$ , then  $R$  is total and functional.

#### PROOF 8.7.2.1.5 ► PROOF OF PROPOSITION 8.7.2.1.4

Item 1: Functoriality

Clear.

Item 2: Adjointness

This follows from ?? of ??.

Item 3: Lax Preservation of Colimits

Omitted.

Item 4: Preservation of Limits

This follows from Item 2 and ?? of ??.

Item 5: Symmetric Lax Monoidality With Respect to Unions

This follows from Item 3.

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from Item 4.

Item 7: Interaction With Inverse Images I

We claim we have an equality

$$R_{-1}(Y \setminus V) = X \setminus R^{-1}(V).$$

Indeed, we have

$$\begin{aligned} R_{-1}(Y \setminus V) &= \{a \in X \mid R(a) \subset Y \setminus V\}, \\ X \setminus R^{-1}(V) &= \{a \in X \mid R(a) \cap V = \emptyset\}. \end{aligned}$$

Taking  $V = Y \setminus V$  then implies the original statement.

Item 8: Interaction With Inverse Images II

Item 8a is clear, while Items 8b and 8c follow from Item 6 of Proposition 8.2.2.1.2.

00S5

**PROPOSITION 8.7.2.1.6 ► PROPERTIES OF THE COINVERSE IMAGE FUNCTION OPERATION**

Let  $R: X \rightarrow Y$  be a relation.

00S6

- Functionality I.* The assignment  $R \mapsto R_{-1}$  defines a function

$$(-)_{-1}: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

00S7

- Functionality II.* The assignment  $R \mapsto R_{-1}$  defines a function

$$(-)_{-1}: \text{Sets}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

00S8

- Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have

$$(\text{id}_X)_{-1} = \text{id}_{\mathcal{P}(X)}.$$

00S9

- Interaction With Composition.* For each pair of composable relations  $R: X \rightarrow Y$  and  $S: Y \rightarrow C$ , we have

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{S_{-1}} & \mathcal{P}(Y) \\ (S \diamond R)_{-1} = R_{-1} \circ S_{-1}, & \searrow & \downarrow R_{-1} \\ & & \mathcal{P}(X). \end{array}$$

**PROOF 8.7.2.1.7 ► PROOF OF PROPOSITION 8.7.2.1.6**

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

Indeed, we have

$$\begin{aligned} (\chi_X)_{-1}(U) &\stackrel{\text{def}}{=} \{a \in X \mid \chi_X(a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in X \mid \{a\} \subset U\} \\ &= U \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ . Thus  $(\chi_X)_{-1} = \text{id}_{\mathcal{P}(X)}$ .

Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned}
 (S \diamond R)_{-1}(U) &\stackrel{\text{def}}{=} \{a \in X \mid [S \diamond R](a) \subset U\} \\
 &\stackrel{\text{def}}{=} \{a \in X \mid S(R(a)) \subset U\} \\
 &\stackrel{\text{def}}{=} \{a \in X \mid S_!(R(a)) \subset U\} \\
 &= \{a \in X \mid R(a) \subset S_{-1}(U)\} \\
 &\stackrel{\text{def}}{=} R_{-1}(S_{-1}(U)) \\
 &\stackrel{\text{def}}{=} [R_{-1} \circ S_{-1}](U)
 \end{aligned}$$

for each  $U \in \mathcal{P}(C)$ , where we used [Item 2 of Proposition 8.7.2.1.4](#), which implies that the conditions

- We have  $S_!(R(a)) \subset U$ .
- We have  $R(a) \subset S_{-1}(U)$ .

are equivalent. Thus  $(S \diamond R)_{-1} = R_{-1} \circ S_{-1}$ . □

### 00SA 8.7.3 Inverse Images

Let  $X$  and  $Y$  be sets and let  $R: X \rightarrow Y$  be a relation.

00SB

#### DEFINITION 8.7.3.1.1 ► INVERSE IMAGES

The **inverse image function associated to  $R^1$**  is the function

$$R^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$$

defined by<sup>2</sup>

$$R^{-1}(V) \stackrel{\text{def}}{=} \{a \in X \mid R(a) \cap V \neq \emptyset\}$$

for each  $V \in \mathcal{P}(Y)$ .

<sup>1</sup>Further Terminology: Also called simply the **inverse image function associated to  $R$** .

<sup>2</sup>Further Terminology: The set  $R^{-1}(V)$  is called the **inverse image of  $V$  by  $R$**  or simply the **inverse image of  $V$  by  $R$** .

00SC

#### REMARK 8.7.3.1.2 ► UNWINDING DEFINITION 8.7.3.1.1

Identifying subsets of  $Y$  with relations from  $Y$  to pt via [Item 3 of Proposition 4.4.1.1.4](#), we see that the inverse image function associated to  $R$  is equivalently the function

$$R^{-1}: \underbrace{\mathcal{P}(Y)}_{\cong \text{Rel}(Y, \text{pt})} \rightarrow \underbrace{\mathcal{P}(X)}_{\cong \text{Rel}(X, \text{pt})}$$

defined by

$$R^{-1}(V) \stackrel{\text{def}}{=} V \diamond R$$

for each  $V \in \mathcal{P}(X)$ , where  $R \diamond V$  is the composition

$$X \xrightarrow{R} Y \xrightarrow{V} \text{pt.}$$

Explicitly, we have

$$\begin{aligned} R^{-1}(V) &\stackrel{\text{def}}{=} V \diamond R \\ &\stackrel{\text{def}}{=} \int^{b \in Y} V_b^{-1} \times R_{-2}^b. \end{aligned}$$

#### PROOF 8.7.3.1.3 ► PROOF OF REMARK 8.7.3.1.2

We have

$$\begin{aligned} V \diamond R &\stackrel{\text{def}}{=} \int^{b \in Y} V_b^{-1} \times R_{-2}^b \\ &= \left\{ a \in X \mid \int^{b \in Y} V_b^* \times R_a^b = \text{true} \right\} \\ &= \left\{ a \in X \mid \begin{array}{l} \text{there exists } b \in Y \text{ such that the} \\ \text{following conditions hold:} \end{array} \right. \\ &\quad \left. \begin{array}{l} 1. \text{ We have } V_b^* = \text{true} \\ 2. \text{ We have } R_a^b = \text{true} \end{array} \right\} \\ &= \left\{ a \in X \mid \begin{array}{l} \text{there exists } b \in Y \text{ such that the} \\ \text{following conditions hold:} \end{array} \right. \\ &\quad \left. \begin{array}{l} 1. \text{ We have } b \in V \\ 2. \text{ We have } b \in R(a) \end{array} \right\} \\ &= \{a \in X \mid \text{there exists } b \in V \text{ such that } b \in R(a)\} \\ &= \{a \in X \mid R(a) \cap V \neq \emptyset\} \\ &\stackrel{\text{def}}{=} R^{-1}(V) \end{aligned}$$

This finishes the proof. □

00SD

**PROPOSITION 8.7.3.1.4 ► PROPERTIES OF INVERSE IMAGE FUNCTIONS**

Let  $R: X \rightarrow Y$  be a relation.

00SE

- Functoriality.* The assignment  $V \mapsto R^{-1}(V)$  defines a functor

$$R^{-1}: (\mathcal{P}(Y), \subset) \rightarrow (\mathcal{P}(X), \subset)$$

where

- *Action on Objects.* For each  $V \in \mathcal{P}(Y)$ , we have

$$[R^{-1}](V) \stackrel{\text{def}}{=} R^{-1}(V).$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(Y)$ :

- If  $U \subset V$ , then  $R^{-1}(U) \subset R^{-1}(V)$ .

00SF

- Adjointness.* We have an adjunction

$$(R^{-1} \dashv R_*): \quad \mathcal{P}(Y) \begin{array}{c} \xrightarrow{R^{-1}} \\ \perp \\ \xleftarrow{R_*} \end{array} \mathcal{P}(X),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(X)}(R^{-1}(U), V) \cong \text{Hom}_{\mathcal{P}(Y)}(U, R_*(V)),$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ , i.e. such that:

- (★) The following conditions are equivalent:

- We have  $R^{-1}(U) \subset V$ .
- We have  $U \subset R_*(V)$ .

00SG

- Preservation of Colimits.* We have an equality of sets

$$R^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} R^{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} R^{-1}(U) \cup R^{-1}(V) &= R^{-1}(U \cup V), \\ R^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00SH

4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$R^{-1}\left(\bigcap_{i \in I} U_i\right) \subset \bigcap_{i \in I} R^{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} R^{-1}(U \cap V) &\subset R^{-1}(U) \cap R^{-1}(V), \\ R^{-1}(X) &\subset Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00SJ

5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of Item 1 has a symmetric strict monoidal structure

$$(R^{-1}, R^{-1, \otimes}, R_{\mathbb{1}}^{-1, \otimes}) : (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} R_{U,V}^{-1, \otimes} : R^{-1}(U) \cup R^{-1}(V) &\xrightarrow{=} R^{-1}(U \cup V), \\ R_{\mathbb{1}}^{-1, \otimes} : \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00SK

6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of Item 1 has a symmetric oplax monoidal structure

$$(R^{-1}, R^{-1, \otimes}, R_{\mathbb{1}}^{-1, \otimes}) : (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with inclusions

$$\begin{aligned} R_{U,V}^{-1, \otimes} : R^{-1}(U \cap V) &\subset R^{-1}(U) \cap R^{-1}(V), \\ R_{\mathbb{1}}^{-1, \otimes} : R^{-1}(X) &\subset Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(Y)$ .

00SL

7. *Interaction With Coinverse Images I.* We have

$$R^{-1}(V) = X \setminus R_{-1}(Y \setminus V)$$

for each  $V \in \mathcal{P}(Y)$ .

00SM

8. *Interaction With Coinverse Images II.* Let  $R : X \dashrightarrow Y$  be a relation from  $X$  to  $Y$ .

00SN

(a) If  $R$  is a total relation, then we have an inclusion of sets

$$R_{-1}(V) \subset R^{-1}(V)$$

natural in  $V \in \mathcal{P}(Y)$ .

00SP

(b) If  $R$  is total and functional, then the above inclusion is in fact an equality.

00SQ

(c) Conversely, if we have  $R_{-1} = R^{-1}$ , then  $R$  is total and functional.

#### PROOF 8.7.3.1.5 ► PROOF OF PROPOSITION 8.7.3.1.4

Item 1: Functoriality

Clear.

Item 2: Adjointness

This follows from ?? of ??.

Item 3: Preservation of Colimits

This follows from Item 2 and ?? of ??.

Item 4: Oplax Preservation of Limits

Omitted.

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from Item 3.

Item 6: Symmetric Oplax Monoidality With Respect to Intersections

This follows from Item 4.

Item 7: Interaction With Coinverse Images I

This follows from Item 7 of Proposition 8.7.2.1.4.

Item 8: Interaction With Coinverse Images II

This was proved in Item 8 of Proposition 8.7.2.1.4. 

00SR

#### PROPOSITION 8.7.3.1.6 ► PROPERTIES OF THE INVERSE IMAGE FUNCTION OPERATION

Let  $R: X \rightarrow Y$  be a relation.

00SS

1. *Functionality I.* The assignment  $R \mapsto R^{-1}$  defines a function

$$(-)^{-1}: \text{Rel}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

00ST

2. *Functionality II.* The assignment  $R \mapsto R^{-1}$  defines a function

$$(-)^{-1}: \text{Rel}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

00SU

3. *Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have<sup>1</sup>

$$(\chi_X)^{-1} = \text{id}_{\mathcal{P}(X)}.$$

00SV

4. *Interaction With Composition.* For each pair of composable relations  $R: X \rightarrow Y$  and  $S: Y \rightarrow C$ , we have<sup>2</sup>

$$(S \diamond R)^{-1} = R^{-1} \circ S^{-1}, \quad \begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{S^{-1}} & \mathcal{P}(Y) \\ & \searrow (S \diamond R)^{-1} & \downarrow R^{-1} \\ & & \mathcal{P}(X). \end{array}$$

<sup>1</sup>That is, the postcomposition

$$(\chi_X)^{-1}: \text{Rel(pt, } X) \rightarrow \text{Rel(pt, } X)$$

is equal to  $\text{id}_{\text{Rel(pt, } X)}$ .

<sup>2</sup>That is, we have

$$(S \diamond R)^{-1} = R^{-1} \circ S^{-1}, \quad \begin{array}{ccc} \text{Rel(pt, } C) & \xrightarrow{R^{-1}} & \text{Rel(pt, } Y) \\ & \searrow (S \diamond R)^{-1} & \downarrow S^{-1} \\ & & \text{Rel(pt, } X). \end{array}$$

#### PROOF 8.7.3.1.7 ► PROOF OF PROPOSITION 8.7.3.1.6

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from Item 5 of Proposition 11.1.4.1.2.

Item 4: Interaction With Composition

This follows from Item 2 of Proposition 11.1.4.1.2. 

#### 00SW 8.7.4 Codirect Images

Let  $X$  and  $Y$  be sets and let  $R: X \rightarrow Y$  be a relation.

00SX

## DEFINITION 8.7.4.1.1 ► CODIRECT IMAGES

The **codirect image function associated to  $R$**  is the function

$$R_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by<sup>1,2</sup>

$$\begin{aligned} R_*(U) &\stackrel{\text{def}}{=} \left\{ b \in Y \mid \begin{array}{l} \text{for each } a \in X, \text{ if we have} \\ b \in R(a), \text{ then } a \in U \end{array} \right\} \\ &= \{b \in Y \mid R^{-1}(b) \subset U\} \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ .

<sup>1</sup>Further Terminology: The set  $R_*(U)$  is called the **codirect image of  $U$  by  $R$** .

<sup>2</sup>We also have

$$R_*(U) = Y \setminus R_!(X \setminus U);$$

see Item 7 of Proposition 8.7.4.1.4.

00SY

## REMARK 8.7.4.1.2 ► UNWINDING DEFINITION 8.7.4.1.1

Identifying subsets of  $Y$  with relations from  $\text{pt}$  to  $Y$  via Item 3 of Proposition 4.4.1.1.4, we see that the codirect image function associated to  $R$  is equivalently the function

$$R_*: \underbrace{\mathcal{P}(X)}_{\cong \text{Rel}(X, \text{pt})} \rightarrow \underbrace{\mathcal{P}(Y)}_{\cong \text{Rel}(Y, \text{pt})}$$

defined by

$$R_*(U) \stackrel{\text{def}}{=} \text{Ran}_R(U), \quad \begin{array}{ccc} & Y & \\ & \swarrow R \quad \searrow & \\ X & \xrightarrow[U]{+} & \text{pt}, & \downarrow & \text{Ran}_R(U) \end{array}$$

being explicitly computed by

$$\begin{aligned} R_*(U) &\stackrel{\text{def}}{=} \text{Ran}_R(U) \\ &\cong \int_{a \in X} \text{Hom}_{\{\text{t,f}\}}(R_a^{-2}, U_a^{-1}), \end{aligned}$$

where we have used ??.

## PROOF 8.7.4.1.3 ► PROOF OF REMARK 8.7.4.1.2

We have

$$\begin{aligned}
\text{Ran}_R(V) &\cong \int_{a \in X} \text{Hom}_{\{\text{t,f}\}}(R_a^{-2}, U_a^{-1}) \\
&= \left\{ b \in Y \mid \int_{a \in X} \text{Hom}_{\{\text{t,f}\}}(R_a^b, U_a^\star) = \text{true} \right\} \\
&= \left\{ b \in Y \mid \begin{array}{l} \text{for each } a \in X, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\
&\quad \left. \begin{array}{l} 1. \text{ We have } R_a^b = \text{false} \\ 2. \text{ The following conditions hold:} \end{array} \right. \\
&\quad \left. \begin{array}{l} (a) \text{ We have } R_a^b = \text{true} \\ (b) \text{ We have } U_a^\star = \text{true} \end{array} \right\} \\
&= \left\{ b \in Y \mid \begin{array}{l} \text{for each } a \in X, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\
&\quad \left. \begin{array}{l} 1. \text{ We have } b \notin R(X) \\ 2. \text{ The following conditions hold:} \end{array} \right. \\
&\quad \left. \begin{array}{l} (a) \text{ We have } b \in R(a) \\ (b) \text{ We have } a \in U \end{array} \right\} \\
&= \left\{ b \in Y \mid \begin{array}{l} \text{for each } a \in X, \text{ if we have} \\ b \in R(a), \text{ then } a \in U \end{array} \right\} \\
&= \left\{ b \in Y \mid R^{-1}(b) \subset U \right\} \\
&\stackrel{\text{def}}{=} R^{-1}(U).
\end{aligned}$$

This finishes the proof. ■

00SZ

## PROPOSITION 8.7.4.1.4 ► PROPERTIES OF CODIRECT IMAGES

Let  $R: X \rightarrow Y$  be a relation.

00TO

- Functionality.* The assignment  $U \mapsto R_*(U)$  defines a functor

$$R_*: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset)$$

where

- Action on Objects.* For each  $U \in \mathcal{P}(X)$ , we have

$$[R_*](U) \stackrel{\text{def}}{=} R_*(U).$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(X)$ :

– If  $U \subset V$ , then  $R_*(U) \subset R_*(V)$ .

00T1

2. *Adjointness.* We have an adjunction

$$(R^{-1} \dashv R_*): \quad \mathcal{P}(Y) \begin{array}{c} \xrightarrow{R^{-1}} \\ \perp \\ \xleftarrow{R_*} \end{array} \mathcal{P}(X),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(X)}(R^{-1}(U), V) \cong \text{Hom}_{\mathcal{P}(X)}(U, R_*(V)),$$

natural in  $U \in \mathcal{P}(X)$  and  $V \in \mathcal{P}(Y)$ , i.e. such that:

- (★) The following conditions are equivalent:

- We have  $R^{-1}(U) \subset V$ .
- We have  $U \subset R_*(V)$ .

00T2

3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} R_*(U_i) \subset R_*(\bigcup_{i \in I} U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} R_*(U) \cup R_*(V) &\subset R_*(U \cup V), \\ \emptyset &\subset R_*(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00T3

4. *Preservation of Limits.* We have an equality of sets

$$R_*(\bigcap_{i \in I} U_i) = \bigcap_{i \in I} R_*(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} R_*(U \cap V) &= R_*(U) \cap R_*(V), \\ R_*(X) &= Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00T4

5. *Symmetric Lax Monoidality With Respect to Unions.* The codirect image function of Item 1 has a symmetric lax monoidal structure

$$(R_*, R_*^\otimes, R_{!|\perp}^\otimes): (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} R_{!|U,V}^\otimes : R_*(U) \cup R_*(V) &\subset R_*(U \cup V), \\ R_{!|\mathbb{1}}^\otimes : \emptyset &\subset R_*(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00T5

6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of [Item 1](#) has a symmetric strict monoidal structure

$$(R_*, R_*^\otimes, R_{!|\mathbb{1}}^\otimes) : (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with equalities

$$\begin{aligned} R_{!|U,V}^\otimes : R_*(U \cap V) &\xrightarrow{=} R_*(U) \cap R_*(V), \\ R_{!|\mathbb{1}}^\otimes : R_*(X) &\xrightarrow{=} Y, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(X)$ .

00T6

7. *Relation to Direct Images.* We have

$$R_*(U) = Y \setminus R_!(X \setminus U)$$

for each  $U \in \mathcal{P}(X)$ .

#### PROOF 8.7.4.1.5 ► PROOF OF PROPOSITION 8.7.4.1.4

**Item 1: Functoriality**

Clear.

**Item 2: Adjointness**

This follows from ?? of ??.

**Item 3: Lax Preservation of Colimits**

Omitted.

**Item 4: Preservation of Limits**

This follows from [Item 2](#) and ?? of ??.

**Item 5: Symmetric Lax Monoidality With Respect to Unions**

This follows from [Item 3](#).

**Item 6: Symmetric Strict Monoidality With Respect to Intersections**

This follows from **Item 4**.

**Item 7: Relation to Direct Images**

This follows from **Item 7** of [Proposition 8.7.1.1.4](#). Alternatively, we may prove it directly as follows, with the proof proceeding in the same way as in the case of functions ([Item 16](#) of [Proposition 4.6.3.1.7](#)).

We claim that  $R_*(U) = Y \setminus R_!(X \setminus U)$ :

- *The First Implication.* We claim that

$$R_*(U) \subset Y \setminus R_!(X \setminus U).$$

Let  $b \in R_*(U)$ . We need to show that  $b \notin R_!(X \setminus U)$ , i.e. that there is no  $a \in X \setminus U$  such that  $b \in R(a)$ .

This is indeed the case, as otherwise we would have  $a \in R^{-1}(b)$  and  $a \notin U$ , contradicting  $R^{-1}(b) \subset U$  (which holds since  $b \in R_*(U)$ ).

Thus  $b \in Y \setminus R_!(X \setminus U)$ .

- *The Second Implication.* We claim that

$$Y \setminus R_!(X \setminus U) \subset R_*(U).$$

Let  $b \in Y \setminus R_!(X \setminus U)$ . We need to show that  $b \in R_*(U)$ , i.e. that  $R^{-1}(b) \subset U$ .

Since  $b \notin R_!(X \setminus U)$ , there exists no  $a \in X \setminus U$  such that  $b \in R(a)$ , and hence  $R^{-1}(b) \subset U$ .

Thus  $b \in R_*(U)$ .

This finishes the proof. 

00T7

**PROPOSITION 8.7.4.1.6 ► PROPERTIES OF THE CODIRECT IMAGE FUNCTION OPERATION**

Let  $R: X \rightarrow Y$  be a relation.

00T8

1. *Functionality I.* The assignment  $R \mapsto R_*$  defines a function

$$(-)_*: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

00T9

2. *Functionality II.* The assignment  $R \mapsto R_*$  defines a function

$$(-)_*: \text{Sets}(X, Y) \rightarrow \text{Hom}_{\text{Pos}}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

00TA

3. *Interaction With Identities.* For each  $X \in \text{Obj}(\text{Sets})$ , we have

$$(\text{id}_X)_* = \text{id}_{\mathcal{P}(X)}.$$

00TB

4. *Interaction With Composition.* For each pair of composable relations  $R: X \rightarrow Y$  and  $S: Y \rightarrow C$ , we have

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{R_*} & \mathcal{P}(Y) \\ (S \diamond R)_* = S_* \circ R_*, & \searrow & \downarrow S_* \\ (S \diamond R)_* & & \mathcal{P}(C). \end{array}$$

#### PROOF 8.7.4.1.7 ▶ PROOF OF PROPOSITION 8.7.4.1.6

##### Item 1: Functionality I

Clear.

##### Item 2: Functionality II

Clear.

##### Item 3: Interaction With Identities

Indeed, we have

$$\begin{aligned} (\chi_X)_*(U) &\stackrel{\text{def}}{=} \{a \in X \mid \chi_X^{-1}(a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in X \mid \{a\} \subset U\} \\ &= U \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ . Thus  $(\chi_X)_* = \text{id}_{\mathcal{P}(X)}$ .

##### Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned} (S \diamond R)_*(U) &\stackrel{\text{def}}{=} \{c \in C \mid [S \diamond R]^{-1}(c) \subset U\} \\ &\stackrel{\text{def}}{=} \{c \in C \mid S^{-1}(R^{-1}(c)) \subset U\} \\ &= \{c \in C \mid R^{-1}(c) \subset S_*(U)\} \\ &\stackrel{\text{def}}{=} R_*(S_*(U)) \\ &\stackrel{\text{def}}{=} [R_* \circ S_*](U) \end{aligned}$$

for each  $U \in \mathcal{P}(C)$ , where we used [Item 2 of Proposition 8.7.4.1.4](#), which implies that the conditions

- We have  $S^{-1}(R^{-1}(c)) \subset U$ .
- We have  $R^{-1}(c) \subset S_*(U)$ .

are equivalent. Thus  $(S \diamond R)_* = S_* \circ R_*$ . □

**01JA 8.7.5 Functoriality of Powersets**

00TD

**PROPOSITION 8.7.5.1.1 ► FUNCTORIALITY OF POWERSETS I**

The assignment  $X \mapsto \mathcal{P}(X)$  defines functors<sup>1</sup>

$$\begin{aligned}\mathcal{P}_! &: \text{Rel} \rightarrow \text{Sets}, \\ \mathcal{P}_{-1} &: \text{Rel}^{\text{op}} \rightarrow \text{Sets}, \\ \mathcal{P}^{-1} &: \text{Rel}^{\text{op}} \rightarrow \text{Sets}, \\ \mathcal{P}_* &: \text{Rel} \rightarrow \text{Sets}\end{aligned}$$

where

- *Action on Objects.* For each  $X \in \text{Obj}(\text{Rel})$ , we have

$$\begin{aligned}\mathcal{P}_!(X) &\stackrel{\text{def}}{=} \mathcal{P}(X), \\ \mathcal{P}_{-1}(X) &\stackrel{\text{def}}{=} \mathcal{P}(X), \\ \mathcal{P}^{-1}(X) &\stackrel{\text{def}}{=} \mathcal{P}(X), \\ \mathcal{P}_*(X) &\stackrel{\text{def}}{=} \mathcal{P}(X).\end{aligned}$$

- *Action on Morphisms.* For each morphism  $R: X \rightarrow Y$  of Rel, the images

$$\begin{aligned}\mathcal{P}_!(R) &: \mathcal{P}(X) \rightarrow \mathcal{P}(Y), \\ \mathcal{P}_{-1}(R) &: \mathcal{P}(Y) \rightarrow \mathcal{P}(X), \\ \mathcal{P}^{-1}(R) &: \mathcal{P}(Y) \rightarrow \mathcal{P}(X), \\ \mathcal{P}_*(R) &: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)\end{aligned}$$

of  $R$  by  $\mathcal{P}_!$ ,  $\mathcal{P}_{-1}$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_*$  are defined by

$$\begin{aligned}\mathcal{P}_!(R) &\stackrel{\text{def}}{=} R_!, \\ \mathcal{P}_{-1}(R) &\stackrel{\text{def}}{=} R_{-1}, \\ \mathcal{P}^{-1}(R) &\stackrel{\text{def}}{=} R^{-1}, \\ \mathcal{P}_*(R) &\stackrel{\text{def}}{=} R_*,\end{aligned}$$

as in Definitions 8.7.1.1.1, 8.7.2.1.1, 8.7.3.1.1 and 8.7.4.1.1.

---

<sup>1</sup>The functor  $\mathcal{P}_!: \text{Rel} \rightarrow \text{Sets}$  admits a left adjoint; see Item 2 of Proposition 8.2.2.1.2.

**PROOF 8.7.5.1.2 ▶ PROOF OF PROPOSITION 8.7.5.1.1**

This follows from **Items 3 and 4 of Proposition 8.7.1.1.6**, **Items 3 and 4 of Proposition 8.7.2.1.6**, **Items 3 and 4 of Proposition 8.7.3.1.6**, and **Items 3 and 4 of Proposition 8.7.4.1.6**. 

Let  $X$  and  $Y$  be sets and let  $R: X \nrightarrow Y$  be a relation.

00TF

**DEFINITION 8.7.5.1.3 ▶ THE RELATION ON POWERSETS ASSOCIATED TO A RELATION**

The **relation on powersets associated to  $R$**  is the relation


$$\mathcal{P}(R): \mathcal{P}(X) \nrightarrow \mathcal{P}(Y)$$

defined by<sup>1</sup>

$$\mathcal{P}(R)_U^V \stackrel{\text{def}}{=} \mathbf{Rel}(\chi_{\text{pt}}, V \diamond R \diamond U)$$

for each  $U \in \mathcal{P}(X)$  and each  $V \in \mathcal{P}(Y)$ .

<sup>1</sup>Illustration:



00TG

**REMARK 8.7.5.1.4 ▶ UNWINDING DEFINITION 8.7.5.1.3**

In detail, we have  $U \sim_{\mathcal{P}(R)} V$  iff the following equivalent conditions hold:

- We have  $\chi_{\text{pt}} \subset V \diamond R \diamond U$ .
- We have  $(V \diamond R \diamond U)^* = \text{true}$ , i.e. we have

$$\int^{a \in X} \int^{b \in Y} V_b^* \times R_a^b \times U_a^* = \text{true}.$$

- There exists some  $a \in X$  and some  $b \in Y$  such that:
  - We have  $U_a^* = \text{true}$ .
  - We have  $R_a^b = \text{true}$ .
  - We have  $V_b^* = \text{true}$ .
- There exists some  $a \in X$  and some  $b \in Y$  such that:
  - We have  $a \in U$ .
  - We have  $a \sim_R b$ .
  - We have  $b \in V$ .

00TH

**PROPOSITION 8.7.5.1.5 ▶ FUNCTORIALITY OF POWERSETS II**

The assignment  $R \mapsto \mathcal{P}(R)$  defines a functor

$$\mathcal{P} : \mathbf{Rel} \rightarrow \mathbf{Rel}.$$

**PROOF 8.7.5.1.6 ▶ PROOF OF PROPOSITION 8.7.5.1.5**

Omitted. 

00MN

**8.8 The Left Skew Monoidal Structure on  $\mathbf{Rel}(A, B)$** **00MP 8.8.1 The Left Skew Monoidal Product**

Let  $A$  and  $B$  be sets and let  $J : A \nrightarrow B$  be a relation.

00MQ

**DEFINITION 8.8.1.1.1 ▶ THE LEFT  $J$ -SKEW MONOIDAL PRODUCT OF  $\mathbf{Rel}(A, B)$** 

The **left  $J$ -skew monoidal product of  $\mathbf{Rel}(A, B)$**  is the functor

$$\triangleleft_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

where

- *Action on Objects.* For each  $R, S \in \mathbf{Obj}(\mathbf{Rel}(A, B))$ , we have

$$S \triangleleft_J R \stackrel{\text{def}}{=} S \diamond \text{Rift}_J(R), \quad \begin{array}{ccc} A & \xrightarrow{S} & B \\ \text{Rift}_J(R) \swarrow & \nearrow & \downarrow J \\ A & \xrightarrow{R} & B \end{array}$$

- *Action on Morphisms.* For each  $R, S, R', S' \in \mathbf{Obj}(\mathbf{Rel}(A, B))$ , the action on Hom-sets

$$(\triangleleft_J)_{(G,F),(G',F')} : \mathbf{Hom}_{\mathbf{Rel}(A,B)}(S, S') \times \mathbf{Hom}_{\mathbf{Rel}(A,B)}(R, R') \rightarrow \mathbf{Hom}_{\mathbf{Rel}(A,B)}(S \triangleleft_J R, S' \triangleleft_J R')$$

of  $\triangleleft_J$  at  $((R, S), (R', S'))$  is defined by<sup>1</sup>

$$\beta \triangleleft_J \alpha \stackrel{\text{def}}{=} \beta \diamond \text{Rift}_J(\alpha),$$

for each  $\beta \in \text{Hom}_{\mathbf{Rel}(A, B)}(S, S')$  and each  $\alpha \in \text{Hom}_{\mathbf{Rel}(A, B)}(R, R')$ .

<sup>1</sup>Since  $\mathbf{Rel}(A, B)$  is posetal, this is to say that if  $S \subset S'$  and  $R \subset R'$ , then  $S \triangleleft_J R \subset S' \triangleleft_J R'$ .

### 00MR 8.8.2 The Left Skew Monoidal Unit

Let  $A$  and  $B$  be sets and let  $J: A \rightarrow B$  be a relation.

00MS

#### DEFINITION 8.8.2.1.1 ► THE LEFT $J$ -SKEW MONOIDAL UNIT OF $\mathbf{Rel}(A, B)$

The **left  $J$ -skew monoidal unit of  $\mathbf{Rel}(A, B)$**  is the functor

$$\mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A, B)}: \text{pt} \rightarrow \mathbf{Rel}(A, B)$$

picking the object

$$\mathbb{1}_{\mathbf{Rel}(A, B)}^{\triangleleft_J} \stackrel{\text{def}}{=} J$$

of  $\mathbf{Rel}(A, B)$ .

### 00MT 8.8.3 The Left Skew Associators

Let  $A$  and  $B$  be sets and let  $J: A \rightarrow B$  be a relation.

00MU

#### DEFINITION 8.8.3.1.1 ► THE LEFT $J$ -SKEW ASSOCIATOR OF $\mathbf{Rel}(A, B)$

The **left  $J$ -skew associator of  $\mathbf{Rel}(A, B)$**  is the natural transformation

$$\alpha^{\mathbf{Rel}(A, B), \triangleleft_J}: \triangleleft_J \circ (\triangleleft_J \times \text{id}) \Longrightarrow \triangleleft_J \circ (\text{id} \times \triangleleft_J) \circ \alpha_{\mathbf{Rel}(A, B), \mathbf{Rel}(A, B), \mathbf{Rel}(A, B)}^{\text{Cats}},$$

as in the diagram

$$\begin{array}{ccc}
 & \text{Rel}(A, B) \times (\text{Rel}(A, B) \times \text{Rel}(A, B)) & \\
 & \alpha_{\text{Rel}(A, B), \text{Rel}(A, B), \text{Rel}(A, B)}^{\text{Cats}} \nearrow \text{id} & \searrow \text{id} \times \triangleleft_J \\
 (\text{Rel}(A, B) \times \text{Rel}(A, B)) \times \text{Rel}(A, B) & \text{Rel}(A, B) \times \text{Rel}(A, B) & \\
 \downarrow \triangleleft_J \times \text{id} & \alpha^{\text{Rel}(A, B), \triangleleft_J} \swarrow & \downarrow \triangleleft_J \\
 \text{Rel}(A, B) \times \text{Rel}(A, B) & \xrightarrow{\triangleleft_J} & \text{Rel}(A, B),
 \end{array}$$

whose component

$$\alpha_{T, S, R}^{\text{Rel}(A, B), \triangleleft_J} : \underbrace{(T \triangleleft_J S) \triangleleft_J R}_{\stackrel{\text{def}}{=} T \diamond \text{Rift}_J(S) \diamond \text{Rift}_J(R)} \hookrightarrow \underbrace{T \triangleleft_J (S \triangleleft_J R)}_{\stackrel{\text{def}}{=} T \diamond \text{Rift}_J(S \diamond \text{Rift}_J(R))}$$

at  $(T, S, R)$  is given by

$$\alpha_{T, S, R}^{\text{Rel}(A, B), \triangleleft_J} \stackrel{\text{def}}{=} \text{id}_T \diamond \gamma,$$

where

$$\gamma : \text{Rift}_J(S) \diamond \text{Rift}_J(R) \hookrightarrow \text{Rift}_J(S \diamond \text{Rift}_J(R))$$

is the inclusion adjunct to the inclusion

$$\epsilon_S \star \text{id}_{\text{Rift}_J(R)} : \underbrace{J \diamond \text{Rift}_J(S) \diamond \text{Rift}_J(R)}_{\stackrel{\text{def}}{=} J_!(\text{Rift}_J(S) \diamond \text{Rift}_J(R))} \hookrightarrow S \diamond \text{Rift}_J(R)$$

under the adjunction  $J_! \dashv \text{Rift}_J$ , where  $\epsilon : J \diamond \text{Rift}_J \Rightarrow \text{id}_{\text{Rel}(A, B)}$  is the counit of the adjunction  $J_! \dashv \text{Rift}_J$ .

#### 00MV 8.8.4 The Left Skew Left Unitors

Let  $A$  and  $B$  be sets and let  $J : A \rightarrow B$  be a relation.

00MW

##### DEFINITION 8.8.4.1.1 ► THE LEFT $J$ -SKEW LEFT UNITOR OF $\text{Rel}(A, B)$

The **left  $J$ -skew left unitor of  $\text{Rel}(A, B)$**  is the natural transformation

$$\lambda^{\text{Rel}(A, B), \triangleleft_J} : \triangleleft_J \circ (\mathbb{1}_{\triangleleft_J}^{\text{Rel}(A, B)} \times \text{id}) \Rightarrow \lambda_{\text{Rel}(A, B)}^{\text{Cats}_2}$$

as in the diagram

$$\begin{array}{ccc}
 \mathbf{pt} \times \mathbf{Rel}(A, B) & \xrightarrow{\mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A, B)} \times \text{id}} & \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \\
 & \searrow \lambda_{\mathbf{Rel}(A, B), \triangleleft_J}^{\mathbf{Cats}_2} & \downarrow \triangleleft_J \\
 & \swarrow \lambda_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} & \\
 & \mathbf{Rel}(A, B), &
 \end{array}$$

whose component

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleleft_J} : \underbrace{J \triangleleft_J R}_{\stackrel{\text{def}}{=} J \diamond \text{Rift}_J(R)} \hookrightarrow R$$

at  $R$  is given by

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleleft_J} \stackrel{\text{def}}{=} \epsilon_R,$$

where  $\epsilon : J_! \diamond \text{Rift}_J \Rightarrow \text{id}_{\mathbf{Rel}(A, B)}$  is the counit of the adjunction  $J_! \dashv \text{Rift}_J$ .

### 00MX 8.8.5 The Left Skew Right Unitors

Let  $A$  and  $B$  be sets and let  $J : A \nrightarrow B$  be a relation.

00MY

#### DEFINITION 8.8.5.1.1 ► THE LEFT $J$ -SKEW RIGHT UNITOR OF $\mathbf{Rel}(A, B)$

The **left  $J$ -skew right unitor** of  $\mathbf{Rel}(A, B)$  is the natural transformation

$$\rho^{\mathbf{Rel}(A, B), \triangleleft_J} : \rho_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} \Rightarrow \triangleleft_J \circ (\text{id} \times \mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A, B)})$$

as in the diagram

$$\begin{array}{ccc}
 \mathbf{Rel}(A, B) \times \mathbf{pt} & \xrightarrow{\text{id} \times \mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A, B)}} & \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B), \\
 & \searrow \rho_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} & \downarrow \triangleleft_J \\
 & \swarrow \rho^{\mathbf{Rel}(A, B), \triangleleft_J} & \\
 & \mathbf{Rel}(A, B) &
 \end{array}$$

whose component

$$\rho_R^{\mathbf{Rel}(A,B), \triangleleft_J} : R \hookrightarrow \underbrace{R \triangleleft_J J}_{\stackrel{\text{def}}{=} R \diamond \mathbf{Rift}_J(J)}$$

at  $R$  is given by the composition

$$\begin{aligned} R &\xrightarrow{\sim} R \diamond \chi_A \\ &\xrightarrow{\text{id}_R \diamond \eta_{\chi_A}} R \diamond \mathbf{Rift}_J(J_!(\chi_A)) \\ &\stackrel{\text{def}}{=} R \diamond \mathbf{Rift}_J(J \diamond \chi_A) \\ &\xrightarrow{\sim} R \diamond \mathbf{Rift}_J(J) \\ &\stackrel{\text{def}}{=} R \triangleleft_J J, \end{aligned}$$

where  $\eta : \text{id}_{\mathbf{Rel}(A,A)} \Rightarrow \mathbf{Rift}_J \circ J_!$  is the unit of the adjunction  $J_! \dashv \mathbf{Rift}_J$ .

### 00MZ 8.8.6 The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00N0

#### PROPOSITION 8.8.6.1.1 ► THE LEFT $J$ -SKEW MONOIDAL STRUCTURE ON $\mathbf{Rel}(A, B)$

The category  $\mathbf{Rel}(A, B)$  admits a left skew monoidal category structure consisting of

- *The Underlying Category.* The posetal category associated to the poset  $\mathbf{Rel}(A, B)$  of relations from  $A$  to  $B$  of ?? of ??.
- *The Left Skew Monoidal Product.* The left  $J$ -skew monoidal product

$$\triangleleft_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of [Definition 8.8.1.1.1](#).

- *The Left Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Rel}(A,B), \triangleleft_J} : \mathbf{pt} \rightarrow \mathbf{Rel}(A, B)$$

of [Definition 8.8.2.1.1](#).

- *The Left Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Rel}(A,B), \triangleleft_J} : \triangleleft_J \circ (\triangleleft_J \times \text{id}) \Rightarrow \triangleleft_J \circ (\text{id} \times \triangleleft_J) \circ \alpha_{\mathbf{Rel}(A,B), \mathbf{Rel}(A,B), \mathbf{Rel}(A,B)}^{\mathbf{Cats}}$$

of [Definition 8.8.3.1.1](#).

- *The Left Skew Left Unitors.* The natural transformation

$$\lambda^{\mathbf{Rel}(A,B), \triangleleft_J} : \triangleleft_J \circ (\mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A,B)} \times \text{id}) \Rightarrow \lambda_{\mathbf{Rel}(A,B)}^{\mathbf{Cats}_2}$$

of [Definition 8.8.4.1.1](#).

- *The Left Skew Right Unitors.* The natural transformation

$$\rho^{\mathbf{Rel}(A, B), \triangleleft_J} : \rho_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} \implies \triangleleft_J \circ (\text{id} \times \mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A, B)})$$

of Definition 8.8.5.1.1.

#### PROOF 8.8.6.1.2 ▶ PROOF OF PROPOSITION 8.8.6.1.1

Since  $\mathbf{Rel}(A, B)$  is posetal, the commutativity of the pentagon identity, the left skew left triangle identity, the left skew right triangle identity, the left skew middle triangle identity, and the zigzag identity is automatic (?? of ??), and thus  $\mathbf{Rel}(A, B)$  together with the data in the statement forms a left skew monoidal category. 

## 00N1 8.9 The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

Let  $A$  and  $B$  be sets and let  $J : A \rightarrow B$  be a relation.

### 00N2 8.9.1 The Right Skew Monoidal Product

#### 00N3 DEFINITION 8.9.1.1.1 ▶ THE RIGHT $J$ -SKEW MONOIDAL PRODUCT OF $\mathbf{Rel}(A, B)$

The **right  $J$ -skew monoidal product** of  $\mathbf{Rel}(A, B)$  is the functor

$$\triangleright_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

where


- *Action on Objects.* For each  $R, S \in \text{Obj}(\mathbf{Rel}(A, B))$ , we have

$$A \xrightarrow{R} B \dashrightarrow \text{Ran}_J(S) \quad \begin{array}{c} \text{Ran}_J(S) \\ \dashrightarrow \\ \uparrow J \\ \text{Ran}_J(S) \diamond R \\ \uparrow J \\ A \end{array}$$

- *Action on Morphisms.* For each  $R, S, R', S' \in \text{Obj}(\mathbf{Rel}(A, B))$ , the action on Hom-sets

$$(\triangleright_J)_{(S, R), (S', R')} : \text{Hom}_{\mathbf{Rel}(A, B)}(S, S') \times \text{Hom}_{\mathbf{Rel}(A, B)}(R, R') \rightarrow \text{Hom}_{\mathbf{Rel}(A, B)}(S \triangleright_J R, S' \triangleright_J R')$$

of  $\triangleright_J$  at  $((S, R), (S', R'))$  is defined by<sup>1</sup>



for each  $\beta \in \text{Hom}_{\mathbf{Rel}(A, B)}(S, S')$  and each  $\alpha \in \text{Hom}_{\mathbf{Rel}(A, B)}(R, R')$ .

<sup>1</sup>Since  $\mathbf{Rel}(A, B)$  is posetal, this is to say that if  $S \subset S'$  and  $R \subset R'$ , then  $S \triangleright_J R \subset S' \triangleright_J R'$ .

#### 00N4 8.9.2 The Right Skew Monoidal Unit

00N5

##### DEFINITION 8.9.2.1.1 ► THE RIGHT J-SKEW MONOIDAL UNIT OF $\mathbf{Rel}(A, B)$

The **right  $J$ -skew monoidal unit of  $\mathbf{Rel}(A, B)$**  is the functor

$$\mathbb{1}_{\triangleright_J}^{\mathbf{Rel}(A, B)} : \text{pt} \rightarrow \mathbf{Rel}(A, B)$$

picking the object

$$\mathbb{1}_{\mathbf{Rel}(A, B)}^{\triangleright_J} \stackrel{\text{def}}{=} J$$

of  $\mathbf{Rel}(A, B)$ .

#### 00N6 8.9.3 The Right Skew Associators

00N7

##### DEFINITION 8.9.3.1.1 ► THE RIGHT J-SKEW ASSOCIATOR OF $\mathbf{Rel}(A, B)$

The **right  $J$ -skew associator of  $\mathbf{Rel}(A, B)$**  is the natural transformation

$$\alpha^{\mathbf{Rel}(A, B), \triangleright_J} : \triangleright_J \circ (\text{id} \times \triangleright_J) \Longrightarrow \triangleright_J \circ (\triangleright_J \times \text{id}) \circ \alpha_{\mathbf{Rel}(A, B), \mathbf{Rel}(A, B), \mathbf{Rel}(A, B)}^{\text{Cats}, -1},$$

as in the diagram

$$\begin{array}{ccc}
 & (\mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B)) \times \mathbf{Rel}(A, B) & \\
 & \alpha_{\mathbf{Rel}(A, B), \mathbf{Rel}(A, B), \mathbf{Rel}(A, B)}^{\mathbf{Cats}, -1} \nearrow \text{dotted} & \searrow \triangleright_J \times \text{id} \\
 \mathbf{Rel}(A, B) \times (\mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B)) & \xrightarrow{\quad} & \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \\
 & \downarrow \text{id} \times \triangleright_J & \swarrow \alpha_{\mathbf{Rel}(A, B), \triangleright_J}^{\mathbf{Rel}(A, B), \triangleright_J} \\
 & \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) & \xrightarrow[\triangleright_J]{} \mathbf{Rel}(A, B),
 \end{array}$$

whose component

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleright_J} : \underbrace{T \triangleright_J (S \triangleright_J R)}_{\stackrel{\text{def}}{=} \text{Ran}_J(T) \diamond \text{Ran}_J(S) \diamond R} \hookrightarrow \underbrace{(T \triangleright_J S) \triangleright_J R}_{\stackrel{\text{def}}{=} \text{Ran}_J(\text{Ran}_J(T) \diamond S) \diamond R}$$

at  $(T, S, R)$  is given by

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleright} \stackrel{\text{def}}{=} \gamma \diamond \text{id}_R,$$

where

$$\gamma : \text{Ran}_J(T) \diamond \text{Ran}_J(S) \hookrightarrow \text{Ran}_J(\text{Ran}_J(T) \diamond S)$$

is the inclusion adjunct to the inclusion

$$\text{id}_{\text{Ran}_J(T) \diamond S} : \underbrace{\text{Ran}_J(T) \diamond \text{Ran}_J(S) \diamond J}_{\stackrel{\text{def}}{=} J^*(\text{Ran}_J(T) \diamond \text{Ran}_J(S))} \hookrightarrow \text{Ran}_J(T) \diamond S$$

under the adjunction  $J^* \dashv \text{Ran}_J$ , where  $\epsilon : \text{Ran}_J \diamond J \Rightarrow \text{id}_{\mathbf{Rel}(A, B)}$  is the counit of the adjunction  $J^* \dashv \text{Ran}_J$ .

#### 00N8 8.9.4 The Right Skew Left Unitors

##### 00N9 DEFINITION 8.9.4.1.1 ► THE RIGHT $J$ -SKEW LEFT UNITOR OF $\mathbf{Rel}(A, B)$

The **right  $J$ -skew left unitor** of  $\mathbf{Rel}(A, B)$  is the natural transformation

$$\lambda^{\mathbf{Rel}(A, B), \triangleright_J} : \lambda_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} \Rightarrow \triangleright_J \circ (\mathbb{1}_{\triangleright}^{\mathbf{Rel}(A, B)} \times \text{id}),$$

as in the diagram

$$\begin{array}{ccc}
 \mathbf{pt} \times \mathbf{Rel}(A, B) & \xrightarrow{\mathbb{1}_{\triangleright_J}^{\mathbf{Rel}(A, B)} \times \text{id}} & \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \\
 & \searrow \lambda_{\mathbf{Rel}(A, B), \triangleright_J}^{\mathbf{Cats}_2} \quad \swarrow \lambda_{\mathbf{Rel}(A, B), \triangleright_J}^{\mathbf{Rel}(A, B), \triangleright_J} & \downarrow \triangleright_J \\
 & \dashrightarrow \lambda_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} & \mathbf{Rel}(A, B),
 \end{array}$$

whose component

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleright_J} : R \hookrightarrow \underbrace{J \triangleright_J R}_{\stackrel{\text{def}}{=} \text{Ran}_J(J) \diamond R}$$

at  $R$  is given by the composition

$$\begin{aligned}
 R &\xrightarrow{\sim} \chi_B \diamond R \\
 &\xrightarrow{\eta_{\chi_B}} \diamond \text{id}_{\text{Ran}_J(J^*(\chi_A))} \diamond R \\
 &\stackrel{\text{def}}{=} \text{Ran}_J(J^* \diamond \chi_A) \diamond R \\
 &\xrightarrow{\sim} \text{Ran}_J(J) \diamond R \\
 &\stackrel{\text{def}}{=} R \triangleright_J J,
 \end{aligned}$$

where  $\eta : \text{id}_{\mathbf{Rel}(B, B)} \Rightarrow \text{Ran}_J \circ J^*$  is the unit of the adjunction  $J^* \dashv \text{Ran}_J$ .

### 00NA 8.9.5 The Right Skew Right Unitors

00NB

**DEFINITION 8.9.5.1.1 ► THE RIGHT  $J$ -SKEW RIGHT UNITOR OF  $\mathbf{Rel}(A, B)$**

The **right  $J$ -skew right unitor of  $\mathbf{Rel}(A, B)$**  is the natural transformation

$$\rho^{\mathbf{Rel}(A, B), \triangleright_J} : \triangleright_J \circ (\text{id} \times \mathbb{1}_{\triangleright}^{\mathbf{Rel}(A, B)}) \Rightarrow \rho_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2},$$

as in the diagram

$$\begin{array}{ccc}
 \mathbf{Rel}(A, B) \times \mathbf{pt} & \xrightarrow{\text{id} \times \mathbb{1}_{\triangleright_J}^{\mathbf{Rel}(A, B)}} & \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B), \\
 & \searrow \rho_{\mathbf{Rel}(A, B), \triangleright_J}^{\mathbf{Rel}(A, B), \triangleright_J} \quad \swarrow & \downarrow \triangleright_J \\
 & \rho_{\mathbf{Rel}(A, B)}^{\mathbf{Cats}_2} & \mathbf{Rel}(A, B)
 \end{array}$$

whose component

$$\rho_S^{\mathbf{Rel}(A, B), \triangleright_J} : \underbrace{S \triangleright_J J}_{\stackrel{\text{def}}{=} \mathbf{Ran}_J(S) \diamond J} \hookrightarrow S$$

at  $S$  is given by

$$\rho_S^{\mathbf{Rel}(A, B), \triangleright_J} \stackrel{\text{def}}{=} \epsilon_R,$$

where  $\epsilon : J^* \circ \mathbf{Ran}_J \Rightarrow \text{id}_{\mathbf{Rel}(A, B)}$  is the counit of the adjunction  $J^* \dashv \mathbf{Ran}_J$ .

### 00NC 8.9.6 The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00ND

#### PROPOSITION 8.9.6.1.1 ► THE RIGHT $J$ -SKEW MONOIDAL STRUCTURE ON $\mathbf{Rel}(A, B)$

The category  $\mathbf{Rel}(A, B)$  admits a right skew monoidal category structure consisting of

- *The Underlying Category.* The posetal category associated to the poset  $\mathbf{Rel}(A, B)$  of relations from  $A$  to  $B$  of ?? of ??.
- *The Right Skew Monoidal Product.* The right  $J$ -skew monoidal product

$$\triangleleft_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of Definition 8.9.1.1.1.

- *The Right Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Rel}(A, B), \triangleleft_J} : \mathbf{pt} \rightarrow \mathbf{Rel}(A, B)$$

of Definition 8.9.2.1.1.

- *The Right Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Rel}(A, B), \triangleright_J} : \triangleright_J \circ (\text{id} \times \triangleright_J) \Rightarrow \triangleright_J \circ (\triangleright_J \times \text{id}) \circ \alpha_{\mathbf{Rel}(A, B), \mathbf{Rel}(A, B), \mathbf{Rel}(A, B)}^{\mathbf{Cats}, -1}$$

of Definition 8.9.3.1.1.

- *The Right Skew Left Unitors.* The natural transformation

$$\lambda^{\text{Rel}(A,B), \triangleright_J}_{\text{Rel}(A,B)} : \lambda^{\text{Cats}_2}_{\text{Rel}(A,B)} \implies \triangleright_J \circ (\mathbb{1}^{\text{Rel}(A,B)}_{\triangleright} \times \text{id})$$

of Definition 8.9.4.1.1.

- *The Right Skew Right Unitors.* The natural transformation

$$\rho^{\text{Rel}(A,B), \triangleright_J}_{\text{Rel}(A,B)} : \triangleright_J \circ (\text{id} \times \mathbb{1}^{\text{Rel}(A,B)}_{\triangleright}) \implies \rho^{\text{Cats}_2}_{\text{Rel}(A,B)}$$

of Definition 8.9.5.1.1.

#### PROOF 8.9.6.1.2 ► PROOF OF PROPOSITION 8.9.6.1.1

Since  $\text{Rel}(A, B)$  is posetal, the commutativity of the pentagon identity, the right skew left triangle identity, the right skew right triangle identity, the right skew middle triangle identity, and the zigzag identity is automatic (?? of ??), and thus  $\text{Rel}(A, B)$  together with the data in the statement forms a right skew monoidal category. 

# Appendices

## 8.A Other Chapters

### Preliminaries

1. Introduction
2. A Guide to the Literature

### Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

### Relations

8. Relations
9. Constructions With Relations

### 10. Conditions on Relations

### Categories

11. Categories
12. Presheaves and the Yoneda Lemma

### Monoidal Categories

13. Constructions With Monoidal Categories

### Bicategories

14. Types of Morphisms in Bicategories

### Extra Part

15. Notes

# Chapter 9

## Constructions With Relations

**02H3** This chapter contains some material about constructions with relations. Notably, we discuss and explore:

- 029U** 1. The existence or non-existence of Kan extensions and Kan lifts in the 2-category **Rel** ([Old Tag 15.2.1.1.8](#)).
- 029V** 2. The various kinds of constructions involving relations, such as graphs, domains, ranges, unions, intersections, products, converse relations, composition of relations, and collages ([Section 9.2](#)).

This chapter is under revision. TODO:

1. Rename range to image
2. Co/limits in **Rel**.

### Contents

---

9.1	Co/Limits in the Category of Relations.....	691
9.2	More Constructions With Relations.....	692
9.2.1	The Domain and Range of a Relation.....	692
9.2.2	Binary Unions of Relations.....	692
9.2.3	Unions of Families of Relations .....	694
9.2.4	Binary Intersections of Relations.....	695
9.2.5	Intersections of Families of Relations .....	696
9.2.6	Binary Products of Relations .....	697
9.2.7	Products of Families of Relations.....	699
9.2.8	The Collage of a Relation .....	699
9.A	Other Chapters.....	702

---

### **00NF** 9.1 Co/Limits in the Category of Relations

This section is currently just a stub, and will be properly developed later on.

## 00NZ 9.2 More Constructions With Relations

### 00PM 9.2.1 The Domain and Range of a Relation

Let  $A$  and  $B$  be sets.

00PN

#### DEFINITION 9.2.1.1 ► THE DOMAIN AND RANGE OF A RELATION

Let  $R: A \rightarrow B$  be a relation.<sup>1,2</sup>

02AV

1. The **domain of  $R$**  is the subset  $\text{dom}(R)$  of  $A$  defined by

$$\text{dom}(R) \stackrel{\text{def}}{=} \left\{ a \in A \mid \begin{array}{l} \text{there exists some } b \in B \\ \text{such that } a \sim_R b \end{array} \right\}.$$

02AW

2. The **range of  $R$**  is the subset  $\text{range}(R)$  of  $B$  defined by

$$\text{range}(R) \stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in A \\ \text{such that } a \sim_R b \end{array} \right\}.$$

---

<sup>1</sup>Following ??, we may compute the (characteristic functions associated to the) domain and range of a relation using the following colimit formulas:

$$\begin{aligned} \chi_{\text{dom}(R)}(a) &\cong \underset{b \in B}{\text{colim}}(R_a^b) \quad (a \in A) \\ &\cong \bigvee_{b \in B} R_a^b, \\ \chi_{\text{range}(R)}(b) &\cong \underset{a \in A}{\text{colim}}(R_a^b) \quad (b \in B) \\ &\cong \bigvee_{a \in A} R_a^b, \end{aligned}$$

where the join  $\vee$  is taken in the poset  $(\{\text{true}, \text{false}\}, \preceq)$  of Definition 3.2.2.1.3.

<sup>2</sup>Viewing  $R$  as a function  $R: A \rightarrow \mathcal{P}(B)$ , we have

$$\begin{aligned} \text{dom}(R) &\cong \underset{y \in Y}{\text{colim}}(R(y)) \\ &\cong \bigcup_{y \in Y} R(y), \\ \text{range}(R) &\cong \underset{x \in X}{\text{colim}}(R(x)) \\ &\cong \bigcup_{x \in X} R(x), \end{aligned}$$

### 00PP 9.2.2 Binary Unions of Relations

Let  $A$  and  $B$  be sets and let  $R$  and  $S$  be relations from  $A$  to  $B$ .

00PQ

**DEFINITION 9.2.2.1.1 ► BINARY UNIONS OF RELATIONS**

The **union of  $R$  and  $S$** <sup>1</sup> is the relation  $R \cup S$  from  $A$  to  $B$  defined as follows:

- Viewing relations from  $A$  to  $B$  as subsets of  $A \times B$ , we define<sup>2</sup>

$$R \cup S \stackrel{\text{def}}{=} \{(a, b) \in B \times A \mid \text{we have } a \sim_R b \text{ or } a \sim_S b\}.$$

- Viewing relations from  $A$  to  $B$  as functions  $A \rightarrow \mathcal{P}(B)$ , we define

$$[R \cup S](a) \stackrel{\text{def}}{=} R(a) \cup S(a)$$

for each  $a \in A$ .

<sup>1</sup>Further Terminology: Also called the **binary union of  $R$  and  $S$** , for emphasis.

<sup>2</sup>This is the same as the union of  $R$  and  $S$  as subsets of  $A \times B$ .

00PR

**PROPOSITION 9.2.2.1.2 ► PROPERTIES OF BINARY UNIONS OF RELATIONS**

Let  $R$ ,  $S$ ,  $R_1$ , and  $R_2$  be relations from  $A$  to  $B$ , and let  $S_1$  and  $S_2$  be relations from  $B$  to  $C$ .

00PS

1. *Interaction With Convereses.* We have

$$(R \cup S)^\dagger = R^\dagger \cup S^\dagger.$$

00PT

2. *Interaction With Composition.* We have

$$(S_1 \diamond R_1) \cup (S_2 \diamond R_2) \stackrel{\text{POSS.}}{\neq} (S_1 \cup S_2) \diamond (R_1 \cup R_2).$$

**PROOF 9.2.2.1.3 ► PROOF OF PROPOSITION 9.2.2.1.2**

Item 1: Interaction With Convereses

Clear.

Item 2: Interaction With Composition

Unwinding the definitions, we see that:

- The condition for  $(S_1 \diamond R_1) \cup (S_2 \diamond R_2)$  is:
  - There exists some  $b \in B$  such that:
    - \*  $a \sim_{R_1} b$  and  $b \sim_{S_1} c$ ;
    - or
    - \*  $a \sim_{R_2} b$  and  $b \sim_{S_2} c$ ;

- The condition for  $(S_1 \cup S_2) \diamond (R_1 \cup R_2)$  is:

– There exists some  $b \in B$  such that:

\*  $a \sim_{R_1} b$  or  $a \sim_{R_2} b$ ;

and

\*  $b \sim_{S_1} c$  or  $b \sim_{S_2} c$ .

These two conditions may fail to agree (counterexample omitted), and thus the two resulting relations on  $A \times C$  may differ. 

### 00PU 9.2.3 Unions of Families of Relations

Let  $A$  and  $B$  be sets and let  $\{R_i\}_{i \in I}$  be a family of relations from  $A$  to  $B$ .

00PV

#### DEFINITION 9.2.3.1.1 ► THE UNION OF A FAMILY OF RELATIONS

The **union of the family**  $\{R_i\}_{i \in I}$  is the relation  $\bigcup_{i \in I} R_i$  from  $A$  to  $B$  defined as follows:

- Viewing relations from  $A$  to  $B$  as subsets of  $A \times B$ , we define<sup>1</sup>

$$\bigcup_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a, b) \in (A \times B)^{\times I} \mid \begin{array}{l} \text{there exists some } i \in I \\ \text{such that } a \sim_{R_i} b \end{array} \right\}.$$

- Viewing relations from  $A$  to  $B$  as functions  $A \rightarrow \mathcal{P}(B)$ , we define

$$\left[ \bigcup_{i \in I} R_i \right](a) \stackrel{\text{def}}{=} \bigcup_{i \in I} R_i(a)$$

for each  $a \in A$ .

<sup>1</sup>This is the same as the union of  $\{R_i\}_{i \in I}$  as a collection of subsets of  $A \times B$ .

00PW

#### PROPOSITION 9.2.3.1.2 ► PROPERTIES OF UNIONS OF FAMILIES OF RELATIONS

Let  $A$  and  $B$  be sets and let  $\{R_i\}_{i \in I}$  be a family of relations from  $A$  to  $B$ .

00PX

- Interaction With Converses. We have

$$(\bigcup_{i \in I} R_i)^\dagger = \bigcup_{i \in I} R_i^\dagger.$$

**PROOF 9.2.3.1.3 ► PROOF OF PROPOSITION 9.2.3.1.2**

Item 1: Interaction With Converses

Clear.

**00PY 9.2.4 Binary Intersections of Relations**

Let  $A$  and  $B$  be sets and let  $R$  and  $S$  be relations from  $A$  to  $B$ .

**00PZ**

**DEFINITION 9.2.4.1.1 ► BINARY INTERSECTIONS OF RELATIONS**

The **intersection of  $R$  and  $S$** <sup>1</sup> is the relation  $R \cap S$  from  $A$  to  $B$  defined as follows:

- Viewing relations from  $A$  to  $B$  as subsets of  $A \times B$ , we define<sup>2</sup>

$$R \cap S \stackrel{\text{def}}{=} \{(a, b) \in B \times A \mid \text{we have } a \sim_R b \text{ and } a \sim_S b\}.$$

- Viewing relations from  $A$  to  $B$  as functions  $A \rightarrow \mathcal{P}(B)$ , we define

$$[R \cap S](a) \stackrel{\text{def}}{=} R(a) \cap S(a)$$

for each  $a \in A$ .

<sup>1</sup>Further Terminology: Also called the **binary intersection of  $R$  and  $S$** , for emphasis.

<sup>2</sup>This is the same as the intersection of  $R$  and  $S$  as subsets of  $A \times B$ .

**00Q0**

**PROPOSITION 9.2.4.1.2 ► PROPERTIES OF BINARY INTERSECTIONS OF RELATIONS**

Let  $R$ ,  $S$ ,  $R_1$ , and  $R_2$  be relations from  $A$  to  $B$ , and let  $S_1$  and  $S_2$  be relations from  $B$  to  $C$ .

**00Q1**

1. *Interaction With Converses.* We have

$$(R \cap S)^\dagger = R^\dagger \cap S^\dagger.$$

**00Q2**

2. *Interaction With Composition.* We have

$$(S_1 \diamond R_1) \cap (S_2 \diamond R_2) = (S_1 \cap S_2) \diamond (R_1 \cap R_2).$$

**PROOF 9.2.4.1.3 ► PROOF OF PROPOSITION 9.2.4.1.2**

Item 1: Interaction With Converses

Clear.

**Item 2: Interaction With Composition**

Unwinding the definitions, we see that:

- The condition for  $(S_1 \diamond R_1) \cap (S_2 \diamond R_2)$  is:
  - There exists some  $b \in B$  such that:
    - \*  $a \sim_{R_1} b$  and  $b \sim_{S_1} c$ ;
    - and
    - \*  $a \sim_{R_2} b$  and  $b \sim_{S_2} c$ ;
- The condition for  $(S_1 \cap S_2) \diamond (R_1 \cap R_2)$  is:
  - There exists some  $b \in B$  such that:
    - \*  $a \sim_{R_1} b$  and  $a \sim_{R_2} b$ ;
    - and
    - \*  $b \sim_{S_1} c$  and  $b \sim_{S_2} c$ .

These two conditions agree, and thus so do the two resulting relations on  $A \times C$ . 

**00Q3 9.2.5 Intersections of Families of Relations**

Let  $A$  and  $B$  be sets and let  $\{R_i\}_{i \in I}$  be a family of relations from  $A$  to  $B$ .

**00Q4**

**DEFINITION 9.2.5.1.1 ► THE INTERSECTION OF A FAMILY OF RELATIONS**

The **intersection of the family**  $\{R_i\}_{i \in I}$  is the relation  $\bigcup_{i \in I} R_i$  defined as follows:

- Viewing relations from  $A$  to  $B$  as subsets of  $A \times B$ , we define<sup>1</sup>

$$\bigcup_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a, b) \in (A \times B)^{\times I} \mid \begin{array}{l} \text{for each } i \in I, \\ \text{we have } a \sim_{R_i} b \end{array} \right\}.$$

- Viewing relations from  $A$  to  $B$  as functions  $A \rightarrow \mathcal{P}(B)$ , we define

$$\left[ \bigcap_{i \in I} R_i \right](a) \stackrel{\text{def}}{=} \bigcap_{i \in I} R_i(a)$$

for each  $a \in A$ .

---

<sup>1</sup>This is the same as the intersection of  $\{R_i\}_{i \in I}$  as a collection of subsets of  $A \times B$ .

00Q5

**PROPOSITION 9.2.5.1.2 ► PROPERTIES OF INTERSECTIONS OF FAMILIES OF RELATIONS**

Let  $A$  and  $B$  be sets and let  $\{R_i\}_{i \in I}$  be a family of relations from  $A$  to  $B$ .

00Q6

1. *Interaction With Converses.* We have

$$(\bigcap_{i \in I} R_i)^\dagger = \bigcap_{i \in I} R_i^\dagger.$$

**PROOF 9.2.5.1.3 ► PROOF OF PROPOSITION 9.2.5.1.2**

Item 1: Interaction With Converses

Clear. 

**00Q7 9.2.6 Binary Products of Relations**

Let  $A$ ,  $B$ ,  $X$ , and  $Y$  be sets, let  $R: A \rightarrow B$  be a relation from  $A$  to  $B$ , and let  $S: X \rightarrow Y$  be a relation from  $X$  to  $Y$ .

00Q8

**DEFINITION 9.2.6.1.1 ► BINARY PRODUCTS OF RELATIONS**

The **product of  $R$  and  $S$** <sup>1</sup> is the relation  $R \times S$  from  $A \times X$  to  $B \times Y$  defined as follows:

- Viewing relations from  $A \times X$  to  $B \times Y$  as subsets of  $(A \times X) \times (B \times Y)$ , we define  $R \times S$  as the Cartesian product of  $R$  and  $S$  as subsets of  $A \times X$  and  $B \times Y$ .<sup>2</sup>
- Viewing relations from  $A \times X$  to  $B \times Y$  as functions  $A \times X \rightarrow \mathcal{P}(B \times Y)$ , we define  $R \times S$  as the composition

$$A \times X \xrightarrow{R \times S} \mathcal{P}(B) \times \mathcal{P}(Y) \xrightarrow{\mathcal{P}_{B,Y}^\otimes} \mathcal{P}(B \times Y)$$

in Sets, i.e. by

$$[R \times S](a, x) \stackrel{\text{def}}{=} R(a) \times S(x)$$

for each  $(a, x) \in A \times X$ .

<sup>1</sup>Further Terminology: Also called the **binary product of  $R$  and  $S$**  for emphasis. That is,  $R \times S$  is the relation given by declaring  $(a, x) \sim_{R \times S} (b, y)$  iff  $a \sim_R b$  and  $x \sim_S y$ .

00Q9

**PROPOSITION 9.2.6.1.2 ► PROPERTIES OF BINARY PRODUCTS OF RELATIONS**

Let  $A$ ,  $B$ ,  $X$ , and  $Y$  be sets.

00QA

1. *Interaction With Converses.* Let

$$\begin{aligned} R : A &\rightarrow A, \\ S : X &\rightarrow X \end{aligned}$$

We have

$$(R \times S)^\dagger = R^\dagger \times S^\dagger.$$

00QB

2. *Interaction With Composition.* Let

$$\begin{aligned} R_1 : A &\rightarrow B, \\ S_1 : B &\rightarrow C, \\ R_2 : X &\rightarrow Y, \\ S_2 : Y &\rightarrow Z \end{aligned}$$

be relations. We have

$$(S_1 \diamond R_1) \times (S_2 \diamond R_2) = (S_1 \times S_2) \diamond (R_1 \times R_2).$$

#### PROOF 9.2.6.1.3 ► PROOF OF PROPOSITION 9.2.6.1.2

##### Item 1: Interaction With Converses

Unwinding the definitions, we see that:

- We have  $(a, x) \sim_{(R \times S)^\dagger} (b, y)$  iff:
  - We have  $(b, y) \sim_{R \times S} (a, x)$ , i.e. iff:
    - \* We have  $b \sim_R a$ ;
    - \* We have  $y \sim_S x$ ;
- We have  $(a, x) \sim_{R^\dagger \times S^\dagger} (b, y)$  iff:
  - We have  $a \sim_{R^\dagger} b$  and  $x \sim_{S^\dagger} y$ , i.e. iff:
    - \* We have  $b \sim_R a$ ;
    - \* We have  $y \sim_S x$ .

These two conditions agree, and thus the two resulting relations on  $A \times X$  are equal.

##### Item 2: Interaction With Composition

Unwinding the definitions, we see that:

- We have  $(a, x) \sim_{(S_1 \diamond R_1) \times (S_2 \diamond R_2)} (c, z)$  iff:

- We have  $a \sim_{S_1 \diamond R_1} c$  and  $x \sim_{S_2 \diamond R_2} z$ , i.e. iff:
  - \* There exists some  $b \in B$  such that  $a \sim_{R_1} b$  and  $b \sim_{S_1} c$ ;
  - \* There exists some  $y \in Y$  such that  $x \sim_{R_2} y$  and  $y \sim_{S_2} z$ ;
- We have  $(a, x) \sim_{(S_1 \times S_2) \diamond (R_1 \times R_2)} (c, z)$  iff:
  - There exists some  $(b, y) \in B \times Y$  such that  $(a, x) \sim_{R_1 \times R_2} (b, y)$  and  $(b, y) \sim_{S_1 \times S_2} (c, z)$ , i.e. such that:
    - \* We have  $a \sim_{R_1} b$  and  $x \sim_{R_2} y$ ;
    - \* We have  $b \sim_{S_1} c$  and  $y \sim_{S_2} z$ .

These two conditions agree, and thus the two resulting relations from  $A \times X$  to  $C \times Z$  are equal. 

### 00QC 9.2.7 Products of Families of Relations

Let  $\{A_i\}_{i \in I}$  and  $\{B_i\}_{i \in I}$  be families of sets, and let  $\{R_i: A_i \rightarrow B_i\}_{i \in I}$  be a family of relations.

00QD

#### DEFINITION 9.2.7.1.1 ► THE PRODUCT OF A FAMILY OF RELATIONS

The **product of the family**  $\{R_i\}_{i \in I}$  is the relation  $\prod_{i \in I} R_i$  from  $\prod_{i \in I} A_i$  to  $\prod_{i \in I} B_i$  defined as follows:

- Viewing relations as subsets, we define  $\prod_{i \in I} R_i$  as its product as a family of sets, i.e. we have

$$\prod_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a_i, b_i)_{i \in I} \in \prod_{i \in I} (A_i \times B_i) \mid \begin{array}{l} \text{for each } i \in I, \\ \text{we have } a_i \sim_{R_i} b_i \end{array} \right\}.$$

- Viewing relations as functions to powersets, we define

$$\left[ \prod_{i \in I} R_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} R_i(a_i)$$

for each  $(a_i)_{i \in I} \in \prod_{i \in I} R_i$ .

### 00R2 9.2.8 The Collage of a Relation

Let  $A$  and  $B$  be sets and let  $R: A \rightarrow B$  be a relation from  $A$  to  $B$ .

00R3

## DEFINITION 9.2.8.1.1 ► THE COLLEGE OF A RELATION

The **collage** of  $R^1$  is the poset  $\text{Coll}(R) \stackrel{\text{def}}{=} (\text{Coll}(R), \preceq_{\text{Coll}(R)})$  consisting of:

- *The Underlying Set.* The set  $\text{Coll}(R)$  defined by

$$\text{Coll}(R) \stackrel{\text{def}}{=} A \coprod B.$$

- *The Partial Order.* The partial order

$$\preceq_{\text{Coll}(R)} : \text{Coll}(R) \times \text{Coll}(R) \rightarrow \{\text{true}, \text{false}\}$$

on  $\text{Coll}(R)$  defined by

$$\preceq(a, b) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } a = b \text{ or } a \sim_R b, \\ \text{false} & \text{otherwise.} \end{cases}$$

---

<sup>1</sup>Further Terminology: Also called the **cograph** of  $R$ .


02AZ

**NOTATION 9.2.8.1.2 ► NOTATION:**  $\text{Pos}_{/\Delta^1}(A, B)$

We write  $\text{Pos}_{/\Delta^1}(A, B)$  for the category defined as the pullback

$$\text{Pos}_{/\Delta^1}(A, B) \stackrel{\text{def}}{=} \text{pt} \times_{[A], \text{Pos}, \text{ev}_0} \text{Pos}_{/\Delta^1} \times_{\text{ev}_1, \text{Pos}, [B]} \text{pt},$$

as in the diagram



02B0

#### REMARK 9.2.8.1.3 ► UNWINDING NOTATION 9.2.8.1.2

In detail,  $\text{Pos}_{/\Delta^1}(A, B)$  is the category where:

- *Objects.* An object of  $\text{Pos}_{/\Delta^1}(A, B)$  is a pair  $(X, \phi_X)$  consisting of
    - A poset  $X$ ;

- A morphism  $\phi_X: X \rightarrow \Delta^1$ ;

such that we have

$$\begin{aligned}\phi_X^{-1}(0) &= A, \\ \phi_X^{-1}(1) &= B.\end{aligned}$$

- *Morphisms.* A morphism of  $\text{Pos}_{/\Delta^1}(A, B)$  from  $(X, \phi_X)$  to  $(Y, \phi_Y)$  is a morphism of posets  $f: X \rightarrow Y$  making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \phi_X \searrow & & \swarrow \phi_Y \\ & \Delta^1 & \end{array}$$

commute.

00R4

#### PROPOSITION 9.2.8.1.4 ► PROPERTIES OF COLLAGES OF RELATIONS

Let  $A$  and  $B$  be sets and let  $R: A \rightarrow B$  be a relation from  $A$  to  $B$ .

00R5

1. *Functoriality.* The assignment  $R \mapsto \text{Coll}(R)$  defines a functor

$$\text{Coll}: \text{Rel}(A, B) \rightarrow \text{Pos}_{/\Delta^1}(A, B),$$

where

- *Action on Objects.* For each  $R \in \text{Obj}(\text{Rel}(A, B))$ , we have

$$[\text{Coll}](R) \stackrel{\text{def}}{=} (\text{Coll}(R), \phi_R)$$

for each  $R \in \text{Rel}(A, B)$ , where

- The poset  $\text{Coll}(R)$  is the collage of  $R$  of [Definition 9.2.8.1.1](#).
- The morphism  $\phi_R: \text{Coll}(R) \rightarrow \Delta^1$  is given by

$$\phi_R(x) \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x \in A, \\ 1 & \text{if } x \in B \end{cases}$$

for each  $x \in \text{Coll}(R)$ .

- *Action on Morphisms.* For each  $R, S \in \text{Obj}(\text{Rel}(A, B))$ , the action on Hom-sets

$$\text{Coll}_{R,S}: \text{Hom}_{\text{Rel}(A, B)}(R, S) \rightarrow \text{Pos}(\text{Coll}(R), \text{Coll}(S))$$

of  $\mathbf{Coll}$  at  $(R, S)$  is given by sending an inclusion

$$\iota: R \subset S$$

to the morphism

$$\mathbf{Coll}(\iota): \mathbf{Coll}(R) \rightarrow \mathbf{Coll}(S)$$

of posets over  $\Delta^1$  defined by

$$[\mathbf{Coll}(\iota)](x) \stackrel{\text{def}}{=} x$$

for each  $x \in \mathbf{Coll}(R)$ .<sup>1</sup>

00R6 2. *Equivalence.* The functor of **Item 1** is an equivalence of categories.

---

<sup>1</sup>Note that this is indeed a morphism of posets: if  $x \preceq_{\mathbf{Coll}(R)} y$ , then  $x = y$  or  $x \sim_R y$ , so we have either  $x = y$  or  $x \sim_S y$  (as  $R \subset S$ ), and thus  $x \preceq_{\mathbf{Coll}(S)} y$ .

**PROOF 9.2.8.1.5 ▶ PROOF OF PROPOSITION 9.2.8.1.4**

Item 1: Functoriality

Clear.

Item 2: Equivalence

Omitted. 

# Appendices

## 9.A Other Chapters

### Preliminaries

- 1. Introduction
- 2. A Guide to the Literature

### Sets

- 3. Sets
- 4. Constructions With Sets
- 5. Monoidal Structures on the Category of Sets
- 6. Pointed Sets

### 7. Tensor Products of Pointed Sets

### Relations

- 8. Relations
- 9. Constructions With Relations
- 10. Conditions on Relations

### Categories

- 11. Categories
- 12. Presheaves and the Yoneda Lemma

### Monoidal Categories

13. Constructions With Monoidal Categories

gories

**Bicategories**

**Extra Part**

14. Types of Morphisms in Bicate-

15. Notes

# Chapter 10

## Conditions on Relations

**02H3** This chapter contains some material about reflexive, symmetric, transitive, equivalence, and apartness relations.

### Contents

---

10.1 Functional and Total Relations.....	704
10.1.1 Functional Relations.....	704
10.1.2 Total Relations.....	705
10.2 Reflexive Relations.....	706
10.2.1 Foundations .....	706
10.2.2 The Reflexive Closure of a Relation.....	707
10.3 Symmetric Relations.....	709
10.3.1 Foundations .....	709
10.3.2 The Symmetric Closure of a Relation .....	710
10.4 Transitive Relations.....	713
10.4.1 Foundations .....	713
10.4.2 The Transitive Closure of a Relation .....	714
10.5 Equivalence Relations.....	717
10.5.1 Foundations .....	717
10.5.2 The Equivalence Closure of a Relation.....	717
10.6 Quotients by Equivalence Relations .....	720
10.6.1 Equivalence Classes .....	720
10.6.2 Quotients of Sets by Equivalence Relations.....	721
10.A Other Chapters.....	725

---

**02D0 10.1 Functional and Total Relations**

**00JC 10.1.1 Functional Relations**

Let  $A$  and  $B$  be sets.

00JD

**DEFINITION 10.1.1.1 ► FUNCTIONAL RELATIONS**

A relation  $R: A \rightarrow B$  is **functional** if, for each  $a \in A$ , the set  $R(a)$  is either empty or a singleton.

00JE

**PROPOSITION 10.1.1.2 ► PROPERTIES OF FUNCTIONAL RELATIONS**

Let  $R: A \rightarrow B$  be a relation.

00JF

1. *Characterisations.* The following conditions are equivalent:

00JG

(a) The relation  $R$  is functional.

00JH

(b) We have  $R \diamond R^\dagger \subset \chi_B$ .

**PROOF 10.1.1.3 ► PROOF OF PROPOSITION 10.1.1.2****Item 1: Characterisations**

We claim that **Items 1a** and **1b** are indeed equivalent:

- *Item 1a*  $\implies$  *Item 1b*: Let  $(b, b') \in B \times B$ . We need to show that

$$[R \diamond R^\dagger](b, b') \preceq_{\{\text{t,f}\}} \chi_B(b, b'),$$

i.e. that if there exists some  $a \in A$  such that  $b \sim_{R^\dagger} a$  and  $a \sim_R b'$ , then  $b = b'$ . But since  $b \sim_{R^\dagger} a$  is the same as  $a \sim_R b$ , we have both  $a \sim_R b$  and  $a \sim_R b'$  at the same time, which implies  $b = b'$  since  $R$  is functional.

- *Item 1b*  $\implies$  *Item 1a*: Suppose that we have  $a \sim_R b$  and  $a \sim_R b'$  for  $b, b' \in B$ . We claim that  $b = b'$ :

- Since  $a \sim_R b$ , we have  $b \sim_{R^\dagger} a$ .
- Since  $R \diamond R^\dagger \subset \chi_B$ , we have

$$[R \diamond R^\dagger](b, b') \preceq_{\{\text{t,f}\}} \chi_B(b, b'),$$

and since  $b \sim_{R^\dagger} a$  and  $a \sim_R b'$ , it follows that  $[R \diamond R^\dagger](b, b') = \text{true}$ , and thus  $\chi_B(b, b') = \text{true}$  as well, i.e.  $b = b'$ .

This finishes the proof. 

**00JJ 10.1.2 Total Relations**

Let  $A$  and  $B$  be sets.

00JK

**DEFINITION 10.1.2.1.1 ► TOTAL RELATIONS**

A relation  $R: A \rightarrow B$  is **total** if, for each  $a \in A$ , we have  $R(a) \neq \emptyset$ .

00JL

**PROPOSITION 10.1.2.1.2 ► PROPERTIES OF TOTAL RELATIONS**

Let  $R: A \rightarrow B$  be a relation.

00JM

1. *Characterisations.* The following conditions are equivalent:

00JN

(a) The relation  $R$  is total.

00JP

(b) We have  $\chi_A \subset R^\dagger \diamond R$ .

**PROOF 10.1.2.1.3 ► PROOF OF PROPOSITION 10.1.2.1.2****Item 1: Characterisations**

We claim that **Items 1a** and **1b** are indeed equivalent:

- **Item 1a  $\implies$  Item 1b:** We have to show that, for each  $(a, a') \in A$ , we have

$$\chi_A(a, a') \preceq_{\{t,f\}} [R^\dagger \diamond R](a, a'),$$

i.e. that if  $a = a'$ , then there exists some  $b \in B$  such that  $a \sim_R b$  and  $b \sim_{R^\dagger} a'$  (i.e.  $a \sim_R b$  again), which follows from the totality of  $R$ .

- **Item 1b  $\implies$  Item 1a:** Given  $a \in A$ , since  $\chi_A \subset R^\dagger \diamond R$ , we must have

$$\{a\} \subset [R^\dagger \diamond R](a),$$

implying that there must exist some  $b \in B$  such that  $a \sim_R b$  and  $b \sim_{R^\dagger} a$  (i.e.  $a \sim_R b$ ) and thus  $R(a) \neq \emptyset$ , as  $b \in R(a)$ .

This finishes the proof. 

**00TK 10.2 Reflexive Relations****00TL 10.2.1 Foundations**

Let  $A$  be a set.

00TM

**DEFINITION 10.2.1.1.1 ► REFLEXIVE RELATIONS**

A **reflexive relation** is equivalently:<sup>1</sup>

- An  $\mathbb{E}_0$ -monoid in  $(N_\bullet(\text{Rel}(A, A)), \chi_A)$ .

- A pointed object in  $(\mathbf{Rel}(A, A), \chi_A)$ .

<sup>1</sup>Note that since  $\mathbf{Rel}(A, A)$  is posetal, reflexivity is a property of a relation, rather than extra structure.

00TN

**REMARK 10.2.1.1.2 ► UNWINDING DEFINITION 10.2.1.1**

In detail, a relation  $R$  on  $A$  is **reflexive** if we have an inclusion

$$\eta_R : \chi_A \subset R$$

of relations in  $\mathbf{Rel}(A, A)$ , i.e. if, for each  $a \in A$ , we have  $a \sim_R a$ .

00TP

**DEFINITION 10.2.1.3 ► THE PO/SSET OF REFLEXIVE RELATIONS ON A SET**

Let  $A$  be a set.

00TQ

1. The **set of reflexive relations on  $A$**  is the subset  $\mathbf{Rel}^{\text{refl}}(A, A)$  of  $\mathbf{Rel}(A, A)$  spanned by the reflexive relations.

00TR

2. The **poset of relations on  $A$**  is the subposet  $\mathbf{Rel}^{\text{refl}}(A, A)$  of  $\mathbf{Rel}(A, A)$  spanned by the reflexive relations.

00TS

**PROPOSITION 10.2.1.4 ► PROPERTIES OF REFLEXIVE RELATIONS**

Let  $R$  and  $S$  be relations on  $A$ .

00TT

1. *Interaction With Inverses.* If  $R$  is reflexive, then so is  $R^\dagger$ .

00TU

2. *Interaction With Composition.* If  $R$  and  $S$  are reflexive, then so is  $S \diamond R$ .

**PROOF 10.2.1.5 ► PROOF OF PROPOSITION 10.2.1.4**

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

Clear.

**00TV 10.2.2 The Reflexive Closure of a Relation**

Let  $R$  be a relation on  $A$ .

00TW

## DEFINITION 10.2.2.1.1 ► THE REFLEXIVE CLOSURE OF A RELATION

The **reflexive closure** of  $\sim_R$  is the relation  $\sim_R^{\text{refl}}$ <sup>1</sup> satisfying the following universal property:<sup>2</sup>

- (★) Given another reflexive relation  $\sim_S$  on  $A$  such that  $R \subset S$ , there exists an inclusion  $\sim_R^{\text{refl}} \subset \sim_S$ .

<sup>1</sup>Further Notation: Also written  $R^{\text{refl}}$ .

<sup>2</sup>Slogan: The reflexive closure of  $R$  is the smallest reflexive relation containing  $R$ .

00TX

## CONSTRUCTION 10.2.2.1.2 ► THE REFLEXIVE CLOSURE OF A RELATION

Concretely,  $\sim_R^{\text{refl}}$  is the free pointed object on  $R$  in  $(\mathbf{Rel}(A, A), \chi_A)$ <sup>1</sup>, being given by

$$\begin{aligned} R^{\text{refl}} &\stackrel{\text{def}}{=} R \coprod^{\mathbf{Rel}(A, A)} \Delta_A \\ &= R \cup \Delta_A \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } a = b\}. \end{aligned}$$

<sup>1</sup>Or, equivalently, the free  $\mathbb{E}_0$ -monoid on  $R$  in  $(\mathbf{N}_*(\mathbf{Rel}(A, A)), \chi_A)$ .

## PROOF 10.2.2.1.3 ► PROOF OF CONSTRUCTION 10.2.2.1.2

Clear. 

00TY

## PROPOSITION 10.2.2.1.4 ► PROPERTIES OF THE REFLEXIVE CLOSURE OF A RELATION

Let  $R$  be a relation on  $A$ .

00TZ

1. *Adjointness.* We have an adjunction

$$\left( (-)^{\text{refl}} \dashv \overline{\text{忘}} \right): \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{refl}}} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \mathbf{Rel}^{\text{refl}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{refl}}(R^{\text{refl}}, S) \cong \mathbf{Rel}(R, S),$$

natural in  $R \in \text{Obj}(\mathbf{Rel}^{\text{refl}}(A, A))$  and  $S \in \text{Obj}(\mathbf{Rel}(A, A))$ .

00U0

2. *The Reflexive Closure of a Reflexive Relation.* If  $R$  is reflexive, then  $R^{\text{refl}} = R$ .

00U1

3. *Idempotency.* We have

$$(R^{\text{refl}})^{\text{refl}} = R^{\text{refl}}.$$

00U2

4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} & \text{Rel}(A, A) \\ \left(R^\dagger\right)^{\text{refl}} = \left(R^{\text{refl}}\right)^\dagger, & \downarrow (-)^\dagger & \downarrow (-)^\dagger \\ \text{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} & \text{Rel}(A, A). \end{array}$$

00U3

5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\ (S \diamond R)^{\text{refl}} = S^{\text{refl}} \diamond R^{\text{refl}}, & \downarrow (-)^{\text{refl}} \times (-)^{\text{refl}} & \downarrow (-)^{\text{refl}} \\ \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A). \end{array}$$

#### PROOF 10.2.2.1.5 ► PROOF OF PROPOSITION 10.2.2.1.4

**Item 1: Adjointness**

This is a rephrasing of the universal property of the reflexive closure of a relation, stated in [Definition 10.2.2.1.1](#).

**Item 2: The Reflexive Closure of a Reflexive Relation**

Clear.

**Item 3: Idempotency**

This follows from [Item 2](#).

**Item 4: Interaction With Inverses**

Clear.

**Item 5: Interaction With Composition**

This follows from [Item 2](#) of [Proposition 10.2.1.1.4](#). 

## 00U4 10.3 Symmetric Relations

### 00U5 10.3.1 Foundations

Let  $A$  be a set.

00U6

#### DEFINITION 10.3.1.1.1 ► SYMMETRIC RELATIONS

A relation  $R$  on  $A$  is **symmetric** if we have  $R^\dagger = R$ .

00U7

**REMARK 10.3.1.1.2 ► UNWINDING DEFINITION 10.3.1.1**

In detail, a relation  $R$  is symmetric if it satisfies the following condition:

- (★) For each  $a, b \in A$ , if  $a \sim_R b$ , then  $b \sim_R a$ .

00U8

**DEFINITION 10.3.1.1.3 ► THE PO/SET OF SYMMETRIC RELATIONS ON A SET**

Let  $A$  be a set.

00U9

1. The **set of symmetric relations on  $A$**  is the subset  $\text{Rel}^{\text{symm}}(A, A)$  of  $\text{Rel}(A, A)$  spanned by the symmetric relations.

00UA

2. The **poset of relations on  $A$**  is the subposet  $\text{Rel}^{\text{symm}}(A, A)$  of  $\text{Rel}(A, A)$  spanned by the symmetric relations.

00UB

**PROPOSITION 10.3.1.1.4 ► PROPERTIES OF SYMMETRIC RELATIONS**

Let  $R$  and  $S$  be relations on  $A$ .

00UC

1. *Interaction With Inverses.* If  $R$  is symmetric, then so is  $R^\dagger$ .

00UD

2. *Interaction With Composition.* If  $R$  and  $S$  are symmetric, then so is  $S \diamond R$ .

**PROOF 10.3.1.1.5 ► PROOF OF PROPOSITION 10.3.1.1.4**

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

Clear.

**00UE 10.3.2 The Symmetric Closure of a Relation**

Let  $R$  be a relation on  $A$ .

00UF

**DEFINITION 10.3.2.1.1 ► THE SYMMETRIC CLOSURE OF A RELATION**

The **symmetric closure** of  $\sim_R$  is the relation  $\sim_R^{\text{symm}}$ <sup>1</sup> satisfying the following universal property:<sup>2</sup>

- (★) Given another symmetric relation  $\sim_S$  on  $A$  such that  $R \subset S$ , there exists an inclusion  $\sim_R^{\text{symm}} \subset \sim_S$ .

<sup>1</sup>Further Notation: Also written  $R^{\text{symm}}$ .

<sup>2</sup>Slogan: The symmetric closure of  $R$  is the smallest symmetric relation containing  $R$ .

00UG

**CONSTRUCTION 10.3.2.1.2 ► THE SYMMETRIC CLOSURE OF A RELATION**

Concretely,  $\sim_R^{\text{symm}}$  is the symmetric relation on  $A$  defined by

$$\begin{aligned} R^{\text{symm}} &\stackrel{\text{def}}{=} R \cup R^\dagger \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } b \sim_R a\}. \end{aligned}$$

**PROOF 10.3.2.1.3 ► PROOF OF CONSTRUCTION 10.3.2.1.2**

Clear. 

00UH

**PROPOSITION 10.3.2.1.4 ► PROPERTIES OF THE SYMMETRIC CLOSURE OF A RELATION**

Let  $R$  be a relation on  $A$ .

00UJ

1. *Adjointness.* We have an adjunction

$$((-)^{\text{symm}} \dashv \overline{\vdash}): \quad \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{symm}}} \\ \perp \\ \xleftarrow{\overline{\vdash}} \end{array} \mathbf{Rel}^{\text{symm}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{symm}}(R^{\text{symm}}, S) \cong \mathbf{Rel}(R, S),$$

natural in  $R \in \text{Obj}(\mathbf{Rel}^{\text{symm}}(A, A))$  and  $S \in \text{Obj}(\mathbf{Rel}(A, A))$ .

00UK

2. *The Symmetric Closure of a Symmetric Relation.* If  $R$  is symmetric, then  $R^{\text{symm}} = R$ .

00UL

3. *Idempotency.* We have

$$(R^{\text{symm}})^{\text{symm}} = R^{\text{symm}}.$$

00UM

4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{symm}}} & \mathbf{Rel}(A, A) \\ \left(R^\dagger\right)^{\text{symm}} = \left(R^{\text{symm}}\right)^\dagger, & \downarrow (-)^\dagger & \downarrow (-)^\dagger \\ \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{symm}}} & \mathbf{Rel}(A, A). \end{array}$$

00UN

5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \mathbf{Rel}(A, A) \times \mathbf{Rel}(A, A) & \xrightarrow{\diamond} & \mathbf{Rel}(A, A) \\ (S \diamond R)^{\text{symm}} = S^{\text{symm}} \diamond R^{\text{symm}}, & \downarrow (-)^{\text{symm}} \times (-)^{\text{symm}} & \downarrow (-)^{\text{symm}} \\ \mathbf{Rel}(A, A) \times \mathbf{Rel}(A, A) & \xrightarrow{\diamond} & \mathbf{Rel}(A, A). \end{array}$$



**Item 1: Adjointness**

This is a rephrasing of the universal property of the symmetric closure of a relation, stated in [Definition 10.3.2.1.1](#).

**Item 2: The Symmetric Closure of a Symmetric Relation**

Clear.

**Item 3: Idempotency**

This follows from [Item 2](#).

**Item 4: Interaction With Inverses**

Clear.

**Item 5: Interaction With Composition**

This follows from [Item 2](#) of [Proposition 10.3.1.1.4](#). 

**00UP 10.4 Transitive Relations****00UQ 10.4.1 Foundations**

Let  $A$  be a set.

**00UR****DEFINITION 10.4.1.1.1 ► TRANSITIVE RELATIONS**

A **transitive relation** is equivalently:<sup>1</sup>

- A non-unital  $\mathbb{E}_1$ -monoid in  $(N_*(\text{Rel}(A, A)), \diamond)$ .
- A non-unital monoid in  $(\text{Rel}(A, A), \diamond)$ .

<sup>1</sup>Note that since  $\text{Rel}(A, A)$  is posetal, transitivity is a property of a relation, rather than extra structure.

**00US****REMARK 10.4.1.1.2 ► UNWINDING DEFINITION 10.4.1.1.1**

In detail, a relation  $R$  on  $A$  is **transitive** if we have an inclusion

$$\mu_R: R \diamond R \subset R$$

of relations in  $\text{Rel}(A, A)$ , i.e. if, for each  $a, c \in A$ , the following condition is satisfied:

- (★) If there exists some  $b \in A$  such that  $a \sim_R b$  and  $b \sim_R c$ , then  $a \sim_R c$ .

00UT

**DEFINITION 10.4.1.3 ► THE PO/SET OF TRANSITIVE RELATIONS ON A SET**

Let  $A$  be a set.

00UU

1. The **set of transitive relations from  $A$  to  $B$**  is the subset  $\text{Rel}^{\text{trans}}(A)$  of  $\text{Rel}(A, A)$  spanned by the transitive relations.

00UV

2. The **poset of relations from  $A$  to  $B$**  is the subposet  $\text{Rel}^{\text{trans}}(A)$  of  $\text{Rel}(A, A)$  spanned by the transitive relations.

00UW

**PROPOSITION 10.4.1.4 ► PROPERTIES OF TRANSITIVE RELATIONS**

Let  $R$  and  $S$  be relations on  $A$ .

00UX

1. *Interaction With Inverses.* If  $R$  is transitive, then so is  $R^\dagger$ .

00UY

2. *Interaction With Composition.* If  $R$  and  $S$  are transitive, then  $S \diamond R$  may fail to be transitive.

**PROOF 10.4.1.5 ► PROOF OF PROPOSITION 10.4.1.4**

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

See [MSE 2096272].<sup>1</sup>



<sup>1</sup>Intuition: Transitivity for  $R$  and  $S$  fails to imply that of  $S \diamond R$  because the composition operation for relations intertwines  $R$  and  $S$  in an incompatible way:

- If  $a \sim_{S \diamond R} c$  and  $c \sim_{S \diamond R} e$ , then:
  - There is some  $b \in A$  such that:
    - \*  $a \sim_R b$ ;
    - \*  $b \sim_S c$ ;
  - There is some  $d \in A$  such that:
    - \*  $c \sim_R d$ ;
    - \*  $d \sim_S e$ .

**00UZ 10.4.2 The Transitive Closure of a Relation**

Let  $R$  be a relation on  $A$ .

00V0

**DEFINITION 10.4.2.1.1 ► THE TRANSITIVE CLOSURE OF A RELATION**

The **transitive closure** of  $\sim_R$  is the relation  $\sim_R^{\text{trans}1}$  satisfying the following universal property:<sup>2</sup>

- (★) Given another transitive relation  $\sim_S$  on  $A$  such that  $R \subset S$ , there exists an inclusion  $\sim_R^{\text{trans}} \subset \sim_S$ .

<sup>1</sup>Further Notation: Also written  $R^{\text{trans}}$ .

<sup>2</sup>Slogan: The transitive closure of  $R$  is the smallest transitive relation containing  $R$ .

00V1

#### CONSTRUCTION 10.4.2.1.2 ► THE TRANSITIVE CLOSURE OF A RELATION

Concretely,  $\sim_R^{\text{trans}}$  is the free non-unital monoid on  $R$  in  $(\mathbf{Rel}(A, A), \diamond)$ <sup>1</sup>, being given by

$$\begin{aligned} R^{\text{trans}} &\stackrel{\text{def}}{=} \bigsqcup_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \bigcup_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists some } (x_1, \dots, x_n) \in R^{\times n} \\ \text{such that } a \sim_R x_1 \sim_R \dots \sim_R x_n \sim_R b \end{array} \right\}. \end{aligned}$$

<sup>1</sup>Or, equivalently, the free non-unital  $\mathbb{E}_1$ -monoid on  $R$  in  $(N_*(\mathbf{Rel}(A, A)), \diamond)$ .

#### PROOF 10.4.2.1.3 ► PROOF OF CONSTRUCTION 10.4.2.1.2

Clear. 

00V2

#### PROPOSITION 10.4.2.1.4 ► PROPERTIES OF THE TRANSITIVE CLOSURE OF A RELATION

Let  $R$  be a relation on  $A$ .

00V3

1. *Adjointness.* We have an adjunction

$$((-)^{\text{trans}} \dashv \overline{\text{Res}}) : \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{trans}}} \\ \perp \\ \xleftarrow{\text{Res}} \end{array} \mathbf{Rel}^{\text{trans}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{trans}}(R^{\text{trans}}, S) \cong \mathbf{Rel}(R, S),$$

natural in  $R \in \text{Obj}(\mathbf{Rel}^{\text{trans}}(A, A))$  and  $S \in \text{Obj}(\mathbf{Rel}(A, B))$ .

00V4

2. *The Transitive Closure of a Transitive Relation.* If  $R$  is transitive, then  $R^{\text{trans}} = R$ .

00V5

3. *Idempotency.* We have

$$(R^{\text{trans}})^{\text{trans}} = R^{\text{trans}}.$$

00V6

4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A) \\ \left(R^\dagger\right)^{\text{trans}} = \left(R^{\text{trans}}\right)^\dagger, & \downarrow (-)^\dagger & \downarrow (-)^\dagger \\ \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A). \end{array}$$

00V7

5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\ (S \diamond R)^{\text{trans}} \stackrel{\text{poss.}}{\neq} S^{\text{trans}} \diamond R^{\text{trans}}, & \downarrow (-)^{\text{trans}} \times (-)^{\text{trans}} & \downarrow (-)^{\text{trans}} \\ \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A). \end{array}$$

#### PROOF 10.4.2.1.5 ► PROOF OF PROPOSITION 10.4.2.1.4

##### Item 1: Adjointness

This is a rephrasing of the universal property of the transitive closure of a relation, stated in [Definition 10.4.2.1.1](#).

##### Item 2: The Transitive Closure of a Transitive Relation

Clear.

##### Item 3: Idempotency

This follows from [Item 2](#).

##### Item 4: Interaction With Inverses

We have

$$\begin{aligned} (R^\dagger)^{\text{trans}} &= \bigcup_{n=1}^{\infty} (R^\dagger)^{\diamond n} \\ &= \bigcup_{n=1}^{\infty} (R^{\diamond n})^\dagger \\ &= (\bigcup_{n=1}^{\infty} R^{\diamond n})^\dagger \\ &= (R^{\text{trans}})^\dagger, \end{aligned}$$

where we have used, respectively:

- Construction 10.4.2.1.2.
- ?? of Proposition 8.1.3.1.4.
- ?? of Proposition 9.2.3.1.2.
- Construction 10.4.2.1.2.

This finishes the proof.

Item 5: Interaction With Composition

This follows from Item 2 of Proposition 10.4.1.1.4.



## 00V8 10.5 Equivalence Relations

### 00V9 10.5.1 Foundations

Let  $A$  be a set.

00VA

#### DEFINITION 10.5.1.1.1 ► EQUIVALENCE RELATIONS

A relation  $R$  is an **equivalence relation** if it is reflexive, symmetric, and transitive.<sup>1</sup>

<sup>1</sup>*Further Terminology:* If instead  $R$  is just symmetric and transitive, then it is called a **partial equivalence relation**.

00VB

#### EXAMPLE 10.5.1.1.2 ► THE KERNEL OF A FUNCTION

The **kernel of a function**  $f: A \rightarrow B$  is the equivalence relation  $\sim_{\text{Ker}(f)}$  on  $A$  obtained by declaring  $a \sim_{\text{Ker}(f)} b$  iff  $f(a) = f(b)$ .<sup>1</sup>

<sup>1</sup>The kernel  $\text{Ker}(f): A \rightarrow A$  of  $f$  is the underlying functor of the monad induced by the adjunction  $\text{Gr}(f) \dashv f^{-1}: A \rightleftarrows B$  in **Rel** of Item 4 of Proposition 8.2.2.1.2.

00VC

#### DEFINITION 10.5.1.1.3 ► THE PO/SSET OF EQUIVALENCE RELATIONS ON A SET

Let  $A$  and  $B$  be sets.

00VD

1. The **set of equivalence relations from  $A$  to  $B$**  is the subset  $\text{Rel}^{\text{eq}}(A, B)$  of  $\text{Rel}(A, B)$  spanned by the equivalence relations.

00VE

2. The **poset of relations from  $A$  to  $B$**  is the subposet  $\text{Rel}^{\text{eq}}(A, B)$  of  $\text{Rel}(A, B)$  spanned by the equivalence relations.

## 00VF 10.5.2 The Equivalence Closure of a Relation

Let  $R$  be a relation on  $A$ .

00VG

The **equivalence closure**<sup>1</sup> of  $\sim_R$  is the relation  $\sim_R^{\text{eq}}$ <sup>2</sup> satisfying the following universal property:<sup>3</sup>

- (★) Given another equivalence relation  $\sim_S$  on  $A$  such that  $R \subset S$ , there exists an inclusion  $\sim_R^{\text{eq}} \subset \sim_S$ .

<sup>1</sup>Further Terminology: Also called the **equivalence relation associated to**  $\sim_R$ .

<sup>2</sup>Further Notation: Also written  $R^{\text{eq}}$ .

<sup>3</sup>Slogan: The equivalence closure of  $R$  is the smallest equivalence relation containing  $R$ .

00VH

#### CONSTRUCTION 10.5.2.1.2 ► THE EQUIVALENCE CLOSURE OF A RELATION

Concretely,  $\sim_R^{\text{eq}}$  is the equivalence relation on  $A$  defined by

$$\begin{aligned} R^{\text{eq}} &\stackrel{\text{def}}{=} ((R^{\text{refl}})^{\text{symm}})^{\text{trans}} \\ &= ((R^{\text{symm}})^{\text{trans}})^{\text{refl}} \\ &= \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists } (x_1, \dots, x_n) \in R^{\times n} \text{ satisfying at} \\ \text{least one of the following conditions:} \end{array} \right\} \\ &\quad \left. \begin{array}{l} 1. \text{ The following conditions are satisfied:} \\ \quad \begin{array}{l} (a) \text{ We have } a \sim_R x_1 \text{ or } x_1 \sim_R a; \\ (b) \text{ We have } x_i \sim_R x_{i+1} \text{ or } x_{i+1} \sim_R x_i \\ \quad \text{for each } 1 \leq i \leq n-1; \\ (c) \text{ We have } b \sim_R x_n \text{ or } x_n \sim_R b; \end{array} \\ 2. \text{ We have } a = b. \end{array} \right\}. \end{aligned}$$

#### PROOF 10.5.2.1.3 ► PROOF OF CONSTRUCTION 10.5.2.1.2

From the universal properties of the reflexive, symmetric, and transitive closures of a relation (Definitions 10.2.2.1.1, 10.3.2.1.1 and 10.4.2.1.1), we see that it suffices to prove that:

00VJ

1. The symmetric closure of a reflexive relation is still reflexive.

00VK

2. The transitive closure of a symmetric relation is still symmetric.

which are both clear. 

00VL

#### PROPOSITION 10.5.2.1.4 ► PROPERTIES OF EQUIVALENCE RELATIONS

Let  $R$  be a relation on  $A$ .

00VM

1. *Adjointness.* We have an adjunction

$$((-)^{\text{eq}} \dashv \overline{\text{es}}) : \mathbf{Rel}(A, B) \begin{array}{c} \xrightarrow{(-)^{\text{eq}}} \\ \perp \\ \xleftarrow{\text{es}} \end{array} \mathbf{Rel}^{\text{eq}}(A, B),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{eq}}(R^{\text{eq}}, S) \cong \mathbf{Rel}(R, S),$$

natural in  $R \in \text{Obj}(\mathbf{Rel}^{\text{eq}}(A, B))$  and  $S \in \text{Obj}(\mathbf{Rel}(A, B))$ .

00VN

2. *The Equivalence Closure of an Equivalence Relation.* If  $R$  is an equivalence relation, then  $R^{\text{eq}} = R$ .

00VP

3. *Idempotency.* We have

$$(R^{\text{eq}})^{\text{eq}} = R^{\text{eq}}.$$

#### PROOF 10.5.2.1.5 ► PROOF OF PROPOSITION 10.5.2.1.4

##### Item 1: Adjointness

This is a rephrasing of the universal property of the equivalence closure of a relation, stated in [Definition 10.5.2.1.1](#).

##### Item 2: The Equivalence Closure of an Equivalence Relation

Clear.

##### Item 3: Idempotency

This follows from [Item 2](#). 

## 00VQ 10.6 Quotients by Equivalence Relations

### 00VR 10.6.1 Equivalence Classes

Let  $A$  be a set, let  $R$  be a relation on  $A$ , and let  $a \in A$ .

02B1

#### DEFINITION 10.6.1.1.1 ► EQUIVALENCE CLASSES

The **equivalence class associated to  $a$**  is the set  $[a]$  defined by

$$\begin{aligned} [a] &\stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\} \\ &= \{x \in X \mid a \sim_R x\}. \quad (\text{since } R \text{ is symmetric}) \end{aligned}$$

**02B2 10.6.2 Quotients of Sets by Equivalence Relations**

Let  $A$  be a set and let  $R$  be a relation on  $A$ .

**02B3**

**DEFINITION 10.6.2.1.1 ► QUOTIENTS OF SETS BY EQUIVALENCE RELATIONS**

The **quotient of  $X$  by  $R$**  is the set  $X/\sim_R$  defined by

$$X/\sim_R \stackrel{\text{def}}{=} \{[a] \in \mathcal{P}(X) \mid a \in X\}.$$

**00VV**

**REMARK 10.6.2.1.2 ► WHY USE “EQUIVALENCE” RELATIONS FOR QUOTIENT SETS**

The reason we define quotient sets for equivalence relations only is that each of the properties of being an equivalence relation—reflexivity, symmetry, and transitivity—ensures that the equivalences classes  $[a]$  of  $X$  under  $R$  are well-behaved:

- *Reflexivity.* If  $R$  is reflexive, then, for each  $a \in X$ , we have  $a \in [a]$ .
- *Symmetry.* The equivalence class  $[a]$  of an element  $a$  of  $X$  is defined by

$$[a] \stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\},$$

but we could equally well define

$$[a]' \stackrel{\text{def}}{=} \{x \in X \mid a \sim_R x\}$$

instead. This is not a problem when  $R$  is symmetric, as we then have  $[a] = [a]'$ .<sup>1</sup>

- *Transitivity.* If  $R$  is transitive, then  $[a]$  and  $[b]$  are disjoint iff  $a \not\sim_R b$ , and equal otherwise.

<sup>1</sup>When categorifying equivalence relations, one finds that  $[a]$  and  $[a]'$  correspond to presheaves and copresheaves; see ??.

**00VW**

**PROPOSITION 10.6.2.1.3 ► PROPERTIES OF QUOTIENT SETS**

Let  $f: X \rightarrow Y$  be a function and let  $R$  be a relation on  $X$ .

**02B4**

1. As a *Coequaliser*. We have an isomorphism of sets

$$X/\sim_R^{\text{eq}} \cong \text{CoEq}(R \hookrightarrow X \times X \xrightarrow{\text{pr}_2} X),$$

where  $\sim_R^{\text{eq}}$  is the equivalence relation generated by  $\sim_R$ .

02B5

2. As a Pushout. We have an isomorphism of sets<sup>1</sup>

$$\begin{array}{ccc} X/\sim_R^{\text{eq}} & \xleftarrow{\quad} & X \\ \uparrow \lrcorner & & \uparrow \\ X/\sim_R^{\text{eq}} \cong X \coprod_{\text{Eq}(\text{pr}_1, \text{pr}_2)} X, & & \\ & & \uparrow \\ & & X \leftarrow \text{Eq}(\text{pr}_1, \text{pr}_2). \end{array}$$

where  $\sim_R^{\text{eq}}$  is the equivalence relation generated by  $\sim_R$ .

02B6

3. The First Isomorphism Theorem for Sets. We have an isomorphism of sets<sup>2,3</sup>

$$X/\sim_{\text{Ker}(f)} \cong \text{Im}(f).$$

00W0

4. Descending Functions to Quotient Sets, I. Let  $R$  be an equivalence relation on  $X$ . The following conditions are equivalent:

02B7

(a) There exists a map

$$\bar{f}: X/\sim_R \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \nearrow \exists \pi & \\ X/\sim_R & & \end{array}$$

commute.

02B8

(b) We have  $R \subset \text{Ker}(f)$ .

02B9

(c) For each  $x, y \in X$ , if  $x \sim_R y$ , then  $f(x) = f(y)$ .

00W1

5. Descending Functions to Quotient Sets, II. Let  $R$  be an equivalence relation on  $X$ . If the conditions of Item 4 hold, then  $\bar{f}$  is the unique map making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \nearrow \exists! \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

00W2

6. *Descending Functions to Quotient Sets, III.* Let  $R$  be an equivalence relation on  $X$ . We have a bijection

$$\text{Hom}_{\text{Sets}}(X/\sim_R, Y) \cong \text{Hom}_{\text{Sets}}^R(X, Y),$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ , given by the assignment  $f \mapsto \bar{f}$  of **Items 4** and **5**, where  $\text{Hom}_{\text{Sets}}^R(X, Y)$  is the set defined by

$$\text{Hom}_{\text{Sets}}^R(X, Y) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(X, Y) \middle| \begin{array}{l} \text{for each } x, y \in X, \\ \text{if } x \sim_R y, \text{ then } \\ f(x) = f(y) \end{array} \right\}.$$

02BA

7. *Descending Functions to Quotient Sets, IV.* Let  $R$  be an equivalence relation on  $X$ . If the conditions of **Item 4** hold, then the following conditions are equivalent:

02BB

(a) The map  $\bar{f}$  is an injection.

02BC

(b) We have  $R = \text{Ker}(f)$ .

02BD

(c) For each  $x, y \in X$ , we have  $x \sim_R y$  iff  $f(x) = f(y)$ .

02BE

8. *Descending Functions to Quotient Sets, V.* Let  $R$  be an equivalence relation on  $X$ . If the conditions of **Item 4** hold, then the following conditions are equivalent:

02BF

(a) The map  $f: X \rightarrow Y$  is surjective.

02BG

(b) The map  $\bar{f}: X/\sim_R \rightarrow Y$  is surjective.

02BH

9. *Descending Functions to Quotient Sets, VI.* Let  $R$  be a relation on  $X$  and let  $\sim_R^{\text{eq}}$  be the equivalence relation associated to  $R$ . The following conditions are equivalent:

02BJ

(a) The map  $f$  satisfies the equivalent conditions of **Item 4**:

- There exists a map

$$\bar{f}: X/\sim_R^{\text{eq}} \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \swarrow \exists \bar{f} & \\ X/\sim_R^{\text{eq}} & & \end{array}$$

commute.

02BK

- For each  $x, y \in X$ , if  $x \sim_R^{\text{eq}} y$ , then  $f(x) = f(y)$ .

(b) For each  $x, y \in X$ , if  $x \sim_R y$ , then  $f(x) = f(y)$ .

<sup>1</sup>Dually, we also have an isomorphism of sets

$$\begin{array}{ccc} \text{Eq}(\text{pr}_1, \text{pr}_2) & \longrightarrow & X \\ \downarrow & & \downarrow \\ \text{Eq}(\text{pr}_1, \text{pr}_2) \cong X \times_{X/\sim_R^{\text{eq}}} X, & & \\ & \downarrow & \\ & X & \longrightarrow X/\sim_R^{\text{eq}}. \end{array}$$

<sup>2</sup>Further Terminology: The set  $X/\sim_{\text{Ker}(f)}$  is often called the **coimage of  $f$** , and denoted by  $\text{CoIm}(f)$ .

<sup>3</sup>In a sense this is a result relating the monad in **Rel** induced by  $f$  with the comonad in **Rel** induced by  $f$ , as the kernel and image

$$\begin{aligned} \text{Ker}(f) : X &\dashrightarrow X, \\ \text{Im}(f) &\subset Y \end{aligned}$$

of  $f$  are the underlying functors of (respectively) the induced monad and comonad of the adjunction

$$\left( \text{Gr}(f) \dashv f^{-1} \right) : A \begin{array}{c} \xrightarrow{\quad \perp \quad} \\[-1ex] \dashv \\[-1ex] \xleftarrow{f^{-1}} \end{array} B$$

of Item 4 of Proposition 8.2.2.1.2.

#### PROOF 10.6.2.1.4 ► PROOF OF PROPOSITION 10.6.2.1.3

Item 1: As a Coequaliser

Omitted.

Item 2: As a Pushout

Omitted.

Item 3: The First Isomorphism Theorem for Sets

Clear.

Item 4: Descending Functions to Quotient Sets, I

See [Pro25m].

Item 5: Descending Functions to Quotient Sets, II

See [Pro25z].

Item 6: Descending Functions to Quotient Sets, III

This follows from Items 5 and 6.

Item 7: Descending Functions to Quotient Sets, IV

See [Pro25l].

Item 8: Descending Functions to Quotient Sets, V

See [Pro25k].

Item 9: Descending Functions to Quotient Sets, VI

The implication Item 8a  $\implies$  Item 8b is clear.

Conversely, suppose that, for each  $x, y \in X$ , if  $x \sim_R y$ , then  $f(x) = f(y)$ . Spelling out the definition of the equivalence closure of  $R$ , we see that the condition  $x \sim_R^{\text{eq}} y$  unwinds to the following:

- (★) There exist  $(x_1, \dots, x_n) \in R^{\times n}$  satisfying at least one of the following conditions:

- The following conditions are satisfied:
  - \* We have  $x \sim_R x_1$  or  $x_1 \sim_R x$ ;
  - \* We have  $x_i \sim_R x_{i+1}$  or  $x_{i+1} \sim_R x_i$  for each  $1 \leq i \leq n-1$ ;
  - \* We have  $y \sim_R x_n$  or  $x_n \sim_R y$ ;
- We have  $x = y$ .

Now, if  $x = y$ , then  $f(x) = f(y)$  trivially; otherwise, we have

$$\begin{aligned} f(x) &= f(x_1), \\ f(x_1) &= f(x_2), \\ &\vdots \\ f(x_{n-1}) &= f(x_n), \\ f(x_n) &= f(y), \end{aligned}$$

and  $f(x) = f(y)$ , as we wanted to show. □

## Appendices

### 10.A Other Chapters

#### Preliminaries

1. Introduction
2. A Guide to the Literature

#### Sets

3. Sets

#### 4. Constructions With Sets

5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

**Relations**

- 8. Relations
- 9. Constructions With Relations
- 10. Conditions on Relations

**Categories**

- 11. Categories
- 12. Presheaves and the Yoneda Lemma

**Monoidal Categories**

- 13. Constructions With Monoidal Categories

**Bicategories**

- 14. Types of Morphisms in Bicategories

**Extra Part**

- 15. Notes

## **Part IV**

# **Categories**

# Chapter 11

## Categories

02H3 This chapter contains some elementary material about categories, functors, and natural transformations. Notably, we discuss and explore:

- 02BL 1. Categories ([Section 11.1](#)).
- 02BM 2. Examples of categories ([Section 11.2](#)).
- 02BN 3. The quadruple adjunction  $\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}$  between the category of categories and the category of sets ([Section 11.3](#)).
- 02BP 4. Groupoids, categories in which all morphisms admit inverses ([Section 11.4](#)).
- 02BQ 5. Functors ([Section 11.5](#)).
- 02BR 6. The conditions one may impose on functors in decreasing order of importance:
  - 02BS (a) [Section 11.6](#) introduces the foundationally important conditions one may impose on functors, such as faithfulness, conservativity, essential surjectivity, etc.
  - 02BT (b) [Section 11.7](#) introduces more conditions one may impose on functors that are still important but less omni-present than those of [Section 11.6](#), such as being dominant, being a monomorphism, being pseudomonic, etc.
  - 02BU (c) [Section 11.8](#) introduces some rather rare or uncommon conditions one may impose on functors that are nevertheless still useful to explicit record in this chapter.
- 02BV 7. Natural transformations ([Section 11.9](#)).
- 02BW 8. The various categorical and 2-categorical structures formed by categories, functors, and natural transformations ([Section 11.10](#)).

This chapter is under active revision. TODO:

- Fix categories having an underlying set of objects by having them have an underlying setoid of objects (not necessarily by definition, as that'll likely be bothersome; at least [Section 11.3](#) should be fixed and several remarks should be added at several points). Related: [Warning 11.3.1.1.3](#)

## Contents

---

<b>11.1 Categories.....</b>	<b>731</b>
11.1.1 Foundations .....	731
11.1.2 Subcategories .....	733
11.1.3 Skeletons of Categories.....	734
11.1.4 Precomposition and Postcomposition .....	735
<b>11.2 Examples of Categories.....</b>	<b>739</b>
11.2.1 The Empty Category .....	739
11.2.2 The Punctual Category .....	740
11.2.3 Monoids as One-Object Categories .....	740
11.2.4 Ordinal Categories.....	741
11.2.5 The Walking Arrow .....	742
11.2.6 More Examples of Categories .....	743
11.2.7 Posetal Categories .....	743
<b>11.3 The Quadruple Adjunction With Sets.....</b>	<b>746</b>
11.3.1 Statement.....	746
11.3.2 Connected Components and Connected Cate- gories.....	747
11.3.3 Discrete Categories .....	750
11.3.4 Indiscrete Categories.....	752
<b>11.4 Groupoids.....</b>	<b>754</b>
11.4.1 Isomorphisms .....	754
11.4.2 Groupoids.....	754
11.4.3 The Groupoid Completion of a Category.....	755
11.4.4 The Core of a Category .....	759
<b>11.5 Functors .....</b>	<b>762</b>
11.5.1 Foundations .....	762
11.5.2 Contravariant Functors .....	766
11.5.3 Forgetful Functors .....	768
11.5.4 The Natural Transformation Associated to a Func- tor.....	770
<b>11.6 Conditions on Functors.....</b>	<b>772</b>
11.6.1 Faithful Functors .....	772

---

11.6.2 Full Functors .....	774
11.6.3 Fully Faithful Functors.....	781
11.6.4 Conservative Functors .....	785
11.6.5 Essentially Injective Functors .....	787
11.6.6 Essentially Surjective Functors .....	788
11.6.7 Equivalences of Categories .....	789
11.6.8 Isomorphisms of Categories .....	791
<b>11.7 More Conditions on Functors.....</b>	<b>793</b>
11.7.1 Dominant Functors.....	793
11.7.2 Monomorphisms of Categories .....	794
11.7.3 Epimorphisms of Categories.....	795
11.7.4 Pseudomonadic Functors.....	797
11.7.5 Pseudoepic Functors .....	800
<b>11.8 Even More Conditions on Functors .....</b>	<b>803</b>
11.8.1 Injective on Objects Functors .....	803
11.8.2 Surjective on Objects Functors .....	804
11.8.3 Bijective on Objects Functors .....	804
11.8.4 Functors Representably Faithful on Cores .....	804
11.8.5 Functors Representably Full on Cores .....	805
11.8.6 Functors Representably Fully Faithful on Cores...	806
11.8.7 Functors Corepresentably Faithful on Cores .....	807
11.8.8 Functors Corepresentably Full on Cores .....	808
11.8.9 Functors Corepresentably Fully Faithful on Cores	
809	
<b>11.9 Natural Transformations .....</b>	<b>811</b>
11.9.1 Transformations.....	811
11.9.2 Natural Transformations .....	812
11.9.3 Examples of Natural Transformations .....	813
11.9.4 Vertical Composition of Natural Transformations.	815
11.9.5 Horizontal Composition of Natural Transforma-	
tions .....	819
11.9.6 Properties of Natural Transformations.....	824
11.9.7 Natural Isomorphisms .....	826
<b>11.10 Categories of Categories .....</b>	<b>828</b>
11.10.1 Functor Categories.....	828
11.10.2 The Category of Categories and Functors.....	832
11.10.3 The 2-Category of Categories, Functors, and Nat-	
ural Transformations .....	833
11.10.4 The Category of Groupoids.....	834
11.10.5 The 2-Category of Groupoids .....	834
<b>11.A Other Chapters .....</b>	<b>835</b>

---

**00W9 11.1 Categories**

**00WA 11.1.1 Foundations**

01CZ

DEFINITION 11.1.1.1 ► CATEGORIES

A **category**  $(C, \circ^C, \mathbb{1}^C)$  consists of:

- *Objects.* A class  $\text{Obj}(C)$  of **objects**.
- *Morphisms.* For each  $A, B \in \text{Obj}(C)$ , a class  $\text{Hom}_C(A, B)$ , called the **class of morphisms of  $C$  from  $A$  to  $B$** .
- *Identities.* For each  $A \in \text{Obj}(C)$ , a map of sets

$$\mathbb{1}_A^C : \text{pt} \rightarrow \text{Hom}_C(A, A),$$

called the **unit map of  $C$  at  $A$** , determining a morphism

$$\text{id}_A : A \rightarrow A$$

of  $C$ , called the **identity morphism of  $A$** .

- *Composition.* For each  $A, B, C \in \text{Obj}(C)$ , a map of sets

$$\circ_{A,B,C}^C : \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) \rightarrow \text{Hom}_C(A, C),$$

called the **composition map of  $C$  at  $(A, B, C)$** .

such that the following conditions are satisfied:

1. *Associativity.* The diagram

$$\begin{array}{ccc}
 & \text{Hom}_C(C, D) \times (\text{Hom}_C(B, C) \times \text{Hom}_C(A, B)) & \\
 & \swarrow \alpha_{\text{Hom}_C(C,D), \text{Hom}_C(B,C), \text{Hom}_C(A,B)}^{\text{Sets}} \quad \searrow \text{id}_{\text{Hom}_C(C,D)} \times \circ_{A,B,C}^C & \\
 (\text{Hom}_C(C, D) \times \text{Hom}_C(B, C)) \times \text{Hom}_C(A, B) & & \text{Hom}_C(C, D) \times \text{Hom}_C(A, C) \\
 & \downarrow \circ_{B,C,D}^C \times \text{id}_{\text{Hom}_C(A,B)} & \downarrow \circ_{A,C,D}^C \\
 & \text{Hom}_C(B, D) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,D}^C} \text{Hom}_C(A, D)
 \end{array}$$

commutes, i.e. for each composable triple  $(f, g, h)$  of morphisms of  $C$ , we have

$$(f \circ g) \circ h = f \circ (g \circ h).$$

2. *Left Unitality.* The diagram

$$\begin{array}{ccc}
 \text{pt} \times \text{Hom}_C(A, B) & & \\
 \downarrow \text{id}_B \times \text{id}_{\text{Hom}_C(A, B)} & \nearrow \lambda_{\text{Hom}_C(A, B)}^{\text{Sets}} & \\
 \text{Hom}_C(B, B) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A, B, B}^C} & \text{Hom}_C(A, B)
 \end{array}$$

commutes, i.e. for each morphism  $f: A \rightarrow B$  of  $C$ , we have

$$\text{id}_B \circ f = f.$$

3. *Right Unitality.* The diagram

$$\begin{array}{ccc}
 \text{Hom}_C(A, B) \times \text{pt} & & \\
 \downarrow \text{id}_{\text{Hom}_C(A, B)} \times \text{id}_A^C & \nearrow \rho_{\text{Hom}_C(A, B)}^{\text{Sets}} & \\
 \text{Hom}_C(A, B) \times \text{Hom}_C(A, A) & \xrightarrow{\circ_{A, A, B}^C} & \text{Hom}_C(A, B)
 \end{array}$$

commutes, i.e. for each morphism  $f: A \rightarrow B$  of  $C$ , we have

$$f \circ \text{id}_A = f.$$

00WC

#### NOTATION 11.1.1.2 ► FURTHER NOTATION FOR MORPHISMS IN CATEGORIES

Let  $C$  be a category.

00WD

1. We also write  $C(A, B)$  for  $\text{Hom}_C(A, B)$ .

00WE

2. We write  $\text{Mor}(C)$  for the class of all morphisms of  $C$ .

00WF

#### DEFINITION 11.1.1.3 ► SIZE CONDITIONS ON CATEGORIES

Let  $\kappa$  be a regular cardinal. A category  $C$  is

00WG

1. **Locally small** if, for each  $A, B \in \text{Obj}(C)$ , the class  $\text{Hom}_C(A, B)$  is a set.

00WH

2. **Locally essentially small** if, for each  $A, B \in \text{Obj}(C)$ , the class

$$\text{Hom}_C(A, B)/\{\text{isomorphisms}\}$$

is a set.

00WJ

3. **Small** if  $C$  is locally small and  $\text{Obj}(C)$  is a set.

00WK

4.  **$\kappa$ -Small** if  $C$  is locally small,  $\text{Obj}(C)$  is a set, and we have  $\#\text{Obj}(C) < \kappa$ .

00XC **11.1.2 Subcategories**

Let  $C$  be a category.

00XD

**DEFINITION 11.1.2.1.1 ► SUBCATEGORIES**

A **subcategory** of  $C$  is a category  $\mathcal{A}$  satisfying the following conditions:

1. *Objects*. We have  $\text{Obj}(\mathcal{A}) \subset \text{Obj}(C)$ .
2. *Morphisms*. For each  $A, B \in \text{Obj}(\mathcal{A})$ , we have

$$\text{Hom}_{\mathcal{A}}(A, B) \subset \text{Hom}_C(A, B).$$

3. *Identities*. For each  $A \in \text{Obj}(\mathcal{A})$ , we have

$$\mathbb{1}_A^{\mathcal{A}} = \mathbb{1}_A^C.$$

4. *Composition*. For each  $A, B, C \in \text{Obj}(\mathcal{A})$ , we have

$$\circ_{A,B,C}^{\mathcal{A}} = \circ_{A,B,C}^C.$$

00XE

**DEFINITION 11.1.2.1.2 ► FULL SUBCATEGORIES**

A subcategory  $\mathcal{A}$  of  $C$  is **full** if the canonical inclusion functor  $\mathcal{A} \rightarrow C$  is full, i.e. if, for each  $A, B \in \text{Obj}(\mathcal{A})$ , the inclusion

$$\iota_{A,B}: \text{Hom}_{\mathcal{A}}(A, B) \hookrightarrow \text{Hom}_C(A, B)$$

is surjective (and thus bijective).

00XF

**DEFINITION 11.1.2.1.3 ► STRICTLY FULL SUBCATEGORIES**

A subcategory  $\mathcal{A}$  of a category  $C$  is **strictly full** if it satisfies the following conditions:

1. *Fullness*. The subcategory  $\mathcal{A}$  is full.
2. *Closedness Under Isomorphisms*. The class  $\text{Obj}(\mathcal{A})$  is closed under isomorphisms.<sup>1</sup>

<sup>1</sup>That is, given  $A \in \text{Obj}(\mathcal{A})$  and  $C \in \text{Obj}(C)$ , if  $C \cong A$ , then  $C \in \text{Obj}(\mathcal{A})$ .

00XG

**DEFINITION 11.1.2.1.4 ► WIDE SUBCATEGORIES**

A subcategory  $\mathcal{A}$  of  $C$  is **wide**<sup>1</sup> if  $\text{Obj}(\mathcal{A}) = \text{Obj}(C)$ .

<sup>1</sup>Further Terminology: Also called **lluf**.

00XH

**11.1.3 Skeletons of Categories**

00XJ

**DEFINITION 11.1.3.1.1 ► SKELETONS OF CATEGORIES**

A<sup>1</sup> **skeleton** of a category  $C$  is a full subcategory  $\text{Sk}(C)$  with one object from each isomorphism class of objects of  $C$ .

<sup>1</sup>Due to Item 3 of Proposition 11.1.3.1.3, which states that any two skeletons of a category are equivalent, we often refer to any such full subcategory  $\text{Sk}(C)$  of  $C$  as *the* skeleton of  $C$ .

00XK

**DEFINITION 11.1.3.1.2 ► SKELETAL CATEGORIES**

A category  $C$  is **skeletal** if  $C \cong \text{Sk}(C)$ .<sup>1</sup>

<sup>1</sup>That is,  $C$  is **skeletal** if isomorphic objects of  $C$  are equal.

00XL

**PROPOSITION 11.1.3.1.3 ► PROPERTIES OF SKELETONS OF CATEGORIES**

Let  $C$  be a category.

00XM

1. *Existence.* Assuming the axiom of choice,  $\text{Sk}(C)$  always exists.

00XN

2. *Pseudofunctoriality.* The assignment  $C \mapsto \text{Sk}(C)$  defines a pseudofunctor

$$\text{Sk}: \text{Cats}_2 \rightarrow \text{Cats}_2.$$

00XP

3. *Uniqueness Up to Equivalence.* Any two skeletons of  $C$  are equivalent.

00XQ

4. *Inclusions of Skeletons Are Equivalences.* The inclusion

$$\iota_C: \text{Sk}(C) \hookrightarrow C$$

of a skeleton of  $C$  into  $C$  is an equivalence of categories.

**PROOF 11.1.3.1.4 ► PROOF OF PROPOSITION 11.1.3.1.3**

Item 1: Existence

See [nLab23, Section “Existence of Skeletons of Categories”].

Item 2: Pseudofunctoriality

See [nLab23, Section “Skeletons as an Endo-Pseudofunctor on  $\mathfrak{Cat}$ ”].

Item 3: Uniqueness Up to Equivalence

Omitted.

Item 4: Inclusions of Skeletons Are Equivalences

Omitted. 

#### 00XR 11.1.4 Precomposition and Postcomposition

Let  $C$  be a category and let  $A, B, C, X \in \text{Obj}(C)$ .

00XS

##### DEFINITION 11.1.4.1.1 ► PRECOMPOSITION AND POSTCOMPOSITION FUNCTIONS

Let  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be morphisms of  $C$ .

00XT

1. The **precomposition function associated to  $f$**  is the function

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

defined by

$$f^*(\phi) \stackrel{\text{def}}{=} \phi \circ f$$

for each  $\phi \in \text{Hom}_C(B, X)$ .

00XU

2. The **postcomposition function associated to  $g$**  is the function

$$g_*: \text{Hom}_C(X, B) \rightarrow \text{Hom}_C(X, C)$$

defined by

$$g_*(\phi) \stackrel{\text{def}}{=} g \circ \phi$$

for each  $\phi \in \text{Hom}_C(X, B)$ .

00XV

##### PROPOSITION 11.1.4.1.2 ► PROPERTIES OF PRE/POSTCOMPOSITION

Let  $A, B, C, D, X \in \text{Obj}(C)$ .

00XW

1. *Interaction Between Precomposition and Postcomposition.* Let  $f: A \rightarrow B$  and  $g: X \rightarrow Y$  be morphisms of  $C$ . We have

$$\begin{array}{ccc} \text{Hom}_C(B, X) & \xrightarrow{g^*} & \text{Hom}_C(B, Y) \\ g_* \circ f^* = f^* \circ g_*, & \downarrow f^* & \downarrow f^* \\ \text{Hom}_C(A, X) & \xrightarrow{g_*} & \text{Hom}_C(A, Y). \end{array}$$

00XX

2. *Interaction With Composition I.* Let  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be morphisms of  $C$ . We have

$$\begin{array}{ccc} \text{Hom}_C(X, A) & \xrightarrow{f_*} & \text{Hom}_C(X, B) \\ (g \circ f)_* = g_* \circ f_*, & \searrow (g \circ f)_* & \downarrow g_* \\ & & \text{Hom}_C(X, C), \end{array}$$
  

$$\begin{array}{ccc} \text{Hom}_C(C, X) & \xrightarrow{g^*} & \text{Hom}_C(B, X) \\ (g \circ f)^* = f^* \circ g^*, & \searrow (g \circ f)^* & \downarrow f^* \\ & & \text{Hom}_C(A, X). \end{array}$$

00XY

3. *Interaction With Composition II.* Let  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be morphisms of  $C$ . We have

$$\begin{array}{ccc} \text{pt} & \xrightarrow{[f]} & \text{Hom}_C(A, B) \\ & \searrow [g \circ f] & \downarrow g_* \\ & & \text{Hom}_C(A, C) \end{array} \quad \begin{array}{c} [g \circ f] = g_* \circ [f], \\ [g \circ f] = f^* \circ [g], \end{array} \quad \begin{array}{ccc} \text{pt} & \xrightarrow{[g]} & \text{Hom}_C(B, C) \\ & \searrow [g \circ f] & \downarrow f^* \\ & & \text{Hom}_C(A, C). \end{array}$$

00XZ

4. *Interaction With Composition III.* Let  $f: X \rightarrow A$  and  $g: C \rightarrow D$  be morphisms of  $C$ . We have

$$\begin{array}{ccc} \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ f^* \circ \circ_{A,B,C}^C = \circ_{X,B,C}^C \circ (\text{id} \times f^*), & \downarrow \text{id} \times f^* & \downarrow f^* \\ \text{Hom}_C(B, C) \times \text{Hom}_C(X, B) & \xrightarrow{\circ_{X,B,C}^C} & \text{Hom}_C(X, C), \end{array}$$
  

$$\begin{array}{ccc} \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ g_* \circ \circ_{A,B,C}^C = \circ_{A,B,D}^C \circ (g_* \times \text{id}), & \downarrow g_* \times \text{id} & \downarrow g^* \\ \text{Hom}_C(B, D) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,D}^C} & \text{Hom}_C(A, D). \end{array}$$

00Y0

5. *Interaction With Identities.* We have

$$\begin{aligned} \text{id}_A^* &= \text{id}_{\text{Hom}_C(A, B)}, \\ (\text{id}_B)_* &= \text{id}_{\text{Hom}_C(A, B)}. \end{aligned}$$

**PROOF 11.1.4.1.3 ▶ PROOF OF PROPOSITION 11.1.4.1.2****Item 1: Interaction Between Precomposition and Postcomposition**

For each  $\phi \in \text{Hom}_C(B, X)$ , we have

$$\begin{aligned} [g_* \circ f^*](\phi) &= g_*(\phi \circ f) \\ &= g \circ (\phi \circ f) \\ &= (g \circ \phi) \circ f \\ &= f^*(g \circ \phi) \\ &= [f^* \circ g_*](\phi). \end{aligned}$$

Thus  $g_* \circ f^* = f^* \circ g_*$ .

**Item 2: Interaction With Composition I**

$$(g \circ f)_* = g_* \circ f_*$$

For each  $\phi \in \text{Hom}_C(X, A)$ , we have

$$\begin{aligned} (g \circ f)_*(\phi) &= (g \circ f) \circ \phi \\ &= g \circ (f \circ \phi) \\ &= g \circ f_*(\phi) \\ &= g_*(f_*(\phi)) \\ &= [g_* \circ f_*](\phi). \end{aligned}$$

Thus  $(g \circ f)_* = g_* \circ f_*$ .

$$(g \circ f)^* = g^* \circ f^*$$

For each  $\phi \in \text{Hom}_C(C, X)$ , we have

$$\begin{aligned} (g \circ f)^*(\phi) &= \phi \circ (g \circ f) \\ &= (\phi \circ g) \circ f \\ &= (g^*(\phi)) \circ f \\ &= f^*(g^*(\phi)) \\ &= [f^* \circ g^*](\phi). \end{aligned}$$

Thus  $(g \circ f)^* = g^* \circ f^*$ .

**Item 3: Interaction With Composition II**

It suffices to show the equalities of the maps on  $\star \in \text{pt}$ . We have

$$\begin{aligned} [g \circ f](\star) &= g \circ f \\ &= g_*(f) \\ &= g_*([f](\star)) \\ &= (g_* \circ [f])(\star), \end{aligned}$$

and

$$\begin{aligned} [g \circ f](\star) &= g \circ f \\ &= f^*(g) \\ &= f^*([g](\star)) \\ &= (f^* \circ [g])(\star). \end{aligned}$$

Thus  $[g \circ f] = g_* \circ [f]$  and  $[g \circ f] = f^* \circ [g]$ .

**Item 4: Interaction With Composition III**

$$f^* \circ \circ_{A,B,C}^C = \circ_{X,B,C}^C \circ (\text{id} \times f^*)$$

For each  $(\psi, \phi) \in \text{Hom}_C(B, C) \times \text{Hom}_C(A, B)$ , we have

$$\begin{aligned} [f^* \circ \circ_{A,B,C}^C](\psi, \phi) &= f^*(\psi \circ \phi) \\ &= (\psi \circ \phi) \circ f \\ &= \psi \circ (\phi \circ f) \\ &= \circ_{X,B,C}^C(\psi, \phi \circ f) \\ &= \circ_{X,B,C}^C(\psi, f^*(\phi)) \\ &= [\circ_{X,B,C}^C \circ (\text{id} \times f^*)](\psi, \phi). \end{aligned}$$

Thus  $f^* \circ \circ_{A,B,C}^C = \circ_{X,B,C}^C \circ (\text{id} \times f^*)$ .

$$g_* \circ \circ_{A,B,C}^C = \circ_{A,B,D}^C \circ (g_* \times \text{id})$$

For each  $(\psi, \phi) \in \text{Hom}_C(B, C) \times \text{Hom}_C(A, B)$ , we have

$$\begin{aligned} [g_* \circ \circ_{A,B,C}^C](\psi, \phi) &= g_*(\psi \circ \phi) \\ &= g \circ (\psi \circ \phi) \\ &= (g \circ \psi) \circ \phi \end{aligned}$$

$$\begin{aligned}
 &= \circ_{A,B,D}^C(g \circ \psi, \phi) \\
 &= \circ_{A,B,D}^C(g_*(\psi), \phi) \\
 &= [\circ_{A,B,D}^C \circ (g_* \times \text{id})](\psi, \phi).
 \end{aligned}$$

Thus  $g_* \circ \circ_{A,B,C}^C = \circ_{A,B,D}^C \circ (g_* \times \text{id})$ .

#### Item 5: Interaction With Identities

We have

$$\begin{aligned}
 \text{id}_A^*(\phi) &= \phi \circ \text{id}_A \\
 &= \phi \\
 &= \text{id}_{\text{Hom}_C(A,B)}(\phi)
 \end{aligned}$$

and

$$\begin{aligned}
 (\text{id}_B)_*(\phi) &= \text{id}_B \circ \phi \\
 &= \phi \\
 &= \text{id}_{\text{Hom}_C(A,B)}(\phi)
 \end{aligned}$$

for each  $\phi \in \text{Hom}_C(A, B)$ . ■

## 00WL 11.2 Examples of Categories

### 01TT 11.2.1 The Empty Category

00WP

#### EXAMPLE 11.2.1.1.1 ► THE EMPTY CATEGORY

The **empty category** is the category  $\emptyset_{\text{cat}}$  where

- *Objects.* We have

$$\text{Obj}(\emptyset_{\text{cat}}) \stackrel{\text{def}}{=} \emptyset.$$

- *Morphisms.* We have

$$\text{Mor}(\emptyset_{\text{cat}}) \stackrel{\text{def}}{=} \emptyset.$$

- *Identities and Composition.* Having no objects,  $\emptyset_{\text{cat}}$  has no unit nor composition maps.

**01TU 11.2.2 The Punctual Category**

00WM

**EXAMPLE 11.2.2.1.1 ► THE PUNCTUAL CATEGORY**

The **punctual category**<sup>1</sup> is the category pt where

- *Objects.* We have

$$\text{Obj}(\text{pt}) \stackrel{\text{def}}{=} \{\star\}.$$

- *Morphisms.* The unique Hom-set of pt is defined by

$$\text{Hom}_{\text{pt}}(\star, \star) \stackrel{\text{def}}{=} \{\text{id}_\star\}.$$

- *Identities.* The unit map

$$\text{id}_\star^{\text{pt}} : \text{pt} \rightarrow \text{Hom}_{\text{pt}}(\star, \star)$$

of pt at  $\star$  is defined by

$$\text{id}_\star^{\text{pt}} \stackrel{\text{def}}{=} \text{id}_\star.$$

- *Composition.* The composition map

$$\circ_{\star, \star, \star}^{\text{pt}} : \text{Hom}_{\text{pt}}(\star, \star) \times \text{Hom}_{\text{pt}}(\star, \star) \rightarrow \text{Hom}_{\text{pt}}(\star, \star)$$

of pt at  $(\star, \star, \star)$  is given by the bijection  $\text{pt} \times \text{pt} \cong \text{pt}$ .

<sup>1</sup>Further Terminology: Also called the **singleton category**.

**01TV 11.2.3 Monoids as One-Object Categories**

00WN

**EXAMPLE 11.2.3.1.1 ► MONOIDS AS ONE-OBJECT CATEGORIES**

We have an isomorphism of categories<sup>1</sup>

$$\begin{array}{ccc} \text{Mon} & \longrightarrow & \text{Cats} \\ \downarrow & \lrcorner & \downarrow \text{Obj} \\ \text{Mon} \cong \text{pt} \times_{\text{Sets}} \text{Cats}, & & \\ \downarrow & & \downarrow \\ \text{pt} & \xrightarrow{[\text{pt}]} & \text{Sets} \end{array}$$

via the delooping functor  $B : \text{Mon} \rightarrow \text{Cats}$  of ?? of ??, exhibiting monoids as exactly those categories having a single object.

<sup>1</sup>This can be enhanced to an isomorphism of 2-categories

$$\begin{array}{ccc}
 \text{Mon}_{2\text{disc}} & \longrightarrow & \text{Cats}_{2,*} \\
 \downarrow & \lrcorner & \downarrow \text{Obj} \\
 \text{pt}_{\text{bi}} \times_{\text{Sets}_{2\text{disc}}} \text{Cats}_{2,*}, & & \\
 \downarrow & & \downarrow \\
 \text{pt}_{\text{bi}} & \xrightarrow{[\text{pt}]} & \text{Sets}_{2\text{disc}}
 \end{array}$$

between the discrete 2-category  $\text{Mon}_{2\text{disc}}$  on  $\text{Mon}$  and the 2-category of pointed categories with one object.

**PROOF 11.2.3.1.2 ▶ PROOF OF EXAMPLE 11.2.3.1.1**

Omitted. 

## 01TW 11.2.4 Ordinal Categories

00WQ

**EXAMPLE 11.2.4.1.1 ▶ ORDINAL CATEGORIES**

The  $n$ th ordinal category is the category  $\mathbb{n}$  where<sup>1</sup>

- *Objects.* We have

$$\text{Obj}(\mathbb{n}) \stackrel{\text{def}}{=} \{[0], \dots, [n]\}.$$

- *Morphisms.* For each  $[i], [j] \in \text{Obj}(\mathbb{n})$ , we have

$$\text{Hom}_{\mathbb{n}}([i], [j]) \stackrel{\text{def}}{=} \begin{cases} \{\text{id}_{[i]}\} & \text{if } [i] = [j], \\ \{[i] \rightarrow [j]\} & \text{if } [j] < [i], \\ \emptyset & \text{if } [j] > [i]. \end{cases}$$

- *Identities.* For each  $[i] \in \text{Obj}(\mathbb{n})$ , the unit map

$$\mathbb{1}_{[i]}^{\mathbb{n}} : \text{pt} \rightarrow \text{Hom}_{\mathbb{n}}([i], [i])$$

of  $\mathbb{n}$  at  $[i]$  is defined by

$$\text{id}_{[i]}^{\mathbb{n}} \stackrel{\text{def}}{=} \text{id}_{[i]}.$$

- *Composition.* For each  $[i], [j], [k] \in \text{Obj}(\mathbb{n})$ , the composition map

$$\circ_{[i], [j], [k]}^{\mathbb{n}} : \text{Hom}_{\mathbb{n}}([j], [k]) \times \text{Hom}_{\mathbb{n}}([i], [j]) \rightarrow \text{Hom}_{\mathbb{n}}([i], [k])$$

of  $\mathbb{n}$  at  $([i], [j], [k])$  is defined by

$$\begin{aligned}
 \text{id}_{[i]} \circ \text{id}_{[i]} &= \text{id}_{[i]}, \\
 ([j] \rightarrow [k]) \circ ([i] \rightarrow [j]) &= ([i] \rightarrow [k]). 
 \end{aligned}$$

<sup>1</sup>In other words,  $\mathbb{m}$  is the category associated to the poset

$$[0] \rightarrow [1] \rightarrow \cdots \rightarrow [n-1] \rightarrow [n].$$

The category  $\mathbb{m}$  for  $n \geq 2$  may also be defined in terms of  $\mathbb{0}$  and joins (??): we have isomorphisms of categories

$$\begin{aligned} \mathbb{1} &\cong \mathbb{0} \star \mathbb{0}, \\ \mathbb{2} &\cong \mathbb{1} \star \mathbb{0} \\ &\cong (\mathbb{0} \star \mathbb{0}) \star \mathbb{0}, \\ \mathbb{3} &\cong \mathbb{2} \star \mathbb{0} \\ &\cong (\mathbb{1} \star \mathbb{0}) \star \mathbb{0} \\ &\cong ((\mathbb{0} \star \mathbb{0}) \star \mathbb{0}) \star \mathbb{0}, \\ \mathbb{4} &\cong \mathbb{3} \star \mathbb{0} \\ &\cong (\mathbb{2} \star \mathbb{0}) \star \mathbb{0} \\ &\cong ((\mathbb{1} \star \mathbb{0}) \star \mathbb{0}) \star \mathbb{0} \\ &\cong (((\mathbb{0} \star \mathbb{0}) \star \mathbb{0}) \star \mathbb{0}) \star \mathbb{0}, \end{aligned}$$

and so on.

### 01TX 11.2.5 The Walking Arrow

01TY

#### DEFINITION 11.2.5.1.1 ► THE WALKING ARROW

The **walking arrow** is the category  $\mathbb{1}$  defined as the first ordinal category.

01TZ

#### REMARK 11.2.5.1.2 ► UNWINDING DEFINITION 11.2.5.1.1

In detail, the walking arrow is the category  $\mathbb{1}$  where:

- *Objects.* We have  $\text{Obj}(\mathbb{1}) = \{0, 1\}$ .
- *Morphisms.* We have

$$\begin{aligned} \text{Hom}_{\mathbb{1}}(0, 0) &= \{\text{id}_0\}, \\ \text{Hom}_{\mathbb{1}}(1, 1) &= \{\text{id}_1\}, \\ \text{Hom}_{\mathbb{1}}(0, 1) &= \{f_{01}\}, \\ \text{Hom}_{\mathbb{1}}(1, 0) &= \emptyset. \end{aligned}$$

- *Identities and Composition.* The identities and composition of  $\mathbb{1}$  are completely determined by the unitality and associativity axioms for  $\mathbb{1}$ .

### 01U0 11.2.6 More Examples of Categories

00WR

#### EXAMPLE 11.2.6.1.1 ► MORE EXAMPLES OF CATEGORIES

Here we list some of the other categories appearing throughout this work.

00WS

1. The category  $\text{Sets}_*$  of pointed sets of [Definition 6.1.3.1.1](#).

00WT

2. The category  $\text{Rel}$  of sets and relations of [Definition 8.3.2.1.1](#).

00WU

3. The category  $\text{Span}(A, B)$  of spans from a set  $A$  to a set  $B$  of [??](#).

00WV

4. The category  $\text{ISets}(K)$  of  $K$ -indexed sets of [??](#).

00WW

5. The category  $\text{ISets}$  of indexed sets of [??](#).

00WX

6. The category  $\text{FibSets}(K)$  of  $K$ -fibred sets of [??](#).

00WY

7. The category  $\text{FibSets}$  of fibred sets of [??](#).

00WZ

8. Categories of functors  $\text{Fun}(C, \mathcal{D})$  as in [Definition 11.10.1.1.1](#).

00X0

9. The category of categories  $\text{Cats}$  of [Definition 11.10.2.1.1](#).

00X1

10. The category of groupoids  $\text{Grpd}$  of [Definition 11.10.4.1.1](#).

### 00X2 11.2.7 Posetal Categories

00X3

#### DEFINITION 11.2.7.1.1 ► POSETAL CATEGORIES

Let  $(X, \preceq_X)$  be a poset.

00X4

1. The **posetal category associated to**  $(X, \preceq_X)$  is the category  $X_{\text{pos}}$  where

- *Objects.* We have

$$\text{Obj}(X_{\text{pos}}) \stackrel{\text{def}}{=} X.$$

- *Morphisms.* For each  $a, b \in \text{Obj}(X_{\text{pos}})$ , we have

$$\text{Hom}_{X_{\text{pos}}}(a, b) \stackrel{\text{def}}{=} \begin{cases} \text{pt} & \text{if } a \preceq_X b, \\ \emptyset & \text{otherwise.} \end{cases}$$

- *Identities.* For each  $a \in \text{Obj}(X_{\text{pos}})$ , the unit map

$$\mathbb{1}_a^{X_{\text{pos}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{pos}}}(a, a)$$

of  $X_{\text{pos}}$  at  $a$  is given by the identity map.

- *Composition.* For each  $a, b, c \in \text{Obj}(X_{\text{pos}})$ , the composition map

$$\circ^{X_{\text{pos}}}_{a,b,c} : \text{Hom}_{X_{\text{pos}}}(b, c) \times \text{Hom}_{X_{\text{pos}}}(a, b) \rightarrow \text{Hom}_{X_{\text{pos}}}(a, c)$$

of  $X_{\text{pos}}$  at  $(a, b, c)$  is defined as either the inclusion  $\emptyset \hookrightarrow \text{pt}$  or the identity map of  $\text{pt}$ , depending on whether we have  $a \preceq_X b$ ,  $b \preceq_X c$ , and  $a \preceq_X c$ .

- 00X5** 2. A category  $C$  is **posetal**<sup>1</sup> if  $C$  is equivalent to  $X_{\text{pos}}$  for some poset  $(X, \preceq_X)$ .

<sup>1</sup>Further Terminology: Also called a **thin** category or a **(0, 1)-category**.

**00X6**

#### PROPOSITION 11.2.7.1.2 ► PROPERTIES OF POSETAL CATEGORIES

Let  $(X, \preceq_X)$  be a poset and let  $C$  be a category.

**00X7**

1. *Functoriality.* The assignment  $(X, \preceq_X) \mapsto X_{\text{pos}}$  defines a functor

$$(-)_{\text{pos}} : \text{Pos} \rightarrow \text{Cats}.$$

where:

- *Action on Objects.* For each  $X \in \text{Obj}(\text{Pos})$ , we have

$$[(-)_{\text{pos}}](X) \stackrel{\text{def}}{=} X_{\text{pos}},$$

where  $X_{\text{pos}}$  is the category of of Item 1 of Definition 11.2.7.1.1.

- *Action on Morphisms.* For each morphism of posets  $f: X \rightarrow Y$  in  $\text{Pos}$ , the image

$$f_{\text{pos}}: X_{\text{pos}} \rightarrow Y_{\text{pos}}$$

of  $f$  by  $(-)_{\text{pos}}$  is the functor defined as follows:

- *The Action of  $f_{\text{pos}}$  on Objects.* For each  $x \in \text{Obj}(X_{\text{pos}})$ , we have

$$f_{\text{pos}}(x) \stackrel{\text{def}}{=} f(x).$$

- *The Action of  $f_{\text{pos}}$  on Morphisms.* For each  $x, y \in \text{Obj}(X_{\text{pos}})$ , the action

$$f_{\text{pos}|x,y}: \text{Hom}_{X_{\text{pos}}}(x, y) \rightarrow \text{Hom}_{Y_{\text{pos}}}(f(x), f(y))$$

of  $f$  at  $(x, y)$  is given by

$$f_{\text{pos}|x,y}(\text{pt}_{\text{Hom}_{X_{\text{pos}}}(x, y)}) \stackrel{\text{def}}{=} \text{pt}_{\text{Hom}_{Y_{\text{pos}}}(f(x), f(y))}$$

if  $x \preceq_X y$  or, otherwise, by the inclusion of the empty set into  $\text{Hom}_{Y_{\text{pos}}}(f(x), f(y))$ .

00X8

2. *Fully Faithfulness.* The functor  $(-)_\text{pos}$  of [Item 1](#) is fully faithful.

00X9

3. *Characterisations.* The following conditions are equivalent:

00XA

(a) The category  $C$  is posetal.

00XB

(b) For each  $A, B \in \text{Obj}(C)$  and each  $f, g \in \text{Hom}_C(A, B)$ , we have  $f = g$ .

02BX

4. *Automatic Commutativity of Diagrams.* Every diagram in a posetal category commutes.

#### PROOF 11.2.7.1.3 ► PROOF OF PROPOSITION 11.2.7.1.2

##### Item 1: Functoriality

First, note that given a morphism of posets  $f: X \rightarrow Y$ , the corresponding functor  $f_\text{pos}: X_\text{pos} \rightarrow Y_\text{pos}$  is indeed a functor: since all morphisms in the Hom-sets of  $Y_\text{pos}$  are equal, it preserves identities and compositions trivially.

Next, we claim that  $(-)_\text{pos}$  is indeed a functor:

- *Preservation of Identities.* Let  $X \in \text{Obj}(\text{Pos})$ . Given  $x, y \in X$  with  $x \preceq_X y$ , we have

$$\begin{aligned} (\text{id}_X)_\text{pos}(x) &= \text{id}_X(x) \\ &= \text{id}_{X_\text{pos}}(x), \end{aligned}$$

so  $(\text{id}_X)_\text{pos}$  acts like the identity functor of  $X_\text{pos}$  on objects, and

$$\begin{aligned} (\text{id}_X)_\text{pos}(\text{pt}_{\text{Hom}_{X_\text{pos}}(x,y)}) &= \text{pt}_{\text{Hom}_{X_\text{pos}}((\text{id}_X)_\text{pos}(x), (\text{id}_X)_\text{pos}(y))} \\ &= \text{pt}_{\text{Hom}_{X_\text{pos}}(a,b)}, \end{aligned}$$

so the same holds for morphisms. Thus  $(\text{id}_X)_\text{pos} = \text{id}_{X_\text{pos}}$ .

- *Preservation of Composition.* Let  $X, Y, Z \in \text{Obj}(\text{Pos})$ . Given morphisms of posets  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$ , we need to show

$$(g \circ f)_\text{pos} = g_\text{pos} \circ f_\text{pos}.$$

Indeed, given  $x \in X$ , we have

$$\begin{aligned} (g \circ f)_\text{pos}(x) &= (g \circ f)(x) \\ &= g(f(x)) \end{aligned}$$

$$\begin{aligned} &= g_{\text{pos}}(f_{\text{pos}}(x)) \\ &= [g_{\text{pos}} \circ f_{\text{pos}}](x), \end{aligned}$$

so the identity holds on objects. Since  $Z_{\text{pos}}$  is a posetal category, the identity automatically holds on morphisms since

$$\begin{aligned} (g \circ f)_{\text{pos}}(\text{pt}_{\text{Hom}_{X_{\text{pos}}}(x,y)}) &= \text{pt}_{\text{Hom}_{Z_{\text{pos}}}(g_{\text{pos}}(f_{\text{pos}}(x)), g_{\text{pos}}(f_{\text{pos}}(y)))} \\ &= [g_{\text{pos}} \circ f_{\text{pos}}](\text{pt}_{\text{Hom}_{X_{\text{pos}}}(x,y)}) \end{aligned}$$

for each  $x, y \in X$  with  $x \preceq_X y$ .

Thus  $(-)^{\text{pos}}$  is indeed a functor.

Item 2: Fully Faithfulness

Omitted.

Item 3: Characterisations

Omitted.

Item 4: Automatic Commutativity of Diagrams

This follows from the fact that if  $C$  is posetal, then there is at most one morphism between any two objects, namely  $\text{pt}$ . 


## 00Y1 11.3 The Quadruple Adjunction With Sets

### 00Y2 11.3.1 Statement

Let  $C$  be a category.

#### 00Y3 PROPOSITION 11.3.1.1.1 ► THE QUADRUPLE ADJUNCTION BETWEEN Sets AND Cats

We have a quadruple adjunction



witnessed by bijections of sets

$$\text{Hom}_{\text{Sets}}(\pi_0(C), X) \cong \text{Hom}_{\text{Cats}}(C, X_{\text{disc}}),$$

$$\text{Hom}_{\text{Cats}}(X_{\text{disc}}, C) \cong \text{Hom}_{\text{Sets}}(X, \text{Obj}(C)),$$

$$\text{Hom}_{\text{Sets}}(\text{Obj}(C), X) \cong \text{Hom}_{\text{Cats}}(C, X_{\text{indisc}}),$$

natural in  $C \in \text{Obj}(\text{Cats})$  and  $X \in \text{Obj}(\text{Sets})$ , where

- The functor

$$\pi_0: \text{Cats} \rightarrow \text{Sets},$$

the **connected components functor**, is the functor sending a category to its set of connected components of [Definition 11.3.2.2.1](#).

- The functor

$$(-)_{\text{disc}}: \text{Sets} \rightarrow \text{Cats},$$

the **discrete category functor**, is the functor sending a set to its associated discrete category of [Item 1](#).

- The functor

$$\text{Obj}: \text{Cats} \rightarrow \text{Sets},$$

the **object functor**, is the functor sending a category to its set of objects.

- The functor

$$(-)_{\text{indisc}}: \text{Sets} \rightarrow \text{Cats},$$

the **indiscrete category functor**, is the functor sending a set to its associated indiscrete category of [Item 1](#).

#### PROOF 11.3.1.1.2 ► PROOF OF PROPOSITION 11.3.1.1.1

Omitted. 

**WARNING 11.3.1.1.3 ► PROPOSITION 11.3.1.1.1 CANNOT BE ENHANCED TO A 2-CATEGORICAL ADJUNCTION**



(This is a stub, to be revised and expanded upon later.)

The discrete category functor of [Proposition 11.3.1.1.1](#) lifts to a 2-functor, but it fails to preserve 2-categorical colimits, and hence lacks a right 2-adjoint. For instance, the 2-pushout of  $\text{pt} \leftarrow S^0 \rightarrow \text{pt}$  in  $\text{Sets}_{\text{disc}}$  is  $\text{pt}$ , but in  $\text{Cats}_2$  it is given by  $B\mathbb{Z}$ .

### 00Y4 11.3.2 Connected Components and Connected Categories

#### 00Y5 11.3.2.1 Connected Components of Categories

Let  $C$  be a category.

00Y6

**DEFINITION 11.3.2.1.1 ► CONNECTED COMPONENTS OF CATEGORIES**

A **connected component** of  $C$  is a full subcategory  $\mathcal{I}$  of  $C$  satisfying the following conditions:<sup>1</sup>

1. *Non-Emptiness.* We have  $\text{Obj}(\mathcal{I}) \neq \emptyset$ .
2. *Connectedness.* There exists a zigzag of arrows between any two objects of  $\mathcal{I}$ .

<sup>1</sup>In other words, a **connected component** of  $C$  is an element of the set  $\text{Obj}(C)/\sim$  with  $\sim$  the equivalence relation generated by the relation  $\sim'$  obtained by declaring  $A \sim' B$  iff there exists a morphism of  $C$  from  $A$  to  $B$ .

00Y7 11.3.2.2 Sets of Connected Components of Categories

Let  $C$  be a category.

00Y8

**DEFINITION 11.3.2.2.1 ► SETS OF CONNECTED COMPONENTS OF CATEGORIES**

The **set of connected components** of  $C$  is the set  $\pi_0(C)$  whose elements are the connected components of  $C$ .

00Y9

**PROPOSITION 11.3.2.2.2 ► PROPERTIES OF SETS OF CONNECTED COMPONENTS**

Let  $C$  be a category.

00YA

1. *Functoriality.* The assignment  $C \mapsto \pi_0(C)$  defines a functor

$$\pi_0: \text{Cats} \rightarrow \text{Sets}.$$

00YB

2. *Adjointness.* We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}): \text{Sets} \rightleftarrows \text{Cats.}$$

```

    \begin{array}{ccc}
    \pi_0 & & \\
    \downarrow & \swarrow & \searrow \\
    (-)_{\text{disc}} & & \\
    \downarrow & \swarrow & \searrow \\
    \text{Sets} & \rightleftarrows & \text{Cats.} \\
    \downarrow & \swarrow & \searrow \\
    (-)_{\text{indisc}} & & 
    \end{array}
  
```

00YC

3. *Interaction With Groupoids.* If  $C$  is a groupoid, then we have an isomorphism of categories

$$\pi_0(C) \cong K(C),$$

where  $K(C)$  is the set of isomorphism classes of  $C$  of ??.

00YD

4. *Preservation of Colimits.* The functor  $\pi_0$  of [Item 1](#) preserves colimits. In particular, we have bijections of sets

$$\begin{aligned}\pi_0(C \coprod \mathcal{D}) &\cong \pi_0(C) \coprod \pi_0(\mathcal{D}), \\ \pi_0(C \coprod_{\mathcal{E}} \mathcal{D}) &\cong \pi_0(C) \coprod_{\pi_0(\mathcal{E})} \pi_0(\mathcal{D}), \\ \pi_0(\text{CoEq}(C \xrightarrow[G]{F} \mathcal{D})) &\cong \text{CoEq}(\pi_0(C) \xrightarrow[\pi_0(G)]{\pi_0(F)} \pi_0(\mathcal{D})),\end{aligned}$$

natural in  $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$ .

00YE

5. *Symmetric Strong Monoidality With Respect to Coproducts.* The connected components functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\pi_0, \pi_0^{\coprod}, \pi_{0|1\!\!1}): (\text{Cats}, \coprod, \emptyset_{\text{cat}}) \rightarrow (\text{Sets}, \coprod, \emptyset),$$

being equipped with isomorphisms

$$\begin{aligned}\pi_{0|C,\mathcal{D}}^{\coprod}: \pi_0(C) \coprod \pi_0(\mathcal{D}) &\xrightarrow{\sim} \pi_0(C \coprod \mathcal{D}), \\ \pi_{0|1\!\!1}^{\coprod}: \emptyset &\xrightarrow{\sim} \pi_0(\emptyset_{\text{cat}}),\end{aligned}$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ .

00YF

6. *Symmetric Strong Monoidality With Respect to Products.* The connected components functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\pi_0, \pi_0^{\times}, \pi_{0|1\!\!1}^{\times}): (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt}),$$

being equipped with isomorphisms

$$\begin{aligned}\pi_{0|C,\mathcal{D}}^{\times}: \pi_0(C) \times \pi_0(\mathcal{D}) &\xrightarrow{\sim} \pi_0(C \times \mathcal{D}), \\ \pi_{0|1\!\!1}^{\times}: \text{pt} &\xrightarrow{\sim} \pi_0(\text{pt}),\end{aligned}$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ .

#### PROOF 11.3.2.2.3 ► PROOF OF PROPOSITION 11.3.2.2.2

Item 1: Functoriality

Omitted.

Item 2: Adjointness

This is proved in [Proposition 11.3.1.1.1](#).

Item 3: Interaction With Groupoids

Omitted.

Item 4: Preservation of Colimits

This follows from [Item 2](#) and ?? of ??.

Item 5: Symmetric Strong Monoidality With Respect to Coproducts

Omitted.

Item 6: Symmetric Strong Monoidality With Respect to Products

Omitted.



### 00YG 11.3.2.3 Connected Categories

00YH

#### DEFINITION 11.3.2.3.1 ► CONNECTED CATEGORIES

A category  $C$  is **connected** if  $\pi_0(C) \cong \text{pt}$ .<sup>1,2</sup>

<sup>1</sup>Further Terminology: A category is **disconnected** if it is not connected.

<sup>2</sup>Example: A groupoid is connected iff any two of its objects are isomorphic.

### 00YJ 11.3.3 Discrete Categories

00YK

#### DEFINITION 11.3.3.1.1 ► DISCRETE CATEGORIES

Let  $X$  be a set.

00YL

1. The **discrete category on  $X$**  is the category  $X_{\text{disc}}$  where

- *Objects.* We have

$$\text{Obj}(X_{\text{disc}}) \stackrel{\text{def}}{=} X.$$

- *Morphisms.* For each  $A, B \in \text{Obj}(X_{\text{disc}})$ , we have

$$\text{Hom}_{X_{\text{disc}}}(A, B) \stackrel{\text{def}}{=} \begin{cases} \text{id}_A & \text{if } A = B, \\ \emptyset & \text{if } A \neq B. \end{cases}$$

- *Identities.* For each  $A \in \text{Obj}(X_{\text{disc}})$ , the unit map

$$\mathbb{1}_A^{X_{\text{disc}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{disc}}}(A, A)$$

of  $X_{\text{disc}}$  at  $A$  is defined by

$$\text{id}_A^{X_{\text{disc}}} \stackrel{\text{def}}{=} \text{id}_A.$$

- *Composition.* For each  $A, B, C \in \text{Obj}(X_{\text{disc}})$ , the composition map

$$\circ_{A,B,C}^{X_{\text{disc}}} : \text{Hom}_{X_{\text{disc}}}(B, C) \times \text{Hom}_{X_{\text{disc}}}(A, B) \rightarrow \text{Hom}_{X_{\text{disc}}}(A, C)$$

of  $X_{\text{disc}}$  at  $(A, B, C)$  is defined by

$$\text{id}_A \circ \text{id}_A \stackrel{\text{def}}{=} \text{id}_A.$$

**00YM** 2. A category  $C$  is **discrete** if it is equivalent to  $X_{\text{disc}}$  for some set  $X$ .

**00YN**

### PROPOSITION 11.3.3.1.2 ► PROPERTIES OF DISCRETE CATEGORIES ON SETS

Let  $X$  be a set.

**00YP**

1. *Functionality.* The assignment  $X \mapsto X_{\text{disc}}$  defines a functor

$$(-)_{\text{disc}} : \text{Sets} \rightarrow \text{Cats}.$$

**00YQ**

2. *Adjointness.* We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}) : \text{Sets} \rightleftarrows \text{Cats.}$$

**00YR**

3. *Symmetric Strong Monoidality With Respect to Coproducts.* The functor of Item 1 has a symmetric strong monoidal structure

$$((-)_{\text{disc}}, (-)_{\text{disc}}^{\coprod}, (-)_{\text{disc}|\mathbb{1}}^{\coprod}) : (\text{Sets}, \coprod, \emptyset) \rightarrow (\text{Cats}, \coprod, \emptyset_{\text{cat}}),$$

being equipped with isomorphisms

$$(-)_{\text{disc}|X,Y}^{\coprod} : X_{\text{disc}} \coprod Y_{\text{disc}} \xrightarrow{\sim} (X \coprod Y)_{\text{disc}},$$

$$(-)_{\text{disc}|\mathbb{1}}^{\coprod} : \emptyset_{\text{cat}} \xrightarrow{\sim} \emptyset_{\text{disc}},$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

**00YS**

4. *Symmetric Strong Monoidality With Respect to Products.* The functor of Item 1 has a symmetric strong monoidal structure

$$((-)_{\text{disc}}, (-)_{\text{disc}}^{\times}, (-)_{\text{disc}|\mathbb{1}}^{\times}) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Cats}, \times, \text{pt}),$$

being equipped with isomorphisms

$$(-)_{\text{disc}|X,Y}^{\times} : X_{\text{disc}} \times Y_{\text{disc}} \xrightarrow{\sim} (X \times Y)_{\text{disc}},$$

$$(-)_{\text{disc}|\mathbb{1}}^{\times} : \text{pt} \xrightarrow{\sim} \text{pt}_{\text{disc}},$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

#### PROOF 11.3.3.1.3 ▶ PROOF OF PROPOSITION 11.3.3.1.2

Item 1: Functoriality

Omitted.

Item 2: Adjointness

This is proved in [Proposition 11.3.1.1.1](#).

Item 3: Symmetric Strong Monoidality With Respect to Coproducts

Omitted.

Item 4: Symmetric Strong Monoidality With Respect to Products

Omitted.



### 00YT 11.3.4 Indiscrete Categories

00YU

#### DEFINITION 11.3.4.1.1 ▶ INDISCRETE CATEGORIES

Let  $X$  be a set.

00YV

1. The **indiscrete category on  $X^1$**  is the category  $X_{\text{indisc}}$  where

- *Objects.* We have

$$\text{Obj}(X_{\text{indisc}}) \stackrel{\text{def}}{=} X.$$

- *Morphisms.* For each  $A, B \in \text{Obj}(X_{\text{indisc}})$ , we have

$$\begin{aligned} \text{Hom}_{X_{\text{disc}}}(A, B) &\stackrel{\text{def}}{=} \{[A] \rightarrow [B]\} \\ &\cong \text{pt}. \end{aligned}$$

- *Identities.* For each  $A \in \text{Obj}(X_{\text{indisc}})$ , the unit map

$$\mathbb{1}_A^{X_{\text{indisc}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{indisc}}}(A, A)$$

of  $X_{\text{indisc}}$  at  $A$  is defined by

$$\text{id}_A^{X_{\text{indisc}}} \stackrel{\text{def}}{=} \{[A] \rightarrow [A]\}.$$

- *Composition.* For each  $A, B, C \in \text{Obj}(X_{\text{indisc}})$ , the composition map

$$\circ^{X_{\text{indisc}}}_{A,B,C} : \text{Hom}_{X_{\text{indisc}}}(B, C) \times \text{Hom}_{X_{\text{indisc}}}(A, B) \rightarrow \text{Hom}_{X_{\text{indisc}}}(A, C)$$

of  $X_{\text{disc}}$  at  $(A, B, C)$  is defined by

$$([B] \rightarrow [C]) \circ ([A] \rightarrow [B]) \stackrel{\text{def}}{=} ([A] \rightarrow [C]).$$

- 00YW** 2. A category  $C$  is **indiscrete** if it is equivalent to  $X_{\text{indisc}}$  for some set  $X$ .

<sup>1</sup>Further Terminology: Sometimes called the **chaotic category on  $X$** .

**00YX**

#### PROPOSITION 11.3.4.1.2 ► PROPERTIES OF INDISCRETE CATEGORIES ON SETS

Let  $X$  be a set.

**00YY**

1. *Functoriality.* The assignment  $X \mapsto X_{\text{indisc}}$  defines a functor

$$(-)_{\text{indisc}} : \text{Sets} \rightarrow \text{Cats}.$$

**00YZ**

2. *Adjointness.* We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}) : \text{Sets} \begin{array}{c} \xleftarrow{\quad \perp \quad} \\[-1ex] \xleftarrow{\quad (-)_{\text{disc}} \quad} \\[-1ex] \xleftarrow{\quad \perp \quad} \\[-1ex] \xleftarrow{\quad \text{Obj} \quad} \\[-1ex] \xleftarrow{\quad \perp \quad} \\[-1ex] \xleftarrow{\quad (-)_{\text{indisc}} \quad} \end{array} \text{Cats.}$$

**00Z0**

3. *Symmetric Strong Monoidality With Respect to Products.* The functor of **Item 1** has a symmetric strong monoidal structure

$$((-)_{\text{indisc}}, (-)_{\text{indisc}}^{\times}, (-)_{\text{indisc}|1}^{\times}) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Cats}, \times, \text{pt}),$$

being equipped with isomorphisms

$$(-)_{\text{indisc}|X,Y}^{\times} : X_{\text{indisc}} \times Y_{\text{indisc}} \xrightarrow{\sim} (X \times Y)_{\text{indisc}},$$

$$(-)_{\text{indisc}|1}^{\times} : \text{pt} \xrightarrow{\sim} \text{pt}_{\text{indisc}},$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

**PROOF 11.3.4.1.3 ► PROOF OF PROPOSITION 11.3.4.1.2**

Item 1: Functoriality

Omitted.

Item 2: Adjointness

This is proved in [Proposition 11.3.1.1.1](#).

Item 3: Symmetric Strong Monoidality With Respect to Products

Omitted.

**00Z1 11.4 Groupoids****01U1 11.4.1 Isomorphisms**

Let  $C$  be a category.

00Z3

**DEFINITION 11.4.1.1.1 ► ISOMORPHISMS**

A morphism  $f: A \rightarrow B$  of  $C$  is an **isomorphism** if there exists a morphism  $f^{-1}: B \rightarrow A$  of  $C$  such that

$$\begin{aligned} f \circ f^{-1} &= \text{id}_B, \\ f^{-1} \circ f &= \text{id}_A. \end{aligned}$$

00Z4

**NOTATION 11.4.1.1.2 ► THE SET OF ISOMORPHISMS BETWEEN TWO OBJECTS IN A CATEGORY**

We write  $\text{Iso}_C(A, B)$  for the set of all isomorphisms in  $C$  from  $A$  to  $B$ .

**01U2 11.4.2 Groupoids**

00Z5

**DEFINITION 11.4.2.1.1 ► GROUPOIDS**

A **groupoid** is a category in which every morphism is an isomorphism.

01U3

**EXAMPLE 11.4.2.1.2 ► GROUPS AS ONE-OBJECT GROUPOIDS**

The isomorphism of categories of Example 11.2.3.1.1 restricts to an isomorphism

$$\begin{array}{ccc} \text{Grp} & \longrightarrow & \text{Grpd} \\ \text{Grp} \cong \underset{\text{Sets}}{\text{pt} \times \text{Grpd}}, & \downarrow & \downarrow \text{Obj} \\ \text{pt} & \xrightarrow{[\text{pt}]} & \text{Sets} \end{array}$$

where Grpd is the full subcategory of Cats spanned by the groupoids. In other words, we have an identification

$$\{\text{Groups}\} \cong \{\text{One-object groupoids}\}.$$

### 00Z6 11.4.3 The Groupoid Completion of a Category

Let  $C$  be a category.

#### 00Z7 DEFINITION 11.4.3.1.1 ► THE GROUPOID COMPLETION OF A CATEGORY

The **groupoid completion** of  $C$ <sup>1</sup> is the pair  $(K_0(C), \iota_C)$  consisting of

- A groupoid  $K_0(C)$ ;
- A functor  $\iota_C: C \rightarrow K_0(C)$ ;

satisfying the following universal property:<sup>2</sup>

(UP) Given another such pair  $(\mathcal{G}, i)$ , there exists a unique functor  $K_0(C) \xrightarrow{\exists!} \mathcal{G}$  making the diagram

$$\begin{array}{ccc} & K_0(C) & \\ \iota_C \nearrow & \downarrow \exists! & \\ C & \xrightarrow{i} & \mathcal{G} \end{array}$$

commute.

<sup>1</sup>Further Terminology: Also called the **Grothendieck groupoid** of  $C$  or the **Grothendieck groupoid completion** of  $C$ . See Item 5 of Proposition 11.4.3.1.4 for an explicit construction.

00Z8

**CONSTRUCTION 11.4.3.1.2 ► CONSTRUCTION OF THE GROUPOID COMPLETION OF A CATEGORY**

Concretely, the groupoid completion of  $C$  is the Gabriel–Zisman localisation  $\text{Mor}(C)^{-1}C$  of  $C$  at the set  $\text{Mor}(C)$  of all morphisms of  $C$ ; see ??.

(To be expanded upon later on.)

**PROOF 11.4.3.1.3 ► PROOF OF CONSTRUCTION 11.4.3.1.2**

Omitted. 

00Z9

**PROPOSITION 11.4.3.1.4 ► PROPERTIES OF GROUPOID COMPLETION**

Let  $C$  be a category.

00ZA

- Functoriality.* The assignment  $C \mapsto K_0(C)$  defines a functor

$$K_0 : \text{Cats} \rightarrow \text{Grpd}.$$

00ZB

- 2-Functoriality.* The assignment  $C \mapsto K_0(C)$  defines a 2-functor

$$K_0 : \text{Cats}_2 \rightarrow \text{Grpd}_2.$$

00ZC

- Adjointness.* We have an adjunction

$$(K_0 \dashv \iota) : \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \xleftarrow{\iota} \end{array} \text{Grpd},$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

natural in  $C \in \text{Obj}(\text{Cats})$  and  $\mathcal{G} \in \text{Obj}(\text{Grpd})$ , forming, together with the functor  $\text{Core}$  of Item 1 of Proposition 11.4.4.1.5, a triple adjunction

$$(K_0 \dashv \iota \dashv \text{Core}) : \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \xleftarrow{\iota} \xrightarrow{\perp} \\ \text{Core} \end{array} \text{Grpd},$$

witnessed by bijections of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$  and  $\mathcal{G} \in \text{Obj}(\text{Grpd})$ .

00ZD

4. *2-Adjointness.* We have a 2-adjunction

$$(K_0 \dashv \iota): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp_2 \\ \xleftarrow{\iota} \end{array} \text{Grpd},$$

witnessed by an isomorphism of categories

$$\text{Fun}(K_0(C), \mathcal{G}) \cong \text{Fun}(C, \mathcal{G}),$$

natural in  $C \in \text{Obj}(\text{Cats})$  and  $\mathcal{G} \in \text{Obj}(\text{Grpd})$ , forming, together with the 2-functor Core of [Item 2 of Proposition 11.4.4.1.5](#), a triple 2-adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp_2 \\ \xleftarrow{\iota} \xrightarrow{\text{Core}} \\ \perp_2 \end{array} \text{Grpd},$$

witnessed by isomorphisms of categories

$$\text{Fun}(K_0(C), \mathcal{G}) \cong \text{Fun}(C, \mathcal{G}),$$

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$  and  $\mathcal{G} \in \text{Obj}(\text{Grpd})$ .

00ZE

5. *Interaction With Classifying Spaces.* We have an isomorphism of groupoids

$$K_0(C) \cong \Pi_{\leq 1}(|N_\bullet(C)|),$$

natural in  $C \in \text{Obj}(\text{Cats})$ ; i.e. the diagram

$$\begin{array}{ccc} \text{Cats} & \xrightarrow{K_0} & \text{Grp} \\ N_\bullet \downarrow & \Downarrow \gamma & \uparrow \Pi_{\leq 1} \\ \text{sSets} & \xrightarrow{|-|} & \Pi \end{array}$$

commutes up to natural isomorphism.

00ZF

6. *Symmetric Strong Monoidality With Respect to Coproducts.* The groupoid completion functor of [Item 1](#) has a symmetric strong monoidal structure

$$(K_0, K_0^{\coprod}, K_{0|1}^{\coprod}): (\text{Cats}, \coprod, \emptyset_{\text{cat}}) \rightarrow (\text{Grpd}, \coprod, \emptyset_{\text{cat}})$$

being equipped with isomorphisms

$$\begin{aligned} K_0^{\coprod}_{|C, \mathcal{D}} : K_0(C) \coprod K_0(\mathcal{D}) &\xrightarrow{\sim} K_0(C \coprod \mathcal{D}), \\ K_0^{\coprod}_{|\mathbb{1}} : \emptyset_{\text{cat}} &\xrightarrow{\sim} K_0(\emptyset_{\text{cat}}), \end{aligned}$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ .

00ZG

7. *Symmetric Strong Monoidality With Respect to Products.* The groupoid completion functor of [Item 1](#) has a symmetric strong monoidal structure

$$(K_0, K_0^{\times}, K_0^{\times}_{|\mathbb{1}}) : (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Grpd}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned} K_0^{\times}_{|C, \mathcal{D}} : K_0(C) \times K_0(\mathcal{D}) &\xrightarrow{\sim} K_0(C \times \mathcal{D}), \\ K_0^{\times}_{|\mathbb{1}} : \text{pt} &\xrightarrow{\sim} K_0(\text{pt}), \end{aligned}$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ .

#### PROOF 11.4.3.1.5 ▶ PROOF OF PROPOSITION 11.4.3.1.4

Item 1: Functoriality

Omitted.

Item 2: 2-Functoriality

Omitted.

Item 3: Adjointness

Omitted.

Item 4: 2-Adjointness

Omitted.

Item 5: Interaction With Classifying Spaces

See Corollary 18.33 of <https://web.ma.utexas.edu/users/dafm/M392C-2012/Notes/lecture18.pdf>.

Item 6: Symmetric Strong Monoidality With Respect to Coproducts

Omitted.

Item 7: Symmetric Strong Monoidality With Respect to Products

Omitted.



**00ZH 11.4.4 The Core of a Category**

Let  $C$  be a category.

**00ZJ**

**DEFINITION 11.4.4.1.1 ► THE CORE OF A CATEGORY**

The **core** of  $C$  is the pair  $(\text{Core}(C), \iota_C)$  consisting of

- A groupoid  $\text{Core}(C)$ ;
- A functor  $\iota_C: \text{Core}(C) \hookrightarrow C$ ;

satisfying the following universal property:

(UP) Given another such pair  $(G, i)$ , there exists a unique functor  $G \xrightarrow{\exists!} \text{Core}(C)$  making the diagram

$$\begin{array}{ccc} & \text{Core}(C) & \\ & \nearrow \exists! & \downarrow \iota_C \\ G & \xrightarrow{i} & C \end{array}$$

commute.

**00ZK**

**NOTATION 11.4.4.1.2 ► ALTERNATIVE NOTATION FOR THE CORE OF A CATEGORY**

We also write  $C^\simeq$  for  $\text{Core}(C)$ .

**00ZL**

**CONSTRUCTION 11.4.4.1.3 ► CONSTRUCTION OF THE CORE OF A CATEGORY**

The core of  $C$  is the wide subcategory of  $C$  spanned by the isomorphisms of  $C$ , i.e. the category  $\text{Core}(C)$  where<sup>1</sup>

1. *Objects.* We have

$$\text{Obj}(\text{Core}(C)) \stackrel{\text{def}}{=} \text{Obj}(C).$$

2. *Morphisms.* The morphisms of  $\text{Core}(C)$  are the isomorphisms of  $C$ .

---

<sup>1</sup>*Slogan:* The groupoid  $\text{Core}(C)$  is the maximal subgroupoid of  $C$ .

**PROOF 11.4.4.1.4 ► PROOF OF CONSTRUCTION 11.4.4.1.3**

This follows from the fact that functors preserve isomorphisms ([Item 1 of Proposition 11.5.1.1.8](#)). 

00ZM

**PROPOSITION 11.4.4.1.5 ► PROPERTIES OF THE CORE OF A CATEGORY**

Let  $C$  be a category.

00ZN

1. *Functoriality.* The assignment  $C \mapsto \text{Core}(C)$  defines a functor

$$\text{Core}: \text{Cats} \rightarrow \text{Grpd}.$$

00ZP

2. *2-Functoriality.* The assignment  $C \mapsto \text{Core}(C)$  defines a 2-functor

$$\text{Core}: \text{Cats}_2 \rightarrow \text{Grpd}_2.$$

00ZQ

3. *Adjointness.* We have an adjunction

$$(\iota \dashv \text{Core}): \text{Grpd} \begin{array}{c} \xrightarrow{\iota} \\ \perp \\ \xleftarrow{\text{Core}} \end{array} \text{Cats},$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in  $\mathcal{G} \in \text{Obj}(\text{Grpd})$  and  $\mathcal{D} \in \text{Obj}(\text{Cats})$ , forming, together with the functor  $K_0$  of [Item 1 of Proposition 11.4.3.1.4](#), a triple adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xleftarrow{\iota} \\ \perp \\ \xrightarrow{\text{Core}} \\ \perp \\ \xleftarrow{K_0} \end{array} \text{Grpd},$$

witnessed by bijections of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$  and  $\mathcal{G} \in \text{Obj}(\text{Grpd})$ .

00ZR

4. *2-Adjointness.* We have an adjunction

$$(\iota \dashv \text{Core}): \text{Grpd} \begin{array}{c} \xrightarrow{\iota} \\ \perp_2 \\ \xleftarrow{\text{Core}} \end{array} \text{Cats},$$

witnessed by an isomorphism of categories

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in  $\mathcal{G} \in \text{Obj}(\text{Grpd})$  and  $\mathcal{D} \in \text{Obj}(\text{Cats})$ , forming, together with the 2-functor  $K_0$  of [Item 2 of Proposition 11.4.3.1.4](#), a triple 2-adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xleftarrow{\quad \iota \quad} \\[-1ex] \xrightarrow{\quad K_0 \quad} \\[-1ex] \xleftarrow{\quad \text{Core} \quad} \end{array} \text{Grpd},$$

witnessed by isomorphisms of categories

$$\text{Fun}(K_0(C), \mathcal{G}) \cong \text{Fun}(C, \mathcal{G}),$$

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$  and  $\mathcal{G} \in \text{Obj}(\text{Grpd})$ .

**00ZS**

5. *Symmetric Strong Monoidality With Respect to Products.* The core functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\text{Core}, \text{Core}^\times, \text{Core}_{\mathbb{1}}^\times): (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Grpd}, \times, \text{pt})$$

being equipped with isomorphisms

$$\text{Core}_{C, \mathcal{D}}^\times: \text{Core}(C) \times \text{Core}(\mathcal{D}) \xrightarrow{\sim} \text{Core}(C \times \mathcal{D}),$$

$$\text{Core}_{\mathbb{1}}^\times: \text{pt} \xrightarrow{\sim} \text{Core}(\text{pt}),$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ .

**00ZT**

6. *Symmetric Strong Monoidality With Respect to Coproducts.* The core functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\text{Core}, \text{Core}^{\coprod}, \text{Core}_{\mathbb{1}}^{\coprod}): (\text{Cats}, \coprod, \emptyset_{\text{cat}}) \rightarrow (\text{Grpd}, \coprod, \emptyset_{\text{cat}})$$

being equipped with isomorphisms

$$\text{Core}_{C, \mathcal{D}}^{\coprod}: \text{Core}(C) \coprod \text{Core}(\mathcal{D}) \xrightarrow{\sim} \text{Core}(C \coprod \mathcal{D}),$$

$$\text{Core}_{\mathbb{1}}^{\coprod}: \emptyset_{\text{cat}} \xrightarrow{\sim} \text{Core}(\emptyset_{\text{cat}}),$$

natural in  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ .

**PROOF 11.4.4.1.6 ► PROOF OF PROPOSITION 11.4.4.1.5**

**Item 1: Functoriality**

Omitted.

**Item 2: 2-Functoriality**

Omitted.

**Item 3: Adjointness**

Omitted.

**Item 4: 2-Adjointness**

Omitted.

**Item 5: Symmetric Strong Monoidality With Respect to Products**

Omitted.

**Item 6: Symmetric Strong Monoidality With Respect to Coproducts**

Omitted.

**00ZU 11.5 Functors****00ZV 11.5.1 Foundations**

Let  $C$  and  $\mathcal{D}$  be categories.

00ZW

**DEFINITION 11.5.1.1.1 ► FUNCTORS**

A **functor**  $F: C \rightarrow \mathcal{D}$  from  $C$  to  $\mathcal{D}$ <sup>1</sup> consists of:

1. *Action on Objects.* A map of sets

$$F: \text{Obj}(C) \rightarrow \text{Obj}(\mathcal{D}),$$

called the **action on objects** of  $F$ .

2. *Action on Morphisms.* For each  $A, B \in \text{Obj}(C)$ , a map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F(A), F(B)),$$

called the **action on morphisms** of  $F$  at  $(A, B)$ <sup>2</sup>.

satisfying the following conditions:

1. *Preservation of Identities.* For each  $A \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccc} & \text{pt} & \\ & \downarrow \mathbb{1}_A^C & \searrow \mathbb{1}_{F(A)}^{\mathcal{D}} \\ \text{Hom}_C(A, A) & \xrightarrow{F_{A,A}} & \text{Hom}_{\mathcal{D}}(F(A), F(A)) \end{array}$$

commutes, i.e. we have

$$F(\text{id}_A) = \text{id}_{F(A)}.$$

2. *Preservation of Composition.* For each  $A, B, C \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccc} \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ F_{B,C} \times F_{A,B} \downarrow & & \downarrow F_{A,C} \\ \text{Hom}_{\mathcal{D}}(F(B), F(C)) \times \text{Hom}_{\mathcal{D}}(F(A), F(B)) & \xrightarrow{\circ_{F(A), F(B), F(C)}^{\mathcal{D}}} & \text{Hom}_{\mathcal{D}}(F(A), F(C)) \end{array}$$

commutes, i.e. for each composable pair  $(g, f)$  of morphisms of  $C$ , we have

$$F(g \circ f) = F(g) \circ F(f).$$

<sup>1</sup>Further Terminology: Also called a **covariant functor**.

<sup>2</sup>Further Terminology: Also called **action on Hom-sets of  $F$  at  $(A, B)$** .

00ZX

#### NOTATION 11.5.1.1.2 ► SUBSCRIPT AND SUPERSCRIPT NOTATION FOR FUNCTORS

Let  $C$  and  $\mathcal{D}$  be categories, and write  $C^{\text{op}}$  for the opposite category of  $C$  of ??.

00ZY

1. Given a functor

$$F: C \rightarrow \mathcal{D},$$

we also write  $F_A$  for  $F(A)$ .

00ZZ

2. Given a functor

$$F: C^{\text{op}} \rightarrow \mathcal{D},$$

we also write  $F^A$  for  $F(A)$ .

0100

3. Given a functor

$$F: C \times C \rightarrow \mathcal{D},$$

we also write  $F_{A,B}$  for  $F(A, B)$ .

0101

4. Given a functor

$$F: C^{\text{op}} \times C \rightarrow \mathcal{D},$$

we also write  $F_B^A$  for  $F(A, B)$ .

We employ a similar notation for morphisms, writing e.g.  $F_f$  for  $F(f)$  given a functor  $F: C \rightarrow \mathcal{D}$ .

0102

**NOTATION 11.5.1.1.3 ► ADDITIONAL NOTATION FOR FUNCTORS**

Following the notation  $\llbracket x \mapsto f(x) \rrbracket$  for a function  $f: X \rightarrow Y$  introduced in [Notation 3.1.1.1.2](#), we will sometimes denote a functor  $F: C \rightarrow \mathcal{D}$  by

$$F \stackrel{\text{def}}{=} \llbracket A \mapsto F(A) \rrbracket,$$

specially when the action on morphisms of  $F$  is clear from its action on objects.

0103

**EXAMPLE 11.5.1.1.4 ► IDENTITY FUNCTORS**

The **identity functor** of a category  $C$  is the functor  $\text{id}_C: C \rightarrow C$  where

1. *Action on Objects.* For each  $A \in \text{Obj}(C)$ , we have

$$\text{id}_C(A) \stackrel{\text{def}}{=} A.$$

2. *Action on Morphisms.* For each  $A, B \in \text{Obj}(C)$ , the action on morphisms

$$(\text{id}_C)_{A,B}: \text{Hom}_C(A, B) \rightarrow \underbrace{\text{Hom}_C(\text{id}_C(A), \text{id}_C(B))}_{\stackrel{\text{def}}{=} \text{Hom}_C(A, B)}$$

of  $\text{id}_C$  at  $(A, B)$  is defined by

$$(\text{id}_C)_{A,B} \stackrel{\text{def}}{=} \text{id}_{\text{Hom}_C(A, B)}.$$

**PROOF 11.5.1.1.5 ► PROOF OF EXAMPLE 11.5.1.1.4****Preservation of Identities**

We have  $\text{id}_C(\text{id}_A) \stackrel{\text{def}}{=} \text{id}_A$  for each  $A \in \text{Obj}(C)$  by definition.

**Preservation of Compositions**

For each composable pair  $A \xrightarrow{f} B \xrightarrow{g} B$  of morphisms of  $C$ , we have

$$\begin{aligned} \text{id}_C(g \circ f) &\stackrel{\text{def}}{=} g \circ f \\ &\stackrel{\text{def}}{=} \text{id}_C(g) \circ \text{id}_C(f). \end{aligned}$$

This finishes the proof. 

0104

**DEFINITION 11.5.1.1.6 ► COMPOSITION OF FUNCTORS**

The **composition** of two functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  is the functor  $G \circ F$  where

- *Action on Objects.* For each  $A \in \text{Obj}(\mathcal{C})$ , we have

$$[G \circ F](A) \stackrel{\text{def}}{=} G(F(A)).$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(\mathcal{C})$ , the action on morphisms

$$(G \circ F)_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{E}}(G_{F_A}, G_{F_B})$$

of  $G \circ F$  at  $(A, B)$  is defined by

$$[G \circ F](f) \stackrel{\text{def}}{=} G(F(f)).$$

**PROOF 11.5.1.1.7 ► PROOF OF DEFINITION 11.5.1.1.6****Preservation of Identities**

For each  $A \in \text{Obj}(\mathcal{C})$ , we have

$$\begin{aligned} G_{F_{\text{id}_A}} &= G_{\text{id}_{F_A}} && (\text{functoriality of } F) \\ &= \text{id}_{G_{F_A}}. && (\text{functoriality of } G) \end{aligned}$$

**Preservation of Composition**

For each composable pair  $(g, f)$  of morphisms of  $\mathcal{C}$ , we have

$$\begin{aligned} G_{F_{g \circ f}} &= G_{F_g \circ F_f} && (\text{functoriality of } F) \\ &= G_{F_g} \circ G_{F_f}. && (\text{functoriality of } G) \end{aligned}$$

This finishes the proof. 

0105

**PROPOSITION 11.5.1.1.8 ► ELEMENTARY PROPERTIES OF FUNCTORS**

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a functor.

0106

1. *Preservation of Isomorphisms.* If  $f$  is an isomorphism in  $\mathcal{C}$ , then  $F(f)$  is an isomorphism in  $\mathcal{D}$ .<sup>1</sup>

<sup>1</sup>When the converse holds, we call  $F$  *conservative*, see Definition 11.6.4.1.1.

**PROOF 11.5.1.1.9 ▶ PROOF OF PROPOSITION 11.5.1.1.8****Item 1: Preservation of Isomorphisms**

Indeed, we have

$$\begin{aligned} F(f)^{-1} \circ F(f) &= F(f^{-1} \circ f) \\ &= F(\text{id}_A) \\ &= \text{id}_{F(A)} \end{aligned}$$

and

$$\begin{aligned} F(f) \circ F(f)^{-1} &= F(f \circ f^{-1}) \\ &= F(\text{id}_B) \\ &= \text{id}_{F(B)}, \end{aligned}$$

showing  $F(f)$  to be an isomorphism. 

**0107 11.5.2 Contravariant Functors**

Let  $C$  and  $\mathcal{D}$  be categories, and let  $C^{\text{op}}$  denote the opposite category of  $C$  of  $??$ .

**0108****DEFINITION 11.5.2.1.1 ▶ CONTRAVARIANT FUNCTORS**

A **contravariant functor** from  $C$  to  $\mathcal{D}$  is a functor from  $C^{\text{op}}$  to  $\mathcal{D}$ .

**0109****REMARK 11.5.2.1.2 ▶ UNWINDING DEFINITION 11.5.2.1.1**

In detail, a **contravariant functor** from  $C$  to  $\mathcal{D}$  consists of:

1. *Action on Objects.* A map of sets

$$F: \text{Obj}(C) \rightarrow \text{Obj}(\mathcal{D}),$$

called the **action on objects of  $F$** .

2. *Action on Morphisms.* For each  $A, B \in \text{Obj}(C)$ , a map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F(B), F(A)),$$

called the **action on morphisms of  $F$  at  $(A, B)$** .

satisfying the following conditions:

1. *Preservation of Identities.* For each  $A \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccc} \text{pt} & & \\ \downarrow \mathbb{1}_A^C & \searrow \mathbb{1}_{F(A)}^D & \\ \text{Hom}_C(A, A) & \xrightarrow{F_{A,A}} & \text{Hom}_D(F(A), F(A)) \end{array}$$

commutes, i.e. we have

$$F(\text{id}_A) = \text{id}_{F(A)}.$$

2. *Preservation of Composition.* For each  $A, B, C \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccccc} & & \text{Hom}_D(F(C), F(B)) \times \text{Hom}_D(F(B), F(A)) & & \\ & \nearrow F_{B,C} \times F_{A,B} & & \searrow \sigma_{\text{Hom}_D(F(C), F(B)), \text{Hom}_D(F(B), F(A))}^{\text{Sets}} & \\ \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & & \text{Hom}_D(F(B), F(A)) \times \text{Hom}_D(F(C), F(B)) & & \\ \downarrow \circ_{A,B,C}^C & & & \downarrow \circ_{F(C), F(B), F(A)}^D & \\ \text{Hom}_C(A, C) & \xrightarrow{F_{A,C}} & \text{Hom}_D(F(C), F(A)) & & \end{array}$$

commutes, i.e. for each composable pair  $(g, f)$  of morphisms of  $C$ , we have

$$F(g \circ f) = F(f) \circ F(g).$$

010A

#### REMARK 11.5.2.1.3 ► ON THE TERM CONTRAVARIANT FUNCTOR

Throughout this work we will not use the term “contravariant” functor, speaking instead simply of functors  $F: C^{\text{op}} \rightarrow D$ . We will usually, however, write

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_D(F(B), F(A))$$

for the action on morphisms

$$F_{A,B}: \text{Hom}_{C^{\text{op}}}(A, B) \rightarrow \text{Hom}_D(F(A), F(B))$$

of  $F$ , as well as write  $F(g \circ f) = F(f) \circ F(g)$ .

## 010B 11.5.3 Forgetful Functors

010C

## DEFINITION 11.5.3.1.1 ► FORGETFUL FUNCTORS

There isn't a precise definition of a **forgetful functor**.

010D

## REMARK 11.5.3.1.2 ► UNWINDING DEFINITION 11.5.3.1.1

Despite there not being a formal or precise definition of a forgetful functor, the term is often very useful in practice, similarly to the word “canonical”. The idea is that a “forgetful functor” is a functor that forgets structure or properties, and is best explained through examples, such as the ones below (see Examples 11.5.3.1.3 and 11.5.3.1.4).

010E

## EXAMPLE 11.5.3.1.3 ► FORGETFUL FUNCTORS THAT FORGET STRUCTURE

Examples of forgetful functors that forget structure include:

010F

1. *Forgetting Group Structures.* The functor  $\text{Grp} \rightarrow \text{Sets}$  sending a group  $(G, \mu_G, \eta_G)$  to its underlying set  $G$ , forgetting the multiplication and unit maps  $\mu_G$  and  $\eta_G$  of  $G$ .

010G

2. *Forgetting Topologies.* The functor  $\text{Tl} \rightarrow \text{Sets}$  sending a topological space  $(X, \mathcal{T}_X)$  to its underlying set  $X$ , forgetting the topology  $\mathcal{T}_X$ .

010H

3. *Forgetting Fibrations.* The functor  $\text{FibSets}(K) \rightarrow \text{Sets}$  sending a  $K$ -fibred set  $\phi_X : X \rightarrow K$  to the set  $X$ , forgetting the map  $\phi_X$  and the base set  $K$ .

010J

## EXAMPLE 11.5.3.1.4 ► FORGETFUL FUNCTORS THAT FORGET PROPERTIES

Examples of forgetful functors that forget properties include:

010K

1. *Forgetting Commutativity.* The inclusion functor  $\iota : \text{CMon} \hookrightarrow \text{Mon}$  which forgets the property of being commutative.

010L

2. *Forgetting Inverses.* The inclusion functor  $\iota : \text{Grp} \hookrightarrow \text{Mon}$  which forgets the property of having inverses.

010M

## NOTATION 11.5.3.1.5 ► NOTATION FOR FORGETFUL FUNCTORS THAT FORGET STRUCTURE

Throughout this work, we will denote forgetful functors that forget structure by 忘, e.g. as in

忘:  $\text{Grp} \rightarrow \text{Sets}$ .

The symbol 忘, pronounced *wasureru* (see Item 1 of Remark 11.5.3.1.6 below), means *to forget*, and is a kanji found in the following words in Japanese and Chinese:

- 010N 1. 忘れる, transcribed as *wasureru*, meaning *to forget*.
- 010P 2. 忘却関手, transcribed as *boukyaku kanshu*, meaning *forgetful functor*.
- 010Q 3. 忘记 or 忘記, transcribed as *wàngjì*, meaning *to forget*.
- 010R 4. 遗忘函子 or 遺忘函子, transcribed as *yíwàng hánzì*, meaning *forgetful functor*.

#### 010S REMARK 11.5.3.1.6 ► PRONUNCIATION OF THE WORDS IN NOTATION 11.5.3.1.5

Here we collect the pronunciation of the words in Notation 11.5.3.1.5 for accuracy and completeness.

- 010T 1. Pronunciation of 忘れる:
  - See [here](#).
  - IPA broad transcription: [wäsureru].
  - IPA narrow transcription: [u̯päsi̯l̩r̩eu̯p̩].
- 010U 2. Pronunciation of 忘却関手: Pronunciation:
  - See [here](#).
  - IPA broad transcription: [bø:kjäku kã:ŋçeu].
  - IPA narrow transcription: [bø:kjäkup̩ kã:ŋçeu].
- 010V 3. Pronunciation of 忘记:
  - See [here](#).
  - Broad IPA transcription: [waŋtei].
  - Sinological IPA transcription: [wan<sup>51-53</sup>fci<sup>51</sup>].
- 010W 4. Pronunciation of 遗忘函子:
  - See [here](#).
  - Broad IPA transcription: [iwaŋ xänfszi].
  - Sinological IPA transcription: [i<sup>35</sup>wan<sup>51</sup>xän<sup>35</sup>fsz<sup>214-21(4)</sup>].

**010X 11.5.4 The Natural Transformation Associated to a Functor**

**010Y**

**DEFINITION 11.5.4.1.1 ► THE NATURAL TRANSFORMATION ASSOCIATED TO A FUNCTOR**

Every functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  defines a natural transformation<sup>1</sup>

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} \times \mathcal{C} & \xrightarrow{F^{\text{op}} \times F} & \mathcal{D}^{\text{op}} \times \mathcal{D} \\ F^\dagger: \text{Hom}_{\mathcal{C}} \Rightarrow \text{Hom}_{\mathcal{D}} \circ (F^{\text{op}} \times F), & \swarrow \quad \searrow & \swarrow \quad \searrow \\ \text{Hom}_{\mathcal{C}} & \cong_{F^\dagger} & \text{Hom}_{\mathcal{D}} \\ & & \text{Sets}, \end{array}$$

called the **natural transformation associated to  $F$** , consisting of the collection

$$\left\{ F_{A,B}^\dagger: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B) \right\}_{(A,B) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})}$$

with

$$F_{A,B}^\dagger \stackrel{\text{def}}{=} F_{A,B}.$$

<sup>1</sup>This is the 1-categorical version of ?? of ??.

**PROOF 11.5.4.1.2 ► PROOF OF DEFINITION 11.5.4.1.1**

The naturality condition for  $F^\dagger$  is the requirement that for each morphism

$$(\phi, \psi): (X, Y) \rightarrow (A, B)$$

of  $\mathcal{C}^{\text{op}} \times \mathcal{C}$ , the diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(X, Y) & \xrightarrow{\phi^* \circ \psi_* = \psi_* \circ \phi^*} & \text{Hom}_{\mathcal{C}}(A, B) \\ F_{X,Y} \downarrow & & \downarrow F_{A,B} \\ \text{Hom}_{\mathcal{D}}(F_X, F_Y) & \xrightarrow[F(\phi)^* \circ F(\psi)_* = F(\psi)_* \circ F(\phi)^*]{} & \text{Hom}_{\mathcal{D}}(F_A, F_B), \end{array}$$

acting on elements as

$$\begin{array}{ccc} f & \longmapsto & \psi \circ f \circ \phi \\ \downarrow & & \downarrow \\ F(f) & \longmapsto & F(\psi) \circ F(f) \circ F(\phi) = F(\psi \circ f \circ \phi) \end{array}$$

commutes, which follows from the functoriality of  $F$ . 

**PROPOSITION 11.5.4.1.3 ► PROPERTIES OF NATURAL TRANSFORMATIONS ASSOCIATED TO FUNCTORS**

010Z

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  be functors.

0110

1. *Interaction With Natural Isomorphisms.* The following conditions are equivalent:

0111

(a) The natural transformation  $F^\dagger: \text{Hom}_{\mathcal{C}} \Rightarrow \text{Hom}_{\mathcal{D}} \circ (F^{\text{op}} \times F)$  associated to  $F$  is a natural isomorphism.

0112

(b) The functor  $F$  is fully faithful.

0113

2. *Interaction With Composition.* We have an equality of pasting diagrams

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} \times \mathcal{C} & \xrightarrow{F^{\text{op}} \times F} & \mathcal{D}^{\text{op}} \times \mathcal{D} & \xrightarrow{G^{\text{op}} \times G} & \mathcal{E}^{\text{op}} \times \mathcal{E} \\ \swarrow \text{Hom}_{\mathcal{C}} \quad \cong F^\dagger \quad \downarrow \text{Hom}_{\mathcal{D}} \quad \cong G^\dagger \quad \searrow \text{Hom}_{\mathcal{E}} \\ \text{Sets} & & & & \end{array} = \begin{array}{ccc} \mathcal{C}^{\text{op}} \times \mathcal{C} & \xrightarrow{(G \circ F)^{\text{op}} \times (G \circ F)} & \mathcal{E}^{\text{op}} \times \mathcal{E} \\ \swarrow \text{Hom}_{\mathcal{C}} \quad \cong (G \circ F)^\dagger \quad \searrow \text{Hom}_{\mathcal{E}} \\ \text{Sets} & & \end{array}$$

in  $\text{Cats}_2$ , i.e. we have

$$(G \circ F)^\dagger = (G^\dagger \star \text{id}_{F^{\text{op}} \times F}) \circ F^\dagger.$$

0114

3. *Interaction With Identities.* We have

$$\text{id}_C^\dagger = \text{id}_{\text{Hom}_{\mathcal{C}}(-1, -2)},$$

i.e. the natural transformation associated to  $\text{id}_C$  is the identity natural transformation of the functor  $\text{Hom}_{\mathcal{C}}(-1, -2)$ .

**PROOF 11.5.4.1.4 ► PROOF OF PROPOSITION 11.5.4.1.3**

Item 1: Interaction With Natural Isomorphisms

Omitted.

Item 2: Interaction With Composition

Omitted.

Item 3: Interaction With Identities

Omitted. 

## 0115 11.6 Conditions on Functors

### 0116 11.6.1 Faithful Functors

Let  $C$  and  $\mathcal{D}$  be categories.

#### 0117 DEFINITION 11.6.1.1 ► FAITHFUL FUNCTORS

A functor  $F: C \rightarrow \mathcal{D}$  is **faithful** if, for each  $A, B \in \text{Obj}(C)$ , the action on morphisms

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of  $F$  at  $(A, B)$  is injective.

#### 0118 PROPOSITION 11.6.1.2 ► PROPERTIES OF FAITHFUL FUNCTORS

Let  $F: C \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  be functors.

01U4 1. *Interaction With Composition.* If  $F$  and  $G$  are faithful, then so is  $G \circ F$ .

0119 2. *Interaction With Postcomposition.* The following conditions are equivalent:

011A (a) The functor  $F: C \rightarrow \mathcal{D}$  is faithful.

011B (b) For each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(X, C) \rightarrow \text{Fun}(X, \mathcal{D})$$

is faithful.

011C (c) The functor  $F: C \rightarrow \mathcal{D}$  is a representably faithful morphism in  $\text{Cats}_2$  in the sense of [Definition 14.1.1.1](#).

011D 3. *Interaction With Precomposition I.* Let  $F: C \rightarrow \mathcal{D}$  be a functor.

011E (a) If  $F$  is faithful, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(C, X)$$

*can fail* to be faithful.

011F (b) Conversely, if the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(C, X)$$

is faithful, then  $F$  *can fail* to be faithful.

011G

4. *Interaction With Precomposition II.* If  $F$  is essentially surjective, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is faithful.

011H

5. *Interaction With Precomposition III.* The following conditions are equivalent:

011J

- (a) For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is faithful.

011K

- (b) For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is conservative.

011L

- (c) For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is monadic.

011M

- (d) The functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a corepresentably faithful morphism in  $\text{Cats}_2$  in the sense of [Definition 14.2.1.1.1](#).

011N

- (e) The components

$$\eta_G: G \Rightarrow \text{Ran}_F(G \circ F)$$

of the unit

$$\eta: \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})} \Rightarrow \text{Ran}_F \circ F^*$$

of the adjunction  $F^* \dashv \text{Ran}_F$  are all monomorphisms.

011P

- (f) The components

$$\epsilon_G: \text{Lan}_F(G \circ F) \Rightarrow G$$

of the counit

$$\epsilon: \text{Lan}_F \circ F^* \Rightarrow \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})}$$

of the adjunction  $\text{Lan}_F \dashv F^*$  are all epimorphisms.

011Q

- (g) The functor  $F$  is dominant ([Definition 11.7.1.1.1](#)), i.e. every object of  $\mathcal{D}$  is a retract of some object in  $\text{Im}(F)$ :

- (★) For each  $B \in \text{Obj}(\mathcal{D})$ , there exist:

- An object  $A$  of  $\mathcal{C}$ ;
- A morphism  $s: B \rightarrow F(A)$  of  $\mathcal{D}$ ;
- A morphism  $r: F(A) \rightarrow B$  of  $\mathcal{D}$ ;

such that  $r \circ s = \text{id}_B$ .

**PROOF 11.6.1.1.3 ▶ PROOF OF PROPOSITION 11.6.1.1.2****Item 1: Interaction With Composition**

Since the map

$$(G \circ F)_{A,B} : \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(G_{F_A}, G_{F_B}),$$

defined as the composition

$$\text{Hom}_C(A, B) \xrightarrow{F_{A,B}} \text{Hom}_{\mathcal{D}}(F_A, F_B) \xrightarrow{G_{F(A),F(B)}} \text{Hom}_{\mathcal{D}}(G_{F_A}, G_{F_B}),$$

is a composition of injective functions, it follows from ?? that it is also injective. Therefore  $G \circ F$  is faithful.

**Item 2: Interaction With Postcomposition**

Omitted.

**Item 3: Interaction With Precomposition I**

See [MSE 733163] for Item 3a. Item 3b follows from Item 4 and the fact that there are essentially surjective functors that are not faithful.

**Item 4: Interaction With Precomposition II**

Omitted, but see [https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp\\_fully\\_faithful.html](https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp_fully_faithful.html) for a formalised proof.

**Item 5: Interaction With Precomposition III**

We claim Items 5a to 5g are equivalent:

- *Items 5a and 5d Are Equivalent:* This is true by the definition of corepresentably faithful morphism; see Definition 14.2.1.1.1.
- *Items 5a to 5c and 5g Are Equivalent:* See [Adá+01, Proposition 4.1] or alternatively [Fre09, Lemmas 3.1 and 3.2] for the equivalence between Items 5a and 5g.
- *Items 5a, 5e and 5f Are Equivalent:* See ?? of ??.

This finishes the proof. 

**011R 11.6.2 Full Functors**

Let  $C$  and  $\mathcal{D}$  be categories.

011S

**DEFINITION 11.6.2.1.1 ► FULL FUNCTORS**

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **full** if, for each  $A, B \in \text{Obj}(\mathcal{C})$ , the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of  $F$  at  $(A, B)$  is surjective.

011T

**PROPOSITION 11.6.2.1.2 ► PROPERTIES OF FULL FUNCTORS**

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  be functors.

01U5

1. *Interaction With Composition.* If  $F$  and  $G$  are full, then so is  $G \circ F$ .

01U6

2. *Interaction With Postcomposition I.* If  $F$  is full, then the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

*can fail* to be full.

01U7

3. *Interaction With Postcomposition II.* If, for each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is full, then  $F$  is also full.

011Y

4. *Interaction With Precomposition I.* If  $F$  is full, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

*can fail* to be full.

011Z

5. *Interaction With Precomposition II.* If, for each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is full, then  $F$  *can fail* to be full.

0120

6. *Interaction With Precomposition III.* If  $F$  is essentially surjective and full, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is full (and also faithful by [Item 4 of Proposition 11.6.1.1.2](#)).

0121

7. *Interaction With Precomposition IV.* The following conditions are equivalent:

0122

(a) For each  $X \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(C, X)$$

is full.

0123

(b) The functor  $F: C \rightarrow \mathcal{D}$  is a corepresentably full morphism in  $\text{Cats}_2$  in the sense of [Definition 14.2.1.1.1](#).

0124

(c) The components

$$\eta_G: G \Rightarrow \text{Ran}_F(G \circ F)$$

of the unit

$$\eta: \text{id}_{\text{Fun}(\mathcal{D}, X)} \Rightarrow \text{Ran}_F \circ F^*$$

of the adjunction  $F^* \dashv \text{Ran}_F$  are all retractions/split epimorphisms.

0125

(d) The components

$$\epsilon_G: \text{Lan}_F(G \circ F) \Rightarrow G$$

of the counit

$$\epsilon: \text{Lan}_F \circ F^* \Rightarrow \text{id}_{\text{Fun}(\mathcal{D}, X)}$$

of the adjunction  $\text{Lan}_F \dashv F^*$  are all sections/split monomorphisms.

0126

(e) For each  $B \in \text{Obj}(\mathcal{D})$ , there exist:

- An object  $A_B$  of  $C$ ;
- A morphism  $s_B: B \rightarrow F(A_B)$  of  $\mathcal{D}$ ;
- A morphism  $r_B: F(A_B) \rightarrow B$  of  $\mathcal{D}$ ;

satisfying the following condition:

(★) For each  $A \in \text{Obj}(C)$  and each pair of morphisms

$$r: F(A) \rightarrow B,$$

$$s: B \rightarrow F(A)$$

of  $\mathcal{D}$ , we have

$$[(A_B, s_B, r_B)] = [(A, s, r \circ s_B \circ r_B)]$$

$$\text{in } \int^{A \in C} h_{F_A}^{B'} \times h_B^{F_A}.$$

**PROOF 11.6.2.1.3 ▶ PROOF OF PROPOSITION 11.6.2.1.2****Item 1: Interaction With Composition**

Since the map

$$(G \circ F)_{A,B} : \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(G_{F_A}, G_{F_B}),$$

defined as the composition

$$\text{Hom}_C(A, B) \xrightarrow{F_{A,B}} \text{Hom}_{\mathcal{D}}(F_A, F_B) \xrightarrow{G_{F(A),F(B)}} \text{Hom}_{\mathcal{D}}(G_{F_A}, G_{F_B}),$$

is a composition of surjective functions, it follows from ?? that it is also surjective. Therefore  $G \circ F$  is full.

**Item 2: Interaction With Postcomposition I**

We follow the proof (completely formalised in cubical Agda!) given by Naïm Camille Favier in [favier:postcompose-not-full]. Let  $C$  be the category where:

- *Objects.* We have  $\text{Obj}(C) = \{A, B\}$ .
- *Morphisms.* We have

$$\begin{aligned}\text{Hom}_C(A, A) &= \{e_A, \text{id}_A\}, \\ \text{Hom}_C(B, B) &= \{e_B, \text{id}_B\}, \\ \text{Hom}_C(A, B) &= \{f, g\}, \\ \text{Hom}_C(B, A) &= \emptyset.\end{aligned}$$

- *Composition.* The nontrivial compositions in  $C$  are the following:

$$\begin{aligned}e_A \circ e_A &= \text{id}_A, & f \circ e_A &= g, & e_B \circ f &= f, \\ e_B \circ e_B &= \text{id}_B, & g \circ e_A &= f, & e_B \circ g &= g.\end{aligned}$$

We may picture  $C$  as follows:

$$A \begin{array}{c} \nearrow e_A \\ \xrightarrow[g]{f} \\ \searrow e_B \end{array} B$$

Next, let  $\mathcal{D}$  be the walking arrow category  $\mathbb{1}$  of [Definition 11.2.5.1.1](#) and let  $F: C \rightarrow \mathbb{1}$  be the functor given on objects by

$$F(A) = 0,$$

$$F(B) = 1$$

and on non-identity morphisms by

$$\begin{aligned} F(f) &= f_{01}, & F(e_A) &= \text{id}_0, \\ F(g) &= f_{01}, & F(e_B) &= \text{id}_1. \end{aligned}$$

Finally, let  $X = B\mathbb{Z}/2$  be the walking involution and let  $\iota_A, \iota_B : B\mathbb{Z}/2 \rightrightarrows C$  be the inclusion functors from  $B\mathbb{Z}/2$  to  $C$  with

$$\begin{aligned} \iota_A(\bullet) &= A, \\ \iota_B(\bullet) &= B. \end{aligned}$$

Since every morphism in  $\mathbb{1}$  has a preimage in  $C$  by  $F$ , the functor  $F$  is full. Now, for  $F_*$  to be full, the map

$$\begin{aligned} F_*|_{\iota_A, \iota_B} : \text{Nat}(\iota_A, \iota_B) &\longrightarrow \text{Nat}(F \circ \iota_A, F \circ \iota_B) \\ \alpha &\longmapsto \text{id}_F \star \alpha \end{aligned}$$

would need to be surjective. However, as we will show next, we have

$$\begin{aligned} \text{Nat}(\iota_A, \iota_B) &= \emptyset, \\ \text{Nat}(F \circ \iota_A, F \circ \iota_B) &\cong \text{pt}, \end{aligned}$$

so this is impossible:

- *Proof of  $\text{Nat}(\iota_A, \iota_B) = \emptyset$ :* A natural transformation  $\alpha : \iota_A \Rightarrow \iota_B$  consists of a morphism

$$\alpha : \underbrace{\iota_A(\bullet)}_{=A} \rightarrow \underbrace{\iota_B(\bullet)}_{=B}$$

in  $C$  making the diagram

$$\begin{array}{ccc} \iota_A(\bullet) & \xrightarrow{\iota_A(e)} & \iota_A(\bullet) \\ \alpha \downarrow & & \downarrow \alpha \\ \iota_B(\bullet) & \xrightarrow{\iota_B(e)} & \iota_B(\bullet) \end{array}$$

commute for each  $e \in \text{Hom}_{B\mathbb{Z}/2}(\bullet, \bullet) \cong \mathbb{Z}/2$ . We have two cases:

02BY

1. If  $\alpha = f$ , the naturality diagram for the unique nonidentity element of  $\mathbb{Z}/2$  is given by

$$\begin{array}{ccc} A & \xrightarrow{e_A} & A \\ f \downarrow & & \downarrow f \\ B & \xrightarrow{e_B} & B. \end{array}$$

However,  $e_B \circ f = f$  and  $f \circ e_A = g$ , so this diagram does not commute.

02BZ

2. If  $\alpha = g$ , the naturality diagram for the unique nonidentity element of  $\mathbb{Z}/2$  is given by

$$\begin{array}{ccc} A & \xrightarrow{e_A} & A \\ g \downarrow & & \downarrow g \\ B & \xrightarrow{e_B} & B. \end{array}$$

However,  $e_B \circ g = g$  and  $g \circ e_A = f$ , so this diagram does not commute.

As a result, there are no natural transformations from  $\iota_A$  to  $\iota_B$ .

- *Proof of  $\text{Nat}(F \circ \iota_A, F \circ \iota_B) \cong pt$ :* A natural transformation

$$\beta : F \circ \iota_A \Rightarrow F \circ \iota_B$$

consists of a morphism

$$\beta : \underbrace{[F \circ \iota_A](\bullet)}_{=0} \rightarrow \underbrace{[F \circ \iota_B](\bullet)}_{=1}$$

in  $\mathbb{1}$  making the diagram

$$\begin{array}{ccc} [F \circ \iota_A](\bullet) & \xrightarrow{[F \circ \iota_A](e)} & [F \circ \iota_A](\bullet) \\ \beta \downarrow & & \downarrow \beta \\ [F \circ \iota_B](\bullet) & \xrightarrow{[F \circ \iota_B](e)} & [F \circ \iota_B](\bullet) \end{array}$$

commute for each  $e \in \text{Hom}_{\mathcal{B}\mathbb{Z}/2}(\bullet, \bullet) \cong \mathbb{Z}/2$ . Since the only morphism from 0 to 1 in  $\mathbb{1}$  is  $f_{01}$ , we must have  $\beta = f_{01}$  if such a

transformation were to exist, and in fact it indeed does, as in this case the naturality diagram above becomes

$$\begin{array}{ccc} 0 & \xrightarrow{\text{id}_0} & 0 \\ f_{01} \downarrow & & \downarrow f_{01} \\ 1 & \xrightarrow{\text{id}_1} & 1 \end{array}$$

for each  $e \in \mathbb{Z}/2$ , and this diagram indeed commutes, making  $\beta$  into a natural transformation.

This finishes the proof.

#### Item 3: Interaction With Postcomposition II

Taking  $X = \text{pt}$ , it follows by assumption that the functor

$$F_* : \text{Fun}(\text{pt}, C) \rightarrow \text{Fun}(\text{pt}, \mathcal{D})$$

is full. However, by [Item 5 of Proposition 11.10.1.1.2](#), we have isomorphisms of categories

$$\begin{aligned} \text{Fun}(\text{pt}, C) &\cong C, \\ \text{Fun}(\text{pt}, \mathcal{D}) &\cong \mathcal{D} \end{aligned}$$

and the diagram

$$\begin{array}{ccc} \text{Fun}(\text{pt}, C) & \xrightarrow{F_*} & \text{Fun}(\text{pt}, \mathcal{D}) \\ \downarrow \wr & & \downarrow \wr \\ C & \xrightarrow{F} & \mathcal{D} \end{array}$$

commutes. It then follows from [Item 1](#) that  $F$  is full.

#### Item 4: Interaction With Precomposition I

Omitted.

#### Item 5: Interaction With Precomposition II

See [\[BS10, p. 47\]](#).

#### Item 6: Interaction With Precomposition III

Omitted, but see [https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp\\_fully\\_faithful.html](https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp_fully_faithful.html) for a formalised proof.

Item 7: Interaction With Precomposition IV

We claim [Items 7a](#) to [7e](#) are equivalent:

- [Items 7a and 7b Are Equivalent](#): This is true by the definition of corepresentably full morphism; see [Definition 14.2.2.1.1](#).
- [Items 7a, 7c and 7d Are Equivalent](#): See ?? of ??.
- [Items 7a and 7e Are Equivalent](#): See [[Adá+01](#), Item (b) of Remark 4.3].

This finishes the proof. □

0127

**QUESTION 11.6.2.1.4 ► BETTER CHARACTERISATIONS OF FUNCTORS WITH FULL PRECOMPOSITION**

[Item 7](#) of [Proposition 11.6.2.1.2](#) gives a characterisation of the functors  $F$  for which  $F^*$  is full, but the characterisations given there are really messy. Are there better ones?

This question also appears as [[MO 468121b](#)].

0128

### 11.6.3 Fully Faithful Functors

Let  $C$  and  $\mathcal{D}$  be categories.

0129

**DEFINITION 11.6.3.1.1 ► FULLY FAITHFUL FUNCTORS**

A functor  $F: C \rightarrow \mathcal{D}$  is **fully faithful** if  $F$  is full and faithful, i.e. if, for each  $A, B \in \text{Obj}(C)$ , the action on morphisms

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of  $F$  at  $(A, B)$  is bijective.

012A

**PROPOSITION 11.6.3.1.2 ► PROPERTIES OF FULLY FAITHFUL FUNCTORS**

Let  $F: C \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  be functors.

012B

1. *Characterisations.* The following conditions are equivalent:

012C

- (a) The functor  $F$  is fully faithful.

012D

(b) We have a pullback square

$$\begin{array}{ccc} \text{Arr}(C) & \xrightarrow{\text{Arr}(F)} & \text{Arr}(\mathcal{D}) \\ \text{Arr}(C) \cong (C \times C) \times_{\mathcal{D} \times \mathcal{D}} \text{Arr}(\mathcal{D}), & \text{src} \times \text{tgt} \downarrow & \downarrow \text{src} \times \text{tgt} \\ C \times C & \xrightarrow{F \times F} & \mathcal{D} \times \mathcal{D} \end{array}$$

in  $\text{Cats}$ .

012U8

2. *Interaction With Composition.* If  $F$  and  $G$  are fully faithful, then so is  $G \circ F$ .

012E

3. *Conservativity.* If  $F$  is fully faithful, then  $F$  is conservative.

012F

4. *Essential Injectivity.* If  $F$  is fully faithful, then  $F$  is essentially injective.

012G

5. *Interaction With Co/Limits.* If  $F$  is fully faithful, then  $F$  reflects co/limits.

012H

6. *Interaction With Postcomposition.* The following conditions are equivalent:

012J

- (a) The functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is fully faithful.

012K

- (b) For each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(X, \mathcal{C}) \rightarrow \text{Fun}(X, \mathcal{D})$$

is fully faithful.

012L

- (c) The functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a representably fully faithful morphism in  $\text{Cats}_2$  in the sense of [Definition 14.1.3.1.1](#).

012M

7. *Interaction With Precomposition I.* If  $F$  is fully faithful, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(\mathcal{C}, X)$$

*can fail* to be fully faithful.

012N

8. *Interaction With Precomposition II.* If the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(\mathcal{C}, X)$$

is fully faithful, then  $F$  *can fail* to be fully faithful (and in fact it can also fail to be either full or faithful).

012P

9. *Interaction With Precomposition III.* If  $F$  is essentially surjective and full, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is fully faithful.

012Q

10. *Interaction With Precomposition IV.* The following conditions are equivalent:

012R

- (a) For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is fully faithful.

012S

- (b) The precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \text{Sets}) \rightarrow \text{Fun}(\mathcal{C}, \text{Sets})$$

is fully faithful.

012T

- (c) The functor

$$\text{Lan}_F: \text{Fun}(\mathcal{C}, \text{Sets}) \rightarrow \text{Fun}(\mathcal{D}, \text{Sets})$$

is fully faithful.

012U

- (d) The functor  $F$  is a corepresentably fully faithful morphism in  $\text{Cats}_2$  in the sense of [Definition 14.2.3.1.1](#).

012V

- (e) The functor  $F$  is absolutely dense.

012W

- (f) The components

$$\eta_G: G \Rightarrow \text{Ran}_F(G \circ F)$$

of the unit

$$\eta: \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})} \Rightarrow \text{Ran}_F \circ F^*$$

of the adjunction  $F^* \dashv \text{Ran}_F$  are all isomorphisms.

012X

- (g) The components

$$\epsilon_G: \text{Lan}_F(G \circ F) \Rightarrow G$$

of the counit

$$\epsilon: \text{Lan}_F \circ F^* \Rightarrow \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})}$$

of the adjunction  $\text{Lan}_F \dashv F^*$  are all isomorphisms.

012Y

(h) The natural transformation

$$\alpha: \text{Lan}_{h_F}(h^F) \Rightarrow h$$

with components

$$\alpha_{B',B}: \int^{A \in C} h_{F_A}^{B'} \times h_B^{F_A} \rightarrow h_B^{B'}$$

given by

$$\alpha_{B',B}([\phi, \psi]) = \psi \circ \phi$$

is a natural isomorphism.

012Z

(i) For each  $B \in \text{Obj}(\mathcal{D})$ , there exist:

- An object  $A_B$  of  $C$ ;
- A morphism  $s_B: B \rightarrow F(A_B)$  of  $\mathcal{D}$ ;
- A morphism  $r_B: F(A_B) \rightarrow B$  of  $\mathcal{D}$ ;

satisfying the following conditions:

0130

- i. The triple  $(F(A_B), r_B, s_B)$  is a retract of  $B$ , i.e. we have  $r_B \circ s_B = \text{id}_B$ .

0131

- ii. For each morphism  $f: B' \rightarrow B$  of  $\mathcal{D}$ , we have

$$[(A_B, s_{B'}, f \circ r_{B'})] = [(A_B, s_B \circ f, r_B)]$$

$$\text{in } \int^{A \in C} h_{F_A}^{B'} \times h_B^{F_A}.$$

**PROOF 11.6.3.1.3 ► PROOF OF PROPOSITION 11.6.3.1.2**

Item 1: Characterisations

Omitted.

Item 2: Interaction With Composition

Since the map

$$(G \circ F)_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(G_{F_A}, G_{F_B}),$$

defined as the composition

$$\text{Hom}_C(A, B) \xrightarrow{F_{A,B}} \text{Hom}_{\mathcal{D}}(F_A, F_B) \xrightarrow{G_{F(A), F(B)}} \text{Hom}_{\mathcal{D}}(G_{F_A}, G_{F_B}),$$

is a composition of bijective functions, it follows from ?? that it is also bijective. Therefore  $G \circ F$  is fully faithful.

**Item 3: Conservativity**

This is a repetition of [Item 2 of Proposition 11.6.4.1.2](#), and is proved there.

**Item 4: Essential Injectivity**

Omitted.

**Item 5: Interaction With Co/Limits**

Omitted.

**Item 6: Interaction With Postcomposition**

This follows from [Item 2 of Proposition 11.6.1.1.2](#) and [Item 1 of Proposition 11.6.2.1.2](#).

**Item 7: Interaction With Precomposition I**

See [[MSE 733161](#)] for an example of a fully faithful functor whose precomposition with which fails to be full.

**Item 8: Interaction With Precomposition II**

See [[MSE 749304](#), Item 3].

**Item 9: Interaction With Precomposition III**

Omitted, but see [https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp\\_fully\\_faithful.html](https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp_fully_faithful.html) for a formalised proof.

**Item 10: Interaction With Precomposition IV**

We claim [Items 10a](#) to [10i](#) are equivalent:

- [Items 10a and 10d Are Equivalent](#): This is true by the definition of corepresentably fully faithful morphism; see [Definition 14.2.3.1.1](#).
- [Items 10a, 10f and 10g Are Equivalent](#): See ?? of ??.
- [Items 10a to 10c Are Equivalent](#): This follows from [[Low15](#), Proposition A.1.5].
- [Items 10a, 10e, 10h and 10i Are Equivalent](#): See [[Fre09](#), Theorem 4.1] and [[Adá+01](#), Theorem 1.1].

This finishes the proof. 

## 0132 11.6.4 Conservative Functors

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

0133

**DEFINITION 11.6.4.1.1 ► CONSERVATIVE FUNCTORS**

A functor  $F: C \rightarrow \mathcal{D}$  is **conservative** if it satisfies the following condition:<sup>1</sup>

- (★) For each  $f \in \text{Mor}(C)$ , if  $F(f)$  is an isomorphism in  $\mathcal{D}$ , then  $f$  is an isomorphism in  $C$ .

<sup>1</sup>Slogan: A functor  $F$  is **conservative** if it reflects isomorphisms.

0134

**PROPOSITION 11.6.4.1.2 ► PROPERTIES OF CONSERVATIVE FUNCTORS**

Let  $F: C \rightarrow \mathcal{D}$  be a functor.

0135

1. *Characterisations.* The following conditions are equivalent:

0136

- (a) The functor  $F$  is conservative.

0137

- (b) For each  $f \in \text{Mor}(C)$ , the morphism  $F(f)$  is an isomorphism in  $\mathcal{D}$  iff  $f$  is an isomorphism in  $C$ .

0138

2. *Interaction With Fully Faithfulness.* Every fully faithful functor is conservative.

0139

3. *Interaction With Precomposition.* The following conditions are equivalent:

013A

- (a) For each  $X \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(C, X)$$

is conservative.

013B

- (b) The equivalent conditions of [Item 5 of Proposition 11.6.1.1.2](#) are satisfied.

**PROOF 11.6.4.1.3 ► PROOF OF PROPOSITION 11.6.4.1.2****Item 1: Characterisations**

This follows from [Item 1 of Proposition 11.5.1.1.8](#).

**Item 2: Interaction With Fully Faithfulness**

Let  $F: C \rightarrow \mathcal{D}$  be a fully faithful functor, let  $f: A \rightarrow B$  be a morphism of  $C$ , and suppose that  $F_f$  is an isomorphism. We have

$$\begin{aligned} F(\text{id}_B) &= \text{id}_{F(B)} \\ &= F(f) \circ F(f)^{-1} \end{aligned}$$

$$= F(f \circ f^{-1}).$$

Similarly,  $F(\text{id}_A) = F(f^{-1} \circ f)$ . But since  $F$  is fully faithful, we must have

$$\begin{aligned} f \circ f^{-1} &= \text{id}_B, \\ f^{-1} \circ f &= \text{id}_A, \end{aligned}$$

showing  $f$  to be an isomorphism. Thus  $F$  is conservative. 

013C

#### QUESTION 11.6.4.1.4 ► CHARACTERISATIONS OF FUNCTORS WITH CONSERVATIVE PRE-/POSTCOMPOSITION

Is there a characterisation of functors  $F: C \rightarrow \mathcal{D}$  satisfying the following condition:

- (★) For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is conservative?

This question also appears as [MO 468121a].

013D 11.6.5 Essentially Injective Functors

Let  $C$  and  $\mathcal{D}$  be categories.

013E

#### DEFINITION 11.6.5.1.1 ► ESSENTIALLY INJECTIVE FUNCTORS

A functor  $F: C \rightarrow \mathcal{D}$  is **essentially injective** if it satisfies the following condition:

- (★) For each  $A, B \in \text{Obj}(C)$ , if  $F(A) \cong F(B)$ , then  $A \cong B$ .

013F

#### QUESTION 11.6.5.1.2 ► CHARACTERISATIONS OF FUNCTORS WITH ESSENTIALLY INJECTIVE PRE/POSTCOMPOSITION

Is there a characterisation of functors  $F: C \rightarrow \mathcal{D}$  such that:

013G

1. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is essentially injective, i.e. if  $\phi \circ F \cong \psi \circ F$ , then  $\phi \cong \psi$  for all functors  $\phi$  and  $\psi$ ?

013H

2. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is essentially injective, i.e. if  $F \circ \phi \cong F \circ \psi$ , then  $\phi \cong \psi$ ?

This question also appears as [MO 468121a].

### 013J 11.6.6 Essentially Surjective Functors

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

013K

#### DEFINITION 11.6.6.1.1 ► ESSENTIALLY SURJECTIVE FUNCTORS

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **essentially surjective**<sup>1</sup> if it satisfies the following condition:

- (★) For each  $D \in \text{Obj}(\mathcal{D})$ , there exists some object  $A$  of  $\mathcal{C}$  such that  $F(A) \cong D$ .

<sup>1</sup>Further Terminology: Also called an **eso** functor, meaning *essentially surjective on objects*.

013L

#### QUESTION 11.6.6.1.2 ► CHARACTERISATIONS OF FUNCTORS WITH ESSENTIALLY SURJECTIVE PRE/POSTCOMPOSITION

Is there a characterisation of functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  such that:

013M

1. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is essentially surjective?

013N

2. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is essentially surjective?

This question also appears as [MO 468121a].

**013P 11.6.7 Equivalences of Categories**

013Q

**DEFINITION 11.6.7.1.1 ► EQUIVALENCES OF CATEGORIES**

Let  $C$  and  $\mathcal{D}$  be categories.

013R

1. An **equivalence of categories** between  $C$  and  $\mathcal{D}$  consists of a pair of functors

$$\begin{aligned} F: C &\rightarrow \mathcal{D}, \\ G: \mathcal{D} &\rightarrow C \end{aligned}$$

together with natural isomorphisms

$$\begin{aligned} \eta: \text{id}_C &\xrightarrow{\sim} G \circ F, \\ \epsilon: F \circ G &\xrightarrow{\sim} \text{id}_{\mathcal{D}}. \end{aligned}$$

013S

2. An **adjoint equivalence of categories** between  $C$  and  $\mathcal{D}$  is an equivalence  $(F, G, \eta, \epsilon)$  between  $C$  and  $\mathcal{D}$  which is also an adjunction.

013T

**PROPOSITION 11.6.7.1.2 ► PROPERTIES OF EQUIVALENCES OF CATEGORIES**

Let  $F: C \rightarrow \mathcal{D}$  be a functor.

013U

1. *Characterisations.* If  $C$  and  $\mathcal{D}$  are small<sup>1</sup>, then the following conditions are equivalent:<sup>2</sup>

013V

(a) The functor  $F$  is an equivalence of categories.

013W

(b) The functor  $F$  is fully faithful and essentially surjective.

013X

(c) The induced functor

$$\uparrow F\text{Sk}(C): \text{Sk}(C) \rightarrow \text{Sk}(\mathcal{D})$$

is an *isomorphism* of categories.

013Y

- (d) For each  $X \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(C, X)$$

is an equivalence of categories.

013Z

- (e) For each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(X, C) \rightarrow \text{Fun}(X, \mathcal{D})$$

is an equivalence of categories.

0140

2. *Two-Out-of-Three.* Let

$$\begin{array}{ccc} C & \xrightarrow{G \circ F} & \mathcal{E} \\ & \searrow F & \nearrow G \\ & \mathcal{D} & \end{array}$$

be a diagram in Cats. If two out of the three functors among  $F$ ,  $G$ , and  $G \circ F$  are equivalences of categories, then so is the third.

0141

3. *Stability Under Composition.* Let

$$\begin{array}{ccccc} C & \xleftarrow[G]{F} & \mathcal{D} & \xleftarrow[G']{F'} & \mathcal{E} \end{array}$$

be a diagram in Cats. If  $(F, G)$  and  $(F', G')$  are equivalences of categories, then so is their composite  $(F' \circ F, G' \circ G)$ .

0142

4. *Equivalences vs. Adjoint Equivalences.* Every equivalence of categories can be promoted to an adjoint equivalence.<sup>3</sup>

0143

5. *Interaction With Groupoids.* If  $C$  and  $\mathcal{D}$  are groupoids, then the following conditions are equivalent:

0144

(a) The functor  $F$  is an equivalence of groupoids.

0145

(b) The following conditions are satisfied:

0146

i. The functor  $F$  induces a bijection

$$\pi_0(F): \pi_0(C) \rightarrow \pi_0(\mathcal{D})$$

of sets.

0147

ii. For each  $A \in \text{Obj}(C)$ , the induced map

$$F_{x,x}: \text{Aut}_C(A) \rightarrow \text{Aut}_{\mathcal{D}}(F_A)$$

is an isomorphism of groups.

<sup>1</sup>Otherwise there will be size issues. One can also work with large categories and universes, or require  $F$  to be *constructively* essentially surjective; see [MSE 1465107].

<sup>2</sup>In ZFC, the equivalence between Item 1a and Item 1b is equivalent to the axiom of choice; see [MO 119454].

In Univalent Foundations, this is true without requiring neither the axiom of choice nor the law of excluded middle.

<sup>3</sup>More precisely, we can promote an equivalence of categories  $(F, G, \eta, \epsilon)$  to adjoint equivalences  $(F, G, \eta', \epsilon)$  and  $(F, G, \eta, \epsilon')$ .

## PROOF 11.6.7.1.3 ► PROOF OF PROPOSITION 11.6.7.1.2

## Item 1: Characterisations

We claim that **Items 1a** to **1e** are indeed equivalent:

1. *Item 1a*  $\implies$  *Item 1b*: Omitted.
2. *Item 1b*  $\implies$  *Item 1a*: Since  $F$  is essentially surjective and  $C$  and  $\mathcal{D}$  are small, we can choose, using the axiom of choice, for each  $B \in \text{Obj}(\mathcal{D})$ , an object  $j_B$  of  $C$  and an isomorphism  $i_B: B \rightarrow F_{j_B}$  of  $\mathcal{D}$ .  
Since  $F$  is fully faithful, we can extend the assignment  $B \mapsto j_B$  to a *unique* functor  $j: \mathcal{D} \rightarrow C$  such that the isomorphisms  $i_B: B \rightarrow F_{j_B}$  assemble into a natural isomorphism  $\eta: \text{id}_{\mathcal{D}} \xrightarrow{\sim} F \circ j$ , with a similar natural isomorphism  $\epsilon: \text{id}_C \xrightarrow{\sim} j \circ F$ . Hence  $F$  is an equivalence.
3. *Item 1a*  $\implies$  *Item 1c*: This follows from **Item 4** of **Proposition 11.1.3.1.3**.
4. *Item 1c*  $\implies$  *Item 1a*: Omitted.
5. *Items 1a, 1d and 1e Are Equivalent*: This follows from ??.

This finishes the proof of **Item 1**.

## Item 2: Two-Out-of-Three

Omitted.

## Item 3: Stability Under Composition

Omitted.

## Item 4: Equivalences vs. Adjoint Equivalences

See [Rie16, Proposition 4.4.5].

## Item 5: Interaction With Groupoids

See [nLa25a, Proposition 4.4].



## 0148 11.6.8 Isomorphisms of Categories

## 0149 DEFINITION 11.6.8.1.1 ► ISOMORPHISMS OF CATEGORIES

An **isomorphism of categories** is a pair of functors

$$\begin{aligned} F: C &\rightarrow \mathcal{D}, \\ G: \mathcal{D} &\rightarrow C \end{aligned}$$

such that we have

$$G \circ F = \text{id}_C,$$

$$F \circ G = \text{id}_{\mathcal{D}}.$$

014A

**EXAMPLE 11.6.8.1.2 ► EQUIVALENT BUT NON-ISOMORPHIC CATEGORIES**

Categories can be equivalent but non-isomorphic. For example, the category consisting of two isomorphic objects is equivalent to pt, but not isomorphic to it.

014B

**PROPOSITION 11.6.8.1.3 ► PROPERTIES OF ISOMORPHISMS OF CATEGORIES**

Let  $F: C \rightarrow \mathcal{D}$  be a functor.

014C

1. *Characterisations.* If  $C$  and  $\mathcal{D}$  are small, then the following conditions are equivalent:

014D

(a) The functor  $F$  is an isomorphism of categories.

014E

(b) The functor  $F$  is fully faithful and bijective on objects.

014F

(c) For each  $X \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(C, X)$$

is an isomorphism of categories.

014G

- (d) For each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(X, C) \rightarrow \text{Fun}(X, \mathcal{D})$$

is an isomorphism of categories.

**PROOF 11.6.8.1.4 ► PROOF OF PROPOSITION 11.6.8.1.3****Item 1: Characterisations**

We claim that **Items 1a** to **1d** are indeed equivalent:

1. *Items 1a and 1b Are Equivalent:* Omitted, but similar to **Item 1** of Proposition 11.6.7.1.2.
2. *Items 1a, 1c and 1d Are Equivalent:* This follows from ??.

This finishes the proof. 

## 014H 11.7 More Conditions on Functors

### 014J 11.7.1 Dominant Functors

Let  $C$  and  $\mathcal{D}$  be categories.

#### 014K DEFINITION 11.7.1.1 ► DOMINANT FUNCTORS

A functor  $F: C \rightarrow \mathcal{D}$  is **dominant** if every object of  $\mathcal{D}$  is a retract of some object in  $\text{Im}(F)$ , i.e.:

(★) For each  $B \in \text{Obj}(\mathcal{D})$ , there exist:

- An object  $A$  of  $C$ ;
- A morphism  $r: F(A) \rightarrow B$  of  $\mathcal{D}$ ;
- A morphism  $s: B \rightarrow F(A)$  of  $\mathcal{D}$ ;

such that we have

$$\begin{array}{ccc} B & \xrightarrow{s} & F(A) \\ r \circ s = \text{id}_B, & \searrow \text{id}_B & \downarrow r \\ & & B. \end{array}$$

#### 014L PROPOSITION 11.7.1.2 ► PROPERTIES OF DOMINANT FUNCTORS

Let  $F, G: C \Rightarrow \mathcal{D}$  be functors and let  $I: \mathcal{X} \rightarrow C$  be a functor.

014M 1. *Interaction With Right Whiskering.* If  $I$  is full and dominant, then the map

$$-\star \text{id}_I: \text{Nat}(F, G) \rightarrow \text{Nat}(F \circ I, G \circ I)$$

is a bijection.

014N 2. *Interaction With Adjunctions.* Let  $(F, G): C \rightleftarrows \mathcal{D}$  be an adjunction.

014P (a) If  $F$  is dominant, then  $G$  is faithful.

014Q (b) The following conditions are equivalent:

- The functor  $G$  is full.
- The restriction

$$\upharpoonright G\text{Im}_F: \text{Im}(F) \rightarrow C$$

of  $G$  to  $\text{Im}(F)$  is full.

**PROOF 11.7.1.1.3 ▶ PROOF OF PROPOSITION 11.7.1.1.2**

Item 1: Interaction With Right Whiskering

See [DFH75, Proposition 1.4].

Item 2: Interaction With Adjunctions

See [DFH75, Proposition 1.7]. 

**QUESTION 11.7.1.1.4 ▶ CHARACTERISATIONS OF FUNCTORS WITH DOMINANT PRE/POST-COMPOSITION**

014T

Is there a characterisation of functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  such that:

014U

- For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is dominant?

014V

- For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is dominant?

This question also appears as [MO 468121a].

014W **11.7.2 Monomorphisms of Categories**

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

014X

**DEFINITION 11.7.2.1.1 ▶ MONOMORPHISMS OF CATEGORIES**

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a **monomorphism of categories** if it is a monomorphism in  $\text{Cats}$  (see ??).

014Y

**PROPOSITION 11.7.2.1.2 ▶ PROPERTIES OF MONOMORPHISMS OF CATEGORIES**

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a functor.

014Z

- Characterisations.* The following conditions are equivalent:

0150

- The functor  $F$  is a monomorphism of categories.

0151

- (b) The functor  $F$  is injective on objects and morphisms, i.e.  $F$  is injective on objects and the map

$$F: \text{Mor}(\mathcal{C}) \rightarrow \text{Mor}(\mathcal{D})$$

is injective.

**PROOF 11.7.2.1.3 ► PROOF OF PROPOSITION 11.7.2.1.2**

**Item 1: Characterisations**

Omitted. 

0152

**QUESTION 11.7.2.1.4 ► CHARACTERISATIONS OF FUNCTORS WITH MONIC PRE/POSTCOMPOSITION**

Is there a characterisation of functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  such that:

0153

1. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is a monomorphism of categories?

0154

2. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is a monomorphism of categories?

This question also appears as [MO 468121a].

**0155 11.7.3 Epimorphisms of Categories**

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

0156

**DEFINITION 11.7.3.1.1 ► EPIMORPHISMS OF CATEGORIES**

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a **epimorphism of categories** if it is a epimorphism in  $\text{Cats}$  (see ??).

0157

**PROPOSITION 11.7.3.1.2 ► PROPERTIES OF EPIMORPHISMS OF CATEGORIES**

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a functor.


0158

1. *Characterisations.* The following conditions are equivalent:<sup>1</sup>

0159

(a) The functor  $F$  is an epimorphism of categories.

015A

(b) For each morphism  $f: A \rightarrow B$  of  $\mathcal{D}$ , we have a diagramin  $\mathcal{D}$  satisfying the following conditions:

015B

i. We have  $f = \alpha_0 \circ \phi_1$ .

015C

ii. We have  $f = \psi_m \circ \alpha_{2m}$ .

015D

iii. For each  $0 \leq i \leq 2m$ , we have  $\alpha_i \in \text{Mor}(\text{Im}(F))$ .

015E

2. *Surjectivity on Objects.* If  $F$  is an epimorphism of categories, then  $F$  is surjective on objects.<sup>1</sup>*Further Terminology:* This statement is known as **Isbell's zigzag theorem**.**PROOF 11.7.3.1.3 ▶ PROOF OF PROPOSITION 11.7.3.1.2****Item 1: Characterisations**

See [Isb68].

**Item 2: Surjectivity on Objects**

Omitted.

**QUESTION 11.7.3.1.4 ▶ CHARACTERISATIONS OF FUNCTORS WITH EPIC PRE/POSTCOMPOSITION**

015F

Is there a characterisation of functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  such that:

015G

1. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is an epimorphism of categories?

015H

2. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is an epimorphism of categories?

This question also appears as [MO 468121a].

#### 015J 11.7.4 Pseudomonic Functors

Let  $C$  and  $\mathcal{D}$  be categories.

015K

##### DEFINITION 11.7.4.1.1 ► PSEUDOMONIC FUNCTORS

A functor  $F: C \rightarrow \mathcal{D}$  is **pseudomonic** if it satisfies the following conditions:

015L

- For all diagrams of the form

$$X \xrightarrow[\psi]{\alpha \parallel \beta} C \xrightarrow{F} \mathcal{D},$$

if we have

$$\text{id}_F \star \alpha = \text{id}_F \star \beta,$$

then  $\alpha = \beta$ .

015M

- For each  $X \in \text{Obj}(\text{Cats})$  and each natural isomorphism

$$\beta: F \circ \phi \xrightarrow{\sim} F \circ \psi, \quad X \xrightarrow[\substack{F \circ \psi \\ \beta \downarrow}]{} \mathcal{D},$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \xrightarrow[\substack{\psi \\ \alpha \downarrow}]{} C$$

such that we have an equality

$$X \xrightarrow[\substack{\psi \\ \alpha \downarrow}]{} C \xrightarrow{F} \mathcal{D} = X \xrightarrow[\substack{F \circ \psi \\ \beta \downarrow}]{} \mathcal{D}$$

of pasting diagrams, i.e. such that we have

$$\beta = \text{id}_F \star \alpha.$$

015N

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a functor.

015P 1. *Characterisations.* The following conditions are equivalent:

015Q (a) The functor  $F$  is pseudomonic.

015R (b) The functor  $F$  satisfies the following conditions:

015S i. The functor  $F$  is faithful, i.e. for each  $A, B \in \text{Obj}(\mathcal{C})$ , the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of  $F$  at  $(A, B)$  is injective.

015T ii. For each  $A, B \in \text{Obj}(\mathcal{C})$ , the restriction

$$F_{A,B}^{\text{iso}}: \text{Iso}_{\mathcal{C}}(A, B) \rightarrow \text{Iso}_{\mathcal{D}}(F_A, F_B)$$

of the action on morphisms of  $F$  at  $(A, B)$  to isomorphisms is surjective.

015U (c) We have an isocomma square of the form

$$\begin{array}{ccc} C & \xrightarrow{\text{id}_C} & C \\ C \xrightarrow{\text{eq.}} C \times_{\mathcal{D}} C & \downarrow \text{id}_C & \downarrow F \\ & \lrcorner \nearrow \lrcorner & \\ C & \xrightarrow{F} & \mathcal{D} \end{array}$$

in  $\text{Cats}_2$  up to equivalence.

015V (d) We have an isocomma square of the form

$$\begin{array}{ccc} C & \hookrightarrow \text{Arr}(C) & \\ C \xrightarrow{\text{eq.}} C \times_{\text{Arr}(\mathcal{D})} \mathcal{D} & \downarrow F & \downarrow \text{Arr}(F) \\ & \lrcorner \nearrow \lrcorner & \\ \mathcal{D} & \hookrightarrow \text{Arr}(\mathcal{D}) & \end{array}$$

in  $\text{Cats}_2$  up to equivalence.

015W (e) For each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition<sup>1</sup> functor

$$F_*: \text{Fun}(X, \mathcal{C}) \rightarrow \text{Fun}(X, \mathcal{D})$$

is pseudomonic.

015X 2. *Conservativity.* If  $F$  is pseudomonic, then  $F$  is conservative.

015Y 3. *Essential Injectivity.* If  $F$  is pseudomonic, then  $F$  is essentially injective.

<sup>1</sup>Asking the precomposition functors

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(\mathcal{C}, X)$$

to be pseudomonic leads to pseudoepic functors; see Item 1b of Item 1 of Proposition 11.7.5.1.2.

**PROOF 11.7.4.1.3 ▶ PROOF OF PROPOSITION 11.7.4.1.2**

Item 1: Characterisations

Omitted.

Item 2: Conservativity

Omitted.

Item 3: Essential Injectivity

Omitted.

**015Z 11.7.5 Pseudoepic Functors**

Let  $C$  and  $\mathcal{D}$  be categories.

**0160 DEFINITION 11.7.5.1.1 ▶ PSEUDOEPIC FUNCTORS**

A functor  $F: C \rightarrow \mathcal{D}$  is **pseudoepic** if it satisfies the following conditions:

**0161** 1. For all diagrams of the form

$$C \xrightarrow{F} \mathcal{D} \begin{array}{c} \nearrow \phi \\ \alpha \parallel \beta \\ \searrow \psi \end{array} X,$$

if we have

$$\alpha \star \text{id}_F = \beta \star \text{id}_F,$$

then  $\alpha = \beta$ .

**0162** 2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ F \xrightarrow{\sim} \psi \circ F, \quad C \begin{array}{c} \nearrow \phi \circ F \\ \beta \parallel \\ \searrow \psi \circ F \end{array} X$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{D} \begin{array}{c} \nearrow \phi \\ \alpha \parallel \\ \searrow \psi \end{array} X$$

of  $C$  such that we have an equality

$$C \xrightarrow{F} \mathcal{D} \begin{array}{c} \nearrow \phi \\ \alpha \parallel \\ \searrow \psi \end{array} X = C \begin{array}{c} \nearrow \phi \circ F \\ \beta \parallel \\ \searrow \psi \circ F \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_F.$$

0163

**PROPOSITION 11.7.5.1.2 ► PROPERTIES OF PSEUDOEPIC FUNCTORS**

Let  $F: C \rightarrow \mathcal{D}$  be a functor.

0164

1. *Characterisations.* The following conditions are equivalent:

0165

(a) The functor  $F$  is pseudoepic.

0166

(b) For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

given by precomposition by  $F$  is pseudomonic.

0167

(c) We have an isococomma square of the form

$$\begin{array}{ccc} & \mathcal{D} & \\ & \xleftarrow{\text{id}_{\mathcal{D}}} & \\ \mathcal{D} \stackrel{\text{eq.}}{\cong} \mathcal{D} \amalg_C \mathcal{D} & \xleftrightarrow{\quad} & \mathcal{D} \\ \text{id}_{\mathcal{D}} \uparrow & \swarrow \approx & \uparrow F \\ \mathcal{D} & \xleftarrow{F} & C \end{array}$$

in  $\text{Cats}_2$  up to equivalence.

0168

2. *Dominance.* If  $F$  is pseudoepic, then  $F$  is dominant ([Definition 11.7.1.1.1](#)).

**PROOF 11.7.5.1.3 ► PROOF OF PROPOSITION 11.7.5.1.2**

**Item 1: Characterisations**

Omitted.

**Item 2: Dominance**

If  $F$  is pseudoepic, then

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is pseudomonic for all  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , and thus in particular faithful. By [Item 5g](#) of [Item 5](#) of [Proposition 11.6.1.1.2](#), this is equivalent to requiring  $F$  to be dominant. 

0169

**QUESTION 11.7.5.1.4 ► CHARACTERISATIONS OF PSEUDOEPIC FUNCTORS**

Is there a nice characterisation of the pseudoepic functors, similarly to the characterisation of pseudomonic functors given in [Item 1b](#) of [Item 1](#) of [Proposition 11.7.4.1.2](#)?

This question also appears as [[MO 321971](#)].

016A

A pseudomonic and pseudoepic functor is dominant, faithful, essentially injective, and full on isomorphisms. Is it necessarily an equivalence of categories? If not, how bad can this fail, i.e. how far can a pseudomonic and pseudoepic functor be from an equivalence of categories?  
This question also appears as [MO 468334].

016B

**QUESTION 11.7.5.1.6 ► CHARACTERISATIONS OF FUNCTORS WITH PSEUDOEPIC PRE/POST-COMPOSITION**

Is there a characterisation of functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  such that:

016C

1. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is pseudoepic?

016D

2. For each  $\mathcal{X} \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is pseudoepic?

This question also appears as [MO 468121a].

016E

## 11.8 Even More Conditions on Functors

016F

### 11.8.1 Injective on Objects Functors

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

016G

**DEFINITION 11.8.1.1.1 ► INJECTIVE ON OBJECTS FUNCTORS**

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **injective on objects** if the action on objects

$$F: \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$$

of  $F$  is injective.

016H

**PROPOSITION 11.8.1.1.2 ► PROPERTIES OF INJECTIVE ON OBJECTS FUNCTORS**

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a functor.

016J

1. *Characterisations.* The following conditions are equivalent:

016K

- (a) The functor  $F$  is injective on objects.

016L

(b) The functor  $F$  is an isocofibration in  $\text{Cats}_2$ .

**PROOF 11.8.1.1.3 ▶ PROOF OF PROPOSITION 11.8.1.1.2**

Item 1: Characterisations

Omitted.



**016M 11.8.2 Surjective on Objects Functors**

Let  $C$  and  $\mathcal{D}$  be categories.

016N

**DEFINITION 11.8.2.1.1 ▶ SURJECTIVE ON OBJECTS FUNCTORS**

A functor  $F: C \rightarrow \mathcal{D}$  is **surjective on objects** if the action on objects

$$F: \text{Obj}(C) \rightarrow \text{Obj}(\mathcal{D})$$

of  $F$  is surjective.

**016P 11.8.3 Bijective on Objects Functors**

Let  $C$  and  $\mathcal{D}$  be categories.

016Q

**DEFINITION 11.8.3.1.1 ▶ BIJECTIVE ON OBJECTS FUNCTORS**

A functor  $F: C \rightarrow \mathcal{D}$  is **bijective on objects**<sup>1</sup> if the action on objects

$$F: \text{Obj}(C) \rightarrow \text{Obj}(\mathcal{D})$$

of  $F$  is a bijection.

<sup>1</sup>Further Terminology: Also called a **bo** functor.

**016R 11.8.4 Functors Representably Faithful on Cores**

Let  $C$  and  $\mathcal{D}$  be categories.

016S

**DEFINITION 11.8.4.1.1 ▶ FUNCTORS REPRESENTABLY FAITHFUL ON CORES**

A functor  $F: C \rightarrow \mathcal{D}$  is **representably faithful on cores** if, for each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition by  $F$  functor

$$F_*: \text{Core}(\text{Fun}(X, C)) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is faithful.

016T

**REMARK 11.8.4.1.2 ► UNWINDING DEFINITION 11.8.4.1.1**

In detail, a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **representably faithful on cores** if, given a diagram of the form

$$\mathcal{X} \xrightarrow[\psi]{\alpha \parallel \beta} \mathcal{C} \xrightarrow{F} \mathcal{D},$$

if  $\alpha$  and  $\beta$  are natural isomorphisms and we have

$$\text{id}_F \star \alpha = \text{id}_F \star \beta,$$

then  $\alpha = \beta$ .

016U

**QUESTION 11.8.4.1.3 ► CHARACTERISATION OF FUNCTORS REPRESENTABLY FAITHFUL ON CORES**

Is there a characterisation of functors representably faithful on cores?

016V

**11.8.5 Functors Representably Full on Cores**

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

016W

**DEFINITION 11.8.5.1.1 ► FUNCTORS REPRESENTABLY FULL ON CORES**

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **representably full on cores** if, for each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition by  $F$  functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is full.

016X

**REMARK 11.8.5.1.2 ► UNWINDING DEFINITION 11.8.5.1.1**

In detail, a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **representably full on cores** if, for each  $X \in \text{Obj}(\text{Cats})$  and each natural isomorphism

$$\beta: F \circ \phi \xrightarrow{\sim} F \circ \psi, \quad X \xrightarrow[\substack{F \circ \psi \\ \beta \downarrow}]{} \mathcal{D},$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \xrightarrow[\psi]{\alpha \Downarrow \beta} C$$

such that we have an equality

$$X \xrightarrow[\psi]{\alpha \Downarrow \beta} C \xrightarrow{F} D = X \xrightarrow[\psi]{F \circ \phi \Downarrow F \circ \beta} D$$

of pasting diagrams in  $\text{Cats}_2$ , i.e. such that we have

$$\beta = \text{id}_F \star \alpha.$$

016Y

#### QUESTION 11.8.5.1.3 ► CHARACTERISATION OF FUNCTORS REPRESENTABLY FULL ON CORES

Is there a characterisation of functors representably full on cores?  
This question also appears as [MO 468121a].

016Z

#### 11.8.6 Functors Representably Fully Faithful on Cores

Let  $C$  and  $D$  be categories.

0170

#### DEFINITION 11.8.6.1.1 ► FUNCTORS REPRESENTABLY FULLY FAITHFUL ON CORES

A functor  $F: C \rightarrow D$  is **representably fully faithful on cores** if, for each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition by  $F$  functor

$$F_*: \text{Core}(\text{Fun}(X, C)) \rightarrow \text{Core}(\text{Fun}(X, D))$$

is fully faithful.

0171

#### REMARK 11.8.6.1.2 ► UNWINDING DEFINITION 11.8.6.1.1

In detail, a functor  $F: C \rightarrow D$  is **representably fully faithful on cores** if it satisfies the conditions in [Remarks 11.8.4.1.2](#) and [11.8.5.1.2](#), i.e.:

0172

1. For all diagrams of the form

$$X \xrightarrow[\psi]{\alpha \Downarrow \beta} C \xrightarrow{F} D,$$

with  $\alpha$  and  $\beta$  natural isomorphisms, if we have  $\text{id}_F \star \alpha = \text{id}_F \star \beta$ , then  $\alpha = \beta$ .

0173

2. For each  $X \in \text{Obj}(\text{Cats})$  and each natural isomorphism

$$\beta: F \circ \phi \xrightarrow{\sim} F \circ \psi, \quad X \xrightarrow[\substack{\beta \Downarrow \\ F \circ \psi}]{} \mathcal{D}$$

of  $C$ , there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \xrightarrow[\substack{\alpha \Downarrow \\ \psi}]{} C$$

of  $C$  such that we have an equality

$$X \xrightarrow[\substack{\phi \\ \alpha \Downarrow \\ \psi}]{} C \xrightarrow{F} \mathcal{D} = X \xrightarrow[\substack{\beta \Downarrow \\ F \circ \psi}]{} \mathcal{D}$$

of pasting diagrams in  $\text{Cats}_2$ , i.e. such that we have

$$\beta = \text{id}_F \star \alpha.$$

0174

#### QUESTION 11.8.6.1.3 ► CHARACTERISATION OF FUNCTORS REPRESENTABLY FULLY FAITHFUL ON CORES

Is there a characterisation of functors representably fully faithful on cores?

0175

### 11.8.7 Functors Corepresentably Faithful on Cores

Let  $C$  and  $\mathcal{D}$  be categories.

0176

#### DEFINITION 11.8.7.1.1 ► FUNCTORS COREPRESENTABLY FAITHFUL ON CORES

A functor  $F: C \rightarrow \mathcal{D}$  is **corepresentably faithful on cores** if, for each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition by  $F$  functor

$$F_*: \text{Core}(\text{Fun}(X, C)) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is faithful.

0177

**REMARK 11.8.7.1.2 ► UNWINDING DEFINITION 11.8.7.1.1**

In detail, a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **corepresentably faithful on cores** if, given a diagram of the form

$$\mathcal{C} \xrightarrow{F} \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \parallel \beta \\[-1ex] \xrightarrow{\psi} \end{array} X,$$

if  $\alpha$  and  $\beta$  are natural isomorphisms and we have

$$\alpha \star \text{id}_F = \beta \star \text{id}_F,$$

then  $\alpha = \beta$ .

0178

**QUESTION 11.8.7.1.3 ► CHARACTERISATION OF FUNCTORS COREPRESENTABLY FAITHFUL ON CORES**

Is there a characterisation of functors corepresentably faithful on cores?

0179

**11.8.8 Functors Corepresentably Full on Cores**

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.

017A

**DEFINITION 11.8.8.1.1 ► FUNCTORS COREPRESENTABLY FULL ON CORES**

A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **corepresentably full on cores** if, for each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition by  $F$  functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is full.

017B

**REMARK 11.8.8.1.2 ► UNWINDING DEFINITION 11.8.8.1.1**

In detail, a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is **corepresentably full on cores** if, for each  $X \in \text{Obj}(\text{Cats})$  and each natural isomorphism

$$\beta: \phi \circ F \xrightarrow{\sim} \psi \circ F, \quad \mathcal{C} \begin{array}{c} \xrightarrow{\phi \circ F} \\[-1ex] \beta \parallel \\[-1ex] \xrightarrow{\psi \circ F} \end{array} X,$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \xrightarrow[\psi]{\alpha \Downarrow} \\[-1ex] \xrightarrow{\psi} \end{array} X$$

such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \xrightarrow[\psi]{\alpha \Downarrow} \\[-1ex] \xrightarrow{\psi} \end{array} C \xrightarrow{F} \mathcal{D} = X \begin{array}{c} \xrightarrow{F \circ \phi} \\[-1ex] \xrightarrow[\psi]{\beta \Downarrow} \\[-1ex] \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of pasting diagrams in  $\text{Cats}_2$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_F.$$

017C

#### QUESTION 11.8.8.1.3 ► CHARACTERISATION OF FUNCTORS COREPRESENTABLY FULL ON CORES

Is there a characterisation of functors corepresentably full on cores?  
This question also appears as [MO 468121a].

017D

#### 11.8.9 Functors Corepresentably Fully Faithful on Cores

Let  $C$  and  $\mathcal{D}$  be categories.

017E

#### DEFINITION 11.8.9.1.1 ► FUNCTORS COREPRESENTABLY FULLY FAITHFUL ON CORES

A functor  $F: C \rightarrow \mathcal{D}$  is **corepresentably fully faithful on cores** if, for each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition by  $F$  functor

$$F_*: \text{Core}(\text{Fun}(X, C)) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is fully faithful.

017F

#### REMARK 11.8.9.1.2 ► UNWINDING DEFINITION 11.8.9.1.1

In detail, a functor  $F: C \rightarrow \mathcal{D}$  is **corepresentably fully faithful on cores** if it satisfies the conditions in Remarks 11.8.7.1.2 and 11.8.8.1.2, i.e.:

017G

1. For all diagrams of the form

$$C \xrightarrow{F} \mathcal{D} \xrightarrow{\phi} X,$$

$\alpha \Downarrow \beta$

if  $\alpha$  and  $\beta$  are natural isomorphisms and we have

$$\alpha \star \text{id}_F = \beta \star \text{id}_F,$$

then  $\alpha = \beta$ .

017H

2. For each  $X \in \text{Obj}(\text{Cats})$  and each natural isomorphism

$$\beta: \phi \circ F \xrightarrow{\sim} \psi \circ F, \quad C \xrightarrow[\psi \circ F]{\phi \circ F} X,$$

$\beta \Downarrow$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{D} \xrightarrow[\psi]{\phi} X$$

$\alpha \Downarrow$

such that we have an equality

$$X \xrightarrow[\psi]{\phi} C \xrightarrow{F} \mathcal{D} = X \xrightarrow[\psi \circ F]{F \circ \phi} \mathcal{D}$$

of pasting diagrams in  $\text{Cats}_2$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_F.$$

017J

**QUESTION 11.8.9.1.3 ► CHARACTERISATION OF FUNCTORS COREPRESENTABLY FULLY FAITHFUL ON CORES**

Is there a characterisation of functors corepresentably fully faithful on cores?

**017K 11.9 Natural Transformations****017L 11.9.1 Transformations**

Let  $C$  and  $\mathcal{D}$  be categories and let  $F, G: C \Rightarrow \mathcal{D}$  be functors.

017M

**DEFINITION 11.9.1.1.1 ► TRANSFORMATIONS**

A **transformation**<sup>1</sup>  $\alpha: F \Rightarrow G$  from  $F$  to  $G$  is a collection

$$\{\alpha_A: F(A) \rightarrow G(A)\}_{A \in \text{Obj}(C)}$$

of morphisms of  $\mathcal{D}$ .

<sup>1</sup>*Further Terminology:* Also called an **unnatural transformation** for emphasis.

017N

**NOTATION 11.9.1.1.2 ► THE SET OF TRANSFORMATIONS BETWEEN TWO FUNCTORS**

We write  $\text{Trans}(F, G)$  for the set of transformations from  $F$  to  $G$ .

01U9

**REMARK 11.9.1.1.3 ► THE SET OF TRANSFORMATIONS AS A PRODUCT**

We have an isomorphism

$$\text{Trans}(F, G) \cong \prod_{A \in C} \text{Hom}_{\mathcal{D}}(F_A, G_A).$$

**PROOF 11.9.1.1.4 ► PROOF OF REMARK 11.9.1.1.3**

Omitted. 

017P

**11.9.2 Natural Transformations**

Let  $C$  and  $\mathcal{D}$  be categories and  $F, G: C \rightrightarrows \mathcal{D}$  be functors.

017Q

**DEFINITION 11.9.2.1.1 ► NATURAL TRANSFORMATIONS**

A **natural transformation**  $\alpha: F \Rightarrow G$  from  $F$  to  $G$  is a transformation

$$\{\alpha_A: F(A) \rightarrow G(A)\}_{A \in \text{Obj}(C)}$$

from  $F$  to  $G$  such that, for each morphism  $f: A \rightarrow B$  of  $C$ , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \alpha_A \downarrow & & \downarrow \alpha_B \\ G(A) & \xrightarrow{G(f)} & G(B) \end{array}$$

commutes.

01UA

**REMARK 11.9.2.1.2 ► FURTHER TERMINOLOGY AND NOTATION FOR NATURAL TRANSFORMATIONS**

01UB

Let  $\alpha: F \Rightarrow G$  be a natural transformation.

01UC

1. For each  $A \in \text{Obj}(C)$ , the morphism  $\alpha_A: F_A \rightarrow G_A$  is called the **component of  $\alpha$  at  $A$** .
2. We denote natural transformations such as  $\alpha$  in diagrams as

$$\begin{array}{ccc} C & \xrightarrow{\quad \alpha \Downarrow \quad} & \mathcal{D}. \\ \downarrow & \nearrow & \downarrow \\ G & & \end{array}$$

017S

**NOTATION 11.9.2.1.3 ► THE SET OF NATURAL TRANSFORMATIONS BETWEEN TWO FUNCTIONS**

We write  $\text{Nat}(F, G)$  for the set of natural transformations from  $F$  to  $G$ .

017U

**DEFINITION 11.9.2.1.4 ► EQUALITY OF NATURAL TRANSFORMATIONS**

Two natural transformations  $\alpha, \beta: F \Rightarrow G$  are **equal** if we have

$$\alpha_A = \beta_A$$

for each  $A \in \text{Obj}(C)$ .

01UD

**11.9.3 Examples of Natural Transformations**

017T

**EXAMPLE 11.9.3.1.1 ► IDENTITY NATURAL TRANSFORMATIONS**

The **identity natural transformation**  $\text{id}_F: F \Rightarrow F$  of  $F$  is the natural transformation consisting of the collection

$$\{(\text{id}_F)_A: F(A) \rightarrow F(A)\}_{A \in \text{Obj}(C)}$$

defined by

$$(\text{id}_F)_A \stackrel{\text{def}}{=} \text{id}_{F(A)}$$

for each  $A \in \text{Obj}(C)$ .

**PROOF 11.9.3.1.2 ▶ PROOF OF EXAMPLE 11.9.3.1.1**

The naturality condition for  $\text{id}_F$  is the requirement that, for each morphism  $f: A \rightarrow B$  of  $C$ , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \text{id}_{F(A)} \downarrow & & \downarrow \text{id}_{F(B)} \\ F(A) & \xrightarrow[F(f)]{} & F(B) \end{array}$$

commutes. This follows from unitality of the composition of  $\mathcal{D}$ , as we have

$$\begin{aligned} F(f) \circ \text{id}_{F(A)} &= F(f) \\ &= \text{id}_{F(B)} \circ F(f), \end{aligned}$$

where we have applied unitality twice. ■

01UE

**EXAMPLE 11.9.3.1.3 ▶ NATURAL TRANSFORMATIONS BETWEEN MORPHISMS OF MONOIDS**

Let  $A$  and  $B$  be monoids and let  $f, g: A \Rightarrow B$  be morphisms of monoids. Applying the delooping construction of ??, we obtain functors  $Bf, Bg: BA \Rightarrow BB$ . We then have

$$\text{Nat}(Bf, Bg) \cong \left\{ b \in B \middle| \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } bf(a) = g(a)b \end{array} \right\}.$$

**PROOF 11.9.3.1.4 ▶ PROOF OF EXAMPLE 11.9.3.1.3**

Unwinding the definitions in this case, we see that a transformation  $\alpha$  from  $Bf$  to  $Bg$  consists of a collection

$$\{\alpha_{\bullet}: \bullet \rightarrow \bullet\}_{\bullet \in \text{Obj}(BA)}$$

of morphisms of  $BB$  indexed by  $\text{Obj}(BA)$ . Since  $\text{Obj}(BA) = \text{pt}$  and the morphisms of  $BB$  are precisely the elements of  $B$ , it follows that  $\alpha$  corresponds precisely to the data of an element  $b \in B$ . Now, a transformation  $[b]: Bf \Rightarrow Bg$  is natural precisely if, for each  $a \in \text{Hom}_{BA}(\bullet, \bullet) \stackrel{\text{def}}{=} A$ , the

diagram

$$\begin{array}{ccc} \mathbf{B}f(\bullet) & \xrightarrow{\mathbf{B}f(a)} & \mathbf{B}f(\bullet) \\ [b]_\bullet \downarrow & & \downarrow [b]_\bullet \\ \mathbf{B}g(\bullet) & \xrightarrow{\mathbf{B}g(a)} & \mathbf{B}g(\bullet) \end{array}$$

commutes. Unwinding the definitions, we see that this diagram is given by

$$\begin{array}{ccc} \bullet & \xrightarrow{f(a)} & \bullet \\ b \downarrow & & \downarrow b \\ \bullet & \xrightarrow{g(a)} & \bullet, \end{array}$$

and hence corresponds precisely to the condition  $g(a)b = bf(a)$ . ■

#### 017V 11.9.4 Vertical Composition of Natural Transformations

017W

##### DEFINITION 11.9.4.1.1 ► VERTICAL COMPOSITION OF NATURAL TRANSFORMATIONS

The **vertical composition** of two natural transformations  $\alpha: F \Rightarrow G$  and  $\beta: G \Rightarrow H$  as in the diagram

$$\begin{array}{ccc} & F & \\ & \alpha \Downarrow & \\ C & \xrightarrow{G} & \mathcal{D} \\ & \beta \Downarrow & \\ & H & \end{array}$$

is the natural transformation  $\beta \circ \alpha: F \Rightarrow H$  consisting of the collection

$$\{(\beta \circ \alpha)_A: F(A) \rightarrow H(A)\}_{A \in \text{Obj}(C)}$$

with

$$(\beta \circ \alpha)_A \stackrel{\text{def}}{=} \beta_A \circ \alpha_A$$

for each  $A \in \text{Obj}(C)$ .

## PROOF 11.9.4.1.2 ► PROOF OF DEFINITION 11.9.4.1.1

The naturality condition for  $\beta \circ \alpha$  is the requirement that the boundary of the diagram

$$\begin{array}{ccc}
 F(A) & \xrightarrow{F(f)} & F(B) \\
 \alpha_A \downarrow & (1) & \downarrow \alpha_B \\
 G(A) & \xrightarrow{G(f)} & G(B) \\
 \beta_A \downarrow & (2) & \downarrow \beta_B \\
 H(A) & \xrightarrow{H(f)} & H(B)
 \end{array}$$

commutes. Since

- Subdiagram (1) commutes by the naturality of  $\alpha$ .
- Subdiagram (2) commutes by the naturality of  $\beta$ .

so does the boundary diagram. Hence  $\beta \circ \alpha$  is a natural transformation.



## PROPOSITION 11.9.4.1.3 ► PROPERTIES OF VERTICAL COMPOSITION OF NATURAL TRANSFORMATIONS

017X

Let  $C$ ,  $\mathcal{D}$ , and  $\mathcal{E}$  be categories.

017Y

1. *Functionality.* The assignment  $(\beta, \alpha) \mapsto \beta \circ \alpha$  defines a function

$$\circ_{F,G,H} : \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H).$$

017Z

2. *Associativity.* Let  $F, G, H, K : C \rightrightarrows \mathcal{D}$  be functors. The diagram

$$\begin{array}{ccc}
 & \text{Nat}(H, K) \times (\text{Nat}(G, H) \times \text{Nat}(F, G)) & \\
 & \swarrow \alpha_{\text{Nat}(H,K), \text{Nat}(G,H), \text{Nat}(F,G)}^{\text{Sets}} \quad \searrow \text{id}_{\text{Nat}(H,K)} \times \circ_{F,G,H} & \\
 (\text{Nat}(H, K) \times \text{Nat}(G, H)) \times \text{Nat}(F, G) & & \text{Nat}(H, K) \times \text{Nat}(F, H) \\
 & \swarrow \circ_{G,H,K} \times \text{id}_{\text{Nat}(F,G)} & \searrow \circ_{F,H,K} \\
 & \text{Nat}(G, K) \times \text{Nat}(F, G) & \xrightarrow{\circ_{F,G,K}} \text{Nat}(F, K)
 \end{array}$$

commutes, i.e. given natural transformations

$$F \xrightarrow{\alpha} G \xrightarrow{\beta} H \xrightarrow{\gamma} K,$$

we have

$$(\gamma \circ \beta) \circ \alpha = \gamma \circ (\beta \circ \alpha).$$

0180

3. *Unitality.* Let  $F, G: C \Rightarrow \mathcal{D}$  be functors.

(a) *Left Unitality.* The diagram

$$\begin{array}{ccc} \text{pt} \times \text{Nat}(F, G) & & \\ \downarrow [\text{id}_G] \times \text{id}_{\text{Nat}(F, G)} & \nearrow \lambda_{\text{Nat}(F, G)}^{\text{Sets}} & \\ \text{Nat}(G, G) \times \text{Nat}(F, G) & \xrightarrow{\circ_{F, G, G}} & \text{Nat}(F, G) \end{array}$$

commutes, i.e. given a natural transformation  $\alpha: F \Rightarrow G$ , we have

$$\text{id}_G \circ \alpha = \alpha.$$

(b) *Right Unitality.* The diagram

$$\begin{array}{ccc} \text{Nat}(F, G) \times \text{pt} & & \\ \downarrow \text{id}_{\text{Nat}(F, G)} \times [\text{id}_F] & \nearrow \rho_{\text{Nat}(F, G)}^{\text{Sets}} & \\ \text{Nat}(F, G) \times \text{Nat}(F, F) & \xrightarrow{\circ_{F, F, G}^C} & \text{Nat}(F, G) \end{array}$$

commutes, i.e. given a natural transformation  $\alpha: F \Rightarrow G$ , we have

$$\alpha \circ \text{id}_F = \alpha.$$

0181

4. *Middle Four Exchange.* Let  $F_1, F_2, F_3: C \rightarrow \mathcal{D}$  and  $G_1, G_2, G_3: \mathcal{D} \rightarrow \mathcal{E}$  be functors. The diagram

$$\begin{array}{ccc} (\text{Nat}(G_2, G_3) \times \text{Nat}(G_1, G_2)) \times (\text{Nat}(F_2, F_3) \times \text{Nat}(F_1, F_2)) & \xleftarrow{\mu_4} & (\text{Nat}(G_2, G_3) \times \text{Nat}(F_2, F_3)) \times (\text{Nat}(G_1, G_2) \times \text{Nat}(F_1, F_2)) \\ \downarrow \circ_{G_1, G_2, G_3} \times \circ_{F_1, F_2, F_3} & & \downarrow \star_{F_2, F_3, G_2, G_3} \times \star_{F_1, F_2, G_1, G_2} \\ \text{Nat}(G_1, G_3) \times \text{Nat}(F_1, F_3) & & \text{Nat}(G_2 \circ F_2, G_3 \circ F_3) \times \text{Nat}(G_1 \circ F_1, G_2 \circ F_2) \\ & \searrow \star_{F_1, F_3, G_1, G_3} & \swarrow \circ_{G_1 \circ F_1, G_2 \circ F_2, G_3 \circ F_3} \\ & \text{Nat}(G_1 \circ F_1, G_3 \circ F_3) & \end{array}$$

commutes, i.e. given a diagram

$$\begin{array}{ccccc}
 & F_1 & & G_1 & \\
 & \alpha \Downarrow & & \beta \Downarrow & \\
 C & \xrightarrow{F_2} & \mathcal{D} & \xrightarrow{G_2} & \mathcal{E} \\
 \alpha' \Downarrow & & & \beta' \Downarrow & \\
 & F_3 & & G_3 &
 \end{array}$$

in  $\text{Cats}_2$ , we have

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

#### PROOF 11.9.4.1.4 ► PROOF OF PROPOSITION 11.9.4.1.3

**Item 1: Functionality**

Omitted.

**Item 2: Associativity**

Indeed, we have

$$\begin{aligned}
 ((\gamma \circ \beta) \circ \alpha)_A &\stackrel{\text{def}}{=} (\gamma \circ \beta)_A \circ \alpha_A \\
 &\stackrel{\text{def}}{=} (\gamma_A \circ \beta_A) \circ \alpha_A \\
 &= \gamma_A \circ (\beta_A \circ \alpha_A) \\
 &\stackrel{\text{def}}{=} \gamma_A \circ (\beta \circ \alpha)_A \\
 &\stackrel{\text{def}}{=} (\gamma \circ (\beta \circ \alpha))_A
 \end{aligned}$$

for each  $A \in \text{Obj}(C)$ , showing the desired equality.

**Item 3: Unitality**

We have

$$\begin{aligned}
 (\text{id}_G \circ \alpha)_A &= \text{id}_G \circ \alpha_A \\
 &= \alpha_A, \\
 (\alpha \circ \text{id}_F)_A &= \alpha_A \circ \text{id}_F \\
 &= \alpha_A
 \end{aligned}$$

for each  $A \in \text{Obj}(C)$ , showing the desired equality.

**Item 4: Middle Four Exchange**

This is proved in **Item 4** of [Proposition 11.9.5.1.4](#).



**0182 11.9.5 Horizontal Composition of Natural Transformations**

**0183**

**DEFINITION 11.9.5.1.1 ► HORIZONTAL COMPOSITION OF NATURAL TRANSFORMATIONS**

The **horizontal composition**<sup>1,2</sup> of two natural transformations  $\alpha: F \Rightarrow G$  and  $\beta: H \Rightarrow K$  as in the diagram

$$\begin{array}{ccc} C & \xrightarrow[F]{\alpha \Downarrow} & \mathcal{D} \\ & \Downarrow G & \\ & \xrightarrow[H]{\beta \Downarrow} & \mathcal{E} \\ & \Downarrow K & \end{array}$$

of  $\alpha$  and  $\beta$  is the natural transformation

$$\beta \star \alpha: (H \circ F) \Rightarrow (K \circ G),$$

as in the diagram

$$\begin{array}{ccc} C & \xrightarrow[H \circ F]{\beta \star \alpha \Downarrow} & \mathcal{E}, \\ & \Downarrow K \circ G & \end{array}$$

consisting of the collection

$$\{(\beta \star \alpha)_A: H(F(A)) \rightarrow K(G(A))\}_{A \in \text{Obj}(C)},$$

of morphisms of  $\mathcal{E}$  with

$$\begin{array}{ccc} (\beta \star \alpha)_A & \stackrel{\text{def}}{=} & \beta_{G(A)} \circ H(\alpha_A) \\ & = & K(\alpha_A) \circ \beta_{F(A)}, \\ & & \end{array} \quad \begin{array}{ccc} H(F(A)) & \xrightarrow{H(\alpha_A)} & H(G(A)) \\ \beta_{F(A)} \downarrow & & \downarrow \beta_{G(A)} \\ K(F(A)) & \xrightarrow{K(\alpha_A)} & K(G(A)). \end{array}$$

<sup>1</sup>Further Terminology: Also called the **Godement product** of  $\alpha$  and  $\beta$ .

<sup>2</sup>Horizontal composition forms a map

$$\star_{(F,H),(G,K)}: \text{Nat}(H,K) \times \text{Nat}(F,G) \rightarrow \text{Nat}(H \circ F, K \circ G).$$

## PROOF 11.9.5.1.2 ▶ PROOF OF DEFINITION 11.9.5.1.1

First, we claim that we indeed have

$$\begin{array}{ccc} H(F(A)) & \xrightarrow{H(\alpha_A)} & H(G(A)) \\ \beta_{G(A)} \circ H(\alpha_A) = K(\alpha_A) \circ \beta_{F(A)}, & \beta_{F(A)} \downarrow & \downarrow \beta_{G(A)} \\ K(F(A)) & \xrightarrow{K(\alpha_A)} & K(G(A)). \end{array}$$

This is, however, simply the naturality square for  $\beta$  applied to the morphism  $\alpha_A: F(A) \rightarrow G(A)$ . Next, we check the naturality condition for  $\beta \star \alpha$ , which is the requirement that the boundary of the diagram

$$\begin{array}{ccc} H(F(A)) & \xrightarrow{H(F(f))} & H(F(B)) \\ H(\alpha_A) \downarrow & (1) & \downarrow H(\alpha_B) \\ H(G(A)) & \xrightarrow{H(G(f))} & H(G(B)) \\ \beta_{G(A)} \downarrow & (2) & \downarrow \beta_{G(B)} \\ K(G(A)) & \xrightarrow{K(G(f))} & K(G(B)) \end{array}$$

commutes. Since

- Subdiagram (1) commutes by the naturality of  $\alpha$ .
- Subdiagram (2) commutes by the naturality of  $\beta$ .

so does the boundary diagram. Hence  $\beta \circ \alpha$  is a natural transformation.<sup>1</sup>



<sup>1</sup>Reference: [Bor94, Proposition 1.3.4].

0184

## DEFINITION 11.9.5.1.3 ▶ WHISKERING OF FUNCTORS WITH NATURAL TRANSFORMATIONS

Let

$$\mathcal{X} \xrightarrow{F} \mathcal{C} \xrightarrow{\alpha \Downarrow \phi} \mathcal{D} \xrightarrow{G} \mathcal{Y}$$

be a diagram in  $\text{Cats}_2$ .

0185

1. The **left whiskering of  $\alpha$  with  $G$**  is the natural transformation<sup>1</sup>

$$\text{id}_G \star \alpha : G \circ \phi \Rightarrow G \circ \psi.$$

0186

2. The **right whiskering of  $\alpha$  with  $F$**  is the natural transformation<sup>2</sup>

$$\alpha \star \text{id}_F : \phi \circ F \Rightarrow \psi \circ F.$$

<sup>1</sup>*Further Notation:* Also written  $G\alpha$  or  $G \star \alpha$ , although we won't use either of these notations in this work.

<sup>2</sup>*Further Notation:* Also written  $\alpha F$  or  $\alpha \star F$ , although we won't use either of these notations in this work.

0187

#### PROPOSITION 11.9.5.1.4 ► PROPERTIES OF HORIZONTAL COMPOSITION OF NATURAL TRANSFORMATIONS

Let  $C$ ,  $\mathcal{D}$ , and  $\mathcal{E}$  be categories.

0188

1. *Functionality.* The assignment  $(\beta, \alpha) \mapsto \beta \star \alpha$  defines a function

$$\star_{(F,G),(H,K)} : \text{Nat}(H, K) \times \text{Nat}(F, G) \rightarrow \text{Nat}(H \circ F, K \circ G).$$

0189

2. *Associativity.* Let

$$C \xrightarrow[G_1]{F_1} \mathcal{D} \xrightarrow[G_2]{F_2} \mathcal{E} \xrightarrow[G_3]{F_3} \mathcal{F}$$

be a diagram in  $\text{Cats}_2$ . The diagram

$$\begin{array}{ccc} \text{Nat}(F_3, G_3) \times \text{Nat}(F_2, G_2) \times \text{Nat}(F_1, G_1) & \xrightarrow{\star_{(F_2, G_2), (F_3, G_3)} \times \text{id}} & \text{Nat}(F_3 \circ F_2, G_3 \circ G_2) \times \text{Nat}(F_1, G_1) \\ \downarrow \text{id} \times \star_{(F_1, G_1), (F_2, G_2)} & & \downarrow \star_{(F_3 \circ F_2), (G_3 \circ G_2, F_1, G_1)} \\ \text{Nat}(F_3, G_3) \times \text{Nat}(F_2 \circ F_1, G_2 \circ G_1) & \xrightarrow[\star_{(F_2 \circ F_1), (G_2 \circ G_1, F_3, G_3)}]{} & \text{Nat}(F_3 \circ F_2 \circ F_1, G_3 \circ G_2 \circ G_1) \end{array}$$

commutes, i.e. given natural transformations

$$C \xrightarrow[G_1]{\alpha \Downarrow} \mathcal{D} \xrightarrow[G_2]{\beta \Downarrow} \mathcal{E} \xrightarrow[G_3]{\gamma \Downarrow} \mathcal{F},$$

we have

$$(\gamma \star \beta) \star \alpha = \gamma \star (\beta \star \alpha).$$

018A

3. *Interaction With Identities.* Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  be functors. The diagram

$$\begin{array}{ccc} \text{pt} \times \text{pt} & \xrightarrow{[\text{id}_G] \times [\text{id}_F]} & \text{Nat}(G, G) \times \text{Nat}(F, F) \\ \uparrow & & \downarrow \star_{(F,F),(G,G)} \\ \text{pt} & \xrightarrow{[\text{id}_{G \circ F}]} & \text{Nat}(G \circ F, G \circ F) \end{array}$$

commutes, i.e. we have

$$\text{id}_G \star \text{id}_F = \text{id}_{G \circ F}.$$

018B

4. *Middle Four Exchange.* Let  $F_1, F_2, F_3: \mathcal{C} \rightarrow \mathcal{D}$  and  $G_1, G_2, G_3: \mathcal{D} \rightarrow \mathcal{E}$  be functors. The diagram

$$\begin{array}{ccc} (\text{Nat}(G_2, G_3) \times \text{Nat}(G_1, G_2)) \times (\text{Nat}(F_2, F_3) \times \text{Nat}(F_1, F_2)) & \xleftarrow{\mu_4} & (\text{Nat}(G_2, G_3) \times \text{Nat}(F_2, F_3)) \times (\text{Nat}(G_1, G_2) \times \text{Nat}(F_1, F_2)) \\ \downarrow \circ_{G_1, G_2, G_3} \times \circ_{F_1, F_2, F_3} & & \downarrow \star_{F_2, F_3, G_2, G_3} \times \star_{F_1, F_2, G_1, G_2} \\ \text{Nat}(G_1, G_3) \times \text{Nat}(F_1, F_3) & & \text{Nat}(G_2 \circ F_2, G_3 \circ F_3) \times \text{Nat}(G_1 \circ F_1, G_2 \circ F_2) \\ & \searrow \star_{F_1, F_3, G_1, G_3} & \swarrow \circ_{G_1 \circ F_1, G_2 \circ F_2, G_3 \circ F_3} \\ & \text{Nat}(G_1 \circ F_1, G_3 \circ F_3) & \end{array}$$

commutes, i.e. given a diagram

$$\begin{array}{ccccc} & F_1 & & G_1 & \\ & \alpha \Downarrow & & \beta \Downarrow & \\ C & \xrightarrow{F_2} & \mathcal{D} & \xrightarrow{G_2} & \mathcal{E} \\ & \alpha' \Downarrow & \uparrow & \beta' \Downarrow & \\ & F_3 & & G_3 & \end{array}$$

in  $\text{Cats}_2$ , we have

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

**PROOF 11.9.5.1.5 ▶ PROOF OF PROPOSITION 11.9.5.1.4**

Item 1: Functionality

Omitted.

**Item 2: Associativity**

Omitted.

**Item 3: Interaction With Identities**

We have

$$\begin{aligned} (\text{id}_G \star \text{id}_F)_A &\stackrel{\text{def}}{=} (\text{id}_G)_{F_A} \circ G(\text{id}_F)_A \\ &\stackrel{\text{def}}{=} \text{id}_{G_{F_A}} \circ G\text{id}_{F_A} \\ &= \text{id}_{G_{F_A}} \circ \text{id}_{G_{F_A}} \\ &= \text{id}_{G_{F_A}} \\ &\stackrel{\text{def}}{=} (\text{id}_{G \circ F})_A \end{aligned}$$

for each  $A \in \text{Obj}(C)$ , showing the desired equality.

**Item 4: Middle Four Exchange**

Let  $A \in \text{Obj}(C)$  and consider the diagram

$$\begin{array}{ccccc}
 & & G_1(F_3(A)) & & \\
 & \nearrow G_1(\alpha'_A) & & \searrow \beta_{F_3(A)} & \\
 G_1(F_1(A)) & \xrightarrow{G_1(\alpha_A)} & G_1(F_2(A)) & (1) & G_2(F_3(A)) \xrightarrow{\beta'_{F_3(A)}} G_3(F_3(A)). \\
 & \searrow \beta_{F_2(A)} & & \nearrow G_2(\alpha'_A) & \\
 & & G_2(F_2(A)) & &
 \end{array}$$

The top composition

$$\begin{array}{ccccc}
 & & G_1(F_3(A)) & & \\
 & \nearrow G_1(\alpha'_A) & & \searrow \beta_{F_3(A)} & \\
 G_1(F_1(A)) & \xrightarrow{G_1(\alpha_A)} & G_1(F_2(A)) & (1) & G_2(F_3(A)) \xrightarrow{\beta'_{F_3(A)}} G_3(F_3(A)). \\
 & \searrow \beta_{F_2(A)} & & \nearrow G_2(\alpha'_A) & \\
 & & G_2(F_2(A)) & &
 \end{array}$$

is given by  $((\beta' \circ \beta) \star (\alpha' \circ \alpha))_A$ , while the bottom composition

$$\begin{array}{ccccc}
 & & G_1(F_3(A)) & & \\
 & \nearrow G_1(\alpha'_A) & & \searrow \beta_{F_3(A)} & \\
 G_1(F_1(A)) & \xrightarrow{G_1(\alpha_A)} & G_1(F_2(A)) & (1) & G_2(F_3(A)) \xrightarrow{\beta'_{F_3(A)}} G_3(F_3(A)). \\
 & \searrow \beta_{F_2(A)} & & \nearrow G_2(\alpha'_A) & \\
 & & G_2(F_2(A)) & &
 \end{array}$$

is given by  $((\beta' \star \alpha') \circ (\beta \star \alpha))_A$ . Now, Subdiagram (1) corresponds to the naturality condition

$$\begin{array}{ccc}
 G_1(F_2(A)) & \xrightarrow{G_1(\alpha'_A)} & G_1(F_3(A)) \\
 G_2(\alpha'_A) \circ \beta_{F_2(A)} = \beta_{F_3(A)} \circ G_1(\alpha'_A), & \beta_{F_2(A)} \downarrow & \downarrow \beta_{F_3(A)} \\
 G_2(F_2(A)) & \xrightarrow[G_2(\alpha'_A)]{} & G_2(F_3(A))
 \end{array}$$

for  $\beta: G_1 \Rightarrow G_2$  at  $\alpha'_A: F_2(A) \rightarrow F_3(A)$ , and thus commutes. Thus we have

$$((\beta' \circ \beta) \star (\alpha' \circ \alpha))_A = ((\beta' \star \alpha') \circ (\beta \star \alpha))_A$$

for each  $A \in \text{Obj}(C)$  and therefore

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

This finishes the proof. □

### 018C 11.9.6 Properties of Natural Transformations

018D

#### PROPOSITION 11.9.6.1.1 ► NATURAL TRANSFORMATIONS AS CATEGORICAL HOMOTOPIES

Let  $F, G: C \rightrightarrows \mathcal{D}$  be functors. The following data are equivalent:<sup>1</sup>

018E

1. A natural transformation  $\alpha: F \Rightarrow G$ .

018F

2. A functor  $[\alpha]: C \rightarrow \mathcal{D}^{\mathbb{1}}$  filling the diagram

$$\begin{array}{ccc}
 & \mathcal{D} & \\
 F \nearrow & \uparrow ev_0 & \\
 C & \xrightarrow{[\alpha]} & \mathcal{D}^{\mathbb{1}}. \\
 G \searrow & \downarrow ev_1 & \\
 & \mathcal{D} &
 \end{array}$$

018G

3. A functor  $[\alpha]: C \times \mathbb{1} \rightarrow \mathcal{D}$  filling the diagram

$$\begin{array}{ccc}
 C & & \\
 \uparrow ev_0 & \searrow F & \\
 C \times \mathbb{1} & \xrightarrow{[\alpha]} & \mathcal{D}. \\
 \downarrow ev_1 & \nearrow G & \\
 C & &
 \end{array}$$

<sup>1</sup>Taken from [MO 64365].

## PROOF 11.9.6.1.2 ► PROOF OF PROPOSITION 11.9.6.1.1

From Item 1 to Item 2 and Back

We may identify  $\mathcal{D}^{\mathbb{1}}$  with  $\text{Arr}(\mathcal{D})$ . Given a natural transformation  $\alpha: F \Rightarrow G$ , we have a functor

$$\begin{aligned}
 [\alpha]: C &\longrightarrow \mathcal{D}^{\mathbb{1}} \\
 A &\longmapsto \alpha_A \\
 (f: A \rightarrow B) &\longmapsto \left( \begin{array}{ccc} F_A & \xrightarrow{F_f} & F_B \\ \alpha_A \downarrow & & \downarrow \alpha_B \\ G_A & \xrightarrow{G_f} & G_B \end{array} \right)
 \end{aligned}$$

making the diagram in [Item 2](#) commute. Conversely, every such functor gives rise to a natural transformation from  $F$  to  $G$ , and these constructions are inverse to each other.

[From Item 2 to Item 3 and Back](#)

This follows from [Item 3](#) of [Proposition 11.10.1.1.2](#). 

### 018H 11.9.7 Natural Isomorphisms

Let  $C$  and  $\mathcal{D}$  be categories and let  $F, G: C \Rightarrow \mathcal{D}$  be functors.

#### 018J DEFINITION 11.9.7.1.1 ► NATURAL ISOMORPHISMS

A natural transformation  $\alpha: F \Rightarrow G$  is a **natural isomorphism** if there exists a natural transformation  $\alpha^{-1}: G \Rightarrow F$  such that

$$\begin{aligned}\alpha^{-1} \circ \alpha &= \text{id}_F, \\ \alpha \circ \alpha^{-1} &= \text{id}_G.\end{aligned}$$

#### 018K PROPOSITION 11.9.7.1.2 ► PROPERTIES OF NATURAL ISOMORPHISMS

Let  $\alpha: F \Rightarrow G$  be a natural transformation.

018L 1. *Characterisations.* The following conditions are equivalent:

- 018M (a) The natural transformation  $\alpha$  is a natural isomorphism.  
018N (b) For each  $A \in \text{Obj}(C)$ , the morphism  $\alpha_A: F_A \rightarrow G_A$  is an isomorphism.

018P 2. *Componentwise Inverses of Natural Transformations Assemble Into Natural Transformations.* Let  $\alpha^{-1}: G \Rightarrow F$  be a transformation such that, for each  $A \in \text{Obj}(C)$ , we have

$$\begin{aligned}\alpha_A^{-1} \circ \alpha_A &= \text{id}_{F(A)}, \\ \alpha_A \circ \alpha_A^{-1} &= \text{id}_{G(A)}.\end{aligned}$$

Then  $\alpha^{-1}$  is a natural transformation.

#### PROOF 11.9.7.1.3 ► PROOF OF PROPOSITION 11.9.7.1.2

[Item 1: Characterisations](#)

The implication [Item 1a](#)  $\implies$  [Item 1b](#) is clear, whereas the implication

**Item 1b**  $\implies$  **Item 1a** follows from **Item 2**.

## Item 2: Componentwise Inverses of Natural Transformations Assemblies

The naturality condition for  $\alpha^{-1}$  corresponds to the commutativity of the diagram

$$\begin{array}{ccc} G(A) & \xrightarrow{G(f)} & G(B) \\ \alpha_A^{-1} \downarrow & & \downarrow \sigma_B^{-1} \\ F(A) & \xrightarrow[F(f)]{} & F(B) \end{array}$$

for each  $A, B \in \text{Obj}(C)$  and each  $f \in \text{Hom}_C(A, B)$ . Considering the diagram

$$\begin{array}{ccc}
 G(A) & \xrightarrow{G(f)} & G(B) \\
 \alpha_A^{-1} \downarrow & (1) & \downarrow \alpha_B^{-1} \\
 F(A) & \xrightarrow{F(f)} & F(B) \\
 \alpha_A \downarrow & (2) & \downarrow \alpha_B \\
 G(A) & \xrightarrow{G(f)} & G(B),
 \end{array}$$

where the boundary diagram as well as Subdiagram (2) commute, we have

$$\begin{aligned} G(f) &= G(f) \circ \text{id}_{G(A)} \\ &= G(f) \circ \alpha_A \circ \alpha_A^{-1} \\ &= \alpha_B \circ F(f) \circ \alpha_A^{-1}. \end{aligned}$$

Postcomposing both sides with  $\alpha_B^{-1}$ , we get

$$\begin{aligned}\alpha_B^{-1} \circ G(f) &= \alpha_B^{-1} \circ \alpha_B \circ F(f) \circ \alpha_A^{-1} \\ &= \text{id}_{F(B)} \circ F(f) \circ \alpha_A^{-1} \\ &= F(f) \circ \alpha_A^{-1},\end{aligned}$$

which is the naturality condition we wanted to show. Thus  $\alpha^{-1}$  is a natural transformation. 

**018Q 11.10 Categories of Categories****018R 11.10.1 Functor Categories**

Let  $C$  be a category and  $\mathcal{D}$  be a small category.

018S

## DEFINITION 11.10.1.1.1 ► FUNCTOR CATEGORIES

The **category of functors from  $C$  to  $\mathcal{D}$** <sup>1</sup> is the category  $\text{Fun}(C, \mathcal{D})$ <sup>2</sup> where

- *Objects.* The objects of  $\text{Fun}(C, \mathcal{D})$  are functors from  $C$  to  $\mathcal{D}$ .
- *Morphisms.* For each  $F, G \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$ , we have

$$\text{Hom}_{\text{Fun}(C, \mathcal{D})}(F, G) \stackrel{\text{def}}{=} \text{Nat}(F, G).$$

- *Identities.* For each  $F \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$ , the unit map

$$\mathbb{1}_F^{\text{Fun}(C, \mathcal{D})} : \text{pt} \rightarrow \text{Nat}(F, F)$$

of  $\text{Fun}(C, \mathcal{D})$  at  $F$  is given by

$$\text{id}_F^{\text{Fun}(C, \mathcal{D})} \stackrel{\text{def}}{=} \text{id}_F,$$

where  $\text{id}_F : F \Rightarrow F$  is the identity natural transformation of  $F$  of [Example 11.9.3.1.1](#).

- *Composition.* For each  $F, G, H \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$ , the composition map

$$\circ_{F, G, H}^{\text{Fun}(C, \mathcal{D})} : \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H)$$

of  $\text{Fun}(C, \mathcal{D})$  at  $(F, G, H)$  is given by

$$\beta \circ_{F, G, H}^{\text{Fun}(C, \mathcal{D})} \alpha \stackrel{\text{def}}{=} \beta \circ \alpha,$$

where  $\beta \circ \alpha$  is the vertical composition of  $\alpha$  and  $\beta$  of [Item 1](#) of [Proposition 11.9.4.1.3](#).

<sup>1</sup>Further Terminology: Also called the **functor category**  $\text{Fun}(C, \mathcal{D})$ .

<sup>2</sup>Further Notation: Also written  $\mathcal{D}^C$  and  $[C, \mathcal{D}]$ .

018T

## PROPOSITION 11.10.1.1.2 ► PROPERTIES OF FUNCTOR CATEGORIES

Let  $C$  and  $\mathcal{D}$  be categories and let  $F : C \rightarrow \mathcal{D}$  be a functor.

018U

1. *Functionality.* The assignments  $C, \mathcal{D}, (C, \mathcal{D}) \mapsto \text{Fun}(C, \mathcal{D})$  define functors

$$\begin{aligned} \text{Fun}(C, -) : & \quad \text{Cats} \rightarrow \text{Cats}, \\ \text{Fun}(-, \mathcal{D}) : & \quad \text{Cats}^{\text{op}} \rightarrow \text{Cats}, \\ \text{Fun}(-_1, -_2) : & \quad \text{Cats}^{\text{op}} \times \text{Cats} \rightarrow \text{Cats}. \end{aligned}$$

018V

2. *2-Functionality.* The assignments  $C, \mathcal{D}, (C, \mathcal{D}) \mapsto \text{Fun}(C, \mathcal{D})$  define 2-functors

$$\begin{aligned}\text{Fun}(C, -) : \quad \text{Cats}_2 &\rightarrow \text{Cats}_2, \\ \text{Fun}(-, \mathcal{D}) : \quad \text{Cats}_2^{\text{op}} &\rightarrow \text{Cats}_2, \\ \text{Fun}(-_1, -_2) : \quad \text{Cats}_2^{\text{op}} \times \text{Cats}_2 &\rightarrow \text{Cats}_2.\end{aligned}$$

018W

3. *Adjointness.* We have adjunctions

$$\begin{aligned}(C \times - \dashv \text{Fun}(C, -)) : \quad \text{Cats} &\begin{array}{c} \xrightarrow{C \times -} \\ \perp \\ \xleftarrow{\text{Fun}(C, -)} \end{array} \text{Cats}, \\ (- \times \mathcal{D} \dashv \text{Fun}(\mathcal{D}, -)) : \quad \text{Cats} &\begin{array}{c} \xrightarrow{- \times \mathcal{D}} \\ \perp \\ \xleftarrow{\text{Fun}(\mathcal{D}, -)} \end{array} \text{Cats},\end{aligned}$$

witnessed by bijections of sets

$$\begin{aligned}\text{Hom}_{\text{Cats}}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Hom}_{\text{Cats}}(\mathcal{D}, \text{Fun}(C, \mathcal{E})), \\ \text{Hom}_{\text{Cats}}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Hom}_{\text{Cats}}(C, \text{Fun}(\mathcal{D}, \mathcal{E})),\end{aligned}$$

natural in  $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$ .

018X

4. *2-Adjointness.* We have 2-adjunctions

$$\begin{aligned}(C \times - \dashv \text{Fun}(C, -)) : \quad \text{Cats}_2 &\begin{array}{c} \xrightarrow{C \times -} \\ \perp_2 \\ \xleftarrow{\text{Fun}(C, -)} \end{array} \text{Cats}_2, \\ (- \times \mathcal{D} \dashv \text{Fun}(\mathcal{D}, -)) : \quad \text{Cats}_2 &\begin{array}{c} \xrightarrow{- \times \mathcal{D}} \\ \perp_2 \\ \xleftarrow{\text{Fun}(\mathcal{D}, -)} \end{array} \text{Cats}_2,\end{aligned}$$

witnessed by isomorphisms of categories

$$\begin{aligned}\text{Fun}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Fun}(\mathcal{D}, \text{Fun}(C, \mathcal{E})), \\ \text{Fun}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Fun}(C, \text{Fun}(\mathcal{D}, \mathcal{E})),\end{aligned}$$

natural in  $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats}_2)$ .

018Y

5. *Interaction With Punctual Categories.* We have a canonical isomorphism of categories

$$\text{Fun}(\text{pt}, C) \cong C,$$

natural in  $C \in \text{Obj}(\text{Cats})$ .

018Z

6. *Objectwise Computation of Co/Limits.* Let

$$D: \mathcal{I} \rightarrow \text{Fun}(C, \mathcal{D})$$

be a diagram in  $\text{Fun}(C, \mathcal{D})$ . We have isomorphisms

$$\lim(D)_A \cong \lim_{i \in \mathcal{I}}(D_i(A)),$$

$$\text{colim}(D)_A \cong \text{colim}_{i \in \mathcal{I}}(D_i(A)),$$

naturally in  $A \in \text{Obj}(C)$ .

0190

7. *Interaction With Co/Completeness.* If  $\mathcal{E}$  is co/complete, then so is  $\text{Fun}(C, \mathcal{E})$ .

0191

8. *Monomorphisms and Epimorphisms.* Let  $\alpha: F \Rightarrow G$  be a morphism of  $\text{Fun}(C, \mathcal{D})$ . The following conditions are equivalent:

0192

(a) The natural transformation

$$\alpha: F \Rightarrow G$$

is a monomorphism (resp. epimorphism) in  $\text{Fun}(C, \mathcal{D})$ .

0193

(b) For each  $A \in \text{Obj}(C)$ , the morphism

$$\alpha_A: F_A \rightarrow G_A$$

is a monomorphism (resp. epimorphism) in  $\mathcal{D}$ .

#### PROOF 11.10.1.1.3 ► PROOF OF PROPOSITION 11.10.1.1.2

Item 1: Functoriality

Omitted.

Item 2: 2-Functoriality

Omitted.

Item 3: Adjointness

Omitted.

Item 4: 2-Adjointness

Omitted.

Item 5: Interaction With Punctual Categories

Omitted.

Item 6: Objectwise Computation of Co/Limits

Omitted.

Item 7: Interaction With Co/Completeness

This follows from ??.

Item 8: Monomorphisms and Epimorphisms

Omitted.



### 0194 11.10.2 The Category of Categories and Functors

0195

#### DEFINITION 11.10.2.1.1 ► THE CATEGORY OF CATEGORIES AND FUNCTORS

The **category of (small) categories and functors** is the category  $\text{Cats}$  where

- *Objects.* The objects of  $\text{Cats}$  are small categories.
- *Morphisms.* For each  $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ , we have

$$\text{Hom}_{\text{Cats}}(C, \mathcal{D}) \stackrel{\text{def}}{=} \text{Obj}(\text{Fun}(C, \mathcal{D})).$$

- *Identities.* For each  $C \in \text{Obj}(\text{Cats})$ , the unit map

$$1_C^{\text{Cats}}: \text{pt} \rightarrow \text{Hom}_{\text{Cats}}(C, C)$$

of  $\text{Cats}$  at  $C$  is defined by

$$\text{id}_C^{\text{Cats}} \stackrel{\text{def}}{=} \text{id}_C,$$

where  $\text{id}_C: C \rightarrow C$  is the identity functor of  $C$  of [Example 11.5.1.1.4](#).

- *Composition.* For each  $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$ , the composition map

$$\circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}}: \text{Hom}_{\text{Cats}}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}}(C, \mathcal{D}) \rightarrow \text{Hom}_{\text{Cats}}(C, \mathcal{E})$$

of  $\text{Cats}$  at  $(C, \mathcal{D}, \mathcal{E})$  is given by

$$G \circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}} F \stackrel{\text{def}}{=} G \circ F,$$

where  $G \circ F: C \rightarrow \mathcal{E}$  is the composition of  $F$  and  $G$  of [Definition 11.5.1.1.6](#).

0196

**PROPOSITION 11.10.2.1.2 ► PROPERTIES OF THE CATEGORY Cats**

Let  $\mathcal{C}$  be a category.

0197

1. *Co/Completeness*. The category Cats is complete and cocomplete.

0198

2. *Cartesian Monoidal Structure*. The quadruple  $(\text{Cats}, \times, \text{pt}, \text{Fun})$  is a Cartesian closed monoidal category.

**PROOF 11.10.2.1.3 ► PROOF OF PROPOSITION 11.10.2.1.2**

Item 1: Co/Completeness

Omitted.

Item 2: Cartesian Monoidal Structure

Omitted. 

### 0199 11.10.3 The 2-Category of Categories, Functors, and Natural Transformations

019A

**DEFINITION 11.10.3.1.1 ► THE 2-CATEGORY OF CATEGORIES**

The **2-category of (small) categories, functors, and natural transformations** is the 2-category  $\text{Cats}_2$  where

- *Objects*. The objects of  $\text{Cats}_2$  are small categories.
- *Hom-Categories*. For each  $\mathcal{C}, \mathcal{D} \in \text{Obj}(\text{Cats}_2)$ , we have

$$\text{Hom}_{\text{Cats}_2}(\mathcal{C}, \mathcal{D}) \stackrel{\text{def}}{=} \text{Fun}(\mathcal{C}, \mathcal{D}).$$

- *Identities*. For each  $\mathcal{C} \in \text{Obj}(\text{Cats}_2)$ , the unit functor

$$\mathbb{1}_C^{\text{Cats}_2} : \text{pt} \rightarrow \text{Fun}(\mathcal{C}, \mathcal{C})$$

of  $\text{Cats}_2$  at  $\mathcal{C}$  is the functor picking the identity functor  $\text{id}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$  of  $\mathcal{C}$ .

- *Composition*. For each  $\mathcal{C}, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats}_2)$ , the composition bifunctor

$$\circ_{\mathcal{C}, \mathcal{D}, \mathcal{E}}^{\text{Cats}_2} : \text{Hom}_{\text{Cats}_2}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}_2}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Hom}_{\text{Cats}_2}(\mathcal{C}, \mathcal{E})$$

of  $\text{Cats}_2$  at  $(\mathcal{C}, \mathcal{D}, \mathcal{E})$  is the functor where

- *Action on Objects*. For each object  $(G, F) \in \text{Obj}(\text{Hom}_{\text{Cats}_2}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}_2}(\mathcal{C}, \mathcal{D}))$ , we have

$$\circ_{\mathcal{C}, \mathcal{D}, \mathcal{E}}^{\text{Cats}_2}(G, F) \stackrel{\text{def}}{=} G \circ F.$$

- *Action on Morphisms.* For each morphism  $(\beta, \alpha) : (K, H) \Rightarrow (G, F)$  of  $\text{Hom}_{\text{Cats}_2}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}_2}(\mathcal{C}, \mathcal{D})$ , we have

$$\circ_{\mathcal{C}, \mathcal{D}, \mathcal{E}}^{\text{Cats}_2}(\beta, \alpha) \stackrel{\text{def}}{=} \beta \star \alpha,$$

where  $\beta \star \alpha$  is the horizontal composition of  $\alpha$  and  $\beta$  of [Definition 11.9.5.1.1](#).

019B

**PROPOSITION 11.10.3.1.2 ► PROPERTIES OF THE 2-CATEGORY  $\text{Cats}_2$** 

Let  $C$  be a category.

019C

1. *2-Categorical Co/Completeness.* The 2-category  $\text{Cats}_2$  is complete and cocomplete as a 2-category, having all 2-categorical and bicategorical co/limits.

**PROOF 11.10.3.1.3 ► PROOF OF PROPOSITION 11.10.3.1.2**

Item 1: Co/Completeness

Omitted. 

019D **11.10.4 The Category of Groupoids**

019E

**DEFINITION 11.10.4.1.1 ► THE CATEGORY OF SMALL GROUPOIDS**

The **category of (small) groupoids** is the full subcategory  $\text{Grpd}$  of  $\text{Cats}$  spanned by the groupoids.

019F **11.10.5 The 2-Category of Groupoids**

019G

**DEFINITION 11.10.5.1.1 ► THE 2-CATEGORY OF SMALL GROUPOIDS**

The **2-category of (small) groupoids** is the full sub-2-category  $\text{Grpd}_2$  of  $\text{Cats}_2$  spanned by the groupoids.

# Appendices

## 11.A Other Chapters

<b>Preliminaries</b>	10. Conditions on Relations
1. Introduction	
2. A Guide to the Literature	
<b>Sets</b>	<b>Categories</b>
3. Sets	11. Categories
4. Constructions With Sets	
5. Monoidal Structures on the Category of Sets	12. Presheaves and the Yoneda Lemma
6. Pointed Sets	
7. Tensor Products of Pointed Sets	
<b>Relations</b>	<b>Monoidal Categories</b>
8. Relations	13. Constructions With Monoidal Categories
9. Constructions With Relations	
	<b>Bicategories</b>
	14. Types of Morphisms in Bicategories
	<b>Extra Part</b>
	15. Notes

## Chapter 12

# Presheaves and the Yoneda Lemma

**02H3** This chapter contains some material about presheaves and the Yoneda lemma.

This chapter is under revision. TODO:

1. Subsection properties of categories of copresheaves
2. Adjointness of tensor product of functors
3. Limit of category of elements (instead of colimit)
4. Category of elements where objects are natural transformations  $\mathcal{F} \Rightarrow h_x$  instead of the other way around. Is this related to Isbell duality?
5. Motivate the proof of the Yoneda lemma as in Martin's comment here:  
[https://mathoverflow.net/questions/130883/are-there-proofs-that-you-feel-you-did-not-understand-for-a-long-time#comment360113\\_131050](https://mathoverflow.net/questions/130883/are-there-proofs-that-you-feel-you-did-not-understand-for-a-long-time#comment360113_131050)
6. Add discussion of universal properties
7. Add  $h_{g \circ f} = h_g \circ h_f$  to properties of representable natural transformations

## Contents

---

12.1 Presheaves .....	837
12.1.1 Foundations .....	837
12.1.2 Representable Presheaves.....	838
12.1.3 Representable Natural Transformations .....	840
12.1.4 The Yoneda Embedding .....	840
12.1.5 The Yoneda Lemma .....	843
12.1.6 Properties of Categories of Presheaves.....	848
12.2 Copresheaves .....	849
12.2.1 Foundations .....	849

12.2.2 Corepresentable Copresheaves .....	850
12.2.3 Corepresentable Natural Transformations .....	852
12.2.4 The Contravariant Yoneda Embedding .....	852
12.2.5 The Contravariant Yoneda Lemma .....	854
<b>12.3 Restricted Yoneda Embeddings and Yoneda Extensions .....</b>	<b>854</b>
12.3.1 Foundations .....	854
12.3.2 The Yoneda Extension Functor .....	858
<b>12.4 Functor Tensor Products .....</b>	<b>861</b>
12.4.1 The Tensor Product of Presheaves With Copresheaves	
861	
12.4.2 The Tensor of a Presheaf With a Functor .....	864
12.4.3 The Tensor of a Copresheaf With a Functor .....	865
<b>12.A Other Chapters .....</b>	<b>867</b>

## 02H4 12.1 Presheaves

### 02H5 12.1.1 Foundations

Let  $C$  be a category.

#### 02H6 DEFINITION 12.1.1.1 ► PRESHEAVES ON A CATEGORY

A **presheaf on  $C$**  is a functor  $\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets}$ .

#### 02H7 EXAMPLE 12.1.1.2 ► PRESHEAVES ON ONE-OBJECT CATEGORIES

Presheaves on the delooping BA of a monoid  $A$  are precisely the left  $A$ -sets; see ??.

#### 02H8 DEFINITION 12.1.1.3 ► MORPHISMS OF PRESHEAVES

A **morphism of presheaves** on  $C$  from  $\mathcal{F}$  to  $\mathcal{G}$  is a natural transformation  $\alpha: \mathcal{F} \Rightarrow \mathcal{G}$ .

#### 02H9 DEFINITION 12.1.1.4 ► THE CATEGORY OF PRESHEAVES ON A CATEGORY

The **category of presheaves on  $C$**  is the category  $\text{PSh}(C)$ <sup>1</sup> defined by

$$\text{PSh}(C) \stackrel{\text{def}}{=} \text{Fun}(C^{\text{op}}, \text{Sets}).$$

<sup>1</sup>Further Notation: Also written  $\widehat{C}$  in some parts of the literature.

02HA

**REMARK 12.1.1.5 ▶ UNWINDING DEFINITION 12.1.1.4**

In detail, the **category of presheaves on  $C$**  is the category  $\text{PSh}(C)$  where

- *Objects.* The objects of  $\text{PSh}(C)$  are presheaves on  $C$  as in [Definition 12.1.1.1](#).
- *Morphisms.* The morphisms of  $\text{PSh}(C)$  are morphisms of presheaves as in [Definition 12.1.1.3](#), i.e. we have

$$\text{Hom}_{\text{PSh}(C)}(\mathcal{F}, \mathcal{G}) \stackrel{\text{def}}{=} \text{Nat}(\mathcal{F}, \mathcal{G})$$

for each  $\mathcal{F}, \mathcal{G} \in \text{Obj}(\text{PSh}(C))$ .

- *Identities.* For each  $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$ , the unit map

$$1_{\mathcal{F}}^{\text{PSh}(C)} : \text{pt} \rightarrow \text{Nat}(\mathcal{F}, \mathcal{F})$$

of  $\text{PSh}(C)$  at  $\mathcal{F}$  is defined by

$$\text{id}_{\mathcal{F}}^{\text{PSh}(C)} \stackrel{\text{def}}{=} \text{id}_{\mathcal{F}},$$

where  $\text{id}_{\mathcal{F}} : \mathcal{F} \Rightarrow \mathcal{F}$  is the identity natural transformation of [Example 11.9.3.1.1](#).

- *Composition.* For each  $\mathcal{F}, \mathcal{G}, \mathcal{H} \in \text{Obj}(\text{PSh}(C))$ , the composition map

$$\circ^{\text{PSh}(C)}_{\mathcal{F}, \mathcal{G}, \mathcal{H}} : \text{Nat}(\mathcal{G}, \mathcal{H}) \times \text{Nat}(\mathcal{F}, \mathcal{G}) \rightarrow \text{Nat}(\mathcal{F}, \mathcal{H})$$

of  $\text{PSh}(C)$  at  $(\mathcal{F}, \mathcal{G}, \mathcal{H})$  is defined by

$$\beta \circ^{\text{PSh}(C)}_{\mathcal{F}, \mathcal{G}, \mathcal{H}} \alpha \stackrel{\text{def}}{=} \beta \circ \alpha,$$

where  $\beta \circ \alpha : \mathcal{F} \Rightarrow \mathcal{H}$  is the vertical composition of  $\alpha$  and  $\beta$  of [Definition 11.9.4.1.1](#).

02HB **12.1.2 Representable Presheaves**

Let  $C$  be a category.

02HC

**DEFINITION 12.1.2.1.1 ▶ REPRESENTABLE PRESHEAVES**

Let  $A \in \text{Obj}(C)$ .

02HD

1. The **representable presheaf associated to  $A$**  is the presheaf

$$h_A: \mathcal{C}^{\text{op}} \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each  $X \in \text{Obj}(\mathcal{C})$ , we have

$$h_A(X) \stackrel{\text{def}}{=} \text{Hom}_{\mathcal{C}}(X, A).$$

- *Action on Morphisms.* For each  $X, Y \in \text{Obj}(\mathcal{C})$ , the action on morphisms

$$h_{A|X,Y}: \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\text{Sets}}(h_A(Y), h_A(X))$$

of  $h_A$  at  $(X, Y)$  is given by sending a morphism

$$f: X \rightarrow Y$$

of  $\mathcal{C}$  to the map of sets

$$h_A(f): \underbrace{h_A(Y)}_{\stackrel{\text{def}}{=} \text{Hom}_{\mathcal{C}}(Y, A)} \rightarrow \underbrace{h_A(X)}_{\stackrel{\text{def}}{=} \text{Hom}_{\mathcal{C}}(X, A)}$$

defined by

$$h_A(f) \stackrel{\text{def}}{=} f^*,$$

where  $f^*$  is the precomposition by  $f$  morphism of **Item 1** of **Definition 11.1.4.1.1**.

02HE

2. A **representing object** for a presheaf  $\mathcal{F}: \mathcal{C}^{\text{op}} \rightarrow \text{Sets}$  on  $\mathcal{C}$  is an object  $A$  of  $\mathcal{C}$  such that we have  $\mathcal{F} \cong h_A$ .

02HF

3. A presheaf  $\mathcal{F}: \mathcal{C}^{\text{op}} \rightarrow \text{Sets}$  on  $\mathcal{C}$  is **representable** if  $\mathcal{F}$  admits a representing object.

02HG

#### EXAMPLE 12.1.2.1.2 ► REPRESENTABLE PRESHEAVES ON ONE-OBJECT CATEGORIES

The representable presheaf on the delooping  $\text{BA}$  of a monoid  $A$  associated to the unique object  $\bullet$  of  $\text{BA}$  is the left regular representation of  $A$  of  $\text{??}$ .

02HH

#### PROPOSITION 12.1.2.1.3 ► UNIQUENESS OF REPRESENTING OBJECTS UP TO ISOMORPHISM

Let  $\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets}$  be a presheaf. If there exist  $A, B \in \text{Obj}(C)$  such that we have natural isomorphisms

$$\begin{aligned} h_A &\cong \mathcal{F}, \\ h_B &\cong \mathcal{F}, \end{aligned}$$

then  $A \cong B$ .

#### PROOF 12.1.2.1.4 ► PROOF OF PROPOSITION 12.1.2.1.3

By composing the isomorphisms  $h_A \cong \mathcal{F} \cong h_B$ , we get a natural isomorphism  $h_A \cong h_B$ . By Item 2 of Proposition 12.1.4.1.3, we have  $A \cong B$ .



### 02HJ 12.1.3 Representable Natural Transformations

Let  $C$  be a category, let  $A, B \in \text{Obj}(C)$ , and let  $f: A \rightarrow B$  be a morphism of  $C$ .

#### 02HK DEFINITION 12.1.3.1.1 ► REPRESENTABLE NATURAL TRANSFORMATIONS

The **representable natural transformation associated to  $f$**  is the natural transformation

$$h_f: h_A \Rightarrow h_B$$

consisting of the collection

$$\left\{ h_{f|X}: \underbrace{h_A(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(X, A)} \rightarrow \underbrace{h_B(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(X, B)} \right\}_{X \in \text{Obj}(C)}$$

with

$$h_{f|X} \stackrel{\text{def}}{=} f_*,$$

where  $f_*$  is the postcomposition by  $f$  morphism of Item 2 of Definition 11.1.4.1.1.

### 02HL 12.1.4 The Yoneda Embedding

#### 02HM DEFINITION 12.1.4.1.1 ► THE YONEDA EMBEDDING

The **Yoneda embedding** of  $C^{\text{op}}$  is the functor<sup>2</sup>

$$\mathfrak{y}_C: C \hookrightarrow \text{PSh}(C)$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(C)$ , we have

$$\mathfrak{J}_C(A) \stackrel{\text{def}}{=} h_A.$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(C)$ , the action on morphisms

$$\mathfrak{J}_{C|A,B} : \text{Hom}_C(A, B) \rightarrow \text{Nat}(h_A, h_B)$$

of  $\mathfrak{J}_C$  at  $(A, B)$  is given by

$$\mathfrak{J}_{C|A,B}(f) \stackrel{\text{def}}{=} h_f$$

for each  $f \in \text{Hom}_C(A, B)$ , where  $h_f$  is the representable natural transformation associated to  $f$  of [Definition 12.1.3.1.1](#).

<sup>1</sup>Further Terminology: Also called the **covariant Yoneda embedding** to distinguish it from the contravariant Yoneda embedding of [Theorem 12.2.5.1.1](#).

<sup>2</sup>Further Notation: Also written  $h_{(-)}$ , or simply  $\mathfrak{J}$ .

02HN

#### REMARK 12.1.4.1.2 ► ON THE USAGE OF $\mathfrak{J}$ TO DENOTE THE YONEDA EMBEDDING

The notation  $\mathfrak{J}$  for the Yoneda embedding was first introduced in [\[JS17\]](#). The symbol  $\mathfrak{J}$  is the [hiragana for yo](#), and comes from “Yoneda” in Nobuo Yoneda (米田信夫).

It is pronounced *yo* but without letting the “o” in *yo* sound like an o-u diphthong:

- See [here](#).
- IPA transcription: [jø].

02HP

#### PROPOSITION 12.1.4.1.3 ► PROPERTIES OF THE YONEDA EMBEDDING

Let  $C$  be a category.

02HQ

1. *Fully Faithfulness.* The Yoneda embedding

$$\mathfrak{J}_C : C \hookrightarrow \text{PSh}(C)$$

is fully faithful.

02HR

2. *Preservation and Reflection of Isomorphisms.* The Yoneda embedding

$$\mathfrak{J}_C : C \hookrightarrow \text{PSh}(C)$$

preserves and reflects isomorphisms, i.e. given  $A, B \in \text{Obj}(C)$ , the following conditions are equivalent:

02HS

(a) We have  $A \cong B$ .

02HT

(b) We have  $h_A \cong h_B$ .

02HU

3. *Density.* The Yoneda embedding

$$\jmath_C : C \hookrightarrow \mathbf{PSh}(C)$$

is dense.

02HV

4. *Interaction With Density Comonads.* We have

$$\begin{array}{ccc} & \mathbf{PSh}(C) & \\ \text{Lan}_{\jmath_C}(\jmath_C) \cong \text{id}_{\mathbf{PSh}(C)}, & \nearrow \jmath_C & \downarrow \text{Lan}_{\jmath_C}(\jmath_C) \\ C & \xrightarrow{\quad \jmath_C \quad} & \mathbf{PSh}(C). \end{array}$$

02HW

5. *Interaction With Codensity Monads.* We have

$$\text{Ran}_{\jmath_C}(\jmath_C) \cong \text{Spec} \circ \text{O},$$

where  $\text{Spec}$  and  $\text{O}$  are the functors of ??.

## PROOF 12.1.4.1.4 ► PROOF OF PROPOSITION 12.1.4.1.3

## Item 1: Fully Faithfulness

Let  $A, B \in \text{Obj}(C)$ . Applying the Yoneda lemma ([Theorem 12.1.5.1.1](#)) to the functor  $h_B$  (i.e. in the case  $\mathcal{F} = h_B$ ), we have

$$\text{Hom}_C(A, B) \cong \text{Nat}(h_A, h_B),$$

and the natural isomorphism

$$\xi_{A,B} : h_B(A) \Rightarrow \text{Nat}(h_A, h_B)$$

witnessing this bijection is given by

$$\begin{aligned} \xi_{A,B}(g)_X &\stackrel{\text{def}}{=} h_B^X(g) \\ &\stackrel{\text{def}}{=} g_* \end{aligned}$$

for each  $X \in \text{Obj}(C)$  and each  $g \in h_B^X$ , i.e. we have  $\xi_{A,B} = \jmath_{C|A,B}$ . Thus  $\jmath_C$  is fully faithful.

Item 2: Preservation and Reflection of Isomorphisms

This follows from [Item 1 of Proposition 11.5.1.1.8](#) and [Item 3 of Proposition 11.6.3.1.2](#).

Item 3: Density

Omitted.

Item 4: Interaction With Density Comonads

Omitted.

Item 5: Interaction With Codensity Monads

Omitted.



### 02HX 12.1.5 The Yoneda Lemma

Let  $\mathcal{F}: \mathcal{C}^{\text{op}} \rightarrow \text{Sets}$  be a presheaf on  $\mathcal{C}$ .

02HY

#### THEOREM 12.1.5.1.1 ► THE YONEDA LEMMA

We have a bijection

$$\text{Nat}(h_A, \mathcal{F}) \cong \mathcal{F}(A),$$

natural in  $A \in \text{Obj}(\mathcal{C})$ , determining a natural isomorphism of functors

$$\text{Nat}(h_{(-)}, \mathcal{F}) \cong \mathcal{F}.$$

#### PROOF 12.1.5.1.2 ► PROOF OF THEOREM 12.1.5.1.1

The Transformation  $\text{ev}: \text{Nat}(h_{(-)}, \mathcal{F}) \Rightarrow \mathcal{F}$

Let

$$\text{ev}: \text{Nat}(h_{(-)}, \mathcal{F}) \Rightarrow \mathcal{F}$$

be the transformation consisting of the collection

$$\{\text{ev}_A: \text{Nat}(h_A, \mathcal{F}) \rightarrow \mathcal{F}(A)\}_{A \in \text{Obj}(\mathcal{C})}$$

with

$$\text{ev}_A(\alpha) = \alpha_A(\text{id}_A)$$

for each  $\alpha \in \text{Nat}(h_A, \mathcal{F})$ , where  $\alpha_A$  is the component

$$\alpha_A: \text{Hom}_{\mathcal{C}}(A, A) \rightarrow \mathcal{F}(A)$$

of  $\alpha$  at  $A$ .

The Transformation  $\xi: \mathcal{F} \Rightarrow \text{Nat}(h_{(-)}, \mathcal{F})$

Let

$$\xi: \mathcal{F} \Rightarrow \text{Nat}(h_{(-)}, \mathcal{F})$$

be the transformation consisting of the collection

$$\{\xi_A: \mathcal{F}(A) \rightarrow \text{Nat}(h_A, \mathcal{F})\}_{A \in \text{Obj}(C)},$$

where  $\xi_A$  is the map sending an element  $\phi \in \mathcal{F}(A)$  to the transformation

$$\xi_A(\phi): h_A \Rightarrow \mathcal{F}$$

(which we will show is natural in a bit) consisting of the collection

$$\{\xi_A(\phi)_X: h_A(X) \rightarrow \mathcal{F}(X)\}_{X \in \text{Obj}(C)},$$

with

$$\xi_A(\phi)_X(f) \stackrel{\text{def}}{=} [\mathcal{F}(f)](\phi)$$

for each  $f \in h_A(X)$ , where

$$\mathcal{F}(f): \mathcal{F}(A) \rightarrow \mathcal{F}(X)$$

is the image of  $f$  by  $\mathcal{F}$ .

Naturality of  $\xi_A(\phi): h_A \Rightarrow \mathcal{F}$

The transformation

$$\xi_A(\phi): h_A \Rightarrow \mathcal{F}$$

is indeed natural, as the diagram

$$\begin{array}{ccc} h_A^Y & \xrightarrow{f^*} & h_A^X \\ \xi_A(\phi)_Y \downarrow & & \downarrow \xi_A(\phi)_X \\ \mathcal{F}(Y) & \xrightarrow[\mathcal{F}(f)]{} & \mathcal{F}(X) \end{array}$$

commutes for each morphism  $f: X \rightarrow Y$  of  $C$ , acting on elements as

$$\begin{array}{ccc} h & & h \mapsto h \circ f \\ \downarrow & & \downarrow \\ [\mathcal{F}(h)](\phi) & \mapsto & [\mathcal{F}(f)][[\mathcal{F}(h)](\phi)] & & [\mathcal{F}(h \circ f)](\phi), \end{array}$$

where we have

$$[\mathcal{F}(f)][[\mathcal{F}(h)](\phi)] = [\mathcal{F}(h \circ f)](\phi)$$

by the functoriality of  $\mathcal{F}$ .

Naturality of  $\text{ev}$ :  $\text{Nat}(h_{(-)}, \mathcal{F}) \Rightarrow \mathcal{F}$

Let  $f: X \rightarrow Y$  be a morphism of  $C$ . We claim the naturality diagram

$$\begin{array}{ccc} \text{Nat}(h_Y, \mathcal{F}) & \xrightarrow{(h_f)^*} & \text{Nat}(h_X, \mathcal{F}) \\ \text{ev}_Y \downarrow & & \downarrow \text{ev}_X \\ \mathcal{F}(Y) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(X) \end{array}$$

for  $\text{ev}$  at  $f$ , acting on elements as

$$\begin{array}{ccc} \alpha & & \alpha \longmapsto \alpha \circ h_f \\ \downarrow & & \downarrow \\ \alpha_Y(\text{id}_Y) \longmapsto [\mathcal{F}(f)](\alpha_Y(\text{id}_Y)) & & [\alpha \circ h_f]_X(\text{id}_X), \end{array}$$

commutes. Indeed:

- We have

$$\begin{aligned} [\alpha \circ h_f]_X(\text{id}_X) &\stackrel{\text{def}}{=} [\alpha_X \circ h_{f|X}](\text{id}_X) \\ &\stackrel{\text{def}}{=} [\alpha_X \circ f_*](\text{id}_X) \\ &\stackrel{\text{def}}{=} \alpha_X(f_*(\text{id}_X)) \\ &\stackrel{\text{def}}{=} \alpha_X(f). \end{aligned}$$

- Applying the naturality diagram

$$\begin{array}{ccc} h_Y^Y & \xrightarrow{f^*} & h_X^X \\ \alpha_Y \downarrow & & \downarrow \alpha_X \\ \mathcal{F}(Y) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(X) \end{array}$$

of  $\alpha: h_Y \Rightarrow \mathcal{F}$  at  $f: X \rightarrow Y$  to the element  $\text{id}_Y$  of  $h_Y^Y$ , we have

$$\begin{array}{ccc} \text{id}_Y & & \text{id}_Y \longmapsto f \\ \downarrow & & \downarrow \\ \alpha_Y(\text{id}_Y) \longmapsto [\mathcal{F}(f)](\alpha_Y(\text{id}_Y)) & & \alpha_X(f), \end{array}$$

showing that we have

$$[\mathcal{F}(f)](\alpha_Y(\text{id}_Y)) = \alpha_X(f).$$

Thus the naturality diagram for  $\text{ev}$  at  $f$  commutes, and  $\text{ev}$  is natural.

Naturality of  $\xi: \mathcal{F} \Rightarrow \text{Nat}(h_{(-)}, \mathcal{F})$

Let  $f: X \rightarrow Y$  be a morphism of  $C$ . We claim the naturality diagram

$$\begin{array}{ccc} \mathcal{F}(Y) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(X) \\ \xi_Y \downarrow & & \downarrow \xi_X \\ \text{Nat}(h_Y, \mathcal{F}) & \xrightarrow{(h_f)^*} & \text{Nat}(h_X, \mathcal{F}) \end{array}$$

for  $\xi$  at  $f$ , acting on elements as

$$\begin{array}{ccc} \phi & & \phi \longmapsto [\mathcal{F}(f)](\phi) \\ \downarrow & & \downarrow \\ \xi_Y(\phi) \longmapsto \xi_Y(\phi) \circ h_f & & \xi_X([\mathcal{F}(f)](\phi)) \end{array}$$

commutes. Indeed, for each  $X \in \text{Obj}(C)$  and each  $g \in h_X^A$ , we have

$$\begin{aligned} [\xi_Y(\phi) \circ h_f]_X(g) &\stackrel{\text{def}}{=} [\xi_Y(\phi)_X \circ h_{f|X}](g) \\ &\stackrel{\text{def}}{=} [\xi_Y(\phi)_X \circ f_*](g) \\ &\stackrel{\text{def}}{=} \xi_Y(\phi)_X(f_*(g)) \\ &\stackrel{\text{def}}{=} \xi_Y(\phi)_X(f \circ g) \\ &\stackrel{\text{def}}{=} [\mathcal{F}(f \circ g)](\phi) \end{aligned}$$

and

$$\begin{aligned} [\xi_X([\mathcal{F}(f)](\phi))]_X(g) &\stackrel{\text{def}}{=} \mathcal{F}(g)([\mathcal{F}(f)](\phi)) \\ &= [\mathcal{F}(f \circ g)](\phi), \end{aligned}$$

where we have used the functoriality of  $\mathcal{F}$ . Thus  $\xi_Y(\phi) \circ h_f$  and  $\xi_X([\mathcal{F}(f)](\phi))$  are equal, and the naturality diagram for  $\xi$  at  $f$  above commutes, showing  $\xi$  to be natural.

Invertibility I:  $\text{ev} \circ \xi = \text{id}_{\mathcal{F}}$

We claim that  $\text{ev} \circ \xi = \text{id}_{\mathcal{F}}$ , i.e. that we have

$$(\text{ev} \circ \xi)_A = \text{id}_{\mathcal{F}(A)}$$

for each  $A \in \text{Obj}(C)$ . Indeed, we have

$$\begin{aligned} [\text{ev} \circ \xi]_A(\phi) &\stackrel{\text{def}}{=} [\text{ev}_A \circ \xi_A](\phi) \\ &\stackrel{\text{def}}{=} \text{ev}_A(\xi_A(\phi)) \\ &\stackrel{\text{def}}{=} \xi_A(\phi)_A(\text{id}_A) \\ &\stackrel{\text{def}}{=} [\mathcal{F}(\text{id}_A)](\phi) \\ &= [\text{id}_{\mathcal{F}(A)}](\phi) \end{aligned}$$

for each  $\phi \in \mathcal{F}(A)$ .

Invertibility II:  $\xi \circ \text{ev} = \text{id}_{\text{Nat}(h_{(-)}, \mathcal{F})}$

We claim that  $\xi \circ \text{ev} = \text{id}_{\text{Nat}(h_{(-)}, \mathcal{F})}$ , i.e. that we have

$$(\xi \circ \text{ev})_A = \text{id}_{\text{Nat}(h_A, \mathcal{F})}$$

for each  $A \in \text{Obj}(C)$ . Indeed:

- We have

$$\begin{aligned} [\xi \circ \text{ev}]_A(\alpha) &\stackrel{\text{def}}{=} [\xi_A \circ \text{ev}_A](\alpha) \\ &\stackrel{\text{def}}{=} \xi_A(\text{ev}_A(\alpha)) \\ &\stackrel{\text{def}}{=} \xi_A(\alpha_A(\text{id}_A)) \end{aligned}$$

for each  $\alpha \in \text{Nat}(h_A, \mathcal{F})$ .

- For each  $X \in \text{Obj}(C)$ , we have

$$\xi_A(\alpha_A(\text{id}_A))_X = \alpha_X,$$

since we have

$$\begin{aligned} \xi_A(\alpha_A(\text{id}_A))_X(f) &\stackrel{\text{def}}{=} [\mathcal{F}(f)](\alpha_A(\text{id}_A)) \\ &\stackrel{(\dagger)}{=} \alpha_X(f) \end{aligned}$$

for each  $f \in h_A(X)$ , where the equality marked with  $(\dagger)$  follows from the commutativity of the naturality diagram

$$\begin{array}{ccc} h_A^A & \xrightarrow{f_*} & h_X^A \\ \alpha_A \downarrow & & \downarrow \alpha_X \\ \mathcal{F}(A) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(X) \end{array}$$

of  $\alpha$  at  $f: A \rightarrow X$ , which acts on  $\text{id}_A$  as

$$\begin{array}{ccc} \text{id}_A & \xrightarrow{\quad} & f \\ \downarrow & & \downarrow \\ \alpha_A(\text{id}_A) & \longmapsto & [\mathcal{F}(f)](\alpha_A(\text{id}_A)) = \alpha_X(f). \end{array}$$

This finishes the proof. 

### 02HZ 12.1.6 Properties of Categories of Presheaves

02J0

#### PROPOSITION 12.1.6.1.1 ► PROPERTIES OF CATEGORIES OF PRESHEAVES

Let  $C$  be a category.

02J1

1. *Functoriality.* The assignment  $C \mapsto \text{PSh}(C)$  defines a functor

$$\text{PSh}: \text{Cats} \rightarrow \text{Cats}$$

up to some set-theoretic considerations.<sup>1</sup>

02J2

2. *Interaction With Slice Categories.* Let  $X \in \text{Obj}(C)$ . We have an equivalence of categories

$$\text{PSh}(C/X) \xrightarrow{\text{eq.}} \text{PSh}(C)_{/h_X}.$$

02J3

3. *Interaction With Categories of Elements.* Let  $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$ . We have an equivalence of categories

$$\text{PSh}(\int_C \mathcal{F}) \xrightarrow{\text{eq.}} \text{PSh}(C)_{/\mathcal{F}}.$$

<sup>1</sup>For instance:

- The Cats in the source of PSh could be small categories, and then the Cats in the right would be locally small categories.
- The Cats in the source of PSh could be locally small categories, and then the Cats on the right would be large categories.

In general, one can systematise and formalise this using Grothendieck universes.

#### PROOF 12.1.6.1.2 ► PROOF OF PROPOSITION 12.1.6.1.1

Item 1: Functoriality

Omitted.

Item 2: Interaction With Slice Categories

Omitted.

Item 3: Interaction With Categories of Elements

Omitted.



## 02J4 12.2 Copresheaves

### 02J5 12.2.1 Foundations

Let  $C$  be a category.

02J6

#### DEFINITION 12.2.1.1.1 ► COPRESHEAVES ON A CATEGORY

A **copresheaf** on  $C$  is a functor  $F: C \rightarrow \text{Sets}$ .

02J7

#### EXAMPLE 12.2.1.1.2 ► COPRESHEAVES ON ONE-OBJECT CATEGORIES

Copresheaves on the delooping  $BA$  of a monoid  $A$  are precisely the right  $A$ -sets; see ??.

02J8

#### DEFINITION 12.2.1.1.3 ► MORPHISMS OF COPRESHEAVES

A **morphism of copresheaves** on  $C$  from  $F$  to  $G$  is a natural transformation  $\alpha: F \Rightarrow G$ .

02J9

#### DEFINITION 12.2.1.1.4 ► THE CATEGORY OF COPRESHEAVES ON A CATEGORY

The **category of copresheaves** on  $C$  is the category  $\text{CoPSh}(C)$  defined by

$$\text{CoPSh}(C) \stackrel{\text{def}}{=} \text{Fun}(C, \text{Sets}).$$

02JA

#### REMARK 12.2.1.1.5 ► UNWINDING DEFINITION 12.2.1.1.4

In detail, the **category of copresheaves** on  $C$  is the category  $\text{CoPSh}(C)$  where

- *Objects.* The objects of  $\text{CoPSh}(C)$  are copresheaves on  $C$  as in [Definition 12.2.1.1.1](#).
- *Morphisms.* The morphisms of  $\text{CoPSh}(C)$  are morphisms of copresheaves as in [Definition 12.2.1.1.3](#), i.e. we have

$$\text{Hom}_{\text{CoPSh}(C)}(F, G) \stackrel{\text{def}}{=} \text{Nat}(F, G)$$

for each  $F, G \in \text{Obj}(\text{CoPSh}(C))$ .

- *Identities.* For each  $F \in \text{Obj}(\text{CoPSh}(C))$ , the unit map

$$\mathbb{1}_F^{\text{CoPSh}(C)} : \text{pt} \rightarrow \text{Nat}(F, F)$$

of  $\text{CoPSh}(C)$  at  $F$  is defined by

$$\text{id}_F^{\text{CoPSh}(C)} \stackrel{\text{def}}{=} \text{id}_F,$$

where  $\text{id}_F : F \Rightarrow F$  is the identity natural transformation of [Example 11.9.3.1.1](#).

- *Composition.* For each  $F, G, H \in \text{Obj}(\text{CoPSh}(C))$ , the composition map

$$\circ_{F,G,H}^{\text{CoPSh}(C)} : \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H)$$

of  $\text{CoPSh}(C)$  at  $(F, G, H)$  is defined by

$$\beta \circ_{F,G,H}^{\text{CoPSh}(C)} \alpha \stackrel{\text{def}}{=} \beta \circ \alpha,$$

where  $\beta \circ \alpha : F \Rightarrow H$  is the vertical composition of  $\alpha$  and  $\beta$  of [Definition 11.9.4.1.1](#).

### 02JB 12.2.2 Corepresentable Copresheaves

Let  $C$  be a category.

02JC

#### DEFINITION 12.2.2.1.1 ► COREPRESENTABLE COPRESHEAVES

Let  $A \in \text{Obj}(C)$ .

02JD

1. The **corepresentable copresheaf associated to  $A$**  is the copresheaf

$$h^A : C \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each  $X \in \text{Obj}(C)$ , we have

$$h^A(X) \stackrel{\text{def}}{=} \text{Hom}_C(A, X).$$

- *Action on Morphisms.* For each  $X, Y \in \text{Obj}(C)$ , the action on morphisms

$$h_{X,Y}^A : \text{Hom}_C(X, Y) \rightarrow \text{Hom}_{\text{Sets}}(h^A(X), h^A(Y))$$

of  $h^A$  at  $(X, Y)$  is given by sending a morphism

$$f: X \rightarrow Y$$

of  $C$  to the map of sets

$$h^A(f): \underbrace{h^A(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(A, X)} \rightarrow \underbrace{h^A(Y)}_{\stackrel{\text{def}}{=} \text{Hom}_C(A, Y)}$$

defined by

$$h^A(f) \stackrel{\text{def}}{=} f_*,$$

where  $f_*$  is the postcomposition by  $f$  morphism of **Item 2** of [Definition 11.1.4.1.1](#).

- 02JE** 2. A **corepresenting object** for a copresheaf  $F: C \rightarrow \text{Sets}$  on  $C$  is an object  $A$  of  $C$  such that we have  $F \cong h^A$ .
- 02JF** 3. A copresheaf  $F: C^{\text{op}} \rightarrow \text{Sets}$  on  $C$  is **corepresentable** if  $F$  admits a corepresenting object.

#### 02JG EXAMPLE 12.2.2.1.2 ► COREPRESENTABLE COPRESHEAVES ON ONE-OBJECT CATEGORIES

The corepresentable copresheaf on the delooping  $\text{BA}$  of a monoid  $A$  associated to the unique object  $\bullet$  of  $\text{BA}$  is the right regular representation of  $A$  of  $\bullet$ .

#### 02JH PROPOSITION 12.2.2.1.3 ► UNIQUENESS OF COREPRESENTING OBJECTS UP TO ISOMORPHISM

Let  $F: C \rightarrow \text{Sets}$  be a copresheaf. If there exist  $A, B \in \text{Obj}(C)$  such that we have natural isomorphisms

$$\begin{aligned} h^A &\cong F, \\ h^B &\cong F, \end{aligned}$$

then  $A \cong B$ .

#### PROOF 12.2.2.1.4 ► PROOF OF PROPOSITION 12.2.2.1.3

By composing the isomorphisms  $h^A \cong F \cong h^B$ , we get a natural isomorphism  $h^A \cong h^B$ . By **Item 2** of [Proposition 12.2.4.1.2](#), we have  $A \cong B$ .

**02JJ 12.2.3 Corepresentable Natural Transformations**

Let  $C$  be a category, let  $A, B \in \text{Obj}(C)$ , and let  $f: A \rightarrow B$  be a morphism of  $C$ .

**02JK**

**DEFINITION 12.2.3.1.1 ► COREPRESENTABLE NATURAL TRANSFORMATIONS**

The **corepresentable natural transformation associated to  $f$**  is the natural transformation

$$h^f: h^B \Rightarrow h^A$$

consisting of the collection

$$\left\{ h_X^f: \underbrace{h^B(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(B, X)} \rightarrow \underbrace{h^A(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(A, X)} \right\}_{X \in \text{Obj}(C)}$$

with

$$h_X^f \stackrel{\text{def}}{=} f^*,$$

where  $f_*$  is the precomposition by  $f$  morphism of **Item 1** of [Definition 11.1.4.1.1](#).

**02JL 12.2.4 The Contravariant Yoneda Embedding**

02JM

## DEFINITION 12.2.4.1.1 ► THE CONTRAVARIANT YONEDA EMBEDDING

The **contravariant Yoneda embedding** of  $C$  is the functor<sup>1</sup>

$$\mathfrak{P}_C: C^{\text{op}} \hookrightarrow \text{CoPSh}(C)$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(C)$ , we have

$$\mathfrak{P}_C(A) \stackrel{\text{def}}{=} h^A.$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(C)$ , the action on morphisms

$$\mathfrak{P}_{C|A,B}: \text{Hom}_C(A, B) \rightarrow \text{Nat}(h^B, h^A)$$

of  $\mathfrak{P}_C$  at  $(A, B)$  is given by

$$\mathfrak{P}_{C|A,B}(f) \stackrel{\text{def}}{=} h^f$$

for each  $f \in \text{Hom}_C(A, B)$ , where  $h^f$  is the corepresentable natural transformation associated to  $f$  of [Definition 12.2.3.1.1](#).

<sup>1</sup>Further Notation: Also written  $h^{(-)}$ , or simply  $\mathfrak{P}$ .

02JN

## PROPOSITION 12.2.4.1.2 ► PROPERTIES OF THE CONTRAVARIANT YONEDA EMBEDDING

Let  $C$  be a category.

02JP

1. *Fully Faithfulness.* The contravariant Yoneda embedding

$$\mathfrak{P}_C: C^{\text{op}} \hookrightarrow \text{CoPSh}(C)$$

is fully faithful.

02JQ

2. *Preservation and Reflection of Isomorphisms.* The contravariant Yoneda embedding

$$\mathfrak{P}_C: C^{\text{op}} \hookrightarrow \text{CoPSh}(C)$$

preserves and reflects isomorphisms, i.e. given  $A, B \in \text{Obj}(C)$ , the following conditions are equivalent:

02JR

(a) We have  $A \cong B$ .

02JS

(b) We have  $h^A \cong h^B$ .

**PROOF 12.2.4.1.3 ▶ PROOF OF PROPOSITION 12.2.4.1.2**

**Item 1: Fully Faithfulness**

The proof is dual to that of **Item 1** of [Proposition 12.1.4.1.3](#), and is therefore omitted.

**Item 2: Preservation and Reflection of Isomorphisms**

This follows from **Item 1** of [Proposition 11.5.1.1.8](#) and **Item 3** of [Proposition 11.6.3.1.2](#). 

**02JT 12.2.5 The Contravariant Yoneda Lemma**

Let  $F: C \rightarrow \text{Sets}$  be a copresheaf on  $C$ .

**02JU****THEOREM 12.2.5.1.1 ▶ THE CONTRAVARIANT YONEDA LEMMA**

We have a bijection

$$\text{Nat}(h^A, F) \cong F(A),$$

natural in  $A \in \text{Obj}(C)$ , determining a natural isomorphism of functors

$$\text{Nat}(h^{(-)}, F) \cong F.$$

**PROOF 12.2.5.1.2 ▶ PROOF OF THEOREM 12.2.5.1.1**

The proof is dual to that of [Theorem 12.1.5.1.1](#), and is therefore omitted. 

**02JV 12.3 Restricted Yoneda Embeddings and Yoneda Extensions****02JW 12.3.1 Foundations**

let  $F: C \rightarrow \mathcal{D}$  be a functor.

**02JX****DEFINITION 12.3.1.1.1 ▶ THE RESTRICTED YONEDA EMBEDDING ASSOCIATED TO A FUNCTOR**

The **restricted Yoneda embedding associated to  $F$**  is the functor

$$\mathfrak{J}_F: \mathcal{D} \hookrightarrow \text{PSh}(C)$$

defined as the composition

$$\mathcal{D} \xrightarrow{\mathfrak{J}_{\mathcal{D}}} \text{PSh}(\mathcal{D}) \xrightarrow{F^{\text{op},*}} \text{PSh}(C).$$

02JY

In detail, the **restricted Yoneda embedding associated to  $F$**  is the functor

$$\mathfrak{y}_F: \mathcal{D} \hookrightarrow \mathbf{PSh}(C)$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(\mathcal{D})$ , we have

$$\begin{aligned}\mathfrak{y}_F(A) &\stackrel{\text{def}}{=} h_A \circ F^{\text{op}} \\ &\stackrel{\text{def}}{=} h_A^{F(-)}.\end{aligned}$$

- *Action on Morphisms.* For each  $A, B \in \text{Obj}(\mathcal{D})$ , the action on morphisms

$$\mathfrak{y}_{F|A,B}: \text{Hom}_{\mathcal{D}}(A, B) \rightarrow \text{Nat}(h_A^{F(-)}, h_B^{F(-)})$$

of  $\mathfrak{y}_F$  at  $(A, B)$  is given by

$$\begin{aligned}\mathfrak{y}_{F|A,B}(f) &\stackrel{\text{def}}{=} h_f^{F(-)} \\ &\stackrel{\text{def}}{=} h_f \star \text{id}_{F^{\text{op}}}\end{aligned}$$

for each  $f \in \text{Hom}_{\mathcal{D}}(A, B)$ , where  $h_f$  is the representable natural transformation associated to  $f$  of [Definition 12.1.3.1.1](#).

02JZ

#### EXAMPLE 12.3.1.3 ► EXAMPLES OF RESTRICTED YONEDA EMBEDDINGS

Here are some examples of restricted Yoneda embeddings.

02K0

1. *The Nerve Functor.* Let

$$\iota: \Delta \hookrightarrow \mathbf{Cats}$$

be the functor given by  $[n] \rightarrow \mathbb{n}$ . Then the restricted Yoneda embedding

$$\mathfrak{y}_\iota: \mathbf{Cats} \rightarrow \overbrace{\mathbf{PSh}(\Delta)}^{\stackrel{\text{def}}{=} \mathbf{sSets}}$$

of  $\iota$  is given by the nerve functor  $N_\bullet$  of [??](#).

02K1

2. *The Singular Simplicial Set Associated to a Topological Space.* Let

$$\iota: \Delta \hookrightarrow \mathbb{T}$$

be the functor given by  $[n] \rightarrow |\Delta^n|$ . Then the restricted Yoneda embedding

$$\mathfrak{y}_\iota: \mathbb{T} \rightarrow \overbrace{\mathbf{PSh}(\Delta)}^{\stackrel{\text{def}}{=} \mathbf{sSets}}$$

of  $\iota$  is given by the singular simplicial set functor  $\text{Sing}_\bullet$  of ??.

02K2

3. *The Coherent Nerve Functor.* Let

$$\iota: \Delta \hookrightarrow \text{sCats}$$

be the functor given by  $[n] \rightarrow \text{Path}(\Delta^n)$ , where  $\text{Path}(\Delta^n)$  is the simplicial category of ???. Then the restricted Yoneda embedding

$$\underline{\jmath}_\iota: \text{sCats} \xrightarrow{\sim} \underset{\text{def}}{\overbrace{\text{PSh}(\Delta)}} \\ \text{def} \\ \text{sSets}$$

of  $\iota$  is given by the coherent nerve functor  $N_\bullet^{\text{hc}}$  of ??.

02K3

4. *Kan's Ex Functor.* Let

$$\text{sd}: \Delta \hookrightarrow \text{sSets}$$

be the functor given by  $[n] \rightarrow \text{Sd}(\Delta^n)$ , where  $\text{Sd}(\Delta^n)$  is the barycentric subdivision of  $\Delta^n$  of ???. Then the restricted Yoneda embedding

$$\underline{\jmath}_{\text{sd}}: \text{sSets} \xrightarrow{\sim} \underset{\text{def}}{\overbrace{\text{PSh}(\Delta)}} \\ \text{def} \\ \text{sSets}$$

of  $\text{sd}$  is given by Kan's Ex functor of ??.

02K4

#### PROPOSITION 12.3.1.1.4 ► PROPERTIES OF THE RESTRICTED YONEDA EMBEDDING

let  $F: C \rightarrow \mathcal{D}$  be a functor.

02K5

1. *Interaction With Fully Faithfulness.* The following conditions are equivalent:

02K6

(a) The restricted Yoneda embedding  $\underline{\jmath}_F$  is fully faithful.

02K7

(b) The functor  $F$  is dense (??).

02K8

2. *As a Left Kan Extension.* We have a natural isomorphism of functors

$$\underline{\jmath}_F \cong \text{Lan}_F(\underline{\jmath}),$$

**PROOF 12.3.1.1.5 ▶ PROOF OF PROPOSITION 12.3.1.1.4**

**Item 1: Interaction With Fully Faithfulness**

Omitted.

**Item 2: As a Left Kan Extension**

Omitted. 

**02K9 12.3.2 The Yoneda Extension Functor**

Let  $F: C \rightarrow \mathcal{D}$  be a functor with  $C$  small and  $\mathcal{D}$  cocomplete.

**02KA****DEFINITION 12.3.2.1.1 ▶ THE YONEDA EXTENSION FUNCTOR**

The **Yoneda extension functor associated to  $F$**  is the left Kan extension

$$\text{Lan}_{\mathbb{1}}(F): \text{PSh}(C) \rightarrow \mathcal{D}, \quad \begin{array}{ccc} & \text{PSh}(C) & \\ & \downarrow \text{Lan}_{\mathbb{1}}(F) & \\ C & \xrightarrow{F} & \mathcal{D}. \end{array}$$

$\text{Lan}_{\mathbb{1}}(F)$

$\text{PSh}(C)$

$\downarrow \text{Lan}_{\mathbb{1}}(F)$

$C \xrightarrow{F} \mathcal{D}.$

**02KB****EXAMPLE 12.3.2.1.2 ▶ EXAMPLES OF YONEDA EXTENSIONS**

Here are some examples of Yoneda extensions.

**02KC**

1. *The Homotopy Category Functor.* Let

$$\iota: \Delta \hookrightarrow \text{Cats}$$

be the functor given by  $[n] \rightarrow \text{Cat}$ . Then the Yoneda extension

$$\text{Lan}_{\mathbb{1}}(\iota): \overbrace{\text{PSh}(\Delta)}^{\text{def}=\text{sSets}} \rightarrow \text{Cats}$$

of  $\iota$  is given by the homotopy category functor  $\text{Ho}$  of ??.

**02KD**

2. *The Geometric Realisation Functor.* Let

$$\iota: \Delta \hookrightarrow \mathbb{T}$$

be the functor given by  $[n] \rightarrow |\Delta^n|$ . Then the Yoneda extension

$$\text{Lan}_{\mathbb{1}}(\iota): \overbrace{\text{PSh}(\Delta)}^{\text{def}=\text{sSets}} \rightarrow \mathbb{T}$$

of  $\iota$  is given by the geometric realisation functor  $|-|$  of ??.

02KE

3. *The Path Simplicial Category Functor.* Let

$$\iota: \Delta \hookrightarrow \text{sCats}$$

be the functor given by  $[n] \rightarrow \text{Path}(\Delta^n)$ , where  $\text{Path}(\Delta^n)$  is the simplicial category of  $\Delta^n$ . Then the Yoneda extension

$$\text{Lan}_{\mathcal{J}}(\iota): \underbrace{\text{PSh}(\Delta)}_{\substack{\text{def} \\ \equiv \text{sSets}}} \rightarrow \text{sCats}$$

of  $\iota$  is given by the path simplicial category functor  $\text{Path}$ .

02KF

4. *The Barycentric Subdivision Functor.* Let

$$\text{sd}: \Delta \hookrightarrow \text{sSets}$$

be the functor given by  $[n] \rightarrow \text{Sd}(\Delta^n)$ , where  $\text{Sd}(\Delta^n)$  is the barycentric subdivision of  $\Delta^n$ . Then the Yoneda extension

$$\text{Lan}_{\mathcal{J}}(\text{sd}): \underbrace{\text{PSh}(\Delta)}_{\substack{\text{def} \\ \equiv \text{sSets}}} \rightarrow \text{sSets}$$

of  $\text{sd}$  is given by the barycentric subdivision functor  $\text{Sd}$ .

02KG

**PROPOSITION 12.3.2.1.3 ► PROPERTIES OF YONEDA EXTENSIONS**

Let  $F: C \rightarrow \mathcal{D}$  be a functor with  $C$  small and  $\mathcal{D}$  cocomplete.

02KH

1. *Functoriality.* The assignment  $F \mapsto \text{Lan}_{\mathcal{J}}(F)$  defines a functor

$$\text{Lan}_{\mathcal{J}}: \text{Fun}(C, \mathcal{D}) \rightarrow \text{Fun}(\text{PSh}(C), \mathcal{D}).$$

02KJ

2. *Adjointness.* We have an adjunction<sup>1</sup>

$$(\text{Lan}_{\mathcal{J}}(F) \dashv \mathcal{J}_F): \text{PSh}(C) \begin{array}{c} \xrightarrow{\text{Lan}_{\mathcal{J}}(F)} \\ \perp \\ \xleftarrow{\mathcal{J}_F} \end{array} \mathcal{D},$$

witnessed by a bijection

$$\text{Hom}_{\mathcal{D}}([\text{Lan}_{\mathcal{J}}(F)](\mathcal{F}), D) \cong \text{Nat}(\mathcal{F}, \mathcal{J}_F(D)),$$

natural in  $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$  and  $D \in \text{Obj}(\mathcal{D})$ .

02KK

3. *Interaction With the Yoneda Embedding.* We have a natural isomorphism of functors

$$\text{Lan}_{\mathcal{Y}}(F) \circ \mathcal{Y}_C \cong F, \quad \begin{array}{ccc} & \mathcal{Y}_C & \nearrow \text{Lan}_{\mathcal{Y}}(F) \\ C & \xrightarrow{F} & \mathcal{D}. \end{array}$$

02KL

4. *As a Coend.* We have

$$\begin{aligned} [\text{Lan}_{\mathcal{Y}}(F)](\mathcal{F}) &\cong \int^{A \in C} \text{Nat}(h_A, \mathcal{F}) \odot F(A) \\ &\cong \int^{A \in C} \mathcal{F}(A) \odot F(A) \end{aligned}$$

for each  $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$ .

02KM

5. *Interaction With Tensors of Presheaves With Functors.* We have a natural isomorphism

$$\text{Lan}_{\mathcal{Y}}(F) \cong (-) \odot_C F,$$

natural in  $F \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$ .

02KN

6. *Interaction With Finite Limits.* Let  $F: C \rightarrow \text{Sets}$  be a functor. The following conditions are equivalent:

02KP

(a) The functor  $F$  preserves finite limits.

02KQ

(b) The functor  $\text{Lan}_{\mathcal{Y}}(F)$  preserves finite limits.

02KR

(c) The category of elements  $\int_C F$  of  $F$  is cofiltered.

<sup>1</sup>Applying Item 2 of Proposition 12.3.1.4, we see that this adjunction has the form  $\text{Lan}_{\mathcal{Y}}(F) \dashv \text{Lan}_F(\mathcal{Y})$ .

PROOF 12.3.2.1.4 ► PROOF OF PROPOSITION 12.3.2.1.3

Item 1: Functoriality

This follows from ?? of ??.

Item 2: Adjointness

Omitted.

Item 3: Interaction With the Yoneda Embedding

This follows from ?? of ??.

Item 4: As a Coend

This follows from ?? of ?? and [Theorem 12.1.5.1.1](#).

Item 5: Interaction With Tensors of Presheaves With Functors

This follows from [Item 4](#).

Item 6: Interaction With Finite Limits

See [[coend-calculus](#)].



## 02LR 12.4 Functor Tensor Products

### 02LS 12.4.1 The Tensor Product of Presheaves With Copresheaves

Let  $C$  be a category, let  $\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets}$  be a presheaf on  $C$ , and let  $G: C \rightarrow \text{Sets}$  be a copresheaf on  $C$ .

02LT

#### DEFINITION 12.4.1.1.1 ► THE TENSOR PRODUCT OF PRESHEAVES WITH COPRESHEAVES

The **tensor product** of  $\mathcal{F}$  with  $G$  is the set  $\mathcal{F} \boxtimes_C G^1$  defined by

$$\mathcal{F} \boxtimes_C G \stackrel{\text{def}}{=} \int^{A \in C} \mathcal{F}(A) \times G(A).$$

<sup>1</sup>Further Notation: Also written simply  $\mathcal{F} \boxtimes G$ .

02LU

#### REMARK 12.4.1.1.2 ► UNWINDING DEFINITION 12.4.1.1

In other words, the tensor product of  $\mathcal{F}$  with  $G$  is the set  $\mathcal{F} \boxtimes_C G$  defined as the coend of the functor

$$C^{\text{op}} \times C \xrightarrow{\mathcal{F} \times G} \text{Sets} \times \text{Sets} \xrightarrow{\times} \text{Sets},$$

which is equivalently the composition

$$\begin{array}{ccc} C & \xrightarrow{F} & \text{pt} \\ & \searrow \times \circ (\mathcal{F} \times G) & \downarrow \mathcal{F} \\ & & C \end{array}$$

in  $\text{Prof}$ .

02LV

**EXAMPLE 12.4.1.1.3 ► THE TENSOR PRODUCT OF PRESHEAVES WITH COPRESHEAVES ON ONE OBJECT CATEGORIES**

02LW

**PROPOSITION 12.4.1.1.4 ► PROPERTIES OF TENSOR PRODUCTS OF PRESHEAVES WITH Co-PRESHEAVES**

02LY

Let  $C$  be a category.

02LX

1. *Functionality.* The assignments  $\mathcal{F}, G, (\mathcal{F}, G) \mapsto \mathcal{F} \boxtimes_C G$  define functors

$$\begin{aligned}\mathcal{F} \boxtimes_C - : \quad \text{PSh}(C) &\rightarrow \text{Sets}, \\ - \boxtimes_C G : \quad \text{CoPSh}(C) &\rightarrow \text{Sets}, \\ -_1 \boxtimes_C -_2 : \quad \text{PSh}(C) \times \text{CoPSh}(C) &\rightarrow \text{Sets}.\end{aligned}$$

2. As a Composition of Profunctors. Let  $C$  be a category and let:

- $\mathcal{F} : \text{pt} \nrightarrow C$  be a presheaf on  $C$ , viewed as a profunctor.
- $F : C \nrightarrow \text{pt}$  be a copresheaf on  $C$ , viewed as a profunctor.

We have a natural isomorphism of profunctors

$$\begin{array}{ccc}\mathcal{F} \boxtimes_C F & \cong & F \diamond \mathcal{F}, \\ \text{pt} & \xrightarrow{\quad \perp \quad} & \text{pt},\end{array} \quad \begin{array}{c} C \\ \swarrow \mathcal{F} \quad \downarrow F \\ \text{pt} \end{array}$$

natural in  $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$  and  $F \in \text{Obj}(\text{CoPSh}(C))$ .

02LZ

3. *Interaction With Representable Presheaves.* Let  $\mathcal{F}$  be a presheaf on  $C$ . We have a bijection of sets

$$\mathcal{F} \boxtimes_C h^X \cong \mathcal{F}(X),$$

natural in  $X \in \text{Obj}(C)$ , giving a natural isomorphism of functors

$$\begin{array}{ccc}\mathcal{F} \boxtimes_C h^{(-)} & \cong & \mathcal{F}, \\ C^{\text{op}} & \xrightarrow{\quad \mathcal{F} \quad} & \text{Sets}.\end{array} \quad \begin{array}{c} \text{CoPSh}(C) \\ \swarrow \mathcal{F}_C \quad \downarrow \mathcal{F} \boxtimes_C - \\ \text{Sets.} \end{array}$$

02M0

4. *Interaction With Corepresentable Copresheaves.* Let  $G$  be a copresheaf on  $C$ . We have a bijection of sets

$$h_X \boxtimes_C G \cong G(X),$$

natural in  $X \in \text{Obj}(C)$ , giving a natural isomorphism of functors

$$\begin{array}{ccc} & \mathsf{PSh}(C) & \\ h_{(-)} \boxtimes_C G \cong G, & \begin{array}{c} \xrightarrow{\mathfrak{x}_C} \\ \Downarrow \end{array} & \downarrow \dashv \boxtimes_C G \\ C & \xrightarrow[G]{} & \mathsf{Sets}. \end{array}$$

02M1

5. *Interaction With Yoneda Extensions.* Let  $G: C \rightarrow \mathsf{Sets}$  be a copresheaf on  $C$ . We have a natural isomorphism

$$\begin{array}{ccc} & \mathsf{PSh}(C) & \\ \text{Lan}_{\mathfrak{x}}(G) \cong (-) \boxtimes_C G, & \begin{array}{c} \xrightarrow{\mathfrak{x}_C} \\ \Downarrow \end{array} & \downarrow \dashv (-) \boxtimes_C G \\ C & \xrightarrow[G]{} & \mathsf{Sets}, \end{array}$$

natural in  $G \in \text{Obj}(\mathsf{CoPSh}(C))$ .

02M2

6. *Interaction With Contravariant Yoneda Extensions.* Let  $\mathcal{F}: C^{\text{op}} \rightarrow \mathsf{Sets}$  be a presheaf on  $C$ . We have a natural isomorphism

$$\begin{array}{ccc} & \mathsf{CoPSh}(C) & \\ \text{Lan}_{\mathfrak{F}}(\mathcal{F}) \cong \mathcal{F} \boxtimes_C (-), & \begin{array}{c} \xrightarrow{\mathfrak{F}_C} \\ \Downarrow \end{array} & \downarrow \dashv \mathcal{F} \boxtimes_C (-) \\ C^{\text{op}} & \xrightarrow[\mathcal{F}]{} & \mathsf{Sets}, \end{array}$$

natural in  $\mathcal{F} \in \text{Obj}(\mathsf{PSh}(C))$ .

#### PROOF 12.4.1.1.5 ▶ PROOF OF PROPOSITION 12.4.1.1.4

Item 1: Functoriality

Omitted.

Item 2: As a Composition of Profunctors

Clear.

Item 3: Interaction With Representable Presheaves

This follows from ??.

Item 4: Interaction With Corepresentable Copresheaves

This follows from ??.

Item 5: Interaction With Yoneda Extensions

This is a special case of **Item 5 of Proposition 12.3.2.1.3**.

Item 6: Interaction With Contravariant Yoneda Extensions

This is a special case of ?? of ??.



### 02M3 12.4.2 The Tensor of a Presheaf With a Functor

Let  $C$  be a category, let  $\mathcal{D}$  be a category with coproducts, let  $\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets}$  be a presheaf on  $C$ , and let  $G: C \rightarrow \mathcal{D}$  be a functor.

02M4

#### DEFINITION 12.4.2.1.1 ► THE TENSOR OF A PRESHEAF WITH A FUNCTOR

The **tensor** of  $\mathcal{F}$  with  $G$  is the object  $\mathcal{F} \odot_C G$ <sup>1</sup> of  $\mathcal{D}$  defined by

$$\mathcal{F} \odot_C G \stackrel{\text{def}}{=} \int^{A \in C} \mathcal{F}(A) \odot G(A).$$

<sup>1</sup>Further Notation: Also written simply  $\mathcal{F} \odot G$ .

02M5

#### REMARK 12.4.2.1.2 ► UNWINDING DEFINITION 12.4.2.1.1

In other words, the tensor of  $\mathcal{F}$  with  $G$  is the object  $\mathcal{F} \odot_C G$  of  $\mathcal{D}$  defined as the coend of the functor

$$C^{\text{op}} \times C \xrightarrow{\mathcal{F} \times G} \text{Sets} \times \mathcal{D} \xrightarrow{\odot} \mathcal{D}.$$

02M6

#### PROPOSITION 12.4.2.1.3 ► PROPERTIES OF TENSORS OF PRESHEAVES WITH FUNCTORS

Let  $C$  be a category.

02M7

1. *Functionality.* The assignments  $\mathcal{F}, G, (\mathcal{F}, G) \mapsto \mathcal{F} \odot_C G$  define functors

$$\begin{aligned} \mathcal{F} \odot_C -: & \quad \text{PSh}(C) & \rightarrow \mathcal{D}, \\ - \odot_C G: & \quad \text{Fun}(C, \mathcal{D}) & \rightarrow \mathcal{D}, \\ -_1 \odot_C -_2: & \quad \text{PSh}(C) \times \text{Fun}(C, \mathcal{D}) & \rightarrow \mathcal{D}. \end{aligned}$$

02M8

2. *Interaction With Corepresentable Copresheaves.* We have an isomorphism

$$h_X \odot_C G \cong G(X),$$

natural in  $X \in \text{Obj}(C)$ , giving a natural isomorphism of functors

$$h_{(-)} \odot_C G \cong G.$$

02M9

3. *Interaction With Yoneda Extensions.* We have a natural isomorphism

$$\text{Lan}_{\mathcal{Y}}(G) \cong (-) \odot_C G,$$

natural in  $G \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$ .

#### PROOF 12.4.2.1.4 ► PROOF OF PROPOSITION 12.4.2.1.3

Item 1: Functoriality

Omitted.

???: Interaction With Corepresentable Copresheaves

This follows from ??.

Item 3: Interaction With Yoneda Extensions

This is a repetition of Item 5 of Proposition 12.3.2.1.3, and is proved there. 

### 02MA 12.4.3 The Tensor of a Copresheaf With a Functor

Let  $C$  be a category, let  $\mathcal{D}$  be a category with coproducts, let  $F: C \rightarrow \text{Sets}$  be a copresheaf on  $C$ , and let  $G: C^{\text{op}} \rightarrow \mathcal{D}$  be a functor.

02MB

#### DEFINITION 12.4.3.1.1 ► THE TENSOR OF A COPRESHEAF WITH A FUNCTOR

The **tensor** of  $F$  with  $G$  is the set  $F \odot_C G$ <sup>1</sup> defined by

$$F \odot_C G \stackrel{\text{def}}{=} \int^{A \in C} F(A) \odot G(A).$$

<sup>1</sup>Further Notation: Also written simply  $F \odot G$ .

02MC

#### REMARK 12.4.3.1.2 ► UNWINDING DEFINITION 12.4.3.1.1

In other words, the tensor of  $F$  with  $G$  is the object  $F \odot_C G$  of  $\mathcal{D}$  defined as the coend of the functor

$$C^{\text{op}} \times C \xrightarrow{\sim} C \times C^{\text{op}} \xrightarrow{F \times G} \text{Sets} \times \mathcal{D} \xrightarrow{\odot} \mathcal{D}.$$

02MD

#### PROPOSITION 12.4.3.1.3 ► PROPERTIES OF TENSORS OF COPRESHEAVES WITH FUNCTORS

Let  $C$  be a category.

02ME

1. *Functionality.* The assignments  $F, G, (F, G) \mapsto F \odot_C G$  define functors

$$\begin{aligned} F \odot_C - : \quad \text{CoPSh}(C) &\rightarrow \mathcal{D}, \\ - \odot_C \mathcal{G} : \quad \text{Fun}(C^{\text{op}}, \mathcal{D}) &\rightarrow \mathcal{D}, \\ -_1 \odot_C -_2 : \quad \text{Fun}(C^{\text{op}}, \mathcal{D}) \times \text{CoPSh}(C) &\rightarrow \mathcal{D}. \end{aligned}$$

02MF

2. *Interaction With Corepresentable Copresheaves.* We have an isomorphism

$$h^X \odot_C G \cong G(X),$$

natural in  $X \in \text{Obj}(C)$ , giving a natural isomorphism of functors

$$h^{(-)} \odot_C G \cong G.$$

02MG

3. *Interaction With Contravariant Yoneda Extensions.* We have a natural isomorphism

$$\text{Lan}_{\mathfrak{P}}(G) \cong G \odot_C (-),$$

natural in  $G \in \text{Obj}(\text{Fun}(C^{\text{op}}, \mathcal{D}))$ .

#### PROOF 12.4.3.1.4 ► PROOF OF PROPOSITION 12.4.3.1.3

Item 1: Functionality

Omitted.

??: Interaction With Representable Presheaves

This follows from ??.

??: Interaction With Corepresentable Copresheaves

This follows from ??.

??: Interaction With Yoneda Extensions

Omitted.

Item 3: Interaction With Contravariant Yoneda Extensions

Omitted.



## Appendices

## 12.A Other Chapters

<b>Preliminaries</b>	10. Conditions on Relations
1. Introduction	
2. A Guide to the Literature	
<b>Sets</b>	<b>Categories</b>
3. Sets	11. Categories
4. Constructions With Sets	
5. Monoidal Structures on the Category of Sets	12. Presheaves and the Yoneda Lemma
6. Pointed Sets	
7. Tensor Products of Pointed Sets	
<b>Relations</b>	<b>Monoidal Categories</b>
8. Relations	13. Constructions With Monoidal Categories
9. Constructions With Relations	
	<b>Bicategories</b>
	14. Types of Morphisms in Bicategories
	<b>Extra Part</b>
	15. Notes

**Part V**

**Monoidal Categories**

## Chapter 13

# Constructions With Monoidal Categories

**02H3** This chapter contains some material on constructions with monoidal categories.

### Contents

---

13.1 Moduli Categories of Monoidal Structures .....	869
13.1.1 The Moduli Category of Monoidal Structures on a Category .....	869
13.1.2 The Moduli Category of Braided Monoidal Struc- tures on a Category .....	882
13.1.3 The Moduli Category of Symmetric Monoidal Structures on a Category .....	882
13.2 Moduli Categories of Closed Monoidal Structures.....	882
13.3 Moduli Categories of Refinements of Monoidal Struc- tures .....	882
13.3.1 The Moduli Category of Braided Refinements of a Monoidal Structure .....	882
13.A Other Chapters .....	883

---

**01UG 13.1 Moduli Categories of Monoidal Structures**

**01UH 13.1.1 The Moduli Category of Monoidal Structures on a Category**

Let  $C$  be a category.

01UJ

**DEFINITION 13.1.1.1 ► THE MODULI CATEGORY OF MONOIDAL STRUCTURES ON A CATEGORY**

The **moduli category of monoidal structures on  $C$**  is the category  $\mathcal{M}_{\mathbb{E}_1}(C)$  defined by

$$\begin{array}{ccc} \mathcal{M}_{\mathbb{E}_1}(C) & \longrightarrow & \text{MonCats} \\ \downarrow & \lrcorner & \downarrow \text{忘} \\ \text{pt} & \xrightarrow{[C]} & \text{Cats.} \end{array}$$

01UK

**REMARK 13.1.1.2 ► UNWINDING DEFINITION 13.1.1.1, I**

In detail, the **moduli category of monoidal structures on  $C$**  is the category  $\mathcal{M}_{\mathbb{E}_1}(C)$  where:

- **Objects.** The objects of  $\mathcal{M}_{\mathbb{E}_1}(C)$  are monoidal categories  $(C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C)$  whose underlying category is  $C$ .
- **Morphisms.** A morphism from  $(C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C)$  to  $(C, \boxtimes_C, \mathbb{1}'_C, \alpha^{C,'}, \lambda^{C,'}, \rho^{C,'})$  is a strong monoidal functor structure

$$\begin{aligned} \text{id}_C^\otimes: A \boxtimes_C B &\xrightarrow{\sim} A \otimes_C B, \\ \text{id}_{\mathbb{1}|C}^\otimes: \mathbb{1}'_C &\xrightarrow{\sim} \mathbb{1}_C \end{aligned}$$

on the identity functor  $\text{id}_C: C \rightarrow C$  of  $C$ .

- **Identities.** For each  $M \stackrel{\text{def}}{=} (C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C) \in \text{Obj}(\mathcal{M}_{\mathbb{E}_1}(C))$ , the unit map

$$\mathbb{1}_{M,M}^{\mathcal{M}_{\mathbb{E}_1}(C)}: \text{pt} \rightarrow \text{Hom}_{\mathcal{M}_{\mathbb{E}_1}(C)}(M, M)$$

of  $\mathcal{M}_{\mathbb{E}_1}(C)$  at  $M$  is defined by

$$\text{id}_M^{\mathcal{M}_{\mathbb{E}_1}(C)} \stackrel{\text{def}}{=} (\text{id}_C^\otimes, \text{id}_{\mathbb{1}|C}^\otimes),$$

where  $(\text{id}_C^\otimes, \text{id}_{\mathbb{1}|C}^\otimes)$  is the identity monoidal functor of  $C$  of ??.

- **Composition.** For each  $M, N, P \in \text{Obj}(\mathcal{M}_{\mathbb{E}_1}(C))$ , the composition map

$$\circ_{M,N,P}^{\mathcal{M}_{\mathbb{E}_1}(C)}: \text{Hom}_{\mathcal{M}_{\mathbb{E}_1}(C)}(N, P) \times \text{Hom}_{\mathcal{M}_{\mathbb{E}_1}(C)}(M, N) \rightarrow \text{Hom}_{\mathcal{M}_{\mathbb{E}_1}(C)}(M, P)$$

of  $\mathcal{M}_{\mathbb{E}_1}(C)$  at  $(M, N, P)$  is defined by

$$(\text{id}_C^{\otimes,'}, \text{id}_{\mathbb{1}|C}^{\otimes,'}) \circ_{M,N,P}^{\mathcal{M}_{\mathbb{E}_1}(C)} (\text{id}_C^\otimes, \text{id}_{\mathbb{1}|C}^\otimes) \stackrel{\text{def}}{=} (\text{id}_C^{\otimes,'} \circ \text{id}_C^\otimes, \text{id}_{\mathbb{1}|C}^{\otimes,'} \circ \text{id}_{\mathbb{1}|C}^\otimes).$$

01UL

In particular, a morphism in  $\mathcal{M}_{\mathbb{E}_1}(C)$  from  $(C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C)$  to  $(C, \boxtimes_C, \mathbb{1}'_C, \alpha^{C,'}, \lambda^{C,'}, \rho^{C,'})$  satisfies the following conditions:

01UM

1. *Naturality.* For each pair  $f: A \rightarrow X$  and  $g: B \rightarrow Y$  of morphisms of  $C$ , the diagram

$$\begin{array}{ccc} A \boxtimes_C B & \xrightarrow{f \boxtimes_C g} & X \boxtimes_C Y \\ \downarrow \text{id}_{A,B}^\otimes & & \downarrow \text{id}_{X,Y}^\otimes \\ A \otimes_C B & \xrightarrow{f \otimes_C g} & X \otimes_C Y \end{array}$$

commutes.

01UN

2. *Monoidality.* For each  $A, B, C \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccc} & (A \boxtimes_C B) \boxtimes_C C & \\ \text{id}_{A,B}^\otimes \boxtimes_C \text{id}_C & \swarrow & \searrow \alpha_{A,B,C}^{C,'} \\ (A \otimes_C B) \boxtimes_C C & & A \boxtimes_C (B \otimes_C C) \\ \downarrow \text{id}_{A \otimes_C B, C}^\otimes & & \downarrow \text{id}_A \boxtimes_C \text{id}_{B,C}^\otimes \\ (A \otimes_C B) \otimes_C C & & A \boxtimes_C (B \otimes_C C) \\ \downarrow \alpha_{A,B,C}^C & \searrow & \swarrow \text{id}_{A,B \otimes_C C}^\otimes \\ A \otimes_C (B \otimes_C C) & & \end{array}$$

commutes.

01UP

3. *Left Monoidal Unity.* For each  $A \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccc} \mathbb{1}_C \boxtimes_C A & \xrightarrow{\text{id}_{\mathbb{1}_C,A}^\otimes} & \mathbb{1}_C \otimes_C A \\ \uparrow \text{id}_{\mathbb{1}}^\otimes \boxtimes_C \text{id}_A & & \downarrow \lambda_A^C \\ \mathbb{1}'_C \boxtimes_C A & \xrightarrow{\lambda_A^{C,'}} & A \end{array}$$

commutes.

01UQ

4. *Right Monoidal Unity.* For each  $A \in \text{Obj}(C)$ , the diagram

$$\begin{array}{ccc}
 A \boxtimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{A,\mathbb{1}_C}^\otimes} & A \otimes_C \mathbb{1}_C \\
 \text{id}_A \boxtimes_C \text{id}_{\mathbb{1}}^\otimes \nearrow & & \searrow \rho_A^C \\
 A \boxtimes_C \mathbb{1}'_C & \xrightarrow{\rho_{A,C}^{C,'}} & A
 \end{array}$$

commutes.

01UR

**PROPOSITION 13.1.1.4 ► PROPERTIES OF THE MODULI CATEGORY OF MONOIDAL STRUCTURES ON A CATEGORY**

Let  $C$  be a category.

01US

1. *Extra Monoidality Conditions.* Let  $(\text{id}_C^\otimes, \text{id}_{\mathbb{1}|C}^\otimes)$  be a morphism of  $\mathcal{M}_{\mathbb{E}_1}(C)$  from  $(C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C)$  to  $(C, \boxtimes_C, \mathbb{1}'_C, \alpha^{C,'}, \lambda^{C,'}, \rho^{C,'})$ .

01UT

(a) The diagram

$$\begin{array}{ccc}
 (A \boxtimes_C B) \boxtimes_C C & \xrightarrow{\text{id}_{A,B}^\otimes \boxtimes_C \text{id}_C} & (A \otimes_C B) \boxtimes_C C \\
 \text{id}_{A \boxtimes_C B,C}^\otimes \downarrow & & \downarrow \text{id}_{A \otimes_C B,C}^\otimes \\
 (A \boxtimes_C B) \otimes_C C & \xrightarrow{\text{id}_{A,B}^\otimes \otimes_C \text{id}_C} & (A \otimes_C B) \otimes_C C
 \end{array}$$

commutes.

01UU

(b) The diagram

$$\begin{array}{ccc}
 A \boxtimes_C (B \boxtimes_C C) & \xrightarrow{\text{id}_A \boxtimes_C \text{id}_{B,C}^\otimes} & A \boxtimes_C (B \otimes_C C) \\
 \text{id}_{A,B \boxtimes_C C}^\otimes \downarrow & & \downarrow \text{id}_{A,B \otimes_C C}^\otimes \\
 A \otimes_C (B \boxtimes_C C) & \xrightarrow{\text{id}_A \otimes_C \text{id}_{B,C}^\otimes} & A \otimes_C (B \otimes_C C)
 \end{array}$$

commutes.

01WB

2. *Extra Monoidal Unity Constraints.* Let  $(\text{id}_C^\otimes, \text{id}_{\mathbb{1}|C}^\otimes)$  be a morphism of  $\mathcal{M}_{\mathbb{E}_1}(C)$  from  $(C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C)$  to  $(C, \boxtimes_C, \mathbb{1}'_C, \alpha^{C,'}, \lambda^{C,'}, \rho^{C,'})$ .

01WC

(a) The diagram

$$\begin{array}{ccc} \mathbb{1}_C \otimes_C \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}_C \boxtimes_C \mathbb{1}'_C \\ \lambda_{\mathbb{1}'_C}^C \downarrow & & \downarrow \rho_{\mathbb{1}'_C}^C \\ \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}}^\otimes} & \mathbb{1}_C \end{array}$$

commutes.

01WD

(b) The diagram

$$\begin{array}{ccc} \mathbb{1}'_C \otimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}'_C \boxtimes_C \mathbb{1}_C \\ \rho_{\mathbb{1}'_C}^C \downarrow & & \downarrow \lambda_{\mathbb{1}'_C}^C \\ \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}}^\otimes} & \mathbb{1}_C \end{array}$$

commutes.

01WE

(c) The diagram

$$\begin{array}{ccc} \mathbb{1}'_C \boxtimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^\otimes} & \mathbb{1}'_C \otimes_C \mathbb{1}_C \\ \lambda_{\mathbb{1}_C}^{C,\prime} \downarrow & & \downarrow \rho_{\mathbb{1}'_C}^C \\ \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}}^{\otimes, -1}} & \mathbb{1}'_C \end{array}$$

commutes.

01WF

(d) The diagram

$$\begin{array}{ccc} \mathbb{1}_C \boxtimes_C \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^\otimes} & \mathbb{1}_C \otimes_C \mathbb{1}'_C \\ \rho_{\mathbb{1}_C}^{C,\prime} \downarrow & & \downarrow \lambda_{\mathbb{1}'_C}^C \\ \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}}^{\otimes, -1}} & \mathbb{1}'_C \end{array}$$

commutes.

01UV

3. *Mixed Associators.* Let  $(C, \otimes_C, \mathbb{1}_C, \alpha^C, \lambda^C, \rho^C)$  and  $(C, \boxtimes_C, \mathbb{1}'_C, \alpha^{C,\prime}, \lambda^{C,\prime}, \rho^{C,\prime})$  be monoidal structures on  $C$  and let

$$\text{id}_{-1, -2}^\otimes : -_1 \boxtimes_C -_2 \rightarrow -_1 \otimes_C -_2$$

be a natural transformation.

01UW

(a) If there exists a natural transformation

$$\alpha_{A,B,C}^{\otimes}: (A \otimes_C B) \boxtimes_C C \rightarrow A \otimes_C (B \boxtimes_C C)$$

making the diagrams

$$\begin{array}{ccc} (A \otimes_C B) \boxtimes_C C & \xrightarrow{\alpha_{A,B,C}^{\otimes}} & A \otimes_C (B \boxtimes_C C) \\ \downarrow \text{id}_{A \otimes_C B, C}^{\otimes} & & \downarrow \text{id}_A \otimes_C \text{id}_{B,C}^{\otimes} \\ (A \otimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^C} & A \otimes_C (B \otimes_C C) \end{array}$$

and

$$\begin{array}{ccc} (A \boxtimes_C B) \boxtimes_C C & \xrightarrow{\alpha_{A,B,C}^{C,*}} & A \boxtimes_C (B \boxtimes_C C) \\ \downarrow \text{id}_{A,B \boxtimes_C C}^{\otimes} & & \downarrow \text{id}_{A,B \boxtimes_C C} \\ (A \otimes_C B) \boxtimes_C C & \xrightarrow{\alpha_{A,B,C}^{\otimes}} & A \otimes_C (B \boxtimes_C C) \end{array}$$

commute, then the natural transformation  $\text{id}^{\otimes}$  satisfies the monoidality condition of Item 2 of Remark 13.1.1.3.

01UX

(b) If there exists a natural transformation

$$\alpha_{A,B,C}^{\boxtimes}: (A \boxtimes_C B) \otimes_C C \rightarrow A \boxtimes_C (B \otimes_C C)$$

making the diagrams

$$\begin{array}{ccc} (A \boxtimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^{\boxtimes}} & A \boxtimes_C (B \otimes_C C) \\ \downarrow \text{id}_{A,B \otimes_C C}^{\otimes} & & \downarrow \text{id}_{A,B \otimes_C C}^{\otimes} \\ (A \otimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^C} & A \otimes_C (B \otimes_C C) \end{array}$$

and

$$\begin{array}{ccc} (A \boxtimes_C B) \boxtimes_C C & \xrightarrow{\alpha_{A,B,C}^{C,*}} & A \boxtimes_C (B \boxtimes_C C) \\ \downarrow \text{id}_{A \boxtimes_C B, C}^{\otimes} & & \downarrow \text{id}_A \boxtimes_C \text{id}_{B,C}^{\otimes} \\ (A \boxtimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^{\boxtimes}} & A \boxtimes_C (B \otimes_C C) \end{array}$$

commute, then the natural transformation  $\text{id}^{\otimes}$  satisfies the monoidality condition of Item 2 of Remark 13.1.1.3.

01UY

(c) If there exists a natural transformation

$$\alpha_{A,B,C}^{\boxtimes,\otimes}: (A \boxtimes_C B) \otimes_C C \rightarrow A \otimes_C (B \boxtimes_C C)$$

making the diagrams

$$\begin{array}{ccc} (A \boxtimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^{\boxtimes,\otimes}} & A \otimes_C (B \boxtimes_C C) \\ \downarrow \text{id}_{A,B}^{\otimes} \otimes_C \text{id}_C & & \downarrow \text{id}_A \otimes_C \text{id}_{B,C}^{\otimes} \\ (A \otimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^C} & A \otimes_C (B \otimes_C C) \end{array}$$

and

$$\begin{array}{ccc} (A \boxtimes_C B) \boxtimes_C C & \xrightarrow{\alpha_{A,B,C}^{C,\prime}} & A \boxtimes_C (B \boxtimes_C C) \\ \downarrow \text{id}_{A \boxtimes_C B,C}^{\otimes} & & \downarrow \text{id}_{A,B \boxtimes_C C}^{\otimes} \\ (A \boxtimes_C B) \otimes_C C & \xrightarrow{\alpha_{A,B,C}^{\boxtimes,\otimes}} & A \otimes_C (B \boxtimes_C C) \end{array}$$

commute, then the natural transformation  $\text{id}^{\otimes}$  satisfies the monoidality condition of **Item 2** of **Remark 13.1.1.1.3**.**PROOF 13.1.1.5 ▶ PROOF OF PROPOSITION 13.1.1.4****Item 1: Extra Monoidality Conditions**We claim that **Items 1a** and **1b** are indeed true:

1. *Proof of Item 1a:* This follows from the naturality of  $\text{id}^{\otimes}$  with respect to the morphisms  $\text{id}_{A,B}^{\otimes}$  and  $\text{id}_C$ .
2. *Proof of Item 1b:* This follows from the naturality of  $\text{id}^{\otimes}$  with respect to the morphisms  $\text{id}_A$  and  $\text{id}_{B,C}^{\otimes}$ .

This finishes the proof.

**Item 2: Extra Monoidal Unity Constraints**We claim that **Items 2a** and **2b** are indeed true:

1. *Proof of Item 1a:* Indeed, consider the diagram

$$\begin{array}{ccccc}
 \mathbb{1}_C \otimes_C \mathbb{1}'_C & \xrightarrow{\quad \text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1} \quad} & \mathbb{1}_C \boxtimes_C \mathbb{1}'_C & \xrightarrow{\quad} & \\
 \downarrow \lambda_{\mathbb{1}'_C}^C & \searrow \text{id}_{\mathbb{1}_C} \otimes_C \text{id}_{\mathbb{1}}^\otimes & & \downarrow \text{id}_{\mathbb{1}_C} \boxtimes_C \text{id}_{\mathbb{1}}^\otimes & \\
 & (1) & & & \\
 & \mathbb{1}_C \otimes_C \mathbb{1}_C & \xrightarrow{\quad \text{id}_{\mathbb{1}_C, \mathbb{1}_C}^{\otimes, -1} \quad} & \mathbb{1}_C \boxtimes_C \mathbb{1}_C & \\
 & \downarrow \text{id}_{\mathbb{1}_C, \mathbb{1}_C}^\otimes & \searrow \text{id}_{\mathbb{1}_C, \mathbb{1}_C}^\otimes & \downarrow \text{id}_{\mathbb{1}_C, \mathbb{1}_C}^\otimes & \\
 & (2) & & (4) & \\
 & \mathbb{1}_C \otimes_C \mathbb{1}_C & \xrightarrow{\quad \text{id}_{\mathbb{1}_C, \mathbb{1}_C}^\otimes \quad} & \mathbb{1}_C & \\
 \downarrow \text{id}_{\mathbb{1}}^\otimes & (3) & \downarrow \lambda_{\mathbb{1}_C}^C = \rho_{\mathbb{1}_C}^{C,*} & \downarrow \lambda_{\mathbb{1}_C}^C = \rho_{\mathbb{1}_C}^C & \\
 & & & & 
 \end{array}$$

whose boundary diagram is the diagram whose commutativity we wish to prove. Since:

- Subdiagram (1) commutes by the naturality of  $\text{id}_C^{\otimes, -1}$ ;
- Subdiagram (2) commutes trivially;
- Subdiagram (3) commutes by the naturality of  $\lambda^C$ , where the equality  $\rho_{\mathbb{1}_C}^C = \lambda_{\mathbb{1}_C}^C$  comes from ??;

- Subdiagram (4) commutes by the right monoidal unity of  $(\text{id}_C, \text{id}_C^\otimes, \text{id}_{C|1})$ ;

so does the boundary diagram, and we are done.

2. *Proof of Item 1b:* Indeed, consider the diagram

$$\begin{array}{ccccc}
 \mathbb{1}'_C \otimes_C \mathbb{1}_C & \xrightarrow{\quad \text{id}_{\mathbb{1}'_C, \mathbb{1}_C}^{\otimes, -1} \quad} & \mathbb{1}'_C \boxtimes_C \mathbb{1}_C & \xrightarrow{\quad} & \\
 \downarrow \rho_{\mathbb{1}'_C}^C & \searrow \text{id}_1^\otimes \otimes_C \text{id}_{\mathbb{1}_C} & \downarrow \text{id}_1^\otimes \boxtimes_C \text{id}_{\mathbb{1}_C} & \downarrow \text{id}_{\mathbb{1}_C, \mathbb{1}_C}^\otimes & \downarrow \lambda_{\mathbb{1}_C}^{C, \prime} \\
 & (1) & & (2) & \\
 & \mathbb{1}_C \otimes_C \mathbb{1}_C & \xrightarrow{-\text{id}_{\mathbb{1}_C, \mathbb{1}_C}^{\otimes, -1}} & \mathbb{1}_C \boxtimes_C \mathbb{1}_C & \\
 \downarrow & \swarrow & \downarrow & \downarrow & \\
 & (3) & & (4) & \\
 & \text{id}_1^\otimes & & \rho_{\mathbb{1}_C}^C = \lambda_{\mathbb{1}_C}^C & \\
 & \searrow & & \downarrow & \\
 & & & \mathbb{1}_C & 
 \end{array}$$

whose boundary diagram is the diagram whose commutativity we wish to prove. Since:

- Subdiagram (1) commutes by the naturality of  $\text{id}_C^{\otimes, -1}$ ;

- Subdiagram (2) commutes trivially;
- Subdiagram (3) commutes by the naturality of  $\rho^C$ , where the equality  $\rho_{\mathbb{1}_C}^C = \lambda_{\mathbb{1}_C}^C$  comes from ??;
- Subdiagram (4) commutes by the left monoidal unity of  $(\text{id}_C, \text{id}_C^\otimes, \text{id}_{C|1}^\otimes)$ ;

so does the boundary diagram, and we are done.

3. *Proof of Item 2c:* Indeed, consider the diagram

$$\begin{array}{ccccc}
 \mathbb{1}'_C \otimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}'_C \boxtimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^\otimes} & \mathbb{1}'_C \otimes_C \mathbb{1}_C \\
 \downarrow \rho_{\mathbb{1}'_C}^C & & \downarrow \lambda_{\mathbb{1}_C}^{C, \prime} & & \downarrow \rho_{\mathbb{1}'_C}^C \\
 \mathbb{1}'_C & \xrightarrow{\text{id}_\mathbb{1}^\otimes} & \mathbb{1}_C & \xrightarrow{\text{id}_\mathbb{1}^{\otimes, -1}} & \mathbb{1}'_C.
 \end{array}$$

Since:

- The boundary diagram commutes trivially;
- Subdiagram (1) commutes by **Item 1b**;

it follows that the diagram

$$\begin{array}{ccccc}
 \mathbb{1}'_C \otimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}'_C \boxtimes_C \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^\otimes} & \mathbb{1}'_C \otimes_C \mathbb{1}_C \\
 \downarrow \lambda_{\mathbb{1}_C}^{C, \prime} & & \downarrow (\dagger) & & \downarrow \rho_{\mathbb{1}'_C}^C \\
 \mathbb{1}_C & \xrightarrow{\text{id}_\mathbb{1}^{\otimes, -1}} & \mathbb{1}'_C & &
 \end{array}$$

commutes. But since  $\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}$  is an isomorphism, it follows that the diagram  $(\dagger)$  also commutes, and we are done.

4. *Proof of Item 2d:* Indeed, consider the diagram

$$\begin{array}{ccccc}
 \mathbb{1}_C \otimes_C \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}_C \boxtimes_C \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}_C \otimes_C \mathbb{1}'_C \\
 \downarrow \lambda_{\mathbb{1}'_C}^C & & \downarrow \rho_{\mathbb{1}_C}^{C, \prime} & & \downarrow \lambda_{\mathbb{1}'_C}^C \\
 \mathbb{1}'_C & \xrightarrow{\text{id}_\mathbb{1}^\otimes} & \mathbb{1}_C & \xrightarrow{\text{id}_\mathbb{1}^{\otimes, -1}} & \mathbb{1}_C
 \end{array}$$

Since:

- The boundary diagram commutes trivially;
- Subdiagram (1) commutes by [Item 1a](#);

it follows that the diagram

$$\begin{array}{ccccc}
 \mathbb{1}_C \otimes_C \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}_C \boxtimes_C \mathbb{1}'_C & \xrightarrow{\text{id}_{\mathbb{1}_C, \mathbb{1}'_C}^{\otimes, -1}} & \mathbb{1}_C \otimes_C \mathbb{1}'_C \\
 \downarrow \rho_{\mathbb{1}_C}^{C,*} & & \downarrow (\dagger) & & \downarrow \lambda_{\mathbb{1}'_C}^C \\
 \mathbb{1}_C & \xrightarrow{\text{id}_{\mathbb{1}}^{\otimes, -1}} & \mathbb{1}_C & &
 \end{array}$$

commutes. But since  $\text{id}_{\mathbb{1}}^{\otimes, -1}$  is an isomorphism, it follows that the diagram  $(\dagger)$  also commutes, and we are done.

This finishes the proof.

### Item 3: Mixed Associators

We claim that [Items 3a](#) to [3c](#) are indeed true:

01UZ

1. *Proof of Item 3a:* We may partition the monoidality diagram for  $\text{id}^\otimes$  of [Item 2 of Remark 13.1.1.1.3](#) as follows:

$$\begin{array}{ccccc}
 & & (A \boxtimes_C B) \boxtimes_C C & & \\
 & \swarrow \text{id}_{A,B}^\otimes \boxtimes_C \text{id}_C & \downarrow & \searrow \alpha_{A,B,C}^C & \\
 (A \otimes_C B) \boxtimes_C C & & \text{id}_{A \otimes_C B, C}^\otimes & & A \boxtimes_C (B \boxtimes_C C) \\
 \downarrow \text{id}_{A \otimes_C B, C}^\otimes & \quad (1) \quad & \downarrow & \quad (2) \quad & \downarrow \text{id}_A \boxtimes_C \text{id}_{B,C}^\otimes \\
 & \swarrow \text{id}_{A,B}^\otimes \otimes_C \text{id}_C & & \searrow \alpha_{A,B,C}^\otimes & \\
 (A \otimes_C B) \otimes_C C & & \text{id}_{A,B \otimes_C C}^\otimes & & A \boxtimes_C (B \otimes_C C) \\
 \downarrow \alpha_{A,B,C}^{C,*} & \quad (3) \quad & \downarrow & & \downarrow \text{id}_{A,B \otimes_C C}^\otimes \\
 & \swarrow & & \searrow & \\
 & A \otimes_C (B \otimes_C C) & & &
 \end{array}$$

Since:

- Subdiagram (1) commutes by [Item 1a](#) of [Item 1](#).

- Subdiagram (2) commutes by assumption.
- Subdiagram (3) commutes by assumption.

it follows that the boundary diagram also commutes, i.e.  $\text{id}^\otimes$  satisfies the monoidality condition of **Item 2 of Remark 13.1.1.3**.

**01V0** 2. *Proof of Item 3b:* We may partition the monoidality diagram for  $\text{id}^\otimes$  of **Item 2 of Remark 13.1.1.3** as follows:

$$\begin{array}{ccccc}
 & & (A \boxtimes_C B) \boxtimes_C C & & \\
 & \swarrow \text{id}_{A,B}^\otimes \boxtimes_C \text{id}_C & & \searrow \alpha_{A,B,C}^C & \\
 (A \otimes_C B) \boxtimes_C C & & (1) & & A \boxtimes_C (B \boxtimes_C C) \\
 \downarrow \text{id}_{A \otimes_C B,C}^\otimes & \swarrow \alpha_{A,B,C}^\otimes & & \downarrow \text{id}_{A,B \otimes_C C}^\otimes & \downarrow \text{id}_A \boxtimes_C \text{id}_{B,C}^\otimes \\
 & A \otimes_C (B \boxtimes_C C) & & & \\
 \downarrow \text{id}_A \otimes_C \text{id}_{B,C}^\otimes & & (2) & & \downarrow \text{id}_A \boxtimes_C (B \otimes_C C) \\
 (A \otimes_C B) \otimes_C C & & & & A \boxtimes_C (B \otimes_C C) \\
 \downarrow \alpha_{A,B,C}^{C,\prime} & & & & \downarrow \text{id}_{A,B \otimes_C C}^\otimes \\
 & A \otimes_C (B \otimes_C C) & & &
 \end{array}$$

Since:

- Subdiagram (1) commutes by assumption.
- Subdiagram (2) commutes by assumption.
- Subdiagram (3) commutes by **Item 1b of Item 1**.

it follows that the boundary diagram also commutes, i.e.  $\text{id}^\otimes$  satisfies the monoidality condition of **Item 2 of Remark 13.1.1.3**.

**01V1** 3. *Proof of Item 3c:* We may partition the monoidality diagram for

$\text{id}^\otimes$  of Item 2 of Remark 13.1.1.1.3 as follows:

$$\begin{array}{ccccc}
& & (A \otimes_C B) \otimes_C C & & \\
& \swarrow & & \searrow & \\
\text{id}_{A,B}^{\otimes} \otimes_C \text{id}_C & & & & \alpha_{A,B,C}^C \\
& \nwarrow & & & \\
(A \otimes_C B) \otimes_C C & & & & A \otimes_C (B \otimes_C C) \\
& \downarrow & & & \downarrow \text{id}_A \otimes_C \text{id}_{B,C}^{\otimes} \\
& & (1) & & \\
& & \alpha_{A,B,C}^{\otimes, \otimes} & & \\
& \searrow & & \swarrow & \\
& & (2) & & \\
& & & & \\
(A \otimes_C B) \otimes_C C & & & & A \otimes_C (B \otimes_C C) \\
& \swarrow & & \searrow & \\
& \alpha_{A,B,C}^{C,\prime} & & & \text{id}_{A,B \otimes_C C}^{\otimes} \\
& & & & \\
& & & & A \otimes_C (B \otimes_C C)
\end{array}$$

Since subdiagrams (1) and (2) commute by assumption, it follows that the boundary diagram also commutes, i.e.  $\text{id}^\otimes$  satisfies the monoidality condition of [Item 2 of Remark 13.1.1.1.3](#).

This finishes the proof.

## 01V2 13.1.2 The Moduli Category of Braided Monoidal Structures on a Category

### 01V3 13.1.3 The Moduli Category of Symmetric Monoidal Structures on a Category

## 01V4 13.2 Moduli Categories of Closed Monoidal Structures

## 01v5 13.3 Moduli Categories of Refinements of Monoidal Structures

### 01V6 13.3.1 The Moduli Category of Braided Refinements of a Monoidal Structure

## Appendices

## 13.A Other Chapters

<b>Preliminaries</b>	10. Conditions on Relations
1. Introduction	
2. A Guide to the Literature	
<b>Sets</b>	<b>Categories</b>
3. Sets	11. Categories
4. Constructions With Sets	
5. Monoidal Structures on the Category of Sets	12. Presheaves and the Yoneda Lemma
6. Pointed Sets	
7. Tensor Products of Pointed Sets	
<b>Relations</b>	<b>Monoidal Categories</b>
8. Relations	13. Constructions With Monoidal Categories
9. Constructions With Relations	
	<b>Bicategories</b>
	14. Types of Morphisms in Bicategories
	<b>Extra Part</b>
	15. Notes

## **Part VI**

# **Bicategories**

## Chapter 14

# Types of Morphisms in Bicategories

**02H3** In this chapter, we study special kinds of morphisms in bicategories:

1. *Monomorphisms and Epimorphisms in Bicategories* ([Sections 14.1 and 14.2](#)).

There is a large number of different notions capturing the idea of a “monomorphism” or of an “epimorphism” in a bicategory.

Arguably, the notion that best captures these concepts is that of a *pseudomonic morphism* ([Definition 14.1.10.1.1](#)) and of a *pseudoepic morphism* ([Definition 14.2.10.1.1](#)), although the other notions introduced in [Sections 14.1](#) and [14.2](#) are also interesting on their own.

### Contents

---

14.1 Monomorphisms in Bicategories .....	886
14.1.1 Representably Faithful Morphisms .....	886
14.1.2 Representably Full Morphisms.....	887
14.1.3 Representably Fully Faithful Morphisms.....	888
14.1.4 Morphisms Representably Faithful on Cores .....	889
14.1.5 Morphisms Representably Full on Cores .....	890
14.1.6 Morphisms Representably Fully Faithful on Cores	
891	
14.1.7 Representably Essentially Injective Morphisms....	892
14.1.8 Representably Conservative Morphisms.....	893
14.1.9 Strict Monomorphisms.....	893
14.1.10 Pseudomonic Morphisms.....	894
14.2 Epimorphisms in Bicategories.....	896
14.2.1 Corepresentably Faithful Morphisms.....	896
14.2.2 Corepresentably Full Morphisms.....	897
14.2.3 Corepresentably Fully Faithful Morphisms .....	898

14.2.4 Morphisms Corepresentably Faithful on Cores ....	901
14.2.5 Morphisms Corepresentably Full on Cores .....	901
14.2.6 Morphisms Corepresentably Fully Faithful on Cores .....	902
14.2.7 Corepresentably Essentially Injective Morphisms.	904
14.2.8 Corepresentably Conservative Morphisms.....	904
14.2.9 Strict Epimorphisms.....	905
14.2.10 Pseudoepic Morphisms .....	906
14.A Other Chapters .....	908

## 019J 14.1 Monomorphisms in Bicategories

### 019K 14.1.1 Representably Faithful Morphisms

Let  $C$  be a bicategory.

#### 019L DEFINITION 14.1.1.1 ► REPRESENTABLY FAITHFUL MORPHISMS

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably faithful**<sup>1</sup> if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is faithful.

<sup>1</sup>Further Terminology: Also called simply a **faithful morphism**, based on Item 1 of Example 14.1.1.3.

#### 019M REMARK 14.1.1.2 ► UNWINDING DEFINITION 14.1.1.1

In detail,  $f$  is representably faithful if, for all diagrams in  $C$  of the form

$$\begin{array}{ccc} X & \xrightarrow{\quad \phi \quad} & A \\ \alpha \parallel \beta & \nearrow \psi & \downarrow f \\ & & B \end{array}$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

#### 019N EXAMPLE 14.1.1.3 ► EXAMPLES OF REPRESENTABLY FAITHFUL MORPHISMS

Here are some examples of representably faithful morphisms.

019P

1. *Representably Faithful Morphisms in  $\text{Cats}_2$* . The representably faithful morphisms in  $\text{Cats}_2$  are precisely the faithful functors; see [Item 2 of Proposition 11.6.1.1.2](#).

019Q

2. *Representably Faithful Morphisms in  $\text{Rel}$* . Every morphism of  $\text{Rel}$  is representably faithful; see [Old Tag 15.2.1.1.14](#) of [Old Tag 15.2.1.1.13](#).

### 019R 14.1.2 Representably Full Morphisms

Let  $C$  be a bicategory.

019S

#### DEFINITION 14.1.2.1.1 ► REPRESENTABLY FULL MORPHISMS

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably full**<sup>1</sup> if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is full.

<sup>1</sup>*Further Terminology:* Also called simply a **full morphism**, based on [Item 1 of Example 14.1.2.1.3](#).

019T

#### REMARK 14.1.2.1.2 ► UNWINDING DEFINITION 14.1.2.1.1

In detail,  $f$  is representably full if, for each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: f \circ \phi \Rightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\[-1ex] \beta \Downarrow \\[-1ex] \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \psi \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \psi \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\[-1ex] \beta \Downarrow \\[-1ex] f \circ \psi \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

019U

**EXAMPLE 14.1.2.1.3 ► EXAMPLES OF REPRESENTABLY FULL MORPHISMS**

Here are some examples of representably full morphisms.

019V

1. *Representably Full Morphisms in  $\text{Cats}_2$ .* The representably full morphisms in  $\text{Cats}_2$  are precisely the full functors; see Item 1 of Proposition 11.6.2.1.2.

019W

2. *Representably Full Morphisms in  $\text{Rel}$ .* The representably full morphisms in  $\text{Rel}$  are characterised in Old Tag 15.2.1.1.15 of Old Tag 15.2.1.1.13.

**019X 14.1.3 Representably Fully Faithful Morphisms**

Let  $C$  be a bicategory.

019Y

**DEFINITION 14.1.3.1.1 ► REPRESENTABLY FULLY FAITHFUL MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably fully faithful**<sup>1</sup> if the following equivalent conditions are satisfied:

019Z

1. The 1-morphism  $f$  is representably faithful (Definition 14.1.1.1.1) and representably full (Definition 14.1.2.1.1).

01A0

2. For each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is fully faithful.

<sup>1</sup>Further Terminology: Also called simply a **fully faithful morphism**, based on Item 1 of Example 14.1.3.1.3.

01A1

**REMARK 14.1.3.1.2 ► UNWINDING REPRESENTABLY FULLY FAITHFUL MORPHISMS**

In detail,  $f$  is representably fully faithful if the conditions in Remark 14.1.1.1.2 and Remark 14.1.2.1.2 hold:

1. For all diagrams in  $C$  of the form

$$X \xrightarrow[\psi]{\alpha \parallel \beta} A \xrightarrow{f} B,$$

$\phi$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: f \circ \phi \Rightarrow f \circ \psi, \quad X \xrightarrow[\psi]{\beta} \xrightarrow[\phi]{f \circ \phi} B$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \xrightarrow[\psi]{\alpha} \xrightarrow[\phi]{\phi} A$$

of  $C$  such that we have an equality

$$X \xrightarrow[\psi]{\alpha} \xrightarrow[\phi]{\phi} A \xrightarrow{f} B = X \xrightarrow[\psi]{\beta} \xrightarrow[\phi]{f \circ \phi} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

01A2

#### EXAMPLE 14.1.3.1.3 ► EXAMPLES OF REPRESENTABLY FULLY FAITHFUL MORPHISMS

Here are some examples of representably fully faithful morphisms.

01A3

1. *Representably Fully Faithful Morphisms in  $\text{Cats}_2$* . The representably fully faithful morphisms in  $\text{Cats}_2$  are precisely the fully faithful functors; see Item 6 of Proposition 11.6.3.1.2.

01A4

2. *Representably Fully Faithful Morphisms in  $\text{Rel}$* . The representably fully faithful morphisms of  $\text{Rel}$  coincide (Old Tag 15.2.1.1.22 of Old Tag 15.2.1.1.13) with the representably full morphisms in  $\text{Rel}$ , which are characterised in Old Tag 15.2.1.1.15 of Old Tag 15.2.1.1.13.

01A5

#### 14.1.4 Morphisms Representably Faithful on Cores

Let  $C$  be a bicategory.

01A6

#### DEFINITION 14.1.4.1.1 ► MORPHISMS REPRESENTABLY FAITHFUL ON CORES

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably faithful on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by  $f$  is faithful.

01A7

**REMARK 14.1.4.1.2 ► UNWINDING DEFINITION 14.1.4.1.1**

In detail,  $f$  is representably faithful on cores if, for all diagrams in  $C$  of the form

$$\begin{array}{ccc} X & \xrightarrow{\phi} & A \\ \alpha \parallel \beta \downarrow \downarrow & \nearrow \psi & \xrightarrow{f} \\ & & B \end{array}$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

**01A8 14.1.5 Morphisms Representably Full on Cores**

Let  $C$  be a bicategory.

01A9

**DEFINITION 14.1.5.1.1 ► MORPHISMS REPRESENTABLY FULL ON CORES**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably full on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by  $f$  is full.

01AA

**REMARK 14.1.5.1.2 ► UNWINDING DEFINITION 14.1.5.1.1**

In detail,  $f$  is representably full on cores if, for each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: f \circ \phi \xrightarrow{\sim} f \circ \psi, \quad \begin{array}{ccc} X & \xrightarrow{\beta} & B \\ \beta \downarrow & \nearrow f \circ \phi & \\ & & f \circ \phi \end{array}$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \psi \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \psi \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\[-1ex] \beta \Downarrow \\[-1ex] f \circ \psi \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

### 01AB 14.1.6 Morphisms Representably Fully Faithful on Cores

Let  $C$  be a bicategory.

01AC

#### DEFINITION 14.1.6.1.1 ► MORPHISMS REPRESENTABLY FULLY FAITHFUL ON CORES

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably fully faithful on cores** if the following equivalent conditions are satisfied:

01AD

1. The 1-morphism  $f$  is representably faithful on cores (Definition 14.1.5.1.1) and representably full on cores (Definition 14.1.4.1.1).

01AE

2. For each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by  $f$  is fully faithful.

01AF

#### REMARK 14.1.6.1.2 ► UNWINDING DEFINITION 14.1.6.1.1

In detail,  $f$  is representably fully faithful on cores if the conditions in Remark 14.1.4.1.2 and Remark 14.1.5.1.2 hold:

1. For all diagrams in  $C$  of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \beta \Downarrow \\[-1ex] \psi \end{array} A \xrightarrow{f} B,$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: f \circ \phi \xrightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\[-1ex] \beta \Downarrow \\[-1ex] \xrightarrow{f \circ \psi} \end{array} B$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \xrightarrow{\psi} \end{array} A$$

of  $C$  such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \psi \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\[-1ex] \beta \Downarrow \\[-1ex] \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

### 01AG 14.1.7 Representably Essentially Injective Morphisms

Let  $C$  be a bicategory.

#### 01AH DEFINITION 14.1.7.1.1 ► REPRESENTABLY ESSENTIALLY INJECTIVE MORPHISMS

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably essentially injective** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is essentially injective.

#### 01AJ REMARK 14.1.7.1.2 ► UNWINDING DEFINITION 14.1.7.1.1

In detail,  $f$  is representably essentially injective if, for each pair of morphisms  $\phi, \psi: X \rightrightarrows A$  of  $C$ , the following condition is satisfied:

(★) If  $f \circ \phi \cong f \circ \psi$ , then  $\phi \cong \psi$ .

### 01AK 14.1.8 Representably Conservative Morphisms

Let  $C$  be a bicategory.

01AL

#### DEFINITION 14.1.8.1.1 ► REPRESENTABLY CONSERVATIVE MORPHISMS

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **representably conservative** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is conservative.

01AM

#### REMARK 14.1.8.1.2 ► UNWINDING DEFINITION 14.1.8.1.1

In detail,  $f$  is representably conservative if, for each pair of morphisms  $\phi, \psi: X \rightrightarrows A$  and each 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \Downarrow \alpha \\[-1ex] \xrightarrow{\psi} \end{array} A$$

of  $C$ , if the 2-morphism

$$\text{id}_f \star \alpha: f \circ \phi \Rightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\[-1ex] \Downarrow \text{id}_f \star \alpha \\[-1ex] \xrightarrow{f \circ \psi} \end{array} B$$

is a 2-isomorphism, then so is  $\alpha$ .

### 01AN 14.1.9 Strict Monomorphisms

Let  $C$  be a bicategory.

01AP

#### DEFINITION 14.1.9.1.1 ► STRICT MONOMORPHISMS

A 1-morphism  $f: A \rightarrow B$  of  $C$  is a **strict monomorphism** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is injective on objects, i.e. its action on objects

$$f_*: \text{Obj}(\text{Hom}_C(X, A)) \rightarrow \text{Obj}(\text{Hom}_C(X, B))$$

is injective.

01AQ

**REMARK 14.1.9.1.2 ► UNWINDING DEFINITION 14.1.9.1.1**

In detail,  $f$  is a strict monomorphism in  $C$  if, for each diagram in  $C$  of the form

$$X \xrightarrow[\psi]{\phi} A \xrightarrow{f} B,$$

if  $f \circ \phi = f \circ \psi$ , then  $\phi = \psi$ .

01AR

**EXAMPLE 14.1.9.1.3 ► EXAMPLES OF STRICT MONOMORPHISMS**

Here are some examples of strict monomorphisms.

01AS

1. *Strict Monomorphisms in Cats<sub>2</sub>*. The strict monomorphisms in Cats<sub>2</sub> are precisely the functors which are injective on objects and injective on morphisms; see Item 1 of Proposition 11.7.2.1.2.

01AT

2. *Strict Monomorphisms in Rel*. The strict monomorphisms in Rel are characterised in Old Tag 15.2.1.1.11.

**01AU 14.1.10 Pseudomonic Morphisms**

Let  $C$  be a bicategory.

01AV

**DEFINITION 14.1.10.1.1 ► PSEUDOMONIC MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **pseudomonic** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by  $f$  is pseudomonic.

01AW

**REMARK 14.1.10.1.2 ► UNWINDING DEFINITION 14.1.10.1.1**

In detail, a 1-morphism  $f: A \rightarrow B$  of  $C$  is pseudomonic if it satisfies the following conditions:

01AX

1. For all diagrams in  $C$  of the form

$$X \xrightarrow[\psi]{\alpha \parallel \beta} A \xrightarrow{f} B,$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then  $\alpha = \beta$ .

01AY

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: f \circ \phi \xrightarrow{\sim} f \circ \psi, \quad X \xrightarrow[\psi]{\beta \Downarrow} B$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \xrightarrow[\psi]{\alpha \Downarrow} A$$

of  $C$  such that we have an equality

$$X \xrightarrow[\psi]{\alpha \Downarrow} A \xrightarrow{f} B = X \xrightarrow[\psi]{\beta \Downarrow} B$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

01AZ

#### PROPOSITION 14.1.10.1.3 ► PROPERTIES OF PSEUDOMONIC MORPHISMS

Let  $f: A \rightarrow B$  be a 1-morphism of  $C$ .

01B0

1. *Characterisations.* The following conditions are equivalent:

01B1

(a) The morphism  $f$  is pseudomonic.

01B2

(b) The morphism  $f$  is representably full on cores and representably faithful.

01B3

(c) We have an isocomma square of the form

$$\begin{array}{ccc} A & \xrightarrow{\text{id}_A} & A \\ A \xrightarrow{\text{eq.}} A \times_B A, & \downarrow \text{id}_A & \downarrow F \\ & \lrcorner \swarrow \nearrow \lrcorner & \\ A & \xrightarrow{F} & B \end{array}$$

in  $C$  up to equivalence.

01B4

2. *Interaction With Cotensors.* If  $C$  has cotensors with  $\mathbb{1}$ , then the following conditions are equivalent:

- (a) The morphism  $f$  is pseudomonic.
- (b) We have an isocomma square of the form

$$\begin{array}{ccc} A & \hookrightarrow & \mathbb{1} \pitchfork A \\ A \xrightarrow{\text{eq.}} A \times_{\mathbb{1} \pitchfork F} B, & \downarrow F & \downarrow \mathbb{1} \pitchfork F \\ & \lrcorner \swarrow \nearrow \lrcorner & \\ B & \hookrightarrow & \mathbb{1} \pitchfork B \end{array}$$

in  $C$  up to equivalence.

**PROOF 14.1.10.1.4 ► PROOF OF PROPOSITION 14.1.10.1.3**

Item 1: Characterisations

Omitted.

Item 2: Interaction With Cotensors

Omitted. 

## 01B5 14.2 Epimorphisms in Bicategories

### 01B6 14.2.1 Corepresentably Faithful Morphisms

Let  $C$  be a bicategory.

01B7

**DEFINITION 14.2.1.1.1 ► COREPRESENTABLY FAITHFUL MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably faithful** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is faithful.

01B8

## REMARK 14.2.1.1.2 ▶ UNWINDING DEFINITION 14.2.1.1.1

In detail,  $f$  is corepresentably faithful if, for all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \beta \\[-1ex] \xleftarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

01B9

## EXAMPLE 14.2.1.1.3 ▶ EXAMPLES OF COREPRESENTABLY FAITHFUL MORPHISMS

Here are some examples of corepresentably faithful morphisms.

01BA

1. *Corepresentably Faithful Morphisms in  $\text{Cats}_2$* . The corepresentably faithful morphisms in  $\text{Cats}_2$  are characterised in Item 5 of Proposition 11.6.1.1.2.

01BB

2. *Corepresentably Faithful Morphisms in  $\text{Rel}$* . Every morphism of  $\text{Rel}$  is corepresentably faithful; see Old Tag 15.2.1.1.27 of Old Tag 15.2.1.1.26.

## 01BC 14.2.2 Corepresentably Full Morphisms

Let  $C$  be a bicategory.

01BD

## DEFINITION 14.2.2.1.1 ▶ COREPRESENTABLY FULL MORPHISMS

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably full** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is full.

01BE

## REMARK 14.2.2.1.2 ▶ UNWINDING DEFINITION 14.2.2.1.1

In detail,  $f$  is corepresentably full if, for each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: \phi \circ f \Rightarrow \psi \circ f, \quad A \xrightarrow[\psi \circ f]{\beta \Downarrow} X$$

$\phi \circ f$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad B \xrightarrow[\psi]{\alpha \Downarrow} X$$

$\phi$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \xrightarrow[\psi]{\alpha \Downarrow} X = A \xrightarrow[\psi \circ f]{\beta \Downarrow} X$$

$\phi$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01BF

**EXAMPLE 14.2.2.1.3 ► EXAMPLES OF COREPRESENTABLY FULL MORPHISMS**

Here are some examples of corepresentably full morphisms.

01BG

1. *Corepresentably Full Morphisms in  $\text{Cats}_2$ .* The corepresentably full morphisms in  $\text{Cats}_2$  are characterised in Item 7 of Proposition 11.6.2.1.2.

01BH

2. *Corepresentably Full Morphisms in  $\text{Rel}$ .* The corepresentably full morphisms in  $\text{Rel}$  are characterised in ?? of Old Tag 15.2.1.1.13.

**01BJ 14.2.3 Corepresentably Fully Faithful Morphisms**

Let  $C$  be a bicategory.

01BK

**DEFINITION 14.2.3.1.1 ► COREPRESENTABLY FULLY FAITHFUL MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably fully faithful**<sup>1</sup> if the following equivalent conditions are satisfied:

01BL

1. The 1-morphism  $f$  is corepresentably full (Definition 14.2.2.1.1) and corepresentably faithful (Definition 14.2.1.1.1).

01BM

2. For each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is fully faithful.

<sup>1</sup>*Further Terminology:* Corepresentably fully faithful morphisms have also been called **lax epimorphisms** in the literature (e.g. in [Adá+01]), though we will always use the name “corepresentably fully faithful morphism” instead in this work.

01BN

#### REMARK 14.2.3.1.2 ► UNWINDING DEFINITION 14.2.3.1.1

In detail,  $f$  is corepresentably fully faithful if the conditions in Remark 14.2.1.1.2 and Remark 14.2.2.1.2 hold:

1. For all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \beta \\ \searrow \psi \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-morphism

$$\beta: \phi \circ f \Rightarrow \psi \circ f, \quad A \begin{array}{c} \nearrow \phi \circ f \\ \beta \Downarrow \\ \searrow \psi \circ f \end{array} X$$

of  $C$ , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \\ \searrow \psi \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \\ \searrow \psi \end{array} X = A \begin{array}{c} \nearrow \phi \circ f \\ \beta \Downarrow \\ \searrow \psi \circ f \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01BP

Here are some examples of corepresentably fully faithful morphisms.

01BQ

1. *Corepresentably Fully Faithful Morphisms in  $\text{Cats}_2$* . The fully faithful epimorphisms in  $\text{Cats}_2$  are characterised in [Item 10 of Proposition 11.6.3.1.2](#).

01BR

2. *Corepresentably Fully Faithful Morphisms in  $\text{Rel}$* . The corepresentably fully faithful morphisms of  $\text{Rel}$  coincide ([Old Tag 15.2.1.1.35](#) of [Old Tag 15.2.1.1.26](#)) with the corepresentably full morphisms in  $\text{Rel}$ , which are characterised in [Old Tag 15.2.1.1.28](#) of [Old Tag 15.2.1.1.26](#).

#### 01BS 14.2.4 Morphisms Corepresentably Faithful on Cores

Let  $C$  be a bicategory.

01BT

##### DEFINITION 14.2.4.1.1 ► MORPHISMS COREPRESENTABLY FAITHFUL ON CORES

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably faithful on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by  $f$  is faithful.

01BU

##### REMARK 14.2.4.1.2 ► UNWINDING DEFINITION 14.2.4.1.1

In detail,  $f$  is corepresentably faithful on cores if, for all diagrams in  $C$  of the form

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \alpha \Downarrow \beta & \Downarrow \psi \\ & \phi & \end{array}$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

#### 01BV 14.2.5 Morphisms Corepresentably Full on Cores

Let  $C$  be a bicategory.

01BW

**DEFINITION 14.2.5.1.1 ► MORPHISMS COREPRESENTABLY FULL ON CORES**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably full on cores** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by  $f$  is full.

01BX

**REMARK 14.2.5.1.2 ► UNWINDING DEFINITION 14.2.5.1.1**

In detail,  $f$  is corepresentably full on cores if, for each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ f \xrightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\[-1ex] \beta \Downarrow \\[-1ex] \xrightarrow{\psi \circ f} \end{array} X$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \xrightarrow{\psi} \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \alpha \Downarrow \\[-1ex] \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\[-1ex] \beta \Downarrow \\[-1ex] \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

**01BY 14.2.6 Morphisms Corepresentably Fully Faithful on Cores**

Let  $C$  be a bicategory.

01BZ

**DEFINITION 14.2.6.1.1 ► MORPHISMS COREPRESENTABLY FULLY FAITHFUL ON CORES**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably fully faithful on cores** if the following equivalent conditions are satisfied:

01C0

1. The 1-morphism  $f$  is corepresentably full on cores (Definition 14.2.5.1.1) and corepresentably faithful on cores (Definition 14.2.5.1.2).

tion 14.2.1.1.1).

**01C1** 2. For each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by  $f$  is fully faithful.

**01C2**

**REMARK 14.2.6.1.2 ► UNWINDING DEFINITION 14.2.6.1.1**

In detail,  $f$  is corepresentably fully faithful on cores if the conditions in [Remark 14.2.4.1.2](#) and [Remark 14.2.5.1.2](#) hold:

1. For all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \xrightarrow{\phi} X, \quad \begin{array}{c} \alpha \parallel \beta \\ \downarrow \end{array}$$

if  $\alpha$  and  $\beta$  are 2-isomorphisms and we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ f \xrightarrow{\sim} \psi \circ f, \quad A \xrightarrow{\phi \circ f} X, \quad \begin{array}{c} \beta \parallel \\ \downarrow \end{array}$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \xrightarrow{\phi} X, \quad \begin{array}{c} \alpha \parallel \\ \downarrow \end{array}$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \xrightarrow{\phi} X = A \xrightarrow{\psi \circ f} X, \quad \begin{array}{c} \alpha \parallel \\ \downarrow \end{array}$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

**01C3 14.2.7 Corepresentably Essentially Injective Morphisms**

Let  $C$  be a bicategory.

01C4

**DEFINITION 14.2.7.1.1 ► COREPRESENTABLY ESSENTIALLY INJECTIVE MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably essentially injective** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is essentially injective.

01C5

**REMARK 14.2.7.1.2 ► UNWINDING DEFINITION 14.2.7.1.1**

In detail,  $f$  is corepresentably essentially injective if, for each pair of morphisms  $\phi, \psi: B \rightrightarrows X$  of  $C$ , the following condition is satisfied:

- (★) If  $\phi \circ f \cong \psi \circ f$ , then  $\phi \cong \psi$ .

**01C6 14.2.8 Corepresentably Conservative Morphisms**

Let  $C$  be a bicategory.

01C7

**DEFINITION 14.2.8.1.1 ► COREPRESENTABLY CONSERVATIVE MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **corepresentably conservative** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is conservative.

01C8

**REMARK 14.2.8.1.2 ► UNWINDING DEFINITION 14.2.8.1.1**

In detail,  $f$  is corepresentably conservative if, for each pair of morphisms  $\phi, \psi: B \rightrightarrows X$  and each 2-morphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\[-1ex] \Downarrow \alpha \\[-1ex] \xrightarrow{\psi} \end{array} X$$

of  $C$ , if the 2-morphism

$$\alpha \star \text{id}_f : \phi \circ f \Longrightarrow \psi \circ f, \quad A \begin{array}{c} \nearrow \phi \circ f \\ \Downarrow \alpha \star \text{id}_f \\ \searrow \psi \circ f \end{array} X$$

is a 2-isomorphism, then so is  $\alpha$ .

### 01C9 14.2.9 Strict Epimorphisms

Let  $C$  be a bicategory.

01CA

#### DEFINITION 14.2.9.1.1 ► STRICT EPIMORPHISMS

A 1-morphism  $f: A \rightarrow B$  is a **strict epimorphism** in  $C$  if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is injective on objects, i.e. its action on objects

$$f_*: \text{Obj}(\text{Hom}_C(B, X)) \rightarrow \text{Obj}(\text{Hom}_C(A, X))$$

is injective.

01CB

#### REMARK 14.2.9.1.2 ► UNWINDING DEFINITION 14.2.9.1.1

In detail,  $f$  is a strict epimorphism if, for each diagram in  $C$  of the form

$$A \xrightarrow{f} B \rightrightarrows X,$$

if  $\phi \circ f = \psi \circ f$ , then  $\phi = \psi$ .

01CC

#### EXAMPLE 14.2.9.1.3 ► EXAMPLES OF STRICT EPIMORPHISMS

Here are some examples of strict epimorphisms.

01CD

1. *Strict Epimorphisms in  $\text{Cats}_2$* . The strict epimorphisms in  $\text{Cats}_2$  are characterised in Item 1 of Proposition 11.7.3.1.2.

01CE

2. *Strict Epimorphisms in  $\text{Rel}$* . The strict epimorphisms in  $\text{Rel}$  are characterised in Old Tag 15.2.1.1.23.

**01CF 14.2.10 Pseudoepic Morphisms**

Let  $C$  be a bicategory.

01CG

**DEFINITION 14.2.10.1.1 ► PSEUDOEPIC MORPHISMS**

A 1-morphism  $f: A \rightarrow B$  of  $C$  is **pseudoepic** if, for each  $X \in \text{Obj}(C)$ , the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by  $f$  is pseudomonic.

01CH

**REMARK 14.2.10.1.2 ► UNWINDING DEFINITION 14.2.10.1.1**

In detail, a 1-morphism  $f: A \rightarrow B$  of  $C$  is pseudoepic if it satisfies the following conditions:

01CJ

1. For all diagrams in  $C$  of the form

$$A \xrightarrow{f} B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \beta \\ \searrow \psi \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then  $\alpha = \beta$ .

01CK

2. For each  $X \in \text{Obj}(C)$  and each 2-isomorphism

$$\beta: \phi \circ f \xrightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \nearrow \phi \circ f \\ \beta \Downarrow \\ \searrow \psi \circ f \end{array} X$$

of  $C$ , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \\ \searrow \psi \end{array} X$$

of  $C$  such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \\ \searrow \psi \end{array} X = A \begin{array}{c} \nearrow \phi \circ f \\ \beta \Downarrow \\ \searrow \psi \circ f \end{array} X$$

of pasting diagrams in  $C$ , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01CL

Let  $f: A \rightarrow B$  be a 1-morphism of  $C$ .

**01CM** 1. *Characterisations.* The following conditions are equivalent:

- 01CN** (a) The morphism  $f$  is pseudoepic.  
**01CP** (b) The morphism  $f$  is corepresentably full on cores and corepresentably faithful.  
**01CQ** (c) We have an isococomma square of the form

$$\begin{array}{ccc} B & \xleftarrow{\text{id}_B} & B \\ \uparrow \text{id}_B & \nearrow \lrcorner \urcorner & \uparrow F \\ B & \xleftarrow{F} & A \end{array}$$

$B \xrightleftharpoons{\text{eq.}} B \coprod_A B,$

in  $C$  up to equivalence.

**PROOF** 14.2.10.1.4 ► **PROOF OF PROPOSITION** 14.2.10.1.3

Item 1: Characterisations

Omitted. 

## Appendices

### 14.A Other Chapters

#### Preliminaries

1. Introduction
2. A Guide to the Literature

#### Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

#### Relations

8. Relations
9. Constructions With Relations
10. Conditions on Relations

#### Categories

11. Categories
12. Presheaves and the Yoneda Lemma

#### Monoidal Categories

13. Constructions With Monoidal Categories

**Bicategories**

14. Types of Morphisms in Bicategories

**Extra Part**

15. Notes

**Part VII**

**Extra Part**

# Chapter 15

## Notes

**02H3** This chapter contains some notes.

### Contents

---

15.1	TikZ Code for Commutative Diagrams.....	911
15.1.1	Product Diagram With Circular Arrows .....	912
15.1.2	Coproduct Diagram With Circular Arrows.....	914
15.1.3	Cube Diagram.....	916
15.1.4	Cube Diagram With Labelled Faces.....	917
15.1.5	Pentagon Diagram .....	920
15.1.6	Hexagon Diagram.....	921
15.1.7	Double Square Diagram.....	922
15.1.8	Double Hexagon Diagram.....	924
15.2	Retired Tags.....	926
15.2.1	Relations .....	926
15.2.2	Pointed Sets .....	930
15.2.3	Tensor Products of Pointed Sets.....	930
15.2.4	Categories.....	931
15.3	Miscellany .....	932
15.3.1	List of Things To Explore/Add.....	932
15.A	Other Chapters.....	978

---

### **01VR** 15.1 TikZ Code for Commutative Diagrams

In this section we gather some useful examples of `tikzcd` code for commutative diagrams.

**02C1 15.1.1 Product Diagram With Circular Arrows**

Define

```
\newlength{\DL}
\setlength{\DL}{0.9em}
```

in the preamble, as well as

```
\tikzcdset{
    productArrows/.style args={#1#2#3}={
        execute at end picture={
            % FIRST ARROW
            % Step 1: Draw arrow body
            \begin{scope}
                \clip (\tikzcdmatrixname-1-2.east) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-3.north) -- (\tikzcdmatrixname-1-3.center) -- cycle;
                \path[draw, line width=rule_thickness] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=0,radius=#1];
            \end{scope}
            % Step 2: Draw arrow head
            % Step 2.1: Find the point at which to place the arrowhead
            \path[name path=curve-1-a] (\tikzcdmatrixname-1-2.east) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-3.north) -- (\tikzcdmatrixname-1-3.center) -- cycle;
            \path[name path=curve-1-b] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=0,radius=#1];
            \fill [name intersections={of=curve-1-a and curve-1-b}] (intersection-2);
            % Step 2.2: Find the angle at which to place the arrowhead
            \coordinate (arc-start) at (\tikzcdmatrixname-1-2.east);
            \coordinate (arc-center) at (\tikzcdmatrixname-2-2.center);
            \draw let
                \p1 = ($(intersection-2) - (arc-center)$), % \p1 is the vector from the arc
                \n1 = {atan2(\y1, \x1)}, % \n1 is the angle of that vector in degrees
                \n2 = {\n1 - 90} % \n2 is the angle of the tangent (90 degrees from the radius)
            in [->] (intersection-2) -- +(\n2:0.1pt);
            % SECOND ARROW
            % Step 1: Draw arrow body
            \begin{scope}
                \clip (\tikzcdmatrixname-1-2.west) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-1.north) -- (\tikzcdmatrixname-1-1.center) -- cycle;
                \path[draw, line width=rule_thickness] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=180,radius=#1];
            
```

```

        \end{scope}
    % Step 2: Draw arrow head
    % Step 2.1: Find the point at which to place the arrowhead
    \path[name path=curve-2-a] (\tikzcdmatrixname-1-2.west) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-1.north) -- (\tikzcdmatrixname-1-1.center) -- cycle;
    \path[name path=curve-2-b] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=180, radius=1pt]
    \fill [name intersections={of=curve-2-a and curve-2-b}] (intersection-2);
    % Step 2.2: Find the angle at which to place the arrowhead
    \coordinate (arc-start) at (\tikzcdmatrixname-1-2.west);
    \coordinate (arc-center) at (\tikzcdmatrixname-2-2.center);
    \draw let
        \p1 = {$(\text{intersection-2}) - (\text{arc-center})$}, % \p1 is the vector from the arc
        \n1 = {atan2(\y1, \x1)}, % \n1 is the angle of that vector in degrees
        \n2 = {\n1 - 90} % \n2 is the angle of the tangent (90 degrees from the radius)
    in [<-] (intersection-2) -- ++(\n2:0.1pt);
    % Labels
    \path (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=180, radius=1pt] node[above]
    \path (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=0, radius=1pt] node[below]
}
}
}

```


The code

```

\begin{tikzcd}[row sep={4.5*\the\DL,between origins}, column sep={4.5*\the\DL,between origins}]
    {}% Don't remove this line, it's important!
    \&
    P
    \arrow[d,"\phi"\{pos=0.475}, "\exists!"\{pos=0.475}, dashed]
    \&
    {}% Don't remove this line, it's important!
    \\
    A
    \&
    A\times B
    \arrow[l,"\text{pr}_{\{1\}}"\{pos=0.425}, two heads]
    \arrow[r,"\text{pr}_{\{2\}}"\{pos=0.425}, two heads]
    \&
    B
\end{tikzcd}

```

will then produce the following diagram:



### 02C2 15.1.2 Coproduct Diagram With Circular Arrows

Define

```
\newlength{\DL}
\setlength{\DL}{0.9em}
```

in the preamble, as well as

```
\tikzcdset{
    coproductArrows/.style args={#1#2#3}={
        execute at end picture={
            % FIRST ARROW
            % Step 1: Draw arrow body
            \begin{scope}
                \clip (\tikzcdmatrixname-1-2.east) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-3.north) -- (\tikzcdmatrixname-1-3.center) -- cycle;
                \path[draw, line width=rule_thickness] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=0,radius=#1];
            \end{scope}
            % Step 2: Draw arrow head
            % Step 2.1: Find the point at which to place the arrowhead
            \path[name path=curve-1-a] (\tikzcdmatrixname-1-2.east) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-3.north) -- (\tikzcdmatrixname-1-3.center) -- cycle;
            \path[name path=curve-1-b] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=0,radius=#1];
            \fill [name intersections={of=curve-1-a and curve-1-b}] (intersection-1);
            % Step 2.2: Find the angle at which to place the arrowhead
            \coordinate (arc-start) at (\tikzcdmatrixname-1-2.east);
            \coordinate (arc-center) at (\tikzcdmatrixname-2-2.center);
            \draw let
                \p1 = {($(\tikzcdmatrixname-1-2) - (\tikzcdmatrixname-2-2)$)}, % \p1 is the vector from the arc
                \n1 = {atan2(\y1, \x1)}, % \n1 is the angle of that vector in degrees
                \n2 = {\n1 - 90} % \n2 is the angle of the tangent (90 degrees from the rad
            in
```

```

    in [<-] (intersection-1) -- ++(\n2:0.1pt);
% SECOND ARROW
% Step 1: Draw arrow body
\begin{scope}
    \clip (\tikzcdmatrixname-1-2.west) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-1.north) -- (\tikzcdmatrixname-1-1.center) -- cycle;
    \path[draw, line width=rule_thickness] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=180,radius=#1];
\end{scope}
% Step 2: Draw arrow head
% Step 2.1: Find the point at which to place the arrowhead
\path[name path=curve-2-a] (\tikzcdmatrixname-1-2.west) -- (\tikzcdmatrixname-2-2.center) -- (\tikzcdmatrixname-2-1.north) -- (\tikzcdmatrixname-1-1.center) -- cycle;
\path[name path=curve-2-b] (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=\n1]
\fill [name intersections={of=curve-2-a and curve-2-b}] (intersection-1);
% Step 2.2: Find the angle at which to place the arrowhead
\coordinate (arc-start) at (\tikzcdmatrixname-1-2.west);
\coordinate (arc-center) at (\tikzcdmatrixname-2-2.center);
\draw let
    \p1 = {$(\text{intersection-1}) - (arc-center)$}, % \p1 is the vector from the arc
2 for the 2nd intersection)
    \n1 = {atan2(\y1, \x1)}, % \n1 is the angle of that vector in degrees
    \n2 = {\n1 - 90} % \n2 is the angle of the tangent (90 degrees from the radius)
in [->] (intersection-1) -- ++(\n2:0.1pt);
% Labels
\path (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=180,radius=#1] node[above]
\path (\tikzcdmatrixname-1-2) arc[start angle=90,end angle=0,radius=#1] node[below]
}
}
}

```

The code

```

\begin{tikzcd}[row sep={4.5*\the\DL,between origins}, column sep={4.5*\the\DL,between origins}]
    {}% Don't remove this line, it's important!
    \&
    C
    \arrow[from=d,"\phi","\exists!", dashed]
    \&
    {}% Don't remove this line, it's important!
    \\
    A


```

```

\&
A \icoprod B
\arrow[from=l, "\inj_1", hook]
\arrow[from=r, "\inj_2", hook']
\&
B
\end{tikzcd}

```

will then produce the following diagram:



## 01VS 15.1.3 Cube Diagram

## Define

```
\newlength{\DL}  
\setlength{\DL}{0.9em}
```

### The code


```
\begin{tikzcd}[row sep={4.0*\the\DL,between origins}, column sep={4.0*\the\DL,between o  
1  
\&  
\&  
2  
\&  
\&  
1'  
\&  
\&  
2'  
\&  
3  
\&  
\&  
4  
\&  
\&  
\&
```

```

3'
\&
\&
4'
% 1-Arrows
% First Square
\arrow[from=1-1,to=3-1,"f'"]%
\arrow[from=3-1,to=3-3,"h"\{description,pos=0.25\}]%
\arrow[from=1-1,to=1-3,"g"]%
\arrow[from=1-3,to=3-3,"k"\{description,pos=0.25\}]%
% Second Square
\arrow[from=2-2,to=4-2,"f'"\{description,pos=0.3\},crossing over]%
\arrow[from=4-2,to=4-4,"h'']\%
\arrow[from=2-2,to=2-4,"g'"\{description,pos=0.3\},crossing over]%
\arrow[from=2-4,to=4-4,"k'"]%
% Connecting Arrows
\arrow[from=1-1,to=2-2,"c_{\{1\}}"\{description\}]%
\arrow[from=1-3,to=2-4,"c_{\{2\}}"]\%
\arrow[from=3-1,to=4-2,"c_{\{3\}}'"]\%
\arrow[from=3-3,to=4-4,"c_{\{4\}}"\{description\}]%
\end{tikzcd}

```

will produce the following diagram:



#### 01VT 15.1.4 Cube Diagram With Labelled Faces

Define

```
\newlength{\DL}
\setlength{\DL}{0.9em}
```

The code

```

\begin{tikzcd}[row sep={4.0*\the\DL,between origins}, column sep={4.0*\the\DL,between o
  1
  \&
  \&
  2
  \&
  \\
  \&
  1'
  \&
  \&
  2'
  \\
  3
  \&
  \&
  \&
  \\
  \&
  3'
  \&
  \&
  4'
% 1-Arrows
% First Square
\arrow[from=1-1,to=3-1,"f'"]
\arrow[from=1-1,to=1-3,"g"]
% Second Square
\arrow[from=2-2,to=4-2,"f'{description},crossing over"]
\arrow[from=4-2,to=4-4,"h'"]
\arrow[from=2-2,to=2-4,"g'{description},crossing over"]
\arrow[from=2-4,to=4-4,"k'"]
% Connecting Arrows
\arrow[from=1-1,to=2-2,"c_{1}"description]
\arrow[from=1-3,to=2-4,"c_{2}"]
\arrow[from=3-1,to=4-2,"c_{3}"]
% Subdiagrams
\arrow[from=2-2,to=1-3,"{\scriptstyle(1)}"\{rotate=-0.3,xslant=-0.903569337,yslant=0,xscale=7.0341,yscale=4.4454,xscale=0.225,yscale=0.225},phantom]
\arrow[from=3-1,to=2-2,"{\scriptstyle(2)}"\{rotate=-44.6,xslant=-0.965688775,yslant=0,xscale=8.6931,yscale=8.2852,xscale=0.15,yscale=0.15},phantom]
\arrow[from=4-2,to=2-4,"{\scriptstyle(3)}"\{rotate=0,xslant=0,yslant=0,xscale=1.5,yscale=1.5},phantom]
\end{tikzcd}
\qqquad


```

```

\begin{tikzcd}[row sep={4.0*\the\DL,between origins}, column sep={4.0*\the\DL,between o
  1
  \&
  \&
  2
  \&
  \\
  \&
  \&
  \&
  \&
  2'
  \\
  3
  \&
  \&
  4
  \&
  \\
  \&
  \&
  3'
  \&
  \&
  4'
% 1-Arrows
% First Square
\arrow[from=1-1,to=3-1,"f'"]%
\arrow[from=3-1,to=3-3,"h"\{description}\}]%
\arrow[from=1-1,to=1-3,"g"]%
\arrow[from=1-3,to=3-3,"k"\{description}\}]%
% Second Square
\arrow[from=4-2,to=4-4,"h'']%
\arrow[from=2-4,to=4-4,"k'"]%
% Connecting Arrows
\arrow[from=1-3,to=2-4,"c_{2}"]%
\arrow[from=3-1,to=4-2,"c_{3}"]%
\arrow[from=3-3,to=4-4,"c_{4}"description]\}
% Subdiagrams
\arrow[from=1-1,to=3-3,"{\scriptstyle(4)}"\{rotate=0,xslant=0,yslant=0,xscale=1.5,yscale=1.5,y
\arrow[from=3-3,to=2-4,"{\scriptstyle(5)}"\{rotate=-44.6,xslant=-0.965688775,yslant=0,xscale=8.6931,yscale=8.2852,xscale=0.15,yscale=0.15},phantom]%
\arrow[from=4-2,to=3-3,"{\scriptstyle(6)}"\{rotate=-0.3,xslant=-0.903569337,yslant=0,xscale=7.0341,yscale=4.4454,xscale=0.225,yscale=0.225},phantom]%
\end{tikzcd}

```

will produce the following diagram:



### 01VU 15.1.5 Pentagon Diagram

Define


```
\newlength{\ThreeCm}
\setlength{\ThreeCm}{3.0cm}
```

The code

```
\begin{tikzcd}[row sep={0*\the\DL,between origins}, column sep={0*\the\DL,between origins}
    &[0.30901699437\ThreeCm]
    &[0.5\ThreeCm]
    A\otimes_{\{R\}}(A\otimes_{\{R\}}A)
    &[0.5\ThreeCm]
    \&[0.30901699437\ThreeCm]
    \\\&[0.58778525229\ThreeCm]
    (A\otimes_{\{R\}}A)\otimes_{\{R\}}A
    &[0.30901699437\ThreeCm]
    \&[0.5\ThreeCm]
    &[0.5\ThreeCm]
    \&[0.30901699437\ThreeCm]
    A\otimes_{\{R\}}A
    \\\&[0.95105651629\ThreeCm]
    \&[0.30901699437\ThreeCm]
    A\otimes_{\{R\}}A
    \&[0.5\ThreeCm]
    \&[0.5\ThreeCm]
    A
    \&[0.30901699437\ThreeCm]
    % 1-Arrows
    % Left Boundary
```

```
\arrow[from=2-1,to=1-3,"\alpha^{\mathrm{Mod}_R}_{A,A,A}]{pos=0.4125}%
\arrow[from=1-3,to=2-5,"\mathrm{id}_A \otimes_R \mu^{\times}{\times}_A]{pos=0.6}%
\arrow[from=2-5,to=3-4,"{\mu}^{\times}{\times}_A]{pos=0.425}%
% Right Boundary
\arrow[from=2-1,to=3-2,"{\mu}^{\times}{\times}_A \otimes_R \mathrm{id}_A"!]{pos=0.425}%
\arrow[from=3-2,to=3-4,"{\mu}^{\times}{\times}_A"!]%
\end{tikzcd}
```

will produce the following pentagon diagram:



To make the diagram larger, one could use e.g.

```
\newlength{\FourCm}
\setlength{\FourCm}{2.0cm}
```

and replace all instances of `\ThreeCm` with `\FourCm` in the code above.

### 01VV 15.1.6 Hexagon Diagram

Define

```
\newlength{\OneCmPlusHalf}
\setlength{\OneCmPlusHalf}{1.5cm}
```


The code

```
\begin{tikzcd}[row sep={0.0*\the\DL,between origins}, column sep={0.0*\the\DL,between origins}]
&[0.86602540378\OneCmPlusHalf]
1 &[0.86602540378\OneCmPlusHalf]
&\backslash[0.5\OneCmPlusHalf]
2 &[0.86602540378\OneCmPlusHalf]
&\&[0.86602540378\OneCmPlusHalf]
3

```

```
\\"[\OneCmPlusHalf]
4
\&[0.86602540378\OneCmPlusHalf]
\&[0.86602540378\OneCmPlusHalf]
5
\\"[0.5\OneCmPlusHalf]
\&[0.86602540378\OneCmPlusHalf]
6
\&[0.86602540378\OneCmPlusHalf]
% 1-Arrows
% Left Boundary
\arrow[from=1-2,to=2-1,"L_{1}"]%
\arrow[from=2-1,to=3-1,"L_{2}"']%
\arrow[from=3-1,to=4-2,"L_{3}"]%
% Right Boundary
\arrow[from=1-2,to=2-3,"R_{1}"]%
\arrow[from=2-3,to=3-3,"R_{2}"]%
\arrow[from=3-3,to=4-2,"R_{3}"]%
\end{tikzcd}
```

will produce the following hexagon diagram:



To make the diagram larger, one could use e.g.

```
\newlength{\TwoCm}
\setlength{\TwoCm}{2.0cm}
```

and replace all instances of \OneCmPlusHalf with \TwoCm in the code above.

### 02C3 15.1.7 Double Square Diagram

Define

```
\newlength{\DL}
\setlength{\DL}{0.9cm}
```

The code


```

\begin{tikzcd}[row sep={10.0*\the\DL,between origins}, column sep={10.0*\the\DL,between
\bullet
\&
\&
\&
\bullet
\\
\&
\&
\bullet
\&
\&
\bullet
\&
\&
\&
\&
\bullet
\&
\&
\bullet
\&
\&
\&
\&
\bullet
\&
\&
\&
\&
\&
\bullet
% Arrows
% Outer Square
\arrow[from=1-1,to=1-4]%
\arrow[from=1-4,to=4-4]%
%
\arrow[from=1-1,to=4-1]%
\arrow[from=4-1,to=4-4]%
% Inner Square
\arrow[from=2-2,to=2-3]%
\arrow[from=2-3,to=3-3]%
%
\arrow[from=2-2,to=3-2]%
\arrow[from=3-2,to=3-3]%
% Connecting Arrows
\arrow[from=1-1,to=2-2]%
\arrow[from=1-4,to=2-3]%
\arrow[from=3-2,to=4-1]%
\arrow[from=3-3,to=4-4]%
% Subdiagrams
\arrow[from=2-2,to=3-3,"{\scriptstyle(1)}",phantom,yshift=10.0*\the\DL]%

```

```
\arrow[from=2-2,to=3-2,"\scriptstyle(2)",phantom,xshift=-5.0*\the\DL]%
\arrow[from=2-2,to=3-3,"\scriptstyle(3)",phantom]%
\arrow[from=2-3,to=3-3,"\scriptstyle(4)",phantom,xshift=5.0*\the\DL]%
\arrow[from=2-2,to=3-3,"\scriptstyle(5)",phantom,yshift=-10.0*\the\DL]%
\end{tikzcd}
```

will produce the following double square diagram:



### 01WH 15.1.8 Double Hexagon Diagram

Define

```
\newlength{\OneCm}
\setlength{\OneCm}{1.0cm}
```


The code

```
\begin{tikzcd}[row sep={0.0*\the\DL,between origins}, column sep={0.0*\the\DL,between o
  \&[1.73205081*\OneCm]
  \&[1.73205081*\OneCm]
  \text{1-3}
  \&[1.73205081*\OneCm]
  \&[1.73205081*\OneCm]
  \\\text{2-1}
  \&[1.73205081*\OneCm]
  \&[1.73205081*\OneCm]
  \text{2-3}
  \&[1.73205081*\OneCm]
  \&[1.73205081*\OneCm]
  \text{2-5}
  \\\text{1-5}
```

```
\&[1.73205081*\OneCm]
\text{3-2}
\&[1.73205081*\OneCm]
\&[1.73205081*\OneCm]
\text{3-4}
\&[1.73205081*\OneCm]
\\[2.0*\OneCm]
\&[1.73205081*\OneCm]
\text{4-2}
\&[1.73205081*\OneCm]
\&[1.73205081*\OneCm]
\text{4-4}
\&[1.73205081*\OneCm]
\\[1.0*\OneCm]
\text{5-1}
\&[1.73205081*\OneCm]
\&[1.73205081*\OneCm]
\text{5-3}
\&[1.73205081*\OneCm]
\&[1.73205081*\OneCm]
\text{5-5}
\\[2.0*\OneCm]
\&[1.73205081*\OneCm]
\&[1.73205081*\OneCm]
\text{6-3}
\&[1.73205081*\OneCm]
\&[1.73205081*\OneCm]
% Arrows
\arrow[from=1-3,to=2-1,"1'"]%
\arrow[from=2-1,to=5-1,"2'"]%
\arrow[from=5-1,to=6-3,"3'"]%
%
\arrow[from=1-3,to=2-5,"4"]%
\arrow[from=2-5,to=5-5,"5"]%
\arrow[from=5-5,to=6-3,"6"]%
%
\arrow[from=2-3,to=3-2,"1"description]%
\arrow[from=3-2,to=4-2,"2"description]%
\arrow[from=4-2,to=5-3,"3"description]%
%
\arrow[from=2-3,to=3-4,"4"description]%
\arrow[from=3-4,to=4-4,"5"description]%
\arrow[from=4-4,to=5-3,"6"description]%
%
```

```
\arrow[from=1-3,to=2-3,"a"description]%
\arrow[from=2-1,to=3-2,"b"description]%
\arrow[from=2-5,to=3-4,"c"description]%
\arrow[from=5-1,to=4-2,"d"description]%
\arrow[from=5-5,to=4-4,"e"description]%
\arrow[from=5-3,to=6-3,"f"description]%
\end{tikzcd}
```

will produce the following double hexagon diagram:



To make the diagram larger, one could use e.g.

```
\newlength{\TwoCm}
\setlength{\TwoCm}{2.0cm}
```

and replace all instances of \OneCm with \TwoCm in the code above.

## 01VC 15.2 Retired Tags

### 01VD 15.2.1 Relations

01VE OLD TAG 15.2.1.1.1 ► EQUIVALENT DEFINITIONS OF RELATIONS

The content of this tag has been moved to [Definition 8.1.1.1.1](#).

00R1	<b>OLD TAG 15.2.1.1.2 ► INTERACTION BETWEEN COMPOSITION AND CHARACTERISTIC RELATIONS</b>
	The original statement of this tag was false.
00QQ	<b>OLD TAG 15.2.1.1.3 ► INTERACTION BETWEEN COMPOSITION AND CHARACTERISTIC RELATIONS</b>
	The original statement of this tag was false.
00NN	<b>OLD TAG 15.2.1.1.4 ► EXPLICIT DESCRIPTION OF INTERNAL LEFT KAN EXTENSIONS ALONG FUNCTIONS</b>
	This was a question. Now an explicit description is available as ??.
00NU	<b>OLD TAG 15.2.1.1.5 ► EXPLICIT DESCRIPTION OF INTERNAL LEFT KAN LIFTS ALONG FUNCTIONS</b>
	This was a question. Now an explicit description is available as ??.
00MG	<b>OLD TAG 15.2.1.1.6 ► INTERNAL KAN EXTENSIONS AND LIFTS</b>
	This tag is obsolete; see <a href="#">Sections 8.5.13</a> to <a href="#">8.5.16</a> instead.
00MH	<b>OLD TAG 15.2.1.1.7 ► INTERNAL KAN EXTENSIONS AND LIFTS</b>
	This tag is obsolete; see <a href="#">Sections 8.5.13</a> to <a href="#">8.5.16</a> instead.
00NG	<b>OLD TAG 15.2.1.1.8 ► INTERNAL KAN EXTENSIONS AND LIFTS</b>
	This tag is obsolete; see <a href="#">Sections 8.5.13</a> to <a href="#">8.5.16</a> instead.
00LV	<b>OLD TAG 15.2.1.1.9 ► BETTER CHARACTERISATIONS OF REPRESENTABLY FULL MORPHISMS IN Rel</b>
	This was originally a question. It has been answered in ??.
00MD	<b>OLD TAG 15.2.1.1.10 ► BETTER CHARACTERISATIONS OF COREPRESENTABLY FULL MORPHISMS IN Rel</b>
	This was originally a question. It has been answered in <a href="#">Section 8.5.11</a> .

**00LE** OLD TAG 15.2.1.1.11 ► CHARACTERISATION OF MONOMORPHISMS IN Rel

Superseded by ??.

**00LJ** OLD TAG 15.2.1.1.12 ► CHARACTERISATION OF 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LK** OLD TAG 15.2.1.1.13 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LL** OLD TAG 15.2.1.1.14 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LM** OLD TAG 15.2.1.1.15 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LN** OLD TAG 15.2.1.1.16 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LP** OLD TAG 15.2.1.1.17 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LQ** OLD TAG 15.2.1.1.18 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LR** OLD TAG 15.2.1.1.19 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LS** OLD TAG 15.2.1.1.20 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

**00LT** OLD TAG 15.2.1.1.21 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

00LU OLD TAG 15.2.1.1.22 ► 2-CATEGORICAL MONOMORPHISMS IN Rel

Superseded by ??.

00LX OLD TAG 15.2.1.1.23 ► CHARACTERISATION OF EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

02SD OLD TAG 15.2.1.1.24 ► CHARACTERISATION OF EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M2 OLD TAG 15.2.1.1.25 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M3 OLD TAG 15.2.1.1.26 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M4 OLD TAG 15.2.1.1.27 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M5 OLD TAG 15.2.1.1.28 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M6 OLD TAG 15.2.1.1.29 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M7 OLD TAG 15.2.1.1.30 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M8 OLD TAG 15.2.1.1.31 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

00M9 OLD TAG 15.2.1.1.32 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11.](#)

**00MA** OLD TAG 15.2.1.1.33 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11](#).

**00MB** OLD TAG 15.2.1.1.34 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11](#).

**00MC** OLD TAG 15.2.1.1.35 ► 2-CATEGORICAL EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11](#).

**029S** OLD TAG 15.2.1.1.36 ► EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11](#).

**029T** OLD TAG 15.2.1.1.37 ► EPIMORPHISMS IN Rel

Superseded by [Section 8.5.11](#).

## 01VF 15.2.2 Pointed Sets

**009F** OLD TAG 15.2.2.1.1 ► THE UNDERLYING POINTED SET OF A SEMIMODULE

The **underlying pointed set** of a semimodule  $(M, \alpha_M)$  is the pointed set  $(M, 0_M)$ .

**009G** OLD TAG 15.2.2.1.2 ► THE UNDERLYING POINTED SET OF A MODULE

The **underlying pointed set** of a module  $(M, \alpha_M)$  is the pointed set  $(M, 0_M)$ .

## 01WG 15.2.3 Tensor Products of Pointed Sets

**00H2** OLD TAG 15.2.3.1.1 ► SECTION ON UNIVERSAL PROPERTIES OF THE SMASH PRODUCT OF POINTED SETS I

Absorbed into [Section 7.5.10](#).

**00H4** OLD TAG 15.2.3.1.2 ► SECTION ON UNIVERSAL PROPERTIES OF THE SMASH PRODUCT OF POINTED SETS II

Absorbed into [Section 7.5.10](#).

	OLD TAG 15.2.3.1.3 ► UNIVERSAL PROPERTIES OF THE SMASH PRODUCT OF POINTED SETS
00H3	I
	OLD TAG 15.2.3.1.4 ► UNIVERSAL PROPERTIES OF THE SMASH PRODUCT OF POINTED SETS
00H5	II

Absorbed into [Section 7.5.10](#).

Absorbed into [Section 7.5.10](#).

#### 01VJ 15.2.4 Categories

017R OLD TAG 15.2.4.1.1 ► PICTURING NATURAL TRANSFORMATIONS IN DIAGRAMS

We denote natural transformations in diagrams as

$$\begin{array}{ccc} C & \begin{array}{c} \xrightarrow{F} \\ \Downarrow \alpha \\ \xrightarrow{G} \end{array} & \mathcal{D}. \end{array}$$

(This tag has been removed and is now part of [Remark 11.9.2.1.2](#).)

01VL OLD TAG 15.2.4.1.2 ► INTERACTION BETWEEN FULLNESS AND POSTCOMPOSITION FUNCTORS

(This Tag was an item of [Proposition 11.6.2.1.2](#), but has since been removed because its statement is incorrect. Naïm Camille Favier provided a counterexample, and the corrected statements now appear as [Items 2 and 3 of Proposition 11.6.2.1.2](#).)

01VM 1. *Interaction With Postcomposition.* The following conditions are equivalent:

01VN (a) The functor  $F: C \rightarrow \mathcal{D}$  is full.

01VP (b) For each  $X \in \text{Obj}(\text{Cats})$ , the postcomposition functor

$$F_*: \text{Fun}(X, C) \rightarrow \text{Fun}(X, \mathcal{D})$$

is full.

01VQ (c) The functor  $F: C \rightarrow \mathcal{D}$  is a representably full morphism in  $\text{Cats}_2$  in the sense of [Definition 14.1.2.1.1](#).

## 02C4 15.3 Miscellany

### 01CV 15.3.1 List of Things To Explore/Add

Here we list things to be explored in or added to this work in the future. This is a very quick and dirty list; some items may not be fully intelligible.

01CW

#### REMARK 15.3.1.1 ► THINGS TO EXPLORE/ADD

Set Theory:

1. <https://math.stackexchange.com/questions/200389/show-that-the-set-of-all-finite-subsets-of-mathbbn-is-countable>
2. <https://mathoverflow.net/a/479528>
3. <https://www.maths.ed.ac.uk/~tl/ast/ast.pdf>

Type Theory:

1. <https://mathoverflow.net/questions/497570/universes-don-t-need-to-be-indexed-by-natural-numbers>

Pointed sets:

1. Universal properties (plural!) of the left tensor product of pointed sets
2. Universal properties (plural!) of the right tensor product of pointed sets

Relations:

1. Internal fibrations in **Rel**, like discrete fibrations and Street fibrations
2. Return to Eilenberg–Moore and Kleisli objects in **Rel** once the general theory has been set up for internal monads

Spans:

1. <https://arxiv.org/abs/2505.22832>
2. Spans: study certain compositions of spans like composing  $B \xleftarrow{f} A = A = A \xleftarrow{g} B$  into a span  $B \xleftarrow{f} A \xleftarrow{g} B$
3. Comparison *double functor* from Span to Rel and vice versa

4. Apartness composition for spans and alternate compositions for spans in general
5. non-Cartesian analogue of spans
  - (a) View spans as morphisms  $S \rightarrow A \times B$  and consider instead morphisms  $S \rightarrow A \otimes_C B$
6. Record the universal property of the bicategory of spans of <https://ncatlab.org/nlab/show/span>
7. <https://ncatlab.org/nlab/show/span+trace>
8. Cospans.
9. Multispans.

Un/Straightening for Indexed and Fibred Sets:

1. Analogue of adjoints for Grothendieck construction for indexed and fibred sets
2. Write proper sections on straightening for lax functors from Sets to Rel or Span (displayed sets)
3. co/units for un/straightening adjunction

Categories:

1. <https://www.numdam.org/actas/SE/>, <https://www.numdam.org/journals/CTGDC/>
2. [https://www.numdam.org/item/CTGDC\\_1966\\_\\_8\\_\\_A5\\_0.pdf](https://www.numdam.org/item/CTGDC_1966__8__A5_0.pdf)
3. <https://mathoverflow.net/questions/493931/is-the-category-of-posets-locally-cartesian-closed>
4. From Keith: Presheaves on a topological space  $X$  valued in  $\{t, f\}$ 
  - (a) They are the same as collections of open subsets of  $X$
  - (b) They are sheaves iff that collection is closed under union
  - (c) Their sheafification is the closure of that collection under unions
5. <https://arxiv.org/abs/2504.20949>
6. Notion of equality that is weaker than equivalence but stronger than adjunction

7. Tangent categories, Beck modules, categorical derivations
8. Flat functors
9. Is the classifying space of a category isomorphic to  $\text{Ex}^\infty$  of the nerve of the category? If so, an intuition for having an initial/terminal object implying being homotopically contractible is that taking the free  $\infty$ -groupoid generated by that identifies every object with the terminal one.
10. [https://en.wikipedia.org/wiki/Category\\_algebra](https://en.wikipedia.org/wiki/Category_algebra)
11. simple objects
12. <https://mathoverflow.net/questions/442212/properties-of-categorical-zeta-function>
13. Polynomial functors, <https://ncatlab.org/nlab/show/polynomial+functor>, <https://arxiv.org/abs/2312.00990>
14. <https://ncatlab.org/nlab/show/simple+object>
15. <https://mathoverflow.net/questions/442212/properties-of-categorical-zeta-function>
16. <https://arxiv.org/abs/2409.17489>
17. <https://mathoverflow.net/a/478644>
18. Posetal category associated to a poset as a right adjoint
19. “Presetal category” associated to a preordered set
20. Vopenka’s principle simplifies stuff in the theory of locally presentable categories. If we build categories using type theory or HoTT, what stuff from vopenka holds?
21. Are pseudoepic functors those functors whose restricted Yoneda embedding is pseudomonadic and Yoneda preserves absolute colimits?
22. Absolutely dense functors enriched over  $\mathbb{R}^+$  apparently reduce to topological density
23. Is there a reasonable notion of category homology? It is very common for the geometric realisation of a category to be contractible (e.g. having an initial or terminal object), but maybe some notion of directed homology could work here

24. Nerves of categories:

- (a) Dihedral and symmetric nerves of categories via groupoids  
(define them first for groupoids and then Kan extend along  
 $\text{Grpd} \hookrightarrow \text{Cats}$ )
  - i. Same applies to twisted nerves
- (b) Cyclic nerve of a category
- (c) Crossed Simplicial Group Categorical Nerves, <https://arxiv.org/abs/1603.08768>

25. Define contractible categories and add a discussion of universal properties as stating that certain categories are contractible. (Example of non-unique isomorphisms as e.g. being a group of order 5 corresponds to all objects being isomorphic but the category not being contractible)

26. Expand [Construction 11.4.3.1.2](#) and add a proof to it.

27. Sections and retractions; retracts, <https://ncatlab.org/nlab/show/retract>.

28. Groupoid cardinality

- (a) <https://mathoverflow.net/questions/376175/category-theory-and-arithmetical-identities/376223#376223>
- (b) <https://mathoverflow.net/questions/420088/groupoid-cardinality-of-the-class-of-abelian-p-groups?rq=1>
- (c) <https://mathoverflow.net/questions/363292/what-is-the-groupoid-cardinality-of-the-category-of-vector-spaces-over-a-finite>
- (d) The groupoid cardinality of the core of the category of finite sets is  $e$ . What is the groupoid cardinality of the core of  $\text{FinSets}_G$ ?
- (e) groupoid cardinality of the core of the category of finite G-sets, <https://www.arxiv.org/pdf/2502.03585>
- (f) <https://ncatlab.org/nlab/show/groupoid+cardinality>
- (g) <https://arxiv.org/abs/2104.11399>
- (h) <https://terrytao.wordpress.com/2017/04/13/counting-objects-up-to-isomorphism-groupoid-cardinality/>
- (i) <https://arxiv.org/abs/0809.2130>
- (j) <https://qchu.wordpress.com/2012/11/08/groupoid-cardinality/>

(k) <https://mathoverflow.net/questions/363292/what-is-the-groupoid-cardinality-of-the-category-of-vector-spaces-over-a-finite>

29. combinatorial species

- (a) <https://ncatlab.org/nlab/show/Schur+functor>
  - i. Equivalence between twisted commutative algebras and algebras on categories of polynomial functors, <https://mathweb.ucsd.edu/~ssam/talks/2014/ihp-tca.pdf>
- (b) <https://mathoverflow.net/questions/22462/what-are-some-examples-of-interesting-uses-of-the-theory-of-combinatorial-species>
- (c) [https://en.wikipedia.org/wiki/Combinatorial\\_species](https://en.wikipedia.org/wiki/Combinatorial_species)

30. Leinster's the eventual image, <https://arxiv.org/abs/2210.00302>

- (a) Telescope notation  $\text{tel}_\phi(X) \stackrel{\text{def}}{=} \text{colim}(X \xrightarrow{\phi} X \xrightarrow{\phi} \cdots)$  introduced in <https://arxiv.org/abs/2505.06979>

31. <https://ncatlab.org/nlab/show/separable+functor>

32. Dagger categories:

- (a) [https://en.wikipedia.org/wiki/Dagger\\_category](https://en.wikipedia.org/wiki/Dagger_category)
- (b) <https://ncatlab.org/nlab/show/dagger+category>
- (c) Dagger compact categories, [https://en.wikipedia.org/wiki/Dagger\\_compact\\_category](https://en.wikipedia.org/wiki/Dagger_compact_category)
- (d) <https://mathoverflow.net/questions/220032/are-dagger-categories-truly-evil>
- (e) generalisation of dagger categories to categories with duality, i.e. categories  $C$  together with a functor  $\dagger: C^{\text{op}} \rightarrow C$ 
  - i. Perhaps with the additional condition that  $\dagger \circ \dagger = \text{id}$
  - ii. categories with involutions in general

Regular Categories:

1. <https://arxiv.org/pdf/2004.08964.pdf>.
2. Internal relations

Types of Morphisms in Categories:

1. <https://mathoverflow.net/questions/490476/duality-of-f-injectivity-surjectivity-of-precomposition-map> for motivation of monomorphisms/epimorphisms
2. Characterisation of epimorphisms in the category of fields, <https://math.stackexchange.com/q/4941660>
3. Strong epimorphisms
4. Behaviour in  $\text{Fun}(C, \mathcal{D})$ , e.g. pointwise sections vs. sections in  $\text{Fun}(C, \mathcal{D})$ .
5. Faithful functors from balanced categories are conservative
6. Natural cotransformations:
  - (a) If there is a natural transformation between functors between categories, taking nerves gives a homotopy equivalence (or something like that). What happens for natural cotransformations?
  - (b) Natural transformations come with a vertical composition map

$$\circ : \coprod_{G \in \text{Fun}(C, \mathcal{D})} \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H).$$

As Morgan Rogers shows [here](#), there's no vertical cocomposition map of the form

$$\text{CoNat}(F, H) \rightarrow \prod_{G \in \text{Fun}(C, \mathcal{D})} \text{CoNat}(G, H) \times \text{CoNat}(F, G)$$

or of the form

$$\text{CoNat}(F, H) \rightarrow \prod_{G \in \text{Fun}(C, \mathcal{D})} \text{CoNat}(G, H) \coprod \text{CoNat}(F, G)$$

for natural cotransformations.

- (c) Cap product for CoNat and Nat
  - i. recovers map  $Z(G) \times Cl(G) \rightarrow Cl(G)$ .
- (d) What is the geometric realisation of  $\text{CoTrans}(F, G)$ ?
  - i. Related: <https://mathoverflow.net/questions/89753/geometric-realization-of-hochschild-complex>
- (e) What is the totalisation of  $\text{Trans}(F, G)$ ?

- i. If we view sets as discrete topological spaces, what are the homotopy/homology groups of it? The nLab says this (<https://ncatlab.org/nlab/show/totalization>):

The homotopy groups of the totalization of a cosimplicial space are computed by a Bousfield-Kan spectral sequence.

The homology groups by an Eilenberg-Moore spectral sequence.

(f) Abstract

Adjunctions:

1. Relative adjunctions: message Alyssa asking for her notes
2. Adjunctions, units, counits, and fully faithfulness as in <https://mathoverflow.net/questions/100808/properties-of-functors-and-their-adjoints>.
3. Morphisms between adjunctions and bicategory  $\text{Adj}(C)$ .
4. <https://ncatlab.org/nlab/show/transformation+of+adjoints>

Presheaves and the Yoneda Lemma:

1. <https://mathoverflow.net/questions/498069/products-and-coproducts-in-the-category-of-elements-of-a-presheaf>
2. Yoneda extension along  $\mathcal{Y}_{\mathcal{D}} \circ F: C \rightarrow \text{PSh}(\mathcal{D})$ , giving a functor left adjoint to the precomposition functor  $F^*: \text{PSh}(\mathcal{D}) \rightarrow \text{PSh}(C)$ .
3. Consider the diagram

$$\begin{array}{ccccc} & & \text{PSh}(C) & & \\ & \nearrow & \downarrow & \searrow & \\ C & \longrightarrow & \mathcal{D} & \hookrightarrow & \text{PSh}(\mathcal{D}) \end{array}$$

4. Does the functor tensor product admit a right adjoint (“Hom”) in some sense?
5. Yoneda embedding preserves limits
6. universal objects and universal elements
7. adjoints to the Yoneda embedding and total categories

8. The co-Yoneda lemma: co/presheaves are colimits of co/representables
9. Properties of categories of copresheaves
10. Contravariant restricted Yoneda embedding
11. Contravariant Yoneda extensions
12. Make table of  $\text{Lift}_\circ(\circ)$ ,  $\text{Ran}_\circ(\circ)$ ,  $\text{Ran}_\circ(\circ)$ , etc.
13. Properties of restricted Yoneda embedding, e.g. if the restricted Yoneda embedding is full, then what can we conclude? Related: <https://qchu.wordpress.com/2015/05/17/generators/>
14. Tensor product of functors and relation to profunctors
15. rifts and rans and lifts and lans involving yoneda in Cats and Prof
16. Tensor product of functors and relation to rifts and rans of profunctors

Isbell Duality:

1. enriched Isbell over walking chain complex
2. Isbell self-dual presheaves for Lawvere metric spaces; when

$$f(x) = \sup_{x \in X} \left( \left| f(x) - \sup_{y \in X} (|f(y) - d_X(y, x)|) \right| \right)$$

holds.

3. <https://ncatlab.org/nlab/show/Fr%C3%B6licher+spaces+and+Isbell+envelopes>
4. <https://ncatlab.org/nlab/show/envelope+of+an+adjunction>
5. <https://ncatlab.org/nlab/show/nucleus+of+a+profunctor>
6. <https://ncatlab.org/nlab/show/nuclear+adjunction>
7. <https://ncatlab.org/nlab/show/fixed+point+of+an+adjunction>
8. **Important:** I should reconsider going with the notation O and Spec. Although a bit common in the (somewhat scarce) literature on Isbell duality, I have doubts regarding how useful/nice of a choice O and Spec are, and whether there are better choices of notation for them.

9. Interaction with  $\times$ , Hom,  $F_!$ ,  $F^*$ , and  $F_*$
10. Interactions between presheaves and copresheaves:
  - (a) Natural transformations from a presheaf to a copresheaf and vice versa
  - (b) Mixed Day convolution?
11. Isbell duality for monoids:
  - (a) Set up a dictionary between properties of  $\text{Sets}_A^L$  or  $\text{Sets}_A^R$  and properties of  $A$
  - (b) Do the same for  $O$  given by  $A \mapsto \text{Sets}_A^L(X, A)$
  - (c) Do the same for  $\text{Spec}$  given by  $A \mapsto \text{Sets}_A^R(X, A)$
  - (d) Do the same for  $O \circ \text{Spec}$
  - (e) Do the same for  $\text{Spec} \circ O$
  - (f) Algebras for  $\text{Spec} \circ O$
  - (g) Coalgebras for  $O \circ \text{Spec}$
12. Properties of  $\text{Spec}$  (e.g. fully faithfulness) vs. properties of  $C$
13. Properties of  $O$  (e.g. fully faithfulness) vs. properties of  $C$
14. co/unit being monomorphism/epimorphism
15. reflexive completion
16. Isbell duality for simplicial sets; what's the reflexive completion?
17. Isbell envelope
18. What does Isbell duality look like, when  $\text{Cat}(\text{Aop}, \text{Set})$  is identified with the category of discrete opfibrations over  $A$ , using A.5.14?
19. Generalizations of Isbell duality:
  - (a) Monoidal Isbell duality: monoidality for Isbell adjunction with day convolution (6.3 of coend cofriend)
  - (b) Isbell duality with sheaves
  - (c) Isbell duality with Lawvere theories, product preserving functors or whatever
  - (d) Isbell duality for profunctors
    - i. In view of ?? of ??, can we just use right Kan lifts/extensions?

- ii. Right Kan lift/extension of Hom functors (there's probably a version of the Yoneda lemma here)
- What is  $\text{Rift}_F(\text{Hom}_C)$
  - What is  $\text{Ran}_F(\text{Hom}_C)$
  - What is  $\text{Rift}_{\text{Hom}_C}(F)$
  - What is  $\text{Ran}_{\text{Hom}_C}(F)$
  - What is  $\text{Lift}_F(\text{Hom}_C)$
  - What is  $\text{Lan}_F(\text{Hom}_C)$
  - What is  $\text{Lift}_{\text{Hom}_C}(F)$
  - What is  $\text{Lan}_{\text{Hom}_C}(F)$

20. Tensor product of functors and Isbell duality

- What is  $\mathcal{F} \boxtimes_C \mathcal{O}(\mathcal{F})$ ?
- What is  $\text{Spec}(F) \boxtimes_C F$ ?
- I think there is a canonical morphism

$$\mathcal{F} \boxtimes_C \mathcal{O}(\mathcal{F}) \rightarrow \text{Tr}(C).$$

By the way, what is  $\text{Tr}(\Delta)$ ? What is  $\text{Tr}(BA)$ ? What about  $\text{Nat}(\text{id}_C, \text{id}_C)$  for  $C = BA$  or  $C = \Delta$

21. Isbell with coends:

- $\text{Hom}(F(A), h_A)$  but it's a coend
- Conatural transformations and all that

22. Co/limit preservation for  $\mathcal{O}/\text{Spec}$

23. Isbell duality for  $N$  vs.  $N + N$

24. What do we get if we replace  $\mathcal{O} \stackrel{\text{def}}{=} \text{Nat}(-, h_X)$  by  $\text{Nat}^{[W]}(-, h_X)$ , and in particular by  $\text{DiNat}(-, h_X)$ ?

Species:

- Joyal–Street's  $q$ -species; via promonoidal structures <https://arxiv.org/pdf/1201.2991.pdf#page=22>
- associators, braidings, unitors;  $\mathbb{F}_q^n \rightarrow \mathbb{F}_q^n$  centre of  $\text{GL}_n(\mathbb{F}_q)$  trick
- group completion of  $\mathcal{GL}(\mathbb{F}_q)$  as algebraic k-theory

Constructions With Categories:

- <https://arxiv.org/abs/2504.21764>

2. Comparison between pseudopullbacks and isocomma categories: the “evident” functor  $C \times_{\mathcal{E}}^{\text{ps}} \mathcal{D} \rightarrow C \times_{\mathcal{E}} \mathcal{D}$  is essentially surjective and full, but not faithful in general.
3. Quotients of categories by actions of monoidal categories
  - (a) Quotients of categories by actions of monoids  $\mathbf{BA}$
  - (b) Quotients of categories by actions of monoids  $A_{\text{disc}}$
  - (c) Lax, oplax, pseudo, strict, etc. quotients of categories
  - (d) lax Kan extensions along  $\mathbf{BC} \rightarrow \mathbf{BD}$  for  $C \rightarrow \mathcal{D}$  a monoidal functor
4. Quotient of  $\text{Fun}(\mathbf{BA}, C)$  by the  $A$ -action.
  - (a) This is used to build the cycle and  $p$ -cycle categories from the paracycle category.
  - (b) The quotient of  $\text{Fun}(\mathbf{BN}, C)$  by the  $\mathbb{N}$ -action should act as a kind of cyclic directed loop space of  $C$
5.  $\text{Fun}(\mathbf{BN}, C)$  as a homotopy pullback in  $\text{Cats}_2$ 
  - (a)  $\text{Fun}(\mathbf{BZ}, C)$  as a homotopy pullback in  $\text{Grpd}_2$
  - (b) Free loop space objects

Limits and colimits:

1. adjunction between co/product and diagonal; abstract version of [Item 3](#) and [Item 2](#)
2. Examples of kan extensions along functors of the form  $\text{FinSets} \hookrightarrow \text{Sets}$
3. Initial/terminal objects as left/right adjoints to  $!_C : C \rightarrow \text{pt}$ .
4. A small cocomplete category is a poset, <https://mathoverflow.net/questions/108737/small-categories-and-completeness>
5. Co/limits in  $\mathbf{BA}$ , including e.g. co/equalisers in  $\mathbf{BA}$
6. Add the characterisations of absolutely dense functors given in ?? to [Item 10](#).
7. Absolutely dense functors, <https://ncatlab.org/nlab/show/absolutely+dense+functor>. Also theorem 1.1 here: <http://www.tac.mta.ca/tac/volumes/8/n20/n20.pdf>.

8. Dense functors, codense functors, and absolutely codense functors.
9. van kampen colimits

Completions and cocompletions:

1. <https://mathoverflow.net/questions/429003/manifolds-a-s-cauchy-completed-objects>
2. what is the conservative cocompletion of smooth manifolds? Is it related to diffeological spaces?
3. what is the conservative completion of smooth manifolds? Is it related to diffeological spaces?
4. what is the conservative bicompletion of smooth manifolds? Is it related to diffeological spaces?
5. completion of a category under exponentials
6. <https://mathoverflow.net/questions/468897/cocompletion-without-cocontinuous-functors>
7. The free cocompletion of a category;
8. The free completion of a category;
9. The free completion under finite products;
10. The free cocompletion under finite coproducts;
11. The free bicompletion of a category;
12. The free bicompletion of a category under nonempty products and nonempty coproducts (<https://ncatlab.org/nlab/show/free+bicompletion>);
13. Cauchy completions
14. Dedekind–MacNeille completions
15. Isbell completion (<https://ncatlab.org/nlab/show/reflexive+completion>)
16. Isbell envelope

Ends and Coends:

1. motivate co/ends as co/limits of profunctors

2. Ask Fosco about whether composition of dinatural transformations into higher dinaturals could be useful for  
<https://arxiv.org/abs/2409.10237>

3. Cyclic co/ends

- (a) Try to mimic the construction given in Haugseng for the cycle, paracycle, cube, etc. categories
- (b) cyclotomic stuff for cyclic co/ends
  - i. Check out Ayala–Mazel-Gee–Rozenblyum’s *Symmetries of the cyclic nerve*
  - ii. isogenetic  $\mathbb{N}^\times$ -action (what the fuck does this mean?)

4. After stating the co/ends

$$\int_{A \in C}^{A \in C} h_A \odot \mathcal{F}^A, \quad \int_{A \in C}^{A \in C} \text{Sets}(h_A, \mathcal{F}^A),$$

$$\int_{A \in C}^{A \in C} h^A \odot F_A, \quad \int_{A \in C}^{A \in C} \text{Sets}(h^A, F_A)$$

in the co/end version of the Yoneda lemma, add a remark explaining what the co/ends

$$\int_{A \in C}^{A \in C} h_A \odot \mathcal{F}^A, \quad \int_{A \in C}^{A \in C} \text{Sets}(h_A, \mathcal{F}^A),$$

$$\int_{A \in C}^{A \in C} h^A \odot F_A, \quad \int_{A \in C}^{A \in C} \text{Sets}(h^A, F_A)$$

and the co/ends

$$\int_{A \in C}^{A \in C} \mathcal{F}^A \odot h_A, \quad \int_{A \in C}^{A \in C} \text{Sets}(\mathcal{F}^A, h_A),$$

$$\int_{A \in C}^{A \in C} F_A \odot h^A, \quad \int_{A \in C}^{A \in C} \text{Sets}(F_A, h^A),$$

$$\int_{A \in C}^{A \in C} \mathcal{F}^A \odot h_A, \quad \int_{A \in C}^{A \in C} \text{Sets}(\mathcal{F}^A, h_A),$$

$$\int_{A \in C}^{A \in C} F_A \odot h^A, \quad \int_{A \in C}^{A \in C} \text{Sets}(F_A, h^A)$$

are.

5. ends  $C \rightarrow \mathcal{D}$  with  $\odot$  is a special case of ends for a certain enrichment over  $\mathcal{D}$

6. try to figure out what the end/coend

$$\int^{X \in C} h_X^A \times h_B^X, \quad \int_{X \in C} h_X^A \times h_B^X$$

are for  $C = BA$ . (I think the coend is like tensor product of  $A$  as a left  $A$ -set with it as a right  $A$ -set)

- 7. Cyclic ends
- 8. Dihedral ends
- 9. Does Haugseng's constructions give a way to define cyclic co/homology with coefficients in a bimodule?
- 10. Category of elements of dinatural transformation classifier
- 11. Examples of co/ends: <https://mathoverflow.net/a/461814>
- 12. Cofinality for co/ends, <https://mathoverflow.net/questions/353876>
- 13. “Fourier transforms” as in <https://arxiv.org/pdf/1501.02503#page=168> or <https://tetrapharmakon.github.io/stuff/itaca.pdf>

Weighted/diagonal category theory:

- 1. co/ends as centre/trace-infused co/limits: compare the co/end of  $\text{Hom}_C$  with the co/limit of  $\text{Hom}_C$
- 2. Codensity  $W$ -weighted monads,  $\text{Ran}_F^{[W]}(F)$ ;
- 3. Codensity diagonal monads,  $\text{DiRan}_F(F)$ ;

Profunctors:

- 1. Apartness defines a composition for relations, but its analogue

$$\mathfrak{q} \square \mathfrak{p} \stackrel{\text{def}}{=} \int_{A \in C} \mathfrak{p}_A^{-1} \amalg \mathfrak{q}_{-2}^A$$

fails to be unital for profunctors with the unit  $h_-^A$ . Is it unital for some other unit? Is there a less obvious analogue of apartness composition for profunctors? Or maybe does Prof equipped with  $\square$  and units  $h_-^A$  form a skew bicategory?

Is  $\Delta_\emptyset$  a unit?

2. Figure what monoidal category structures on Sets induce associative and unital compositions on Prof.
3. <https://mathoverflow.net/questions/470213/a-distributor-between-categories-induces-a-distributor-between-their-categories>
4. Different compositions for profunctors from monoidal structures on the category of sets (e.g. <https://mathoverflow.net/questions/155939/what-other-monoidal-structures-exist-on-the-category-of-sets>)
5. Nucleus of a profunctor;
6. Isbell duality for profunctors:
  - (a) <https://mathoverflow.net/questions/259525/isbell-duality-for-profunctors>
  - (b) <https://mathoverflow.net/questions/260322/the-mathfrak-l-functor-on-textsfprof>
  - (c) <https://mathoverflow.net/questions/262462/again-on-the-mathfrak-l-functor-on-mathsfpf>

Centres and Traces of Categories:

1.  $K_0(\text{Fun}(\mathbb{N}, C))$  vs.  $\pi_0(\text{Fun}(\mathbb{N}, C))$  vs.  $\text{Tr}(C)$ , and how these are generalisations of conjugacy classes for monoids
2. Explicitly work out the trace and  $\pi_0 \text{Fun}(\mathbb{N}, -)$  for monoids with few elements.
3.  $[1_A]$  can contain more than one element. An example is  $\text{Sets}(\mathbb{N}, \mathbb{N})$  and the maps given by

$$\begin{aligned}\{0, 1, 2, 3, \dots\} &\mapsto \{0, 0, 1, 2, \dots\}, \\ \{0, 1, 2, 3, \dots\} &\mapsto \{2, 3, 4, 5, \dots\}.\end{aligned}$$

Show also that if  $c \in [1_A]$ , then  $c$  is idempotent.

4. Drinfeld centre
5. trace of the symmetric simplex category; it's probably different from that of FinSets
6. Trace of  $\text{Rep}_G$  and interaction with induction, restriction, etc.

7.  $\pi_0(\text{BN}, \text{BA})$ ,  $K(\text{BN}, \text{BA})$ , and  $\text{Tr}(\text{BN}, \text{BA})$  as concepts of conjugacy for monoids, their equivalents for categories, and comparison with traces
8. Comparison between  $\pi_0(\text{Fun}(\text{BN}, C))$  and  $K(\text{Fun}(\text{BN}, C))$
9. Lax, oplax, pseudo, and strict trace of simplex 2-category
10. duality over  $\Gamma$  might give a map from product of a monoid with a set to  $\text{Tr}(\Gamma)$
11. Studying the set  $\text{Nat}(\text{id}_C, F)$  as a notion of categorical trace:
  - (a) Ganter–Kapranov define the trace of a 1-endomorphism  $f: A \rightarrow A$  in a 2-category  $C$  to be the set  $\text{Hom}_C(\text{id}_A, f)$ ;
    - i. <https://arxiv.org/abs/math/0602510>
    - ii. <https://golem.ph.utexas.edu/string/archives/000757.html>
    - iii. <https://ncatlab.org/nlab/show/categorical+trace>

We should study this notion in detail, and also study  $\text{Nat}(F, \text{id}_C)$  as well as  $\text{CoNat}(\text{id}_C, F)$  and  $\text{CoNat}(F, \text{id}_C)$ .
12. Centre of bicategories
13. Lax centres and lax traces
14. Examples of traces:
  - (a) Discrete categories
  - (b) Posets
    - i.  $\text{Open}(X)$
  - (c) Trace of small but non-finite categories:
    - i. Sets
    - ii.  $\text{Rep}(G)$
    - iii. category of finite groups
    - iv. category of finite abelian groups
    - v. category of finite  $p$ -groups for fixed  $p$
    - vi. category of finite  $p$ -groups for all  $p$
    - vii. category of finite fields
    - viii. category of finite topological spaces
    - ix. category of finite [insert a mathematical object here]
15. When is the trace of a groupoid just the disjoint sum of sets of conjugacy classes?

16. Set-theoretical issues when defining traces

- (a) Sets is a large category, and yet we can speak of its centre

$$\begin{aligned} Z(\text{Sets}) &\stackrel{\text{def}}{=} \int_{A \in \text{Sets}} \text{Sets}(X, X) \\ &\cong \text{Nat}(\text{id}_{\text{Sets}}, \text{id}_{\text{Sets}}) \\ &\cong \text{pt}. \end{aligned}$$

Is there a way to do the same for the trace of sets, or otherwise work with traces of large categories?

17. Understand how traces are defined via universal properties in Xinwen Zhu's [Geometric Satake, categorical traces, and arithmetic of Shimura varieties](#).

18. trace as an  $\text{Obj}(C)$ -indexed set

- (a) properties, functoriality, etc.

19. Maybe actually call  $\text{Fun}(\text{BN}, C)$  the categorical directed loop space of  $C$ ?

20. Cyclic version of  $\text{Fun}(\text{BN}, C)$

21. Traces of categories, nerves of categories, and the cycle category

Categorical Hochschild Homology:

1. To any functor we have an associated natural transformation ([Definition 11.5.4.1.1](#)). Do we have sharp transformations associated to natural transformation?
2. build Hochschild co/simplicial set and study its homotopy groups
3.  $\text{Fun}(\text{BN}, X_\bullet)$  vs.  $\text{Fun}(\Delta^1 / \partial \Delta^1, X_\bullet)$ 
  - (a) Their  $\pi_0$ 's vs. the  $\pi_0$ 's of  $\text{Hom}_{X_\bullet}(x, x)$ , of  $\text{Hom}_{X_\bullet}^L(x, x)$ , and  $\text{Hom}_{X_\bullet}^R(x, x)$ .

Monoidal Categories:

1. <https://mathoverflow.net/questions/380302>
2. Analogue of Picard rings for dualisable objects
3. Moduli of associators, braidings, etc. for species,  $q$ -species
4. When is the left Kan extension along a fully faithful functor of monoidal categories a strong monoidal functor?

5. Interaction between Day convolution and Isbell duality
6. general theory for lifting pseudomonads from Cat to Prof along the equipment embedding
7. definition of prostrength on a functor between promonoidal categories, differential 2-rigs fosco
8. Promonoidal structure in <https://arxiv.org/pdf/1201.2991#page=22>
9. Day convolution as a colimit over category of factorizations  $F(A) \otimes_C G(B) \rightarrow V$
10. Day convolution with respect to Cartesian monoidal structure is Cartesian monoidal. There's an easy proof of this with coend Yoneda
11. <https://mathoverflow.net/questions/491234>
12. <https://mathoverflow.net/questions/488426/adjunction-of-monoidal-closed-categories>
13. <https://arxiv.org/abs/2502.02532>
14. Does the forgetful functor  $\text{IdemMon}(C) \rightarrow \text{Mon}(C)$  admit a left adjoint? What about  $\text{IdemMon}(C) \rightarrow C$ ?
15. Clifford algebras in monoidal categories
16. Exterior algebras in monoidal categories
  - (a) <https://mathoverflow.net/questions/70607/exterior-powers-in-tensor-categories>
  - (b) <https://mathoverflow.net/questions/127476/analogy-between-the-exterior-power-and-the-power-set>
  - (c) <https://mathoverflow.net/questions/182476/delignes-exterior-power>
  - (d) martin brandenburg's phd thesis
17. Different monoidal products in  $\text{Fun}(C, C)$  and their distributivity
  - (a) Composition
  - (b) Pointwise product
  - (c) Day convolution
  - (d) Relative monad version of Day convolution

18. Classification of monoidal structures on  $\Delta$
19. Classification of monoidal structures on  $\Lambda$
20. Tensor Categories, 8.5.4
21. <https://ncatlab.org/nlab/show/monoidal+action+of+a+monoidal+category>
22. <https://arxiv.org/abs/2203.16351>
23. Para construction
24. Drinfeld center; Symmetric center; JY's books on bimonoidal categories
25. Picard and Brauer 2-groups
  - (a) Differential Picard and Brauer Groups via  $\text{Fun}(\text{B}\mathbb{N}, \text{Mod}_R)$ .
  - (b) Brauer and Picard groups of  $(\text{Fun}(C, C), \circ, \text{id}_C)$
  - (c) Brauer and Picard groups of  $\text{Rep}(G)$
  - (d) Brauer and Picard groups of Sets
  - (e) Brauer and Picard groups of  $\text{Ch}_{\mathbb{Z}}(R)$
  - (f) Brauer and Picard groups of  $\text{Shv}(X)$
  - (g) Brauer and Picard groups of  $\text{dgMod}_R$
26. Explore examples in which Day convolution gives weird things, like  $\text{Fun}(\text{B}\mathbb{Z}_{/n}, \text{Sets})$ .
27. Day convolution is a left Kan extension; explore the right Kan extension
28. Further develop the theory of moduli categories of monoidal structures
29. Picard group
  - (a) Picard group for Day convolution. A special case is one of Kaplansky's conjectures, [https://en.wikipedia.org/wiki/Kaplansky%27s\\_conjectures](https://en.wikipedia.org/wiki/Kaplansky%27s_conjectures), about units of group rings
30. Day convolution between representable and an arbitrary presheaf  $\mathcal{F}$  — can we prove something nice using the colimit formula for  $\mathcal{F}$  in terms of representables?
31. Notion of braided monoidal categories in which the braiding is not an isomorphism. Relation to <https://arxiv.org/abs/1307.5969>

32. Proving a certain diagram between free monoidal categories commutes involves Fermat's little theorem. Can we reverse this and prove Fermat's little theorem from the commutativity of that diagram?
33. <https://nilesjohnson.net/notes/grPic-P2S.pdf>
34. Proof that monoidal equivalences  $F$  of monoidal categories automatically admit monoidal natural isomorphisms  $\text{id}_C \cong F^{-1} \circ F$  and  $\text{id}_{\mathcal{D}} \cong F \circ F^{-1}$ .
35. Proof that category with products is monoidal under the Cartesian monoidal structure, [MO 382264].
36. Explore 2-categorical algebra:
- (a) Find a construction of the free 2-group on a monoidal category. Apply it to the multiplicative structure on the category of finite sets and permutations, as well as to the multiplicative structure on the 1-truncation of the sphere spectrum, and try to figure out whether this looks like a categorification of  $\mathbb{Q}$ .
  - (b) What is the free 2-group on  $(\Delta, \oplus, [0])$ ?
37. Categorify the preorder  $\leq$  on  $\mathbb{N}$  to a promonad  $\mathfrak{p}$  on the groupoid of finite sets and permutations  $\mathbb{F}$ :
- (a) A preorder is a monad in  $\text{Rel}$
  - (b) A promonad is a monad in  $\text{Prof}$ .
  - (c) There's a promonad  $\mathfrak{p}$  in  $\mathbb{F}$  defined by
- $$\mathfrak{p}(m, n) \stackrel{\text{def}}{=} \{\text{surjections from } \{1, \dots, m\} \text{ to } \{1, \dots, n\}\}$$
- This promonad categorifies  $\leq$  in that its values are the witnesses to the fact that  $m$  is bigger than  $n$  (i.e. surjections).
- (d) Figure out whether this promonad extends to the 1-truncation of the sphere spectrum, and perhaps to other categorified analogues of monoids/groups/rings.
38. <https://arxiv.org/abs/1307.5969>
39. <https://arxiv.org/abs/1306.3215>
40. <https://mathoverflow.net/questions/477219/reference-for-the-monoidal-category-structure-x-otimes-y-x-y-x-times-y>

41. Include an explicit proof of [Item 14](#)
42. Include an explicit proof of [Item 6](#)
43. [Remark 4.1.3.1.6](#)
44. obstruction theory for braided enhancements of monoidal categories, using the “moduli category of braided enhancements”
45. Define symmetric and exterior algebras internal to braided monoidal categories
  - (a) <https://mathoverflow.net/questions/471372/is-there-an-alternating-power-functor-on-braided-monoidal-categories>
  - (b) <https://arxiv.org/abs/math/0504155>
46. <https://mathoverflow.net/q/382364>
47. <https://mathoverflow.net/q/471490>
48. Concepts of bicategories applied to monoidal categories (e.g. internal adjunctions lead to dualisable objects)
49. Involutive Category Theory
50. <https://mathoverflow.net/questions/474662/the-analogy-between-dualizable-categories-and-compact-hausdorff-spaces>

Bimonoidal Categories:

1. Bimonoidal structures on the category of species
2. Include an explicit proof of [Item 15](#)

Six Functor Formalisms:

1. Michael Shulman:

A lot of the “six functor formalism” makes sense in the context of an arbitrary indexed monoidal category (= monoidal fibration), particularly with cartesian base. In particular, I studied the external tensor product in this generality in my paper on Framed bicategories and monoidal fibrations.

The internal-hom of powersets in particular, with  $\emptyset$  as a dualizing object, is well-known in constructive mathematics and topos theory, where powersets are in general a Heyting algebra rather than a Boolean algebra.

Morgan Rogers:

I second this: you're discovering (and making pleasingly explicit, I might add) a special case of "thin category theory": a lot of what you've discovered will work for posets, with the powerset replaced with the frame of downsets :D

2. A six functor formalism for monoids
3. <https://mathoverflow.net/questions/258159/yoga-of-six-functors-for-group-representations>
4. Is the 1-categorical analogue of six functor formalisms given by Mann interesting?
  - (a) Mann defines:  
A six functor formalism is an  $\infty$ -functor  $f: \text{Corr}(C, E) \rightarrow \text{Cats}_\infty$  such that  $- \otimes A$ ,  $f^*$ , and  $f_!$  admit right adjoints
  - (b) Is the notion  
A 1-categorical six functor formalism is a (lax?) 2-functor  $f: \text{Corr}(C, E) \rightarrow \text{Cats}_2$  (or should Cats be the target?) such that  $- \otimes A$ ,  $f^*$ , and  $f_!$  admit right adjoints  
interesting?
5. Interaction of the six functors with Kan extensions (e.g. how the left Kan extension of  $- \otimes A$  may interact with the other functors)
6. Contexts like Wirthmuller Grothendieck etc
7. formalisation by cisinski and deglise
8. How do the following examples fit?
  - (a) base change between  $C_X$  and  $C_Y$
  - (b)  $f_! \dashv f_* \dashv f^*$  adjunction between powersets
  - (c)  $f_! \dashv f_* \dashv f^*$  adjunction between  $\text{Span}(\text{pt}, A)$  and  $\text{Span}(\text{pt}, B)$
  - (d) quadruple adjunction between powersets induced by a relation
  - (e) adjunctions between categories of presheaves induced by a functor or a profunctor
  - (f) Adjunction between left  $A$ -sets and left  $B$ -sets

Do they have exceptional  $f^!$ ? Is there a notion of Fourier–Mukai transform for them? What kind of compatibility conditions (proper base change, etc.) do we have?

Skew Monoidal Categories:

1. <https://arxiv.org/abs/2506.06847>
2. Try to come up with examples of skew monoidal categories by twisting a tensor product  $A \otimes B$  into  $T(A) \otimes B$ . Related idea: product of  $G$ -sets but twisted on the left by an automorphism of  $G$ , so that  $(ag, b) \sim (a, gb)$  becomes  $(a\phi(g), b) \sim (a, gb)$ .
3. Skew monoidal category induced from  $G$ -sets in analogy to Rel
4. Free monoidal category on a skew monoidal category
5. Skew monoidal structures associated to a locally Cartesian closed category
6. Does the  $\mathbb{E}_1$  tensor product of monoids admit a skew monoidal category structure?
7. Is there a (right?) skew monoidal category structure on  $\text{Fun}(C, \mathcal{D})$  using right Kan extensions instead of left Kan extensions?
8. Similarly, are there skew monoidal category structures on the subcategory of  $\text{Rel}(A, B)$  spanned by the functions using left Kan extensions and left Kan lifts?
9. Add example:  $C$  with coproducts, take  $C_{X/}$  and define

$$(X \xrightarrow{f} A) \oplus (X \xrightarrow{g} B) \stackrel{\text{def}}{=} [X \rightarrow X \coprod X \xrightarrow{f \coprod g} A \coprod B]$$

10. Duals:

- (a) Dualisable objects in monoidal categories and traces of endomorphisms of them, including also examples for monoidal categories which are not autonomous/rigid, such as  $(\text{Fun}(C, C), \circ, \text{id}_C)$ .
- (b) compact closed categories
- (c) star autonomous categories
- (d) Chu construction
- (e) Balanced monoidal categories, <https://ncatlab.org/nlab/show/balanced+monoidal+category>

(f) Traced monoidal categories, <https://ncatlab.org/nlab/show/traced+monoidal+category>

11. Invertible objects and Picard groupoids
12. <https://mathoverflow.net/questions/155939/what-other-monoidal-structures-exist-on-the-category-of-sets>
13. Free braided monoidal category with a braided monoid: <https://ncatlab.org/nlab/show/vine>
14. [https://golem.ph.utexas.edu/category/2024/08/skew\\_monoidal\\_categories\\_throu.html](https://golem.ph.utexas.edu/category/2024/08/skew_monoidal_categories_throu.html)

Fibred Category Theory:

1. <https://arxiv.org/abs/2402.11644>
2. <https://categorytheory.zulipchat.com/#narrow/channel1/29136-theory.3A-category-theory/topic/A.20.22change.20of.20variables.22.20for.20the.20Grothendieck.20construction/near/495776958>
3. Internal **Hom** in categories of co/Cartesian fibrations.
4. *Tensor structures on fibered categories* by Luca Terenzi: <https://arxiv.org/abs/2401.13491>. Check also the other papers by Luca Terenzi.
5. <https://ncatlab.org/nlab/show/cartesian+natural+transformation> (this is a cartesian morphism in  $\text{Fun}(C, \mathcal{D})$  apparently)
6. CoCartesian fibration classifying  $\text{Fun}(F, G)$ , <https://mathoverflow.net/questions/457533/cocartesian-fibration-classifying-mathrmfunf-g>

Operads and Multicategories:

1. [Simplicial lists in operad theory I](#)

Monads:

1. Relative monads: message Alyssa asking for her notes
2. <https://ncatlab.org/nlab/show/adjoint+monad>
3. Kantorovich monad (<https://ncatlab.org/nlab/show/Kantorovich+monad>) and probability monads in general, <https://ncatlab.org/nlab/show/monads+of+probability%2C+measures%2C+and+valuations>.

Enriched Categories:

1.  $\mathcal{V}$ -matrices

Bicategories:

1. Bicategories of Lax Fractions, <https://arxiv.org/abs/2507.12044>
2. Linear bicategories, <https://ncatlab.org/nlab/show/linear+bicategory>
  - (a) Linearly distributive category, <https://ncatlab.org/nlab/show/linearly+distributive+category>
  - (b) Diagrammatic Algebra of First Order Logic
  - (c) Constructing linear bicategories
  - (d) Introduction to linear bicategories
3. Allegories, <https://ncatlab.org/nlab/show/allegory>
4. Skew bicategories
5. Bigroupoid cardinality
6. Bicategory where objects are groups and a morphism  $G \rightarrow H$  is a representation of  $G^{\text{op}} \times H$ . (I.e. functors  $BG^{\text{op}} \times BH \rightarrow \text{Vect}_k$ ).
7. Relative monads internal to a bicategory
8. Bicategory of monoid actions
9. <https://arxiv.org/abs/0809.1760>
10.  $\text{Rel}_G \stackrel{\text{def}}{=} \text{Fun}(BG, \text{Rel})$
11. Rel but for Ab, where morphisms are pairings of the form  $A \otimes_{\mathbb{Z}} B \rightarrow \mathbb{Z}$ .
12. 2-dimensional co/limits in 2-category of categories and adjoint functors
13. Category of equivalence classes
  - (a) Given a category  $C$ , we have a set  $K_0(C)$  of isomorphism classes of objects
  - (b) Given a bicategory  $C$ , there should be a category  $K_0(C)$  with  $\text{Hom}_{K_0(C)}(A, B) \stackrel{\text{def}}{=} K_0(\text{Hom}_C(A, B))$

(c) The set  $K_0^{\text{eq}}(C)$  of equivalence classes of objects of  $C$  should then satisfy

$$K_0^{\text{eq}}(C) \cong K_0(K_0(C)).$$

- 14. bicategory of chain complexes, section “Second Example: Differential Complexes of an Abelian Category” on Gabriel–Zisman’s calculus of fractions
- 15. 2-vector spaces
- 16. Morita equivalence is equivalence internal to bimod
- 17. <https://mathoverflow.net/questions/478867/2-category-structure-on-modR>
- 18. Bicategories of matrices, as in Street’s Variation through enrichment, also <https://arxiv.org/abs/2410.18877>
- 19. <https://mathoverflow.net/a/86933>
- 20. What are the internal 2-adjunctions in the fundamental 2-groupoid of a space?
- 21. 2-category structure on  $\text{Mod}_R$ , where a 2-morphism is a commutative square. Characterisation of adjunctions therein
- 22. Cook up a very large list of examples of bicategories, like the ones I made for the AI problems. In particular, find an interesting bicategory of representations qualitatively different from the one I described in the Epoch AI problem
- 23. 2-category structure on category of  $R$ -algebras as enriched  $\text{Mod}_R$ -categories
- 24. Let  $C$  be a bicategory, let  $A, B \in \text{Obj}(C)$ , and let  $F, G \in \text{Obj}(\text{Hom}_C(A, B))$ .
  - (a) Does precomposition with  $\lambda_{A|F}^C : \text{id}_A \circ F \Rightarrow F$  induce an isomorphism of sets
 
$$\text{Hom}_{\text{Hom}_C(A, B)}(F, G) \cong \text{Hom}_{\text{Hom}_C(A, B)}(F \circ \text{id}_A, G)$$
 for each  $F, G \in \text{Obj}(\text{Hom}_C(A, B))$ ?
  - (b) Similarly, do we have an induced isomorphism of the form
 
$$\text{Hom}_{\text{Hom}_C(A, B)}(F, G) \cong \text{Hom}_{\text{Hom}_C(A, B)}(F, \text{id}_B \circ G)$$
 and so on?

25. Are there two Duskin nerve functors? (lax/oplax/etc.?)
26. Interaction with cotransformations:
- (a) Can we abstract the structure provided to  $\text{Cats}_2$  by natural cotransformations?
  - (b) Are there analogues of cotransformations for **Rel**, **Span**, **BiMod**, **MonAct**, etc.?
  - (c) Perhaps this might also make sense as a 1-categorical definition, e.g. comorphisms of groups from  $A$  to  $B$  as  $\text{Sets}(A, B)$  quotiented by  $f(ab) \sim f(a)f(b)$ .
27. Consider developing the analogue of traces for endomorphisms of dualisable objects in monoidal categories to the setting of bicategories, including e.g. the trace of a category as a trace internal to **Prof**.
28. Centres of bicategories (lax, strict, etc.)
29. Concepts of monoidal categories applied to bicategories (e.g. traces)
30. Internal adjunctions in **Mod** as in [JY21, Section 6.3]; see [JY21, Example 6.2.6].
31. Comonads in the bicategory of profunctors.
32. 2-limit of  $\text{id}, \text{id} : \text{Sets} \Rightarrow \text{Sets}$  is  $\text{B}\mathbb{Z}$ , [https://mathoverflow.net/questions/209904/van-kampen-colimits?rq=1#comment520288\\_209904](https://mathoverflow.net/questions/209904/van-kampen-colimits?rq=1#comment520288_209904)
33. <https://mathoverflow.net/questions/473527/universal-property-of-2-presheaves-and-pseudo-lax-colax-natural-transformations>
34. <https://mathoverflow.net/questions/473526/free-cocompletion-of-a-2-category-under-pseudo-colimits-lax-colimits-and-colax>

Types of Morphisms in Bicategories:

1. Behaviour in 2-categories of pseudofunctors (or lax functors, etc.), e.g. pointwise pseudoepic morphisms in vs. pseudoepic morphisms in 2-categories of pseudofunctors.
2. Statements like “coequifiers are lax epimorphisms”, Item 2 of Examples 2.4 of <https://arxiv.org/abs/2109.09836>, along with most of the other statements/examples there.

3. Dense, absolutely dense, etc. morphisms in bicategories

Internal adjunctions:

1. <https://www.google.com/search?q=mate+of+an+adjunction>
2. Moreover, by uniqueness of adjoints (?? of ??), this implies also that  $S = f^{-1}$ .
3. define bicategory  $\text{Adj}(C)$
4. walking monad
5. proposition: 2-functors preserve unitors and associators
6. <https://ncatlab.org/nlab/show/2-category+of+adjunctions>. Is there a 3-category too?
7. <https://ncatlab.org/nlab/show/free+monad>
8. <https://ncatlab.org/nlab/show/CatAdj>
9. <https://ncatlab.org/nlab/show/Adj>
10.  $\text{Adj}(\text{Adj}(C))$
11. Examples of internal adjunctions
  - (a) Internal adjunctions in  $\text{Mod}$ .
  - (b) Internal adjunctions in  $\text{PseudoFun}(C, \mathcal{D})$ .
  - (c) Internal adjunctions in  $\text{LaxFun}(C, \mathcal{D})$ .
  - (d) Internal adjunctions in 2-categories related to fibrations.

2-Categorical Limits:

1. <https://sorilee.github.io/posts/strict-bilimit-and-its-proper-examples>

Double Categories:

1. Ehresmann
2. <https://arxiv.org/abs/2505.08766>
3. <https://arxiv.org/abs/2504.18065>
4. <https://arxiv.org/abs/2504.11099>
5. Pinwheel/Yojouhan diagrams and compositionality, section on nLab at <https://ncatlab.org/nlab/show/double+category>

Homological Algebra:

1. <https://arxiv.org/abs/2505.08321>
2. <https://mathoverflow.net/questions/418676/derived-functor-or-of-functor-tensor-product>
3. <https://math.stackexchange.com/questions/3665036/higher-chain-homotopies>

Topos theory:

1. <https://arxiv.org/abs/2505.08766>
2. <https://arxiv.org/abs/2304.05338>
3. <https://arxiv.org/abs/2503.20664>
4. <https://arxiv.org/abs/2204.08351>
5. <https://arxiv.org/abs/2404.12313>
6. <https://www.teses.usp.br/teses/disponiveis/45/45131/tde-31082023-163143/en.php>
7. <https://teses.usp.br/teses/disponiveis/45/45131/tde-24042019-195658/pt-br.php>
8. <https://mathoverflow.net/q/479496>
9. Grothendieck topologies on BA
10. Enriched Grothendieck topologies
  - (a) Borceux–Quintero, [https://www.numdam.org/item/CTGDC\\_1996\\_\\_37\\_2\\_145\\_0/](https://www.numdam.org/item/CTGDC_1996__37_2_145_0/)
  - (b) <https://arxiv.org/abs/2405.19529>
11. Cotoptos theory:
  - (a) Copresheaves and copresheaf cotoptoi
  - (b) Elementary cotoptoi
    - i. <https://mathoverflow.net/questions/474287/intuition-for-the-internal-logic-of-a-cotoptos>
    - ii. <https://mathoverflow.net/questions/394098/what-is-a-cotoptos>

In case you haven't seen it yet, Grothendieck studies (pseudo) cotoptos in [pursuing stacks](#)

Formal category theory:

1. Yosegi boxes <https://arxiv.org/abs/1901.01594>

Homotopical Algebra:

1. <https://arxiv.org/abs/2109.07803>

Simplicial stuff:

1. <https://arxiv.org/abs/2507.15341>
2. <https://arxiv.org/abs/2503.13663>
3. [https://www.math.univ-paris13.fr/~harpaz/quasi\\_unital.pdf](https://www.math.univ-paris13.fr/~harpaz/quasi_unital.pdf)

- (a) slogan: geometric definition of  $\infty$ -categories should be geometric for identities too
- (b) In an  $\infty$ -category, define a **quasi-unit** to be a 1-morphism  $f$  such that

$$[f]_* : \text{Hom}_{\text{Ho}(\text{Spaces})}(\text{Hom}_{\mathcal{C}}(X, A) \text{Hom}_{\mathcal{C}}(X, B)),$$

$$[f]^* : \text{Hom}_{\text{Ho}(\text{Spaces})}(\text{Hom}_{\mathcal{C}}(B, X) \text{Hom}_{\mathcal{C}}(A, X))$$

are the identity in  $\text{Ho}(\text{Spaces})$ . Explore equivalent conditions,

- (c) <https://arxiv.org/abs/1606.05669>
- (d) <https://arxiv.org/abs/1702.08696>
- 4. <https://arxiv.org/abs/math/0507116>, <https://arxiv.org/abs/2503.11338>
- 5. <https://arxiv.org/abs/2302.02484> and <https://arxiv.org/abs/2411.19751>
- 6. Internal adjunctions in  $\Delta$  are the same as Galois connections between  $[n]$  and  $[m]$ .
- 7. <https://mathoverflow.net/q/478461>
- 8. draw coherence for lax functors using the diagram for  $\Delta^2$
- 9. characterisation of simplicial sets such that left, right, and two-sided homotopies agree
- 10. every continuous simplicial set arises as the nerve of a poset.

11. Functor  $\text{sd}$  is convolution of  $\times_{\Delta}$  with itself; see <https://arxiv.org/pdf/1501.02503.pdf#page=109>

12. Extra degeneracies

(a) <https://www.google.com/search?client=firefox-b-d&q=augmented+simplicial+objects+with+extra+degeneracies>

(b) [https://leanprover-community.github.io/mathlib\\_docs/algebraic\\_topology/extr\\_degeneracy.html](https://leanprover-community.github.io/mathlib_docs/algebraic_topology/extr_degeneracy.html)

13. Comparison between  $\Delta^1/\partial\Delta^1$  and  $B\mathbb{N}$

$\infty$ -Categories:

1. <https://arxiv.org/abs/2505.22640>

2. <https://arxiv.org/abs/2410.17102>

3. <https://arxiv.org/abs/2410.02578>, [https://scholar.colorado.edu/concern/graduate\\_thesis\\_or\\_dissertations/st74cr650](https://scholar.colorado.edu/concern/graduate_thesis_or_dissertations/st74cr650), <https://arxiv.org/abs/2206.00849>

4. <https://mathoverflow.net/questions/479716/non-strictly-unital-functors-of-infinity-categories>

5. <https://mathoverflow.net/questions/472253/whats-the-localization-of-the-infty-category-of-categories-under-inverting-f>

Condensed Mathematics:

1. [https://golem.ph.utexas.edu/category/2020/03/pyknoticity\\_vs\\_cohesivenes.html#c057724](https://golem.ph.utexas.edu/category/2020/03/pyknoticity_vs_cohesivenes.html#c057724)

2. [https://golem.ph.utexas.edu/category/2020/03/pyknoticity\\_vs\\_cohesivenes.html#c057810](https://golem.ph.utexas.edu/category/2020/03/pyknoticity_vs_cohesivenes.html#c057810)

3. <https://maths.anu.edu.au/news-events/events/universal-property-category-condensed-sets>

4. <https://grossack.site/2024/07/03/life-in-johnstones-topological-topos>

5. <https://grossack.site/2024/07/03/topological-topos-2-algebras>

6. <https://grossack.site/2024/07/03/topological-topos-3-bonus-axioms>

7. <https://terrytao.wordpress.com/2025/04/23/stonean-space-s-projective-objects-the-riesz-representation-theorem-and-possibly-condensed-mathematics/>

Monoids:

1. <https://mathoverflow.net/questions/278429/>
2. Homological algebra of  $A$ -sets, <https://arxiv.org/abs/1503.02309>
3. Catalan monoids, <https://arxiv.org/abs/1309.6120>
4. <https://mathoverflow.net/questions/438305/grothendieck-group-of-the-fibonacci-monoid>
5. <https://math.stackexchange.com/questions/2662005/how-much-of-a-group-g-is-determined-by-the-category-of-g-sets>
6. <https://math.stackexchange.com/a/4996051/603207>, <https://arxiv.org/abs/1006.5687>
7. Six functor formalism for monoids, following [Section 4.6.4](#), but in which  $\cap$  and  $[-, -]$  are replaced with Day convolution.
8. Monoid  $(\{1, \dots, n\} \cup \infty, \text{gcd})$ . The element  $\infty$  can be replaced by  $p_1^{\min(e_1^1, \dots, e_1^m)} \cdots p_k^{\min(e_k^1, \dots, e_k^m)}$ .
9. Universal property of localisation of monoids as a left adjoint to the forgetful functor  $C \rightarrow \mathcal{D}$ , where:
  - $C$  is the category whose objects are pairs  $(A, S)$  with  $A$  a monoid and  $S$  a submonoid of  $A$ .
  - $\mathcal{D}$  is the category whose objects are pairs  $(A, S)$  with  $A$  a monoid and  $S$  a submonoid of  $A$  which is also a group.

Explore this also for localisations of rings

Explore if we can define field spectra with an approach like this

10. Adjunction between monoids and monoids with zero corresponding to  $(-)^- \dashv (-)^+$
11. Rock paper scissors as an example of a non-associative operation
12. <https://mathoverflow.net/questions/438305/grothendieck-group-of-the-fibonacci-monoid>

13. Witt monoid, <https://www.google.com/search?q=Witt+monoid>
14. semi-direct product of monoids, <https://ncatlab.org/nlab/show/semidirect+product+group>
15. morphisms of monoids as natural transformation between left  $A$ -sets over  $A$  and  $B_A$ .
16. Figure out if 2-morphisms of monoids coming from  $\text{Fun}^\otimes(A_{\text{disc}}, B_{\text{disc}})$ ,  $\text{PseudoFun}(BA, BB)$ , etc. are interesting
17. Write sections on the quotient and set of fixed points of a set by a monoid action
18. Isbell's zigzag theorem for semigroups: the following conditions are equivalent:
  - (a) A morphism  $f: A \rightarrow B$  of semigroups is an epimorphism.
  - (b) For each  $b \in B$ , one of the following conditions is satisfied:
    - We have  $f(a) = b$ .
    - There exist some  $m \in \mathbb{N}_{\geq 1}$  and two factorisations

$$\begin{aligned} b &= a_0 y_1, \\ b &= x_m a_{2m} \end{aligned}$$

connected by relations

$$\begin{aligned} a_0 &= x_1 a_1, \\ a_1 y_1 &= a_2 y_2, \\ x_1 a_2 &= x_2 a_3, \\ a_{2m-1} y_m &= a_{2m} \end{aligned}$$

such that, for each  $1 \leq i \leq m$ , we have  $a_i \in \text{Im}(f)$ .

Wikipedia says in [https://en.wikipedia.org/wiki/Isbell%27s\\_zigzag\\_theorem](https://en.wikipedia.org/wiki/Isbell%27s_zigzag_theorem):

For monoids, this theorem can be written more concisely:

19. Representation theory of monoids
  - (a) <https://mathoverflow.net/questions/37115/why-arent-representations-of-monoids-studied-so-much>
  - (b) Representation theory of groups associated to monoids (groups of units, group completions, etc.)

Monoid Actions:

1. <https://link.springer.com/book/10.1007/978-3-642-11297-3>
2. [https://ncatlab.org/schreiber/files/EquivariantInfinit\\_yBundles\\_220809.pdf](https://ncatlab.org/schreiber/files/EquivariantInfinit_yBundles_220809.pdf) has some interesting things, like a fully faithful embedding of  $\text{Mon}(\text{Sets}_A^L)$  into  $\text{Mon}_{/A}$  whose essential image is given by those monoids of the form  $X \rtimes_\alpha A$ .
3.  $f_! \dashv f^* \dashv f_*$  adjunction
  - (a) Is it related to the Kan extensions adjunction for  $f: BA \rightarrow BB$  and the categories  $\text{Sets}_A^L \cong \text{PSh}(BA^{\text{op}}, \text{Sets})$  and  $\text{Sets}_B^L \cong \text{PSh}(BB^{\text{op}}, \text{Sets})$ ?
  - (b) Is it related to the cobase change adjunction of <https://ncatlab.org/nlab/show/base+change>? Maybe we can take a morphism of monoids  $f: A \rightarrow B$  and consider  $B_A^L$  as a left  $A$ -set, and then  $(\text{Sets}_A^L)_{A/}$  and  $(\text{Sets}_A^L)_{B_A^L/}$
4. <https://arxiv.org/abs/2112.10198>
5. double category of monoid actions
6. Analogue of Brauer groups for  $A$ -sets
7. Hochschild homology for  $A$ -sets

Group Theory:

1. <https://mathoverflow.net/questions/45651/is-there-a-q-analog-to-the-braid-group>
2. <https://johncarlosbaez.wordpress.com/2025/03/27/the-mcg-e-group/>
3. <https://bookstore.ams.org/memo-1-2/>
4. <https://link.springer.com/book/10.1007/978-3-662-59144-4>
5. [https://en.wikipedia.org/wiki/Tits\\_group](https://en.wikipedia.org/wiki/Tits_group)
6. [https://en.wikipedia.org/wiki/Group\\_of\\_Lie\\_type](https://en.wikipedia.org/wiki/Group_of_Lie_type)
7. <https://mathoverflow.net/questions/251769/what-meaning-s-does-chevalley-group-have>
8. [https://encyclopediaofmath.org/wiki/Chevalley\\_group](https://encyclopediaofmath.org/wiki/Chevalley_group)

9. [https://en.wikipedia.org/wiki/Group\\_of\\_Lie\\_type](https://en.wikipedia.org/wiki/Group_of_Lie_type)
10. MO: cardinality of  $\text{Cl}(\text{Aut}(\text{GL}_n(\mathbb{F}_q)))$
11. <https://math.stackexchange.com/questions/4419869/do-the-groups-operatornamesl-operatornamepgl-and-operatornamepsl>
12. [https://groupprops.subwiki.org/wiki/Order\\_formulas\\_for\\_linear\\_groups](https://groupprops.subwiki.org/wiki/Order_formulas_for_linear_groups)
13. [https://groupprops.subwiki.org/wiki/Order\\_of\\_semidirect\\_product\\_is\\_product\\_of\\_orders](https://groupprops.subwiki.org/wiki/Order_of_semidirect_product_is_product_of_orders)
14. [https://groupprops.subwiki.org/wiki/Central\\_automorphism\\_group\\_of\\_general\\_linear\\_group](https://groupprops.subwiki.org/wiki/Central_automorphism_group_of_general_linear_group)
15. [https://groupprops.subwiki.org/wiki/Automorphism\\_group\\_of\\_general\\_linear\\_group\\_over\\_a\\_field](https://groupprops.subwiki.org/wiki/Automorphism_group_of_general_linear_group_over_a_field)
16. [https://groupprops.subwiki.org/wiki/Inner-centralizing\\_automorphism](https://groupprops.subwiki.org/wiki/Inner-centralizing_automorphism)
17. <https://math.stackexchange.com/questions/2519372/number-of-conjugacy-classes-for-the-modular-group>
18.  $\text{GL}_n(K)$  for  $K$  a skew field
19. <https://arxiv.org/abs/1212.6157>, <https://arxiv.org/abs/0708.1608>, [https://en.wikipedia.org/wiki/Wild\\_problem](https://en.wikipedia.org/wiki/Wild_problem), <https://www.google.com/search?q=matrix+pair+problem>, <https://arxiv.org/abs/2007.09242>, <https://mathoverflow.net/questions/291815/rational-canonical-form-over-mathbbz-pk-mathbbz>
20. <https://link.springer.com/book/10.1007/978-981-13-2895-4>
21. <https://ysharifi.wordpress.com/2022/09/14/automorphisms-of-dihedral-groups/>
22. [https://en.wikipedia.org/wiki/PSL\(2,7\)](https://en.wikipedia.org/wiki/PSL(2,7))
23. <https://arxiv.org/abs/2304.08617>
24. <https://johncarlosbaez.wordpress.com/2016/03/22/the-inolute-of-a-cubical-parabola/#comment-78884>

25. <https://arxiv.org/abs/0904.1876>
26. finite subgroups of  $SU(2)$ , and viewing them as groups of rotations and such
27. <https://arxiv.org/abs/1201.2363>
28. [https://ncatlab.org/nlab/show/group+extension#Schreier Theory](https://ncatlab.org/nlab/show/group+extension#Schreier_Theory), <https://ncatlab.org/nlab/show/nonabelian+cohomology>, <https://ncatlab.org/nlab/show/nonabelian+group+cohomology>
29. [https://en.wikipedia.org/wiki/Fibonacci\\_group](https://en.wikipedia.org/wiki/Fibonacci_group)
30. Study the functoriality properties of  $G \mapsto \text{Aut}(G)$  via functoriality of ends
31. Is  $\sum_{[g] \in \text{Cl}(G)} \frac{1}{|g|}$  an interesting invariant of  $G$ ?
32. Idempotent endomorphism  $f: A \rightarrow A$  is the same as a decomposition  $A \cong B \oplus C$  via  $B \cong \text{Im}(f)$  and  $C \cong \text{Ker}(f)$ .
  - (a) <https://mathstrek.blog/2015/03/02/idempotents-and-decomposition/>
33. <https://math.stackexchange.com/questions/34271/order-of-f-general-and-special-linear-groups-over-finite-fields>

Linear Algebra:

1. Size of conjugacy class  $[A]$  of  $A \in \text{GL}_n(\mathbb{F}_q)$  is given by  $\#\text{GL}_n(\mathbb{F}_q)$  divided by the centralizer  $Z_{\text{GL}_n(\mathbb{F}_q)}(A)$  of  $A$  in  $\text{GL}_n(\mathbb{F}_q)$ , whose order is given by

$$\begin{aligned} \#\text{Z}_{\text{GL}_n(\mathbb{F}_q)}(A) &= \prod_{i=1}^k \#\text{GL}_{r_i}(\mathbb{F}_q) \\ &= q^{\sum_{i=1}^k \binom{r_i}{2}} \prod_{i=1}^k \prod_{j=0}^{r_i-1} (q^{r_i-j} - 1) \end{aligned}$$

if  $A$  is diagonalisable with eigenvalues  $\lambda_1, \dots, \lambda_k$  having multiplicities  $r_1, \dots, r_k$ . More generally, see [https://groupprops.subwiki.org/wiki/Conjugacy\\_class\\_size\\_formula\\_in\\_general\\_linear\\_group\\_over\\_a\\_finite\\_field](https://groupprops.subwiki.org/wiki/Conjugacy_class_size_formula_in_general_linear_group_over_a_finite_field)

2. [https://en.wikipedia.org/wiki/Semilinear\\_map](https://en.wikipedia.org/wiki/Semilinear_map)
3. conjugacy for  $\text{GL}_n(\mathbb{F}_q)$ , <https://mathoverflow.net/a/104457>

4. [https://en.wikipedia.org/wiki/Dieudonné\\_determinant](https://en.wikipedia.org/wiki/Dieudonné_determinant), <https://ncatlab.org/nlab/show/Dieudonné+determinant#Dieudonne>
5. <https://ncatlab.org/nlab/show/Pfaffian>
6. <https://math.stackexchange.com/questions/1715249/the-number-of-subspaces-over-a-finite-field>
7. <https://math.stackexchange.com/questions/70801/how-many-k-dimensional-subspaces-are-there-in-n-dimensional-vector-space-over>
8. [https://en.wikipedia.org/wiki/Gaussian\\_binomial\\_coefficient](https://en.wikipedia.org/wiki/Gaussian_binomial_coefficient)
9. [https://en.wikipedia.org/wiki/List\\_of\\_q-analogs](https://en.wikipedia.org/wiki/List_of_q-analogs)

Noncommutative Algebra:

1. <https://arxiv.org/abs/1608.08140>
2. <https://arxiv.org/abs/2401.12884>
3. <https://ncatlab.org/nlab/show/dihedral+homology>
4. <https://www.sciencedirect.com/science/article/pii/0022404995000836>
5. <https://arxiv.org/abs/2008.11569>, <https://www.lakeheadu.ca/sites/default/files/uploads/77/docs/Cox%20Daniel.pdf>

Commutative Algebra:

1. If  $M \in \text{Pic}(R)$ , then  $\text{Aut}(M) \cong R^\times$ .
2. <https://math.stackexchange.com/questions/637918/>
3. <https://categorytheory.zulipchat.com/#narrow/stream/411257-theory.3A-mathematics/topic/Big.20Witt.20ring>
4. <https://math.stackexchange.com/questions/535623/how-many-irreducible-factors-does-xn-1-have-over-finite-field>
5. Derivations between morphisms of  $R$ -algebras, after <https://mathoverflow.net/questions/434488>

(a) Namely, a derivation from a morphism  $f: A \rightarrow B$  of  $R$ -algebras to a morphism  $g: A \rightarrow B$  of  $R$ -algebras is a map  $D: B \rightarrow B$  such that we have

$$D(ab) = g(a)D(b) + D(a)f(b)$$

for each  $a, b \in B$ .

Hyper Algebra:

1. <https://arxiv.org/abs/2205.02362>
2. [http://www.numdam.org/item/SD\\_1959-1960\\_\\_13\\_1\\_A9\\_0/](http://www.numdam.org/item/SD_1959-1960__13_1_A9_0/)
3. <https://www.worldscientific.com/worldscibooks/10.1142/13652#t=aboutBook>

Coalgebra:

1. <https://mathoverflow.net/questions/483668/textrepd-4-and-its-three-fiber-functors>

Topological Algebra:

1. [https://golem.ph.utexas.edu/category/2014/08/holy\\_crap\\_do\\_you\\_know\\_what\\_a\\_c.html](https://golem.ph.utexas.edu/category/2014/08/holy_crap_do_you_know_what_a_c.html)
2. <https://categorytheory.zulipchat.com/#narrow/channel/411257-theory.3A-mathematics/topic/topological.20rings.20and.20fields>
3. <https://mathoverflow.net/q/477757>
4. <https://math.stackexchange.com/questions/2593556/galois-theory-for-topological-fields>

Differential Graded Algebras:

1. <https://mathoverflow.net/questions/476150/constructing-an-adjunction-between-algebras-and-differential-grade-d-algebras>

Topology:

1. <https://arxiv.org/abs/2507.18418>
2. Topologies on  $\mathcal{P}(\mathcal{P}(X))$ , <https://mathoverflow.net/questions/496630/topological-analogues-of-gromov-hausdorff-convergence>

3. <https://mathoverflow.net/questions/255912/what-is-the-structure-associated-to-almost-everywhere-convergence>
4. <https://arxiv.org/abs/2504.12965>
5. <https://mathoverflow.net/questions/485669/exponential-law-for-topological-spaces-for-the-topology-of-pointwise-convergence> and comments therein
6. This paper has some cool references on convergence spaces: <https://arxiv.org/abs/2410.18245>
7. <https://arxiv.org/abs/2402.12316>
8. Write about the 6-functor formalism for sheaves on topological spaces and for topological stacks, with lots of examples.
  - (a) MO question titled *6-functor formalism for topological stacks*: <https://mathoverflow.net/q/471758>

Measure Theory:

1. <https://mathoverflow.net/questions/126994/beck-chevalley-for-measures>
2. <https://mathoverflow.net/questions/483726>
3. [https://en.wikipedia.org/wiki/Valuation\\_%28measure\\_theory%29](https://en.wikipedia.org/wiki/Valuation_%28measure_theory%29)
4. There's a theorem saying that there does not exist an infinite-dimensional "Lebesgue" measure, i.e. (from [https://en.wikipedia.org/wiki/Infinite-dimensional\\_Lebesgue\\_measure](https://en.wikipedia.org/wiki/Infinite-dimensional_Lebesgue_measure)):

Let  $X$  be an infinite-dimensional, separable Banach space. Then, the only locally finite and translation invariant Borel measure  $\mu$  on  $X$  is a trivial measure. Equivalently, there is no locally finite, strictly positive, and translation invariant measure on  $X$ .

What kind of measures exist/not exist that satisfy all conditions above except being locally finite?

5. <https://ncatlab.org/nlab/show/categories+of+measure+theory>
6. Functions  $f_!$ ,  $f^*$ , and  $f_*$  between spaces of (probability) measures on probability/measurable spaces, mimicking how a map of sets  $f: X \rightarrow Y$  induces morphisms of sets  $f_!$ ,  $f^*$ , and  $f_*$  between  $\mathcal{P}(X)$  and  $\mathcal{P}(Y)$ .

7. Analogies between representable presheaves and the Yoneda lemma on the one hand and Dirac probability measures on the other hand

- (a) Universal property of the embedding of a space  $X$  into the space of probability measures on  $X$
- (b) Same question but for distributions
- (c) non-symmetric metric on space of probability measures where we define  $d(\mu, \nu)$  to be the measure given by

$$U \mapsto \int_U \rho_\mu d\nu,$$

where  $\rho_\mu$  is the probability density of  $\mu$ . Can we make this idea work?

- 8. <https://arxiv.org/abs/0801.2250>
- 9. <https://mathoverflow.net/questions/325861>

*In particular, I came across a PhD thesis by Martial Aguech. I thought it was interesting because it explicitly investigated the geodesics of Wasserstein space to produce solutions to a type of parabolic PDE.*

Probability Theory:

- 1. [https://en.wikipedia.org/wiki/Wiener\\_sausage](https://en.wikipedia.org/wiki/Wiener_sausage)
- 2. <https://link.springer.com/book/10.1007/978-3-319-20828-2>
- 3. <https://arxiv.org/abs/2406.10676>
- 4. Lévy's forgery theorem
- 5. <https://www.epatters.org/wiki/stats-ml/categorical-probability-theory>
- 6. <https://ncatlab.org/nlab/show/category-theoretic+approaches+to+probability+theory>
- 7. Categorical probability theory
- 8. [https://golem.ph.utexas.edu/category/2024/08/introduction\\_to\\_categorical\\_pr.html](https://golem.ph.utexas.edu/category/2024/08/introduction_to_categorical_pr.html)
- 9. <https://arxiv.org/abs/1109.1880>

10. Connection between fractional differential operators and stochastic processes with jumps

Statistics:

1. <https://towardsdatascience.com/t-test-from-application-to-theory-5e5051b0f9dc>

Metric Spaces:

1. Lawvere metric spaces: object of  $\mathcal{V}$ -natural transformations corresponds to  $\inf(d(f(x), g(x)))$ .
2. Does the assignment  $d(x, y) \mapsto d(x, y)/(1 + d(x, y))$  constructing a bounded metric from a metric be given a universal property?
3. Explore Lawvere metric spaces in a comprehensive manner
4. metric  $\text{lcm}(x, y)/\text{gcd}(x, y)$  on  $\mathbb{N}$ , <https://mathoverflow.net/questions/461588/>. What shape do balls on  $\mathbb{N} \times \mathbb{N}$  have with respect to this metric?
5. [https://golem.ph.utexas.edu/category/2023/05/metric\\_spaces\\_as\\_enriched\\_categories\\_ii.html](https://golem.ph.utexas.edu/category/2023/05/metric_spaces_as_enriched_categories_ii.html)
6. Simon Willerton's work on the Legendre–Fenchel transform:
  - (a) [https://golem.ph.utexas.edu/category/2014/04/enrichment\\_and\\_the\\_legendrefen.html](https://golem.ph.utexas.edu/category/2014/04/enrichment_and_the_legendrefen.html)
  - (b) [https://golem.ph.utexas.edu/category/2014/05/enrichment\\_and\\_the\\_legendrefen\\_1.html](https://golem.ph.utexas.edu/category/2014/05/enrichment_and_the_legendrefen_1.html)
  - (c) <https://arxiv.org/abs/1501.03791>

Special Functions:

1. [https://en.wikipedia.org/wiki/Dickson\\_polynomial](https://en.wikipedia.org/wiki/Dickson_polynomial)

$p$ -Adic Analysis:

1. <https://arxiv.org/abs/2503.08909>
2. Analysis of functions  $\mathbb{Z}_p \rightarrow \mathbb{Q}_q$ ,  $\mathbb{Q}_p \rightarrow \mathbb{Q}_q$ ,  $\mathbb{Z}_p \rightarrow \mathbb{C}_q$ , etc.
  - (a) <https://siegelmaxwellc.wordpress.com/publications-re-prints/>

Partial Differential Equations:

1. Moduli of PDEs

- (a) <https://arxiv.org/abs/2312.05226>, <https://arxiv.org/abs/2406.16825>
- (b) <https://arxiv.org/abs/2304.08671>, <https://arxiv.org/abs/2404.07931>
- (c) <https://arxiv.org/abs/2507.07937>
- 2. [https://en.wikipedia.org/wiki/Homotopy\\_principle](https://en.wikipedia.org/wiki/Homotopy_principle)
- 3. <https://mathoverflow.net/questions/125166/wild-solutions-of-the-heat-equation-how-to-graph-them>
- 4. <https://math.stackexchange.com/questions/2112841/difference-between-linear-semilinear-and-quasilinear-pdes/5036699#5036699>
- 5. Proof of the smoothing property of the heat equation via:
  - (a) Feynman–Kac formula
  - (b) Radon–Nikodym + Wiener process has Gaussian as PDF
  - (c) Convolution of locally integrable with smooth is smooth
- 6. Geometry of PDEs:
  - (a) <https://mathoverflow.net/questions/457268/pdes-and-algebraic-varieties>
  - (b) Can we build a kind of algebraic geometry of PDEs starting with the notion of the zero locus of a differential operator?
    - i. <https://ncatlab.org/nlab/show/difiety>

Functional Analysis:

- 1. [https://www.numdam.org/item/SE\\_1957-1958\\_\\_1\\_\\_A3\\_0/](https://www.numdam.org/item/SE_1957-1958__1__A3_0/)
- 2. <https://thenumb.at/Functions-are-Vectors/>
- 3. Tate vector spaces
- 4. Analytic sheaves, <https://mathoverflow.net/questions/484408/literature-on-fr%c3%a9chet-quasi-coherent-sheaves>
- 5. <https://mathscinet.ams.org/mathscinet/article?mr=1257171>
- 6. Vidav–Palmer theorem
- 7. In the Hilbert space  $\ell^2(\mathbb{N}; \mathbb{C})$ , the operator  $(x_n)_{n \in \mathbb{N}} \mapsto (x_{n+1})_{n \in \mathbb{N}}$  admits  $(x_n)_{n \in \mathbb{N}} \mapsto (0, x_0, x_1, \dots)$  as its adjoint.

8. <https://arxiv.org/abs/2110.06300>

Lie algebras:

1. Pre-Lie algebras
2. Post-Lie algebras
3. <https://arxiv.org/abs/2504.05929>

Modular Representation Theory:

1. [https://en.wikipedia.org/wiki/Deligne%20%93Lusztig\\_theory](https://en.wikipedia.org/wiki/Deligne%20%93Lusztig_theory)
2. <https://math.stackexchange.com/questions/167979/representation-of-cyclic-group-over-finite-field>
3. <https://math.stackexchange.com/questions/153429/irreducible-representations-of-a-cyclic-group-over-a-field-of-prime-order>

Homotopy theory:

1. <https://mathoverflow.net/questions/495229>
2. <https://ncatlab.org/nlab/show/Moore+path+category>, <https://mathoverflow.net/questions/486905/has-the-path-category-of-a-topological-space-been-studied>/487212#487212
3. <https://ncatlab.org/nlab/show/group+actions+on+spheres>,  
<https://www.maths.ed.ac.uk/~v1ranick/papers/wall7.pdf>,  
<https://math.stackexchange.com/questions/1575798/which-groups-act-freely-on-sn>, <https://arxiv.org/abs/math/0212280>.
4. Pascal's triangle via homology of  $n$ -tori, [https://topospaces.subwiki.org/wiki/Homology\\_of\\_torus](https://topospaces.subwiki.org/wiki/Homology_of_torus)
5. Conditions on morphisms of spaces  $f: X \rightarrow Y$  such that  $f^*: [Y, K] \rightarrow [X, K]$  or  $f_*: [K, X] \rightarrow [K, Y]$  are injective/surjective (so, epi/monomorphisms in  $\text{Ho}(\mathcal{T})$ ) or other conditions.

Algebraic Geometry:

1. Galois points, [https://bdtd.ibict.br/vufind/Record/USP\\_c5e6638812a74657c40fc402a894514](https://bdtd.ibict.br/vufind/Record/USP_c5e6638812a74657c40fc402a894514)
2. <https://arxiv.org/abs/2407.09256>

Differential Geometry:

1. [https://en.wikipedia.org/wiki/Spherical\\_3-manifold](https://en.wikipedia.org/wiki/Spherical_3-manifold)
2. functor of points approach to differential geometry

Number Theory:

1. <https://math.stackexchange.com/questions/10233/uses-of-quadratic-reciprocity-theorem/10719#10719>
2. <https://mathoverflow.net/questions/120067/what-do-theta-functions-have-to-do-with-quadratic-reciprocity>

Classical Mechanics:

1. Koopman–von Neumann formalism
2. Relativistic Lagrangian and Hamiltonian mechanics

Quantum Mechanics:

1. <https://ncatlab.org/nlab/show/geometrical+formulation+of+quantum+mechanics>

Quantum Field Theory:

1. <https://arxiv.org/abs/2309.15913> and <https://arxiv.org/abs/2311.09284>
2. The current ongoing work on higher gauge theory, specially Christian Saemann's
3. The recent work about determining the value of the strong coupling constant in the long-distance range, some pointers and keywords for this are available at [this scientific american article](#).

Combinatorics:

1. Catalan numbers, <https://mathstrek.blog/2012/02/19/power-series-and-generating-functions-ii-formal-power-series/>

Other:

1. <https://arxiv.org/abs/2202.00084>
2. Are sedenions and higher useful for anything?
3. <https://mathstodon.xyz/@pschwahn/113388126188923908>

4. Tambara functors, <https://arxiv.org/abs/2410.23052>
5. 2-vector spaces
6. 2-term chain complexes. They form a 2-category and middle-four exchange holds, the proof using the fact that we have

$$h_1 \circ \alpha + \beta \circ g_2 = k_1 \circ \alpha + \beta \circ f_2,$$

which uses the chain homotopy identities

$$\begin{aligned} d_V \circ \alpha &= g_2 - f_2, \\ -\beta \circ d_V &= h_1 - k_1. \end{aligned}$$

Can we modify this to work for usual chain complexes, seeking an answer to <https://mathoverflow.net/questions/424268>? What seems to make things go wrong in that case is that the chain homotopy identities are replaced with

$$\begin{aligned} \alpha_{n+1} \circ d_n^V + d_{n-1}^W \circ \alpha_n &= g_n - f_n, \\ \beta_{n+1} \circ d_n^V + d_{n-1}^W \circ \beta_n &= k_n - h_n. \end{aligned}$$

7. <https://arxiv.org/abs/1402.2600>
8. <https://grossack.site/blog>
9. Classifying space of  $\mathbb{Q}_p$
10. <https://www.valth.eu/proc.htm>
11. Construction of  $\mathbb{R}$  via slopes:
  - (a) <http://maths.mq.edu.au/~street/EffR.pdf>
  - (b) <https://arxiv.org/abs/math/0301015>
  - (c) Pierre Colmez's comment "Et si on remplace  $\mathbb{Z}$  par  $\mathbb{Q}$ , on obtient les adèles."
  - (d) I wonder if one could apply an analogue of this construction to the sphere spectrum and obtain a kind of spectral version of the real numbers, as in e.g. following the spirit of <https://mathoverflow.net/questions/443018>.
12. <https://arxiv.org/abs/2406.04936>
13. <https://mathoverflow.net/a/471510>

14. <https://mathoverflow.net/questions/279478/the-category-of-span-enriched-categories-2-segal-spaces/448523#448523>
15. The works of David Kern, <https://dskern.github.io/writings>
16. <https://qchu.wordpress.com/>
17. <https://aroundtoposes.com/>
18. <https://ncatlab.org/nlab/show/essentially+surjective+and+full+functor>
19. <https://mathoverflow.net/questions/415363/objects-whose-representable-presheaf-is-a-fibration>
20. <https://mathoverflow.net/questions/460146/universal-property-of-isbell-duality>
21. [\(Isbell conjugacy and the reflexive completion\)](http://www.tac.mta.ca/tac/volumes/36/12/36-12abs.html)
22. <https://ncatlab.org/nlab/show/enrichment+versus+internalisation>
23. The works of Philip Saville, <https://philipsaville.co.uk/>
24. [https://golem.ph.utexas.edu/category/2024/02/from\\_cartesian\\_to\\_symmetric\\_mo.html](https://golem.ph.utexas.edu/category/2024/02/from_cartesian_to_symmetric_mo.html)
25. [\(One-object lax transformations\)](https://mathoverflow.net/q/463855)
26. <https://ncatlab.org/nlab/show/analytic+completion+of+a+ring>
27. [https://en.wikipedia.org/wiki/Quaternionic\\_analysis](https://en.wikipedia.org/wiki/Quaternionic_analysis)
28. [\(The Norm Functor over Schemes\)](https://arxiv.org/abs/2401.15051)
29. <https://mathoverflow.net/questions/407291/> (Adjunctions with respect to profunctors)
30. [\(Prof is free completion of Cats under right extensions\)](https://mathoverflow.net/a/462726)
31. there's some cool stuff in <https://arxiv.org/abs/2312.00990> (Polynomial Functors: A Mathematical Theory of Interaction), e.g. on cofunctors.

32. <https://ncatlab.org/nlab/show/adjoint+lifting+theorem>
33. <https://ncatlab.org/nlab/show/Gabriel%20%93Ulmer+du+ality>

General TODO:

1. <https://arxiv.org/abs/2108.11952>
2. <https://mathoverflow.net/questions/483243/is-there-a-theory-of-completions-of-semirings-similar-to-i-adic-completions-of>
3. <https://mathoverflow.net/questions/9218/probabilistic-prroofs-of-analytic-facts>
4. <https://x.com/cihanpoststhms>
5. Special graded rings, [\(a\) https://mathoverflow.net/questions/403448/in-search-of-lost-graded-rings](https://mathoverflow.net/questions/403448/in-search-of-lost-graded-rings)
  - (a) <https://arxiv.org/abs/1209.5122>
6. Counterexamples in category theory
7. <https://math.stackexchange.com/questions/279347/counterexample-math-books>
8. Browse MO questions/answers for interesting ideas/topics
9. Change Longrightarrow to Rightarrow where appropriate
10. Try to minimize the amount of footnotes throughout the project.  
There should be no long footnotes.

## Appendices

### 15.A Other Chapters

#### Preliminaries

1. Introduction
2. A Guide to the Literature

#### Sets

3. Sets

#### 4. Constructions With Sets

5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

**Relations**

- 8. Relations
- 9. Constructions With Relations
- 10. Conditions on Relations

**Categories**

- 11. Categories
- 12. Presheaves and the Yoneda Lemma

**Monoidal Categories**

- 13. Constructions With Monoidal Categories

**Bicategories**

- 14. Types of Morphisms in Bicategories

**Extra Part**

- 15. Notes

# Bibliography

- [MO 119454] user30818. *Category and the axiom of choice*. MathOverflow. URL: <https://mathoverflow.net/q/119454> (cit. on p. 790).
- [MO 321971] Ivan Di Liberti. *Characterization of pseudo monomorphisms and pseudo epimorphisms in Cat*. MathOverflow. URL: <https://mathoverflow.net/q/321971> (cit. on pp. 18, 801).
- [MO 382264] Neil Strickland. *Proof that a cartesian category is monoidal*. MathOverflow. URL: <https://mathoverflow.net/q/382264> (cit. on p. 951).
- [MO 460693] Tim Campion. *Answer to “Existence and characterisations of left Kan extensions and liftings in the bicategory of relations I”*. MathOverflow. URL: <https://mathoverflow.net/q/460693> (cit. on p. 633).
- [MO 461592] Emily. *Existence and characterisations of left Kan extensions and liftings in the bicategory of relations II*. MathOverflow. URL: <https://mathoverflow.net/q/461592> (cit. on pp. 17, 635, 636).
- [MO 467527] Emily. *What are the 2-categorical mono/epimorphisms in the 2-category of relations?* MathOverflow. URL: <https://mathoverflow.net/q/467527> (cit. on p. 17).
- [MO 468121a] Emily. *Characterisations of functors  $F$  such that  $F^*$  or  $F_*$  is [property], e.g. faithful, conservative, etc.* MathOverflow. URL: <https://mathoverflow.net/q/468125> (cit. on pp. 17, 18, 787, 788, 794, 795, 797, 803, 806, 809).
- [MO 468121b] Emily. *Looking for a nice characterisation of functors  $F$  whose precomposition functor  $F^*$  is full.* MathOverflow. URL: <https://mathoverflow.net/q/468121> (cit. on pp. 17, 781).
- [MO 468334] Emily. *Is a pseudomonic and pseudoepic functor necessarily an equivalence of categories?* MathOverflow. URL: <https://mathoverflow.net/q/468334> (cit. on pp. 18, 803).

- [MO 484647] **Zhen Lin.** *Reference for Basic Multicategory Theory.* MathOverflow. URL: <https://mathoverflow.net/q/484647> (cit. on p. 19).
- [MO 490557] **Maxime Ramzi.** *Answer to “What theorems or insights are well known for  $\infty$ -categories but not well known for categories?”* MathOverflow. URL: <https://mathoverflow.net/a/490557> (cit. on p. 18).
- [MO 494959] **Emily.** *Gaps in the category theory literature you’d like to see filled.* MathOverflow. URL: <https://mathoverflow.net/q/494959> (cit. on p. 18).
- [MO 495003] **Fosco.** *Answer to “Gaps in the category theory literature you’d like to see filled”.* MathOverflow. URL: <https://mathoverflow.net/a/495003> (cit. on p. 22).
- [MO 497309] **Noah Snyder.** *Answer to “Gaps in the category theory literature you’d like to see filled”.* MathOverflow. URL: <https://mathoverflow.net/q/497309> (cit. on p. 22).
- [MO 497419] **Nick Hu.** *Answer to “Gaps in the category theory literature you’d like to see filled”.* MathOverflow. URL: <https://mathoverflow.net/q/497419> (cit. on p. 22).
- [MO 64365] **Giorgio Mossa.** *Natural transformations as categorical homotopies.* MathOverflow. URL: <https://mathoverflow.net/q/64365> (cit. on p. 825).
- [MSE 1465107] **kilian.** *Equivalence of categories and axiom of choice.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/1465107> (cit. on p. 790).
- [MSE 2096272] **Akiva Weinberger.** *Is composition of two transitive relations transitive? If not, can you give me a counterexample?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2096272> (cit. on p. 714).
- [MSE 267365] **J. B.** *Show that the powerset partial order is a cartesian closed category.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/267365> (cit. on p. 181).
- [MSE 267469] **Zhen Lin.** *Show that the powerset partial order is a cartesian closed category.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/267469> (cit. on p. 144).
- [MSE 2719059] **Vinny Chase.**  *$\mathcal{P}(X)$  with symmetric difference as addition as a vector space over  $\mathbb{Z}_2$ .* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2719059> (cit. on p. 160).

- [MSE 2855868] **Qiaochu Yuan.** *Is the category of pointed sets Cartesian closed?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2855868> (cit. on pp. 336, 346).
- [MSE 350788] **Qiaochu Yuan.** *Mono's and epi's in the category Rel?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/350788> (cit. on pp. 600, 616).
- [MSE 733161] **Stefan Hamcke.** *Precomposition with a faithful functor.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/733161> (cit. on p. 785).
- [MSE 733163] **Zhen Lin.** *Precomposition with a faithful functor.* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/733163> (cit. on p. 774).
- [MSE 749304] **Zhen Lin.** *If the functor on presheaf categories given by precomposition by  $F$  is ff, is  $F$  full? faithful?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/749304> (cit. on p. 785).
- [MSE 884460] **Martin Brandenburg.** *Why are the category of pointed sets and the category of sets and partial functions “essentially the same”?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/884460> (cit. on p. 337).
- [Adá+01] Jiří Adámek, Robert El Bashir, Manuela Sobral, and Jiří Velebil. “On Functors Which Are Lax Epimorphisms”. In: *Theory Appl. Categ.* 8 (2001), pp. 509–521. ISSN: 1201-561X (cit. on pp. 774, 781, 785, 899).
- [Bor94] Francis Borceux. *Handbook of Categorical Algebra I.* Vol. 50. Encyclopedia of Mathematics and its Applications. Basic Category Theory. Cambridge University Press, Cambridge, 1994, pp. xvi+345. ISBN: 0-521-44178-1 (cit. on p. 820).
- [BS10] John C. Baez and Michael Shulman. “Lectures on  $n$ -Categories and Cohomology”. In: *Towards higher categories.* Vol. 152. IMA Vol. Math. Appl. Springer, New York, 2010, pp. 1–68. DOI: [10.1007/978-1-4419-1524-5\\_1](https://doi.org/10.1007/978-1-4419-1524-5_1). URL: [https://doi.org/10.1007/978-1-4419-1524-5\\_1](https://doi.org/10.1007/978-1-4419-1524-5_1) (cit. on pp. 29, 30, 780).
- [Cie97] Krzysztof Ciesielski. *Set Theory for the Working Mathematician.* Vol. 39. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1997, pp. xii+236. ISBN: 0-521-59441-3; 0-521-59465-0. DOI: [10.1017/CBO9781139173131](https://doi.org/10.1017/CBO9781139173131). URL: <https://doi.org/10.1017/CBO9781139173131> (cit. on p. 101).

- [DFH75] Aristide Deleanu, Armin Frei, and Peter Hilton. “Idempotent Triples and Completion”. In: *Math. Z.* 143 (1975), pp. 91–104. ISSN: 0025-5874,1432-1823. DOI: [10.1007/BF01173053](https://doi.org/10.1007/BF01173053). URL: <https://doi.org/10.1007/BF01173053> (cit. on p. 794).
- [DS06] Eduardo Dubuc and Ross Street. “Dinatural Transformations”. In: *Reports of the Midwest Category Seminar IV*. Springer. 2006, pp. 126–137 (cit. on p. 19).
- [Fre09] Jonas Frey. *On the 2-Categorical Duals of (Full and) Faithful Functors*. <https://citeseervx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4c289321d622f8fcf947e7a7cf1bdf75c95ca33>. Archived at <https://web.archive.org/web/20240331195546/https://citeseervx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4c289321d622f8fcf947e7a7cf1bdf75c95ca33>. July 2009. URL: <https://citeseervx.ist.psu.edu/document?repid=rep1%5C&type=pdf%5C&doi=4c289321d622f8fcf947e7a7cf1bdf75c95ca33> (cit. on pp. 774, 785).
- [GJO24] Nick Gurski, Niles Johnson, and Angélica M. Osorno. “The Symmetric Monoidal 2-Category of Permutative Categories”. In: *High. Struct.* 8.1 (2024), pp. 244–320. ISSN: 2209-0606 (cit. on p. 19).
- [Isb68] John R. Isbell. “Epimorphisms and Dominions. III”. In: *Amer. J. Math.* 90 (1968), pp. 1025–1030. ISSN: 0002-9327,1080-6377. DOI: [10.2307/2373286](https://doi.org/10.2307/2373286). URL: <https://doi.org/10.2307/2373286> (cit. on p. 796).
- [JS17] Theo Johnson-Freyd and Claudia Scheimbauer. “(Op)lax Natural Transformations, Twisted Quantum Field Theories, and “Even Higher” Morita Categories”. In: *Adv. Math.* 307 (2017), pp. 147–223. ISSN: 0001-8708,1090-2082. DOI: [10.1016/j.aim.2016.11.014](https://doi.org/10.1016/j.aim.2016.11.014). URL: <https://doi.org/10.1016/j.aim.2016.11.014> (cit. on p. 841).
- [JY21] Niles Johnson and Donald Yau. *2-Dimensional Categories*. Oxford University Press, Oxford, 2021, pp. xix+615. ISBN: 978-0-19-887138-5; 978-0-19-887137-8. DOI: [10.1093/oso/9780198871378.001.0001](https://doi.org/10.1093/oso/9780198871378.001.0001). URL: <https://doi.org/10.1093/oso/9780198871378.001.0001> (cit. on pp. 3, 9, 958).
- [Low15] Zhen Lin Low. *Notes on Homotopical Algebra*. Nov. 2015. URL: <https://zll22.user.srccf.net/writing/homotopical-algebra/2015-11-10-Main.pdf> (cit. on p. 785).
- [Lur25] Jacob Lurie. *Kerodon*. <https://kerodon.net>. 2025 (cit. on p. 10).

- [Mac98] Saunders Mac Lane. *Categories for the Working Mathematician*. Second. Vol. 5. Graduate Texts in Mathematics. Springer-Verlag, New York, 1998, pp. xii+314. ISBN: 0-387-98403-8 (cit. on p. 9).
- [nLa25a] nLab Authors. *Groupoid*. <https://ncatlab.org/nlab/show/groupoid>. Oct. 2025 (cit. on p. 791).
- [nLa25b] nLab Authors. *Interactions of Images and Pre-images with Unions and Intersections*. <https://ncatlab.org/nlab/show/interactions+of+images+and+pre-images+with+unions+and+intersections>. Oct. 2025 (cit. on p. 243).
- [nLab23] nLab Authors. *Skeleton*. 2025. URL: <https://ncatlab.org/nlab/show/skeleton> (cit. on pp. 734, 735).
- [Pro25a] Proof Wiki Contributors. *Cartesian Product Distributes Over Set Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Cartesian\\_Product\\_Distributes\\_over\\_Set\\_Difference](https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Set_Difference) (cit. on p. 54).
- [Pro25b] Proof Wiki Contributors. *Cartesian Product Distributes Over Symmetric Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Cartesian\\_Product\\_Distributes\\_over\\_Symmetric\\_Difference](https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Symmetric_Difference) (cit. on p. 54).
- [Pro25c] Proof Wiki Contributors. *Cartesian Product Distributes Over Union — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Cartesian\\_Product\\_Distributes\\_over\\_Union](https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Union) (cit. on p. 54).
- [Pro25d] Proof Wiki Contributors. *Cartesian Product of Intersections — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Cartesian\\_Product\\_of\\_Intersections](https://proofwiki.org/wiki/Cartesian_Product_of_Intersections) (cit. on p. 54).
- [Pro25e] Proof Wiki Contributors. *Characteristic Function of Intersection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Characteristic\\_Function\\_of\\_Intersection](https://proofwiki.org/wiki/Characteristic_Function_of_Intersection) (cit. on p. 144).
- [Pro25f] Proof Wiki Contributors. *Characteristic Function of Set Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Characteristic\\_Function\\_of\\_Set\\_Difference](https://proofwiki.org/wiki/Characteristic_Function_of_Set_Difference) (cit. on p. 150).
- [Pro25g] Proof Wiki Contributors. *Characteristic Function of Symmetric Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Characteristic\\_Function\\_of\\_Symmetric\\_Difference](https://proofwiki.org/wiki/Characteristic_Function_of_Symmetric_Difference) (cit. on p. 159).

- [Pro25h] Proof Wiki Contributors. *Characteristic Function of Union — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Characteristic\\_Function\\_of\\_Union](https://proofwiki.org/wiki/Characteristic_Function_of_Union) (cit. on p. 139).
- [Pro25i] Proof Wiki Contributors. *Complement of Complement — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Complement\\_of\\_Complement](https://proofwiki.org/wiki/Complement_of_Complement) (cit. on p. 152).
- [Pro25j] Proof Wiki Contributors. *Complement of Preimage equals Preimage of Complement — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Complement\\_of\\_Preimage\\_equals\\_Preimage\\_of\\_Complement](https://proofwiki.org/wiki/Complement_of_Preimage_equals_Preimage_of_Complement) (cit. on p. 227).
- [Pro25k] Proof Wiki Contributors. *Condition For Mapping from Quotient Set To Be A Surjection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Condition\\_for\\_Mapping\\_from\\_Quotient\\_Set\\_to\\_be\\_Surjection](https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Surjection) (cit. on p. 725).
- [Pro25l] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be An Injection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Condition\\_for\\_Mapping\\_from\\_Quotient\\_Set\\_to\\_be\\_Injection](https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Injection) (cit. on p. 725).
- [Pro25m] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be Well-Defined — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Condition\\_for\\_Mapping\\_from\\_Quotient\\_Set\\_to\\_be\\_Well-Defined](https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Well-Defined) (cit. on p. 724).
- [Pro25n] Proof Wiki Contributors. *De Morgan's Laws (Set Theory) — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/De\\_Morgan%5C27s\\_Laws\\_\(Set\\_Theory\)](https://proofwiki.org/wiki/De_Morgan%5C27s_Laws_(Set_Theory)) (cit. on pp. 148, 152).
- [Pro25o] Proof Wiki Contributors. *De Morgan's Laws (Set Theory)/Set Difference/Difference with Union — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/De\\_Morgan%5C27s\\_Laws\\_\(Set\\_Theory\)/Set\\_Difference/Difference\\_with\\_Union](https://proofwiki.org/wiki/De_Morgan%5C27s_Laws_(Set_Theory)/Set_Difference/Difference_with_Union) (cit. on p. 149).
- [Pro25p] Proof Wiki Contributors. *Equivalence of Definitions of Symmetric Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Equivalence\\_of\\_Definitions\\_of\\_Symmetric\\_Difference](https://proofwiki.org/wiki/Equivalence_of_Definitions_of_Symmetric_Difference) (cit. on p. 158).
- [Pro25q] Proof Wiki Contributors. *Image of Intersection Under Mapping — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Image\\_of\\_Intersection\\_under\\_Mapping](https://proofwiki.org/wiki/Image_of_Intersection_under_Mapping) (cit. on pp. 145, 217, 218).
- [Pro25r] Proof Wiki Contributors. *Image of Set Difference Under Mapping — Proof Wiki*. 2025. URL: <https://proofwiki.org/>

- wiki/Image\_of\_Set\_Difference\_under\_Mapping (cit. on pp. 150, 217).
- [Pro25s] Proof Wiki Contributors. *Image of Union Under Mapping — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Image\\_of\\_Union\\_under\\_Mapping](https://proofwiki.org/wiki/Image_of_Union_under_Mapping) (cit. on pp. 139, 217, 219).
- [Pro25t] Proof Wiki Contributors. *Intersection Distributes Over Symmetric Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_Distributes\\_over\\_Symmetric\\_Difference](https://proofwiki.org/wiki/Intersection_Distributes_over_Symmetric_Difference) (cit. on p. 159).
- [Pro25u] Proof Wiki Contributors. *Intersection Is Associative — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_is\\_Associative](https://proofwiki.org/wiki/Intersection_is_Associative) (cit. on p. 144).
- [Pro25v] Proof Wiki Contributors. *Intersection Is Commutative — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_is\\_Commutative](https://proofwiki.org/wiki/Intersection_is_Commutative) (cit. on p. 144).
- [Pro25w] Proof Wiki Contributors. *Intersection With Empty Set — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_with\\_Empty\\_Set](https://proofwiki.org/wiki/Intersection_with_Empty_Set) (cit. on p. 144).
- [Pro25x] Proof Wiki Contributors. *Intersection With Set Difference Is Set Difference With Intersection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_with\\_Set\\_Difference\\_is\\_Set\\_Difference\\_with\\_Intersection](https://proofwiki.org/wiki/Intersection_with_Set_Difference_is_Set_Difference_with_Intersection) (cit. on p. 149).
- [Pro25y] Proof Wiki Contributors. *Intersection With Subset Is Subset — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_with\\_Subset\\_is\\_Subset](https://proofwiki.org/wiki/Intersection_with_Subset_is_Subset) (cit. on p. 144).
- [Pro25z] Proof Wiki Contributors. *Mapping From Quotient Set When Defined Is Unique — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Mapping\\_from\\_Quotient\\_Set\\_whenDefined\\_is\\_Unique](https://proofwiki.org/wiki/Mapping_from_Quotient_Set_whenDefined_is_Unique) (cit. on p. 724).
- [Pro25aa] Proof Wiki Contributors. *Preimage of Intersection Under Mapping — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Preimage\\_of\\_Intersection\\_under\\_Mapping](https://proofwiki.org/wiki/Preimage_of_Intersection_under_Mapping) (cit. on pp. 145, 227, 228).
- [Pro25ab] Proof Wiki Contributors. *Preimage of Set Difference Under Mapping — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Preimage\\_of\\_Set\\_Difference\\_under\\_Mapping](https://proofwiki.org/wiki/Preimage_of_Set_Difference_under_Mapping) (cit. on pp. 150, 227).

- [Pro25ac] Proof Wiki Contributors. *Preimage of Union Under Mapping — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Preimage\\_of\\_Union\\_under\\_Mapping](https://proofwiki.org/wiki/Preimage_of_Union_under_Mapping) (cit. on pp. 139, 227, 228).
- [Pro25ad] Proof Wiki Contributors. *Quotient Mapping Is Coequalizer — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Quotient\\_Mapping\\_is\\_Coequalizer](https://proofwiki.org/wiki/Quotient_Mapping_is_Coequalizer) (cit. on p. 93).
- [Pro25ae] Proof Wiki Contributors. *Set Difference as Intersection With Complement — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_as\\_Intersection\\_with\\_Complement](https://proofwiki.org/wiki/Set_Difference_as_Intersection_with_Complement) (cit. on p. 149).
- [Pro25af] Proof Wiki Contributors. *Set Difference as Symmetric Difference With Intersection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_as\\_Symmetric\\_Difference\\_with\\_Intersection](https://proofwiki.org/wiki/Set_Difference_as_Symmetric_Difference_with_Intersection) (cit. on p. 149).
- [Pro25ag] Proof Wiki Contributors. *Set Difference Is Right Distributive Over Union — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_is\\_Right\\_Distributive\\_over\\_Union](https://proofwiki.org/wiki/Set_Difference_is_Right_Distributive_over_Union) (cit. on p. 149).
- [Pro25ah] Proof Wiki Contributors. *Set Difference Over Subset — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_over\\_Subset](https://proofwiki.org/wiki/Set_Difference_over_Subset) (cit. on p. 148).
- [Pro25ai] Proof Wiki Contributors. *Set Difference With Empty Set Is Self — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_with\\_Empty\\_Set\\_is\\_Self](https://proofwiki.org/wiki/Set_Difference_with_Empty_Set_is_Self) (cit. on p. 149).
- [Pro25aj] Proof Wiki Contributors. *Set Difference With Self Is Empty Set — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_with\\_Self\\_is\\_Empty\\_Set](https://proofwiki.org/wiki/Set_Difference_with_Self_is_Empty_Set) (cit. on p. 149).
- [Pro25ak] Proof Wiki Contributors. *Set Difference With Set Difference Is Union of Set Difference With Intersection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_with\\_Set\\_Difference\\_is\\_Union\\_of\\_Set\\_Difference\\_with\\_Intersection](https://proofwiki.org/wiki/Set_Difference_with_Set_Difference_is_Union_of_Set_Difference_with_Intersection) (cit. on p. 149).
- [Pro25al] Proof Wiki Contributors. *Set Difference With Subset Is Superset of Set Difference — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_with\\_Subset\\_is\\_Superset\\_of\\_Set\\_Difference](https://proofwiki.org/wiki/Set_Difference_with_Subset_is_Superset_of_Set_Difference) (cit. on p. 148).

- [Pro25am] Proof Wiki Contributors. *Set Difference With Union — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Difference\\_with\\_Union](https://proofwiki.org/wiki/Set_Difference_with_Union) (cit. on p. 149).
- [Pro25an] Proof Wiki Contributors. *Set Intersection Distributes Over Union — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Intersection\\_Distributes\\_over\\_Union](https://proofwiki.org/wiki/Intersection_Distributes_over_Union) (cit. on pp. 138, 144).
- [Pro25ao] Proof Wiki Contributors. *Set Intersection Is Idempotent — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Intersection\\_is\\_Idempotent](https://proofwiki.org/wiki/Set_Intersection_is_Idempotent) (cit. on p. 144).
- [Pro25ap] Proof Wiki Contributors. *Set Intersection Preserves Subsets — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Intersection\\_Preserves\\_Subsets](https://proofwiki.org/wiki/Set_Intersection_Preserves_Subsets) (cit. on p. 144).
- [Pro25aq] Proof Wiki Contributors. *Set Union Is Idempotent — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Union\\_is\\_Idempotent](https://proofwiki.org/wiki/Set_Union_is_Idempotent) (cit. on p. 139).
- [Pro25ar] Proof Wiki Contributors. *Set Union Preserves Subsets — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Set\\_Union\\_Preserves\\_Subsets](https://proofwiki.org/wiki/Set_Union_Preserves_Subsets) (cit. on p. 138).
- [Pro25as] Proof Wiki Contributors. *Symmetric Difference Is Associative — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_is\\_Associative](https://proofwiki.org/wiki/Symmetric_Difference_is_Associative) (cit. on p. 159).
- [Pro25at] Proof Wiki Contributors. *Symmetric Difference Is Commutative — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_is\\_Commutative](https://proofwiki.org/wiki/Symmetric_Difference_is_Commutative) (cit. on p. 159).
- [Pro25au] Proof Wiki Contributors. *Symmetric Difference of Complements — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_of\\_Complements](https://proofwiki.org/wiki/Symmetric_Difference_of_Complements) (cit. on p. 159).
- [Pro25av] Proof Wiki Contributors. *Symmetric Difference on Power Set Forms Abelian Group — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_on\\_Power\\_Set\\_forms\\_Abelian\\_Group](https://proofwiki.org/wiki/Symmetric_Difference_on_Power_Set_forms_Abelian_Group) (cit. on p. 160).
- [Pro25aw] Proof Wiki Contributors. *Symmetric Difference With Complement — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_with\\_Complement](https://proofwiki.org/wiki/Symmetric_Difference_with_Complement) (cit. on p. 159).
- [Pro25ax] Proof Wiki Contributors. *Symmetric Difference With Empty Set — Proof Wiki*. 2025. URL: <https://proofwiki.org/>

- wiki/Symmetric\_Difference\_with\_Empty\_Set (cit. on p. 159).
- [Pro25ay] Proof Wiki Contributors. *Symmetric Difference With Intersection Forms Ring — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_with\\_Intersection\\_forms\\_Ring](https://proofwiki.org/wiki/Symmetric_Difference_with_Intersection_forms_Ring) (cit. on p. 160).
- [Pro25az] Proof Wiki Contributors. *Symmetric Difference With Self Is Empty Set — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_with\\_Self\\_is\\_Empty\\_Set](https://proofwiki.org/wiki/Symmetric_Difference_with_Self_is_Empty_Set) (cit. on p. 159).
- [Pro25ba] Proof Wiki Contributors. *Symmetric Difference With Union Does Not Form Ring — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_with\\_Union\\_does\\_not\\_form\\_Ring](https://proofwiki.org/wiki/Symmetric_Difference_with_Union_does_not_form_Ring) (cit. on p. 158).
- [Pro25bb] Proof Wiki Contributors. *Symmetric Difference With Universe — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Symmetric\\_Difference\\_with\\_Universe](https://proofwiki.org/wiki/Symmetric_Difference_with_Universe) (cit. on p. 159).
- [Pro25bc] Proof Wiki Contributors. *Union as Symmetric Difference With Intersection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Union\\_as\\_Symmetric\\_Difference\\_with\\_Intersection](https://proofwiki.org/wiki/Union_as_Symmetric_Difference_with_Intersection) (cit. on p. 139).
- [Pro25bd] Proof Wiki Contributors. *Union Distributes Over Intersection — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Union\\_Distributes\\_over\\_Intersection](https://proofwiki.org/wiki/Union_Distributes_over_Intersection) (cit. on pp. 138, 144).
- [Pro25be] Proof Wiki Contributors. *Union Is Associative — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Union\\_is\\_Associative](https://proofwiki.org/wiki/Union_is_Associative) (cit. on p. 138).
- [Pro25bf] Proof Wiki Contributors. *Union Is Commutative — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Union\\_is\\_Commutative](https://proofwiki.org/wiki/Union_is_Commutative) (cit. on p. 138).
- [Pro25bg] Proof Wiki Contributors. *Union of Symmetric Differences — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Union\\_of\\_Symmetric\\_Differences](https://proofwiki.org/wiki/Union_of_Symmetric_Differences) (cit. on p. 159).
- [Pro25bh] Proof Wiki Contributors. *Union With Empty Set — Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/Union\\_with\\_Empty\\_Set](https://proofwiki.org/wiki/Union_with_Empty_Set) (cit. on p. 138).
- [PS19] Maximilien Péroux and Brooke Shipley. “Coalgebras in Symmetric Monoidal Categories of Spectra”. In: *Homology Homotopy Appl.* 21.1 (2019), pp. 1–18. ISSN: 1532-0073. doi:

[10.4310/HHA.2019.v21.n1.a1](https://doi.org/10.4310/HHA.2019.v21.n1.a1). URL: <https://doi.org/10.4310/HHA.2019.v21.n1.a1> (cit. on p. 517).

- [Rie16] Emily Riehl. *Category Theory in Context*. Aurora Dover Modern Math Originals. Dover Publications, Inc., Mineola, NY, 2016, pp. xvii+240. ISBN: 978-0-486-80903-8; 0-486-80903-X (cit. on pp. 5, 9, 791).
- [Wik25] Wikipedia Contributors. *Multivalued Function — Wikipedia, The Free Encyclopedia*. 2025. URL: [https://en.wikipedia.org/wiki/Multivalued\\_function](https://en.wikipedia.org/wiki/Multivalued_function) (cit. on p. 554).

# Index of Notation

## A

- $[a]$ , 707
- $(A, B)$ , 94
- $A \coprod B$ , 72
- $A \coprod_C B$ , 75
- $A \coprod_{f,c,g} B$ , 79
- $A \coprod_C^{f,g} B$ , 79
- $\alpha F$ , 803
- $\alpha \star F$ , 803
- $A \odot X$ , 382
- $a \odot x$ , 384
- $A \odot (X, x_0)$ , 382
- $A \pitchfork X$ , 390
- $A \pitchfork (X, x_0)$ , 390
- $A \times B$ , 40
- $A \times_C B$ , 50
- $A \times_{f,C,g} B$ , 52
- $A \times_C^{f,g} B$ , 52

## B

- $\beta \circ \alpha$ , 797
- $\beta \star \alpha$ , 801

## C

- $C(A, B)$ , 717
- $\text{Cats}$ , 814
- $\text{Cats}_2$ , 815
- $[C, \mathcal{D}]$ , 811
- $\dot{\mathcal{Q}}$ , 819
- $\chi_{(-)}$ , 201
- $\chi_{-2}^{-1}$ , 199
- $\chi_U$ , 193

$\chi^x$ , 198

$\chi_x$ , 198

$\chi_X(-_1, -_2)$ , 198

$\chi_X(-, U)$ , 193

$\chi_X(-, x)$ , 198

$\chi_X(U, -)$ , 193

$\chi_X(x, -)$ , 198

$\text{CoEq}(f, g)$ , 84

$\text{CoIm}(f)$ , 711

$\operatorname{colim}_{\substack{\longrightarrow \\ \alpha \in I}} (X_\alpha)$ , 89

$\text{Coll}(R)$ , 686

$\text{Coll}(R)$ , 686

$\coprod_{i \in I} A_i$ , 70

$\text{CoPSh}(C)$ , 831

$\text{Core}(C)$ , 743

$C^\simeq$ , 744

## D

- $\mathcal{D}^C$ , 811
- $\text{dom}(R)$ , 679

## E

- $\emptyset_{\text{cat}}$ , 725
- $\emptyset$ , 69, 94
- $\text{Eq}(f, g)$ , 61
- $\text{eq}(f, g)$ , 61

## F

- $F\alpha$ , 803
- $\mathcal{F} \boxtimes_C G$ , 841
- $\mathcal{F} \boxtimes G$ , 841
- $f^{-1}$ , 216

$f^{-1}(V)$ , 216  
 $F \odot_C G$ , 845  
 $\mathcal{F} \odot_C G$ , 844  
 $F \odot G$ , 845  
 $\mathcal{F} \odot G$ , 844  
 $\mathbb{F}_1$ , 324  
 $f^!$ , 203  
 $f_!$ , 225  
 $f_!(U)$ , 225  
 $f^*$ , 720  
 $F \star \alpha$ , 803  
 $\alpha \star F$ , 803  
 $f_{*,\text{cp}}(U)$ , 227  
 $f_{*,\text{im}}(U)$ , 227  
 $f(U)$ , 203  
 $\text{Fun}(C, \mathcal{D})$ , 811

## G

$\text{Gr}(A)$ , 542  
 $\text{Grpd}$ , 816  
 $\text{Grpd}_2$ , 816

## H

$h^A$ , 832  
 $h_A$ , 820  
 $\text{Hom}_{\text{Sets}_*}^{\otimes, \text{L}}(X \times Y, Z)$ , 379  
 $\text{Hom}_{\text{Sets}_*}^{\otimes, \text{R}}(X \times Y, Z)$ , 380  
 $\text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z)$ , 381  
 $\text{Hom}_{\mathcal{P}(X)}(U, V)$ , 173  
 $\text{Hom}_{\text{Sets}}(A, B)$ , 95  
 $\text{Hom}_{\text{Sets}_*}(X, Y)$ , 462

## I

$\mathsf{I}$ , 190  
 $\text{id}_C$ , 749  
 $\text{id}_F$ , 795  
 $\text{id}_F \star \alpha$ , 803  
 $\bigcup_{i \in I} R_i$ , 683  
 $\bigcap_{U \in \mathcal{U}} U$ , 112  
 $\text{Iso}_C(A, B)$ , 739

## K

$\text{K}_0(C)$ , 740

## L

$\lhd$ , 397  
 $[-, -]_{\text{Sets}_*}^\lhd$ , 403  
 ${}^\lhd \text{Sets}_*$ , 398  
 $\lim_{\leftarrow \alpha \in I} (X_\alpha)$ , 66

## M

$\mathcal{M}_{\mathbb{E}_1}(C)$ , 849  
 $\text{Mor}(C)$ , 717

## N

$\mathbb{N}$ , 726  
 $\langle n \rangle$ , 324  
 $\nabla_X$ , 540  
 $\text{Nat}(F, G)$ , 795

## P

$\mathcal{P}_0(X)$ , 154  
 $\mathcal{P}_{\text{fin}}(X)$ , 154  
 $\pi_0(C)$ , 733  
 $\mathcal{P}^{-1}$ , 157, 665  
 $\star_X$ , 361  
 $\text{pr}_1^{A \times_C B}$ , 51  
 $\text{pr}_2^{A \times_C B}$ , 51  
 $\prod_{i \in I} A_i$ , 37, 333  
 $\text{PSh}(C)$ , 819  
 $\mathcal{P}_!$ , 158, 665  
 $\mathcal{P}_!$ , 157  
 $\mathcal{P}_*$ , 665  
 $\text{pt}$ , 36, 324, 332, 346  
 $\text{pt}$ , 725  
 $\mathcal{P}(X)$ , 154

## R

$\text{range}(R)$ , 679  
 $\text{Ran}_R(F)$ , 623  
 $R^\dagger$ , 534  
 $\text{Rel}$ , 549  
 $\text{Rel}$ , 556, 565

$\text{Rel}(A, B)$ , 517	$\sim_R^{\text{eq}}$ , 706
$\text{Rel}(A, B)$ , 517, 548	$\sim_R^{\text{trans}}$ , 701
$\text{Rel}^\square$ , 564	$\sim_{\text{triv}}$ , 538
$\text{Rel}^{\text{dbl}}$ , 557	$S^0$ , 324
$\text{Rel}^{\text{eq}}(A, B)$ , 704	
$\text{Rel}^{\text{eq}}(A, B)$ , 704	
$\text{Rel}^{\text{refl}}(A, A)$ , 694	
$\text{Rel}^{\text{refl}}(A, A)$ , 694	
$\text{Rel}^{\text{symm}}(A, A)$ , 697	
$\text{Rel}^{\text{symm}}(A, A)$ , 697	
$\text{Rel}^{\text{trans}}(A)$ , 701	
$\text{Rel}^{\text{trans}}(A)$ , 701	
$R^{\text{eq}}$ , 706	
$\triangleright$ , 425	
$[-, -]_{\text{Sets}_*}^\triangleright$ , 430	
$\triangleright_{\text{Sets}_*}$ , 425	
$\text{Rift}_R(F)$ , 628	
$R \cap S$ , 681	
$R^{-1}$ , 653	
$R_{-1}$ , 647	
$R^{-1}(V)$ , 654	
$R_{-1}(V)$ , 647	
$R^{\text{refl}}$ , 694	
$R^!$ , 641	
$R_*$ , 659	
$R_*(U)$ , 659	
$R^{\text{symm}}$ , 697	
$R \times S$ , 684	
$R^{\text{trans}}$ , 702	
$R(U)$ , 641	
$R \cup S$ , 679	
<b>S</b>	
$S \square R$ , 528	
$S \diamond R$ , 520	
$\text{Sets}(A, B)$ , 95	
$\text{Sets}_*^{\text{actv}}$ , 329	
$\text{Sets}_*$ , 325	
$\text{Sets}_*((X, x_0), (Y, y_0))$ , 462	
$\sim_{\text{cotriv}}$ , 539	
$\sim_{\text{id}}$ , 199	
$\sim_R^{\text{refl}}$ , 694	
$\sim_R^{\text{symm}}$ , 697	
<b>T</b>	
$\{\text{t}, \text{f}\}$ , 28	
$\prod_{i \in I} R_i$ , 686	
$\text{Trans}(F, G)$ , 794	
$\{\text{true}, \text{false}\}$ , 27	
<b>U</b>	
$U \boxtimes_{X \times Y} V$ , 155	
$U^c$ , 143	
$U \cap V$ , 132	
$\bigcup_{i \in I} R_i$ , 681	
$\bigcup_{U \in \mathcal{U}} U$ , 98	
$U \Delta V$ , 146	
$U \cup V$ , 126	
$[U, V]_X$ , 173	
<b>V</b>	
$\bigvee_{i \in I} X_i$ , 347	
<b>W</b>	
$\overline{\exists}$ , 753	
$\bigwedge_{i \in I} X_i$ , 510	
<b>X</b>	
$\{X\}$ , 94	
$X_{\text{disc}}$ , 735	
$X_{\text{indisc}}$ , 737	
$x \triangleleft_{\text{Sets}_*} y$ , 400	
$x \triangleleft y$ , 400	
$X^-$ , 369	
$X \setminus Y$ , 138	
$X / \sim_R$ , 708	
$X \otimes_{\mathbb{F}_1} Y$ , 451	
$X^+$ , 361	
$X_{\text{pos}}$ , 728	
$x \triangleright y$ , 427	

$x \triangleright_{\text{Sets}_*} y$ , 427

$x \wedge y$ , 454

$X \wedge Y$ , 451

$\{X, Y\}$ , 94

# Y

$\mathfrak{P}$ , 834

$\mathfrak{P}$ , 822

$\mathfrak{P}_F$ , 836

# Index of Set Theory

## B

bilinear morphism  
    of pointed sets, 380  
bilinear morphism of pointed sets  
    left, 378  
    right, 379  
binary intersection, 132

## C

Cartesian product, 37, 39  
category of relations, 549  
    associator of, 550  
    internal Hom of, 553  
    left unit of, 551  
    monoidal product of, 549  
    monoidal unit of, 550  
    right unit of, 552  
    symmetry of, 552  
    with apartness composition,  
        564  
characteristic embedding, 201  
characteristic function  
    of a subset, 193  
    of an element, 198  
characteristic relation, 198  
coequaliser of sets, 84  
coimage, 711  
complement of a set, 143  
copresheaf  
    category of, 831  
coproduct, 72  
    of a family of pointed sets,  
        347

    of a family of sets, 69  
coproduct of sets  
    associator, 293  
    left unit, 297  
    monoidal unit of, 293  
    right unit, 299  
    symmetry, 301

## D

difference of sets, 138  
direct colimit of sets, 88  
disjoint union, 72  
    of a family of sets, 69

## E

empty set, 94  
equaliser of sets, 61  
equivalence class, 707  
equivalence relation  
    kernel, 704

## F

fibre coproduct of sets, 75  
fibre product of sets, 50  
field with one element  
    module over, 323  
    module over, morphism of,  
        324  
function, 25  
    associated codirect image  
        function, 225

associated codirect image function, complement part, 227  
associated codirect image function, image part, 227  
associated direct image function, 203  
associated inverse image function, 216  
coimage of, 711  
graph of, 542  
inverse of, 545  
kernel of, 704

## I

indicator function, *see* characteristic function  
initial pointed set, 346  
initial set, 69  
injective, 246  
intersection of a family of sets, 112  
inverse limit of sets, 66  
Isbell function, 190

## L

left tensor product of pointed sets diagonal, 415  
left skew unit of, 406  
skew associator, 406  
skew left unit, 409  
skew right unit, 413

## M

( $-1$ )-category, 27  
( $-2$ )-category, 27  
module  
underlying pointed set of, 908  
morphism of pointed sets, 324  
active, 329  
inert, 329  
multivalued function

see relation, 515

## O

ordered pairing, 94

## P

pairing of two sets, 94  
pointed function, *see* pointed set, morphism of pointed set, 323  
category of, 325  
coequaliser of, 358  
copower by a set, *see* pointed set, tensor by a set  
coproduct of, 349  
cotensor by a set, 390  
deleting the basepoint, 369  
equaliser of, 343  
 $\mathbb{F}_1$ , 324  
free, 361  
internal Hom of, 462  
left internal Hom of, 403  
left tensor product, 397  
morphism of, 324  
of morphisms of pointed sets, 462  
power by a set, *see* pointed set, cotensor by a set  
product of, 335  
pullback of, 338  
pushout of, 352  
right internal Hom of, 430  
right tensor product, 425  
standard pointed set with  $n + 1$  elements, 324  
tensor by a set, 382  
trivial, 324  
underlying a module, 908  
underlying a semimodule, 908  
wedge sum, 349  
wedge sum of a family of pointed sets, 347

- 0-sphere, 324  
 poset  
   of  $(-1)$ -categories, 28  
   of truth values, 27  
 powerset, 154  
   internal Hom of, 173  
 presheaf  
   category of, 819  
 product of a family of pointed sets, 333  
 product of a family of sets, 37  
 product of sets, 39  
   associator, 260  
   diagonal, 268  
   left annihilator, 316  
   left distributor, 311  
   left unitor, 262  
   monoidal unit of, 259  
   right annihilator, 317  
   right distributor, 313  
   right unitor, 264  
   symmetry, 266  
 pullback of sets, 50  
 pushout of sets, 75
- Q**
- quotient  
   by an equivalence relation, 708
- R**
- relation, 515  
   antidiagonal, 540  
   apartness composition of, 528  
   associated codirect image  
     function, 659  
   associated coinverse image  
     function, 647  
   associated direct image  
     function, 641  
   associated inverse image  
     function, 653  
   category of, 548
- category of, 549  
 closed symmetric monoidal  
   category of, 555  
 collage of, 686  
 composition of, 520  
 converse of, 534  
 corepresentable, 547  
 cotrivial, 539  
 domain of, 679  
 double category of, 557  
 equivalence closure of, 706  
 equivalence relation, 704  
 equivalence, set of, 704  
 functional, 691  
 intersection of, 681  
 intersection of a family of, 683  
 linear bicategory of, 567  
 on powersets associated to a relation, 666  
 partial equivalence relation, 704  
 poset of, 697, 701, 704  
 product of, 684  
 product of a family of, 686  
 range of, 679  
 reflexive, 693  
 reflexive closure of, 694  
 reflexive, poset of, 694  
 reflexive, set of, 694  
 representable, 547  
 symmetric, 696  
 symmetric closure of, 697  
 symmetric, set of, 697  
 total, 692  
 transitive, 700  
 transitive closure of, 701  
 transitive, set of, 701  
 trivial, 538  
 2-category of, 556  
   with apartness composition, 565  
 union of, 679  
 union of a family of, 681

right tensor product of pointed sets  
diagonal, 441  
right skew unit of, 432  
skew associator, 433  
skew left unit, 436  
skew right unit, 438

## S

semimodule  
underlying pointed set of, 908  
set  
Hom, 95  
of maps, 95  
set of bilinear morphisms of pointed sets, 381  
left, 379  
right, 380  
singleton set, 94  
smash product  
of a family of pointed sets, 510  
of pointed sets, 451

smash product of pointed sets  
associator, 464  
diagonal, 474  
left unit, 466  
monoidal unit of, 464  
right unit, 469  
symmetry, 472  
surjective, 252  
symmetric difference of sets, 146

## T

terminal pointed set, 332  
terminal set, 36

## U

union, 126  
of a family of sets, 98

## Z

0-category, 31  
0-groupoid, 31

# Index of Category Theory

## A

adjoint equivalence of categories,  
774

corepresenting object of, 833  
morphism of, 831  
corepresentable natural  
transformation, 834

## C

category, 716  
adjoint equivalence of, 774  
connected, 735  
connected component, 733  
connected component, set of,  
733  
core of, 743  
disconnected, 735  
discrete, 736  
empty, 725  
equivalence of, 773  
groupoid completion of, 740  
indiscrete, 738  
isomorphism of, 776  
 $\kappa$ -small, 718  
locally essentially small, 718  
locally small, 717  
of small groupoids, 816  
skeletal, 719  
skeleton of, 719  
small, 718  
thin, 729  
category of categories, 814  
contravariant functor, 751  
copresheaf, 831  
corepresentable, 833  
corepresentable associated to  
an object, 832

## D

discrete category  
on a set, 735

## E

equivalence of categories, 773

## F

forgetful functor, 753  
functor, 747  
bijective on objects, 787  
bo, 787  
composition of, 750  
conservative, 770  
contravariant, 751  
corepresentably faithful on  
cores, 791  
corepresentably full on cores,  
791  
corepresentably fully faithful  
on cores, 792  
dominant, 777  
epimorphism, 780  
eso, 773  
essentially injective, 772  
essentially surjective, 773  
faithful, 757

forgetful, 753  
full, 760  
fully faithful, 766  
identity, 749  
injective on objects, 786  
monomorphism, 779  
pseudoepic, 784  
pseudomonic, 781  
representably faithful on cores, 788  
representably full on cores, 788  
representably fully faithful on cores, 789  
surjective on objects, 787  
transformation of, 794  
functor category, 811  
functor tensor product, 841

## G

Godement product, *see* natural transformation,  
horizontal composition  
Grothendieck groupoid  
of a category, 740  
groupoid, 740  
groupoid completion, 740

## I

indiscrete category  
on a set, 737  
Isbell's zigzag theorem, 781  
isomorphism, 739  
isomorphism of categories, 776

## M

middle four exchange  
in Cats, 799, 804  
moduli category  
of monoidal structures, 849

## N

natural isomorphism, 808  
natural transformation, 794  
associated to a functor, 755  
component of, 795  
equality of, 795  
horizontal composition, 801  
identity natural  
transformation, 795  
vertical composition, 797

## O

ordinal category, 726

## P

posetal category, 729  
associated to a poset, 728  
postcomposition, 720  
precomposition, 720  
presheaf, 818  
morphism of, 819  
representable, 820  
representable associated to an object, 820  
representing object of, 820  
punctual category, 725

## R

representable natural  
transformation, 822  
restricted Yoneda embedding, 836

## S

singleton category, *see* punctual category  
subcategory, 718  
full, 718  
lluf, 719  
strictly full, 719  
wide, 719

## T

tensor of a copresheaf with a functor, 845  
tensor of a presheaf with a functor, 844  
tensor product of functors, 841  
transformation between functors, 794  
2-category  
    of small categories, 815  
    of small groupoids, 816

## W

walking arrow, 727  
whiskering

left, 803  
right, 803

## Y

Yoneda embedding, 822  
contravariant, 834  
restricted, 836  
Yoneda extension, 838  
Yoneda lemma, 825  
contravariant, 835

## Z

(0, 1)-category, 729

# Index of Higher Category Theory

## D

double category  
of relations, 557

## E

epimorphism  
strict, 883

## M

monomorphism  
strict, 872

## O

1-morphism  
corepresentably conservative,  
882  
corepresentably essentially  
injective, 882  
corepresentably faithful, 875  
corepresentably faithful on  
cores, 879

corepresentably full, 875  
corepresentably full on cores,  
880  
corepresentably fully faithful,  
876  
corepresentably fully faithful  
on cores, 880  
pseudoepic, 884  
pseudomonic, 872  
representably conservative,  
871  
representably essentially  
injective, 871  
representably faithful, 864  
representably faithful on  
cores, 868  
representably full, 865  
representably full on cores,  
868  
representably fully faithful,  
866  
representably fully faithful on  
cores, 869