

Constructions With Sets

The Clowder Project Authors

July 22, 2025

- 000J This chapter develops some material relating to constructions with sets with an eye towards its categorical and higher-categorical counterparts to be introduced later in this work. Of particular interest are perhaps the following:
- 01YT 1. Explicit descriptions of the major types of co/limits in Sets, including in particular explicit descriptions of pushouts and coequalisers (see [Definitions 4.2.4.1.1](#), [4.2.4.1.3](#), [4.2.5.1.1](#) and [4.2.5.1.3](#)).
- 01YU 2. A discussion of powersets as decategorifications of categories of presheaves, including in particular results such as:
- 01YV (a) A discussion of the internal Hom of a powerset ([Section 4.4.7](#)).
- 01YW (b) A o-categorical version of the Yoneda lemma ([Presheaves and the Yoneda Lemma](#), [Definition 12.1.5.1.1](#)), which we term the *Yoneda lemma for sets* ([Definition 4.5.5.1.1](#)).
- 01YX (c) A characterisation of powersets as free cocompletions ([Section 4.4.5](#)), mimicking the corresponding statement for categories of presheaves (??).
- 01YY (d) A characterisation of powersets as free completions ([Section 4.4.6](#)), mimicking the corresponding statement for categories of copresheaves (??).
- 01YZ (e) A (-1) -categorical version of un/straightening ([Item 2 of Definition 4.5.1.1.4](#) and [Definition 4.5.1.1.5](#)).
- 01Z0 (f) A o-categorical form of Isbell duality internal to powersets ([Section 4.4.8](#)).

01Z1 3. A lengthy discussion of the adjoint triple

$$f_! \dashv f^{-1} \dashv f_*: \mathcal{P}(A) \xrightarrow{\cong} \mathcal{P}(B)$$

of functors (i.e. morphisms of posets) between $\mathcal{P}(A)$ and $\mathcal{P}(B)$ induced by a map of sets $f: A \rightarrow B$, including in particular:

01Z2 (a) How f^{-1} can be described as a precomposition while $f_!$ and f_* can be described as Kan extensions ([Definitions 4.6.1.1.4](#), [4.6.2.1.2](#) and [4.6.3.1.4](#)).

01Z3 (b) An extensive list of the properties of $f_!$, f^{-1} , and f_* ([Definitions 4.6.1.1.5](#), [4.6.1.1.6](#), [4.6.2.1.3](#), [4.6.2.1.4](#), [4.6.3.1.7](#) and [4.6.3.1.8](#)).

01Z4 (c) How the functors $f_!$, f^{-1} , f_* , along with the functors

$$\begin{aligned} -_1 \cap -_2: \mathcal{P}(X) \times \mathcal{P}(X) &\rightarrow \mathcal{P}(X), \\ [-_1, -_2]_X: \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) &\rightarrow \mathcal{P}(X) \end{aligned}$$

may be viewed as a six-functor formalism with the empty set \emptyset as the dualising object ([Section 4.6.4](#)).

Contents

4.1 Limits of Sets	4
4.1.1 The Terminal Set	4
4.1.2 Products of Families of Sets	4
4.1.3 Binary Products of Sets	7
4.1.4 Pullbacks	18
4.1.5 Equalisers	30
4.1.6 Inverse Limits	35
4.2 Colimits of Sets	38
4.2.1 The Initial Set	38
4.2.2 Coproducts of Families of Sets	39
4.2.3 Binary Coproducts	41
4.2.4 Pushouts	44
4.2.5 Coequalisers	53
4.2.6 Direct Colimits	58

4.3 Operations With Sets	63
4.3.1 The Empty Set	63
4.3.2 Singleton Sets	63
4.3.3 Pairings of Sets	63
4.3.4 Ordered Pairs	64
4.3.5 Sets of Maps	64
4.3.6 Unions of Families of Subsets	67
4.3.7 Intersections of Families of Subsets	81
4.3.8 Binary Unions	95
4.3.9 Binary Intersections	101
4.3.10 Differences	107
4.3.11 Complements	111
4.3.12 Symmetric Differences	114
4.4 Powersets	122
4.4.1 Foundations	122
4.4.2 Functoriality of Powersets	125
4.4.3 Adjointness of Powersets I	127
4.4.4 Adjointness of Powersets II	127
4.4.5 Powersets as Free Cocompletions	128
4.4.6 Powersets as Free Completions	135
4.4.7 The Internal Hom of a Powerset	141
4.4.8 Isbell Duality for Sets	159
4.5 Characteristic Functions	161
4.5.1 The Characteristic Function of a Subset	161
4.5.2 The Characteristic Function of a Point	166
4.5.3 The Characteristic Relation of a Set	167
4.5.4 The Characteristic Embedding of a Set	168
4.5.5 The Yoneda Lemma for Sets	169
4.6 The Adjoint Triple $f_! \dashv f^{-1} \dashv f_*$	170
4.6.1 Direct Images	170
4.6.2 Inverse Images	182
4.6.3 Codirect Images	191
4.6.4 A Six-Functor Formalism for Sets	204
A Other Chapters	213

000K 4.1 Limits of Sets

000L 4.1.1 The Terminal Set

000M **Definition 4.1.1.1.** The **terminal set** is the terminal object of **Sets** as in Limits and Colimits, ??.

01DB **Construction 4.1.1.2.** Concretely, the terminal set is the pair $(\text{pt}, \{!_A\}_{A \in \text{Obj}(\text{Sets})})$ consisting of:

01DC 1. *The Limit.* The punctual set $\text{pt} \stackrel{\text{def}}{=} \{\star\}$.

01DD 2. *The Cone.* The collection of maps

$$\{!_A : A \rightarrow \text{pt}\}_{A \in \text{Obj}(\text{Sets})}$$

defined by

$$!_A(a) \stackrel{\text{def}}{=} \star$$

for each $a \in A$ and each $A \in \text{Obj}(\text{Sets})$.

Proof. We claim that pt is the terminal object of **Sets**. Indeed, suppose we have a diagram of the form

$$A \quad \text{pt}$$

in **Sets**. Then there exists a unique map $\phi : A \rightarrow \text{pt}$ making the diagram

$$A \xrightarrow[\exists!]{\phi} \text{pt}$$

commute, namely $!_A$. □

000N 4.1.2 Products of Families of Sets

Let $\{A_i\}_{i \in I}$ be a family of sets.

000P **Definition 4.1.2.1.** The **product**¹ of $\{A_i\}_{i \in I}$ is the product of $\{A_i\}_{i \in I}$ in **Sets** as in Limits and Colimits, ??.

¹*Further Terminology:* Also called the **Cartesian product** of $\{A_i\}_{i \in I}$.

01DE Construction 4.1.2.1.2. Concretely, the product of $\{A_i\}_{i \in I}$ is the pair $\left(\prod_{i \in I} A_i, \{\text{pr}_i\}_{i \in I}\right)$ consisting of:

01DF 1. *The Limit.* The set $\prod_{i \in I} A_i$ defined by

$$\prod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ f \in \text{Sets} \left(I, \bigcup_{i \in I} A_i \right) \left| \begin{array}{l} \text{for each } i \in I, \text{ we} \\ \text{have } f(i) \in A_i \end{array} \right. \right\}.$$

01DG 2. *The Cone.* The collection

$$\left\{ \text{pr}_i : \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

of maps given by

$$\text{pr}_i(f) \stackrel{\text{def}}{=} f(i)$$

for each $f \in \prod_{i \in I} A_i$ and each $i \in I$.

Proof. We claim that $\prod_{i \in I} A_i$ is the categorical product of $\{A_i\}_{i \in I}$ in **Sets**. Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} P & & \\ & \searrow p_i & \\ \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} & A_i \end{array}$$

in **Sets**. Then there exists a unique map $\phi: P \rightarrow \prod_{i \in I} A_i$ making the diagram

$$\begin{array}{ccc} P & & \\ \downarrow \phi \quad \exists! & \searrow p_i & \\ \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} & A_i \end{array}$$

commute, being uniquely determined by the condition $\text{pr}_i \circ \phi = p_i$ for each $i \in I$ via

$$\phi(x) = (p_i(x))_{i \in I}$$

for each $x \in P$. □

01DH Remark 4.1.2.1.3. Less formally, we may think of Cartesian products and projection maps as follows:

01DJ 1. We think of $\prod_{i \in I} A_i$ as the set whose elements are I -indexed collections $(a_i)_{i \in I}$ with $a_i \in A_i$ for each $i \in I$.

01DK 2. We view the projection maps

$$\left\{ \text{pr}_i : \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

as being given by

$$\text{pr}_i \left((a_j)_{j \in I} \right) \stackrel{\text{def}}{=} a_i$$

for each $(a_j)_{j \in I} \in \prod_{i \in I} A_i$ and each $i \in I$.

000Q Proposition 4.1.2.1.4. Let $\{A_i\}_{i \in I}$ be a family of sets.

000R 1. *Functoriality.* The assignment $\{A_i\}_{i \in I} \mapsto \prod_{i \in I} A_i$ defines a functor

$$\prod_{i \in I} : \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, we have

$$\left[\prod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} A_i$$

- *Action on Morphisms.* For each $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, the action on Hom-sets

$$\left(\prod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left(\prod_{i \in I} A_i, \prod_{i \in I} B_i \right)$$

of $\prod_{i \in I}$ at $((A_i)_{i \in I}, (B_i)_{i \in I})$ is defined by sending a map

$$\{f_i : A_i \rightarrow B_i\}_{i \in I}$$

in $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$ to the map of sets

$$\prod_{i \in I} f_i : \prod_{i \in I} A_i \rightarrow \prod_{i \in I} B_i$$

defined by

$$\left[\prod_{i \in I} f_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} (f_i(a_i))_{i \in I}$$

for each $(a_i)_{i \in I} \in \prod_{i \in I} A_i$.

Proof. **Item 1, Functoriality:** This follows from Limits and Colimits, ?? of ??. \square

000S 4.1.3 Binary Products of Sets

Let A and B be sets.

000T Definition 4.1.3.1.1. The **product of A and B** ² is the product of A and B in Sets as in Limits and Colimits, ??.

01DL Construction 4.1.3.1.2. Concretely, the product of A and B is the pair $(A \times B, \{\text{pr}_1, \text{pr}_2\})$ consisting of:

01DM 1. *The Limit.* The set $A \times B$ defined by

$$\begin{aligned} A \times B &\stackrel{\text{def}}{=} \prod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{f \in \text{Sets}(\{0, 1\}, A \cup B) \mid \text{we have } f(0) \in A \text{ and } f(1) \in B\} \\ &\cong \{\{\{a\}, \{a, b\}\} \in \mathcal{P}(\mathcal{P}(A \cup B)) \mid \text{we have } a \in A \text{ and } b \in B\} \\ &\cong \left\{ \begin{array}{l} \text{ordered pairs } (a, b) \text{ with} \\ a \in A \text{ and } b \in B \end{array} \right\}. \end{aligned}$$

01DN 2. *The Cone.* The maps

$$\begin{aligned} \text{pr}_1 : A \times B &\rightarrow A, \\ \text{pr}_2 : A \times B &\rightarrow B \end{aligned}$$

²*Further Terminology:* Also called the **Cartesian product of A and B** .

defined by

$$\begin{aligned} \text{pr}_1(a, b) &\stackrel{\text{def}}{=} a, \\ \text{pr}_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each $(a, b) \in A \times B$.

Proof. We claim that $A \times B$ is the categorical product of A and B in the category of sets. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & P & & \\ & \swarrow p_1 & & \searrow p_2 & \\ A & \xleftarrow{\text{pr}_1} & A \times B & \xrightarrow{\text{pr}_2} & B \end{array}$$

in Sets. Then there exists a unique map $\phi: P \rightarrow A \times B$ making the diagram

$$\begin{array}{ccccc} & & P & & \\ & \swarrow p_1 & \downarrow \phi \mid \exists! & \searrow p_2 & \\ A & \xleftarrow{\text{pr}_1} & A \times B & \xrightarrow{\text{pr}_2} & B \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned} \text{pr}_1 \circ \phi &= p_1, \\ \text{pr}_2 \circ \phi &= p_2 \end{aligned}$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$. □

000U Proposition 4.1.3.1.3. Let A, B, C , and X be sets.

000V 1. *Functoriality.* The assignments $A, B, (A, B) \mapsto A \times B$ define functors

$$\begin{aligned} A \times -: \quad \text{Sets} &\rightarrow \text{Sets}, \\ - \times B: \quad \text{Sets} &\rightarrow \text{Sets}, \\ -_1 \times -_2: \text{Sets} \times \text{Sets} &\rightarrow \text{Sets}, \end{aligned}$$

where $-_1 \times -_2$ is the functor where

- *Action on Objects.* For each $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$, we have

$$[-_1 \times -_2](A, B) \stackrel{\text{def}}{=} A \times B.$$

- *Action on Morphisms.* For each $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\times_{(A,B),(X,Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \times B, X \times Y)$$

of \times at $((A, B), (X, Y))$ is defined by sending (f, g) to the function

$$f \times g : A \times B \rightarrow X \times Y$$

defined by

$$[f \times g](a, b) \stackrel{\text{def}}{=} (f(a), g(b))$$

for each $(a, b) \in A \times B$.

and where $A \times -$ and $- \times B$ are the partial functors of $-_1 \times -_2$ at $A, B \in \text{Obj}(\text{Sets})$.

000W

2. *Adjointness I.* We have adjunctions

$$\begin{aligned} (A \times - \dashv \text{Sets}(A, -)) : \quad & \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Sets}(A, -)} \end{array} \text{Sets}, \\ (- \times B \dashv \text{Sets}(B, -)) : \quad & \text{Sets} \begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Sets}(B, -)} \end{array} \text{Sets}, \end{aligned}$$

witnessed by bijections

$$\text{Sets}(A \times B, C) \cong \text{Sets}(A, \text{Sets}(B, C)),$$

$$\text{Sets}(A \times B, C) \cong \text{Sets}(B, \text{Sets}(A, C)),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

01Z5 3. *Adjointness II.* We have an adjunction

$$(\Delta_{\text{Sets}} \dashv -_1 \times -_2): \text{Sets} \begin{array}{c} \xrightarrow{\Delta_{\text{Sets}}} \\ \perp \\ \xleftarrow{-_1 \times -_2} \end{array} \text{Sets} \times \text{Sets},$$

witnessed by a bijection

$$\text{Hom}_{\text{Sets} \times \text{Sets}}((A, A), (B, C)) \cong \text{Sets}(A, B \times C),$$

natural in $A \in \text{Obj}(\text{Sets})$ and in $(B, C) \in \text{Obj}(\text{Sets} \times \text{Sets})$.

000X 4. *Associativity.* We have an isomorphism of sets

$$\alpha_{A,B,C}^{\text{Sets}}: (A \times B) \times C \xrightarrow{\sim} A \times (B \times C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

000Y 5. *Unitality.* We have isomorphisms of sets

$$\begin{aligned} \lambda_A^{\text{Sets}}: \text{pt} \times A &\xrightarrow{\sim} A, \\ \rho_A^{\text{Sets}}: A \times \text{pt} &\xrightarrow{\sim} A, \end{aligned}$$

natural in $A \in \text{Obj}(\text{Sets})$.

000Z 6. *Commutativity.* We have an isomorphism of sets

$$\sigma_{A,B}^{\text{Sets}}: A \times B \xrightarrow{\sim} B \times A,$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

01DP 7. *Distributivity Over Coproducts.* We have isomorphisms of sets

$$\begin{aligned} \partial_\ell^{\text{Sets}}: A \times (B \amalg C) &\xrightarrow{\sim} (A \times B) \amalg (A \times C), \\ \partial_r^{\text{Sets}}: (A \amalg B) \times C &\xrightarrow{\sim} (A \times C) \amalg (B \times C), \end{aligned}$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

- 0010 8. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{aligned}\zeta_\ell^{\text{Sets}} : \emptyset \times A &\xrightarrow{\sim} \emptyset, \\ \zeta_r^{\text{Sets}} : A \times \emptyset &\xrightarrow{\sim} \emptyset,\end{aligned}$$

natural in $A \in \text{Obj}(\text{Sets})$.

- 0011 9. *Distributivity Over Unions.* Let X be a set. For each $U, V, W \in \mathcal{P}(X)$, we have equalities

$$\begin{aligned}U \times (V \cup W) &= (U \times V) \cup (U \times W), \\ (U \cup V) \times W &= (U \times W) \cup (V \times W)\end{aligned}$$

of subsets of $\mathcal{P}(X \times X)$.

- 0012 10. *Distributivity Over Intersections.* Let X be a set. For each $U, V, W \in \mathcal{P}(X)$, we have equalities

$$\begin{aligned}U \times (V \cap W) &= (U \times V) \cap (U \times W), \\ (U \cap V) \times W &= (U \times W) \cap (V \times W)\end{aligned}$$

of subsets of $\mathcal{P}(X \times X)$.

- 0014 11. *Distributivity Over Differences.* Let X be a set. For each $U, V, W \in \mathcal{P}(X)$, we have equalities

$$\begin{aligned}U \times (V \setminus W) &= (U \times V) \setminus (U \times W), \\ (U \setminus V) \times W &= (U \times W) \setminus (V \times W)\end{aligned}$$

of subsets of $\mathcal{P}(X \times X)$.

- 0015 12. *Distributivity Over Symmetric Differences.* Let X be a set. For each $U, V, W \in \mathcal{P}(X)$, we have equalities

$$\begin{aligned}U \times (V \Delta W) &= (U \times V) \Delta (U \times W), \\ (U \Delta V) \times W &= (U \times W) \Delta (V \times W)\end{aligned}$$

of subsets of $\mathcal{P}(X \times X)$.

0013 13. *Middle-Four Exchange with Respect to Intersections.* The diagram

$$\begin{array}{ccc}
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \xrightarrow{\cap \times \cap} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \mathcal{P}_{X,X}^\times \times \mathcal{P}_{X,X}^\times \downarrow & & \downarrow \mathcal{P}_{X,X}^\times \\
 \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) & \xrightarrow{\cap} & \mathcal{P}(X \times X)
 \end{array}$$

commutes, i.e. we have

$$(U \times V) \cap (W \times T) = (U \cap V) \times (W \cap T).$$

for each $U, V, W, T \in \mathcal{P}(X)$.

0016 14. *Symmetric Monoidality.* The 8-tuple $(\mathbf{Sets}, \times, \text{pt}, \mathbf{Sets}(-1, -2), \alpha^{\mathbf{Sets}}, \lambda^{\mathbf{Sets}}, \rho^{\mathbf{Sets}}, \sigma^{\mathbf{Sets}})$ is a closed symmetric monoidal category.

0017 15. *Symmetric Bimonoidality.* The 18-tuple

$$\begin{aligned}
 & \left(\mathbf{Sets}, \coprod, \times, \emptyset, \text{pt}, \mathbf{Sets}(-1, -2), \alpha^{\mathbf{Sets}}, \lambda^{\mathbf{Sets}}, \rho^{\mathbf{Sets}}, \sigma^{\mathbf{Sets}}, \right. \\
 & \left. \alpha^{\mathbf{Sets}, \coprod}, \lambda^{\mathbf{Sets}, \coprod}, \rho^{\mathbf{Sets}, \coprod}, \sigma^{\mathbf{Sets}, \coprod}, \delta_\ell^{\mathbf{Sets}}, \delta_r^{\mathbf{Sets}}, \zeta_\ell^{\mathbf{Sets}}, \zeta_r^{\mathbf{Sets}} \right),
 \end{aligned}$$

is a symmetric closed bimonoidal category, where $\alpha^{\mathbf{Sets}, \coprod}, \lambda^{\mathbf{Sets}, \coprod}, \rho^{\mathbf{Sets}, \coprod}$, and $\sigma^{\mathbf{Sets}, \coprod}$ are the natural transformations from **Items 3** to **5** of **Definition 4.2.3.1.3**.

Proof. Item 1, Functoriality: This follows from Limits and Colimits, ?? of ??.

Item 2, Adjointness: We prove only that there's an adjunction $- \times B \dashv \mathbf{Sets}(B, -)$, witnessed by a bijection

$$\mathbf{Sets}(A \times B, C) \cong \mathbf{Sets}(A, \mathbf{Sets}(B, C)),$$

natural in $B, C \in \text{Obj}(\mathbf{Sets})$, as the proof of the existence of the adjunction $A \times - \dashv \mathbf{Sets}(A, -)$ follows almost exactly in the same way.

01Z6 • *Map I.* We define a map

$$\Phi_{B,C}: \mathbf{Sets}(A \times B, C) \rightarrow \mathbf{Sets}(A, \mathbf{Sets}(B, C)),$$

by sending a function

$$\xi: A \times B \rightarrow C$$

to the function

$$\begin{aligned} \xi^\dagger: A &\longrightarrow \text{Sets}(B, C), \\ a &\mapsto (\xi_a^\dagger: B \rightarrow C), \end{aligned}$$

where we define

$$\xi_a^\dagger(b) \stackrel{\text{def}}{=} \xi(a, b)$$

for each $b \in B$. In terms of the $\llbracket a \mapsto f(a) \rrbracket$ notation of [Sets, Definition 3.1.1.2](#), we have

$$\xi^\dagger \stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket.$$

0127 • *Map II.* We define a map

$$\Psi_{B,C}: \text{Sets}(A, \text{Sets}(B, C)) \rightarrow \text{Sets}(A \times B, C)$$

given by sending a function

$$\begin{aligned} \xi: A &\longrightarrow \text{Sets}(B, C), \\ a &\mapsto (\xi_a: B \rightarrow C), \end{aligned}$$

to the function

$$\xi^\dagger: A \times B \rightarrow C$$

defined by

$$\begin{aligned} \xi^\dagger(a, b) &\stackrel{\text{def}}{=} \text{ev}_b(\text{ev}_a(\xi)) \\ &\stackrel{\text{def}}{=} \text{ev}_b(\xi_a) \\ &\stackrel{\text{def}}{=} \xi_a(b) \end{aligned}$$

for each $(a, b) \in A \times B$.

- 01Z8 • *Invertibility I.* We claim that

$$\Psi_{A,B} \circ \Phi_{A,B} = \text{id}_{\text{Sets}(A \times B, C)}.$$

Indeed, given a function $\xi: A \times B \rightarrow C$, we have

$$\begin{aligned} [\Psi_{A,B} \circ \Phi_{A,B}](\xi) &= \Psi_{A,B}(\Phi_{A,B}(\xi)) \\ &= \Psi_{A,B}(\Phi_{A,B}(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\ &= \Psi_{A,B}(\llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket) \\ &= \Psi_{A,B}(\llbracket a' \mapsto \llbracket b' \mapsto \xi(a', b') \rrbracket \rrbracket) \\ &= \llbracket (a, b) \mapsto \text{ev}_b(\text{ev}_a(\llbracket a' \mapsto \llbracket b' \mapsto \xi(a', b') \rrbracket) \rrbracket) \rrbracket \\ &= \llbracket (a, b) \mapsto \text{ev}_b(\llbracket b' \mapsto \xi(a, b') \rrbracket) \rrbracket \\ &= \llbracket (a, b) \mapsto \xi(a, b) \rrbracket \\ &= \xi. \end{aligned}$$

- 01Z9 • *Invertibility II.* We claim that

$$\Phi_{A,B} \circ \Psi_{A,B} = \text{id}_{\text{Sets}(A, \text{Sets}(B, C))}.$$

Indeed, given a function

$$\begin{aligned} \xi: A &\longrightarrow \text{Sets}(B, C), \\ a &\mapsto (\xi_a: B \rightarrow C), \end{aligned}$$

we have

$$\begin{aligned} [\Phi_{A,B} \circ \Psi_{A,B}](\xi) &\stackrel{\text{def}}{=} \Phi_{A,B}(\Psi_{A,B}(\xi)) \\ &\stackrel{\text{def}}{=} \Phi_{A,B}(\llbracket (a, b) \mapsto \xi_a(b) \rrbracket) \\ &\stackrel{\text{def}}{=} \Phi_{A,B}(\llbracket (a', b') \mapsto \xi_{a'}(b') \rrbracket) \\ &\stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \text{ev}_{(a,b)}(\llbracket (a', b') \mapsto \xi_{a'}(b') \rrbracket) \rrbracket \\ &\stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \xi_a(b) \rrbracket \rrbracket \\ &\stackrel{\text{def}}{=} \llbracket a \mapsto \xi_a \rrbracket \\ &\stackrel{\text{def}}{=} \xi. \end{aligned}$$

- 01ZA • *Naturality for Φ , Part I.* We need to show that, given a function $g: B \rightarrow$

B' , the diagram

$$\begin{array}{ccc} \text{Sets}(A \times B', C) & \xrightarrow{\Phi_{B',C}} & \text{Sets}(A, \text{Sets}(B', C)), \\ \text{id}_A \times g^* \downarrow & & \downarrow (g^*)_! \\ \text{Sets}(A \times B, C) & \xrightarrow{\Phi_{B,C}} & \text{Sets}(A, \text{Sets}(B, C)) \end{array}$$

commutes. Indeed, given a function

$$\xi: A \times B' \rightarrow C,$$

we have

$$\begin{aligned} [\Phi_{B,C} \circ (\text{id}_A \times g^*)](\xi) &= \Phi_{B,C}([\text{id}_A \times g^*](\xi)) \\ &= \Phi_{B,C}(\xi(-_1, g(-_2))) \\ &= [\xi(-_1, g(-_2))]^\dagger \\ &= \xi_{-1}^\dagger(g(-_2)) \\ &= (g^*)_!(\xi^\dagger) \\ &= (g^*)_!(\Phi_{B',C}(\xi)) \\ &= [(g^*)_! \circ \Phi_{B',C}](\xi). \end{aligned}$$

Alternatively, using the $\llbracket a \mapsto f(a) \rrbracket$ notation of [Sets, Definition 3.1.1.1.2](#), we have

$$\begin{aligned} [\Phi_{B,C} \circ (\text{id}_A \times g^*)](\xi) &= \Phi_{B,C}([\text{id}_A \times g^*](\xi)) \\ &= \Phi_{B,C}([\text{id}_A \times g^*](\llbracket (a, b') \mapsto \xi(a, b') \rrbracket)) \\ &= \Phi_{B,C}(\llbracket (a, b) \mapsto \xi(a, g(b)) \rrbracket) \\ &= \llbracket a \mapsto \llbracket b \mapsto \xi(a, g(b)) \rrbracket \rrbracket \\ &= \llbracket a \mapsto g^*(\llbracket b' \mapsto \xi(a, b') \rrbracket) \rrbracket \\ &= (g^*)_!(\llbracket a \mapsto \llbracket b' \mapsto \xi(a, b') \rrbracket \rrbracket) \\ &= (g^*)_!(\Phi_{B',C}(\llbracket (a, b') \mapsto \xi(a, b') \rrbracket)) \\ &= (g^*)_!(\Phi_{B',C}(\xi)) \\ &= [(g^*)_! \circ \Phi_{B',C}](\xi). \end{aligned}$$

- 01ZB • *Naturality for Φ , Part II.* We need to show that, given a function $b: C \rightarrow C'$, the diagram

$$\begin{array}{ccc} \text{Sets}(A \times B, C) & \xrightarrow{\Phi_{B,C}} & \text{Sets}(A, \text{Sets}(B, C)), \\ \downarrow b_! & & \downarrow (b_!)_! \\ \text{Sets}(A \times B, C') & \xrightarrow{\Phi_{B,C'}} & \text{Sets}(A, \text{Sets}(B, C')) \end{array}$$

commutes. Indeed, given a function

$$\xi: A \times B \rightarrow C,$$

we have

$$\begin{aligned} [\Phi_{B,C} \circ b_!](\xi) &= \Phi_{B,C}(b_!(\xi)) \\ &= \Phi_{B,C}(b_!(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\ &= \Phi_{B,C}(\llbracket (a, b) \mapsto b(\xi(a, b)) \rrbracket) \\ &= \llbracket a \mapsto \llbracket b \mapsto b(\xi(a, b)) \rrbracket \rrbracket \\ &= \llbracket a \mapsto b_!(\llbracket b \mapsto \xi(a, b) \rrbracket) \rrbracket \\ &= (b_!)_!(\llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket) \\ &= (b_!)_!(\Phi_{B,C}(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\ &= (b_!)_!(\Phi_{B,C}(\xi)) \\ &= [(b_!)_! \circ \Phi_{B,C}](\xi). \end{aligned}$$

- 01ZC • *Naturality for Ψ .* Since Φ is natural in each argument and Φ is a componentwise inverse to Ψ in each argument, it follows from **Categories, Item 2** of **Definition 11.9.7.1.2** that Ψ is also natural in each argument.

This finishes the proof.

Item 3, Adjointness II: This follows from the universal property of the product.

Item 4, Associativity: This is proved in the proof of **Monoidal Structures on the Category of Sets, Definition 5.1.4.1.1**.

Item 5, Unitality: This is proved in the proof of **Monoidal Structures on the Category of Sets, Definitions 5.1.5.1.1** and **5.1.6.1.1**.

Item 6, Commutativity: This is proved in the proof of **Monoidal Structures on the Category of Sets, Definition 5.1.7.1.1**.

Item 7, Distributivity Over Coproducts: This is proved in the proof of **Monoidal Structures on the Category of Sets**, Definitions 5.3.1.1.1 and 5.3.2.1.1.

Item 8, Annihilation With the Empty Set: This is proved in the proof of **Monoidal Structures on the Category of Sets**, Definitions 5.3.3.1.1 and 5.3.4.1.1.

Item 9, Distributivity Over Unions: See [Pro25c].

Item 10, Distributivity Over Intersections: See [Pro25d, Corollary 1].

Item 11, Distributivity Over Differences: See [Pro25a].

Item 12, Distributivity Over Symmetric Differences: See [Pro25b].

Item 13, Middle-Four Exchange With Respect to Intersections: See [Pro25d, Corollary 1].

Item 14, Symmetric Monoidality: This is a repetition of **Monoidal Structures on the Category of Sets**, Definition 5.1.9.1.1, and is proved there.

Item 15, Symmetric Bimonoidality: This is a repetition of **Monoidal Structures on the Category of Sets**, Definition 5.3.5.1.1, and is proved there. \square

01DR Remark 4.1.3.1.4. As shown in **Item 1** of Definition 4.1.3.1.3, the Cartesian product of sets defines a functor

$$-_1 \times -_2 : \mathbf{Sets} \times \mathbf{Sets} \rightarrow \mathbf{Sets}.$$

This functor is the $(k, \ell) = (-1, -1)$ case of a family of functors

$$\otimes_{k, \ell} : \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}) \times \mathbf{Mon}_{\mathbb{E}_\ell}(\mathbf{Sets}) \rightarrow \mathbf{Mon}_{\mathbb{E}_{k+\ell}}(\mathbf{Sets})$$

of tensor products of \mathbb{E}_k -monoid objects on **Sets** with \mathbb{E}_ℓ -monoid objects on **Sets**; see ??.

01DS Remark 4.1.3.1.5. We may state the equalities in **Items 9** to **12** of Definition 4.1.3.1.3 as the commutativity of the following diagrams:

$$\begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \Delta_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \swarrow & & \searrow \mu_{\mathbf{Sets}}^{\mathcal{P}} \\
 \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \downarrow \text{id}_{\mathcal{P}(X)} \times \cup & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) & \\
 \downarrow \mathcal{P}_{XX}^\times & \swarrow \cup & \downarrow \cup \\
 & \mathcal{P}(X \times X) &
 \end{array}
 \qquad
 \begin{array}{ccc}
 & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \Delta_{\mathcal{P}(X)} \swarrow & & \searrow \mu_{\mathbf{Sets}}^{\mathcal{P}} \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \downarrow \cup \times \text{id}_{\mathcal{P}(X)} & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) & \\
 \downarrow \mathcal{P}_{XX}^\times & \swarrow \cup & \downarrow \cup \\
 & \mathcal{P}(X \times X) &
 \end{array}$$

$$\begin{array}{ccc}
& (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\Delta_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \nearrow & \mu_4^{\text{Sets}} & \nwarrow \\
\mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\downarrow \text{id}_{\mathcal{P}(X)} \times \cap & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
\mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) \\
& \searrow \mathcal{P}_{XX}^\times & \swarrow \cap \\
& \mathcal{P}(X \times X) &
\end{array}$$

$$\begin{array}{ccc}
& (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \Delta_{\mathcal{P}(X)} \nearrow & \mu_4^{\text{Sets}} & \nwarrow \\
(\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\downarrow \cap \times \text{id}_{\mathcal{P}(X)} & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
\mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) \\
& \searrow \mathcal{P}_{XX}^\times & \swarrow \cap \\
& \mathcal{P}(X \times X) &
\end{array}$$

$$\begin{array}{ccc}
& (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\Delta_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \nearrow & \mu_4^{\text{Sets}} & \nwarrow \\
\mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\downarrow \text{id}_{\mathcal{P}(X)} \times \setminus & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
\mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) \\
& \searrow \mathcal{P}_{XX}^\times & \swarrow \setminus \\
& \mathcal{P}(X \times X) &
\end{array}$$

$$\begin{array}{ccc}
& (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \Delta_{\mathcal{P}(X)} \nearrow & \mu_4^{\text{Sets}} & \nwarrow \\
(\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\downarrow \setminus \times \text{id}_{\mathcal{P}(X)} & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
\mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) \\
& \searrow \mathcal{P}_{XX}^\times & \swarrow \setminus \\
& \mathcal{P}(X \times X) &
\end{array}$$

$$\begin{array}{ccc}
& (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\Delta_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \nearrow & \mu_4^{\text{Sets}} & \nwarrow \\
\mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\downarrow \text{id}_{\mathcal{P}(X)} \times \Delta & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
\mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) \\
& \searrow \mathcal{P}_{XX}^\times & \swarrow \Delta \\
& \mathcal{P}(X \times X) &
\end{array}$$

$$\begin{array}{ccc}
& (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\text{id}_{\mathcal{P}(X)} \times \text{id}_{\mathcal{P}(X)} \times \Delta_{\mathcal{P}(X)} \nearrow & \mu_4^{\text{Sets}} & \nwarrow \\
(\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & (\mathcal{P}(X) \times \mathcal{P}(X)) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
\downarrow \Delta \times \text{id}_{\mathcal{P}(X)} & & \downarrow \mathcal{P}_{XX}^\times \times \mathcal{P}_{XX}^\times \\
\mathcal{P}(X) \times \mathcal{P}(X) & & \mathcal{P}(X \times X) \times \mathcal{P}(X \times X) \\
& \searrow \mathcal{P}_{XX}^\times & \swarrow \Delta \\
& \mathcal{P}(X \times X) &
\end{array}$$

0018 4.1.4 Pullbacks

Let A , B , and C be sets and let $f: A \rightarrow C$ and $g: B \rightarrow C$ be functions.

0019 **Definition 4.1.4.1.1.** The **pullback of A and B over C along f and g** ³ is the pullback of A and B over C along f and g in **Sets** as in Limits and Colimits, ??.

01DT **Construction 4.1.4.1.2.** Concretely, the pullback of A and B over C along f and g is the pair $(A \times_C B, \{\text{pr}_1, \text{pr}_2\})$ consisting of:

³Further Terminology: Also called the **fibre product of A and B over C along f and g** .

01DU 1. *The Limit.* The set $A \times_C B$ defined by

$$A \times_C B \stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\}.$$

01DV 2. *The Cone.* The maps⁴

$$\begin{aligned} \text{pr}_1 &: A \times_C B \rightarrow A, \\ \text{pr}_2 &: A \times_C B \rightarrow B \end{aligned}$$

defined by

$$\begin{aligned} \text{pr}_1(a, b) &\stackrel{\text{def}}{=} a, \\ \text{pr}_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each $(a, b) \in A \times_C B$.

Proof. We claim that $A \times_C B$ is the categorical pullback of A and B over C with respect to (f, g) in **Sets**. First we need to check that the relevant pullback diagram commutes, i.e. that we have

$$f \circ \text{pr}_1 = g \circ \text{pr}_2, \quad \begin{array}{ccc} A \times_C B & \xrightarrow{\text{pr}_2} & B \\ \text{pr}_1 \downarrow & & \downarrow g \\ A & \xrightarrow{f} & C. \end{array}$$


Indeed, given $(a, b) \in A \times_C B$, we have

$$\begin{aligned} [f \circ \text{pr}_1](a, b) &= f(\text{pr}_1(a, b)) \\ &= f(a) \\ &= g(b) \\ &= g(\text{pr}_2(a, b)) \\ &= [g \circ \text{pr}_2](a, b), \end{aligned}$$


where $f(a) = g(b)$ since $(a, b) \in A \times_C B$. Next, we prove that $A \times_C B$ satisfies

⁴*Further Notation:* Also written $\text{pr}_1^{A \times_C B}$ and $\text{pr}_2^{A \times_C B}$.

the universal property of the pullback. Suppose we have a diagram of the form



in Sets. Then there exists a unique map $\phi: P \rightarrow A \times_C B$ making the diagram



commute, being uniquely determined by the conditions

$$\begin{aligned} \text{pr}_1 \circ \phi &= p_1, \\ \text{pr}_2 \circ \phi &= p_2 \end{aligned}$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$, where we note that $(p_1(x), p_2(x)) \in A \times B$ indeed lies in $A \times_C B$ by the condition

$$f \circ p_1 = g \circ p_2,$$

which gives

$$f(p_1(x)) = g(p_2(x))$$

for each $x \in P$, so that $(p_1(x), p_2(x)) \in A \times_C B$. □

01DW Remark 4.1.4.1.3. It is common practice to write $A \times_C B$ for the pullback of A and B over C along f and g , omitting the maps f and g from the notation and instead leaving them implicit, to be understood from the context.

However, the set $A \times_C B$ depends very much on the maps f and g , and sometimes it is necessary or useful to note this dependence explicitly. In such situations, we will write $A \times_{f,C,g} B$ or $A \times_C^{f,g} B$ for $A \times_C B$.

001A Example 4.1.4.1.4. Here are some examples of pullbacks of sets.

001B I. *Unions via Intersections.* Let X be a set. We have

$$A \cap B \cong A \times_{A \cup B} B,$$

$$\begin{array}{ccc} A \cap B & \longrightarrow & B \\ \downarrow & \lrcorner & \downarrow \iota_B \\ A & \xrightarrow{\iota_A} & A \cup B \end{array}$$

for each $A, B \in \mathcal{P}(X)$.

Proof. **Item 1, Unions via Intersections:** Indeed, we have

$$\begin{aligned} A \times_{A \cup B} B &\cong \{(x, y) \in A \times B \mid x = y\} \\ &\cong A \cap B. \end{aligned}$$

This finishes the proof. □

001C Proposition 4.1.4.1.5. Let A, B, C , and X be sets.

001D I. *Functoriality.* The assignment $(A, B, C, f, g) \mapsto A \times_{f,C,g} B$ defines a functor

$$-_1 \times_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where \mathcal{P} is the category that looks like this:

$$\begin{array}{ccc} & & \bullet \\ & & \downarrow \\ \bullet & \longrightarrow & \bullet \end{array}$$

In particular, the action on morphisms of $-_1 \times_{-3} -_1$ is given by sending a morphism

$$\begin{array}{ccccc}
 A \times_C B & \xrightarrow{\quad} & B & & \\
 \downarrow & \lrcorner & \downarrow g & \searrow \psi & \\
 & & A' \times_{C'} B' & \xrightarrow{\quad} & B' \\
 & & \downarrow & & \downarrow g' \\
 A & \xrightarrow{f} & C & & \\
 \searrow \phi & & \searrow \chi & & \\
 & & A' & \xrightarrow{f'} & C'
 \end{array}$$

in $\text{Fun}(\mathcal{P}, \text{Sets})$ to the map $\xi: A \times_C B \xrightarrow{\exists!} A' \times_{C'} B'$ given by

$$\xi(a, b) \stackrel{\text{def}}{=} (\phi(a), \psi(b))$$

for each $(a, b) \in A \times_C B$, which is the unique map making the diagram

$$\begin{array}{ccccc}
 A \times_C B & \xrightarrow{\quad} & B & & \\
 \downarrow & \dashv \lrcorner & \downarrow g & \searrow \psi & \\
 & & A' \times_{C'} B' & \xrightarrow{\quad} & B' \\
 & & \downarrow & & \downarrow g' \\
 A & \xrightarrow{f} & C & & \\
 \searrow \phi & & \searrow \chi & & \\
 & & A' & \xrightarrow{f'} & C'
 \end{array}$$

commute.

01DX 2. *Adjointness I.* We have adjunctions

$$\begin{aligned} (A \times_X - \dashv \mathbf{Sets}_{/X}(A, -)) : \mathbf{Sets}_{/X} &\begin{array}{c} \xrightarrow{A \times_X -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_{/X}(A, -)} \end{array} \mathbf{Sets}_{/X}, \\ (- \times_X B \dashv \mathbf{Sets}_{/X}(B, -)) : \mathbf{Sets}_{/X} &\begin{array}{c} \xrightarrow{- \times_X B} \\ \perp \\ \xleftarrow{\mathbf{Sets}_{/X}(B, -)} \end{array} \mathbf{Sets}_{/X}, \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \mathbf{Sets}_{/X}(A \times_X B, C) &\cong \mathbf{Sets}_{/X}(A, \mathbf{Sets}_{/X}(B, C)), \\ \mathbf{Sets}_{/X}(A \times_X B, C) &\cong \mathbf{Sets}_{/X}(B, \mathbf{Sets}_{/X}(A, C)), \end{aligned}$$

natural in $(A, \phi_A), (B, \phi_B), (C, \phi_C) \in \text{Obj}(\mathbf{Sets}_{/X})$, where $\mathbf{Sets}_{/X}(A, B)$ is the object of $\mathbf{Sets}_{/X}$ consisting of (see Fibred Sets, ??):

- *The Set.* The set $\mathbf{Sets}_{/X}(A, B)$ defined by

$$\mathbf{Sets}_{/X}(A, B) \stackrel{\text{def}}{=} \coprod_{x \in X} \mathbf{Sets}(\phi_A^{-1}(x), \phi_B^{-1}(x))$$

- *The Map to X.* The map

$$\phi_{\mathbf{Sets}_{/X}(A, B)} : \mathbf{Sets}_{/X}(A, B) \rightarrow X$$

defined by

$$\phi_{\mathbf{Sets}_{/X}(A, B)}(x, f) \stackrel{\text{def}}{=} x$$

for each $(x, f) \in \mathbf{Sets}_{/X}(A, B)$.

01ZD 3. *Adjointness II.* We have an adjunction


$$(\Delta_{\mathbf{Sets}_{/X}} \dashv -_1 \times -_2) : \mathbf{Sets}_{/X} \begin{array}{c} \xrightarrow{\Delta_{\mathbf{Sets}_{/X}}} \\ \perp \\ \xleftarrow{-_1 \times -_2} \end{array} \mathbf{Sets}_{/X} \times \mathbf{Sets}_{/X},$$

witnessed by a bijection

$$\text{Hom}_{\mathbf{Sets}_{/X} \times \mathbf{Sets}_{/X}}((A, A), (B, C)) \cong \mathbf{Sets}_{/X}(A, B \times_X C),$$

natural in $A \in \text{Obj}(\mathbf{Sets}_{/X})$ and in $(B, C) \in \text{Obj}(\mathbf{Sets}_{/X} \times \mathbf{Sets}_{/X})$.


001E 4. *Associativity.* Given a diagram




in **Sets**, we have isomorphisms of sets

$$(A \times_X B) \times_Y C \cong (A \times_X B) \times_B (B \times_Y C) \cong A \times_X (B \times_Y C),$$

where these pullbacks are built as in the diagrams



01DY 5. *Interaction With Composition.* Given a diagram




in **Sets**, we have isomorphisms of sets

$$\begin{aligned}
 X \times_K^{f \circ \phi, g \circ \psi} Y &\cong \left(X \times_A^{\phi, q_1} \left(A \times_K^{f, g} B \right) \right) \times_{A \times_K^{f, g} B}^{p_2, p_1} \left(\left(A \times_K^{f, g} B \right) \times_B^{q_2, \psi} Y \right) \\
 &\cong X \times_A^{\phi, p} \left(\left(A \times_K^{f, g} B \right) \times_B^{q_2, \psi} Y \right) \\
 &\cong \left(X \times_A^{\phi, q_1} \left(A \times_K^{f, g} B \right) \right) \times_B^{q, \psi} Y
 \end{aligned}$$

where

$$\begin{aligned}
 q_1 &= \text{pr}_1^{A \times_K^{f,g} B}, & q_2 &= \text{pr}_2^{A \times_K^{f,g} B}, \\
 p_1 &= \text{pr}_1^{(A \times_K^{f,g} B) \times_Y^{q_2, \psi}}, & p_2 &= \text{pr}_2^{X \times_{A \times_K^{f,g} B}^{q_1} (A \times_K^{f,g} B)}, \\
 p &= q_1 \circ \text{pr}_1^{(A \times_K^{f,g} B) \times_B^{q_2, \psi} Y}, & q &= q_2 \circ \text{pr}_2^{X \times_A^{q_1} (A \times_K^{f,g} B)},
 \end{aligned}$$

and where these pullbacks are built as in the following diagrams:



001F 6. *Unitality*. We have isomorphisms of sets

$$\begin{array}{ccc}
 A & \xlongequal{\quad} & A \\
 \downarrow f & \lrcorner & \downarrow f \\
 X & \xlongequal{\quad} & X
 \end{array}
 \quad
 \begin{array}{l}
 \lambda_A^{\mathbf{Sets}/X} : X \times_X A \xrightarrow{\sim} A, \\
 \rho_A^{\mathbf{Sets}/X} : A \times_X X \xrightarrow{\sim} A,
 \end{array}
 \quad
 \begin{array}{ccc}
 A & \xrightarrow{f} & X \\
 \parallel & \lrcorner & \parallel \\
 X & \xrightarrow{f} & X,
 \end{array}$$

natural in $(A, f) \in \mathbf{Obj}(\mathbf{Sets}/X)$.

001G 7. *Commutativity*. We have an isomorphism of sets

$$\begin{array}{ccc}
 A \times_X B & \longrightarrow & B \\
 \downarrow & \lrcorner & \downarrow g \\
 A & \xrightarrow{f} & X,
 \end{array}
 \quad
 \sigma_{A,B}^{\mathbf{Sets}/X} : A \times_X B \xrightarrow{\sim} B \times_X A
 \quad
 \begin{array}{ccc}
 B \times_X A & \longrightarrow & A \\
 \downarrow & \lrcorner & \downarrow f \\
 B & \xrightarrow{g} & X,
 \end{array}$$

natural in $(A, f), (B, g) \in \mathbf{Obj}(\mathbf{Sets}/X)$.

01DZ 8. *Distributivity Over Coproducts*. Let A, B , and C be sets and let $\phi_A : A \rightarrow X$, $\phi_B : B \rightarrow X$, and $\phi_C : C \rightarrow X$ be morphisms of sets. We have isomorphisms of sets

$$\begin{aligned}
 \delta_\ell^{\mathbf{Sets}/X} : A \times_X (B \amalg C) &\xrightarrow{\sim} (A \times_X B) \amalg (A \times_X C), \\
 \delta_r^{\mathbf{Sets}/X} : (A \amalg B) \times_X C &\xrightarrow{\sim} (A \times_X C) \amalg (B \times_X C),
 \end{aligned}$$

as in the diagrams

$$\begin{array}{ccc}
 (A \times_X B) \amalg (A \times_X C) & \longrightarrow & B \amalg C \\
 \downarrow & \lrcorner & \downarrow \phi_B \amalg \phi_C \\
 A & \xrightarrow{\phi_A} & X
 \end{array}
 \quad
 \begin{array}{ccc}
 (A \times_X C) \amalg (B \times_X C) & \longrightarrow & C \\
 \downarrow & \lrcorner & \downarrow \phi_C \\
 A \amalg B & \xrightarrow{\phi_A \amalg \phi_B} & X
 \end{array}$$

natural in $A, B, C \in \mathbf{Obj}(\mathbf{Sets}/X)$.

001H 9. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{array}{ccc}
 \emptyset & \longrightarrow & \emptyset \\
 \downarrow \lrcorner & & \downarrow \\
 A & \xrightarrow{f} & X,
 \end{array}
 \quad
 \begin{array}{l}
 \zeta_\ell^{\mathbf{Sets}/X} : A \times_X \emptyset \xrightarrow{\sim} \emptyset, \\
 \zeta_r^{\mathbf{Sets}/X} : \emptyset \times_X A \xrightarrow{\sim} \emptyset,
 \end{array}
 \quad
 \begin{array}{ccc}
 \emptyset & \longrightarrow & A \\
 \downarrow \lrcorner & & \downarrow f \\
 \emptyset & \longrightarrow & X,
 \end{array}$$

natural in $(A, f) \in \text{Obj}(\mathbf{Sets}/X)$.

001J 10. *Interaction With Products.* We have an isomorphism of sets

$$\begin{array}{ccc}
 A \times_{\text{pt}} B \cong A \times B, & & \begin{array}{ccc} A \times B & \longrightarrow & B \\ \downarrow \lrcorner & & \downarrow !_B \\ A & \xrightarrow{!_A} & \text{pt.} \end{array}
 \end{array}$$

001K II. *Symmetric Monoidality.* The 8-tuple $(\mathbf{Sets}/X, \times_X, X, \mathbf{Sets}/X, \alpha^{\mathbf{Sets}/X}, \lambda^{\mathbf{Sets}/X}, \rho^{\mathbf{Sets}/X}, \sigma^{\mathbf{Sets}/X})$ is a symmetric closed monoidal category.

Proof. Item 1, Functoriality: This is a special case of functoriality of co/limits, Limits and Colimits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pullback diagram.

Item 2, Adjointness I: This is a repetition of Fibred Sets, ?? of ??, and is proved there.

Item 3, Adjointness II: This follows from the universal property of the product (pullbacks are products in \mathbf{Sets}/X).

Item 4, Associativity: We have

$$\begin{aligned}
 (A \times_X B) \times_Y C &\cong \{(a, b), c) \in (A \times_X B) \times C \mid h(b) = k(c)\} \\
 &\cong \{(a, b), c) \in (A \times B) \times C \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\
 &\cong \{(a, (b, c)) \in A \times (B \times_Y C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\
 &\cong \{(a, (b, c)) \in A \times (B \times_Y C) \mid f(a) = g(b)\} \\
 &\cong A \times_X (B \times_Y C)
 \end{aligned}$$

and

$$(A \times_X B) \times_B (B \times_Y C) \cong \{(a, b), (b', c)) \in (A \times_X B) \times (B \times_Y C) \mid b = b'\}$$

$$\begin{aligned}
&\cong \left\{ ((a, b), (b', c)) \in (A \times B) \times (B \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } b(b') = k(c) \end{array} \right\} \\
&\cong \left\{ (a, (b, (b', c))) \in A \times (B \times (B \times C)) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } b(b') = k(c) \end{array} \right\} \\
&\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times B) \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } b(b') = k(c) \end{array} \right\} \\
&\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times_B B) \times C) \mid \begin{array}{l} f(a) = g(b) \text{ and } \\ b(b') = k(c) \end{array} \right\} \\
&\cong \{ (a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } b(b) = k(c) \} \\
&\cong A \times_X (B \times_Y C),
\end{aligned}$$

where we have used **Item 6** for the isomorphism $B \times_B B \cong B$.

Item 5, Interaction With Composition: By **Item 4**, it suffices to construct only the isomorphism

$$X \times_K^{f \circ \phi, g \circ \psi} Y \cong \left(X \times_A^{\phi, q_1} \left(A \times_K^{f, g} B \right) \right) \times_{A \times_K^{f, g} B}^{p_2, p_1} \left(\left(A \times_K^{f, g} B \right) \times_B^{q_2, \psi} Y \right).$$

We have

$$\begin{aligned}
\left(X \times_A^{\phi, q_1} \left(A \times_K^{f, g} B \right) \right) &\stackrel{\text{def}}{=} \left\{ (x, (a, b)) \in X \times \left(A \times_K^{f, g} B \right) \mid \phi(x) = q_1(a, b) \right\} \\
&\stackrel{\text{def}}{=} \left\{ (x, (a, b)) \in X \times \left(A \times_K^{f, g} B \right) \mid \phi(x) = a \right\} \\
&\cong \left\{ (x, (a, b)) \in X \times (A \times B) \mid \phi(x) = a \text{ and } f(a) = g(b) \right\}, \\
\left(\left(A \times_K^{f, g} B \right) \times_B^{q_2, \psi} Y \right) &\stackrel{\text{def}}{=} \left\{ ((a, b), y) \in \left(A \times_K^{f, g} B \right) \times Y \mid q_2(a, b) = \psi(y) \right\} \\
&\stackrel{\text{def}}{=} \left\{ ((a, b), y) \in \left(A \times_K^{f, g} B \right) \times Y \mid b = \psi(y) \right\} \\
&\cong \left\{ ((a, b), y) \in (A \times B) \times Y \mid b = \psi(y) \text{ and } f(a) = g(b) \right\},
\end{aligned}$$

so writing

$$\begin{aligned}
S &= \left(X \times_A^{\phi, q_1} \left(A \times_K^{f, g} B \right) \right) \\
S' &= \left(\left(A \times_K^{f, g} B \right) \times_B^{q_2, \psi} Y \right),
\end{aligned}$$

we have

$$\begin{aligned}
S \times_{A \times_K^{f, g} B}^{p_2, p_1} S' &\stackrel{\text{def}}{=} \left\{ ((x, (a, b)), ((a', b'), y)) \in S \times S' \mid p_1(x, (a, b)) = p_2((a', b'), y) \right\} \\
&\stackrel{\text{def}}{=} \left\{ ((x, (a, b)), ((a', b'), y)) \in S \times S' \mid (a, b) = (a', b') \right\} \\
&\cong \left\{ ((x, a, b, y)) \in X \times A \times B \times Y \mid \phi(x) = a, \psi(y) = b, \text{ and } f(a) = g(b) \right\}
\end{aligned}$$

$$\begin{aligned} &\cong \{((x, a, b, y)) \in X \times A \times B \times Y \mid f(\phi(x)) = g(\psi(y))\} \\ &\stackrel{\text{def}}{=} X \times_K Y. \end{aligned}$$

This finishes the proof.

Item 6, Unitality: We have

$$\begin{aligned} X \times_X A &\cong \{(x, a) \in X \times A \mid f(a) = x\}, \\ A \times_X X &\cong \{(a, x) \in X \times A \mid f(a) = x\}, \end{aligned}$$

which are isomorphic to A via the maps $(x, a) \mapsto a$ and $(a, x) \mapsto a$. The proof of the naturality of $\lambda^{\text{Sets}/X}$ and $\rho^{\text{Sets}/X}$ is omitted.

Item 7, Commutativity: We have

$$\begin{aligned} A \times_C B &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\} \\ &= \{(a, b) \in A \times B \mid g(b) = f(a)\} \\ &\cong \{(b, a) \in B \times A \mid g(b) = f(a)\} \\ &\stackrel{\text{def}}{=} B \times_C A. \end{aligned}$$

The proof of the naturality of $\sigma^{\text{Sets}/X}$ is omitted.

Item 8, Distributivity Over Coproducts: We have

$$\begin{aligned} A \times_X (B \amalg C) &\stackrel{\text{def}}{=} \{(a, z) \in A \times (B \amalg C) \mid \phi_A(a) = \phi_{B \amalg C}(z)\} \\ &= \{(a, z) \in A \times (B \amalg C) \mid z = (0, b) \text{ and } \phi_A(a) = \phi_B(b)\} \\ &\quad \cup \{(a, z) \in A \times (B \amalg C) \mid z = (1, c) \text{ and } \phi_A(a) = \phi_C(c)\} \\ &= \{(a, z) \in A \times (B \amalg C) \mid z = (0, b) \text{ and } \phi_A(a) = \phi_B(b)\} \\ &\quad \cup \{(a, z) \in A \times (B \amalg C) \mid z = (1, c) \text{ and } \phi_A(a) = \phi_C(c)\} \\ &\cong \{(a, b) \in A \times B \mid \phi_A(a) = \phi_B(b)\} \\ &\quad \cup \{(a, c) \in A \times C \mid \phi_A(a) = \phi_C(c)\} \\ &\stackrel{\text{def}}{=} (A \times_X B) \cup (A \times_X C) \\ &\cong (A \times_X B) \amalg (A \times_X C), \end{aligned}$$

with the construction of the isomorphism

$$\delta_r^{\text{Sets}/X} : (A \amalg B) \times_X C \xrightarrow{\sim} (A \times_X C) \amalg (B \times_X C)$$

being similar. The proof of the naturality of $\delta_\ell^{\text{Sets}/X}$ and $\delta_r^{\text{Sets}/X}$ is omitted.

Item 9, Annihilation With the Empty Set: We have

$$\begin{aligned} A \times_X \emptyset &\stackrel{\text{def}}{=} \{(a, b) \in A \times \emptyset \mid f(a) = g(b)\} \\ &= \{k \in \emptyset \mid f(a) = g(b)\} \\ &= \emptyset, \end{aligned}$$

and similarly for $\emptyset \times_X A$, where we have used **Item 8** of **Definition 4.1.3.1.3**. The proof of the naturality of $\zeta_\ell^{\text{Sets}/X}$ and $\zeta_r^{\text{Sets}/X}$ is omitted.

Item 10, Interaction With Products: We have

$$\begin{aligned} A \times_{\text{pt}} B &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid !_A(a) = !_B(b)\} \\ &\stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid \star = \star\} \\ &= \{(a, b) \in A \times B\} \\ &= A \times B. \end{aligned}$$

Item 11, Symmetric Monoidality: Omitted. □

001L 4.1.5 Equalisers

Let A and B be sets and let $f, g: A \rightrightarrows B$ be functions.

001M Definition 4.1.5.1.1. The **equaliser of f and g** is the equaliser of f and g in **Sets** as in Limits and Colimits, ??.

01E0 Construction 4.1.5.1.2. Concretely, the equaliser of f and g is the pair $(\text{Eq}(f, g), \text{eq}(f, g))$ consisting of:

01E1 1. *The Limit.* The set $\text{Eq}(f, g)$ defined by

$$\text{Eq}(f, g) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\}.$$

01E2 2. *The Cone.* The inclusion map

$$\text{eq}(f, g): \text{Eq}(f, g) \rightarrow A.$$

Proof. We claim that $\text{Eq}(f, g)$ is the categorical equaliser of f and g in **Sets**. First we need to check that the relevant equaliser diagram commutes, i.e. that we have

$$f \circ \text{eq}(f, g) = g \circ \text{eq}(f, g),$$

which indeed holds by the definition of the set $\text{Eq}(f, g)$. Next, we prove that $\text{Eq}(f, g)$ satisfies the universal property of the equaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A \xrightleftharpoons[f]{f} B \\ & \nearrow e & \\ E & & \end{array}$$

in Sets. Then there exists a unique map $\phi: E \rightarrow \text{Eq}(f, g)$ making the diagram

$$\begin{array}{ccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A \xrightleftharpoons[f]{f} B \\ \uparrow \phi \quad \exists! & \nearrow e & \\ E & & \end{array}$$

commute, being uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

via

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g)$ by the condition

$$f \circ e = g \circ e,$$

which gives

$$f(e(x)) = g(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g)$. □

001N Proposition 4.1.5.1.3. Let A, B , and C be sets.

001P 1. *Associativity.* We have isomorphisms of sets⁵

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}$$

⁵That is, the following three ways of forming “the” equaliser of (f, g, h) agree:

01ZE 1. Take the equaliser of (f, g, h) , i.e. the limit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B$$

in Sets.

01ZF 2. First take the equaliser of f and g , forming a diagram

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B$$

and then take the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{h} \end{array} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) = \text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))$$

of $\text{Eq}(f, g)$.

01ZG 3. First take the equaliser of g and h , forming a diagram

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{h} \end{array} B$$

and then take the equaliser of the composition

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) = \text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))$$

of $\text{Eq}(g, h)$.

where $\text{Eq}(f, g, b)$ is the limit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{} \\ \xrightarrow{b} \end{array} B$$

in **Sets**, being explicitly given by

$$\text{Eq}(f, g, b) \cong \{a \in A \mid f(a) = g(a) = b(a)\}.$$

001Q 4. *Unitality*. We have an isomorphism of sets

$$\text{Eq}(f, f) \cong A.$$

001R 5. *Commutativity*. We have an isomorphism of sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

001S 6. *Interaction With Composition*. Let

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{} \\ \xrightarrow{k} \end{array} B \begin{array}{c} \xrightarrow{b} \\ \xrightarrow[k]{} \end{array} C$$

be functions. We have an inclusion of sets

$$\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) \subset \text{Eq}(h \circ f, k \circ g),$$

where $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$ is the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{} \\ \xrightarrow{k} \end{array} B \begin{array}{c} \xrightarrow{b} \\ \xrightarrow[k]{} \end{array} C.$$

Proof. **Item 1, Associativity**: We first prove that $\text{Eq}(f, g, b)$ is indeed given by

$$\text{Eq}(f, g, b) \cong \{a \in A \mid f(a) = g(a) = b(a)\}.$$

Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} & \xrightarrow{\text{eq}(f, g, b)} & A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{} \\ \xrightarrow{b} \end{array} B \\ E \nearrow e & & \end{array}$$

in Sets. Then there exists a unique map $\phi: E \rightarrow \text{Eq}(f, g, h)$, uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

being necessarily given by

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g, h)$ by the condition

$$f \circ e = g \circ e = h \circ e,$$

which gives

$$f(e(x)) = g(e(x)) = h(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g, h)$.

We now check the equalities

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) \cong \text{Eq}(f, g, h) \cong \text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)).$$

Indeed, we have

$$\begin{aligned} \text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) &\cong \{x \in \text{Eq}(g, h) \mid [f \circ \text{eq}(g, h)](a) = [g \circ \text{eq}(g, h)](a)\} \\ &\cong \{x \in \text{Eq}(g, h) \mid f(a) = g(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) \text{ and } g(a) = h(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\ &\cong \text{Eq}(f, g, h). \end{aligned}$$

Similarly, we have

$$\begin{aligned} \text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) &\cong \{x \in \text{Eq}(f, g) \mid [f \circ \text{eq}(f, g)](a) = [h \circ \text{eq}(f, g)](a)\} \\ &\cong \{x \in \text{Eq}(f, g) \mid f(a) = h(a)\} \\ &\cong \{x \in A \mid f(a) = h(a) \text{ and } f(a) = g(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\ &\cong \text{Eq}(f, g, h). \end{aligned}$$

Item 4, Unitality: Indeed, we have

$$\begin{aligned} \text{Eq}(f, f) &\stackrel{\text{def}}{=} \{a \in A \mid f(a) = f(a)\} \\ &= A. \end{aligned}$$

Item 5, Commutativity: Indeed, we have

$$\text{Eq}(f, g) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\}$$

$$= \{a \in A \mid g(a) = f(a)\} \\ \stackrel{\text{def}}{=} \text{Eq}(g, f).$$

Item 6, Interaction With Composition: Indeed, we have

$$\begin{aligned} \text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) &\cong \{a \in \text{Eq}(f, g) \mid h(f(a)) = k(g(a))\} \\ &\cong \{a \in A \mid f(a) = g(a) \text{ and } h(f(a)) = k(g(a))\}. \end{aligned}$$

and

$$\text{Eq}(h \circ f, k \circ g) \cong \{a \in A \mid h(f(a)) = k(g(a))\},$$

and thus there's an inclusion from $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$ to $\text{Eq}(h \circ f, k \circ g)$. \square

01E3 4.1.6 Inverse Limits

Let $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I} : (I, \preceq) \rightarrow \text{Sets}$ be an inverse system of sets.

01E4 Definition 4.1.6.1.1. The **inverse limit** of $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ is the inverse limit of $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ in Sets as in Limits and Colimits, ??.

01E5 Construction 4.1.6.1.2. Concretely, the inverse limit of $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ is the pair $(\varprojlim_{\alpha \in I} X_\alpha, \{\text{pr}_\alpha\}_{\alpha \in I})$ consisting of:

01E6 1. *The Limit.* The set $\varprojlim_{\alpha \in I} X_\alpha$ defined by

$$\varprojlim_{\alpha \in I} X_\alpha \stackrel{\text{def}}{=} \left\{ (x_\alpha)_{\alpha \in I} \in \prod_{\alpha \in I} X_\alpha \mid \begin{array}{l} \text{for each } \alpha, \beta \in I, \text{ if } \alpha \preceq \beta, \\ \text{then we have } x_\alpha = f_{\alpha\beta}(x_\beta) \end{array} \right\}.$$

01E7 2. *The Cone.* The collection

$$\left\{ \text{pr}_\gamma : \varprojlim_{\alpha \in I} X_\alpha \rightarrow X_\gamma \right\}_{\gamma \in I}$$

of maps of sets defined as the restriction of the maps

$$\left\{ \text{pr}_\gamma : \prod_{\alpha \in I} X_\alpha \rightarrow X_\gamma \right\}_{\gamma \in I}$$

of **Item 2** of **Definition 4.1.2.1.2** to $\lim_{\longleftarrow \alpha \in I} (X_\alpha)$ and hence given by

$$\text{pr}_\gamma((x_\alpha)_{\alpha \in I}) \stackrel{\text{def}}{=} x_\gamma$$

for each $\gamma \in I$ and each $(x_\alpha)_{\alpha \in I} \in \lim_{\longleftarrow \alpha \in I} (X_\alpha)$.

Proof. We claim that $\lim_{\longleftarrow \alpha \in I} (X_\alpha)$ is the limit of the inverse system of sets $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$.

First we need to check that the limit diagram defined by it commutes, i.e. that we have

$$\begin{array}{ccc} & \lim_{\longleftarrow \alpha \in I} (X_\alpha) & \\ & \swarrow \text{pr}_\alpha & \searrow \text{pr}_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$


$f_{\alpha\beta} \circ \text{pr}_\alpha = \text{pr}_\beta$

for each $\alpha, \beta \in I$ with $\alpha \preceq \beta$. Indeed, given $(x_\gamma)_{\gamma \in I} \in \lim_{\longleftarrow \gamma \in I} (X_\gamma)$, we have


$$\begin{aligned} [f_{\alpha\beta} \circ \text{pr}_\alpha]((x_\gamma)_{\gamma \in I}) &\stackrel{\text{def}}{=} f_{\alpha\beta}(\text{pr}_\alpha((x_\gamma)_{\gamma \in I})) \\ &\stackrel{\text{def}}{=} f_{\alpha\beta}(x_\alpha) \\ &= x_\beta \\ &\stackrel{\text{def}}{=} \text{pr}_\beta((x_\gamma)_{\gamma \in I}), \end{aligned}$$

where the third equality comes from the definition of $\lim_{\longleftarrow \alpha \in I} (X_\alpha)$. Next, we prove that $\lim_{\longleftarrow \alpha \in I} (X_\alpha)$ satisfies the universal property of an inverse limit. Suppose that

we have, for each $\alpha, \beta \in I$ with $\alpha \preceq \beta$, a diagram of the form



in Sets. Then there indeed exists a unique map $\phi: L \xrightarrow{\exists!} \varprojlim_{\alpha \in I} (X_\alpha)$ making the diagram



commute, being uniquely determined by the family of conditions

$$\{p_\alpha = \text{pr}_\alpha \circ \phi\}_{\alpha \in I}$$

via

$$\phi(\ell) = (p_\alpha(\ell))_{\alpha \in I}$$

for each $\ell \in L$, where we note that $(p_\alpha(\ell))_{\alpha \in I} \in \prod_{\alpha \in I} X_\alpha$ indeed lies in $\varprojlim_{\alpha \in I} (X_\alpha)$, as we have

$$\begin{aligned} f_{\alpha\beta}(p_\alpha(\ell)) &\stackrel{\text{def}}{=} [f_{\alpha\beta} \circ p_\alpha](\ell) \\ &\stackrel{\text{def}}{=} p_\beta(\ell) \end{aligned}$$

for each $\beta \in I$ with $\alpha \preceq \beta$ by the commutativity of the diagram for $(L, \{p_\alpha\}_{\alpha \in I})$. \square

01E8 Example 4.1.6.1.3. Here are some examples of inverse limits of sets.

01E9 1. *The p -Adic Integers.* The ring of p -adic integers \mathbb{Z}_p of ?? is the inverse limit

$$\mathbb{Z}_p \cong \varprojlim_{n \in \mathbb{N}} (\mathbb{Z}/p^n);$$

see ??.

01EA 2. *Rings of Formal Power Series.* The ring $R[[t]]$ of formal power series in a variable t is the inverse limit

$$R[[t]] \cong \varprojlim_{n \in \mathbb{N}} (R[t]/t^n R[t]);$$

see ??.

01EB 3. *Profinite Groups.* Profinite groups are inverse limits of finite groups; see ??.

001T 4.2 Colimits of Sets

001U 4.2.1 The Initial Set

001V Definition 4.2.1.1.1. The **initial set** is the initial object of **Sets** as in **Limits and Colimits**, ??.

01EC Construction 4.2.1.1.2. Concretely, the initial set is the pair $(\emptyset, \{\iota_A\}_{A \in \text{Obj}(\text{Sets})})$ consisting of:

01ED 1. *The Colimit.* The empty set \emptyset of **Definition 4.3.1.1.1.**

01EE 2. *The Cocone.* The collection of maps

$$\{\iota_A: \emptyset \rightarrow A\}_{A \in \text{Obj}(\text{Sets})}$$

given by the inclusion maps from \emptyset to A .

Proof. We claim that \emptyset is the initial object of **Sets**. Indeed, suppose we have a

diagram of the form

$$\emptyset \quad A$$

in **Sets**. Then there exists a unique map $\phi: \emptyset \rightarrow A$ making the diagram

$$\emptyset \xrightarrow[\exists!]{\phi} A$$

commute, namely the inclusion map ι_A . □

001W 4.2.2 Coproducts of Families of Sets

Let $\{A_i\}_{i \in I}$ be a family of sets.

001X **Definition 4.2.2.1.1.** The **coproduct** of $\{A_i\}_{i \in I}$ ⁶ is the coproduct of $\{A_i\}_{i \in I}$ in **Sets** as in Limits and Colimits, ??.

01EF **Construction 4.2.2.1.2.** Concretely, the disjoint union of $\{A_i\}_{i \in I}$ is the pair $\left(\coprod_{i \in I} A_i, \{\text{inj}_i\}_{i \in I} \right)$ consisting of:

01EG 1. *The Colimit.* The set $\coprod_{i \in I} A_i$ defined by

$$\coprod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ (i, x) \in I \times \left(\bigcup_{i \in I} A_i \right) \mid x \in A_i \right\}.$$

01EH 2. *The Cocone.* The collection

$$\left\{ \text{inj}_i: A_i \rightarrow \coprod_{i \in I} A_i \right\}_{i \in I}$$

of maps given by

$$\text{inj}_i(x) \stackrel{\text{def}}{=} (i, x)$$

for each $x \in A_i$ and each $i \in I$.

Proof. We claim that $\coprod_{i \in I} A_i$ is the categorical coproduct of $\{A_i\}_{i \in I}$ in **Sets**.

⁶*Further Terminology:* Also called the **disjoint union of the family** $\{A_i\}_{i \in I}$.

Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} & & C \\ & \nearrow \iota_i & \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

in **Sets**. Then there exists a unique map $\phi: \coprod_{i \in I} A_i \rightarrow C$ making the diagram

$$\begin{array}{ccc} & & C \\ & \nearrow \iota_i & \uparrow \phi \exists! \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

commute, being uniquely determined by the condition $\phi \circ \text{inj}_i = \iota_i$ for each $i \in I$ via

$$\phi((i, x)) = \iota_i(x)$$

for each $(i, x) \in \coprod_{i \in I} A_i$. □

001Y Proposition 4.2.2.1.3. Let $\{A_i\}_{i \in I}$ be a family of sets.

001Z I. *Functoriality.* The assignment $\{A_i\}_{i \in I} \mapsto \coprod_{i \in I} A_i$ defines a functor

$$\coprod_{i \in I}: \text{Fun}(I_{\text{disc}}, \mathbf{Sets}) \rightarrow \mathbf{Sets}$$

where

- *Action on Objects.* For each $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \mathbf{Sets}))$, we have

$$\left[\coprod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \coprod_{i \in I} A_i$$

- *Action on Morphisms.* For each $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \mathbf{Sets}))$, the action on Hom-sets

$$\left(\coprod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \mathbf{Sets} \left(\coprod_{i \in I} A_i, \coprod_{i \in I} B_i \right)$$

of $\coprod_{i \in I}$ at $((A_i)_{i \in I}, (B_i)_{i \in I})$ is defined by sending a map

$$\{f_i: A_i \rightarrow B_i\}_{i \in I}$$

in $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$ to the map of sets

$$\coprod_{i \in I} f_i: \coprod_{i \in I} A_i \rightarrow \coprod_{i \in I} B_i$$

defined by

$$\left[\coprod_{i \in I} f_i \right] (i, a) \stackrel{\text{def}}{=} f_i(a)$$

for each $(i, a) \in \coprod_{i \in I} A_i$.

Proof. **Item 1, Functoriality:** This follows from Limits and Colimits, ?? of ??. \square

0020 4.2.3 Binary Coproducts

Let A and B be sets.

0021 Definition 4.2.3.1.1. The **coproduct of A and B** ⁷ is the coproduct of A and B in **Sets** as in Limits and Colimits, ??.

01EJ Construction 4.2.3.1.2. Concretely, the coproduct of A and B is the pair $(A \coprod B, \{\text{inj}_1, \text{inj}_2\})$ consisting of:

01EK 1. *The Colimit.* The set $A \coprod B$ defined by

$$\begin{aligned} A \coprod B &\stackrel{\text{def}}{=} \coprod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{(0, a) \in S \mid a \in A\} \cup \{(1, b) \in S \mid b \in B\}, \end{aligned}$$

where $S = \{0, 1\} \times (A \cup B)$.

01EL 2. *The Cocone.* The maps

$$\begin{aligned} \text{inj}_1: A &\rightarrow A \coprod B, \\ \text{inj}_2: B &\rightarrow A \coprod B, \end{aligned}$$

⁷ *Further Terminology:* Also called the **disjoint union of A and B** .

given by

$$\begin{aligned}\text{inj}_1(a) &\stackrel{\text{def}}{=} (0, a), \\ \text{inj}_2(b) &\stackrel{\text{def}}{=} (1, b),\end{aligned}$$

for each $a \in A$ and each $b \in B$.

Proof. We claim that $A \amalg B$ is the categorical coproduct of A and B in **Sets**. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} & C & \\ \iota_1 \nearrow & & \nwarrow \iota_2 \\ A & \xrightarrow{\text{inj}_1} A \amalg B \xleftarrow{\text{inj}_2} & B \end{array}$$

in **Sets**. Then there exists a unique map $\phi: A \amalg B \rightarrow C$ making the diagram

$$\begin{array}{ccc} & C & \\ \iota_1 \nearrow & \uparrow \phi \mid \exists! & \nwarrow \iota_2 \\ A & \xrightarrow{\text{inj}_1} A \amalg B \xleftarrow{\text{inj}_2} & B \end{array}$$

commute, being uniquely determined by the conditions

$$\begin{aligned}\phi \circ \text{inj}_A &= \iota_A, \\ \phi \circ \text{inj}_B &= \iota_B\end{aligned}$$

via

$$\phi(x) = \begin{cases} \iota_A(a) & \text{if } x = (0, a), \\ \iota_B(b) & \text{if } x = (1, b) \end{cases}$$

for each $x \in A \amalg B$. □

0022 Proposition 4.2.3.1.3. Let A, B, C , and X be sets.

0023 I. *Functoriality.* The assignment $A, B, (A, B) \mapsto A \amalg B$ defines functors

$$\begin{aligned}A \amalg -: \quad \text{Sets} &\rightarrow \text{Sets}, \\ - \amalg B: \quad \text{Sets} &\rightarrow \text{Sets}, \\ -_1 \amalg -_2: \text{Sets} \times \text{Sets} &\rightarrow \text{Sets},\end{aligned}$$

where $-_1 \amalg -_2$ is the functor where

- *Action on Objects.* For each $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$, we have

$$[-_1 \amalg -_2](A, B) \stackrel{\text{def}}{=} A \amalg B.$$

- *Action on Morphisms.* For each $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\amalg_{(A,B),(X,Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \amalg B, X \amalg Y)$$

of \amalg at $((A, B), (X, Y))$ is defined by sending (f, g) to the function

$$f \amalg g : A \amalg B \rightarrow X \amalg Y$$

defined by

$$[f \amalg g](x) \stackrel{\text{def}}{=} \begin{cases} (0, f(a)) & \text{if } x = (0, a), \\ (1, g(b)) & \text{if } x = (1, b), \end{cases}$$

for each $x \in A \amalg B$.

and where $A \amalg -$ and $- \amalg B$ are the partial functors of $-_1 \amalg -_2$ at $A, B \in \text{Obj}(\text{Sets})$.

- 01ZH 2. *Adjointness.* We have an adjunction

$$(-_1 \amalg -_2 \dashv \Delta_{\text{Sets}}) : \text{Sets} \times \text{Sets} \begin{array}{c} \xrightarrow{-_1 \amalg -_2} \\ \perp \\ \xleftarrow{\Delta_{\text{Sets}}} \end{array} \text{Sets},$$

witnessed by a bijection

$$\text{Sets}(A \amalg B, C) \cong \text{Hom}_{\text{Sets} \times \text{Sets}}((A, B), (C, C))$$

natural in $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$ and in $C \in \text{Obj}(\text{Sets})$.

- 0024 3. *Associativity.* We have an isomorphism of sets

$$\alpha_{X,Y,Z}^{\text{Sets}, \amalg} : (X \amalg Y) \amalg Z \xrightarrow{\sim} X \amalg (Y \amalg Z),$$

natural in $X, Y, Z \in \text{Obj}(\text{Sets})$.

0025 4. *Unitality*. We have isomorphisms of sets

$$\begin{aligned}\lambda_X^{\text{Sets}, \coprod} : \emptyset \coprod X &\xrightarrow{\sim} X, \\ \rho_X^{\text{Sets}, \coprod} : X \coprod \emptyset &\xrightarrow{\sim} X,\end{aligned}$$

natural in $X \in \text{Obj}(\text{Sets})$.

0026 5. *Commutativity*. We have an isomorphism of sets

$$\sigma_{X,Y}^{\text{Sets}, \coprod} : X \coprod Y \xrightarrow{\sim} Y \coprod X,$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

0027 6. *Symmetric Monoidality*. The 7-tuple $(\text{Sets}, \coprod, \emptyset, \alpha_{\coprod}^{\text{Sets}}, \lambda_{\coprod}^{\text{Sets}}, \rho_{\coprod}^{\text{Sets}}, \sigma^{\text{Sets}})$ is a symmetric monoidal category.

Proof. **Item 1, Functoriality**: This follows from Limits and Colimits, ?? of ??.

Item 2, Adjointness: This follows from the universal property of the coproduct.

Item 3, Associativity: This is proved in the proof of **Monoidal Structures on the Category of Sets**, Definition 5.2.3.1.1.

Item 4, Unitality: This is proved in the proof of **Monoidal Structures on the Category of Sets**, Definitions 5.2.4.1.1 and 5.2.5.1.1.

Item 5, Commutativity: This is proved in the proof of **Monoidal Structures on the Category of Sets**, Definition 5.2.6.1.1.

Item 6, Symmetric Monoidality: This is a repetition of **Monoidal Structures on the Category of Sets**, Definition 5.2.7.1.1, and is proved there. \square

0028 4.2.4 Pushouts

Let A, B , and C be sets and let $f : C \rightarrow A$ and $g : C \rightarrow B$ be functions.

0029 **Definition 4.2.4.1.1.** The **pushout of A and B over C along f and g** ⁸ is the pushout of A and B over C along f and g in Sets as in Limits and Colimits, ??.

01EM **Construction 4.2.4.1.2.** Concretely, the pushout of A and B over C along f and g is the pair $(A \coprod_C B, \{\text{inj}_1, \text{inj}_2\})$ consisting of:

⁸*Further Terminology*: Also called the **fibre coproduct of A and B over C along f and g** .

01EN 1. *The Colimit.* The set $A \amalg_C B$ defined by

$$A \amalg_C B \stackrel{\text{def}}{=} A \amalg B / \sim_C,$$

where \sim_C is the equivalence relation on $A \amalg B$ generated by $(0, f(c)) \sim_C (1, g(c))$.

01EP 2. *The Cocone.* The maps

$$\text{inj}_1: A \rightarrow A \amalg_C B,$$

$$\text{inj}_2: B \rightarrow A \amalg_C B$$

given by

$$\text{inj}_1(a) \stackrel{\text{def}}{=} [(0, a)]$$

$$\text{inj}_2(b) \stackrel{\text{def}}{=} [(1, b)]$$

for each $a \in A$ and each $b \in B$.

Proof. We claim that $A \amalg_C B$ is the categorical pushout of A and B over C with respect to (f, g) in Sets. First we need to check that the relevant pushout diagram commutes, i.e. that we have


$$\text{inj}_1 \circ f = \text{inj}_2 \circ g,$$

$$\begin{array}{ccc} A \amalg_C B & \xleftarrow{\text{inj}_2} & B \\ \text{inj}_1 \uparrow & & \uparrow g \\ A & \xleftarrow{f} & C. \end{array}$$


Indeed, given $c \in C$, we have

$$\begin{aligned} [\text{inj}_1 \circ f](c) &= \text{inj}_1(f(c)) \\ &= [(0, f(c))] \\ &= [(1, g(c))] \\ &= \text{inj}_2(g(c)) \\ &= [\text{inj}_2 \circ g](c), \end{aligned}$$

where $[(0, f(c))] = [(1, g(c))]$ by the definition of the relation \sim on $A \amalg B$. Next, we prove that $A \amalg_C B$ satisfies the universal property of the pushout. Suppose we have a diagram of the form



in Sets. Then there exists a unique map $\phi: A \amalg_C B \rightarrow P$ making the diagram



commute, being uniquely determined by the conditions

$$\begin{aligned}\phi \circ \text{inj}_1 &= \iota_1, \\ \phi \circ \text{inj}_2 &= \iota_2\end{aligned}$$

via

$$\phi(x) = \begin{cases} \iota_1(a) & \text{if } x = [(0, a)], \\ \iota_2(b) & \text{if } x = [(1, b)] \end{cases}$$

for each $x \in A \amalg_C B$, where the well-definedness of ϕ is guaranteed by the equality $\iota_1 \circ f = \iota_2 \circ g$ and the definition of the relation \sim on $A \amalg B$ as follows:

- 01EQ** 1. *Case 1:* Suppose we have $x = [(0, a)] = [(0, a')]$ for some $a, a' \in A$. Then, by **Definition 4.2.4.1.3**, we have a sequence

$$(0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (0, a').$$

- 01ER** 2. *Case 2:* Suppose we have $x = [(1, b)] = [(1, b')]$ for some $b, b' \in B$. Then, by **Definition 4.2.4.1.3**, we have a sequence

$$(1, b) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b').$$

- 01ES** 3. *Case 3:* Suppose we have $x = [(0, a)] = [(1, b)]$ for some $a \in A$ and $b \in B$. Then, by **Definition 4.2.4.1.3**, we have a sequence

$$(0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b).$$

In all these cases, we declare $x \sim' y$ iff there exists some $c \in C$ such that $x = (0, f(c))$ and $y = (1, g(c))$ or $x = (1, g(c))$ and $y = (0, f(c))$. Then, the equality $\iota_1 \circ f = \iota_2 \circ g$ gives

$$\begin{aligned} \phi([x]) &= \phi([(0, f(c))]) \\ &\stackrel{\text{def}}{=} \iota_1(f(c)) \\ &= \iota_2(g(c)) \\ &\stackrel{\text{def}}{=} \phi([(1, g(c))]) \\ &= \phi([y]), \end{aligned}$$

with the case where $x = (1, g(c))$ and $y = (0, f(c))$ similarly giving $\phi([x]) = \phi([y])$. Thus, if $x \sim' y$, then $\phi([x]) = \phi([y])$. Applying this equality pairwise to the sequences

$$\begin{aligned} (0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (0, a'), \\ (1, b) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b'), \\ (0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b) \end{aligned}$$

gives

$$\begin{aligned} \phi([(0, a)]) &= \phi([(0, a')]), \\ \phi([(1, b)]) &= \phi([(1, b')]), \\ \phi([(0, a)]) &= \phi([(1, b)]), \end{aligned}$$

showing ϕ to be well-defined. \square

- 002A Remark 4.2.4.1.3.** In detail, by **Conditions on Relations**, **Definition 10.5.2.1.2**, the relation \sim of **Definition 4.2.4.1.1** is given by declaring $a \sim b$ iff one of the following conditions is satisfied:

- 01ET 1. We have $a, b \in A$ and $a = b$.
- 01EU 2. We have $a, b \in B$ and $a = b$.
- 01EV 3. There exist $x_1, \dots, x_n \in A \amalg B$ such that $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$, where we declare $x \sim' y$ if one of the following conditions is satisfied:
- 01EW (a) There exists $c \in C$ such that $x = (0, f(c))$ and $y = (1, g(c))$.
- 01EX (b) There exists $c \in C$ such that $x = (1, g(c))$ and $y = (0, f(c))$.

In other words, there exist $x_1, \dots, x_n \in A \amalg B$ satisfying the following conditions:

- 01EY (c) There exists $c_0 \in C$ satisfying one of the following conditions:
- 01ZJ i. We have $a = f(c_0)$ and $x_1 = g(c_0)$.
- 01ZK ii. We have $a = g(c_0)$ and $x_1 = f(c_0)$.
- 01EZ (d) For each $1 \leq i \leq n - 1$, there exists $c_i \in C$ satisfying one of the following conditions:
- 01ZL i. We have $x_i = f(c_i)$ and $x_{i+1} = g(c_i)$.
- 01ZM ii. We have $x_i = g(c_i)$ and $x_{i+1} = f(c_i)$.
- 01F0 (e) There exists $c_n \in C$ satisfying one of the following conditions:
- 01F1 i. We have $x_n = f(c_n)$ and $b = g(c_n)$.
- 01F2 ii. We have $x_n = g(c_n)$ and $b = f(c_n)$.

01ZN **Remark 4.2.4.1.4.** It is common practice to write $A \amalg_C B$ for the pushout of A and B over C along f and g , omitting the maps f and g from the notation and instead leaving them implicit, to be understood from the context.

However, the set $A \amalg_C B$ depends very much on the maps f and g , and sometimes it is necessary or useful to note this dependence explicitly. In such situations, we will write $A \amalg_{f,C,g} B$ or $A \amalg_C^{f,g} B$ for $A \amalg_C B$.

002B **Example 4.2.4.1.5.** Here are some examples of pushouts of sets.

- 002C 1. *Wedge Sums of Pointed Sets.* The wedge sum of two pointed sets of **Pointed Sets, Definition 6.3.3.1.1** is an example of a pushout of sets.

002D 2. *Intersections via Unions.* Let X be a set. We have

$$A \cup B \cong A \coprod_{A \cap B} B,$$

$$\begin{array}{ccc} A \cup B & \longleftarrow & B \\ \uparrow \lrcorner & & \uparrow \\ A & \longleftarrow & A \cap B \end{array}$$

for each $A, B \in \mathcal{P}(X)$.

Proof. Item 1, Wedge Sums of Pointed Sets: This follows by definition, as the wedge sum of two pointed sets is defined as a pushout.


Item 2, Intersections via Unions: Indeed, $A \coprod_{A \cap B} B$ is the quotient of $A \coprod B$ by the equivalence relation obtained by declaring $(0, a) \sim (1, b)$ iff $a = b \in A \cap B$, which is in bijection with $A \cup B$ via the map with $[(0, a)] \mapsto a$ and $[(1, b)] \mapsto b$. \square

002E **Proposition 4.2.4.1.6.** Let A, B, C , and X be sets.

002F 1. *Functoriality.* The assignment $(A, B, C, f, g) \mapsto A \coprod_{f, C, g} B$ defines a functor

$$-_1 \coprod_{-3} -_1: \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where \mathcal{P} is the category that looks like this:




In particular, the action on morphisms of $-_1 \coprod_{-3} -_1$ is given by sending a morphism

$$\begin{array}{ccccc} A \coprod_C B & \longleftarrow & B & & \\ \uparrow \lrcorner & & \uparrow & \searrow \psi & \\ A' \coprod_{C'} B' & \longleftarrow & B' & & \\ \uparrow \lrcorner & & \downarrow g & & \uparrow g' \\ A & \xleftarrow{f} & C & \searrow \chi & \\ \downarrow \phi & & & & \\ A' & \xleftarrow{f'} & C' & & \end{array}$$

in $\text{Fun}(\mathcal{P}, \text{Sets})$ to the map $\xi: A \amalg_C B \xrightarrow{\exists!} A' \amalg_{C'} B'$ given by

$$\xi(x) \stackrel{\text{def}}{=} \begin{cases} \phi(a) & \text{if } x = [(0, a)], \\ \psi(b) & \text{if } x = [(1, b)] \end{cases}$$

for each $x \in A \amalg_C B$, which is the unique map making the diagram



commute.

01ZP 2. *Adjointness.* We have an adjunction


$$\left(-1 \amalg_{X^{-2}} \dashv \Delta_{\text{Sets}_{X/}} \right): \text{Sets}_{X/} \times \text{Sets}_{X/} \xrightleftharpoons[\Delta_{\text{Sets}_{X/}}]{-1 \amalg_{X^{-2}} \dashv} \text{Sets}_{X/},$$

witnessed by a bijection

$$\text{Sets}_{X/}(A \amalg_X B, C), \cong \text{Hom}_{\text{Sets}_{X/} \times \text{Sets}_{X/}}((A, B), (C, C))$$

natural in $(A, B) \in \text{Obj}(\text{Sets}_{X/} \times \text{Sets}_{X/})$ and in $C \in \text{Obj}(\text{Sets}_{X/})$.


002G 3. *Associativity.* Given a diagram



in **Sets**, we have isomorphisms of sets


$$(A \amalg_X B) \amalg_Y C \cong (A \amalg_X B) \amalg_B (B \amalg_Y C) \cong A \amalg_X (B \amalg_Y C)$$

where these pullbacks are built as in the diagrams



01F4

4. *Interaction With Composition.* Given a diagram




in **Sets**, we have isomorphisms of sets

$$\begin{aligned} X \amalg_K^{\phi \circ f, \psi \circ g} Y &\cong \left(X \amalg_A^{\phi, j_1} (A \amalg_K^{f, g} B) \right) \amalg_{A \amalg_K^{f, g} B}^{i_2, i_1} \left((A \amalg_K^{f, g} B) \amalg_B^{j_2, \psi} Y \right) \\ &\cong X \amalg_A^{\phi, i} \left((A \amalg_K^{f, g} B) \amalg_B^{j_2, \psi} Y \right) \\ &\cong \left(X \amalg_A^{\phi, i_1} (A \amalg_K^{f, g} B) \right) \amalg_B^{j, \psi} Y \end{aligned}$$

where

$$\begin{aligned} j_1 &= \text{inj}_1^{A \times_K^{f, g} B}, & j_2 &= \text{inj}_2^{A \times_K^{f, g} B}, \\ i_1 &= \text{inj}_1^{(A \times_K^{f, g} B) \times_Y^{q_2, \psi}}, & i_2 &= \text{inj}_2^{X \times_{A \times_K^{f, g} B}^{\phi, q_1} (A \times_K^{f, g} B)}, \\ i &= j_1 \circ \text{inj}_1^{(A \times_K^{f, g} B) \times_B^{q_2, \psi} Y}, & j &= j_2 \circ \text{inj}_2^{X \times_A^{\phi, q_1} (A \times_K^{f, g} B)}, \end{aligned}$$

and where these pullbacks are built as in the diagrams



002H 5. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc}
 A & \xlongequal{\quad} & A \\
 f \uparrow \lrcorner & & \uparrow f \\
 X & \xlongequal{\quad} & X
 \end{array}
 \quad
 \begin{array}{l}
 \lambda_A^{\mathbf{Sets}_{X/}} : X \amalg_X A \xrightarrow{\sim} A, \\
 \rho_A^{\mathbf{Sets}_{X/}} : A \amalg_X X \xrightarrow{\sim} A,
 \end{array}
 \quad
 \begin{array}{ccc}
 A & \xleftarrow{f} & X \\
 \parallel \lrcorner & & \parallel \\
 X & \xleftarrow{f} & X,
 \end{array}$$

natural in $(A, f) \in \mathbf{Obj}(\mathbf{Sets}_{X/})$.

002J 6. *Commutativity*. We have an isomorphism of sets

$$\begin{array}{ccc} A \amalg_X B & \longleftarrow & B \\ \uparrow \ulcorner & & \uparrow g \\ A & \xleftarrow{f} & X, \end{array} \quad \sigma_A^{\text{Sets}_{X/}} : A \amalg_X B \xrightarrow{\sim} B \amalg_X A \quad \begin{array}{ccc} B \amalg_X A & \longleftarrow & A \\ \uparrow \ulcorner & & \uparrow f \\ B & \xleftarrow{g} & X. \end{array}$$

natural in $(A, f), (B, g) \in \text{Obj}(\text{Sets}_{X/})$.

002K 7. *Interaction With Coproducts*. We have

$$A \amalg_{\emptyset} B \cong A \amalg B, \quad \begin{array}{ccc} A \amalg B & \longleftarrow & B \\ \uparrow \ulcorner & & \uparrow \iota_B \\ A & \xleftarrow{\iota_A} & \emptyset. \end{array}$$

002L 8. *Symmetric Monoidality*. The triple $(\text{Sets}_{X/}, \amalg_X, X)$ is a symmetric monoidal category.

Proof. **Item 1, Functoriality:** This is a special case of functoriality of co/limits, Limits and Colimits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pushout diagram.

Item 2, : Adjointness: This follows from the universal property of the coproduct (pushouts are coproducts in $\text{Sets}_{X/}$).

Item 3, Associativity: Omitted.

Item 4, Interaction With Composition: Omitted.

Item 5, Unitality: Omitted.

Item 6, Commutativity: Omitted.

Item 7, Interaction With Coproducts: Omitted.

Item 8, Symmetric Monoidality: Omitted. \square

002M 4.2.5 Coequalisers

Let A and B be sets and let $f, g: A \rightrightarrows B$ be functions.

002N **Definition 4.2.5.1.1.** The **coequaliser of f and g** is the coequaliser of f and g in Sets as in Limits and Colimits, ??.

01F5 Construction 4.2.5.1.2. Concretely, the coequaliser of f and g is the pair $(\text{CoEq}(f, g), \text{coeq}(f, g))$ consisting of:

01F6 1. *The Colimit.* The set $\text{CoEq}(f, g)$ defined by

$$\text{CoEq}(f, g) \stackrel{\text{def}}{=} B/\sim,$$

where \sim is the equivalence relation on B generated by $f(a) \sim g(a)$.

01F7 2. *The Cocone.* The map

$$\text{coeq}(f, g) : B \twoheadrightarrow \text{CoEq}(f, g)$$

given by the quotient map $\pi : B \twoheadrightarrow B/\sim$ with respect to the equivalence relation generated by $f(a) \sim g(a)$.

Proof. We claim that $\text{CoEq}(f, g)$ is the categorical coequaliser of f and g in **Sets**. First we need to check that the relevant coequaliser diagram commutes, i.e. that we have

$$\text{coeq}(f, g) \circ f = \text{coeq}(f, g) \circ g.$$

Indeed, we have

$$\begin{aligned} [\text{coeq}(f, g) \circ f](a) &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](f(a)) \\ &\stackrel{\text{def}}{=} [f(a)] \\ &= [g(a)] \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](g(a)) \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g) \circ g](a) \end{aligned}$$

for each $a \in A$. Next, we prove that $\text{CoEq}(f, g)$ satisfies the universal property of the coequaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} A & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & B \\ & \searrow c & \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g) \\ & & C \end{array}$$

in **Sets**. Then, since $c(f(a)) = c(g(a))$ for each $a \in A$, it follows from **Conditions on Relations, Items 4 and 5** of **Definition 10.6.2.1.3** that there exists a unique

map $\text{CoEq}(f, g) \xrightarrow{\exists!} C$ making the diagram

$$\begin{array}{ccc} A & \xrightleftharpoons[g]{f} B & \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g) \\ & \searrow c & \downarrow \exists! \\ & & C \end{array}$$

commute. □

002P Remark 4.2.5.1.3. In detail, by [Conditions on Relations](#), [Definition 10.5.2.1.2](#), the relation \sim of [Definition 4.2.5.1.1](#) is given by declaring $a \sim b$ iff one of the following conditions is satisfied:

01ZQ 1. We have $a = b$;

01ZR 2. There exist $x_1, \dots, x_n \in B$ such that $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$, where we declare $x \sim' y$ if one of the following conditions is satisfied:

01ZS (a) There exists $z \in A$ such that $x = f(z)$ and $y = g(z)$.

01ZT (b) There exists $z \in A$ such that $x = g(z)$ and $y = f(z)$.

In other words, there exist $x_1, \dots, x_n \in B$ satisfying the following conditions:

01ZU (a) There exists $z_0 \in A$ satisfying one of the following conditions:

01ZV i. We have $a = f(z_0)$ and $x_1 = g(z_0)$.

01ZW ii. We have $a = g(z_0)$ and $x_1 = f(z_0)$.

01ZX (b) For each $1 \leq i \leq n - 1$, there exists $z_i \in A$ satisfying one of the following conditions:

01ZY i. We have $x_i = f(z_i)$ and $x_{i+1} = g(z_i)$.

01ZZ ii. We have $x_i = g(z_i)$ and $x_{i+1} = f(z_i)$.

0200 (c) There exists $z_n \in A$ satisfying one of the following conditions:

0201 i. We have $x_n = f(z_n)$ and $b = g(z_n)$.

0202 ii. We have $x_n = g(z_n)$ and $b = f(z_n)$.

002Q Example 4.2.5.1.4. Here are some examples of coequalisers of sets.

- 002R I. *Quotients by Equivalence Relations.* Let R be an equivalence relation on a set X . We have a bijection of sets

$$X/\sim_R \cong \text{CoEq}\left(R \rightarrow X \times X \begin{matrix} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} \end{matrix} X\right).$$

Proof. **Item 1, Quotients by Equivalence Relations:** See [Pro25z]. □

002S **Proposition 4.2.5.1.5.** Let A , B , and C be sets.

- 002T I. *Associativity.* We have isomorphisms of sets⁹

$$\underbrace{\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h)}_{=\text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ b)} \cong \text{CoEq}(f, g, h) \cong \underbrace{\text{CoEq}(\text{coeq}(g, b) \circ f, \text{coeq}(g, b) \circ g)}_{=\text{CoEq}(\text{coeq}(g, b) \circ f, \text{coeq}(g, b) \circ h)},$$

⁹That is, the following three ways of forming “the” coequaliser of (f, g, b) agree:

- 0203 I. Take the coequaliser of (f, g, b) , i.e. the colimit of the diagram

$$A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{b} \end{matrix} B$$

in Sets.

- 0204 2. First take the coequaliser of f and g , forming a diagram

$$A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \end{matrix} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g)$$

and then take the coequaliser of the composition

$$A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{b} \end{matrix} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h) = \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ b)$$

of $\text{CoEq}(f, g)$

- 0205 3. First take the coequaliser of g and b , forming a diagram

$$A \begin{matrix} \xrightarrow{g} \\ \xrightarrow{b} \end{matrix} B \xrightarrow{\text{coeq}(g, b)} \text{CoEq}(g, b)$$

where $\text{CoEq}(f, g, h)$ is the colimit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{\quad} \\ \xrightarrow{h} \end{array} B$$

in **Sets**.

002U 4. *Unitality*. We have an isomorphism of sets

$$\text{CoEq}(f, f) \cong B.$$

002V 5. *Commutativity*. We have an isomorphism of sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

002W 6. *Interaction With Composition*. Let

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{\quad} \end{array} B \begin{array}{c} \xrightarrow{h} \\ \xrightarrow[k]{\quad} \end{array} C$$

be functions. We have a surjection

$$\text{CoEq}(h \circ f, k \circ g) \twoheadrightarrow \text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$$

exhibiting $\text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$ as a quotient of $\text{CoEq}(h \circ f, k \circ g)$ by the relation generated by declaring $h(y) \sim k(y)$ for each $y \in B$.

Proof. **Item 1, Associativity**: Omitted.

Item 4, Unitality: Omitted.

Item 5, Commutativity: Omitted.

Item 6, Interaction With Composition: Omitted. □

and then take the coequaliser of the composition

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[g]{\quad} \end{array} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g) = \text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)$$

01F8 4.2.6 Direct Colimits

Let $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I} : (I, \preceq) \rightarrow \Pi$ be a direct system of sets.

01F9 **Definition 4.2.6.1.1.** The **direct colimit** of $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ is the direct colimit of $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ in Sets as in Limits and Colimits, ??.

01FA **Construction 4.2.6.1.2.** Concretely, the direct colimit of $(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}$ is the pair $(\varinjlim_{\alpha \in I} X_\alpha, \{\text{inj}_\alpha\}_{\alpha \in I})$ consisting of:

01FB 1. *The Colimit.* The set $\varinjlim_{\alpha \in I} X_\alpha$ defined by

$$\varinjlim_{\alpha \in I} X_\alpha \stackrel{\text{def}}{=} \left(\coprod_{\alpha \in I} X_\alpha \right) / \sim,$$

where \sim is the equivalence relation on $\coprod_{\alpha \in I} X_\alpha$ generated by declaring $(\alpha, x) \sim (\beta, y)$ iff there exists some $\gamma \in I$ satisfying the following conditions:

- 01FC (a) We have $\alpha \preceq \gamma$.
- 01FD (b) We have $\beta \preceq \gamma$.
- 01FE (c) We have $f_{\alpha\gamma}(x) = f_{\beta\gamma}(y)$.

01FF 2. *The Cocone.* The collection

$$\left\{ \text{inj}_\gamma : X_\gamma \rightarrow \varinjlim_{\alpha \in I} X_\alpha \right\}_{\gamma \in I}$$

of maps of sets defined by

$$\text{inj}_\gamma(x) \stackrel{\text{def}}{=} [(\gamma, x)]$$

for each $\gamma \in I$ and each $x \in X_\gamma$.

Proof. We will prove [Definition 4.2.6.1.2](#) below in a bit, but first we need a lemma (which is interesting in its own right). \square

01FG Lemma 4.2.6.1.3. For each $\alpha, \beta \in I$ and each $x \in X_\alpha$, if $\alpha \preceq \beta$, then we have

$$(\alpha, x) \sim (\beta, f_{\alpha\beta}(x))$$

in $\text{colim}_{\alpha \in I} (X_\alpha)$.

Proof. Taking $\gamma = \beta$, we have $f_{\alpha\gamma} = f_{\alpha\beta}$, we have $f_{\beta\gamma} = f_{\beta\beta} \stackrel{\text{def}}{=} \text{id}_{X_\beta}$, and we have

$$\begin{aligned} f_{\alpha\beta}(x) &= f_{\beta\beta}(f_{\alpha\beta}(x)) \\ &\stackrel{\text{def}}{=} \text{id}_{X_\beta}(f_{\alpha\beta}(x)), \\ &= f_{\alpha\beta}(x). \end{aligned}$$

As a result, since $\alpha \preceq \beta$ and $\beta \preceq \beta$ as well, [Items 1a to 1c](#) of [Definition 4.2.6.1.2](#) are met. Thus we have $(\alpha, x) \sim (\beta, f_{\alpha\beta}(x))$. \square

We can now prove [Definition 4.2.6.1.2](#):

Proof. We claim that $\text{colim}_{\alpha \in I} (X_\alpha)$ is the colimit of the direct system of sets

$$(X_\alpha, f_{\alpha\beta})_{\alpha, \beta \in I}.$$

Commutativity of the Colimit Diagram: First, we need to check that the colimit diagram defined by $\text{colim}_{\alpha \in I} (X_\alpha)$ commutes, i.e. that we have

$$\begin{array}{ccc} & \text{colim}(X_\alpha) & \\ & \xrightarrow{\alpha \in I} & \\ \text{inj}_\alpha = \text{inj}_\beta \circ f_{\alpha\beta} & \nearrow \text{inj}_\alpha & \nwarrow \text{inj}_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$


for each $\alpha, \beta \in I$ with $\alpha \preceq \beta$. Indeed, given $x \in X_\alpha$, we have

$$[\text{inj}_\beta \circ f_{\alpha\beta}](x) \stackrel{\text{def}}{=} \text{inj}_\beta(f_{\alpha\beta}(x))$$


$$\begin{aligned}
&\stackrel{\text{def}}{=} \left[\left(\beta, f_{\alpha\beta}(x) \right) \right] \\
&= [(\alpha, x)] \\
&\stackrel{\text{def}}{=} \text{inj}_\alpha(x),
\end{aligned}$$

where we have used [Definition 4.2.6.1.3](#) for the third equality.

Proof of the Universal Property of the Colimit: Next, we prove that $\text{colim}_{\alpha \in I} (X_\alpha)$ as constructed in [Definition 4.2.6.1.2](#) satisfies the universal property of a direct colimit. Suppose that we have, for each $\alpha, \beta \in I$ with $\alpha \preceq \beta$, a diagram of the form



in Sets. We claim that there exists a unique map $\phi: \text{colim}(X_\alpha) \xrightarrow[\alpha \in I]{\exists!} C$ making the diagram



of $\text{CoEq}(g, h)$.

commute. To this end, first consider the diagram

$$\begin{array}{ccc} \coprod_{\alpha \in I} X_{\alpha} & \xrightarrow{\text{pr}} & \text{colim}_{\alpha \in I} (X_{\alpha}) \\ & \searrow & \uparrow \\ & \coprod_{\alpha \in I} i_{\alpha} & \\ & & C. \end{array}$$

Lemma. If $(\alpha, x) \sim (\beta, y)$, then we have

$$\left[\coprod_{\alpha \in I} i_{\alpha} \right] (x) = \left[\coprod_{\alpha \in I} i_{\alpha} \right] (y).$$

Proof. Indeed, if $(\alpha, x) \sim (\beta, y)$, then there exists some $\gamma \in I$ satisfying the following conditions:

- 0206 1. We have $\alpha \preceq \gamma$.
- 0207 2. We have $\beta \preceq \gamma$.
- 0208 3. We have $f_{\alpha\gamma}(x) = f_{\beta\gamma}(y)$.

We then have

$$\begin{aligned} \left[\coprod_{\alpha \in I} i_{\alpha} \right] (x) &\stackrel{\text{def}}{=} i_{\alpha}(x) \\ &\stackrel{\text{def}}{=} [i_{\gamma} \circ f_{\alpha\gamma}](x) \\ &\stackrel{\text{def}}{=} i_{\gamma}(f_{\alpha\gamma}(x)) \\ &= i_{\gamma}(f_{\beta\gamma}(y)) \\ &\stackrel{\text{def}}{=} [i_{\gamma} \circ f_{\beta\gamma}](y) \\ &= i_{\beta}(y) \\ &\stackrel{\text{def}}{=} \left[\coprod_{\alpha \in I} i_{\alpha} \right] (y). \end{aligned}$$

This finishes the proof of the lemma. Continuing, by **Conditions on Relations**, ?? of **Definition 10.6.2.1.3**, there then exists a map $\phi: \text{colim}(X_\alpha) \xrightarrow{\exists!} C$ making the diagram

$$\begin{array}{ccc} \coprod_{\alpha \in I} X_\alpha & \xrightarrow{\text{pr}} & \text{colim}(X_\alpha) \\ & \searrow & \downarrow \phi \\ & \coprod_{\alpha \in I} i_\alpha & C \end{array}$$

commute. In particular, this implies that the diagram

$$\begin{array}{ccc} X_\alpha & \xrightarrow{\text{inj}_\alpha} & \text{colim}(X_\alpha) \\ & \searrow i_\alpha & \downarrow \phi \\ & & C \end{array}$$

also commutes, and thus so does the diagram

$$\begin{array}{ccc} & C & \\ & \uparrow \phi \mid \exists! & \\ & \text{colim}(X_\alpha) & \\ & \xrightarrow{\alpha \in I} & \\ i_\alpha \swarrow & & \searrow i_\beta \\ X_\alpha & \xrightarrow{f_{\alpha\beta}} & X_\beta \end{array}$$

This finishes the proof.¹⁰

□

¹⁰Incidentally, the conditions

$$\{i_\alpha = \phi \circ \text{inj}_\alpha\}_{\alpha \in I}$$

show that ϕ must be given by

$$\phi([\langle \alpha, x \rangle]) = (i_\alpha(x))_{\alpha \in I}$$

for each $[\langle \alpha, x \rangle] \in \text{colim}_{\alpha \in I} (X_\alpha)$, although we would need to show that this assignment is

01FH Example 4.2.6.I.4. Here are some examples of direct colimits of sets.

01FJ I. *The Prüfer Group.* The Prüfer group $\mathbb{Z}(p^\infty)$ is defined as the direct colimit

$$\mathbb{Z}(p^\infty) \stackrel{\text{def}}{=} \varinjlim_{n \in \mathbb{N}} (\mathbb{Z}/p^n);$$

see ??.

002X 4.3 Operations With Sets

002Y 4.3.I The Empty Set

002Z Definition 4.3.I.I.I. The **empty set** is the set \emptyset defined by

$$\emptyset \stackrel{\text{def}}{=} \{x \in X \mid x \neq x\},$$

where X is the set in the set existence axiom, ?? of ??.

0030 4.3.2 Singleton Sets

Let X be a set.

0031 Definition 4.3.2.I.I. The **singleton set containing** X is the set $\{X\}$ defined by

$$\{X\} \stackrel{\text{def}}{=} \{X, X\},$$

where $\{X, X\}$ is the pairing of X with itself of **Definition 4.3.3.I.I.**

0032 4.3.3 Pairings of Sets

Let X and Y be sets.

0033 Definition 4.3.3.I.I. The **pairing of X and Y** is the set $\{X, Y\}$ defined by

$$\{X, Y\} \stackrel{\text{def}}{=} \{x \in \mathcal{A} \mid x = X \text{ or } x = Y\},$$

where \mathcal{A} is the set in the axiom of pairing, ?? of ??.

0034 4.3.4 Ordered Pairs

Let A and B be sets.

0035 Definition 4.3.4.1.1. The **ordered pair associated to A and B** is the set (A, B) defined by

$$(A, B) \stackrel{\text{def}}{=} \{\{A\}, \{A, B\}\}.$$

0036 Proposition 4.3.4.1.2. Let A and B be sets.

0037 1. *Uniqueness.* Let A, B, C , and D be sets. The following conditions are equivalent:

0038 (a) We have $(A, B) = (C, D)$.

0039 (b) We have $A = C$ and $B = D$.

Proof. *Item 1, Uniqueness:* See [Cie97, Theorem 1.2.3]. □

003A 4.3.5 Sets of Maps

Let A and B be sets.

003B Definition 4.3.5.1.1. The **set of maps from A to B ¹¹** is the set $\text{Sets}(A, B)$ ¹² whose elements are the functions from A to B .

003C Proposition 4.3.5.1.2. Let A and B be sets.

003D 1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto \text{Hom}_{\text{Sets}}(X, Y)$ define functors

$$\begin{aligned} \text{Sets}(X, -) &: \text{Sets} && \rightarrow \text{Sets}, \\ \text{Sets}(-, Y) &: \text{Sets}^{\text{op}} && \rightarrow \text{Sets}, \\ \text{Sets}(-_1, -_2) &: \text{Sets}^{\text{op}} \times \text{Sets} && \rightarrow \text{Sets}. \end{aligned}$$

well-defined were we to prove **Definition 4.2.6.1.2** in this way. Instead, invoking **Conditions on Relations, ??** of **Definition 10.6.2.1.3** gave us a way to avoid having to prove this, leading to a cleaner alternative proof.

¹¹*Further Terminology:* Also called the **Hom set from A to B** .

¹²*Further Notation:* Also written $\text{Hom}_{\text{Sets}}(A, B)$.

01FK 2. *Adjointness.* We have adjunctions

$$(A \times - \dashv \text{Sets}(A, -)): \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Sets}(A, -)} \end{array} \text{Sets},$$

$$(- \times B \dashv \text{Sets}(B, -)): \text{Sets} \begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Sets}(B, -)} \end{array} \text{Sets},$$

witnessed by bijections

$$\begin{aligned} \text{Sets}(A \times B, C) &\cong \text{Sets}(A, \text{Sets}(B, C)), \\ \text{Sets}(A \times B, C) &\cong \text{Sets}(B, \text{Sets}(A, C)), \end{aligned}$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

01FL 3. *Maps From the Punctual Set.* We have a bijection

$$\text{Sets}(\text{pt}, A) \cong A,$$

natural in $A \in \text{Obj}(\text{Sets})$.

01FM 4. *Maps to the Punctual Set.* We have a bijection

$$\text{Sets}(A, \text{pt}) \cong \text{pt},$$

natural in $A \in \text{Obj}(\text{Sets})$.

Proof. **Item 1, Functoriality:** This follows from **Categories, Items 2 and 5** of **Definition II.1.4.1.2**.

Item 2, Adjointness: This is a repetition of **Item 2** of **Definition 4.1.3.1.3** and is proved there.

Item 3, Maps From the Punctual Set: The bijection

$$\Phi_A: \text{Sets}(\text{pt}, A) \xrightarrow{\sim} A$$

is given by

$$\Phi_A(f) \stackrel{\text{def}}{=} f(\star)$$

for each $f \in \text{Sets}(\text{pt}, A)$, admitting an inverse

$$\Phi_A^{-1}: A \xrightarrow{\sim} \text{Sets}(\text{pt}, A)$$

given by

$$\Phi_A^{-1}(a) \stackrel{\text{def}}{=} \llbracket \star \mapsto a \rrbracket$$

for each $a \in A$. Indeed, we have

$$\begin{aligned} [\Phi_A^{-1} \circ \Phi_A](f) &\stackrel{\text{def}}{=} \Phi_A^{-1}(\Phi_A(f)) \\ &\stackrel{\text{def}}{=} \Phi_A^{-1}(f(\star)) \\ &\stackrel{\text{def}}{=} \llbracket \star \mapsto f(\star) \rrbracket \\ &\stackrel{\text{def}}{=} f \\ &\stackrel{\text{def}}{=} [\text{id}_{\text{Sets}(\text{pt}, A)}](f) \end{aligned}$$

for each $f \in \text{Sets}(\text{pt}, A)$ and

$$\begin{aligned} [\Phi_A \circ \Phi_A^{-1}](a) &\stackrel{\text{def}}{=} \Phi_A(\Phi_A^{-1}(a)) \\ &\stackrel{\text{def}}{=} \Phi_A(\llbracket \star \mapsto a \rrbracket) \\ &\stackrel{\text{def}}{=} \text{ev}_\star(\llbracket \star \mapsto a \rrbracket) \\ &\stackrel{\text{def}}{=} a \\ &\stackrel{\text{def}}{=} [\text{id}_A](a) \end{aligned}$$

for each $a \in A$, and thus we have

$$\begin{aligned} \Phi_A^{-1} \circ \Phi_A &= \text{id}_{\text{Sets}(\text{pt}, A)} \\ \Phi_A \circ \Phi_A^{-1} &= \text{id}_A. \end{aligned}$$

To prove naturality, we need to show that the diagram

$$\begin{array}{ccc} \text{Sets}(\text{pt}, A) & \xrightarrow{f!} & \text{Sets}(\text{pt}, B) \\ \Phi_A \downarrow \wr & & \wr \downarrow \Phi_B \\ A & \xrightarrow{f} & B \end{array}$$

commutes. Indeed, we have

$$\begin{aligned} [f \circ \Phi_A](\phi) &\stackrel{\text{def}}{=} f(\Phi_A(\phi)) \\ &\stackrel{\text{def}}{=} f(\phi(\star)) \end{aligned}$$

$$\begin{aligned}
&\stackrel{\text{def}}{=} [f \circ \phi](\star) \\
&\stackrel{\text{def}}{=} \Phi_B(f \circ \phi) \\
&\stackrel{\text{def}}{=} \Phi_B(f_!(\phi)) \\
&\stackrel{\text{def}}{=} [\Phi_B \circ f_!](\phi)
\end{aligned}$$

for each $\phi \in \text{Sets}(\text{pt}, \mathcal{A})$. This finishes the proof.

Item 4, Maps to the Punctual Set: This follows from the universal property of pt as the terminal set, **Definition 4.1.1.1.1**. \square

003E 4.3.6 Unions of Families of Subsets

Let X be a set and let $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

003F Definition 4.3.6.1.1. The **union of** \mathcal{U} is the set $\bigcup_{U \in \mathcal{U}} U$ defined by

$$\bigcup_{U \in \mathcal{U}} U \stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } U \in \mathcal{U} \\ \text{such that we have } x \in U \end{array} \right. \right\}.$$

01FN Proposition 4.3.6.1.2. Let X be a set.

01FP 1. *Functoriality.* The assignment $\mathcal{U} \mapsto \bigcup_{U \in \mathcal{U}} U$ defines a functor

$$\bigcup: (\mathcal{P}(\mathcal{P}(X)), \subset) \rightarrow (\mathcal{P}(X), \subset).$$

In particular, for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$, the following condition is satisfied:

$$(\star) \text{ If } \mathcal{U} \subset \mathcal{V}, \text{ then } \bigcup_{U \in \mathcal{U}} U \subset \bigcup_{V \in \mathcal{V}} V.$$

01FQ 2. *Associativity.* The diagram

$$\begin{array}{ccc}
\mathcal{P}(\mathcal{P}(\mathcal{P}(X))) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \bigcup} & \mathcal{P}(\mathcal{P}(X)) \\
\bigcup \star \text{id}_{\mathcal{P}(X)} \downarrow & & \downarrow \bigcup \\
\mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcup} & \mathcal{P}(X)
\end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \bigcup_{A \in \mathcal{A}} A} U = \bigcup_{A \in \mathcal{A}} \left(\bigcup_{U \in A} U \right)$$

for each $\mathcal{A} \in \mathcal{P}(\mathcal{P}(\mathcal{P}(X)))$.

01FR 3. *Left Unitality*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \chi_{\mathcal{P}(X)} \downarrow & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{V \in \{U\}} V = U$$

for each $U \in \mathcal{P}(X)$.

01FS 4. *Right Unitality*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \mathcal{P}(\chi_X) \downarrow & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{\{u\} \in \chi_X(U)} \{u\} = U$$

for each $U \in \mathcal{P}(X)$.

01FT 5. *Interaction With Unions I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \times \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{W \in \mathcal{U} \cup \mathcal{V}} W = \left(\bigcup_{U \in \mathcal{U}} U \right) \cup \left(\bigcup_{V \in \mathcal{V}} V \right)$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01FU 6. *Interaction With Unions II.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cup -)} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{U \cup -} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cup V)} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{- \cup V} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$\begin{aligned} U \cup \left(\bigcup_{V \in \mathcal{V}} V \right) &= \bigcup_{V \in \mathcal{V}} (U \cup V), \\ \left(\bigcup_{U \in \mathcal{U}} U \right) \cup V &= \bigcup_{U \in \mathcal{U}} (U \cup V) \end{aligned}$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ and each $U, V \in \mathcal{P}(X)$.

01FV 7. *Interaction With Intersections I.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \times \cup \downarrow & \supset & \downarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X), \end{array}$$

with components

$$\bigcup_{W \in \mathcal{U} \cap \mathcal{V}} W \subset \left(\bigcup_{U \in \mathcal{U}} U \right) \cap \left(\bigcup_{V \in \mathcal{V}} V \right)$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01FW 8. *Interaction With Intersections II.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cap -)} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{U \cap -} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cap V)} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{- \cap V} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$\begin{aligned} U \cup \left(\bigcup_{V \in \mathcal{V}} V \right) &= \bigcup_{V \in \mathcal{V}} (U \cup V), \\ \left(\bigcup_{U \in \mathcal{U}} U \right) \cup V &= \bigcup_{U \in \mathcal{U}} (U \cup V) \end{aligned}$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ and each $U, V \in \mathcal{P}(X)$.

01FX 9. *Interaction With Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\setminus} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup \times \cup & \text{X} & \downarrow \cup \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\setminus} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in \mathcal{U} \setminus \mathcal{V}} W \neq \left(\bigcup_{U \in \mathcal{U}} U \right) \setminus \left(\bigcup_{V \in \mathcal{V}} V \right)$$

in general, where $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01FY IO. *Interaction With Complements I.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cup^{\text{op}} & \text{X} & \downarrow \cup \\
 \mathcal{P}(X)^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{U \in \mathcal{U}^c} U \neq \bigcup_{U \in \mathcal{U}} U^c$$

in general, where $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01FZ II. *Interaction With Complements II.* The diagram

$$\begin{array}{ccccc}
 & & \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \\
 & \nearrow \text{id}_{\mathcal{P}(X)} \star (-)^c & & \dashrightarrow \sim & \\
 \mathcal{P}(\mathcal{P}(X)) & & & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\
 \downarrow \cup & & & & \downarrow \cap^{\text{op}} \\
 \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}}, & &
 \end{array}$$

commutes, i.e. we have

$$\left(\bigcup_{U \in \mathcal{U}} U \right)^c = \bigcap_{U \in \mathcal{U}} U^c$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01G0 12. *Interaction With Complements III.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & \\
 \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & & \dashrightarrow \sim \\
 \mathcal{P}(\mathcal{P}(X)) & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\
 \downarrow \cap & & \downarrow \cup^{\text{op}} \\
 \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}},
 \end{array}$$

commutes, i.e. we have

$$\left(\bigcap_{U \in \mathcal{U}} U \right)^c = \bigcup_{U \in \mathcal{U}} U^c$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01G1 13. *Interaction With Symmetric Differences.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\Delta} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cup \times \cup & \text{X} & \downarrow \cup \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\Delta} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in \mathcal{U} \Delta \mathcal{V}} W \neq \left(\bigcup_{U \in \mathcal{U}} U \right) \Delta \left(\bigcup_{V \in \mathcal{V}} V \right)$$

in general, where $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01G2 14. *Interaction With Internal Homs I.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-1, -2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cup^{\text{op}} \times \cup^{\text{op}} & \text{X} & \downarrow \cup \\
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[\bigcup_{U \in \mathcal{U}} U, \bigcup_{V \in \mathcal{V}} V \right]_X$$

in general, where $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01G3 15. *Interaction With Internal Homs II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ \nearrow \sim & & \searrow \cup^{\text{op}} \\ \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\ \downarrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & & \downarrow [-, V]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[\bigcup_{U \in \mathcal{U}} U, V \right]_X = \bigcap_{U \in \mathcal{U}} [U, V]_X$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ and each $V \in \mathcal{P}(X)$.

01G4 16. *Interaction With Internal Homs III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \\ \downarrow \text{id}_{\mathcal{P}(X)} \star [U, -]_X & & \downarrow [U, -]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[U, \bigcup_{V \in \mathcal{V}} V \right]_X = \bigcup_{V \in \mathcal{V}} [U, V]_X$$

for each $U \in \mathcal{P}(X)$ and each $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

- 01G5 17. *Interaction With Direct Images.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f)_!} & \mathcal{P}(\mathcal{P}(Y)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_!(U) = \bigcup_{V \in f_!(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f)_!(\mathcal{U})$.

- 01G6 18. *Interaction With Inverse Images.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{V \in \mathcal{U}} f^{-1}(V) = \bigcup_{U \in f^{-1}(\mathcal{U})} U$$

for each $\mathcal{U} \in \mathcal{P}(Y)$, where $f^{-1}(\mathcal{U}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{U})$.

- 01G7 19. *Interaction With Codirect Images.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_*(U) = \bigcup_{V \in f_*(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$.

01G8 20. *Interaction With Intersections of Families I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \cap} & \mathcal{P}(\mathcal{P}(x)) \\ \downarrow \cup \star \text{id}_{\mathcal{P}(X)} & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{\cap} & X \end{array}$$

commutes, i.e. we have

$$\bigcap_{\substack{U \in \\ A \in \mathcal{A}}} U = \bigcap_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right)$$

for each $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$.


01G9 21. *Interaction With Intersections of Families II.* Let X be a set and consider the compositions

$$\begin{array}{c} \mathcal{P}(\mathcal{P}(\mathcal{P}(X))) \\ \swarrow \cap \star \text{id}_{\mathcal{P}(X)} \quad \downarrow \cup \star \text{id}_{\mathcal{P}(X)} \quad \searrow \text{id}_{\mathcal{P}(X)} \star \cap \quad \searrow \text{id}_{\mathcal{P}(X)} \star \cup \\ \mathcal{P}(\mathcal{P}(X)) \quad \mathcal{P}(\mathcal{P}(X)) \quad \mathcal{P}(\mathcal{P}(X)) \quad \mathcal{P}(\mathcal{P}(X)) \\ \swarrow \cup \quad \searrow \cap \quad \swarrow \cup \quad \searrow \cap \\ \mathcal{P}(X) \end{array}$$

given by

$$\begin{aligned} \mathcal{A} &\mapsto \bigcup_{U \in \bigcap_{A \in \mathcal{A}} A} U, & \mathcal{A} &\mapsto \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U, \\ \mathcal{A} &\mapsto \bigcup_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right), & \mathcal{A} &\mapsto \bigcap_{A \in \mathcal{A}} \left(\bigcup_{U \in A} U \right) \end{aligned}$$

for each $\mathcal{A} \in \mathcal{P}(\mathcal{P}(\mathcal{P}(X)))$. We have the following inclusions:



All other possible inclusions fail to hold in general.

Proof. Item 1, Functoriality: Since $\mathcal{P}(X)$ is posetal, it suffices to prove the condition (\star) . So let $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ with $\mathcal{U} \subset \mathcal{V}$. We claim that

$$\bigcup_{U \in \mathcal{U}} U \subset \bigcup_{V \in \mathcal{V}} V.$$

Indeed, given $x \in \bigcup_{U \in \mathcal{U}} U$, there exists some $U \in \mathcal{U}$ such that $x \in U$, but since $\mathcal{U} \subset \mathcal{V}$, we have $U \in \mathcal{V}$ as well, and thus $x \in \bigcup_{V \in \mathcal{V}} V$, which gives our desired inclusion.

Item 2, Associativity: We have

$$\begin{aligned}
 \bigcup_{A \in \mathcal{A}} U &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } U \in \bigcup_{A \in \mathcal{A}} A \\ \text{such that we have } x \in U \end{array} \right. \right\} \\
 &= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } A \in \mathcal{A} \\ \text{and some } U \in A \text{ such that} \\ \text{we have } x \in U \end{array} \right. \right\} \\
 &= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } A \in \mathcal{A} \\ \text{such that we have } x \in \bigcup_{U \in A} U \end{array} \right. \right\} \\
 &\stackrel{\text{def}}{=} \bigcup_{A \in \mathcal{A}} \left(\bigcup_{U \in A} U \right).
 \end{aligned}$$

This finishes the proof.

Item 3, Left Unitality: We have

$$\begin{aligned}
 \bigcup_{V \in \{U\}} V &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } V \in \{U\} \\ \text{such that we have } x \in V \end{array} \right. \right\} \\
 &= \{x \in X \mid x \in U\} \\
 &= U.
 \end{aligned}$$

This finishes the proof.

Item 4, Right Unitality: We have

$$\begin{aligned}
 \bigcup_{\{u\} \in \chi_X(U)} \{u\} &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } \{u\} \in \chi_X(U) \\ \text{such that we have } x \in \{u\} \end{array} \right. \right\} \\
 &= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } \{u\} \in \chi_X(U) \\ \text{such that we have } x = u \end{array} \right. \right\} \\
 &= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } u \in U \\ \text{such that we have } x = u \end{array} \right. \right\} \\
 &= \{x \in X \mid x \in U\}
 \end{aligned}$$

$$= U.$$

This finishes the proof.

Item 5, Interaction With Unions I: We have

$$\begin{aligned}
 \bigcup_{W \in \mathcal{U} \cup \mathcal{V}} W &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } W \in \mathcal{U} \cup \mathcal{V} \\ \text{such that we have } x \in W \end{array} \right. \right\} \\
 &= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } W \in \mathcal{U} \text{ or some } \\ W \in \mathcal{V} \text{ such that we have } x \in W \end{array} \right. \right\} \\
 &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } W \in \mathcal{U} \\ \text{such that we have } x \in W \end{array} \right. \right\} \\
 &\quad \cup \left\{ x \in X \left| \begin{array}{l} \text{there exists some } W \in \mathcal{V} \\ \text{such that we have } x \in W \end{array} \right. \right\} \\
 &\stackrel{\text{def}}{=} \left(\bigcup_{W \in \mathcal{U}} W \right) \cup \left(\bigcup_{W \in \mathcal{V}} W \right) \\
 &= \left(\bigcup_{U \in \mathcal{U}} U \right) \cup \left(\bigcup_{V \in \mathcal{V}} V \right).
 \end{aligned}$$

This finishes the proof.

Item 6, Interaction With Unions II: Omitted.

Item 7, Interaction With Intersections I: We have

$$\begin{aligned}
 \bigcup_{W \in \mathcal{U} \cap \mathcal{V}} W &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{there exists some } W \in \mathcal{U} \cap \mathcal{V} \\ \text{such that we have } x \in W \end{array} \right. \right\} \\
 &\subset \left\{ x \in X \left| \begin{array}{l} \text{there exists some } U \in \mathcal{U} \text{ and some } V \in \mathcal{V} \\ \text{such that we have } x \in U \text{ and } x \in V \end{array} \right. \right\} \\
 &= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } U \in \mathcal{U} \\ \text{such that we have } x \in U \end{array} \right. \right\} \\
 &\quad \cap \left\{ x \in X \left| \begin{array}{l} \text{there exists some } V \in \mathcal{V} \\ \text{such that we have } x \in V \end{array} \right. \right\} \\
 &\stackrel{\text{def}}{=} \left(\bigcup_{U \in \mathcal{U}} U \right) \cap \left(\bigcup_{V \in \mathcal{V}} V \right).
 \end{aligned}$$

This finishes the proof.

Item 8, Interaction With Intersections II: Omitted.

Item 9, Interaction With Differences: Let $X = \{0, 1\}$, let $\mathcal{U} = \{\{0, 1\}\}$, and let $\mathcal{V} = \{\{0\}\}$. We have

$$\begin{aligned} \bigcup_{W \in \mathcal{U} \setminus \mathcal{V}} W &= \bigcup_{W \in \{\{0, 1\}\}} W \\ &= \{0, 1\}, \end{aligned}$$

whereas

$$\begin{aligned} \left(\bigcup_{U \in \mathcal{U}} U \right) \setminus \left(\bigcup_{V \in \mathcal{V}} V \right) &= \{0, 1\} \setminus \{0\} \\ &= \{1\}. \end{aligned}$$

Thus we have

$$\bigcup_{W \in \mathcal{U} \setminus \mathcal{V}} W = \{0, 1\} \neq \{1\} = \left(\bigcup_{U \in \mathcal{U}} U \right) \setminus \left(\bigcup_{V \in \mathcal{V}} V \right).$$

This finishes the proof.

Item 10, Interaction With Complements I: Let $X = \{0, 1\}$ and let $\mathcal{U} = \{0\}$. We have

$$\begin{aligned} \bigcup_{U \in \mathcal{U}^c} U &= \bigcup_{U \in \{\emptyset, \{1\}, \{0, 1\}\}} U \\ &= \{0, 1\}, \end{aligned}$$

whereas

$$\begin{aligned} \bigcup_{U \in \mathcal{U}} U^c &= \{0\}^c \\ &= \{1\}. \end{aligned}$$

Thus we have

$$\bigcup_{U \in \mathcal{U}^c} U = \{0, 1\} \neq \{1\} = \bigcup_{U \in \mathcal{U}} U^c.$$

This finishes the proof.

Item 11, Interaction With Complements II: Omitted.

Item 12, Interaction With Complements III: Omitted.

Item 13, Interaction With Symmetric Differences: Let $X = \{0, 1\}$, let $\mathcal{U} = \{\{0, 1\}\}$, and let $\mathcal{V} = \{\{0\}, \{0, 1\}\}$. We have

$$\begin{aligned} \bigcup_{W \in \mathcal{U} \Delta \mathcal{V}} W &= \bigcup_{W \in \{\{0\}\}} W \\ &= \{0\}, \end{aligned}$$

whereas

$$\begin{aligned} \left(\bigcup_{U \in \mathcal{U}} U \right) \Delta \left(\bigcup_{V \in \mathcal{V}} V \right) &= \{0, 1\} \Delta \{0, 1\} \\ &= \emptyset, \end{aligned}$$

Thus we have

$$\bigcup_{W \in \mathcal{U} \Delta \mathcal{V}} W = \{0\} \neq \emptyset = \left(\bigcup_{U \in \mathcal{U}} U \right) \Delta \left(\bigcup_{V \in \mathcal{V}} V \right).$$

This finishes the proof.

Item 14, Interaction With Internal Homs I: This is a repetition of *Item 7* of *Definition 4.4.7.1.3* and is proved there.

Item 15, Interaction With Internal Homs II: This is a repetition of *Item 8* of *Definition 4.4.7.1.3* and is proved there.

Item 16, Interaction With Internal Homs III: This is a repetition of *Item 9* of *Definition 4.4.7.1.3* and is proved there.

Item 17, Interaction With Direct Images: This is a repetition of *Item 3* of *Definition 4.6.1.1.5* and is proved there.

Item 18, Interaction With Inverse Images: This is a repetition of *Item 3* of *Definition 4.6.2.1.3* and is proved there.

Item 19, Interaction With Codirect Images: This is a repetition of *Item 3* of *Definition 4.6.3.1.7* and is proved there.

Item 20, Interaction With Intersections of Families I: We have

$$\bigcap_{\substack{U \in \\ A \in \mathcal{A}}} A \stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } U \in \bigcup_{A \in \mathcal{A}} A, \\ \text{we have } x \in U \end{array} \right. \right\}$$

$$\begin{aligned}
&= \left\{ x \in X \mid \begin{array}{l} \text{for each } A \in \mathcal{A} \text{ and each} \\ U \in A, \text{ we have } x \in U \end{array} \right\} \\
&\stackrel{\text{def}}{=} \bigcap_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right).
\end{aligned}$$

This finishes the proof.

Item 21, Interaction With Intersections of Families II: Omitted. \square

003V 4.3.7 Intersections of Families of Subsets

Let X be a set and let $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

003W **Definition 4.3.7.1.1.** The **intersection of \mathcal{U}** is the set $\bigcap_{U \in \mathcal{U}} U$ defined by

$$\bigcap_{U \in \mathcal{U}} U \stackrel{\text{def}}{=} \left\{ x \in X \mid \begin{array}{l} \text{for each } U \in \mathcal{U}, \\ \text{we have } x \in U \end{array} \right\}.$$

01GA **Proposition 4.3.7.1.2.** Let X be a set.

01GB 1. *Functoriality.* The assignment $\mathcal{U} \mapsto \bigcap_{U \in \mathcal{U}} U$ defines a functor

$$\bigcap : (\mathcal{P}(\mathcal{P}(X)), \supset) \rightarrow (\mathcal{P}(X), \subset).$$

In particular, for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$, the following condition is satisfied:

$$(\star) \text{ If } \mathcal{U} \subset \mathcal{V}, \text{ then } \bigcap_{V \in \mathcal{V}} V \subset \bigcap_{U \in \mathcal{U}} U.$$

01GC 2. *Oplax Associativity.* We have a natural transformation

$$\begin{array}{ccc}
\mathcal{P}(\mathcal{P}(\mathcal{P}(X))) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \bigcap} & \mathcal{P}(\mathcal{P}(X)) \\
\bigcap \star \text{id}_{\mathcal{P}(X)} \downarrow & \wr & \downarrow \bigcap \\
\mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcap} & \mathcal{P}(X)
\end{array}$$

with components

$$\bigcap_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right) \subset \bigcap_{U \in \bigcap_{A \in \mathcal{A}} A} U$$

for each $\mathcal{A} \in \mathcal{P}(\mathcal{P}(\mathcal{P}(X)))$.

01GD 3. *Left Unitality*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \downarrow \chi_{\mathcal{P}(X)} & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcap_{V \in \{U\}} V = U.$$

for each $U \in \mathcal{P}(X)$.

01GE 4. *Oplax Right Unitality*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & & \\ \downarrow \mathcal{P}(\chi_X) & \searrow \text{id}_{\mathcal{P}(X)} & \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\bigcap} & \mathcal{P}(X) \end{array}$$

✗

does not commute in general, i.e. we may have

$$\bigcap_{\{x\} \in \chi_X(U)} \{x\} \neq U$$

in general, where $U \in \mathcal{P}(X)$. However, when U is nonempty, we have

$$\bigcap_{\{x\} \in \chi_X(U)} \{x\} \subset U.$$

01GF 5. *Interaction With Unions I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \times \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcap_{W \in \mathcal{U} \cup \mathcal{V}} W = \left(\bigcap_{U \in \mathcal{U}} U \right) \cap \left(\bigcap_{V \in \mathcal{V}} V \right)$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01GG 6. *Interaction With Unions II.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cup -)} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{U \cup -} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cup V)} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{- \cup V} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$\begin{aligned} U \cup \left(\bigcap_{V \in \mathcal{V}} V \right) &= \bigcap_{V \in \mathcal{V}} (U \cup V), \\ \left(\bigcap_{U \in \mathcal{U}} U \right) \cup V &= \bigcap_{U \in \mathcal{U}} (U \cup V) \end{aligned}$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ and each $U, V \in \mathcal{P}(X)$.

01GH 7. *Interaction With Intersections I.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \times \cap \downarrow & \supset & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X), \end{array}$$

with components

$$\left(\bigcap_{U \in \mathcal{U}} U \right) \cap \left(\bigcap_{V \in \mathcal{V}} V \right) \subset \bigcap_{W \in \mathcal{U} \cap \mathcal{V}} W$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01GJ 8. *Interaction With Intersections II.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (U \cap -)} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{U \cap -} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(\mathcal{P}(X))} \times (- \cap V)} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{- \cap V} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have

$$\begin{aligned} U \cup \left(\bigcap_{V \in \mathcal{V}} V \right) &= \bigcap_{V \in \mathcal{V}} (U \cup V), \\ \left(\bigcap_{U \in \mathcal{U}} U \right) \cup V &= \bigcap_{U \in \mathcal{U}} (U \cup V) \end{aligned}$$

for each $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ and each $U, V \in \mathcal{P}(X)$.

01GK 9. *Interaction With Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\setminus} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \times \cap \downarrow & \text{X} & \downarrow \cap \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\setminus} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in \mathcal{U} \setminus \mathcal{V}} W \neq \left(\bigcap_{U \in \mathcal{U}} U \right) \setminus \left(\bigcap_{V \in \mathcal{V}} V \right)$$

in general, where $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01GL IO. *Interaction With Complements I.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(\mathcal{P}(X)) \\
 \cap^{\text{op}} \downarrow & \text{X} & \downarrow \cap \\
 \mathcal{P}(X)^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in \mathcal{U}^c} W \neq \bigcap_{U \in \mathcal{U}} U^c$$

in general, where $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01GM II. *Interaction With Complements II.* The diagram

$$\begin{array}{ccccc}
 & & \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \\
 \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & & \text{---} \sim & & \searrow \\
 \mathcal{P}(\mathcal{P}(X)) & & & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\
 \downarrow \cap & & & & \downarrow \cup^{\text{op}} \\
 \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}}, & &
 \end{array}$$

commutes, i.e. we have

$$\left(\bigcap_{U \in \mathcal{U}} U \right)^c = \bigcup_{U \in \mathcal{U}} U^c$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01GN 12. *Interaction With Complements III.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & \\
 \text{id}_{\mathcal{P}(X)} \star (-)^c \nearrow & & \searrow \sim \\
 \mathcal{P}(\mathcal{P}(X)) & & \mathcal{P}(\mathcal{P}(X))^{\text{op}} \\
 \downarrow \cup & & \downarrow \cap^{\text{op}} \\
 \mathcal{P}(X) & \xrightarrow{(-)^c} & \mathcal{P}(X)^{\text{op}},
 \end{array}$$

commutes, i.e. we have

$$\left(\bigcup_{U \in \mathcal{U}} U \right)^c = \bigcap_{U \in \mathcal{U}} U^c$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01GP 13. *Interaction With Symmetric Differences.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\Delta} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cap \times \cap & \text{X} & \downarrow \cap \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\Delta} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in \mathcal{U} \Delta \mathcal{V}} W \neq \left(\bigcap_{U \in \mathcal{U}} U \right) \Delta \left(\bigcap_{V \in \mathcal{V}} V \right)$$

in general, where $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01GQ 14. *Interaction With Internal Homs I.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-1, -2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cap^{\text{op}} \times \cap^{\text{op}} & \text{X} & \downarrow \cap \\
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[\bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X$$

in general, where $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01GR 15. *Interaction With Internal Homs II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ \nearrow \sim & & \searrow \cap^{\text{op}} \\ \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\ \downarrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & & \downarrow [-, V]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[\bigcap_{U \in \mathcal{U}} U, V \right]_X = \bigcup_{U \in \mathcal{U}} [U, V]_X$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ and each $V \in \mathcal{P}(X)$.

01GS 16. *Interaction With Internal Homs III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \\ \downarrow \text{id}_{\mathcal{P}(X)} \star [U, -]_X & & \downarrow [U, -]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[U, \bigcap_{V \in \mathcal{V}} V \right]_X = \bigcap_{V \in \mathcal{V}} [U, V]_X$$

for each $U \in \mathcal{P}(X)$ and each $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

- 01GT** 17. *Interaction With Direct Images.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f!)_!} & \mathcal{P}(\mathcal{P}(Y)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f!(U) = \bigcap_{V \in f!(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f!(\mathcal{U}) \stackrel{\text{def}}{=} (f!)_!(\mathcal{U})$.

- 01GU** 18. *Interaction With Inverse Images.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcap_{V \in \mathcal{U}} f^{-1}(V) = \bigcap_{U \in f^{-1}(\mathcal{U})} U$$

for each $\mathcal{U} \in \mathcal{P}(Y)$, where $f^{-1}(\mathcal{U}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{U})$.

- 01GV** 19. *Interaction With Codirect Images.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_*(U) = \bigcap_{V \in f_*(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$.

01GW 20. *Interaction With Unions of Families I.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\text{id}_{\mathcal{P}(X)} \star \cap} & \mathcal{P}(\mathcal{P}(x)) \\ \downarrow \cup \star \text{id}_{\mathcal{P}(X)} & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{\cap} & X \end{array}$$

commutes, i.e. we have

$$\bigcap_{\substack{U \in \\ A \in \mathcal{A}}} U = \bigcap_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right)$$

for each $\mathcal{A} \in \mathcal{P}(\mathcal{P}(X))$.

01GX 21. *Interaction With Unions of Families II.* Let X be a set and consider the compositions

$$\begin{array}{c} \mathcal{P}(\mathcal{P}(\mathcal{P}(X))) \\ \swarrow \cap \star \text{id}_{\mathcal{P}(X)} \quad \downarrow \cup \star \text{id}_{\mathcal{P}(X)} \quad \searrow \text{id}_{\mathcal{P}(X)} \star \cap \quad \searrow \text{id}_{\mathcal{P}(X)} \star \cup \\ \mathcal{P}(\mathcal{P}(X)) \quad \mathcal{P}(\mathcal{P}(X)) \quad \mathcal{P}(\mathcal{P}(X)) \quad \mathcal{P}(\mathcal{P}(X)) \\ \swarrow \cup \quad \searrow \cap \quad \swarrow \cup \quad \searrow \cap \\ \mathcal{P}(X) \end{array}$$

given by

$$\begin{aligned} \mathcal{A} &\mapsto \bigcup_{U \in \bigcap_{A \in \mathcal{A}} A} U, & \mathcal{A} &\mapsto \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U, \\ \mathcal{A} &\mapsto \bigcup_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right), & \mathcal{A} &\mapsto \bigcap_{A \in \mathcal{A}} \left(\bigcup_{U \in A} U \right) \end{aligned}$$

for each $\mathcal{A} \in \mathcal{P}(\mathcal{P}(\mathcal{P}(X)))$. We have the following inclusions:

$$\begin{array}{ccc} & \bigcup_{U \in \bigcap_{A \in \mathcal{A}} A} U & \\ & \uparrow & \searrow \\ \bigcup_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right) & & \bigcap_{A \in \mathcal{A}} \left(\bigcup_{U \in A} U \right) \\ & \nwarrow \quad \nearrow & \\ & \bigcap_{U \in \bigcup_{A \in \mathcal{A}} A} U & \end{array}$$

All other possible inclusions fail to hold in general.

Proof. Item 1, Functoriality: Since $\mathcal{P}(X)$ is posetal, it suffices to prove the condition (\star) . So let $\mathcal{U}, \mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$ with $\mathcal{U} \subset \mathcal{V}$. We claim that

$$\bigcap_{V \in \mathcal{V}} V \subset \bigcap_{U \in \mathcal{U}} U.$$

Indeed, if $x \in \bigcap_{V \in \mathcal{V}} V$, then $x \in V$ for all $V \in \mathcal{V}$. But since $\mathcal{U} \subset \mathcal{V}$, it follows that $x \in U$ for all $U \in \mathcal{U}$ as well. Thus $x \in \bigcap_{U \in \mathcal{U}} U$, which gives our desired inclusion.

Item 2, Oplax Associativity: We have

$$\bigcap_{A \in \mathcal{A}} \left(\bigcap_{U \in A} U \right) \stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } A \in \mathcal{A}, \\ \text{we have } x \in \bigcap_{U \in A} U \end{array} \right. \right\}$$

$$\begin{aligned}
&\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } A \in \mathcal{A} \text{ and each} \\ U \in A, \text{ we have } x \in U \end{array} \right. \right\} \\
&= \left\{ x \in X \left| \begin{array}{l} \text{for each } U \in \bigcup_{A \in \mathcal{A}} A, \\ \text{we have } x \in U \end{array} \right. \right\} \\
&\subset \left\{ x \in X \left| \begin{array}{l} \text{for each } U \in \bigcap_{A \in \mathcal{A}} A, \\ \text{we have } x \in U \end{array} \right. \right\} \\
&\stackrel{\text{def}}{=} \bigcap_{U \in \bigcap_{A \in \mathcal{A}} A} U.
\end{aligned}$$

Since $\mathcal{P}(X)$ is posetal, naturality is automatic ([Categories](#), [Item 4](#) of [Definition 11.2.7.1.2](#)). This finishes the proof.

Item 3, Left Unitality: We have

$$\begin{aligned}
\bigcap_{V \in \{U\}} V &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } V \in \{U\}, \\ \text{we have } x \in U \end{array} \right. \right\} \\
&= \{x \in X \mid x \in U\} \\
&= U.
\end{aligned}$$

This finishes the proof.

Item 4, Oplax Right Unitality: If $U = \emptyset$, then we have

$$\begin{aligned}
\bigcap_{\{u\} \in \chi_X(U)} \{u\} &= \bigcap_{\{u\} \in \emptyset} \{u\} \\
&= X,
\end{aligned}$$

so $\bigcap_{\{u\} \in \chi_X(U)} \{u\} = X \neq \emptyset = U$. When U is nonempty, we have two cases:

020B I. If U is a singleton, say $U = \{u\}$, we have

$$\begin{aligned}
\bigcap_{\{u\} \in \chi_X(U)} \{u\} &= \{u\} \\
&\stackrel{\text{def}}{=} U.
\end{aligned}$$

020C 2. If U contains at least two elements, we have

$$\bigcap_{\{u\} \in \mathcal{X}(U)} \{u\} = \emptyset \\ \subset U.$$

This finishes the proof.

Item 5, Interaction With Unions I: We have

$$\begin{aligned} \bigcap_{W \in \mathcal{U} \cup \mathcal{V}} W &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } W \in \mathcal{U} \cup \mathcal{V}, \\ \text{we have } x \in W \end{array} \right. \right\} \\ &= \left\{ x \in X \left| \begin{array}{l} \text{for each } W \in \mathcal{U} \text{ and each} \\ W \in \mathcal{V}, \text{ we have } x \in W \end{array} \right. \right\} \\ &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } W \in \mathcal{U}, \\ \text{we have } x \in W \end{array} \right. \right\} \\ &\quad \cap \left\{ x \in X \left| \begin{array}{l} \text{for each } W \in \mathcal{V}, \\ \text{we have } x \in W \end{array} \right. \right\} \\ &\stackrel{\text{def}}{=} \left(\bigcap_{W \in \mathcal{U}} W \right) \cap \left(\bigcap_{W \in \mathcal{V}} W \right) \\ &= \left(\bigcap_{U \in \mathcal{U}} U \right) \cap \left(\bigcap_{V \in \mathcal{V}} V \right). \end{aligned}$$

This finishes the proof.

Item 6, Interaction With Unions II: Omitted.

Item 7, Interaction With Intersections I: We have

$$\begin{aligned} \left(\bigcap_{U \in \mathcal{U}} U \right) \cap \left(\bigcap_{V \in \mathcal{V}} V \right) &\stackrel{\text{def}}{=} \left\{ x \in X \left| \begin{array}{l} \text{for each } U \in \mathcal{U}, \\ \text{we have } x \in U \end{array} \right. \right\} \\ &\quad \cup \left\{ x \in X \left| \begin{array}{l} \text{for each } V \in \mathcal{V}, \\ \text{we have } x \in V \end{array} \right. \right\} \\ &= \left\{ x \in X \left| \begin{array}{l} \text{for each } W \in \mathcal{U} \cap \mathcal{V}, \\ \text{we have } x \in W \end{array} \right. \right\} \end{aligned}$$

$$\begin{aligned}
& \subset \left\{ x \in X \left| \begin{array}{l} \text{for each } W \in \mathcal{U} \cup \mathcal{V}, \\ \text{we have } x \in W \end{array} \right. \right\} \\
& \stackrel{\text{def}}{=} \bigcap_{W \in \mathcal{U} \cap \mathcal{V}} W.
\end{aligned}$$

Since $\mathcal{P}(X)$ is posetal, naturality is automatic ([Categories, Item 4](#) of [Definition II.2.7.I.2](#)). This finishes the proof.

Item 8, Interaction With Intersections II: Omitted.

Item 9, Interaction With Differences: Let $X = \{0, 1\}$, let $\mathcal{U} = \{\{0\}, \{0, 1\}\}$, and let $\mathcal{V} = \{\{0\}\}$. We have

$$\begin{aligned}
\bigcap_{W \in \mathcal{U} \setminus \mathcal{V}} U &= \bigcap_{W \in \{\{0, 1\}\}} W \\
&= \{0, 1\},
\end{aligned}$$

whereas

$$\begin{aligned}
\left(\bigcap_{U \in \mathcal{U}} U \right) \setminus \left(\bigcap_{V \in \mathcal{V}} V \right) &= \{0\} \setminus \{0\} \\
&= \emptyset.
\end{aligned}$$

Thus we have

$$\bigcap_{W \in \mathcal{U} \setminus \mathcal{V}} W = \{0, 1\} \neq \emptyset = \left(\bigcap_{U \in \mathcal{U}} U \right) \setminus \left(\bigcap_{V \in \mathcal{V}} V \right).$$

This finishes the proof.

Item 10, Interaction With Complements I: Let $X = \{0, 1\}$ and let $\mathcal{U} = \{\{0\}\}$. We have

$$\begin{aligned}
\bigcap_{W \in \mathcal{U}^c} U &= \bigcap_{W \in \{\emptyset, \{1\}, \{0, 1\}\}} W \\
&= \emptyset,
\end{aligned}$$

whereas

$$\bigcap_{U \in \mathcal{U}} U^c = \{0\}^c$$

$$= \{1\}.$$

Thus we have

$$\bigcap_{W \in \mathcal{U}^c} U = \emptyset \neq \{1\} = \bigcap_{U \in \mathcal{U}} U^c.$$

This finishes the proof.

Item 11, Interaction With Complements II: This is a repetition of **Item 12** of **Definition 4.3.6.1.2** and is proved there.

Item 12, Interaction With Complements III: This is a repetition of **Item 11** of **Definition 4.3.6.1.2** and is proved there.

Item 13, Interaction With Symmetric Differences: Let $X = \{0, 1\}$, let $\mathcal{U} = \{\{0, 1\}\}$, and let $\mathcal{V} = \{\{0\}, \{0, 1\}\}$. We have

$$\begin{aligned} \bigcap_{W \in \mathcal{U} \Delta \mathcal{V}} W &= \bigcap_{W \in \{\{0\}\}} W \\ &= \{0\}, \end{aligned}$$

whereas

$$\begin{aligned} \left(\bigcap_{U \in \mathcal{U}} U \right) \Delta \left(\bigcap_{V \in \mathcal{V}} V \right) &= \{0, 1\} \Delta \{0\} \\ &= \emptyset, \end{aligned}$$

Thus we have

$$\bigcap_{W \in \mathcal{U} \Delta \mathcal{V}} W = \{0\} \neq \emptyset = \left(\bigcap_{U \in \mathcal{U}} U \right) \Delta \left(\bigcap_{V \in \mathcal{V}} V \right).$$

This finishes the proof.

Item 14, Interaction With Internal Homs I: This is a repetition of **Item 10** of **Definition 4.4.7.1.3** and is proved there.

Item 15, Interaction With Internal Homs II: This is a repetition of **Item 11** of **Definition 4.4.7.1.3** and is proved there.

Item 16, Interaction With Internal Homs III: This is a repetition of **Item 12** of **Definition 4.4.7.1.3** and is proved there.

Item 17, Interaction With Direct Images: This is a repetition of **Item 4** of **Definition 4.6.1.1.5** and is proved there.

Item 18, Interaction With Inverse Images: This is a repetition of **Item 4** of **Definition 4.6.2.1.3** and is proved there.

Item 19, Interaction With Codirect Images: This is a repetition of **Item 4** of **Definition 4.6.3.1.7** and is proved there.

Item 20, Interaction With Unions of Families I: This is a repetition of **Item 20** of **Definition 4.3.6.1.2** and is proved there.

Item 21, Interaction With Unions of Families II: This is a repetition of **Item 21** of **Definition 4.3.6.1.2** and is proved there. \square

003G 4.3.8 Binary Unions

Let X be a set and let $U, V \in \mathcal{P}(X)$.

003H Definition 4.3.8.1.1. The **union of U and V** is the set $U \cup V$ defined by

$$\begin{aligned} U \cup V &\stackrel{\text{def}}{=} \bigcup_{z \in \{U, V\}} z \\ &\stackrel{\text{def}}{=} \{x \in X \mid x \in U \text{ or } x \in V\}. \end{aligned}$$

003J Proposition 4.3.8.1.2. Let X be a set.

003K 1. *Functoriality.* The assignments $U, V, (U, V) \mapsto U \cup V$ define functors

$$\begin{aligned} U \cup -: & (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ - \cup V: & (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cup -_2: & (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each $U, V, A, B \in \mathcal{P}(X)$:

01GY (a) If $U \subset A$, then $U \cup V \subset A \cup V$.

01GZ (b) If $V \subset B$, then $U \cup V \subset U \cup B$.

01H0 (c) If $U \subset A$ and $V \subset B$, then $U \cup V \subset A \cup B$.

003M 2. *Associativity*. The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \alpha_{\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \nearrow & & \searrow \text{id}_{\mathcal{P}(X)} \times \cup \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \cup \times \text{id}_{\mathcal{P}(X)} \searrow & & \searrow \cup \\
 & \mathcal{P}(X) \times \mathcal{P}(X) \xrightarrow{\cup} \mathcal{P}(X) &
 \end{array}$$

commutes, i.e. we have an equality of sets

$$(U \cup V) \cup W = U \cup (V \cup W)$$

for each $U, V, W \in \mathcal{P}(X)$.

003N 3. *Unitality*. The diagrams

$$\begin{array}{ccc}
 \text{pt} \times \mathcal{P}(X) & \xrightarrow{[\emptyset] \times \text{id}_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \lambda_{\mathcal{P}(X)}^{\text{Sets}} \searrow & & \downarrow \cup \\
 & \mathcal{P}(X) &
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{P}(X) \times \text{pt} & \xrightarrow{\text{id}_{\mathcal{P}(X)} \times [\emptyset]} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \rho_{\mathcal{P}(X)}^{\text{Sets}} \searrow & & \downarrow \cup \\
 & \mathcal{P}(X) &
 \end{array}$$

commute, i.e. we have equalities of sets

$$\emptyset \cup U = U,$$

$$U \cup \emptyset = U$$

for each $U \in \mathcal{P}(X)$.

003P 4. *Commutativity*. The diagram


$$\begin{array}{ccc}
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\sigma_{\mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 & \searrow \cup & \downarrow \cup \\
 & & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cup V = V \cup U$$

for each $U, V \in \mathcal{P}(X)$.

01H1 5. *Annihilation With X*. The diagrams




commute, i.e. we have equalities of sets

$$U \cup X = X,$$

$$X \cup V = X$$

for each $U, V \in \mathcal{P}(X)$.

003R 6. *Distributivity of Unions Over Intersections*. The diagrams




commute, i.e. we have equalities of sets

$$U \cup (V \cap W) = (U \cup V) \cap (U \cup W),$$

$$(U \cap V) \cup W = (U \cup W) \cap (V \cup W)$$

for each $U, V, W \in \mathcal{P}(X)$.

01H2 7. *Distributivity of Intersections Over Unions.* The diagrams




commute, i.e. we have equalities of sets

$$U \cap (V \cup W) = (U \cap V) \cup (U \cap W),$$

$$(U \cup V) \cap W = (U \cap W) \cup (V \cap W)$$

for each $U, V, W \in \mathcal{P}(X)$.

003Q 8. *Idempotency.* The diagram




commutes, i.e. we have an equality of sets

$$U \cup U = U$$

for each $U \in \mathcal{P}(X)$.

003L 9. *Via Intersections and Symmetric Differences.* The diagram



commutes, i.e. we have an equality of sets

$$U \cup V = (U \Delta V) \Delta (U \cap V)$$

for each $U, V \in \mathcal{P}(X)$.

003S 10. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each $U, V \in \mathcal{P}(X)$.

003T 11. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each $U, V \in \mathcal{P}(X)$.

01H3 12. *Interaction With Direct Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U \cup V) = f_!(U) \cup f_!(V)$$

for each $U, V \in \mathcal{P}(X)$.

01H4 13. *Interaction With Inverse Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cup V) = f^{-1}(U) \cup f^{-1}(V)$$

for each $U, V \in \mathcal{P}(Y)$.

- 01H5 14. *Interaction With Codirect Images.* Let $f: X \rightarrow Y$ be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \downarrow \cup & \subset & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U) \cup f_*(V) \subset f_*(U \cup V)$$

indexed by $U, V \in \mathcal{P}(X)$.

- 003U 15. *Interaction With Powersets and Semirings.* The quintuple $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$ is an idempotent commutative semiring.

Proof. **Item 1, Functoriality:** See [Pro25an].

Item 2, Associativity: See [Pro25ba].

Item 3, Unitality: This follows from [Pro25bd] and **Item 4**.

Item 4, Commutativity: See [Pro25bb].

Item 5, Annihilation With X : We have

$$\begin{aligned} U \cup X &\stackrel{\text{def}}{=} \{x \in X \mid x \in U \text{ or } x \in X\} \\ &= \{x \in X \mid x \in X\}, \\ &= X \end{aligned}$$

and

$$\begin{aligned} X \cup V &\stackrel{\text{def}}{=} \{x \in X \mid x \in X \text{ or } x \in V\} \\ &= \{x \in X \mid x \in X\} \\ &= X. \end{aligned}$$

This finishes the proof.

Item 6, Distributivity of Unions Over Intersections: See [Pro25az].

Item 7, Distributivity of Intersections Over Unions: See [Pro25aj].

Item 8, Idempotency: See [Pro25am].

Item 9, Via Intersections and Symmetric Differences: See [Pro25ay].

Item 10, Interaction With Characteristic Functions I: See [Pro25h].

Item 11, Interaction With Characteristic Functions II: See [Pro25h].

Item 12, Interaction With Direct Images: See [Pro25p].

Item 13, Interaction With Inverse Images: See [Pro25y].

Item 14, Interaction With Codirect Images: This is a repetition of *Item 5* of *Definition 4.6.3.1.7* and is proved there.

Item 15, Interaction With Powersets and Semirings: This follows from *Items 2* to *4* and *8* of this proposition and *Items 3* to *6* and *8* of *Definition 4.3.9.1.2*. \square

003X 4.3.9 Binary Intersections

Let X be a set and let $U, V \in \mathcal{P}(X)$.

003Y **Definition 4.3.9.1.1.** The **intersection of U and V** is the set $U \cap V$ defined by

$$\begin{aligned} U \cap V &\stackrel{\text{def}}{=} \bigcap_{z \in \{U, V\}} z \\ &\stackrel{\text{def}}{=} \{x \in X \mid x \in U \text{ or } x \in V\}. \end{aligned}$$

003Z **Proposition 4.3.9.1.2.** Let X be a set.

0040 1. *Functoriality.* The assignments $U, V, (U, V) \mapsto U \cap V$ define functors

$$\begin{aligned} U \cap -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cap V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cap -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each $U, V, A, B \in \mathcal{P}(X)$:

01H6 (a) If $U \subset A$, then $U \cap V \subset A \cap V$.

01H7 (b) If $V \subset B$, then $U \cap V \subset U \cap B$.

01H8 (c) If $U \subset A$ and $V \subset B$, then $U \cap V \subset A \cap B$.

0041 2. *Adjointness.* We have adjunctions

$$\begin{aligned} (U \cap - \dashv [U, -]_X): \mathcal{P}(X) &\overset{U \cap -}{\underset{[U, -]_X}{\rightleftarrows}} \mathcal{P}(X), \\ (- \cap V \dashv [V, -]_X): \mathcal{P}(X) &\overset{- \cap V}{\underset{[V, -]_X}{\rightleftarrows}} \mathcal{P}(X), \end{aligned}$$

witnessed by bijections

$$\mathrm{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \mathrm{Hom}_{\mathcal{P}(X)}(U, [V, W]_X),$$

$$\mathrm{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \mathrm{Hom}_{\mathcal{P}(X)}(V, [U, W]_X),$$

natural in $U, V, W \in \mathcal{P}(X)$, where

$$[-_1, -_2]_X: \mathcal{P}(X)^{\mathrm{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

is the bifunctor of [Section 4.4.7](#). In particular, the following statements hold for each $U, V, W \in \mathcal{P}(X)$:

01H9 (a) The following conditions are equivalent:

01HA i. We have $U \cap V \subset W$.

01HB ii. We have $U \subset [V, W]_X$.

01HC (b) The following conditions are equivalent:

01HD i. We have $U \cap V \subset W$.

01HE ii. We have $V \subset [U, W]_X$.

0042 3. *Associativity*. The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(X) \times (\mathcal{P}(X) \times \mathcal{P}(X)) & \\
 \alpha_{\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\mathrm{Sets}} \swarrow \quad \searrow \mathrm{id}_{\mathcal{P}(X)} \times \cap & & \\
 (\mathcal{P}(X) \times \mathcal{P}(X)) \times \mathcal{P}(X) & & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \cap \times \mathrm{id}_{\mathcal{P}(X)} \searrow & & \searrow \cap \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$(U \cap V) \cap W = U \cap (V \cap W)$$

for each $U, V, W \in \mathcal{P}(X)$.

0043 4. *Unitality*. The diagrams

$$\begin{array}{ccc}
 \text{pt} \times \mathcal{P}(X) & \xrightarrow{[X] \times \text{id}_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \searrow \scriptstyle \sim_{\mathcal{P}(X)}^{\text{Sets}} & & \downarrow \cap \\
 & & \mathcal{P}(X)
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{P}(X) \times \text{pt} & \xrightarrow{\text{id}_{\mathcal{P}(X)} \times [X]} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \searrow \scriptstyle \sim_{\mathcal{P}(X)}^{\text{Sets}} & & \downarrow \cap \\
 & & \mathcal{P}(X)
 \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned}
 X \cap U &= U, \\
 U \cap X &= U
 \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

0044 5. *Commutativity*. The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\sigma_{\mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}}} & \mathcal{P}(X) \times \mathcal{P}(X) \\
 \searrow \cap & & \downarrow \cap \\
 & & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cap V = V \cap U$$

for each $U, V \in \mathcal{P}(X)$.

0047 6. *Annihilation With the Empty Set*. The diagrams

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \text{id}_{\text{pt}} \times \varepsilon_{\mathcal{P}(X)}^{\text{Sets}} \nearrow & & \nwarrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \\
 \text{pt} \times \mathcal{P}(X) & & \text{pt} \\
 \downarrow [\emptyset] \times \text{id}_{\mathcal{P}(X)} & & \downarrow [\emptyset] \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X)
 \end{array}
 \quad
 \begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \varepsilon_{\mathcal{P}(X)}^{\text{Sets}} \times \text{id}_{\text{pt}} \nearrow & & \nwarrow \mu_{4|\mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}} \\
 \mathcal{P}(X) \times \text{pt} & & \text{pt} \\
 \downarrow \text{id}_{\mathcal{P}(X)} \times [\emptyset] & & \downarrow [\emptyset] \\
 \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X)
 \end{array}$$


commute, i.e. we have equalities of sets

$$\begin{aligned}\emptyset \cap X &= \emptyset, \\ X \cap \emptyset &= \emptyset\end{aligned}$$

for each $U \in \mathcal{P}(X)$.

01HF

7. *Distributivity of Unions Over Intersections.* The diagrams




commute, i.e. we have equalities of sets

$$\begin{aligned}U \cup (V \cap W) &= (U \cup V) \cap (U \cup W), \\ (U \cap V) \cup W &= (U \cup W) \cap (V \cup W)\end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

0046

8. *Distributivity of Intersections Over Unions.* The diagrams



commute, i.e. we have equalities of sets

$$\begin{aligned}U \cap (V \cup W) &= (U \cap V) \cup (U \cap W), \\ (U \cup V) \cap W &= (U \cap W) \cup (V \cap W)\end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

0045 9. *Idempotency*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{\Delta_{\mathcal{P}(X)}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ & \searrow \text{id}_{\mathcal{P}(X)} & \downarrow \cap \\ & & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$U \cap U = U$$

for each $U \in \mathcal{P}(X)$.

0048 10. *Interaction With Characteristic Functions I*. We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each $U, V \in \mathcal{P}(X)$.

0049 11. *Interaction With Characteristic Functions II*. We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each $U, V \in \mathcal{P}(X)$.

01HG 12. *Interaction With Direct Images*. Let $f: X \rightarrow Y$ be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & \subset & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U \cap V) \subset f_!(U) \cap f_!(V)$$

indexed by $U, V \in \mathcal{P}(X)$.

- 01HH 13. *Interaction With Inverse Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cap V) = f^{-1}(U) \cap f^{-1}(V)$$

for each $U, V \in \mathcal{P}(Y)$.

- 01HJ 14. *Interaction With Codirect Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U) \cap f_*(V) = f_*(U \cap V)$$

for each $U, V \in \mathcal{P}(X)$.

- 004A 15. *Interaction With Powersets and Monoids With Zero.* The quadruple $((\mathcal{P}(X), \emptyset), \cap, X)$ is a commutative monoid with zero.

- 004B 16. *Interaction With Powersets and Semirings.* The quintuple $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$ is an idempotent commutative semiring.

Proof. **Item 1, Functoriality:** See [Pro25al].

Item 2, Adjointness: See [MSE 267469].

Item 3, Associativity: See [Pro25r].

Item 4, Unitality: This follows from [Pro25v] and **Item 5**.

Item 5, Commutativity: See [Pro25s].

Item 6, Annihilation With the Empty Set: This follows from [Pro25t] and **Item 5**.

Item 7, Distributivity of Unions Over Intersections: See [Pro25az].

Item 8, Distributivity of Intersections Over Unions: See [Pro25aj].

Item 9, Idempotency: See [Pro25ak].

Item 10, Interaction With Characteristic Functions I: See [Pro25e].

Item 11, Interaction With Characteristic Functions II: See [Pro25e].

Item 12, Interaction With Direct Images: See [Pro25n].

Item 13, Interaction With Inverse Images: See [Pro25w].

Item 14, Interaction With Codirect Images: This is a repetition of **Item 6** of **Definition 4.6.3.1.7** and is proved there.

Item 15, Interaction With Powersets and Monoids With Zero: This follows from **Items 3** to **6**.

Item 16, Interaction With Powersets and Semirings: This follows from **Items 2** to **4** and **8** and **Items 3** to **6** and **8** of **Definition 4.3.9.1.2**. \square

004D 4.3.10 Differences

Let X and Y be sets.

004E **Definition 4.3.10.1.1.** The **difference of X and Y** is the set $X \setminus Y$ defined by

$$X \setminus Y \stackrel{\text{def}}{=} \{a \in X \mid a \notin Y\}.$$

004F **Proposition 4.3.10.1.2.** Let X be a set.

004G I. *Functoriality.* The assignments $U, V, (U, V) \mapsto U \cap V$ define functors

$$\begin{aligned} U \setminus -: & \quad (\mathcal{P}(X), \supset) & \rightarrow (\mathcal{P}(X), \subset), \\ - \setminus V: & \quad (\mathcal{P}(X), \subset) & \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \setminus -_2: & \quad (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) & \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each $U, V, A, B \in \mathcal{P}(X)$:

01HK (a) If $U \subset A$, then $U \setminus V \subset A \setminus V$.

01HL (b) If $V \subset B$, then $U \setminus B \subset U \setminus V$.

01HM (c) If $U \subset A$ and $V \subset B$, then $U \setminus B \subset A \setminus V$.

004H 2. *De Morgan's Laws.* We have equalities of sets

$$\begin{aligned} X \setminus (U \cup V) &= (X \setminus U) \cap (X \setminus V), \\ X \setminus (U \cap V) &= (X \setminus U) \cup (X \setminus V) \end{aligned}$$

for each $U, V \in \mathcal{P}(X)$.

004J 3. *Interaction With Unions I.* We have equalities of sets

$$U \setminus (V \cup W) = (U \setminus V) \cap (U \setminus W)$$

for each $U, V, W \in \mathcal{P}(X)$.

004K 4. *Interaction With Unions II.* We have equalities of sets

$$(U \setminus V) \cup W = (U \cup W) \setminus (V \setminus W)$$

for each $U, V, W \in \mathcal{P}(X)$.

004L 5. *Interaction With Unions III.* We have equalities of sets

$$\begin{aligned} U \setminus (V \cup W) &= (U \cup W) \setminus (V \cup W) \\ &= (U \setminus V) \setminus W \\ &= (U \setminus W) \setminus V \end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

004M 6. *Interaction With Unions IV.* We have equalities of sets

$$(U \cup V) \setminus W = (U \setminus W) \cup (V \setminus W)$$

for each $U, V, W \in \mathcal{P}(X)$.

004N 7. *Interaction With Intersections.* We have equalities of sets

$$\begin{aligned} (U \setminus V) \cap W &= (U \cap W) \setminus V \\ &= U \cap (W \setminus V) \end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

- 004P 8. *Interaction With Complements.* We have an equality of sets

$$U \setminus V = U \cap V^c$$

for each $U, V \in \mathcal{P}(X)$.

- 004Q 9. *Interaction With Symmetric Differences.* We have an equality of sets

$$U \setminus V = U \Delta (U \cap V)$$

for each $U, V \in \mathcal{P}(X)$.

- 004R 10. *Triple Differences.* We have

$$U \setminus (V \setminus W) = (U \cap W) \cup (U \setminus V)$$

for each $U, V, W \in \mathcal{P}(X)$.

- 004S 11. *Left Annihilation.* We have

$$\emptyset \setminus U = \emptyset$$

for each $U \in \mathcal{P}(X)$.

- 004T 12. *Right Unitality.* We have

$$U \setminus \emptyset = U$$

for each $U \in \mathcal{P}(X)$.

- 01HN 13. *Right Annihilation.* We have

$$U \setminus X = \emptyset$$

for each $U \in \mathcal{P}(X)$.

- 004U 14. *Invertibility.* We have

$$U \setminus U = \emptyset$$

for each $U \in \mathcal{P}(X)$.

- 004V 15. *Interaction With Containment.* The following conditions are equivalent:

004W (a) We have $V \setminus U \subset W$.

004X (b) We have $V \setminus W \subset U$.

004Y 16. *Interaction With Characteristic Functions.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each $U, V \in \mathcal{P}(X)$.

01HP 17. *Interaction With Direct Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow \setminus & \wr & \downarrow \setminus \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \setminus f_!(V) \subset f_!(U \setminus V)$$

indexed by $U, V \in \mathcal{P}(X)$.

01HQ 18. *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \downarrow \setminus & & \downarrow \setminus \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \setminus V) = f^{-1}(U) \setminus f^{-1}(V)$$

for each $U, V \in \mathcal{P}(X)$.

01HR 19. *Interaction With Codirect Images.* We have a natural transformation

$$\begin{array}{ccc}
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\
 \downarrow & \wr & \downarrow \\
 \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)
 \end{array}$$

with components

$$f_!(U) \setminus f_!(V) \subset f_!(U \setminus V)$$

indexed by $U, V \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** See [Pro25ad] and [Pro25ah].

Item 2, De Morgan's Laws: See [Pro25k].

Item 3, Interaction With Unions I: See [Pro25l].

Item 4, Interaction With Unions II: Omitted.

Item 5, Interaction With Unions III: See [Pro25ai].

Item 6, Interaction With Unions IV: See [Pro25ac].

Item 7, Interaction With Intersections: See [Pro25u].

Item 8, Interaction With Complements: See [Pro25aa].

Item 9, Interaction With Symmetric Differences: See [Pro25ab].

Item 10, Triple Differences: See [Pro25ag].

Item 11, Left Annihilation: Omitted.

Item 12, Right Unitality: See [Pro25ae].

Item 13, Right Annihilation: Omitted.

Item 14, Invertibility: See [Pro25af].

Item 15, Interaction With Containment: Omitted.

Item 16, Interaction With Characteristic Functions: See [Pro25f].

Item 17, Interaction With Direct Images: See [Pro25o].

Item 18, Interaction With Inverse Images: See [Pro25x]. □

004Z 4.3.II Complements

Let X be a set and let $U \in \mathcal{P}(X)$.

0050 Definition 4.3.II.1.1. The **complement of U** is the set U^c defined by

$$\begin{aligned} U^c &\stackrel{\text{def}}{=} X \setminus U \\ &\stackrel{\text{def}}{=} \{a \in X \mid a \notin U\}. \end{aligned}$$

0051 Proposition 4.3.II.1.2. Let X be a set.

0052 1. *Functoriality.* The assignment $U \mapsto U^c$ defines a functor

$$(-)^c: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X).$$

In particular, the following statements hold for each $U, V \in \mathcal{P}(X)$:

(★) If $U \subset V$, then $V^c \subset U^c$.

0053 2. *De Morgan's Laws.* The diagrams

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} & \xrightarrow{\cup^{\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ \downarrow (-)^c \times (-)^c & & \downarrow (-)^c \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array} \quad \begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} & \xrightarrow{\cap^{\text{op}}} & \mathcal{P}(X)^{\text{op}} \\ \downarrow (-)^c \times (-)^c & & \downarrow (-)^c \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commute, i.e. we have equalities of sets

$$\begin{aligned} (U \cup V)^c &= U^c \cap V^c, \\ (U \cap V)^c &= U^c \cup V^c \end{aligned}$$

for each $U, V \in \mathcal{P}(X)$.

0054 3. *Involutority.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{(-)^c} & \mathcal{P}(X) \\ \searrow \text{id}_{\mathcal{P}(X)^{\text{op}}} & & \downarrow (-)^{c, \text{op}} \\ & & \mathcal{P}(X)^{\text{op}} \end{array}$$

commutes, i.e. we have

$$(U^c)^c = U$$

for each $U \in \mathcal{P}(X)$.

0055 4. *Interaction With Characteristic Functions.* We have

$$\chi_{U^c} \equiv 1 - \chi_U \pmod{2}$$

for each $U \in \mathcal{P}(X)$.

01HS 5. *Interaction With Direct Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U^c) = f_*(U)^c$$

for each $U \in \mathcal{P}(X)$.

01HT 6. *Interaction With Inverse Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1, \text{op}}} & \mathcal{P}(X)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U^c) = f^{-1}(U)^c$$

for each $U \in \mathcal{P}(X)$.

01HU 7. *Interaction With Codirect Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U^c) = f_!(U)^c$$

for each $U \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** This follows from **Item 1** of **Definition 4.3.10.1.2**.

Item 2, De Morgan's Laws: See [Pro25k].

Item 3, Involutority: See [Pro25i].

Item 4, Interaction With Characteristic Functions: Omitted.

Item 5, Interaction With Direct Images: This is a repetition of **Item 8** of **Definition 4.6.1.1.5** and is proved there.

Item 6, Interaction With Inverse Images: This is a repetition of **Item 8** of **Definition 4.6.2.1.3** and is proved there.

Item 7, Interaction With Codirect Images: This is a repetition of **Item 7** of **Definition 4.6.3.1.7** and is proved there. \square

0056 4.3.12 Symmetric Differences

Let X be a set and let $U, V \in \mathcal{P}(X)$.

0057 **Definition 4.3.12.1.1.** The **symmetric difference of U and V** is the set $U \Delta V$ defined by¹³


$$U \Delta V \stackrel{\text{def}}{=} (U \setminus V) \cup (V \setminus U).$$

0058 **Proposition 4.3.12.1.2.** Let X be a set.

0059 1. *Lack of Functoriality.* The assignment $(U, V) \mapsto U \Delta V$ **does not** in general define functors

$$\begin{aligned} U \Delta -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \Delta V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \Delta -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

¹³*Illustration:*



005A 2. *Via Unions and Intersections.* We have

$$U \Delta V = (U \cup V) \setminus (U \cap V)$$


for each $U, V \in \mathcal{P}(X)$, as in the Venn diagram



01HV 3. *Symmetric Differences of Disjoint Sets.* If U and V are disjoint, then we have

$$U \Delta V = U \cup V.$$


005B 4. *Associativity.* The diagram




commutes, i.e. we have

$$(U \Delta V) \Delta W = U \Delta (V \Delta W)$$

for each $U, V, W \in \mathcal{P}(X)$, as in the Venn diagram



005D 5. *Unitality.* The diagrams



commute, i.e. we have

$$\begin{aligned} U \Delta \emptyset &= U, \\ \emptyset \Delta U &= U \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

005C 6. *Commutativity.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\sigma_{\mathcal{P}(X), \mathcal{P}(X)}^{\text{Sets}}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ & \searrow \Delta & \downarrow \Delta \\ & & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$U \Delta V = V \Delta U$$

for each $U, V \in \mathcal{P}(X)$.

005E 7. *Invertibility.* We have

$$U \Delta U = \emptyset$$

for each $U \in \mathcal{P}(X)$.

005F 8. *Interaction With Unions.* We have

$$(U \Delta V) \cup (V \Delta T) = (U \cup V \cup W) \setminus (U \cap V \cap W)$$

for each $U, V, W \in \mathcal{P}(X)$.

005G 9. *Interaction With Complements I.* We have

$$U \Delta U^c = X$$

for each $U \in \mathcal{P}(X)$.

005H 10. *Interaction With Complements II.* We have

$$\begin{aligned} U \triangle X &= U^c, \\ X \triangle U &= U^c \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

005J 11. *Interaction With Complements III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\Delta} & \mathcal{P}(X) \\ (-)^c \times (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow[\Delta]{} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$U^c \triangle V^c = U \triangle V$$

for each $U, V \in \mathcal{P}(X)$.

005K 12. *“Transitivity”.* We have

$$(U \triangle V) \triangle (V \triangle W) = U \triangle W$$

for each $U, V, W \in \mathcal{P}(X)$.

005L 13. *The Triangle Inequality for Symmetric Differences.* We have

$$U \triangle W \subset U \triangle V \cup V \triangle W$$

for each $U, V, W \in \mathcal{P}(X)$.

005M 14. *Distributivity Over Intersections.* We have

$$\begin{aligned} U \cap (V \triangle W) &= (U \cap V) \triangle (U \cap W), \\ (U \triangle V) \cap W &= (U \cap W) \triangle (V \cap W) \end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

005N 15. *Interaction With Characteristic Functions.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each $U, V \in \mathcal{P}(X)$.

005P 16. *Bijectivity.* Given $U, V \in \mathcal{P}(X)$, the maps

$$U \Delta -: \mathcal{P}(X) \rightarrow \mathcal{P}(X),$$

$$- \Delta V: \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

are bijections with inverses given by

$$(U \Delta -)^{-1} = - \cup (U \cap -),$$

$$(- \Delta V)^{-1} = - \cup (V \cap -).$$

Moreover, the map

$$\mathcal{P}(X) \longrightarrow \mathcal{P}(X)$$

$$C \longmapsto C \Delta (U \Delta V)$$

is a bijection of $\mathcal{P}(X)$ onto itself sending U to V and V to U .

005Q 17. *Interaction With Powersets and Groups.* Let X be a set.

005R (a) The quadruple $(\mathcal{P}(X), \Delta, \emptyset, \text{id}_{\mathcal{P}(X)})$ is an abelian group.¹⁴

¹⁴Here are some examples:

020H 1. When $X = \emptyset$, we have an isomorphism of groups between $\mathcal{P}(\emptyset)$ and the trivial group:

$$(\mathcal{P}(\emptyset), \Delta, \emptyset, \text{id}_{\mathcal{P}(\emptyset)}) \cong \text{pt.}$$

020J 2. When $X = \text{pt}$, we have an isomorphism of groups between $\mathcal{P}(\text{pt})$ and $\mathbb{Z}/2$:

$$(\mathcal{P}(\text{pt}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\text{pt})}) \cong \mathbb{Z}/2.$$

005S (b) Every element of $\mathcal{P}(X)$ has order 2 with respect to Δ , and thus $\mathcal{P}(X)$ is a *Boolean group* (i.e. an abelian 2-group).

005T 4. *Interaction With Powersets and Vector Spaces I.* The pair $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$ consisting of

- The group $\mathcal{P}(X)$ of **Item 17**;
- The map $\alpha_{\mathcal{P}(X)}: \mathbb{F}_2 \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ defined by

$$\begin{aligned} 0 \cdot U &\stackrel{\text{def}}{=} \emptyset, \\ 1 \cdot U &\stackrel{\text{def}}{=} U; \end{aligned}$$

is an \mathbb{F}_2 -vector space.

005U 5. *Interaction With Powersets and Vector Spaces II.* If X is finite, then:

020L (a) The set of singletons sets on the elements of X forms a basis for the \mathbb{F}_2 -vector space $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$ of **Item 4**.

020M (b) We have

$$\dim(\mathcal{P}(X)) = \#X.$$


005V 6. *Interaction With Powersets and Rings.* The quintuple $(\mathcal{P}(X), \Delta, \cap, \emptyset, X)$ is a commutative ring.¹⁵

01HW 7. *Interaction With Direct Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \supset & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

020K 3. When $X = \{0, 1\}$, we have an isomorphism of groups between $\mathcal{P}(\{0, 1\})$ and $\mathbb{Z}/2 \times \mathbb{Z}/2$:

$$(\mathcal{P}(\{0, 1\}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\{0, 1\})}) \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

¹⁵  **Warning:** The analogous statement replacing intersections by unions (i.e. that the quintuple $(\mathcal{P}(X), \Delta, \cup, \emptyset, X)$ is a ring) is false, however. See [**Pro25aw**] for a proof.

with components

$$f_!(U) \Delta f_!(V) \subset f_!(U \Delta V)$$

indexed by $U, V \in \mathcal{P}(X)$.

01HX 8. *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \Delta \downarrow & & \downarrow \Delta \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

i.e. we have

$$f^{-1}(U) \Delta f^{-1}(V) = f^{-1}(U \Delta V)$$

for each $U, V \in \mathcal{P}(Y)$.

01HY 9. *Interaction With Codirect Images.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \wr & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U \Delta V) \subset f_*(U) \Delta f_*(V)$$

indexed by $U, V \in \mathcal{P}(X)$.

Proof. **Item 1, Lack of Functoriality:** Omitted.

Item 2, Via Unions and Intersections: See [Pro25m].

Item 3, Symmetric Differences of Disjoint Sets: Since U and V are disjoint, we have $U \cap V = \emptyset$, and therefore we have

$$U \Delta V = (U \cup V) \setminus (U \cap V)$$

$$\begin{aligned}
&= (U \cup V) \setminus \emptyset \\
&= U \cup V,
\end{aligned}$$

where we've used [Item 2](#) and [Item 12](#) of [Definition 4.3.10.1.2](#).

Item 4, Associativity: See [[Pro25ao](#)].

Item 5, Unitality: This follows from [Item 6](#) and [[Pro25at](#)].

Item 6, Commutativity: See [[Pro25ap](#)].

Item 7, Invertibility: See [[Pro25av](#)].

Item 8, Interaction With Unions: See [[Pro25bc](#)].

Item 9, Interaction With Complements I: See [[Pro25as](#)].

Item 10, Interaction With Complements II: This follows from [Item 6](#) and [[Pro25ax](#)].

Item 11, Interaction With Complements III: See [[Pro25aq](#)].

Item 12, "Transitivity": We have

$$\begin{aligned}
(U \triangle V) \triangle (V \triangle W) &= U \triangle (V \triangle (V \triangle W)) && \text{(by [Item 4](#))} \\
&= U \triangle ((V \triangle V) \triangle W) && \text{(by [Item 4](#))} \\
&= U \triangle (\emptyset \triangle W) && \text{(by [Item 7](#))} \\
&= U \triangle W. && \text{(by [Item 5](#))}
\end{aligned}$$

This finishes the proof.

Item 13, The Triangle Inequality for Symmetric Differences: This follows from [Items 2](#) and [12](#).

Item 14, Distributivity Over Intersections: See [[Pro25q](#)].

Item 15, Interaction With Characteristic Functions: See [[Pro25g](#)].

Item 16, Bijectivity: Omitted.

Item 17, Interaction With Powersets and Groups: [Item 17a](#) follows from [Items 4](#) to [7](#), while [Item 3b](#) follows from [Item 7](#).¹⁶

Item 4, Interaction With Powersets and Vector Spaces I: See [[MSE 2719059](#)].

Item 5, Interaction With Powersets and Vector Spaces II: See [[MSE 2719059](#)].

Item 6, Interaction With Powersets and Rings: This follows from [Items 6](#) and [15](#) of [Definition 4.3.9.1.2](#) and [Items 14](#) and [17](#).¹⁷

Item 7, Interaction With Direct Images: This is a repetition of [Item 9](#) of [Definition 4.6.1.1.5](#) and is proved there.

Item 8, Interaction With Inverse Images: This is a repetition of [Item 9](#) of [Definition 4.6.2.1.3](#) and is proved there.

Item 9, Interaction With Codirect Images: This is a repetition of [Item 8](#) of [Defini-](#)

¹⁶Reference: [[Pro25ar](#)].

¹⁷Reference: [[Pro25au](#)].

tion 4.6.3.1.7 and is proved there. \square

005W 4.4 Powersets

01HZ 4.4.1 Foundations

Let X be a set.

006P **Definition 4.4.1.1.1.** The **powerset of X** is the set $\mathcal{P}(X)$ defined by

$$\mathcal{P}(X) \stackrel{\text{def}}{=} \{U \in P \mid U \subset X\},$$

where P is the set in the axiom of powerset, ?? of ??.

006Q **Remark 4.4.1.1.2.** Under the analogy that $\{\mathbf{t}, \mathbf{f}\}$ should be the (-1) -categorical analogue of **Sets**, we may view the powerset of a set as a decategorification of the category of presheaves of a category (or of the category of copresheaves):

- The powerset of a set X is equivalently (Item 2 of Definition 4.5.1.1.4) the set

$$\mathbf{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$$

of functions from X to the set $\{\mathbf{t}, \mathbf{f}\}$ of classical truth values.

- The category of presheaves on a category C is the category

$$\mathbf{Fun}(C^{\text{op}}, \mathbf{Sets})$$

of functors from C^{op} to the category **Sets** of sets.

01J0 **Notation 4.4.1.1.3.** Let X be a set.

01J1 1. We write $\mathcal{P}_0(X)$ for the set of nonempty subsets of X .

01J2 2. We write $\mathcal{P}_{\text{fin}}(X)$ for the set of finite subsets of X .

01J3 **Proposition 4.4.1.1.4.** Let X be a set.

01J4 1. *Co/Completeness.* The (posetal) category (associated to) $(\mathcal{P}(X), \subset)$ is complete and cocomplete:

020P (a) *Products.* The products in $\mathcal{P}(X)$ are given by intersection of subsets.

- 020Q (b) *Coproducts*. The coproducts in $\mathcal{P}(X)$ are given by union of subsets.
- 020R (c) *Co/Equalisers*. Being a posetal category, $\mathcal{P}(X)$ only has at most one morphisms between any two objects, so co/equalisers are trivial.
- 01J5 2. *Cartesian Closedness*. The category $\mathcal{P}(X)$ is Cartesian closed.
- 01J6 3. *Powersets as Sets of Relations*. We have bijections

$$\begin{aligned}\mathcal{P}(X) &\cong \text{Rel}(\text{pt}, X), \\ \mathcal{P}(X) &\cong \text{Rel}(X, \text{pt}),\end{aligned}$$

natural in $X \in \text{Obj}(\text{Sets})$.

- 01J7 4. *Interaction With Products I*. The map

$$\begin{aligned}\mathcal{P}(X) \times \mathcal{P}(Y) &\longrightarrow \mathcal{P}(X \amalg Y) \\ (U, V) &\longmapsto U \cup V\end{aligned}$$

is an isomorphism of sets, natural in $X, Y \in \text{Obj}(\text{Sets})$ with respect to each of the functor structures $\mathcal{P}_!$, \mathcal{P}^{-1} , and \mathcal{P}_* on \mathcal{P} of [Definition 4.4.2.1.1](#). Moreover, this makes each of $\mathcal{P}_!$, \mathcal{P}^{-1} , and \mathcal{P}_* into a symmetric monoidal functor.

- 01J8 5. *Interaction With Products II*. The map

$$\begin{aligned}\mathcal{P}(X) \times \mathcal{P}(Y) &\longrightarrow \mathcal{P}(X \amalg Y) \\ (U, V) &\longmapsto U \boxtimes_{X \times Y} V,\end{aligned}$$

where¹⁸

$$U \boxtimes_{X \times Y} V \stackrel{\text{def}}{=} \{(u, v) \in X \times Y \mid u \in U \text{ and } v \in V\}$$

is an inclusion of sets, natural in $X, Y \in \text{Obj}(\text{Sets})$ with respect to each of the functor structures $\mathcal{P}_!$, \mathcal{P}^{-1} , and \mathcal{P}_* on \mathcal{P} of [Definition 4.4.2.1.1](#). Moreover, this makes each of $\mathcal{P}_!$, \mathcal{P}^{-1} , and \mathcal{P}_* into a symmetric monoidal functor.

¹⁸The set $U \boxtimes_{X \times Y} V$ is usually denoted simply $U \times V$. Here we denote it in this somewhat

01J9 6. *Interaction With Products III*. We have an isomorphism

$$\mathcal{P}(X) \otimes \mathcal{P}(Y) \cong \mathcal{P}(X \times Y),$$

natural in $X, Y \in \text{Obj}(\text{Sets})$ with respect to each of the functor structures $\mathcal{P}_!$, \mathcal{P}^{-1} , and \mathcal{P}_* on \mathcal{P} of [Definition 4.4.2.1.1](#), where \otimes denotes the tensor product of suplattices of $\mathbf{2}$. Moreover, this makes each of $\mathcal{P}_!$, \mathcal{P}^{-1} , and \mathcal{P}_* into a symmetric monoidal functor.

Proof. [Item 1](#), *Co/Completeness*: Omitted.

[Item 2](#), *Cartesian Closedness*: See [Section 4.4.7](#).

[Item 3](#), *Powersets as Sets of Relations*: Indeed, we have

$$\begin{aligned} \text{Rel}(\text{pt}, X) &\stackrel{\text{def}}{=} \mathcal{P}(\text{pt} \times X) \\ &\cong \mathcal{P}(X) \end{aligned}$$

and

$$\begin{aligned} \text{Rel}(X, \text{pt}) &\stackrel{\text{def}}{=} \mathcal{P}(X \times \text{pt}) \\ &\cong \mathcal{P}(X), \end{aligned}$$

where we have used [Item 5](#) of [Definition 4.1.3.1.3](#).

[Item 4](#), *Interaction With Products I*: The inverse of the map in the statement is the map

$$\Phi: \mathcal{P}(X \amalg Y) \rightarrow \mathcal{P}(X) \times \mathcal{P}(Y)$$

defined by

$$\Phi(S) \stackrel{\text{def}}{=} (S_X, S_Y)$$

for each $S \in \mathcal{P}(X \amalg Y)$, where

$$\begin{aligned} S_X &\stackrel{\text{def}}{=} \{x \in X \mid (0, x) \in S\} \\ S_Y &\stackrel{\text{def}}{=} \{y \in Y \mid (1, y) \in S\}. \end{aligned}$$

The rest of the proof is omitted.

[Item 5](#), *Interaction With Products II*: Omitted.

[Item 6](#), *Interaction With Products III*: Omitted. □

weird way to highlight the similarity to external tensor products in six-functor formalisms (see also [Section 4.6.4](#)).

01JA 4.4.2 Functoriality of Powersets

01JB **Proposition 4.4.2.I.I.** Let X be a set.

01JC 1. *Functoriality I.* The assignment $X \mapsto \mathcal{P}(X)$ defines a functor

$$\mathcal{P}_! : \mathbf{Sets} \rightarrow \mathbf{Sets},$$

where

- *Action on Objects.* For each $A \in \mathbf{Obj}(\mathbf{Sets})$, we have

$$\mathcal{P}_!(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each $A, B \in \mathbf{Obj}(\mathbf{Sets})$, the action on morphisms

$$\mathcal{P}_{*|A,B} : \mathbf{Sets}(A, B) \rightarrow \mathbf{Sets}(\mathcal{P}(A), \mathcal{P}(B))$$

of $\mathcal{P}_!$ at (A, B) is the map defined by sending a map of sets $f : A \rightarrow B$ to the map

$$\mathcal{P}_!(f) : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\mathcal{P}_!(f) \stackrel{\text{def}}{=} f_!,$$

as in **Definition 4.6.I.I.I.**

01JD 2. *Functoriality II.* The assignment $X \mapsto \mathcal{P}(X)$ defines a functor

$$\mathcal{P}^{-1} : \mathbf{Sets}^{\text{op}} \rightarrow \mathbf{Sets},$$

where

- *Action on Objects.* For each $A \in \mathbf{Obj}(\mathbf{Sets})$, we have

$$\mathcal{P}^{-1}(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each $A, B \in \mathbf{Obj}(\mathbf{Sets})$, the action on

morphisms

$$\mathcal{P}_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(B), \mathcal{P}(A))$$

of \mathcal{P}^{-1} at (A, B) is the map defined by sending a map of sets $f: A \rightarrow B$ to the map

$$\mathcal{P}^{-1}(f): \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by

$$\mathcal{P}^{-1}(f) \stackrel{\text{def}}{=} f^{-1},$$

as in [Definition 4.6.2.1.1](#).

01JE 3. *Functoriality III*. The assignment $X \mapsto \mathcal{P}(X)$ defines a functor

$$\mathcal{P}_*: \text{Sets} \rightarrow \text{Sets},$$

where

- *Action on Objects*. For each $A \in \text{Obj}(\text{Sets})$, we have

$$\mathcal{P}_*(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms*. For each $A, B \in \text{Obj}(\text{Sets})$, the action on morphisms

$$\mathcal{P}_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B))$$

of \mathcal{P}_* at (A, B) is the map defined by sending a map of sets $f: A \rightarrow B$ to the map

$$\mathcal{P}_*(f): \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\mathcal{P}_*(f) \stackrel{\text{def}}{=} f_*,$$

as in [Definition 4.6.3.1.1](#).

Proof. [Item 1, Functoriality I](#): This follows from [Items 3 and 4](#) of [Definition 4.6.1.1.6](#).
[Item 2, Functoriality II](#): This follows from [Items 3 and 4](#) of [Definition 4.6.2.1.4](#).
[Item 3, Functoriality III](#): This follows from [Items 3 and 4](#) of [Definition 4.6.3.1.8](#).

□

01JF 4.4.3 Adjointness of Powersets I

01JG **Proposition 4.4.3.I.I.** We have an adjunction

$$(\mathcal{P}^{-1} \dashv \mathcal{P}^{-1, \text{op}}): \text{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{\mathcal{P}^{-1}} \\ \perp \\ \xleftarrow{\mathcal{P}^{-1, \text{op}}} \end{array} \text{Sets},$$

witnessed by a bijection

$$\underbrace{\text{Sets}^{\text{op}}(\mathcal{P}(X), Y)}_{\stackrel{\text{def}}{=} \text{Sets}(Y, \mathcal{P}(X))} \cong \text{Sets}(X, \mathcal{P}(Y)),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $Y \in \text{Obj}(\text{Sets}^{\text{op}})$.

Proof. We have

$$\begin{aligned} \text{Sets}^{\text{op}}(\mathcal{P}(A), B) &\stackrel{\text{def}}{=} \text{Sets}(B, \mathcal{P}(A)) \\ &\cong \text{Sets}(B, \text{Sets}(A, \{\mathbf{t}, \mathbf{f}\})) && \text{(by Item 2 of Definition 4.5.I.I.4)} \\ &\cong \text{Sets}(A \times B, \{\mathbf{t}, \mathbf{f}\}) && \text{(by Item 2 of Definition 4.I.3.I.3)} \\ &\cong \text{Sets}(A, \text{Sets}(B, \{\mathbf{t}, \mathbf{f}\})) && \text{(by Item 2 of Definition 4.I.3.I.3)} \\ &\cong \text{Sets}(A, \mathcal{P}(B)), && \text{(by Item 2 of Definition 4.5.I.I.4)} \end{aligned}$$

where all bijections are natural in A and B .¹⁹

□

01JH 4.4.4 Adjointness of Powersets II

01JJ **Proposition 4.4.4.I.I.** We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_!): \text{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_!} \end{array} \text{Rel},$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(X), Y) \cong \text{Sets}(X, \mathcal{P}(Y))$$

natural in $X \in \text{Obj}(\text{Sets})$ and $Y \in \text{Obj}(\text{Rel})$, where Gr is the graph functor of **Relations**, **Item 1** of **Definition 8.2.2.I.2** and $\mathcal{P}_!$ is the functor of **Relations**, **Definition 8.7.5.I.I.**

¹⁹Here we are using **Item 3** of **Definition 4.5.I.I.4**.

Proof. We have

$$\begin{aligned}
 \text{Rel}(\text{Gr}(\mathcal{A}), B) &\cong \mathcal{P}(\mathcal{A} \times B) \\
 &\cong \text{Sets}(\mathcal{A} \times B, \{\text{t}, \text{f}\}) && \text{(by Item 2 of Definition 4.5.1.1.4)} \\
 &\cong \text{Sets}(\mathcal{A}, \text{Sets}(B, \{\text{t}, \text{f}\})) && \text{(by Item 2 of Definition 4.1.3.1.3)} \\
 &\cong \text{Sets}(\mathcal{A}, \mathcal{P}(B)), && \text{(by Item 2 of Definition 4.5.1.1.4)}
 \end{aligned}$$

where all bijections are natural in \mathcal{A} , (where we are using Item 3 of Definition 4.5.1.1.4). Explicitly, this isomorphism is given by sending a relation $R: \text{Gr}(\mathcal{A}) \rightarrow B$ to the map $R^\dagger: \mathcal{A} \rightarrow \mathcal{P}(B)$ sending a to the subset $R(a)$ of B , as in Relations, Definition 8.1.1.1.1.

Naturality in B is then the statement that given a relation $R: B \rightarrow B'$, the diagram

$$\begin{array}{ccc}
 \text{Rel}(\text{Gr}(\mathcal{A}), B) & \xrightarrow{R \circ -} & \text{Rel}(\text{Gr}(\mathcal{A}), B') \\
 \vdots \downarrow & & \downarrow \vdots \\
 \text{Sets}(\mathcal{A}, \mathcal{P}(B)) & \xrightarrow{R_!} & \text{Sets}(\mathcal{A}, \mathcal{P}(B'))
 \end{array}$$

commutes, which follows from Relations, Definition 8.7.1.1.3. \square

01JK 4.4.5 Powersets as Free Cocompletions

Let X be a set.

01JL **Proposition 4.4.5.1.1.** The pair $(\mathcal{P}(X), \chi_{(-)})$ consisting of

- The powerset $(\mathcal{P}(X), \subset)$ of X of Definition 4.4.1.1.1;
- The characteristic embedding $\chi_{(-)}: X \rightarrow \mathcal{P}(X)$ of X into $\mathcal{P}(X)$ of Definition 4.5.4.1.1;

satisfies the following universal property:

(★) Given another pair (Y, f) consisting of

- A suplattice (Y, \preceq) ;
- A function $f: X \rightarrow Y$;

there exists a unique morphism of suplattices

$$(\mathcal{P}(X), \subset) \xrightarrow{\exists!} (Y, \preceq)$$

making the diagram

$$\begin{array}{ccc} & & \mathcal{P}(X) \\ & \nearrow \chi_X & \downarrow \exists! \\ X & \xrightarrow{f} & Y \end{array}$$

commute.

Proof. This is a rephrasing of **Definition 4.4.5.1.2**, which we prove below.²⁰ \square

01JM Proposition 4.4.5.1.2. We have an adjunction

$$(\mathcal{P} \dashv \overline{\mathcal{L}}): \text{Sets} \begin{array}{c} \xrightarrow{\mathcal{P}} \\ \perp \\ \xleftarrow{\overline{\mathcal{L}}} \end{array} \text{SupLat},$$

witnessed by a bijection

$$\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) \cong \text{Sets}(X, Y),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $(Y, \preceq) \in \text{Obj}(\text{SupLat})$, where:

- The category SupLat is the category of suplattices of ??.
- The map

$$\chi_X^*: \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

witnessing the above bijection is defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a morphism of suplattices $f: \mathcal{P}(X) \rightarrow Y$ to the composition

$$X \xrightarrow{\chi_X} \mathcal{P}(X) \xrightarrow{f} Y.$$

²⁰Here we only remark that the unique morphism of suplattices in the statement is given by

- The map

$$\text{Lan}_{\chi_X} : \text{Sets}(X, Y) \rightarrow \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$$

witnessing the above bijection is given by sending a function $f : X \rightarrow Y$ to its left Kan extension along χ_X ,

$$\text{Lan}_{\chi_X}(f) : \mathcal{P}(X) \rightarrow Y,$$

Moreover, invoking the bijection $\mathcal{P}(X) \cong \text{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$ of **Item 2** of **Definition 4.5.1.1.4**, $\text{Lan}_{\chi_X}(f)$ can be explicitly computed by

$$\begin{aligned} [\text{Lan}_{\chi_X}(f)](U) &= \int^{x \in X} \chi_{\mathcal{P}(X)}(\chi_x, U) \odot f(x) \\ &= \int^{x \in X} \chi_U(x) \odot f(x) \\ &= \bigvee_{x \in X} (\chi_U(x) \odot f(x)) \\ &= \left(\bigvee_{x \in U} (\chi_U(x) \odot f(x)) \right) \vee \left(\bigvee_{x \in U^c} (\chi_U(x) \odot f(x)) \right) \\ &= \left(\bigvee_{x \in U} f(x) \right) \vee \left(\bigvee_{x \in U^c} \emptyset_Y \right) \\ &= \bigvee_{x \in U} f(x) \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where:

- We have used ?? for the first equality.
- We have used **Definition 4.5.5.1.1** for the second equality.
- We have used ?? for the third equality.

the left Kan extension $\text{Lan}_{\chi_X}(f)$ of f along χ_X .

- The symbol \vee denotes the join in (Y, \preceq) .
- The symbol \odot denotes the tensor of an element of Y by a truth value as in ???. In particular, we have

$$\begin{aligned}\text{true} \odot f(x) &\stackrel{\text{def}}{=} f(x), \\ \text{false} \odot f(x) &\stackrel{\text{def}}{=} \emptyset_Y,\end{aligned}$$

where \emptyset_Y is the bottom element of (Y, \preceq) .

In particular, when $(Y, \preceq_Y) = (\mathcal{P}(B), \subset)$ for some set B , the Kan extension $\text{Lan}_{\chi_X}(f)$ is given by

$$\begin{aligned}[\text{Lan}_{\chi_X}(f)](U) &= \bigvee_{x \in U} f(x) \\ &= \bigcup_{x \in U} f(x)\end{aligned}$$

for each $U \in \mathcal{P}(X)$.

Proof. Map I: We define a map

$$\Phi_{X,Y}: \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

as in the statement, i.e. by


$$\Phi_{X,Y}(f) \stackrel{\text{def}}{=} f \circ \chi_X$$

for each $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$.

Map II: We define a map

$$\Psi_{X,Y}: \text{Sets}(X, Y) \rightarrow \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$$

as in the statement, i.e. by

$$\Psi_{X,Y}(f) \stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f),$$


for each $f \in \text{Sets}(X, Y)$.

Invertibility I: We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))}.$$

We have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](f) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}(f \circ \chi_X) \\ &\stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f \circ \chi_X) \end{aligned}$$

for each $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$. We now claim that

$$\text{Lan}_{\chi_X}(f \circ \chi_X) = f$$

for each $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$. Indeed, we have

$$\begin{aligned} [\text{Lan}_{\chi_X}(f \circ \chi_X)](U) &= \bigvee_{x \in U} f(\chi_X(x)) \\ &= f\left(\bigvee_{x \in U} \chi_X(x)\right) \\ &= f\left(\bigcup_{x \in U} \{x\}\right) \\ &= f(U) \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where we have used that f is a morphism of suplattices and hence preserves joins for the second equality. This proves our claim. Since we have shown that

$$[\Psi_{X,Y} \circ \Phi_{X,Y}](f) = f$$

for each $f \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$, it follows that $\Psi_{X,Y} \circ \Phi_{X,Y}$ must be equal to the identity map $\text{id}_{\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))}$ of $\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$.

Invertibility II: We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X,Y)}.$$

We have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](f) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\text{Lan}_{\chi_X}(f)) \\ &\stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f) \circ \chi_X \end{aligned}$$

for each $f \in \text{Sets}(X, Y)$. We now claim that

$$\text{Lan}_{\chi_X}(f) \circ \chi_X = f$$

for each $f \in \text{Sets}(X, Y)$. Indeed, we have

$$\begin{aligned} [\text{Lan}_{\chi_X}(f) \circ \chi_X](x) &= \bigvee_{y \in \{x\}} f(y) \\ &= f(x) \end{aligned}$$

for each $x \in X$. This proves our claim. Since we have shown that

$$[\Phi_{X,Y} \circ \Psi_{X,Y}](f) = f$$

for each $f \in \text{Sets}(X, Y)$, it follows that $\Phi_{X,Y} \circ \Psi_{X,Y}$ must be equal to the identity map $\text{id}_{\text{Sets}(X,Y)}$ of $\text{Sets}(X, Y)$.

Naturality for Φ , Part I: We need to show that, given a function $f: X \rightarrow X'$, the diagram

$$\begin{array}{ccc} \text{SupLat}((\mathcal{P}(X'), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}(X', Y) \\ \mathcal{P}_!(f)^* \downarrow & & \downarrow f^* \\ \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \end{array}$$

commutes. Indeed, we have

$$\begin{aligned} [\Phi_{X,Y} \circ \mathcal{P}_!(f)^*](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\mathcal{P}_!(f)^*(\xi)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\xi \circ f!) \\ &\stackrel{\text{def}}{=} (\xi \circ f!) \circ \chi_X \\ &= \xi \circ (f! \circ \chi_X) \end{aligned}$$

$$\begin{aligned}
&\stackrel{(\dagger)}{=} \xi \circ (\chi_{X'} \circ f) \\
&= (\xi \circ \chi_{X'}) \circ f \\
&\stackrel{\text{def}}{=} \Phi_{X',Y}(\xi) \circ f \\
&\stackrel{\text{def}}{=} f^*(\Phi_{X',Y}(\xi)) \\
&\stackrel{\text{def}}{=} [f^* \circ \Phi_{X',Y}](\xi),
\end{aligned}$$

for each $\xi \in \text{SupLat}((\mathcal{P}(X'), \subset), (Y, \preceq))$, where we have used **Item 1** of **Definition 4.5.4.1.3** for the fifth equality above.

Naturality for Φ , Part II: We need to show that, given a morphism of suplattices

$$g: (Y, \preceq_Y) \rightarrow (Y', \preceq_{Y'}),$$

the diagram

$$\begin{array}{ccc}
\text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\
\downarrow g! & & \downarrow g! \\
\text{SupLat}((\mathcal{P}(X), \subset), (Y', \preceq)) & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y')
\end{array}$$

commutes. Indeed, we have

$$\begin{aligned}
[\Phi_{X,Y'} \circ g!](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y'}(g!(\xi)) \\
&\stackrel{\text{def}}{=} \Phi_{X,Y'}(g \circ \xi) \\
&\stackrel{\text{def}}{=} (g \circ \xi) \circ \chi_X \\
&= g \circ (\xi \circ \chi_X) \\
&\stackrel{\text{def}}{=} g \circ (\Phi_{X,Y}(\xi)) \\
&\stackrel{\text{def}}{=} g!(\Phi_{X,Y}(\xi)) \\
&\stackrel{\text{def}}{=} [g! \circ \Phi_{X,Y}](\xi).
\end{aligned}$$

for each $\xi \in \text{SupLat}((\mathcal{P}(X), \subset), (Y, \preceq))$.

Naturality for Ψ : Since Φ is natural in each argument and Φ is a componentwise inverse to Ψ in each argument, it follows from **Categories, Item 2** of **Definition II.9.7.1.2** that Ψ is also natural in each argument. \square

01JN Warning 4.4.5.1.3. Although the assignment $X \mapsto \mathcal{P}(X)$ is called the *free cocompletion of X* , it is not an idempotent operation, i.e. we have $\mathcal{P}(\mathcal{P}(X)) \neq \mathcal{P}(X)$.

01JP 4.4.6 Powersets as Free Completions

Let X be a set.

01JQ **Proposition 4.4.6.1.1.** The pair $(\mathcal{P}(X), \chi_{(-)})$ consisting of

- The powerset of X together with reverse inclusion $\mathcal{P}(X)^{\text{op}} = (\mathcal{P}(X), \supseteq)$ of **Definition 4.4.1.1.1**;
- The characteristic embedding $\chi_{(-)}: X \rightarrow \mathcal{P}(X)$ of X into $\mathcal{P}(X)$ of **Definition 4.5.4.1.1**;

satisfies the following universal property:

(★) Given another pair (Y, f) consisting of

- An inflattice (Y, \preceq) ;
- A function $f: X \rightarrow Y$;

there exists a unique morphism of inflattices

$$(\mathcal{P}(X), \supseteq) \xrightarrow{\exists!} (Y, \preceq)$$

making the diagram

$$\begin{array}{ccc} & & \mathcal{P}(X)^{\text{op}} \\ & \nearrow \chi_X & \downarrow \exists! \\ X & \xrightarrow{f} & Y \end{array}$$

commute.

Proof. This is a rephrasing of **Definition 4.4.6.1.2**, which we prove below.²¹ \square

²¹Here we only remark that the unique morphism of inflattices in the statement is given by the right Kan extension $\text{Ran}_{\chi_X}(f)$ of f along χ_X .

01JR Proposition 4.4.6.I.2. We have an adjunction

$$(\mathcal{P} \dashv \text{忘}): \text{Sets} \begin{array}{c} \xrightarrow{\mathcal{P}} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \text{InfLat},$$

witnessed by a bijection

$$\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) \cong \text{Sets}(X, Y),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $(Y, \preceq) \in \text{Obj}(\text{InfLat})$, where:

- The category InfLat is the category of inflattices of ??.
- The map

$$\chi_X^*: \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

witnessing the above bijection is defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a morphism of inflattices $f: \mathcal{P}(X)^{\text{op}} \rightarrow Y$ to the composition

$$X \xrightarrow{\chi_X} \mathcal{P}(X)^{\text{op}} \xrightarrow{f} Y.$$

- The map

$$\text{Ran}_{\chi_X}: \text{Sets}(X, Y) \rightarrow \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$$

witnessing the above bijection is given by sending a function $f: X \rightarrow Y$ to its right Kan extension along χ_X ,

$$\text{Ran}_{\chi_X}(f): \mathcal{P}(X)^{\text{op}} \rightarrow Y,$$

Moreover, invoking the bijection $\mathcal{P}(X) \cong \text{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$ of [Item 2](#) of [Definition 4.5.1.1.4](#), $\text{Ran}_{\chi_X}(f)$ can be explicitly computed by

$$\begin{aligned}
 [\text{Ran}_{\chi_X}(f)](U) &= \int_{x \in X} \chi_{\mathcal{P}(X)^{\text{op}}}(\chi_x, U) \multimap f(x) \\
 &= \int_{x \in X} \chi_{\mathcal{P}(X)}(U, \chi_x) \multimap f(x) \\
 &= \int_{x \in X} \chi_U(x) \multimap f(x) \\
 &= \bigwedge_{x \in X} \chi_U(x) \multimap f(x) \\
 &= \left(\bigwedge_{x \in U} \chi_U(x) \multimap f(x) \right) \wedge \left(\bigwedge_{x \in U^c} \chi_U(x) \multimap f(x) \right) \\
 &= \left(\bigwedge_{x \in U} f(x) \right) \wedge \left(\bigwedge_{x \in U^c} \infty_Y \right) \\
 &= \left(\bigwedge_{x \in U} f(x) \right) \wedge \infty_Y \\
 &= \bigwedge_{x \in U} f(x)
 \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where:

- We have used ?? for the first equality.
- We have used [Definition 4.5.5.1.1](#) for the second equality.
- We have used ?? for the third equality.
- The symbol \bigwedge denotes the meet in (Y, \preceq) .
- The symbol \multimap denotes the cotensor of an element of Y by a truth value as in ?? . In particular, we have

$$\begin{aligned}
 \text{true} \multimap f(x) &\stackrel{\text{def}}{=} f(x), \\
 \text{false} \multimap f(x) &\stackrel{\text{def}}{=} \infty_Y,
 \end{aligned}$$

where ∞_Y is the top element of (Y, \preceq) .

In particular, when $(Y, \preceq_Y) = (\mathcal{P}(B), \subset)$ for some set B , the Kan extension $\text{Ran}_{\chi_X}(f)$ is given by

$$\begin{aligned} [\text{Ran}_{\chi_X}(f)](U) &= \bigwedge_{x \in U} f(x) \\ &= \bigcap_{x \in U} f(x) \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

Proof. Map I: We define a map

$$\Phi_{X,Y}: \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

as in the statement, i.e. by

$$\Phi_{X,Y}(f) \stackrel{\text{def}}{=} f \circ \chi_X$$

for each $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$.

Map II: We define a map

$$\Psi_{X,Y}: \text{Sets}(X, Y) \rightarrow \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$$

as in the statement, i.e. by

$$\Psi_{X,Y}(f) \stackrel{\text{def}}{=} \text{Ran}_{\chi_X}(f),$$

for each $f \in \text{Sets}(X, Y)$.

Invertibility I: We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))}.$$

We have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](f) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}(f \circ \chi_X) \end{aligned}$$

$$\stackrel{\text{def}}{=} \text{Ran}_{\chi_X}(f \circ \chi_X)$$

for each $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$. We now claim that

$$\text{Ran}_{\chi_X}(f \circ \chi_X) = f$$

for each $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$. Indeed, we have

$$\begin{aligned} [\text{Ran}_{\chi_X}(f \circ \chi_X)](U) &= \bigwedge_{x \in U} f(\chi_X(x)) \\ &= f\left(\bigwedge_{x \in U} \chi_X(x)\right) \\ &= f\left(\bigcup_{x \in U} \{x\}\right) \\ &= f(U) \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where we have used that f is a morphism of inflattices and hence preserves meets in $(\mathcal{P}(X), \supset)$ (i.e. joins in $(\mathcal{P}(X), \subset)$) for the second equality. This proves our claim. Since we have shown that

$$[\Psi_{X,Y} \circ \Phi_{X,Y}](f) = f$$

for each $f \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$, it follows that $\Psi_{X,Y} \circ \Phi_{X,Y}$ must be equal to the identity map $\text{id}_{\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))}$ of $\text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$.

Invertibility II: We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X,Y)}.$$

We have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](f) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\text{Ran}_{\chi_X}(f)) \\ &\stackrel{\text{def}}{=} \text{Ran}_{\chi_X}(f) \circ \chi_X \end{aligned}$$

for each $f \in \text{Sets}(X, Y)$. We now claim that

$$\text{Ran}_{\chi_X}(f) \circ \chi_X = f$$

for each $f \in \mathbf{Sets}(X, Y)$. Indeed, we have

$$\begin{aligned} [\mathrm{Ran}_{\chi_X}(f) \circ \chi_X](x) &= \bigwedge_{y \in \{x\}} f(y) \\ &= f(x) \end{aligned}$$

for each $x \in X$. This proves our claim. Since we have shown that

$$[\Phi_{X,Y} \circ \Psi_{X,Y}](f) = f$$

for each $f \in \mathbf{Sets}(X, Y)$, it follows that $\Phi_{X,Y} \circ \Psi_{X,Y}$ must be equal to the identity map $\mathrm{id}_{\mathbf{Sets}(X,Y)}$ of $\mathbf{Sets}(X, Y)$.

Naturality for Φ , Part I: We need to show that, given a function $f: X \rightarrow X'$, the diagram

$$\begin{array}{ccc} \mathrm{InfLat}((\mathcal{P}(X'), \supset), (Y, \preceq)) & \xrightarrow{\Phi_{X',Y}} & \mathbf{Sets}(X', Y) \\ \mathcal{P}_!(f)^* \downarrow & & \downarrow f^* \\ \mathrm{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \mathbf{Sets}(X, Y) \end{array}$$

commutes. Indeed, we have

$$\begin{aligned} [\Phi_{X,Y} \circ \mathcal{P}_!(f)^*](\xi) &\stackrel{\mathrm{def}}{=} \Phi_{X,Y}(\mathcal{P}_!(f)^*(\xi)) \\ &\stackrel{\mathrm{def}}{=} \Phi_{X,Y}(\xi \circ f_!) \\ &\stackrel{\mathrm{def}}{=} (\xi \circ f_!) \circ \chi_X \\ &= \xi \circ (f_! \circ \chi_X) \\ &\stackrel{(\dagger)}{=} \xi \circ (\chi_{X'} \circ f) \\ &= (\xi \circ \chi_{X'}) \circ f \\ &\stackrel{\mathrm{def}}{=} \Phi_{X',Y}(\xi) \circ f \\ &\stackrel{\mathrm{def}}{=} f^*(\Phi_{X',Y}(\xi)) \\ &\stackrel{\mathrm{def}}{=} [f^* \circ \Phi_{X',Y}](\xi), \end{aligned}$$

for each $\xi \in \mathrm{InfLat}((\mathcal{P}(X'), \supset), (Y, \preceq))$, where we have used **Item 1** of **Definition 4.5.4.1.3** for the fifth equality above.

Naturality for Φ , Part II: We need to show that, given a cocontinuous morphism of posets

$$g: (Y, \preceq_Y) \rightarrow (Y', \preceq_{Y'}),$$

the diagram

$$\begin{array}{ccc}
 \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\
 \downarrow g! & & \downarrow g! \\
 \text{InfLat}((\mathcal{P}(X), \supset), (Y', \preceq)) & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y')
 \end{array}$$

commutes. Indeed, we have

$$\begin{aligned}
 [\Phi_{X,Y'} \circ g!](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y'}(g!(\xi)) \\
 &\stackrel{\text{def}}{=} \Phi_{X,Y'}(g \circ \xi) \\
 &\stackrel{\text{def}}{=} (g \circ \xi) \circ \chi_X \\
 &= g \circ (\xi \circ \chi_X) \\
 &\stackrel{\text{def}}{=} g \circ (\Phi_{X,Y}(\xi)) \\
 &\stackrel{\text{def}}{=} g!(\Phi_{X,Y}(\xi)) \\
 &\stackrel{\text{def}}{=} [g! \circ \Phi_{X,Y}](\xi).
 \end{aligned}$$

for each $\xi \in \text{InfLat}((\mathcal{P}(X), \supset), (Y, \preceq))$.

Naturality for Ψ : Since Φ is natural in each argument and Φ is a componentwise inverse to Ψ in each argument, it follows from **Categories, Item 2 of Definition 11.9.7.1.2** that Ψ is also natural in each argument. \square

01JS Warning 4.4.6.1.3. Although the assignment $X \mapsto \mathcal{P}(X)^{\text{op}}$ is called the *free completion of X* , it is not an idempotent operation, i.e. we have $\mathcal{P}(\mathcal{P}(X)^{\text{op}})^{\text{op}} \neq \mathcal{P}(X)^{\text{op}}$.

01JT 4.4.7 The Internal Hom of a Powerset

Let X be a set and let $U, V \in \mathcal{P}(X)$.

01JU Proposition 4.4.7.1.1. The **internal Hom of $\mathcal{P}(X)$ from U to V** is the subset $[U, V]_X$ ²² of X given by

$$\begin{aligned}
 [U, V]_X &= U^c \cup V \\
 &= (U \setminus V)^c
 \end{aligned}$$

where U^c is the complement of U of **Definition 4.3.11.1.1**.

²²*Further Notation:* Also written $\mathbf{Hom}_{\mathcal{P}(X)}(U, V)$.

Proof. Proof of the Equality $U^c \cup V = (U \setminus V)^c$: We have

$$\begin{aligned}
 (U \setminus V)^c &\stackrel{\text{def}}{=} X \setminus (U \setminus V) \\
 &= (X \cap V) \cup (X \setminus U) \\
 &= V \cup (X \setminus U) \\
 &\stackrel{\text{def}}{=} V \cup U^c \\
 &= U^c \cup V,
 \end{aligned}$$

where we have used:

- 020S 1. **Item 10** of **Definition 4.3.10.1.2** for the second equality.
- 020T 2. **Item 4** of **Definition 4.3.9.1.2** for the third equality.
- 020U 3. **Item 4** of **Definition 4.3.8.1.2** for the last equality.

This finishes the proof.

Proof that $U^c \cup V$ Is Indeed the Internal Hom: This follows from **Item 2** of **Definition 4.3.9.1.2**. \square

004C **Remark 4.4.7.1.2.** Henning Makholm suggests the following heuristic intuition for the internal Hom of $\mathcal{P}(X)$ from U to V ([MSE 267365]):

- 01JV 1. Since products in $\mathcal{P}(X)$ are given by binary intersections (**Item 1** of **Definition 4.4.1.1.4**), the right adjoint $\mathbf{Hom}_{\mathcal{P}(X)}(U, -)$ of $U \cap -$ may be thought of as a function type $[U, V]$.
- 01JW 2. Under the Curry–Howard correspondence (??), the function type $[U, V]$ corresponds to implication $U \Rightarrow V$.
- 01JX 3. Implication $U \Rightarrow V$ is logically equivalent to $\neg U \vee V$.
- 01JY 4. The expression $\neg U \vee V$ then corresponds to the set $U^c \cup V$ in $\mathcal{P}(X)$.
- 01JZ 5. The set $U^c \cup V$ turns out to indeed be the internal Hom of $\mathcal{P}(X)$.

01K0 **Proposition 4.4.7.1.3.** Let X be a set.

01K1 I. *Functoriality*. The assignments $U, V, (U, V) \mapsto \mathbf{Hom}_{\mathcal{P}(X)}$ define functors

$$\begin{aligned} [U, -]_X &: (\mathcal{P}(X), \supset) && \rightarrow (\mathcal{P}(X), \subset), \\ [-, V]_X &: (\mathcal{P}(X), \subset) && \rightarrow (\mathcal{P}(X), \subset), \\ [-_1, -_2]_X &: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) && \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

In particular, the following statements hold for each $U, V, A, B \in \mathcal{P}(X)$:

01K2 (a) If $U \subset A$, then $[A, V]_X \subset [U, V]_X$.

01K3 (b) If $V \subset B$, then $[U, V]_X \subset [U, B]_X$.

01K4 (c) If $U \subset A$ and $V \subset B$, then $[A, V]_X \subset [U, B]_X$.

01K5 2. *Adjointness*. We have adjunctions

$$\begin{aligned} (U \cap - \dashv [U, -]_X) &: \mathcal{P}(X) \begin{array}{c} \xrightarrow{U \cap -} \\ \perp \\ \xleftarrow{[U, -]_X} \end{array} \mathcal{P}(X), \\ (- \cap V \dashv [V, -]_X) &: \mathcal{P}(X) \begin{array}{c} \xrightarrow{- \cap V} \\ \perp \\ \xleftarrow{[V, -]_X} \end{array} \mathcal{P}(X), \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \mathrm{Hom}_{\mathcal{P}(X)}(U \cap V, W) &\cong \mathrm{Hom}_{\mathcal{P}(X)}(U, [V, W]_X), \\ \mathrm{Hom}_{\mathcal{P}(X)}(U \cap V, W) &\cong \mathrm{Hom}_{\mathcal{P}(X)}(V, [U, W]_X). \end{aligned}$$

In particular, the following statements hold for each $U, V, W \in \mathcal{P}(X)$:

01K6 (a) The following conditions are equivalent:

01K7 i. We have $U \cap V \subset W$.

01K8 ii. We have $U \subset [V, W]_X$.

01K9 (b) The following conditions are equivalent:

01KA i. We have $U \cap V \subset W$.

01KB ii. We have $V \subset [U, W]_X$.

01KC 3. *Interaction With the Empty Set I.* We have

$$\begin{aligned} [U, \emptyset]_X &= U^c, \\ [\emptyset, V]_X &= X, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

01KD 4. *Interaction With X .* We have

$$\begin{aligned} [U, X]_X &= X, \\ [X, V]_X &= V, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

01KE 5. *Interaction With the Empty Set II.* The functor

$$D_X: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X)$$

defined by

$$\begin{aligned} D_X &\stackrel{\text{def}}{=} [-, \emptyset]_X \\ &= (-)^c \end{aligned}$$

is an involutory isomorphism of categories, making \emptyset into a dualising object for $(\mathcal{P}(X), \cap, X, [-, -]_X)$ in the sense of ???. In particular:

01KF (a) The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{D_X} & \mathcal{P}(X) \\ & \searrow \text{id}_{\mathcal{P}(X)^{\text{op}}} & \downarrow D_X \\ & & \mathcal{P}(X)^{\text{op}} \end{array}$$

commutes, i.e. we have

$$\underbrace{D_X(D_X(U))}_{\stackrel{\text{def}}{=} [[U, \emptyset]_X, \emptyset]_X} = U$$

for each $U \in \mathcal{P}(X)$.

01KG (b) The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} \xrightarrow{\cap^{\text{op}}} \mathcal{P}(X)^{\text{op}} & \\
 \text{id}_{\mathcal{P}(X)^{\text{op}}} \times D_X \nearrow & & \searrow D_X \\
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$\underbrace{D_X(U \cap D_X(V))}_{\stackrel{\text{def}}{=} [U \cap [V, \emptyset]_X, \emptyset]_X} = [U, V]_X$$

for each $U, V \in \mathcal{P}(X)$.

01KH 6. *Interaction With the Empty Set III.* Let $f: X \rightarrow Y$ be a function.

01KJ (a) *Interaction With Direct Images.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\
 D_X \downarrow & & \downarrow D_Y \\
 \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$f_!(D_X(U)) = D_Y(f_*(U))$$

for each $U \in \mathcal{P}(X)$.

01KK (b) *Interaction With Inverse Images.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1, \text{op}}} & \mathcal{P}(X)^{\text{op}} \\
 D_Y \downarrow & & \downarrow D_X \\
 \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$f^{-1}(D_Y(U)) = D_X(f^{-1}(U))$$

for each $U \in \mathcal{P}(X)$.

01KL

(c) *Interaction With Codirect Images*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ D_X \downarrow & & \downarrow D_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(D_X(U)) = D_Y(f_!(U))$$

for each $U \in \mathcal{P}(X)$.

01KM

7. *Interaction With Unions of Families of Subsets I*. The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-1, -2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\ \bigcup^{\text{op}} \times \bigcup^{\text{op}} \downarrow & \text{X} & \downarrow \bigcup \\ \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X), \end{array}$$

does not commute in general, i.e. we may have

$$\bigcup_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[\bigcup_{U \in \mathcal{U}} U, \bigcup_{V \in \mathcal{V}} V \right]_X$$

in general, where $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01KN 8. *Interaction With Unions of Families of Subsets II.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\
 \nearrow \sim & & \searrow \cup^{\text{op}} \\
 \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\
 \downarrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & & \downarrow [-, V]_X \\
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$\left[\bigcup_{U \in \mathcal{U}} U, V \right]_X = \bigcap_{U \in \mathcal{U}} [U, V]_X$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ and each $V \in \mathcal{P}(X)$.

01KP 9. *Interaction With Unions of Families of Subsets III.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \\
 \downarrow \text{id}_{\mathcal{P}(X)} \star [U, -]_X & & \downarrow [U, -]_X \\
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$\left[U, \bigcup_{V \in \mathcal{U}} V \right]_X = \bigcup_{V \in \mathcal{U}} [U, V]_X$$

for each $U \in \mathcal{P}(X)$ and each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01KQ 10. *Interaction With Intersections of Families of Subsets I.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X))^{\text{op}} \times \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{[-_1, -_2]_{\mathcal{P}(X)}} & \mathcal{P}(\mathcal{P}(X)) \\
 \downarrow \cap^{\text{op}} \times \cap^{\text{op}} & \text{X} & \downarrow \cap \\
 \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-_1, -_2]_X} & \mathcal{P}(X),
 \end{array}$$

does not commute in general, i.e. we may have

$$\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W \neq \left[\bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X$$

in general, where $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$.

01KR 11. *Interaction With Intersections of Families of Subsets II.* The diagram

$$\begin{array}{ccc} & \mathcal{P}(\mathcal{P}(X))^{\text{op}} & \\ \nearrow \sim & & \searrow \cap^{\text{op}} \\ \mathcal{P}(\mathcal{P}(X)^{\text{op}}) & & \mathcal{P}(X)^{\text{op}} \\ \downarrow \text{id}_{\mathcal{P}(X)} \star [-, V]_X & & \downarrow [-, V]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cup} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[\bigcap_{U \in \mathcal{U}} U, V \right]_X = \bigcup_{U \in \mathcal{U}} [U, V]_X$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(X))$ and each $V \in \mathcal{P}(X)$.

01KS 12. *Interaction With Intersections of Families of Subsets III.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \\ \downarrow \text{id}_{\mathcal{P}(X)} \star [U, -]_X & & \downarrow [U, -]_X \\ \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{\cap} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\left[U, \bigcap_{V \in \mathcal{V}} V \right]_X = \bigcap_{V \in \mathcal{V}} [U, V]_X$$

for each $U \in \mathcal{P}(X)$ and each $\mathcal{V} \in \mathcal{P}(\mathcal{P}(X))$.

01KT 13. *Interaction With Binary Unions.* We have equalities of sets

$$\begin{aligned}[U \cap V, W]_X &= [U, W]_X \cup [V, W]_X, \\ [U, V \cap W]_X &= [U, V]_X \cap [U, W]_X\end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

01KU 14. *Interaction With Binary Intersections.* We have equalities of sets

$$\begin{aligned}[U \cup V, W]_X &= [U, W]_X \cap [V, W]_X, \\ [U, V \cup W]_X &= [U, V]_X \cup [U, W]_X\end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

01KV 15. *Interaction With Differences.* We have equalities of sets

$$\begin{aligned}[U \setminus V, W]_X &= [U, W]_X \cup [V^c, W]_X \\ &= [U, W]_X \cup [U, V]_X, \\ [U, V \setminus W]_X &= [U, V]_X \setminus (U \cap W)\end{aligned}$$

for each $U, V, W \in \mathcal{P}(X)$.

01KW 16. *Interaction With Complements.* We have equalities of sets

$$\begin{aligned}[U^c, V]_X &= U \cup V, \\ [U, V^c]_X &= U \cap V, \\ [U, V]_X^c &= U \setminus V\end{aligned}$$

for each $U, V \in \mathcal{P}(X)$.

01KX 17. *Interaction With Characteristic Functions.* We have

$$\chi_{[U, V]_{\mathcal{P}(X)}}(x) = \max(1 - \chi_U \pmod{2}, \chi_V)$$

for each $U, V \in \mathcal{P}(X)$.

01KY 18. *Interaction With Direct Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc}\mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow [-1, -2]_X & & \downarrow [-1, -2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)\end{array}$$

commutes, i.e. we have an equality of sets

$$f_!([U, V]_X) = [f_*(U), f_!(V)]_Y,$$

natural in $U, V \in \mathcal{P}(X)$.

- 01KZ** 19. *Interaction With Inverse Images.* Let $f: X \rightarrow Y$ be a function. The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{-1, \text{op}} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \downarrow [-1, -2]_Y & & \downarrow [-1, -2]_X \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$f^{-1}([U, V]_Y) = [f^{-1}(U), f^{-1}(V)]_X,$$

natural in $U, V \in \mathcal{P}(X)$.

- 01L0** 20. *Interaction With Codirect Images.* Let $f: X \rightarrow Y$ be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow [-1, -2]_X & \wr & \downarrow [-1, -2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$[f_!(U), f_*(V)]_Y \subset f_*([U, V]_X)$$

indexed by $U, V \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** Since $\mathcal{P}(X)$ is posetal, it suffices to prove **Items 1a** to **1c**.

020W 1. *Proof of Item 1a:* We have

$$\begin{aligned} [A, V]_X &\stackrel{\text{def}}{=} A^c \cup V \\ &\subset U^c \cup V \\ &\stackrel{\text{def}}{=} [U, V]_X, \end{aligned}$$

where we have used:

020X (a) **Item 1** of **Definition 4.3.II.1.2**, which states that if $U \subset A$, then $A^c \subset U^c$.

020Y (b) **Item 1a** of **Item 1** of **Definition 4.3.II.1.2**, which states that if $A^c \subset U^c$, then $A^c \cup K \subset U^c \cup K$ for any $K \in \mathcal{P}(X)$.

020Z 2. *Proof of Item 1b:* We have

$$\begin{aligned} [U, V]_X &\stackrel{\text{def}}{=} U^c \cup V \\ &\subset U^c \cup B \\ &\stackrel{\text{def}}{=} [U, B]_X, \end{aligned}$$

where we have used **Item 1b** of **Item 1** of **Definition 4.3.II.1.2**, which states that if $V \subset B$, then $K \cup V \subset K \cup B$ for any $K \in \mathcal{P}(X)$.

0210 3. *Proof of Item 1c:* We have

$$\begin{aligned} [A, V]_X &\subset [U, V]_X \\ &\subset [U, B]_X, \end{aligned}$$

where we have used **Items 1a** and **1b**.

This finishes the proof.

Item 2, Adjointness: This is a repetition of **Item 2** of **Definition 4.3.9.1.2** and is proved there.

Item 3, Interaction With the Empty Set I: We have

$$\begin{aligned} [U, \emptyset]_X &\stackrel{\text{def}}{=} U^c \cup \emptyset \\ &= U^c, \end{aligned}$$

where we have used **Item 3** of **Definition 4.3.8.1.2**, and we have

$$[\emptyset, V]_X \stackrel{\text{def}}{=} \emptyset^c \cup V$$

$$\begin{aligned}
&\stackrel{\text{def}}{=} (X \setminus \emptyset) \cup V \\
&= X \cup V \\
&= X,
\end{aligned}$$

where we have used:

- 0211 1. **Item 12** of **Definition 4.3.10.1.2** for the first equality.
- 0212 2. **Item 5** of **Definition 4.3.8.1.2** for the last equality.

Since $\mathcal{P}(X)$ is posetal, naturality is automatic (**Categories**, **Item 4** of **Definition 11.2.7.1.2**).

Item 4, *Interaction With X* : We have

$$\begin{aligned}
[U, X]_X &\stackrel{\text{def}}{=} U^c \cup X \\
&= X,
\end{aligned}$$

where we have used **Item 5** of **Definition 4.3.8.1.2**, and we have

$$\begin{aligned}
[X, V]_X &\stackrel{\text{def}}{=} X^c \cup V \\
&\stackrel{\text{def}}{=} (X \setminus X) \cup V \\
&= \emptyset \cup V \\
&= V,
\end{aligned}$$

where we have used **Item 3** of **Definition 4.3.8.1.2** for the last equality. Since $\mathcal{P}(X)$ is posetal, naturality is automatic (**Categories**, **Item 4** of **Definition 11.2.7.1.2**).

Item 5, *Interaction With the Empty Set II*: We have

$$\begin{aligned}
D_X(D_X(U)) &\stackrel{\text{def}}{=} [[U, \emptyset]_X, \emptyset]_X \\
&= [U^c, \emptyset]_X \\
&= (U^c)^c \\
&= U,
\end{aligned}$$

where we have used:

- 0213 1. **Item 3** for the second and third equalities.
- 0214 2. **Item 3** of **Definition 4.3.11.1.2** for the fourth equality.

Since $\mathcal{P}(X)$ is posetal, naturality is automatic ([Categories](#), [Item 4](#) of [Definition 11.2.7.1.2](#)), and thus we have

$$[[-, \emptyset]_X, \emptyset]_X \cong \text{id}_{\mathcal{P}(X)}$$

This finishes the proof.

Item 6, Interaction With the Empty Set III: Since $D_X = (-)^c$, this is essentially a repetition of the corresponding results for $(-)^c$, namely [Items 5 to 7](#) of [Definition 4.3.11.1.2](#).

Item 7, Interaction With Unions of Families of Subsets I: By [Item 3](#) of [Definition 4.4.7.1.3](#), we have

$$\begin{aligned} [\mathcal{U}, \emptyset]_{\mathcal{P}(X)} &= \mathcal{U}^c, \\ [U, \emptyset]_X &= U^c. \end{aligned}$$

With this, the counterexample given in the proof of [Item 10](#) of [Definition 4.3.6.1.2](#) then applies.

Item 8, Interaction With Unions of Families of Subsets II: We have

$$\begin{aligned} \left[\bigcup_{U \in \mathcal{U}} U, V \right]_X &\stackrel{\text{def}}{=} \left(\bigcup_{U \in \mathcal{U}} U \right)^c \cup V \\ &= \left(\bigcap_{U \in \mathcal{U}} U^c \right) \cup V \\ &= \bigcap_{U \in \mathcal{U}} (U^c \cup V) \\ &\stackrel{\text{def}}{=} \bigcap_{U \in \mathcal{U}} [U, V]_X, \end{aligned}$$

where we have used:

- 0215 1. [Item 11](#) of [Definition 4.3.6.1.2](#) for the second equality.
- 0216 2. [Item 6](#) of [Definition 4.3.7.1.2](#) for the third equality.

This finishes the proof.

Item 9, Interaction With Unions of Families of Subsets III: We have

$$\bigcup_{V \in \mathcal{V}} [U, V]_X \stackrel{\text{def}}{=} \bigcup_{V \in \mathcal{V}} (U^c \cup V)$$

$$\begin{aligned}
&= U^c \cup \left(\bigcup_{V \in \mathcal{V}} V \right) \\
&\stackrel{\text{def}}{=} \left[U, \bigcup_{V \in \mathcal{V}} V \right]_X.
\end{aligned}$$

where we have used **Item 6**. This finishes the proof.

Item 10, Interaction With Intersections of Families of Subsets I: Let $X = \{0, 1\}$, let $\mathcal{U} = \{\{0, 1\}\}$, and let $\mathcal{V} = \{\{0\}, \{0, 1\}\}$. We have

$$\begin{aligned}
\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W &= \bigcap_{W \in \mathcal{P}(X)} W \\
&= \{0, 1\},
\end{aligned}$$

whereas

$$\begin{aligned}
\left[\bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X &= [\{0, 1\}, \{0\}] \\
&= \{0\},
\end{aligned}$$

Thus we have

$$\bigcap_{W \in [\mathcal{U}, \mathcal{V}]_{\mathcal{P}(X)}} W = \{0, 1\} \neq \{0\} = \left[\bigcap_{U \in \mathcal{U}} U, \bigcap_{V \in \mathcal{V}} V \right]_X.$$

This finishes the proof.

Item 11, Interaction With Intersections of Families of Subsets II: We have

$$\begin{aligned}
\left[\bigcap_{U \in \mathcal{U}} U, V \right]_X &\stackrel{\text{def}}{=} \left(\bigcap_{U \in \mathcal{U}} U \right)^c \cup V \\
&= \left(\bigcup_{U \in \mathcal{U}} U^c \right) \cup V \\
&= \bigcup_{U \in \mathcal{U}} (U^c \cup V) \\
&\stackrel{\text{def}}{=} \bigcup_{U \in \mathcal{U}} [U, V]_X,
\end{aligned}$$

where we have used:

0217 1. **Item 12** of **Definition 4.3.6.1.2** for the second equality.

0218 2. **Item 6** of **Definition 4.3.7.1.2** for the third equality.

This finishes the proof.

Item 12, Interaction With Intersections of Families of Subsets III: We have

$$\begin{aligned} \bigcap_{V \in \mathcal{U}} [U, V]_X &\stackrel{\text{def}}{=} \bigcap_{V \in \mathcal{U}} (U^c \cup V) \\ &= U^c \cup \left(\bigcap_{V \in \mathcal{U}} V \right) \\ &\stackrel{\text{def}}{=} \left[U, \bigcap_{V \in \mathcal{U}} V \right]_X. \end{aligned}$$

where we have used **Item 6**. This finishes the proof.

Item 13, Interaction With Binary Unions: We have

$$\begin{aligned} [U \cap V, W]_X &\stackrel{\text{def}}{=} (U \cap V)^c \cup W \\ &= (U^c \cup V^c) \cup W \\ &= (U^c \cup V^c) \cup (W \cup W) \\ &= (U^c \cup W) \cup (V^c \cup W) \\ &\stackrel{\text{def}}{=} [U, W]_X \cup [V, W]_X, \end{aligned}$$

where we have used:

0219 1. **Item 2** of **Definition 4.3.11.1.2** for the second equality.

021A 2. **Item 8** of **Definition 4.3.8.1.2** for the third equality.

021B 3. Several applications of **Items 2** and **4** of **Definition 4.3.8.1.2** and for the fourth equality.

For the second equality in the statement, we have

$$\begin{aligned} [U, V \cap W]_X &\stackrel{\text{def}}{=} U^c \cup (V \cap W) \\ &= (U^c \cup V) \cap (U^c \cup W) \\ &\stackrel{\text{def}}{=} [U, V]_X \cap [U, W]_X, \end{aligned}$$

where we have used **Item 6** of **Definition 4.3.8.1.2** for the second equality.

Item 14, *Interaction With Binary Intersections*: We have

$$\begin{aligned} [U \cup V, W]_X &\stackrel{\text{def}}{=} (U \cup V)^c \cup W \\ &= (U^c \cap V^c) \cup W \\ &= (U^c \cup W) \cap (V^c \cup W) \\ &\stackrel{\text{def}}{=} [U, W]_X \cap [V, W]_X, \end{aligned}$$

where we have used:

021C 1. **Item 2** of **Definition 4.3.11.1.2** for the second equality.

021D 2. **Item 6** of **Definition 4.3.8.1.2** for the third equality.

Now, for the second equality in the statement, we have

$$\begin{aligned} [U, V \cup W]_X &\stackrel{\text{def}}{=} U^c \cup (V \cup W) \\ &= (U^c \cup U^c) \cup (V \cup W) \\ &= (U^c \cup V) \cup (U^c \cup W) \\ &\stackrel{\text{def}}{=} [U, V]_X \cup [U, W]_X, \end{aligned}$$

where we have used:

021E 1. **Item 8** of **Definition 4.3.8.1.2** for the second equality.

021F 2. Several applications of **Items 2** and **4** of **Definition 4.3.8.1.2** and for the third equality.

This finishes the proof.

Item 15, *Interaction With Differences*: We have

$$\begin{aligned} [U \setminus V, W]_X &\stackrel{\text{def}}{=} (U \setminus V)^c \cup W \\ &\stackrel{\text{def}}{=} (X \setminus (U \setminus V)) \cup W \\ &= ((X \cap V) \cup (X \setminus U)) \cup W \\ &= (V \cup (X \setminus U)) \cup W \\ &\stackrel{\text{def}}{=} (V \cup U^c) \cup W \\ &= (V \cup (U^c \cup U^c)) \cup W \\ &= (U^c \cup W) \cup (U^c \cup V) \\ &\stackrel{\text{def}}{=} [U, W]_X \cup [U, V]_X, \end{aligned}$$

where we have used:

- 021G 1. Item 10 of Definition 4.3.10.1.2 for the third equality.
- 021H 2. Item 4 of Definition 4.3.9.1.2 for the fourth equality.
- 021J 3. Item 8 of Definition 4.3.8.1.2 for the sixth equality.
- 021K 4. Several applications of Items 2 and 4 of Definition 4.3.8.1.2 and for the seventh equality.

We also have

$$\begin{aligned}
 [U \setminus V, W]_X &\stackrel{\text{def}}{=} (U \setminus V)^c \cup W \\
 &\stackrel{\text{def}}{=} (X \setminus (U \setminus V)) \cup W \\
 &= ((X \cap V) \cup (X \setminus U)) \cup W \\
 &= (V \cup (X \setminus U)) \cup W \\
 &\stackrel{\text{def}}{=} (V \cup U^c) \cup W \\
 &= (V \cup U^c) \cup (W \cup W) \\
 &= (U^c \cup W) \cup (V \cup W) \\
 &= (U^c \cup W) \cup ((V^c)^c \cup W) \\
 &\stackrel{\text{def}}{=} [U, W]_X \cup [V^c, W]_X,
 \end{aligned}$$

where we have used:

- 021L 1. Item 10 of Definition 4.3.10.1.2 for the third equality.
- 021M 2. Item 4 of Definition 4.3.9.1.2 for the fourth equality.
- 021N 3. Item 8 of Definition 4.3.8.1.2 for the sixth equality.
- 021P 4. Several applications of Items 2 and 4 of Definition 4.3.8.1.2 and for the seventh equality.
- 021Q 5. Item 3 of Definition 4.3.11.1.2 for the eighth equality.

Now, for the second equality in the statement, we have

$$\begin{aligned}
 [U, V \setminus W]_X &\stackrel{\text{def}}{=} U^c \cup (V \setminus W) \\
 &= (V \setminus W) \cup U^c \\
 &= (V \cup U^c) \setminus (W \setminus U^c)
 \end{aligned}$$

$$\begin{aligned}
&\stackrel{\text{def}}{=} (V \cup U^c) \setminus (W \setminus (X \setminus U)) \\
&= (V \cup U^c) \setminus ((W \cap U) \cup (W \setminus X)) \\
&= (V \cup U^c) \setminus ((W \cap U) \cup \emptyset) \\
&= (V \cup U^c) \setminus (W \cap U) \\
&= (V \cup U^c) \setminus (U \cap W) \\
&\stackrel{\text{def}}{=} [U, V]_X \setminus (U \cap W)
\end{aligned}$$

where we have used:

- 021R 1. **Item 4** of **Definition 4.3.8.I.2** for the second equality.
- 021S 2. **Item 4** of **Definition 4.3.10.I.2** for the third equality.
- 021T 3. **Item 10** of **Definition 4.3.10.I.2** for the fifth equality.
- 021U 4. **Item 13** of **Definition 4.3.10.I.2** for the sixth equality.
- 021V 5. **Item 3** of **Definition 4.3.8.I.2** for the seventh equality.
- 021W 6. **Item 5** of **Definition 4.3.9.I.2** for the eighth equality.

This finishes the proof.

Item 16, Interaction With Complements: We have

$$\begin{aligned}
[U^c, V]_X &\stackrel{\text{def}}{=} (U^c)^c \cup V, \\
&= U \cup V,
\end{aligned}$$

where we have used **Item 3** of **Definition 4.3.11.I.2**. We also have

$$\begin{aligned}
[U, V^c]_X &\stackrel{\text{def}}{=} U^c \cup V^c \\
&= U \cap V
\end{aligned}$$

where we have used **Item 2** of **Definition 4.3.11.I.2**. Finally, we have

$$\begin{aligned}
[U, V]_X^c &= ((U \setminus V)^c)^c \\
&= U \setminus V,
\end{aligned}$$

where we have used **Item 2** of **Definition 4.3.11.I.2**.

Item 17, Interaction With Characteristic Functions: We have

$$\begin{aligned}\chi_{[U,V]_{\mathcal{P}(X)}}(x) &\stackrel{\text{def}}{=} \chi_{U^c \cup V}(x) \\ &= \max(\chi_{U^c}, \chi_V) \\ &= \max(1 - \chi_U \pmod{2}, \chi_V),\end{aligned}$$

where we have used:

- 021X 1. *Item 10* of *Definition 4.3.8.1.2* for the second equality.
- 021Y 2. *Item 4* of *Definition 4.3.11.1.2* for the third equality.

This finishes the proof.

Item 18, Interaction With Direct Images: This is a repetition of *Item 10* of *Definition 4.6.1.1.5* and is proved there.

Item 19, Interaction With Inverse Images: This is a repetition of *Item 10* of *Definition 4.6.2.1.3* and is proved there.

Item 20, Interaction With Codirect Images: This is a repetition of *Item 9* of *Definition 4.6.3.1.7* and is proved there. \square

01L1 4.4.8 Isbell Duality for Sets

Let X be a set.

01L2 **Definition 4.4.8.1.1.** The **Isbell function** of X is the map

$$I: \mathcal{P}(X) \rightarrow \text{Sets}(X, \mathcal{P}(X))$$

defined by

$$I(U) \stackrel{\text{def}}{=} \llbracket x \mapsto [U, \{x\}]_X \rrbracket$$

for each $U \in \mathcal{P}(X)$.

01L3 **Remark 4.4.8.1.2.** Recall from *Definition 4.4.1.1.2* that we may view the power-set $\mathcal{P}(X)$ of a set X as the decategorification of the category of presheaves $\text{PSh}(C)$ of a category C . Building upon this analogy, we want to mimic the definition of the Isbell Spec functor, which is given on objects by

$$\text{Spec}(\mathcal{F}) \stackrel{\text{def}}{=} \text{Nat}(\mathcal{F}, h_{(-)})$$

for each $\mathcal{F} \in \text{Obj}(\text{PSh}(C))$. To this end, we could define

$$I(U) \stackrel{\text{def}}{=} [U, \chi_{(-)}]_X,$$

replacing:

- The Yoneda embedding $X \mapsto h_X$ of C into $\mathbf{PSh}(C)$ with the characteristic embedding $x \mapsto \chi_x$ of X into $\mathcal{P}(X)$ of [Definition 4.5.4.1.1](#).
- The internal Hom Nat of $\mathbf{PSh}(C)$ with the internal Hom $[-, -]_X$ of $\mathcal{P}(X)$ of [Definition 4.4.7.1.1](#).

However, since $[U, \chi_x]_X$ is a subset of U instead of a truth value, we get a function

$$! : \mathcal{P}(X) \rightarrow \mathbf{Sets}(X, \mathcal{P}(X))$$

instead of a function

$$! : \mathcal{P}(X) \rightarrow \mathcal{P}(X).$$

This makes some of the properties involving $!$ a bit more cumbersome to state, although we still have an analogue of Isbell duality in that $! \circ !$ evaluates to $\text{id}_{\mathcal{P}(X)}$ in the sense of [Definition 4.4.8.1.3](#).

01L4 Proposition 4.4.8.1.3. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{!} & \mathbf{Sets}(X, \mathcal{P}(X)) \\ & \searrow \Delta_{\Delta_{\text{id}_{\mathcal{P}(X)}}} & \downarrow ! \\ & & \mathbf{Sets}(X, \mathbf{Sets}(X, \mathcal{P}(X))) \end{array}$$

commutes, i.e. we have

$$!(!(U)) = \llbracket x \mapsto \llbracket y \mapsto U \rrbracket \rrbracket$$

for each $U \in \mathcal{P}(X)$.

Proof. We have

$$\begin{aligned} !(!(U)) &\stackrel{\text{def}}{=} !(\llbracket x \mapsto U^c \cup \{x\} \rrbracket) \\ &\stackrel{\text{def}}{=} \llbracket x \mapsto !(U^c \cup \{x\}) \rrbracket \\ &\stackrel{\text{def}}{=} \llbracket x \mapsto \llbracket y \mapsto (U^c \cup \{x\})^c \cup \{x\} \rrbracket \rrbracket \\ &= \llbracket x \mapsto \llbracket y \mapsto (U \cap (X \setminus \{x\})) \cup \{x\} \rrbracket \rrbracket \\ &= \llbracket x \mapsto \llbracket y \mapsto (U \setminus \{x\}) \cup \{x\} \rrbracket \rrbracket \\ &= \llbracket x \mapsto \llbracket y \mapsto U \rrbracket \rrbracket, \end{aligned}$$

where we have used [Item 2](#) of [Definition 4.3.11.1.2](#) for the fourth equality above.

□

01L5 4.5 Characteristic Functions

005X 4.5.1 The Characteristic Function of a Subset

Let X be a set and let $U \in \mathcal{P}(X)$.

005Z **Definition 4.5.1.1.1.** The **characteristic function of U** ²³ is the function $\chi_U: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$ ²⁴ defined by

$$\chi_U(x) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in U, \\ \text{false} & \text{if } x \notin U \end{cases}$$

for each $x \in X$.

01L6 **Remark 4.5.1.1.2.** Under the analogy that $\{\mathbf{t}, \mathbf{f}\}$ should be the (-1) -categorical analogue of **Sets**, we may view a function

$$f: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

as a decategorification of presheaves and copresheaves

$$\begin{aligned} \mathcal{F}: C^{\text{op}} &\rightarrow \mathbf{Sets}, \\ F: C &\rightarrow \mathbf{Sets}. \end{aligned}$$

The characteristic functions χ_U of the subsets of X are then the primordial examples of such functions (and, in fact, all of them).

01L7 **Notation 4.5.1.1.3.** We will often employ the bijection $\{\mathbf{t}, \mathbf{f}\} \cong \{0, 1\}$ to make use of the arithmetical operations defined on $\{0, 1\}$ when discussing characteristic functions.

Examples of this include **Items 4 to 11** of **Definition 4.5.1.1.4** below.

0069 **Proposition 4.5.1.1.4.** Let X be a set.

01L8 1. *Functionality.* The assignment $U \mapsto \chi_U$ defines a function

$$\chi_{(-)}: \mathcal{P}(X) \rightarrow \mathbf{Sets}(X, \{\mathbf{t}, \mathbf{f}\}).$$

01L9 2. *Bijectivity.* The function $\chi_{(-)}$ from **Item 1** is bijective.

²³*Further Terminology:* Also called the **indicator function of U** .

²⁴*Further Notation:* Also written $\chi_X(U, -)$ or $\chi_X(-, U)$.

01LA 3. *Naturality.* The collection

$$\{\chi_{(-)}: \mathcal{P}(X) \rightarrow \mathbf{Sets}(X, \{\mathbf{t}, \mathbf{f}\})\}_{X \in \mathbf{Obj}(\mathbf{Sets})}$$

defines a natural isomorphism between \mathcal{P}^{-1} and $\mathbf{Sets}(-, \{\mathbf{t}, \mathbf{f}\})$. In particular, given a function $f: X \rightarrow Y$, the diagram

$$\begin{array}{ccc} \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \\ \chi_{(-)} \downarrow & & \downarrow \chi_{(-)} \\ \mathbf{Sets}(Y, \{\mathbf{t}, \mathbf{f}\}) & \xrightarrow{f^*} & \mathbf{Sets}(X, \{\mathbf{t}, \mathbf{f}\}) \end{array}$$

commutes, i.e. we have

$$\chi_V \circ f = \chi_{f^{-1}(V)}$$

for each $V \in \mathcal{P}(Y)$.

006B 4. *Interaction With Unions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each $U, V \in \mathcal{P}(X)$.

006C 5. *Interaction With Unions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each $U, V \in \mathcal{P}(X)$.

006D 6. *Interaction With Intersections I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each $U, V \in \mathcal{P}(X)$.

006E 7. *Interaction With Intersections II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each $U, V \in \mathcal{P}(X)$.

006F 8. *Interaction With Differences.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each $U, V \in \mathcal{P}(X)$.

006G 9. *Interaction With Complements.* We have

$$\chi_{U^c} \equiv 1 - \chi_U \pmod{2}$$

for each $U \in \mathcal{P}(X)$.

006H 10. *Interaction With Symmetric Differences.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each $U, V \in \mathcal{P}(X)$.

01LB II. *Interaction With Internal Homs.* We have

$$\chi_{[U, V]_{\mathcal{P}(X)}} = \max(1 - \chi_U \pmod{2}, \chi_V)$$

for each $U, V \in \mathcal{P}(X)$.

Proof. **Item 1, Functionality:** There is nothing to prove.

Item 2, Bijectivity: We proceed in three steps:

021Z I. *The Inverse of $\chi_{(-)}$.* The inverse of $\chi_{(-)}$ is the map

$$\Phi: \text{Sets}(X, \{\text{t}, \text{f}\}) \xrightarrow{\sim} \mathcal{P}(X),$$

defined by

$$\begin{aligned} \Phi(f) &\stackrel{\text{def}}{=} U_f \\ &\stackrel{\text{def}}{=} f^{-1}(\text{true}) \\ &\stackrel{\text{def}}{=} \{x \in X \mid f(x) = \text{true}\} \end{aligned}$$

for each $f \in \text{Sets}(X, \{\text{t}, \text{f}\})$.

0220 2. *Invertibility I.* We have

$$\begin{aligned}
 [\Phi \circ \chi_{(-)}](U) &\stackrel{\text{def}}{=} \Phi(\chi_U) \\
 &\stackrel{\text{def}}{=} \chi_U^{-1}(\text{true}) \\
 &\stackrel{\text{def}}{=} \{x \in X \mid \chi_U(x) = \text{true}\} \\
 &\stackrel{\text{def}}{=} \{x \in X \mid x \in U\} \\
 &= U \\
 &\stackrel{\text{def}}{=} [\text{id}_{\mathcal{P}(X)}](U)
 \end{aligned}$$

for each $U \in \mathcal{P}(X)$. Thus, we have

$$\Phi \circ \chi_{(-)} = \text{id}_{\mathcal{P}(X)}.$$

0221 3. *Invertibility II.* We have

$$\begin{aligned}
 [\chi_{(-)} \circ \Phi](U) &\stackrel{\text{def}}{=} \chi_{\Phi(f)} \\
 &\stackrel{\text{def}}{=} \chi_{f^{-1}(\text{true})} \\
 &\stackrel{\text{def}}{=} \llbracket x \mapsto \begin{cases} \text{true} & \text{if } x \in f^{-1}(\text{true}) \\ \text{false} & \text{otherwise} \end{cases} \rrbracket \\
 &= \llbracket x \mapsto f(x) \rrbracket \\
 &= f \\
 &\stackrel{\text{def}}{=} [\text{id}_{\text{Sets}(X, \{\text{t}, \text{f}\})}](f)
 \end{aligned}$$

for each $f \in \text{Sets}(X, \{\text{t}, \text{f}\})$. Thus, we have

$$\chi_{(-)} \circ \Phi = \text{id}_{\text{Sets}(X, \{\text{t}, \text{f}\})}.$$

This finishes the proof.

Item 3, Naturality: We proceed in two steps:

0222 1. *Naturality of $\chi_{(-)}$.* We have

$$\begin{aligned}
 [\chi_V \circ f](v) &\stackrel{\text{def}}{=} \chi_V(f(v)) \\
 &= \begin{cases} \text{true} & \text{if } f(v) \in V, \\ \text{false} & \text{otherwise} \end{cases}
 \end{aligned}$$

$$\begin{aligned}
&= \begin{cases} \text{true} & \text{if } v \in f^{-1}(V), \\ \text{false} & \text{otherwise} \end{cases} \\
&\stackrel{\text{def}}{=} \chi_{f^{-1}(V)}(v)
\end{aligned}$$

for each $v \in V$.

- 0223 2. *Naturality of Φ* . Since $\chi_{(-)}$ is natural and a componentwise inverse to Φ , it follows from [Categories](#), [Item 2](#) of [Definition 11.9.7.1.2](#) that Φ is also natural in each argument.

This finishes the proof.

Item 4, Interaction With Unions I: This is a repetition of [Item 10](#) of [Definition 4.3.8.1.2](#) and is proved there.

Item 5, Interaction With Unions II: This is a repetition of [Item 11](#) of [Definition 4.3.8.1.2](#) and is proved there.

Item 6, Interaction With Intersections I: This is a repetition of [Item 10](#) of [Definition 4.3.9.1.2](#) and is proved there.

Item 7, Interaction With Intersections II: This is a repetition of [Item 11](#) of [Definition 4.3.9.1.2](#) and is proved there.

Item 8, Interaction With Differences: This is a repetition of [Item 16](#) of [Definition 4.3.10.1.2](#) and is proved there.

Item 9, Interaction With Complements: This is a repetition of [Item 4](#) of [Definition 4.3.11.1.2](#) and is proved there.

Item 10, Interaction With Symmetric Differences: This is a repetition of [Item 15](#) of [Definition 4.3.12.1.2](#) and is proved there.

Item 11, Interaction With Internal Homs: This is a repetition of [Item 17](#) of [Definition 4.4.7.1.3](#) and is proved there. \square

- 0224 **Remark 4.5.1.1.5.** The bijection

$$\mathcal{P}(X) \cong \text{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$$

of [Item 2](#) of [Definition 4.5.1.1.4](#), which

- Takes a subset $U \rightarrow X$ of X and *straightens* it to a function $\chi_U: X \rightarrow \{\text{true}, \text{false}\}$;
- Takes a function $f: X \rightarrow \{\text{true}, \text{false}\}$ and *unstraightens* it to a subset $f^{-1}(\text{true}) \rightarrow X$ of X ;

may be viewed as the (-1) -categorical version of the 0-categorical un/straightening isomorphism between indexed and fibred sets

$$\underbrace{\text{FibSets}_X}_{\stackrel{\text{def}}{=} \text{Sets}/X} \cong \underbrace{\text{ISets}_X}_{\stackrel{\text{def}}{=} \text{Fun}(X_{\text{disc}}, \text{Sets})}$$

of Un/Straightening for Indexed and Fibred Sets, ???. Here we view:

- Subsets $U \rightarrow X$ as being analogous to X -fibred sets $\phi_X: A \rightarrow X$.
- Functions $f: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$ as being analogous to X -indexed sets $A: X_{\text{disc}} \rightarrow \text{Sets}$.

01LC 4.5.2 The Characteristic Function of a Point

Let X be a set and let $x \in X$.

0060 **Definition 4.5.2.1.1.** The **characteristic function of x** is the function²⁵

$$\chi_x: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

defined by

$$\chi_x \stackrel{\text{def}}{=} \chi_{\{x\}},$$

i.e. by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \mathbf{true} & \text{if } x = y, \\ \mathbf{false} & \text{if } x \neq y \end{cases}$$

for each $y \in X$.

01LD **Remark 4.5.2.1.2.** Expanding upon **Definition 4.5.1.1.2**, we may think of the characteristic function

$$\chi_x: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

of an *element* x of X as a decategorification of the representable presheaf and of the representable copresheaf

$$h_X: C^{\text{op}} \rightarrow \text{Sets},$$

$$h^X: C \rightarrow \text{Sets}$$

associated of an *object* X of a category C .

²⁵*Further Notation:* Also written χ^x , $\chi_X(x, -)$, or $\chi_X(-, x)$.

01LE 4.5.3 The Characteristic Relation of a Set

Let X be a set.

0061 **Definition 4.5.3.1.1.** The **characteristic relation on X** ²⁶ is the relation²⁷

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

on X defined by²⁸

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \mathbf{true} & \text{if } x = y, \\ \mathbf{false} & \text{if } x \neq y \end{cases}$$

for each $x, y \in X$.

01LF **Remark 4.5.3.1.2.** Expanding upon **Definitions 4.5.1.1.2** and **4.5.2.1.2**, we may view the characteristic relation

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

of X as a decategorification of the Hom profunctor

$$\text{Hom}_C(-_1, -_2): C^{\text{op}} \times C \rightarrow \text{Sets}$$

of a category C .

01LG **Proposition 4.5.3.1.3.** Let $f: X \rightarrow Y$ be a function.

006A 1. *The Inclusion of Characteristic Relations Associated to a Function.* Let $f: A \rightarrow B$ be a function. We have an inclusion²⁹

$$\chi_B \circ (f \times f) \subset \chi_A,$$

$$\begin{array}{ccc} A \times A & \xrightarrow{f \times f} & B \times B \\ & \searrow \chi_A \quad \supset \quad \swarrow \chi_B & \\ & \{\mathbf{t}, \mathbf{f}\}. & \end{array}$$

Proof. **Item 1, The Inclusion of Characteristic Relations Associated to a Function:** The inclusion $\chi_B(f(a), f(b)) \subset \chi_A(a, b)$ is equivalent to the statement “if $a = b$, then $f(a) = f(b)$ ”, which is true. \square

²⁶*Further Terminology:* Also called the **identity relation on X** .

²⁷*Further Notation:* Also written χ_X^{-1} , or \sim_{id} in the context of relations.

²⁸Under the bijection $\text{Sets}(X \times X, \{\mathbf{t}, \mathbf{f}\}) \cong \mathcal{P}(X \times X)$ of **Item 2** of **Definition 4.5.1.1.4**, the relation χ_X corresponds to the diagonal $\Delta_X \subset X \times X$ of X .

²⁹*Note:* This is the 0-categorical version of **Categories, Definition 11.5.4.1.1**.

01LH 4.5.4 The Characteristic Embedding of a Set

Let X be a set.

0062 **Definition 4.5.4.1.1.** The **characteristic embedding**³⁰ of X into $\mathcal{P}(X)$ is the function

$$\chi(-) : X \rightarrow \mathcal{P}(X)$$

defined by³¹

$$\begin{aligned} \chi(-)(x) &\stackrel{\text{def}}{=} \chi_x \\ &= \{x\} \end{aligned}$$

for each $x \in X$.

01LJ **Remark 4.5.4.1.2.** Expanding upon **Definitions 4.5.1.1.2**, **4.5.2.1.2** and **4.5.3.1.2**, we may view the characteristic embedding

$$\chi(-) : X \rightarrow \mathcal{P}(X)$$

of X into $\mathcal{P}(X)$ as a decategorification of the Yoneda embedding

$$\mathfrak{Y} : C^{\text{op}} \rightarrow \text{PSh}(C)$$

of a category C into $\text{PSh}(C)$.

01LK **Proposition 4.5.4.1.3.** Let $f : X \rightarrow Y$ be a map of sets.

01LL 1. *Interaction With Functions.* We have

$$f! \circ \chi_X = \chi_Y \circ f$$

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \chi_X \downarrow & & \downarrow \chi_Y \\ \mathcal{P}(X) & \xrightarrow{f!} & \mathcal{P}(Y). \end{array}$$

³⁰The name “characteristic *embedding*” is justified by **Definition 4.5.5.1.2**, which gives an analogue of fully faithfulness for $\chi(-)$.

³¹Here we are identifying $\mathcal{P}(X)$ with $\text{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$ as per **Item 2** of **Definition 4.5.1.1.4**.

Proof. **Item 1, Interaction With Functions:** Indeed, we have

$$\begin{aligned}
 [f \circ \chi_X](x) &\stackrel{\text{def}}{=} f(\chi_X(x)) \\
 &\stackrel{\text{def}}{=} f(\{x\}) \\
 &= \{f(x)\} \\
 &\stackrel{\text{def}}{=} \chi_{X'}(f(x)) \\
 &\stackrel{\text{def}}{=} [\chi_{X'} \circ f](x),
 \end{aligned}$$

for each $x \in X$, showing the desired equality. \square

006K 4.5.5 The Yoneda Lemma for Sets

Let X be a set and let $U \subset X$ be a subset of X .

006L Proposition 4.5.5.1.1. We have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$$

for each $x \in X$, giving an equality of functions

$$\chi_{\mathcal{P}(X)}(\chi_{(-)}, \chi_U) = \chi_U,$$

where

$$\chi_{\mathcal{P}(X)}(U, V) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } U \subset V, \\ \text{false} & \text{otherwise.} \end{cases}$$

Proof. We have

$$\begin{aligned}
 \chi_{\mathcal{P}(X)}(\chi_x, \chi_U) &\stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } \{x\} \subset U, \\ \text{false} & \text{otherwise} \end{cases} \\
 &= \begin{cases} \text{true} & \text{if } x \in U \\ \text{false} & \text{otherwise} \end{cases} \\
 &\stackrel{\text{def}}{=} \chi_U(x).
 \end{aligned}$$

This finishes the proof. \square

006M Corollary 4.5.5.1.2. The characteristic embedding is fully faithful, i.e., we have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_y) \cong \chi_X(x, y)$$

for each $x, y \in X$.

Proof. We have

$$\begin{aligned} \chi_{\mathcal{P}(X)}(\chi_x, \chi_y) &= \chi_y(x) \\ &\stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in \{y\} \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if } x = y \\ \text{false} & \text{otherwise} \end{cases} \\ &\stackrel{\text{def}}{=} \chi_X(x, y). \end{aligned}$$

where we have used **Definition 4.5.5.1.1** for the first equality. \square

01LM 4.6 The Adjoint Triple $f_! \dashv f^{-1} \dashv f_*$

007F 4.6.1 Direct Images

Let $f: X \rightarrow Y$ be a function.

007G Definition 4.6.1.1.1. The **direct image function associated to f** is the function³²

$$f_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by³³

$$\begin{aligned} f_!(U) &\stackrel{\text{def}}{=} \left\{ y \in Y \left| \begin{array}{l} \text{there exists some } x \in U \\ \text{such that } y = f(x) \end{array} \right. \right\} \\ &= \{ f(x) \in Y \mid x \in U \} \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

³²*Further Notation:* Also written simply $f: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$.

³³*Further Terminology:* The set $f(U)$ is called the **direct image of U by f** .

007H Notation 4.6.1.1.2. Sometimes one finds the notation

$$\exists_f: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

for $f_!$. This notation comes from the fact that the following statements are equivalent, where $y \in Y$ and $U \in \mathcal{P}(X)$:

- We have $y \in \exists_f(U)$.
- There exists some $x \in U$ such that $f(x) = y$.

We will not make use of this notation elsewhere in Clowder.

0225 Warning 4.6.1.1.3. Notation for direct images between powersets is tricky:

0226 1. Direct images for powersets and presheaves are both adjoint to their corresponding inverse image functors. However, the direct image functor for powersets is a *left* adjoint, while the direct image functor for presheaves is a *right* adjoint:

0227 (a) *Powersets.* Given a function $f: X \rightarrow Y$, we have an inverse image functor

$$f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X).$$

The *left* adjoint of this functor is the usual direct image, defined above in **Definition 4.6.1.1.1**.

0228 (b) *Presheaves.* Given a morphism of topological spaces $f: X \rightarrow Y$, we have an inverse image functor

$$f^{-1}: \mathbf{PSh}(Y) \rightarrow \mathbf{PSh}(X).$$

The *right* adjoint of this functor is the direct image functor of presheaves, defined in ??.

0229 2. The presheaf direct image functor is denoted f_* , but the direct image functor for powersets is denoted $f_!$ (as it's a left adjoint).

022A 3. Adding to the confusion, it's somewhat common for $f_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ to be denoted f_* .

We chose to write $f_!$ for the direct image to keep the notation aligned with the following similar adjoint situations:

SITUATION	ADJOINT STRING
Functoriality of Powersets	$(f_! \dashv f^{-1} \dashv f_*) : \mathcal{P}(X) \xrightarrow{\cong} \mathcal{P}(Y)$
Functoriality of Presheaf Categories	$(f_! \dashv f^{-1} \dashv f_*) : \mathbf{PSh}(X) \xrightarrow{\cong} \mathbf{PSh}(Y)$
Base Change	$(f_! \dashv f^* \dashv f_*) : \mathcal{C}_X \xrightarrow{\cong} \mathcal{C}_Y$
Kan Extensions	$(F_! \dashv F^* \dashv F_*) : \mathbf{Fun}(C, \mathcal{E}) \xrightarrow{\cong} \mathbf{Fun}(D, \mathcal{E})$

007J Remark 4.6.1.1.4. Identifying $\mathcal{P}(X)$ with $\mathbf{Sets}(X, \{\mathbf{t}, \mathbf{f}\})$ via [Item 2 of Definition 4.5.1.1.4](#), we see that the direct image function associated to f is equivalently the function

$$f_! : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by

$$\begin{aligned}
 f_!(\chi_U) &\stackrel{\text{def}}{=} \mathbf{Lan}_f(\chi_U) \\
 &= \text{colim} \left(\left(f \times \underline{(-)} \right) \xrightarrow{\text{pr}} \mathcal{A} \xrightarrow{\chi_U} \{\mathbf{t}, \mathbf{f}\} \right) \\
 &= \text{colim}_{\substack{x \in X \\ f(x) = \mathbf{f}}} (\chi_U(x)) \\
 &= \bigvee_{\substack{x \in X \\ f(x) = \mathbf{f}}} (\chi_U(x)),
 \end{aligned}$$

where we have used ?? for the second equality. In other words, we have

$$\begin{aligned}
 [f_!(\chi_U)](y) &= \bigvee_{\substack{x \in X \\ f(x) = y}} (\chi_U(x)) \\
 &= \begin{cases} \text{true} & \text{if there exists some } x \in X \text{ such} \\ & \text{that } f(x) = y \text{ and } x \in U, \\ \text{false} & \text{otherwise} \end{cases} \\
 &= \begin{cases} \text{true} & \text{if there exists some } x \in U \\ & \text{such that } f(x) = y, \\ \text{false} & \text{otherwise} \end{cases}
 \end{aligned}$$

for each $y \in Y$.

007K Proposition 4.6.I.1.5. Let $f: X \rightarrow Y$ be a function.

007L 1. *Functoriality.* The assignment $U \mapsto f_!(U)$ defines a functor

$$f_!: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset).$$

In particular, for each $U, V \in \mathcal{P}(X)$, the following condition is satisfied:

(★) If $U \subset V$, then $f_!(U) \subset f_!(V)$.

007M 2. *Triple Adjointness.* We have a triple adjunction

$$(f_! \dashv f^{-1} \dashv f_*): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{f_!} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_*} \end{array} \mathcal{P}(Y),$$

witnessed by:

01LN (a) Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\rightarrow f^{-1} \circ f_!, & \text{id}_{\mathcal{P}(Y)} &\rightarrow f_* \circ f^{-1}, \\ f_! \circ f^{-1} &\rightarrow \text{id}_{\mathcal{P}(Y)}, & f^{-1} \circ f_* &\rightarrow \text{id}_{\mathcal{P}(X)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset f^{-1}(f_!(U)), & V &\subset f_*(f^{-1}(V)), \\ f_!(f^{-1}(V)) &\subset V, & f^{-1}(f_*(U)) &\subset U \end{aligned}$$

indexed by $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$.

01LP (b) Bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(Y)}(f_!(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(X)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f_*(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$ and (respectively) $V \in \mathcal{P}(X)$ and $U \in \mathcal{P}(Y)$. In particular:

01LQ i. The following conditions are equivalent:

01LR

A. We have $f_!(U) \subset V$.

01LS

B. We have $U \subset f^{-1}(V)$.

01LT

ii. The following conditions are equivalent:

01LU

A. We have $f^{-1}(U) \subset V$.

01LV

B. We have $U \subset f_*(V)$.

01LW

3. *Interaction With Unions of Families of Subsets.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_!)_!} & \mathcal{P}(\mathcal{P}(Y)) \\
 \cup \downarrow & & \downarrow \cup \\
 \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_!(U) = \bigcup_{V \in f_!(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f_!)_!(\mathcal{U})$.

01LX

4. *Interaction With Intersections of Families of Subsets.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_!)_!} & \mathcal{P}(\mathcal{P}(Y)) \\
 \cap \downarrow & & \downarrow \cap \\
 \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_!(U) = \bigcap_{V \in f_!(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_!(\mathcal{U}) \stackrel{\text{def}}{=} (f_!)_!(\mathcal{U})$.

01LY 5. *Interaction With Binary Unions.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \downarrow \cup & & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U \cup V) = f_!(U) \cup f_!(V)$$

for each $U, V \in \mathcal{P}(X)$.

01LZ 6. *Interaction With Binary Intersections.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_! \times f_!} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \downarrow \cap & \subset & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U \cap V) \subset f_!(U) \cap f_!(V)$$

indexed by $U, V \in \mathcal{P}(X)$.

01M0 7. *Interaction With Differences.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \downarrow \setminus & \supset & \downarrow \setminus \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \setminus f_!(V) \subset f_!(U \setminus V)$$

indexed by $U, V \in \mathcal{P}(X)$.

01M1 8. *Interaction With Complements.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(U^c) = f_*(U)^c$$

for each $U \in \mathcal{P}(X)$.

01M2 9. *Interaction With Symmetric Differences.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \wr & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

with components

$$f_!(U) \Delta f_!(V) \subset f_!(U \Delta V)$$

indexed by $U, V \in \mathcal{P}(X)$.

01M3 10. *Interaction With Internal Homs of Powersets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_!} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ [-1, -2]_X \downarrow & & \downarrow [-1, -2]_Y \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have an equality of sets

$$f_!([U, V]_X) = [f_*(U), f_!(V)]_Y,$$

natural in $U, V \in \mathcal{P}(X)$.

007N II. *Preservation of Colimits.* We have an equality of sets

$$f_! \left(\bigcup_{i \in I} U_i \right) = \bigcup_{i \in I} f_!(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f_!(U) \cup f_!(V) &= f_!(U \cup V), \\ f_!(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

007P 12. *Oplax Preservation of Limits.* We have an inclusion of sets

$$f_! \left(\bigcap_{i \in I} U_i \right) \subset \bigcap_{i \in I} f_!(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} f_!(U \cap V) &\subset f_!(U) \cap f_!(V), \\ f_!(X) &\subset Y, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

007Q 13. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$(f, f_!^\otimes, f_{!1}^\otimes) : (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{!1U,V}^\otimes : f_!(U) \cup f_!(V) &\xrightarrow{=} f_!(U \cup V), \\ f_{!1}^\otimes : \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

007R 14. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct

image function of **Item 1** has a symmetric oplax monoidal structure

$$(f_!, f_!^\otimes, f_{!1}^\otimes): (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with inclusions

$$\begin{aligned} f_{!1U,V}^\otimes: f_!(U \cap V) &\rightarrow f_!(U) \cap f_!(V), \\ f_{!1}^\otimes: f_!(X) &\rightarrow Y, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

- 007S** 15. *Interaction With Coproducts.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be maps of sets. We have

$$(f \amalg g)_!(U \amalg V) = f_!(U) \amalg g_!(V)$$

for each $U \in \mathcal{P}(X)$ and each $V \in \mathcal{P}(Y)$.

- 007T** 16. *Interaction With Products.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be maps of sets. We have

$$(f \boxtimes_{X \times Y} g)_!(U \boxtimes_{X \times Y} V) = f_!(U) \boxtimes_{X' \times Y'} g_!(V)$$

for each $U \in \mathcal{P}(X)$ and each $V \in \mathcal{P}(Y)$.

- 007U** 17. *Relation to Codirect Images.* We have

$$\begin{aligned} f_!(U) &= f_*(U^c)^c \\ &\stackrel{\text{def}}{=} Y \setminus f_*(X \setminus U) \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Triple Adjointness: This follows from **Definition 4.6.1.1.4**, **Definition 4.6.2.1.2**, **Definition 4.6.3.1.4**, and Kan Extensions, ?? of ??.

Item 3, Interaction With Unions of Families of Subsets: We have

$$\bigcup_{V \in f_!(\mathcal{U})} V = \bigcup_{V \in \{f_!(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V$$

$$= \bigcup_{U \in \mathcal{U}} f_!(U).$$

This finishes the proof.

Item 4, Interaction With Intersections of Families of Subsets: We have

$$\begin{aligned} \bigcap_{V \in f_!(\mathcal{U})} V &= \bigcap_{V \in \{f_!(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcap_{U \in \mathcal{U}} f_!(U). \end{aligned}$$

This finishes the proof.

Item 5, Interaction With Binary Unions: See [Pro25p].

Item 6, Interaction With Binary Intersections: See [Pro25n].

Item 7, Interaction With Differences: See [Pro25o].

Item 8, Interaction With Complements: Applying *Item 17* to $X \setminus U$, we have

$$\begin{aligned} f_!(U^c) &= f_!(X \setminus U) \\ &= Y \setminus f_*(X \setminus (X \setminus U)) \\ &= Y \setminus f_*(U) \\ &= f_*(U)^c. \end{aligned}$$

This finishes the proof.

Item 9, Interaction With Symmetric Differences: We have

$$\begin{aligned} f_!(U) \triangle f_!(V) &= (f_!(U) \cup f_!(V)) \setminus (f_!(U) \cap f_!(V)) \\ &\subset (f_!(U) \cup f_!(V)) \setminus (f_!(U \cap V)) \\ &= (f_!(U \cup V)) \setminus (f_!(U \cap V)) \\ &\subset f_!((U \cup V) \setminus (U \cap V)) \\ &= f_!(U \triangle V), \end{aligned}$$

where we have used:

- 022C 1. *Item 2* of *Definition 4.3.12.1.2* for the first equality.
- 022D 2. *Item 6* of this proposition together with *Item 1* of *Definition 4.3.10.1.2* for the first inclusion.

- 022E 3. **Item 5** for the second equality.
- 022F 4. **Item 7** for the second inclusion.
- 022G 5. **Item 2** of **Definition 4.3.12.1.2** for the third equality.

Since $\mathcal{P}(Y)$ is posetal, naturality is automatic (**Categories**, **Item 4** of **Definition 11.2.7.1.2**). This finishes the proof.

Item 10, Interaction With Internal Homs of Powersets: We have

$$\begin{aligned}
 f_!([U, V]_X) &\stackrel{\text{def}}{=} f_!(U^c \cup V) \\
 &= f_!(U^c) \cup f_!(V) \\
 &= f_*(U)^c \cup f_!(V) \\
 &\stackrel{\text{def}}{=} [f_*(U), f_!(V)]_Y,
 \end{aligned}$$

where we have used:

- 022H 1. **Item 5** for the second equality.
- 022J 2. **Item 17** for the third equality.

Since $\mathcal{P}(Y)$ is posetal, naturality is automatic (**Categories**, **Item 4** of **Definition 11.2.7.1.2**). This finishes the proof.

Item 11, Preservation of Colimits: This follows from **Item 2** and ??, ?? of ??.³⁴

Item 12, Oplax Preservation of Limits: The inclusion $f_!(X) \subset Y$ is automatic. See [Pro25n] for the other inclusions.

Item 13, Symmetric Strict Monoidality With Respect to Unions: This follows from **Item 11**.

Item 14, Symmetric Oplax Monoidality With Respect to Intersections: The inclusions in the statement follow from **Item 12**. Since $\mathcal{P}(Y)$ is posetal, the commutativity of the diagrams in the definition of a symmetric oplax monoidal functor is automatic (**Categories**, **Item 4** of **Definition 11.2.7.1.2**).

Item 15, Interaction With Coproducts: Omitted.

Item 16, Interaction With Products: Omitted.

Item 17, Relation to Codirect Images: Applying **Item 16** of **Definition 4.6.3.1.7** to $X \setminus U$, we have

$$f_*(X \setminus U) = B \setminus f_!(X \setminus (X \setminus U))$$

³⁴Reference: [Pro25p].

$$= B \setminus f_!(U).$$

Taking complements, we then obtain

$$\begin{aligned} f_!(U) &= B \setminus (B \setminus f_!(U)), \\ &= B \setminus f_*(X \setminus U), \end{aligned}$$

which finishes the proof. \square

007V Proposition 4.6.1.1.6. Let $f: X \rightarrow Y$ be a function.

007W 1. *Functionality I.* The assignment $f \mapsto f_!$ defines a function

$$(-)_{*|X,Y}: \mathbf{Sets}(X, Y) \rightarrow \mathbf{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

007X 2. *Functionality II.* The assignment $f \mapsto f_!$ defines a function

$$(-)_{*|X,Y}: \mathbf{Sets}(X, Y) \rightarrow \mathbf{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

007Y 3. *Interaction With Identities.* For each $X \in \mathbf{Obj}(\mathbf{Sets})$, we have

$$(\mathrm{id}_X)_! = \mathrm{id}_{\mathcal{P}(X)}.$$

007Z 4. *Interaction With Composition.* For each pair of composable functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, we have

$$(g \circ f)_! = g_! \circ f_!$$

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \\ & \searrow (g \circ f)_! & \downarrow g_! \\ & & \mathcal{P}(Z). \end{array}$$

Proof. **Item 1, Functionality I:** There is nothing to prove.

Item 2, Functionality II: This follows from **Item 1** of **Definition 4.6.1.1.5**.

Item 3, Interaction With Identities: This follows from **Definition 4.6.1.1.4** and Kan Extensions, ?? of ??.

Item 4, Interaction With Composition: This follows from **Definition 4.6.1.1.4** and Kan Extensions, ?? of ??.

\square

0080 4.6.2 Inverse Images

Let $f: X \rightarrow Y$ be a function.

0081 **Definition 4.6.2.1.1.** The **inverse image function associated to f** is the function³⁵

$$f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$$

defined by³⁶

$$f^{-1}(V) \stackrel{\text{def}}{=} \{x \in X \mid \text{we have } f(x) \in V\}$$

for each $V \in \mathcal{P}(Y)$.

0082 **Remark 4.6.2.1.2.** Identifying $\mathcal{P}(Y)$ with $\text{Sets}(Y, \{\text{t}, \text{f}\})$ via **Item 2** of **Definition 4.5.1.1.4**, we see that the inverse image function associated to f is equivalently the function

$$f^*: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$$

defined by

$$f^*(\chi_V) \stackrel{\text{def}}{=} \chi_V \circ f$$

for each $\chi_V \in \mathcal{P}(Y)$, where $\chi_V \circ f$ is the composition

$$X \xrightarrow{f} Y \xrightarrow{\chi_V} \{\text{true}, \text{false}\}$$

in Sets .

0083 **Proposition 4.6.2.1.3.** Let $f: X \rightarrow Y$ be a function.

0084 1. *Functoriality.* The assignment $V \mapsto f^{-1}(V)$ defines a functor

$$f^{-1}: (\mathcal{P}(Y), \subset) \rightarrow (\mathcal{P}(X), \subset).$$

In particular, for each $U, V \in \mathcal{P}(Y)$, the following condition is satisfied:

$$(\star) \text{ If } U \subset V, \text{ then } f^{-1}(U) \subset f^{-1}(V).$$

³⁵*Further Notation:* Also written $f^*: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$.

³⁶*Further Terminology:* The set $f^{-1}(V)$ is called the **inverse image of V by f** .

0085 2. *Triple Adjointness.* We have a triple adjunction

$$(f_! \dashv f^{-1} \dashv f_*): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{f_!} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_*} \end{array} \mathcal{P}(Y),$$

witnessed by:

01M4 (a) Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\rightarrow f^{-1} \circ f_!, & \text{id}_{\mathcal{P}(Y)} &\rightarrow f_* \circ f^{-1}, \\ f_! \circ f^{-1} &\rightarrow \text{id}_{\mathcal{P}(Y)}, & f^{-1} \circ f_* &\rightarrow \text{id}_{\mathcal{P}(X)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset f^{-1}(f_!(U)), & V &\subset f_*(f^{-1}(V)), \\ f_!(f^{-1}(V)) &\subset V, & f^{-1}(f_*(U)) &\subset U \end{aligned}$$

indexed by $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$.

01M5 (b) Bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(Y)}(f_!(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(X)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f_*(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$ and (respectively) $V \in \mathcal{P}(X)$ and $U \in \mathcal{P}(Y)$. In particular:

01M6 i. The following conditions are equivalent:

01M7 A. We have $f_!(U) \subset V$.

01M8 B. We have $U \subset f^{-1}(V)$.

01M9 ii. The following conditions are equivalent:

01MA A. We have $f^{-1}(U) \subset V$.

01MB B. We have $U \subset f_*(V)$.

01MC 3. *Interaction With Unions of Families of Subsets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcup_{V \in \mathcal{U}} f^{-1}(V) = \bigcup_{U \in f^{-1}(\mathcal{U})} U$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(Y))$, where $f^{-1}(\mathcal{U}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{U})$.

01MD 4. *Interaction With Intersections of Families of Subsets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{P}(Y)) & \xrightarrow{(f^{-1})^{-1}} & \mathcal{P}(\mathcal{P}(X)) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\bigcap_{V \in \mathcal{U}} f^{-1}(V) = \bigcap_{U \in f^{-1}(\mathcal{U})} U$$

for each $\mathcal{U} \in \mathcal{P}(\mathcal{P}(Y))$, where $f^{-1}(\mathcal{U}) \stackrel{\text{def}}{=} (f^{-1})^{-1}(\mathcal{U})$.

01ME 5. *Interaction With Binary Unions.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cup \downarrow & & \downarrow \cup \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cup V) = f^{-1}(U) \cup f^{-1}(V)$$

for each $U, V \in \mathcal{P}(Y)$.

01MF 6. *Interaction With Binary Intersections.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{f^{-1} \times f^{-1}} & \mathcal{P}(X) \times \mathcal{P}(X) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \cap V) = f^{-1}(U) \cap f^{-1}(V)$$

for each $U, V \in \mathcal{P}(Y)$.

01MG 7. *Interaction With Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \setminus \downarrow & & \downarrow \setminus \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U \setminus V) = f^{-1}(U) \setminus f^{-1}(V)$$

for each $U, V \in \mathcal{P}(X)$.

01MH 8. *Interaction With Complements.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1, \text{op}}} & \mathcal{P}(X)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$f^{-1}(U^c) = f^{-1}(U)^c$$

for each $U \in \mathcal{P}(X)$.

01MJ 9. *Interaction With Symmetric Differences.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{\text{op}, -1} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ \Delta \downarrow & & \downarrow \Delta \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

i.e. we have

$$f^{-1}(U) \Delta f^{-1}(V) = f^{-1}(U \Delta V)$$

for each $U, V \in \mathcal{P}(Y)$.

01MK 10. *Interaction With Internal Homs of Powersets.* The diagram

$$\begin{array}{ccc} \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{-1, \text{op}} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\ [-1, -2]_Y \downarrow & & \downarrow [-1, -2]_X \\ \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have an equality of sets

$$f^{-1}([U, V]_Y) = [f^{-1}(U), f^{-1}(V)]_X,$$

natural in $U, V \in \mathcal{P}(X)$.

0086 11. *Preservation of Colimits.* We have an equality of sets

$$f^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cup f^{-1}(V) &= f^{-1}(U \cup V), \\ f^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(Y)$.

0087 12. *Preservation of Limits.* We have an equality of sets

$$f^{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(Y)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cap f^{-1}(V) &= f^{-1}(U \cap V), \\ f^{-1}(Y) &= X, \end{aligned}$$

natural in $U, V \in \mathcal{P}(Y)$.

0088 13. *Symmetric Strict Monoidality With Respect to Unions.* The inverse image function of **Item 1** has a symmetric strict monoidal structure

$$(f^{-1}, f^{-1, \otimes}, f_1^{-1, \otimes}): (\mathcal{P}(Y), \cup, \emptyset) \rightarrow (\mathcal{P}(X), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{U, V}^{-1, \otimes}: f^{-1}(U) \cup f^{-1}(V) &\xrightarrow{=} f^{-1}(U \cup V), \\ f_1^{-1, \otimes}: \emptyset &\xrightarrow{=} f^{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(Y)$.

0089 14. *Symmetric Strict Monoidality With Respect to Intersections.* The inverse image function of **Item 1** has a symmetric strict monoidal structure

$$(f^{-1}, f^{-1, \otimes}, f_1^{-1, \otimes}): (\mathcal{P}(Y), \cap, Y) \rightarrow (\mathcal{P}(X), \cap, X),$$

being equipped with equalities

$$\begin{aligned} f_{U, V}^{-1, \otimes}: f^{-1}(U) \cap f^{-1}(V) &\xrightarrow{=} f^{-1}(U \cap V), \\ f_1^{-1, \otimes}: X &\xrightarrow{=} f^{-1}(Y), \end{aligned}$$

natural in $U, V \in \mathcal{P}(Y)$.

- 008A** 15. *Interaction With Coproducts.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be maps of sets. We have

$$(f \amalg g)^{-1}(U' \amalg V') = f^{-1}(U') \amalg g^{-1}(V')$$

for each $U' \in \mathcal{P}(X')$ and each $V' \in \mathcal{P}(Y')$.

- 008B** 16. *Interaction With Products.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be maps of sets. We have

$$(f \boxtimes_{X' \times Y'} g)^{-1}(U' \boxtimes_{X' \times Y'} V') = f^{-1}(U') \boxtimes_{X \times Y} g^{-1}(V')$$

for each $U' \in \mathcal{P}(X')$ and each $V' \in \mathcal{P}(Y')$.

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Triple Adjointness: This follows from **Definition 4.6.1.1.4**, **Definition 4.6.2.1.2**, **Definition 4.6.3.1.4**, and Kan Extensions, ?? of ??.

Item 3, Interaction With Unions of Families of Subsets: We have

$$\begin{aligned} \bigcup_{U \in f^{-1}(\mathcal{V})} U &= \bigcup_{U \in \{f^{-1}(V) \in \mathcal{P}(X) \mid V \in \mathcal{V}\}} U \\ &= \bigcup_{V \in \mathcal{V}} f^{-1}(V). \end{aligned}$$

This finishes the proof.

Item 4, Interaction With Intersections of Families of Subsets: We have

$$\begin{aligned} \bigcap_{U \in f^{-1}(\mathcal{V})} U &= \bigcap_{U \in \{f^{-1}(V) \in \mathcal{P}(X) \mid V \in \mathcal{V}\}} U \\ &= \bigcap_{V \in \mathcal{V}} f^{-1}(V). \end{aligned}$$

This finishes the proof.

Item 5, Interaction With Binary Unions: See **[Pro25y]**.

Item 6, Interaction With Binary Intersections: See **[Pro25w]**.

Item 7, Interaction With Differences: See **[Pro25x]**.

Item 8, Interaction With Complements: See **[Pro25j]**.

Item 9, Interaction With Symmetric Differences: We have

$$f^{-1}(U \triangle V) = f^{-1}((U \cup V) \setminus (U \cap V))$$

$$\begin{aligned}
&= f^{-1}(U \cup V) \setminus f^{-1}(U \cap V) \\
&= f^{-1}(U) \cup f^{-1}(V) \setminus f^{-1}(U \cap V) \\
&= f^{-1}(U) \cup f^{-1}(V) \setminus f^{-1}(U) \cap f^{-1}(V) \\
&= f^{-1}(U) \Delta f^{-1}(V),
\end{aligned}$$

where we have used:

- 022K 1. **Item 2** of **Definition 4.3.12.1.2** for the first equality.
- 022L 2. **Item 7** for the second equality.
- 022M 3. **Item 5** for the third equality.
- 022N 4. **Item 6** for the fourth equality.
- 022P 5. **Item 2** of **Definition 4.3.12.1.2** for the fifth equality.

This finishes the proof.

Item 10, Interaction With Internal Homs of Powersets: We have

$$\begin{aligned}
f^{-1}([U, V]_Y) &\stackrel{\text{def}}{=} f^{-1}(U^c \cup V) \\
&= f^{-1}(U^c) \cup f^{-1}(V) \\
&= f^{-1}(U)^c \cup f^{-1}(V) \\
&\stackrel{\text{def}}{=} [f^{-1}(U), f^{-1}(V)]_X,
\end{aligned}$$

where we have used:

- 022Q 1. **Item 8** for the second equality.
- 022R 2. **Item 5** for the third equality.

Since $\mathcal{P}(Y)$ is posetal, naturality is automatic (**Categories**, **Item 4** of **Definition 11.2.7.1.2**). This finishes the proof.

Item 11, Preservation of Colimits: This follows from **Item 2** and ??, ?? of ??.³⁷

Item 12, Preservation of Limits: This follows from **Item 2** and ??, ?? of ??.³⁸

Item 13, Symmetric Strict Monoidality With Respect to Unions: This follows from **Item 11**.

³⁷Reference: [Pro25y].

³⁸Reference: [Pro25w].

Item 14, Symmetric Strict Monoidality With Respect to Intersections: This follows from *Item 12*.

Item 15, Interaction With Coproducts: Omitted.

Item 16, Interaction With Products: Omitted. \square

008C Proposition 4.6.2.1.4. Let $f: X \rightarrow Y$ be a function.

008D 1. *Functionality I.* The assignment $f \mapsto f^{-1}$ defines a function

$$(-)_{X,Y}^{-1}: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(Y), \mathcal{P}(X)).$$

008E 2. *Functionality II.* The assignment $f \mapsto f^{-1}$ defines a function

$$(-)_{X,Y}^{-1}: \text{Sets}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(Y), \subset), (\mathcal{P}(X), \subset)).$$

008F 3. *Interaction With Identities.* For each $X \in \text{Obj}(\text{Sets})$, we have

$$\text{id}_X^{-1} = \text{id}_{\mathcal{P}(X)}.$$

008G 4. *Interaction With Composition.* For each pair of composable functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, we have

$$(g \circ f)^{-1} = f^{-1} \circ g^{-1},$$

$$\begin{array}{ccc} \mathcal{P}(Z) & \xrightarrow{g^{-1}} & \mathcal{P}(Y) \\ & \searrow (g \circ f)^{-1} & \downarrow f^{-1} \\ & & \mathcal{P}(X). \end{array}$$

Proof. *Item 1, Functionality I:* There is nothing to prove.

Item 2, Functionality II: This follows from *Item 1* of *Definition 4.6.2.1.3*.

Item 3, Interaction With Identities: This follows from *Definition 4.6.2.1.2* and *Categories, Item 5* of *Definition II.1.4.1.2*.

Item 4, Interaction With Composition: This follows from *Definition 4.6.2.1.2* and *Categories, Item 2* of *Definition II.1.4.1.2*. \square

008H 4.6.3 Codirect Images

Let $f: X \rightarrow Y$ be a function.

008J Definition 4.6.3.1.1. The **codirect image function associated to f** is the function

$$f_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

defined by^{39,40}

$$\begin{aligned} f_*(U) &\stackrel{\text{def}}{=} \left\{ y \in Y \mid \begin{array}{l} \text{for each } x \in X, \text{ if we have} \\ f(x) = y, \text{ then } x \in U \end{array} \right\} \\ &= \{ y \in Y \mid \text{we have } f^{-1}(y) \subset U \} \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

008K Notation 4.6.3.1.2. Sometimes one finds the notation

$$\forall_f: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

for $f_!$. This notation comes from the fact that the following statements are equivalent, where $y \in Y$ and $U \in \mathcal{P}(X)$:

- We have $y \in \forall_f(U)$.
- For each $x \in X$, if $y = f(x)$, then $x \in U$.

We will not make use of this notation elsewhere in Clowder.

022V Warning 4.6.3.1.3. See **Definition 4.6.1.1.3**.

008L Remark 4.6.3.1.4. Identifying $\mathcal{P}(X)$ with $\text{Sets}(X, \{t, f\})$ via **Item 2** of **Definition 4.5.1.1.4**, we see that the codirect image function associated to f is equivalently the function

$$f_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

³⁹*Further Terminology:* The set $f_*(U)$ is called the **codirect image of U by f** .

⁴⁰We also have

$$\begin{aligned} f_*(U) &= f_!(U^c)^c \\ &\stackrel{\text{def}}{=} Y \setminus f_!(X \setminus U); \end{aligned}$$

defined by

$$\begin{aligned}
 f_*(\chi_U) &\stackrel{\text{def}}{=} \text{Ran}_f(\chi_U) \\
 &= \lim \left(\left(\underline{(-1)} \times f \right) \xrightarrow{\text{pr}} X \xrightarrow{\chi_U} \{\text{true}, \text{false}\} \right) \\
 &= \lim_{\substack{x \in X \\ f(x) = -1}} (\chi_U(x)) \\
 &= \bigwedge_{\substack{x \in X \\ f(x) = -1}} (\chi_U(x)).
 \end{aligned}$$

where we have used ?? for the second equality. In other words, we have

$$\begin{aligned}
 [f_*(\chi_U)](y) &= \bigwedge_{\substack{x \in X \\ f(x) = y}} (\chi_U(x)) \\
 &= \begin{cases} \text{true} & \text{if, for each } x \in X \text{ such that} \\ & f(x) = y, \text{ we have } x \in U, \\ \text{false} & \text{otherwise} \end{cases} \\
 &= \begin{cases} \text{true} & \text{if } f^{-1}(y) \subset U \\ \text{false} & \text{otherwise} \end{cases}
 \end{aligned}$$

for each $y \in Y$.

see [Item 16](#) of [Definition 4.6.3.1.7](#).

008M Definition 4.6.3.1.5. Let U be a subset of X .^{41,42}

008N 1. The **image part of the codirect image** $f_*(U)$ of U is the set $f_{*,\text{im}}(U)$ defined by

$$\begin{aligned} f_{*,\text{im}}(U) &\stackrel{\text{def}}{=} f_*(U) \cap \text{Im}(f) \\ &= \left\{ y \in Y \left| \begin{array}{l} \text{we have } f^{-1}(y) \subset U \\ \text{and } f^{-1}(y) \neq \emptyset. \end{array} \right. \right\}. \end{aligned}$$

008P 2. The **complement part of the codirect image** $f_*(U)$ of U is the set $f_{*,\text{cp}}(U)$ defined by

$$\begin{aligned} f_{*,\text{cp}}(U) &\stackrel{\text{def}}{=} f_*(U) \cap (Y \setminus \text{Im}(f)) \\ &= Y \setminus \text{Im}(f) \\ &= \left\{ y \in Y \left| \begin{array}{l} \text{we have } f^{-1}(y) \subset U \\ \text{and } f^{-1}(y) = \emptyset. \end{array} \right. \right\} \\ &= \{ y \in Y \mid f^{-1}(y) = \emptyset \}. \end{aligned}$$

008Q Example 4.6.3.1.6. Here are some examples of codirect images.

⁴¹Note that we have

$$f_*(U) = f_{*,\text{im}}(U) \cup f_{*,\text{cp}}(U),$$

as

$$\begin{aligned} f_*(U) &= f_*(U) \cap Y \\ &= f_*(U) \cap (\text{Im}(f) \cup (Y \setminus \text{Im}(f))) \\ &= (f_*(U) \cap \text{Im}(f)) \cup (f_*(U) \cap (Y \setminus \text{Im}(f))) \\ &\stackrel{\text{def}}{=} f_{*,\text{im}}(U) \cup f_{*,\text{cp}}(U). \end{aligned}$$

⁴²In terms of the meet computation of $f_*(U)$ of **Definition 4.6.3.1.4**, namely

$$f_*(\chi_U) = \bigwedge_{\substack{x \in X \\ f(x) = -1}} (\chi_U(x)),$$

we see that $f_{*,\text{im}}$ corresponds to meets indexed over nonempty sets, while $f_{*,\text{cp}}$ corresponds to meets indexed over the empty set.

- 0231 1. *Multiplication by Two.* Consider the function $f: \mathbb{N} \rightarrow \mathbb{N}$ given by

$$f(n) \stackrel{\text{def}}{=} 2n$$

for each $n \in \mathbb{N}$. Since f is injective, we have

$$\begin{aligned} f_{*,\text{im}}(U) &= f_!(U) \\ f_{*,\text{cp}}(U) &= \{\text{odd natural numbers}\} \end{aligned}$$

for any $U \subset \mathbb{N}$. In particular, we have

$$f_*(\{\text{even natural numbers}\}) = \mathbb{N}.$$

- 0232 2. *Parabolas.* Consider the function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) \stackrel{\text{def}}{=} x^2$$

for each $x \in \mathbb{R}$. We have

$$f_{*,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any $U \subset \mathbb{R}$. Moreover, since $f^{-1}(x) = \{-\sqrt{x}, \sqrt{x}\}$, we have e.g.:

$$\begin{aligned} f_{*,\text{im}}([0, 1]) &= \{0\}, \\ f_{*,\text{im}}([-1, 1]) &= [0, 1], \\ f_{*,\text{im}}([1, 2]) &= \emptyset, \\ f_{*,\text{im}}([-2, -1] \cup [1, 2]) &= [1, 4]. \end{aligned}$$

- 0233 3. *Circles.* Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$f(x, y) \stackrel{\text{def}}{=} x^2 + y^2$$

for each $(x, y) \in \mathbb{R}^2$. We have

$$f_{*,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any $U \subset \mathbb{R}^2$, and since

$$f^{-1}(r) = \begin{cases} \text{a circle of radius } r \text{ about the origin} & \text{if } r > 0, \\ \{(0, 0)\} & \text{if } r = 0, \\ \emptyset & \text{if } r < 0, \end{cases}$$

we have e.g.:

$$\begin{aligned} f_{*,\text{im}}([-1, 1] \times [-1, 1]) &= [0, 1], \\ f_{*,\text{im}}(([-1, 1] \times [-1, 1]) \setminus [-1, 1] \times \{0\}) &= \emptyset. \end{aligned}$$

008R Proposition 4.6.3.1.7. Let $f: X \rightarrow Y$ be a function.

008S 1. *Functoriality.* The assignment $U \mapsto f_*(U)$ defines a functor

$$f_*: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset).$$

In particular, for each $U, V \in \mathcal{P}(X)$, the following condition is satisfied:

$$(\star) \text{ If } U \subset V, \text{ then } f_*(U) \subset f_*(V).$$

008T 2. *Triple Adjointness.* We have a triple adjunction

$$(f_! \dashv f^{-1} \dashv f_*): \quad \mathcal{P}(X) \begin{array}{c} \xrightarrow{f_!} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_*} \end{array} \mathcal{P}(Y),$$

witnessed by:

01ML (a) Units and counits of the form

$$\begin{aligned} \text{id}_{\mathcal{P}(X)} &\rightarrow f^{-1} \circ f_!, & \text{id}_{\mathcal{P}(Y)} &\rightarrow f_* \circ f^{-1}, \\ f_! \circ f^{-1} &\rightarrow \text{id}_{\mathcal{P}(Y)}, & f^{-1} \circ f_* &\rightarrow \text{id}_{\mathcal{P}(X)}, \end{aligned}$$

having components of the form

$$\begin{aligned} U &\subset f^{-1}(f_!(U)), & V &\subset f_*(f^{-1}(V)), \\ f_!(f^{-1}(V)) &\subset V, & f^{-1}(f_*(U)) &\subset U \end{aligned}$$

indexed by $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$.

01MM (b) Bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(Y)}(f_!(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(X)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(X)}(U, f_*(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$ and (respectively) $V \in \mathcal{P}(X)$ and $U \in \mathcal{P}(Y)$. In particular:

01MN

i. The following conditions are equivalent:

01MP

A. We have $f_!(U) \subset V$.

01MQ

B. We have $U \subset f^{-1}(V)$.

01MR

ii. The following conditions are equivalent:

01MS

A. We have $f^{-1}(U) \subset V$.

01MT

B. We have $U \subset f_*(V)$.

01MU

3. *Interaction With Unions of Families of Subsets.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\
 \downarrow \cup & & \downarrow \cup \\
 \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$\bigcup_{U \in \mathcal{U}} f_*(U) = \bigcup_{V \in f_*(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$.

01MV

4. *Interaction With Intersections of Families of Subsets.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(\mathcal{P}(X)) & \xrightarrow{(f_*)_*} & \mathcal{P}(\mathcal{P}(Y)) \\
 \downarrow \cap & & \downarrow \cap \\
 \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$\bigcap_{U \in \mathcal{U}} f_*(U) = \bigcap_{V \in f_*(\mathcal{U})} V$$

for each $\mathcal{U} \in \mathcal{P}(X)$, where $f_*(\mathcal{U}) \stackrel{\text{def}}{=} (f_*)_*(\mathcal{U})$.

- 01MW 5. *Interaction With Binary Unions.* Let $f: X \rightarrow Y$ be a function. We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cup \downarrow & \subset & \downarrow \cup \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U) \cup f_*(V) \subset f_*(U \cup V)$$

indexed by $U, V \in \mathcal{P}(X)$.

- 01MX 6. *Interaction With Binary Intersections.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{f_* \times f_*} & \mathcal{P}(Y) \times \mathcal{P}(Y) \\ \cap \downarrow & & \downarrow \cap \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U) \cap f_*(V) = f_*(U \cap V)$$

for each $U, V \in \mathcal{P}(X)$.

- 01MY 7. *Interaction With Complements.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ (-)^c \downarrow & & \downarrow (-)^c \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_*(U^c) = f_!(U)^c$$

for each $U \in \mathcal{P}(X)$.

01MZ 8. *Interaction With Symmetric Differences.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_*^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ \Delta \downarrow & \supset & \downarrow \Delta \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$f_*(U \Delta V) \subset f_*(U) \Delta f_*(V)$$

indexed by $U, V \in \mathcal{P}(X)$.

01N0 9. *Interaction With Internal Homs of Powersets.* We have a natural transformation

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{f_!^{\text{op}} \times f_*} & \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) \\ [-, -]_X \downarrow & \supset & \downarrow [-, -]_Y \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \end{array}$$

with components

$$[f_!(U), f_*(V)]_Y \subset f_*([U, V]_X)$$

indexed by $U, V \in \mathcal{P}(X)$.

008U 10. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} f_*(U_i) \subset f_*\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} f_*(U) \cup f_*(V) &\rightarrow f_*(U \cup V), \\ \emptyset &\rightarrow f_*(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

008V II. *Preservation of Limits.* We have an equality of sets

$$f_* \left(\bigcap_{i \in I} U_i \right) = \bigcap_{i \in I} f_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(X)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U \cap V) &= f_*(U) \cap f^{-1}(V), \\ f_*(X) &= Y, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

008W 12. *Symmetric Lax Monoidality With Respect to Unions.* The codirect image function of **Item I** has a symmetric lax monoidal structure

$$(f_*, f_*^\otimes, f_{*|1}^\otimes): (\mathcal{P}(X), \cup, \emptyset) \rightarrow (\mathcal{P}(Y), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U) \cup f_*(V) &\rightarrow f_*(U \cup V), \\ f_{*|1}^\otimes: \emptyset &\rightarrow f_*(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

008X 13. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item I** has a symmetric strict monoidal structure

$$(f_*, f_*^\otimes, f_{*|1}^\otimes): (\mathcal{P}(X), \cap, X) \rightarrow (\mathcal{P}(Y), \cap, Y),$$

being equipped with equalities

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U \cap V) &\xrightarrow{=} f_*(U) \cap f_*(V), \\ f_{*|1}^\otimes: f_*(X) &\xrightarrow{=} Y, \end{aligned}$$

natural in $U, V \in \mathcal{P}(X)$.

008Y 14. *Interaction With Coproducts.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be maps

of sets. We have

$$(f \amalg g)_*(U \amalg V) = f_*(U) \amalg g_*(V)$$

for each $U \in \mathcal{P}(X)$ and each $V \in \mathcal{P}(Y)$.

- 008Z** 15. *Interaction With Products.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be maps of sets. We have

$$(f \boxtimes_{X \times Y} g)_*(U \boxtimes_{X \times Y} V) = f_*(U) \boxtimes_{X' \times Y'} g_*(V)$$

for each $U \in \mathcal{P}(X)$ and each $V \in \mathcal{P}(Y)$.

- 0090** 16. *Relation to Direct Images.* We have

$$\begin{aligned} f_*(U) &= f_!(U^c)^c \\ &= Y \setminus f_!(X \setminus U) \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

- 0091** 17. *Interaction With Injections.* If f is injective, then we have

$$\begin{aligned} f_{*,\text{im}}(U) &= f_!(U), \\ f_{*,\text{cp}}(U) &= Y \setminus \text{Im}(f), \end{aligned}$$

and so

$$\begin{aligned} f_*(U) &= f_{*,\text{im}}(U) \cup f_{*,\text{cp}}(U) \\ &= f_!(U) \cup (Y \setminus \text{Im}(f)) \end{aligned}$$

for each $U \in \mathcal{P}(X)$.

- 0092** 18. *Interaction With Surjections.* If f is surjective, then we have

$$\begin{aligned} f_{*,\text{im}}(U) &\subset f_!(U), \\ f_{*,\text{cp}}(U) &= \emptyset, \end{aligned}$$

and so

$$f_*(U) \subset f_!(U)$$

for each $U \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Triple Adjointness: This follows from **Definition 4.6.1.1.4**, **Definition 4.6.2.1.2**, **Definition 4.6.3.1.4**, and Kan Extensions, ?? of ??.

Item 3, Interaction With Unions of Families of Subsets: We have

$$\begin{aligned} \bigcup_{V \in f_*(\mathcal{U})} V &= \bigcup_{V \in \{f_*(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcup_{U \in \mathcal{U}} f_*(U). \end{aligned}$$

This finishes the proof.

Item 4, Interaction With Intersections of Families of Subsets: We have

$$\begin{aligned} \bigcap_{V \in f_*(\mathcal{U})} V &= \bigcap_{V \in \{f_*(U) \in \mathcal{P}(X) \mid U \in \mathcal{U}\}} V \\ &= \bigcap_{U \in \mathcal{U}} f_*(U). \end{aligned}$$

This finishes the proof.

Item 5, Interaction With Binary Unions: We have

$$\begin{aligned} f_*(U) \cup f_*(V) &= f_!(U^c)^c \cup f_!(V^c)^c \\ &= (f_!(U^c) \cap f_!(V^c))^c \\ &\subset (f_!(U^c \cap V^c))^c \\ &= f_!((U \cup V)^c)^c \\ &= f_*(U \cup V), \end{aligned}$$

where:

- 023X 1. We have used **Item 16** for the first equality.
- 023Y 2. We have used **Item 2** of **Definition 4.3.11.1.2** for the second equality.
- 023Z 3. We have used **Item 6** of **Definition 4.6.1.1.5** for the third equality.
- 0240 4. We have used **Item 2** of **Definition 4.3.11.1.2** for the fourth equality.
- 0241 5. We have used **Item 16** for the last equality.

This finishes the proof.

Item 6, Interaction With Binary Intersections: This follows from *Item 11*.

Item 7, Interaction With Complements: Omitted.

Item 8, Interaction With Symmetric Differences: Omitted.

Item 9, Interaction With Internal Homs of Powersets: We have

$$\begin{aligned} [f_!(U), f^!(V)]_X &\stackrel{\text{def}}{=} f_!(U)^c \cup f_*(V) \\ &= f_*(U^c) \cup f_*(V) \\ &\subset f_*(U^c \cup V) \\ &\stackrel{\text{def}}{=} f_*([U, V]_X), \end{aligned}$$

where we have used:

- 0242 1. *Item 7* of *Definition 4.6.3.1.7* for the second equality.
- 0243 2. *Item 5* of *Definition 4.6.3.1.7* for the inclusion.

Since $\mathcal{P}(X)$ is posetal, naturality is automatic (*Categories*, *Item 4* of *Definition 11.2.7.1.2*). This finishes the proof.

Item 10, Lax Preservation of Colimits: Omitted.

Item 11, Preservation of Limits: This follows from *Item 2* and ??, ?? of ??.

Item 12, Symmetric Lax Monoidality With Respect to Unions: This follows from *Item 10*.

Item 13, Symmetric Strict Monoidality With Respect to Intersections: This follows from *Item 11*.

Item 14, Interaction With Coproducts: Omitted.

Item 15, Interaction With Products: Omitted.

Item 16, Relation to Direct Images: We claim that $f_*(U) = Y \setminus f_!(X \setminus U)$.

- *The First Implication.* We claim that

$$f_*(U) \subset Y \setminus f_!(X \setminus U).$$

Let $y \in f_*(U)$. We need to show that $y \notin f_!(X \setminus U)$, i.e. that there is no $x \in X \setminus U$ such that $f(x) = y$.

This is indeed the case, as otherwise we would have $x \in f^{-1}(y)$ and $x \notin U$, contradicting $f^{-1}(y) \subset U$ (which holds since $y \in f_*(U)$).

Thus $y \in Y \setminus f_!(X \setminus U)$.

- *The Second Implication.* We claim that

$$Y \setminus f_!(X \setminus U) \subset f_*(U).$$

Let $y \in Y \setminus f_!(X \setminus U)$. We need to show that $y \in f_*(U)$, i.e. that $f^{-1}(y) \subset U$.

Since $y \notin f_!(X \setminus U)$, there exists no $x \in X \setminus U$ such that $y = f(x)$, and hence $f^{-1}(y) \subset U$.

Thus $y \in f_*(U)$.

This finishes the proof of **Item 16**.

Item 17, Interaction With Injections: Omitted.

Item 18, Interaction With Surjections: Omitted. \square

0093 Proposition 4.6.3.1.8. Let $f: X \rightarrow B$ be a function.

0094 1. *Functionality I.* The assignment $f \mapsto f_*$ defines a function

$$(-)_{!|X,Y}: \text{Sets}(X, Y) \rightarrow \text{Sets}(\mathcal{P}(X), \mathcal{P}(Y)).$$

0095 2. *Functionality II.* The assignment $f \mapsto f_*$ defines a function

$$(-)_{!|X,Y}: \text{Sets}(X, Y) \rightarrow \text{Pos}((\mathcal{P}(X), \subset), (\mathcal{P}(Y), \subset)).$$

0096 3. *Interaction With Identities.* For each $X \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_X)_* = \text{id}_{\mathcal{P}(X)}.$$

0097 4. *Interaction With Composition.* For each pair of composable functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, we have

$$(g \circ f)_* = g_* \circ f_*$$

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y) \\ & \searrow (g \circ f)_* & \downarrow g_* \\ & & \mathcal{P}(Z). \end{array}$$

Proof. **Item 1, Functionality I:** There is nothing to prove.

Item 2, Functionality II: This follows from **Item 1** of **Definition 4.6.3.1.7**.

Item 3, Interaction With Identities: This follows from **Definition 4.6.3.1.4** and Kan Extensions, ?? of ??.

Item 4, Interaction With Composition: This follows from **Definition 4.6.3.1.4** and Kan Extensions, ?? of ?? \square

01N1 4.6.4 A Six-Functor Formalism for Sets

01N2 **Remark 4.6.4.1.1.** The assignment $X \mapsto \mathcal{P}(X)$ together with the functors f_* , f^{-1} , and $f_!$ of **Item 1** of **Definition 4.6.1.1.5**, **Item 1** of **Definition 4.6.2.1.3**, and **Item 1** of **Definition 4.6.3.1.7**, and the functors

$$\begin{aligned} -_1 \cap -_2 &: \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow \mathcal{P}(X), \\ [-_1, -_2]_X &: \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X) \end{aligned}$$

of **Item 1** of **Definition 4.3.9.1.2** and **Item 1** of **Definition 4.4.7.1.3** satisfy several properties reminiscent of a six functor formalism in the sense of ??.

We collect these properties in **Definition 4.6.4.1.2** below.⁴³

01N3 **Proposition 4.6.4.1.2.** Let X be a set.

01N4 1. *The Beck–Chevalley Condition.* Let

$$\begin{array}{ccc} X \times_Z Y & \xrightarrow{\text{pr}_2} & Y \\ \text{pr}_1 \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

be a pullback diagram in **Sets**. We have

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{\text{pr}_1^{-1}} & \mathcal{P}(X \times_Z Y) \\ f_! \downarrow & & \downarrow (\text{pr}_2)_! \\ \mathcal{P}(Z) & \xrightarrow{g^{-1}} & \mathcal{P}(Y), \end{array} \quad \begin{array}{l} g^{-1} \circ f_! = (\text{pr}_2)_! \circ \text{pr}_1^{-1}, \\ f^{-1} \circ g_! = (\text{pr}_1)_! \circ \text{pr}_2^{-1}, \end{array} \quad \begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{\text{pr}_2^{-1}} & \mathcal{P}(X \times_Z Y) \\ g_! \downarrow & & \downarrow (\text{pr}_1)_! \\ \mathcal{P}(Z) & \xrightarrow{f^{-1}} & \mathcal{P}(Y). \end{array}$$

⁴³See also [nLaz5].

01N5 2. *The Projection Formula I.* The diagram

$$\begin{array}{ccc}
 & \mathcal{P}(X) \times \mathcal{P}(X) & \\
 \text{id}_{\mathcal{P}(X)} \times f^{-1} \nearrow & & \searrow \cap \\
 \mathcal{P}(X) \times \mathcal{P}(Y) & & \mathcal{P}(X) \\
 f_! \times \text{id}_{\mathcal{P}(Y)} \searrow & & \downarrow f_! \\
 \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{\cap} & \mathcal{P}(Y),
 \end{array}$$

commutes, i.e. we have

$$f_!(U \cap f^{-1}(V)) = f_!(U) \cap V$$

for each $U \in \mathcal{P}(X)$ and each $V \in \mathcal{P}(Y)$.

01N6 3. *The Projection Formula II.* We have a natural transformation

$$\begin{array}{ccc}
 & \mathcal{P}(X) \times \mathcal{P}(X) & \\
 \text{id}_{\mathcal{P}(X)} \times f^{-1} \nearrow & & \searrow \cap \\
 \mathcal{P}(X) \times \mathcal{P}(Y) & & \mathcal{P}(X) \\
 f_* \times \text{id}_{\mathcal{P}(Y)} \searrow & \cup & \downarrow f_* \\
 \mathcal{P}(Y) \times \mathcal{P}(Y) & \xrightarrow{\cap} & \mathcal{P}(Y),
 \end{array}$$

with components

$$f_*(U) \cap V \subset f_*(U \cap f^{-1}(V))$$

indexed by $U \in \mathcal{P}(X)$ and $V \in \mathcal{P}(Y)$.

01N7 4. *Strong Closed Monoidality.* The diagram

$$\begin{array}{ccc}
 \mathcal{P}(Y)^{\text{op}} \times \mathcal{P}(Y) & \xrightarrow{f^{-1, \text{op}} \times f^{-1}} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \\
 \downarrow [-1, -2]_Y & & \downarrow [-1, -2]_X \\
 \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have an equality of sets

$$f^{-1}([U, V]_Y) = [f^{-1}(U), f^{-1}(V)]_X,$$

natural in $U, V \in \mathcal{P}(X)$.

01N8 5. *The External Tensor Product.* We have an external tensor product

$$-_1 \boxtimes_{X \times Y} -_2: \mathcal{P}(X) \times \mathcal{P}(Y) \rightarrow \mathcal{P}(X \times Y)$$

given by

$$\begin{aligned} U \boxtimes_{X \times Y} V &\stackrel{\text{def}}{=} \text{pr}_1^{-1}(U) \cap \text{pr}_2^{-1}(V) \\ &= \{(u, v) \in X \times Y \mid u \in U \text{ and } v \in V\}. \end{aligned}$$

This is the same map as the one in **Item 5** of **Definition 4.4.I.I.4**. Moreover, the following conditions are satisfied:

01N9 (a) *Interaction With Direct Images.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be functions. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(Y) & \xrightarrow{f! \times g!} & \mathcal{P}(X') \times \mathcal{P}(Y') \\ \boxtimes_{X \times Y} \downarrow & & \downarrow \boxtimes_{X' \times Y'} \\ \mathcal{P}(X \times Y) & \xrightarrow{f! \times g!} & \mathcal{P}(X' \times Y') \end{array}$$

commutes, i.e. we have

$$[f! \times g!](U \boxtimes_{X \times Y} V) = f!(U) \boxtimes_{X' \times Y'} g!(V)$$

for each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$.

01NA (b) *Interaction With Inverse Images.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be functions. The diagram

$$\begin{array}{ccc} \mathcal{P}(X') \times \mathcal{P}(Y') & \xrightarrow{f^{-1} \times g^{-1}} & \mathcal{P}(X) \times \mathcal{P}(Y) \\ \boxtimes_{X' \times Y'} \downarrow & & \downarrow \boxtimes_{X \times Y} \\ \mathcal{P}(X' \times Y') & \xrightarrow{f^{-1} \times g^{-1}} & \mathcal{P}(X \times Y) \end{array}$$

commutes, i.e. we have

$$[f^{-1} \times g^{-1}](U \boxtimes_{X' \times Y'} V) = f^{-1}(U) \boxtimes_{X \times Y} g^{-1}(V)$$

for each $(U, V) \in \mathcal{P}(X') \times \mathcal{P}(Y')$.

01NB

(c) *Interaction With Codirect Images.* Let $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$ be functions. The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(Y) & \xrightarrow{f_* \times g_*} & \mathcal{P}(X') \times \mathcal{P}(Y') \\ \boxtimes_{X \times Y} \downarrow & & \downarrow \boxtimes_{X' \times Y'} \\ \mathcal{P}(X \times Y) & \xrightarrow{f_* \times g_*} & \mathcal{P}(X' \times Y') \end{array}$$

commutes, i.e. we have

$$[f_* \times g_*](U \boxtimes_{X \times Y} V) = f_*(U) \boxtimes_{X' \times Y'} g_*(V)$$

for each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$.

01NC

(d) *Interaction With Diagonals.* The diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(X) & \xrightarrow{\boxtimes_{X \times X}} & \mathcal{P}(X \times X) \\ & \searrow \cap & \downarrow \Delta_X^{-1} \\ & & \mathcal{P}(X), \end{array}$$

i.e. we have

$$U \cap V = \Delta_X^{-1}(U \boxtimes_{X \times X} V)$$

for each $U, V \in \mathcal{P}(X)$.

01ND

6. *The Dualisation Functor.* We have a functor

$$D_X: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X)$$

given by

$$D_X(U) \stackrel{\text{def}}{=} [U, \emptyset]_X \\ \stackrel{\text{def}}{=} U^c$$

for each $U \in \mathcal{P}(X)$, as in [Item 5](#) of [Definition 4.4.7.1.3](#), satisfying the following conditions:

01NE (a) *Duality*. We have

$$D_X(D_X(U)) = U,$$

$$\begin{array}{ccc} \mathcal{P}(X) & \xrightarrow{D_X} & \mathcal{P}(X) \\ & \searrow \text{id}_{\mathcal{P}(X)} & \downarrow D_X \\ & & \mathcal{P}(X). \end{array}$$

01NF (b) *Duality*. The diagram

$$\begin{array}{ccc} & \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X)^{\text{op}} \xrightarrow{\cap^{\text{op}}} \mathcal{P}(X)^{\text{op}} & \\ \text{id}_{\mathcal{P}(X)^{\text{op}}} \times D_X \nearrow & & \searrow D_X \\ \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) & \xrightarrow{[-1, -2]_X} & \mathcal{P}(X) \end{array}$$

commutes, i.e. we have

$$\underbrace{D_X(U \cap D_X(V))}_{\stackrel{\text{def}}{=} [U \cap [V, \emptyset]_X, \emptyset]_X} = [U, V]_X$$

for each $U, V \in \mathcal{P}(X)$.

01NG (c) *Interaction With Direct Images*. The diagram

$$\begin{array}{ccc} \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_*^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\ D_X \downarrow & & \downarrow D_Y \\ \mathcal{P}(X) & \xrightarrow{f_!} & \mathcal{P}(Y) \end{array}$$

commutes, i.e. we have

$$f_!(D_X(U)) = D_Y(f_*(U))$$

for each $U \in \mathcal{P}(X)$.

01NH

(d) *Interaction With Inverse Images*. The diagram

$$\begin{array}{ccc}
 \mathcal{P}(Y)^{\text{op}} & \xrightarrow{f^{-1, \text{op}}} & \mathcal{P}(X)^{\text{op}} \\
 D_Y \downarrow & & \downarrow D_X \\
 \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X)
 \end{array}$$

commutes, i.e. we have

$$f^{-1}(D_Y(U)) = D_X(f^{-1}(U))$$

for each $U \in \mathcal{P}(X)$.

01NJ

(e) *Interaction With Codirect Images*. The diagram

$$\begin{array}{ccc}
 \mathcal{P}(X)^{\text{op}} & \xrightarrow{f_!^{\text{op}}} & \mathcal{P}(Y)^{\text{op}} \\
 D_X \downarrow & & \downarrow D_Y \\
 \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(Y)
 \end{array}$$

commutes, i.e. we have

$$f_*(D_X(U)) = D_Y(f_!(U))$$

for each $U \in \mathcal{P}(X)$.*Proof.* **Item 1**, *The Beck–Chevalley Condition*: We have

$$\begin{aligned}
 [g^{-1} \circ f_!](U) &\stackrel{\text{def}}{=} g^{-1}(f_!(U)) \\
 &\stackrel{\text{def}}{=} \{y \in Y \mid g(y) \in f_!(U)\} \\
 &= \left\{ y \in Y \mid \begin{array}{l} \text{there exists some } x \in U \\ \text{such that } f(x) = g(y) \end{array} \right\} \\
 &= \left\{ y \in Y \mid \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid x \in U\} \end{array} \right\}
 \end{aligned}$$

$$\begin{aligned}
&= \left\{ y \in Y \left| \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid x \in U\} \\ \text{such that } y = y \end{array} \right. \right\} \\
&= \left\{ y \in Y \left| \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid x \in U\} \\ \text{such that } \text{pr}_2(x, y) = y \end{array} \right. \right\} \\
&\stackrel{\text{def}}{=} (\text{pr}_2)_! (\{(x, y) \in X \times_Z Y \mid x \in U\}) \\
&= (\text{pr}_2)_! (\{(x, y) \in X \times_Z Y \mid \text{pr}_1(x, y) \in U\}) \\
&\stackrel{\text{def}}{=} (\text{pr}_2)_! (\text{pr}_1^{-1}(U)) \\
&\stackrel{\text{def}}{=} [(\text{pr}_2)_! \circ \text{pr}_1^{-1}](U)
\end{aligned}$$

for each $U \in \mathcal{P}(X)$. Therefore, we have

$$g^{-1} \circ f_! = (\text{pr}_2)_! \circ \text{pr}_1^{-1}.$$

For the second equality, we have

$$\begin{aligned}
[f^{-1} \circ g_!](U) &\stackrel{\text{def}}{=} f^{-1}(g_!(U)) \\
&\stackrel{\text{def}}{=} \{x \in X \mid f(x) \in g_!(V)\} \\
&= \left\{ x \in X \left| \begin{array}{l} \text{there exists some } y \in V \\ \text{such that } f(x) = g(y) \end{array} \right. \right\} \\
&= \left\{ x \in X \left| \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid y \in V\} \end{array} \right. \right\} \\
&= \left\{ x \in X \left| \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid y \in V\} \\ \text{such that } x = x \end{array} \right. \right\} \\
&= \left\{ x \in X \left| \begin{array}{l} \text{there exists some} \\ (x, y) \in \{(x, y) \in X \times_Z Y \mid y \in V\} \\ \text{such that } \text{pr}_1(x, y) = x \end{array} \right. \right\} \\
&\stackrel{\text{def}}{=} (\text{pr}_1)_! (\{(x, y) \in X \times_Z Y \mid y \in V\}) \\
&= (\text{pr}_1)_! (\{(x, y) \in X \times_Z Y \mid \text{pr}_2(x, y) \in V\}) \\
&\stackrel{\text{def}}{=} (\text{pr}_1)_! (\text{pr}_2^{-1}(V))
\end{aligned}$$

$$\stackrel{\text{def}}{=} [(\text{pr}_1)_! \circ \text{pr}_2^{-1}](V)$$

for each $V \in \mathcal{P}(Y)$. Therefore, we have

$$f^{-1} \circ g_! = (\text{pr}_1)_! \circ \text{pr}_2^{-1}.$$

This finishes the proof.

Item 2, The Projection Formula I: We claim that

$$f_!(U) \cap V \subset f_!(U \cap f^{-1}(V)).$$

Indeed, we have

$$\begin{aligned} f_!(U) \cap V &\subset f_!(U) \cap f_!(f^{-1}(V)) \\ &= f_!(U \cap f^{-1}(V)), \end{aligned}$$

where we have used:

024B 1. *Item 2* of **Definition 4.6.1.1.5** for the inclusion.

024C 2. *Item 6* of **Definition 4.6.1.1.5** for the equality.

Conversely, we claim that

$$f_!(U \cap f^{-1}(V)) \subset f_!(U) \cap V.$$

Indeed:

024D 1. Let $y \in f_!(U \cap f^{-1}(V))$.

024E 2. Since $y \in f_!(U \cap f^{-1}(V))$, there exists some $x \in U \cap f^{-1}(V)$ such that $f(x) = y$.

024F 3. Since $x \in U \cap f^{-1}(V)$, we have $x \in U$, and thus $f(x) \in f_!(U)$.

024G 4. Since $x \in U \cap f^{-1}(V)$, we have $x \in f^{-1}(V)$, and thus $f(x) \in V$.

024H 5. Since $f(x) \in f_!(U)$ and $f(x) \in V$, we have $f(x) \in f_!(U) \cap V$.

024J 6. But $y = f(x)$, so $y \in f_!(U) \cap V$.

024K 7. Thus $f_!(U \cap f^{-1}(V)) \subset f_!(U) \cap V$.

This finishes the proof.

Item 3, The Projection Formula II: We have

$$\begin{aligned} f_*(U) \cap V &\subset f_*(U) \cap f_*(f^{-1}(V)) \\ &= f_*(U \cap f^{-1}(V)), \end{aligned}$$

where we have used:

- 024L 1. *Item 2* of *Definition 4.6.3.1.7* for the inclusion.
- 024M 2. *Item 6* of *Definition 4.6.3.1.7* for the equality.

Since $\mathcal{P}(Y)$ is posetal, naturality is automatic (*Categories, Item 4* of *Definition 11.2.7.1.2*).

Item 4, Strong Closed Monoidality: This is a repetition of *Item 19* of *Definition 4.4.7.1.3* and is proved there.

Item 5, The External Tensor Product: We have

$$\begin{aligned} U \boxtimes_{X \times Y} V &\stackrel{\text{def}}{=} \text{pr}_1^{-1}(U) \cap \text{pr}_2^{-1}(V) \\ &\stackrel{\text{def}}{=} \{(x, y) \in X \times Y \mid \text{pr}_1(x, y) \in U\} \\ &\quad \cup \{(x, y) \in X \times Y \mid \text{pr}_2(x, y) \in V\} \\ &= \{(x, y) \in X \times Y \mid x \in U\} \\ &\quad \cup \{(x, y) \in X \times Y \mid y \in V\} \\ &= \{(x, y) \in X \times Y \mid x \in U \text{ and } y \in V\} \\ &\stackrel{\text{def}}{=} U \times V. \end{aligned}$$

Next, we claim that *Items 5a* to *5d* are indeed true:

- 024N 1. *Proof of Item 5a:* This is a repetition of *Item 16* of *Definition 4.6.1.1.5* and is proved there.
- 024P 2. *Proof of Item 5b:* This is a repetition of *Item 16* of *Definition 4.6.2.1.3* and is proved there.
- 024Q 3. *Proof of Item 5c:* This is a repetition of *Item 15* of *Definition 4.6.3.1.7* and is proved there.

024R 4. *Proof of Item 5d:* We have

$$\begin{aligned}\Delta_X^{-1}(U \boxtimes_{X \times X} V) &\stackrel{\text{def}}{=} \{x \in X \mid (x, x) \in U \boxtimes_{X \times X} V\} \\ &= \{x \in X \mid (x, x) \in \{(u, v) \in X \times X \mid u \in U \text{ and } v \in V\}\} \\ &= U \cap V.\end{aligned}$$

This finishes the proof.

Item 6, The Dualisation Functor: This is a repetition of **Items 5** and **6** of **Definition 4.4.7.1.3** and is proved there. \square

Appendices

A Other Chapters

Preliminaries

- I. Introduction
2. A Guide to the Literature

Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

Relations

8. Relations
9. Constructions With Relations

10. Conditions on Relations

Categories

- II. Categories
12. Presheaves and the Yoneda Lemma

Monoidal Categories

13. Constructions With Monoidal Categories

Bicategories

14. Types of Morphisms in Bicategories

Extra Part

15. Notes

References

- [MSE 267365] J. B. *Show that the powerset partial order is a cartesian closed category*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/267365> (cit. on p. 142).
- [MSE 267469] Zhen Lin. *Show that the powerset partial order is a cartesian closed category*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/267469> (cit. on p. 106).
- [MSE 2719059] Vinny Chase. *$\mathcal{P}(X)$ with symmetric difference as addition as a vector space over \mathbb{Z}_2* . Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2719059> (cit. on p. 121).
- [Cie97] Krzysztof Ciesielski. *Set Theory for the Working Mathematician*. Vol. 39. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1997, pp. xii+236. ISBN: 0-521-59441-3; 0-521-59465-0. DOI: [10.1017/CB09781139173131](https://doi.org/10.1017/CB09781139173131). URL: <https://doi.org/10.1017/CB09781139173131> (cit. on p. 64).
- [nLa25] nLab Authors. *Interactions of Images and Pre-images with Unions and Intersections*. <https://ncatlab.org/nlab/show/interactions+of+images+and+pre-images+with+unions+and+intersections>. Oct. 2025 (cit. on p. 204).
- [Pro25a] Proof Wiki Contributors. *Cartesian Product Distributes Over Set Difference — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Set_Difference (cit. on p. 17).
- [Pro25b] Proof Wiki Contributors. *Cartesian Product Distributes Over Symmetric Difference — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Symmetric_Difference (cit. on p. 17).
- [Pro25c] Proof Wiki Contributors. *Cartesian Product Distributes Over Union — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Union (cit. on p. 17).

- [Pro25d] Proof Wiki Contributors. *Cartesian Product of Intersections* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Cartesian_Product_of_Intersections (cit. on p. 17).
- [Pro25e] Proof Wiki Contributors. *Characteristic Function of Intersection* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Intersection (cit. on p. 107).
- [Pro25f] Proof Wiki Contributors. *Characteristic Function of Set Difference* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Set_Difference (cit. on p. 111).
- [Pro25g] Proof Wiki Contributors. *Characteristic Function of Symmetric Difference* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Symmetric_Difference (cit. on p. 121).
- [Pro25h] Proof Wiki Contributors. *Characteristic Function of Union* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Union (cit. on p. 101).
- [Pro25i] Proof Wiki Contributors. *Complement of Complement* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Complement_of_Complement (cit. on p. 114).
- [Pro25j] Proof Wiki Contributors. *Complement of Preimage equals Preimage of Complement* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Complement_of_Preimage_equals_Preimage_of_Complement (cit. on p. 188).
- [Pro25k] Proof Wiki Contributors. *De Morgan's Laws (Set Theory)* — *Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/De_Morgan%27s_Laws_\(Set_Theory\)](https://proofwiki.org/wiki/De_Morgan%27s_Laws_(Set_Theory)) (cit. on pp. 111, 114).
- [Pro25l] Proof Wiki Contributors. *De Morgan's Laws (Set Theory)/Set Difference/Difference with Union* — *Proof Wiki*. 2025. URL: [https://proofwiki.org/wiki/De_Morgan%27s_Laws_\(Set_Theory\)/Set_Difference/Difference_with_Union](https://proofwiki.org/wiki/De_Morgan%27s_Laws_(Set_Theory)/Set_Difference/Difference_with_Union) (cit. on p. 111).

- [Pro25m] Proof Wiki Contributors. *Equivalence of Definitions of Symmetric Difference* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Equivalence_of_Definitions_of_Symmetric_Difference (cit. on p. 120).
- [Pro25n] Proof Wiki Contributors. *Image of Intersection Under Mapping* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Image_of_Intersection_under_Mapping (cit. on pp. 107, 179, 180).
- [Pro25o] Proof Wiki Contributors. *Image of Set Difference Under Mapping* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Image_of_Set_Difference_under_Mapping (cit. on pp. 111, 179).
- [Pro25p] Proof Wiki Contributors. *Image of Union Under Mapping* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Image_of_Union_under_Mapping (cit. on pp. 101, 179, 180).
- [Pro25q] Proof Wiki Contributors. *Intersection Distributes Over Symmetric Difference* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_Distributes_over_Symmetric_Difference (cit. on p. 121).
- [Pro25r] Proof Wiki Contributors. *Intersection Is Associative* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_is_Associative (cit. on p. 106).
- [Pro25s] Proof Wiki Contributors. *Intersection Is Commutative* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_is_Commutative (cit. on p. 107).
- [Pro25t] Proof Wiki Contributors. *Intersection With Empty Set* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_with_Empty_Set (cit. on p. 107).
- [Pro25u] Proof Wiki Contributors. *Intersection With Set Difference Is Set Difference With Intersection* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_with_Set_Difference_is_Set_Difference_with_Intersection (cit. on p. 111).

- [Pro25v] Proof Wiki Contributors. *Intersection With Subset Is Subset* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_with_Subset_is_Subset (cit. on p. 107).
- [Pro25w] Proof Wiki Contributors. *Preimage of Intersection Under Mapping* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Preimage_of_Intersection_under_Mapping (cit. on pp. 107, 188, 189).
- [Pro25x] Proof Wiki Contributors. *Preimage of Set Difference Under Mapping* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Preimage_of_Set_Difference_under_Mapping (cit. on pp. 111, 188).
- [Pro25y] Proof Wiki Contributors. *Preimage of Union Under Mapping* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Preimage_of_Union_under_Mapping (cit. on pp. 101, 188, 189).
- [Pro25z] Proof Wiki Contributors. *Quotient Mapping Is Coequalizer* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Quotient_Mapping_is_Coequalizer (cit. on p. 56).
- [Pro25aa] Proof Wiki Contributors. *Set Difference as Intersection With Complement* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_as_Intersection_with_Complement (cit. on p. 111).
- [Pro25ab] Proof Wiki Contributors. *Set Difference as Symmetric Difference With Intersection* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_as_Symmetric_Difference_with_Intersection (cit. on p. 111).
- [Pro25ac] Proof Wiki Contributors. *Set Difference Is Right Distributive Over Union* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_is_Right_Distributive_over_Union (cit. on p. 111).
- [Pro25ad] Proof Wiki Contributors. *Set Difference Over Subset* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_over_Subset (cit. on p. 111).

- [Pro25ae] Proof Wiki Contributors. *Set Difference With Empty Set Is Self* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_with_Empty_Set_is_Self (cit. on p. III).
- [Pro25af] Proof Wiki Contributors. *Set Difference With Self Is Empty Set* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_with_Self_is_Empty_Set (cit. on p. III).
- [Pro25ag] Proof Wiki Contributors. *Set Difference With Set Difference Is Union of Set Difference With Intersection* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_with_Set_Difference_is_Union_of_Set_Difference_with_Intersection (cit. on p. III).
- [Pro25ah] Proof Wiki Contributors. *Set Difference With Subset Is Superset of Set Difference* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_with_Subset_is_Superset_of_Set_Difference (cit. on p. III).
- [Pro25ai] Proof Wiki Contributors. *Set Difference With Union* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Difference_with_Union (cit. on p. III).
- [Pro25aj] Proof Wiki Contributors. *Set Intersection Distributes Over Union* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Intersection_Distributes_over_Union (cit. on pp. 100, 107).
- [Pro25ak] Proof Wiki Contributors. *Set Intersection Is Idempotent* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Intersection_is_Idempotent (cit. on p. 107).
- [Pro25al] Proof Wiki Contributors. *Set Intersection Preserves Subsets* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Intersection_Preserves_Subsets (cit. on p. 106).
- [Pro25am] Proof Wiki Contributors. *Set Union Is Idempotent* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Union_is_Idempotent (cit. on p. 100).

- [Pro25an] Proof Wiki Contributors. *Set Union Preserves Subsets — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Set_Union_Preserves_Subsets (cit. on p. 100).
- [Pro25ao] Proof Wiki Contributors. *Symmetric Difference Is Associative — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_is_Associative (cit. on p. 121).
- [Pro25ap] Proof Wiki Contributors. *Symmetric Difference Is Commutative — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_is_Commutative (cit. on p. 121).
- [Pro25aq] Proof Wiki Contributors. *Symmetric Difference of Complements — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_of_Complements (cit. on p. 121).
- [Pro25ar] Proof Wiki Contributors. *Symmetric Difference on Power Set Forms Abelian Group — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_on_Power_Set_forms_Abelian_Group (cit. on p. 121).
- [Pro25as] Proof Wiki Contributors. *Symmetric Difference With Complement — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Complement (cit. on p. 121).
- [Pro25at] Proof Wiki Contributors. *Symmetric Difference With Empty Set — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Empty_Set (cit. on p. 121).
- [Pro25au] Proof Wiki Contributors. *Symmetric Difference With Intersection Forms Ring — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Intersection_forms_Ring (cit. on p. 121).
- [Pro25av] Proof Wiki Contributors. *Symmetric Difference With Self Is Empty Set — Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Self_is_Empty_Set (cit. on p. 121).

- [Pro25aw] Proof Wiki Contributors. *Symmetric Difference With Union Does Not Form Ring* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Union_does_not_form_Ring (cit. on p. 119).
- [Pro25ax] Proof Wiki Contributors. *Symmetric Difference With Universe* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Universe (cit. on p. 121).
- [Pro25ay] Proof Wiki Contributors. *Union as Symmetric Difference With Intersection* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Union_as_Symmetric_Difference_with_Intersection (cit. on p. 101).
- [Pro25az] Proof Wiki Contributors. *Union Distributes Over Intersection* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Union_Distributes_over_Intersection (cit. on pp. 100, 107).
- [Pro25ba] Proof Wiki Contributors. *Union Is Associative* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Union_is_Associative (cit. on p. 100).
- [Pro25bb] Proof Wiki Contributors. *Union Is Commutative* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Union_is_Commutative (cit. on p. 100).
- [Pro25bc] Proof Wiki Contributors. *Union of Symmetric Differences* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Union_of_Symmetric_Differences (cit. on p. 121).
- [Pro25bd] Proof Wiki Contributors. *Union With Empty Set* — *Proof Wiki*. 2025. URL: https://proofwiki.org/wiki/Union_with_Empty_Set (cit. on p. 100).