

Conditions on Relations

The Clowder Project Authors

July 22, 2025

00TJ This chapter contains some material about reflexive, symmetric, transitive, equivalence, and apartness relations.

Contents

10.1	Functional and Total Relations	2
10.1.1	Functional Relations	2
10.1.2	Total Relations	3
10.2	Reflexive Relations	3
10.2.1	Foundations	3
10.2.2	The Reflexive Closure of a Relation.....	4
10.3	Symmetric Relations	6
10.3.1	Foundations	6
10.3.2	The Symmetric Closure of a Relation.....	7
10.4	Transitive Relations	8
10.4.1	Foundations	8
10.4.2	The Transitive Closure of a Relation	9
10.5	Equivalence Relations.....	12
10.5.1	Foundations	12
10.5.2	The Equivalence Closure of a Relation.....	12

10.6	Quotients by Equivalence Relations	14
10.6.1	Equivalence Classes	14
10.6.2	Quotients of Sets by Equivalence Relations.....	14
A	Other Chapters	19

02D0 10.1 Functional and Total Relations

00JC 10.1.1 Functional Relations

Let A and B be sets.

00JD **Definition 10.1.1.1.** A relation $R: A \rightarrow B$ is **functional** if, for each $a \in A$, the set $R(a)$ is either empty or a singleton.

00JE **Proposition 10.1.1.2.** Let $R: A \rightarrow B$ be a relation.

00JF 1. *Characterisations.* The following conditions are equivalent:

00JG (a) The relation R is functional.

00JH (b) We have $R \diamond R^\dagger \subset \chi_B$.

Proof. **Item 1, Characterisations:** We claim that **Items 1a** and **1b** are indeed equivalent:

- **Item 1a** \implies **Item 1b:** Let $(b, b') \in B \times B$. We need to show that

$$[R \diamond R^\dagger](b, b') \preceq_{\{t, f\}} \chi_B(b, b'),$$

i.e. that if there exists some $a \in A$ such that $b \sim_{R^\dagger} a$ and $a \sim_R b'$, then $b = b'$. But since $b \sim_{R^\dagger} a$ is the same as $a \sim_R b$, we have both $a \sim_R b$ and $a \sim_R b'$ at the same time, which implies $b = b'$ since R is functional.

- **Item 1b** \implies **Item 1a:** Suppose that we have $a \sim_R b$ and $a \sim_R b'$ for $b, b' \in B$. We claim that $b = b'$:

- Since $a \sim_R b$, we have $b \sim_{R^\dagger} a$.

- Since $R \diamond R^\dagger \subset \chi_B$, we have

$$[R \diamond R^\dagger](b, b') \preceq_{\{t, f\}} \chi_B(b, b'),$$

and since $b \sim_{R^\dagger} a$ and $a \sim_R b'$, it follows that $[R \diamond R^\dagger](b, b') = \text{true}$, and thus $\chi_B(b, b') = \text{true}$ as well, i.e. $b = b'$.

This finishes the proof. □

00JJ 10.1.2 Total Relations

Let A and B be sets.

00JK Definition 10.1.2.1.1. A relation $R: A \rightarrow B$ is **total** if, for each $a \in A$, we have $R(a) \neq \emptyset$.

00JL Proposition 10.1.2.1.2. Let $R: A \rightarrow B$ be a relation.

00JM 1. *Characterisations.* The following conditions are equivalent:

00JN (a) The relation R is total.

00JP (b) We have $\chi_A \subset R^\dagger \diamond R$.

Proof. **Item 1, Characterisations:** We claim that **Items 1a** and **1b** are indeed equivalent:

- **Item 1a** \implies **Item 1b:** We have to show that, for each $(a, a') \in A$, we have

$$\chi_A(a, a') \preceq_{\{t, f\}} [R^\dagger \diamond R](a, a'),$$

i.e. that if $a = a'$, then there exists some $b \in B$ such that $a \sim_R b$ and $b \sim_{R^\dagger} a'$ (i.e. $a \sim_R b$ again), which follows from the totality of R .

- **Item 1b** \implies **Item 1a:** Given $a \in A$, since $\chi_A \subset R^\dagger \diamond R$, we must have

$$\{a\} \subset [R^\dagger \diamond R](a),$$

implying that there must exist some $b \in B$ such that $a \sim_R b$ and $b \sim_{R^\dagger} a$ (i.e. $a \sim_R b$) and thus $R(a) \neq \emptyset$, as $b \in R(a)$.

This finishes the proof. □

00TK 10.2 Reflexive Relations

00TL 10.2.1 Foundations

Let A be a set.

00TM Definition 10.2.1.1.1. A **reflexive relation** is equivalently:¹

- An \mathbb{E}_0 -monoid in $(\mathbf{N}_\bullet(\mathbf{Rel}(A, A)), \chi_A)$.
- A pointed object in $(\mathbf{Rel}(A, A), \chi_A)$.

00TN Remark 10.2.1.1.2. In detail, a relation R on A is **reflexive** if we have an inclusion

$$\eta_R: \chi_A \subset R$$

of relations in $\mathbf{Rel}(A, A)$, i.e. if, for each $a \in A$, we have $a \sim_R a$.

00TP Definition 10.2.1.1.3. Let A be a set.

00TQ 1. The **set of reflexive relations on A** is the subset $\mathbf{Rel}^{\text{refl}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the reflexive relations.

00TR 2. The **poset of relations on A** is the subposet $\mathbf{Rel}^{\text{refl}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the reflexive relations.

00TS Proposition 10.2.1.1.4. Let R and S be relations on A .

00TT 1. *Interaction With Inverses.* If R is reflexive, then so is R^\dagger .

00TU 2. *Interaction With Composition.* If R and S are reflexive, then so is $S \diamond R$.

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: Clear. □

00TV 10.2.2 The Reflexive Closure of a Relation

Let R be a relation on A .

00TW Definition 10.2.2.1.1. The **reflexive closure** of \sim_R is the relation \sim_R^{refl} ² satisfying the following universal property:³

- (★) Given another reflexive relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{refl}} \subset \sim_S$.

¹Note that since $\mathbf{Rel}(A, A)$ is posetal, reflexivity is a property of a relation, rather than extra structure.

²*Further Notation:* Also written R^{refl} .

³*Slogan:* The reflexive closure of R is the smallest reflexive relation containing R .

00TX Construction 10.2.2.1.2. Concretely, \sim_R^{refl} is the free pointed object on R in $(\mathbf{Rel}(A, A), \chi_A)^4$, being given by

$$\begin{aligned} R^{\text{refl}} &\stackrel{\text{def}}{=} R \coprod^{\mathbf{Rel}(A, A)} \Delta_A \\ &= R \cup \Delta_A \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } a = b\}. \end{aligned}$$

Proof. Clear. □

00TY Proposition 10.2.2.1.3. Let R be a relation on A .

00TZ 1. *Adjointness.* We have an adjunction

$$\left((-)^{\text{refl}} \dashv \overline{\omega} \right): \quad \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{refl}}} \\ \perp \\ \xleftarrow{\overline{\omega}} \end{array} \mathbf{Rel}^{\text{refl}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{refl}}(R^{\text{refl}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{refl}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, A))$.

00U0 2. *The Reflexive Closure of a Reflexive Relation.* If R is reflexive, then $R^{\text{refl}} = R$.

00U1 3. *Idempotency.* We have

$$(R^{\text{refl}})^{\text{refl}} = R^{\text{refl}}.$$

00U2 4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} & \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} \mathbf{Rel}(A, A) \\ \left(R^\dagger \right)^{\text{refl}} = \left(R^{\text{refl}} \right)^\dagger, & \downarrow (-)^\dagger & \downarrow (-)^\dagger \\ & \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} \mathbf{Rel}(A, A). \end{array}$$

⁴Or, equivalently, the free \mathbb{E}_0 -monoid on R in $(\mathbf{N}_\bullet(\mathbf{Rel}(A, A)), \chi_A)$.

00U3 5. *Interaction With Composition.* We have

$$\begin{array}{ccc}
 \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\
 (S \diamond R)^{\text{refl}} = S^{\text{refl}} \diamond R^{\text{refl}}, & \begin{array}{c} (-)^{\text{refl}} \times (-)^{\text{refl}} \downarrow \\ \text{Rel}(A, A) \times \text{Rel}(A, A) \end{array} & \begin{array}{c} \downarrow (-)^{\text{refl}} \\ \text{Rel}(A, A) \end{array} \\
 & \xrightarrow{\diamond} & \text{Rel}(A, A).
 \end{array}$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the reflexive closure of a relation, stated in **Definition 10.2.2.1.1**.

Item 2, The Reflexive Closure of a Reflexive Relation: Clear.

Item 3, Idempotency: This follows from **Item 2**.

Item 4, Interaction With Inverses: Clear.

Item 5, Interaction With Composition: This follows from **Item 2** of **Definition 10.2.1.1.4**.

□

00U4 10.3 Symmetric Relations

00U5 10.3.1 Foundations

Let A be a set.

00U6 **Definition 10.3.1.1.1.** A relation R on A is **symmetric** if we have $R^\dagger = R$.

00U7 **Remark 10.3.1.1.2.** In detail, a relation R is symmetric if it satisfies the following condition:

(★) For each $a, b \in A$, if $a \sim_R b$, then $b \sim_R a$.

00U8 **Definition 10.3.1.1.3.** Let A be a set.

00U9 1. The **set of symmetric relations on A** is the subset $\text{Rel}^{\text{symm}}(A, A)$ of $\text{Rel}(A, A)$ spanned by the symmetric relations.

00UA 2. The **poset of relations on A** is the subposet $\text{Rel}^{\text{symm}}(A, A)$ of $\text{Rel}(A, A)$ spanned by the symmetric relations.

00UB **Proposition 10.3.1.1.4.** Let R and S be relations on A .

00UC 1. *Interaction With Inverses.* If R is symmetric, then so is R^\dagger .

00UD 2. *Interaction With Composition.* If R and S are symmetric, then so is $S \diamond R$.

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: Clear.

□

00UE 10.3.2 The Symmetric Closure of a Relation

Let R be a relation on A .

00UF **Definition 10.3.2.1.1.** The **symmetric closure** of \sim_R is the relation \sim_R^{symm} ⁵ satisfying the following universal property:⁶

(★) Given another symmetric relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{symm}} \subset \sim_S$.

00UG **Construction 10.3.2.1.2.** Concretely, \sim_R^{symm} is the symmetric relation on A defined by

$$\begin{aligned} R^{\text{symm}} &\stackrel{\text{def}}{=} R \cup R^\dagger \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } b \sim_R a\}. \end{aligned}$$

Proof. Clear. □

00UH **Proposition 10.3.2.1.3.** Let R be a relation on A .

00UJ 1. *Adjointness.* We have an adjunction

$$((-)^{\text{symm}} \dashv \overline{}): \quad \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{symm}}} \\ \perp \\ \xleftarrow{\overline{}} \end{array} \mathbf{Rel}^{\text{symm}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{symm}}(R^{\text{symm}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{symm}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, A))$.

00UK 2. *The Symmetric Closure of a Symmetric Relation.* If R is symmetric, then $R^{\text{symm}} = R$.

00UL 3. *Idempotency.* We have

$$(R^{\text{symm}})^{\text{symm}} = R^{\text{symm}}.$$

⁵Further Notation: Also written R^{symm} .

⁶Slogan: The symmetric closure of R is the smallest symmetric relation containing R .

00UM 4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} & \text{Rel}(A, A) & \xrightarrow{(-)^{\text{symm}}} \text{Rel}(A, A) \\ \left(R^{\dagger}\right)^{\text{symm}} = \left(R^{\text{symm}}\right)^{\dagger}, & \downarrow (-)^{\dagger} & \downarrow (-)^{\dagger} \\ & \text{Rel}(A, A) & \xrightarrow{(-)^{\text{symm}}} \text{Rel}(A, A). \end{array}$$

00UN 5. *Interaction With Composition.* We have

$$\begin{array}{ccc} & \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} \text{Rel}(A, A) \\ (S \diamond R)^{\text{symm}} = S^{\text{symm}} \diamond R^{\text{symm}}, & \downarrow (-)^{\text{symm}} \times (-)^{\text{symm}} & \downarrow (-)^{\text{symm}} \\ & \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} \text{Rel}(A, A). \end{array}$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the symmetric closure of a relation, stated in **Definition 10.3.2.1.1**.

Item 2, The Symmetric Closure of a Symmetric Relation: Clear.

Item 3, Idempotency: This follows from **Item 2**.

Item 4, Interaction With Inverses: Clear.

Item 5, Interaction With Composition: This follows from **Item 2** of **Definition 10.3.1.1.4**.

□

00UP 10.4 Transitive Relations

00UQ 10.4.1 Foundations

Let A be a set.

00UR **Definition 10.4.1.1.1.** A **transitive relation** is equivalently:⁷

- A non-unital \mathbb{E}_1 -monoid in $(\mathbf{N}_{\bullet}(\mathbf{Rel}(A, A)), \diamond)$.
- A non-unital monoid in $(\mathbf{Rel}(A, A), \diamond)$.

⁷Note that since $\mathbf{Rel}(A, A)$ is posetal, transitivity is a property of a relation, rather than extra structure.

00US Remark 10.4.1.1.2. In detail, a relation R on A is **transitive** if we have an inclusion

$$\mu_R: R \diamond R \subset R$$

of relations in $\text{Rel}(A, A)$, i.e. if, for each $a, c \in A$, the following condition is satisfied:

(★) If there exists some $b \in A$ such that $a \sim_R b$ and $b \sim_R c$, then $a \sim_R c$.

00UT Definition 10.4.1.1.3. Let A be a set.

- 00UU** 1. The **set of transitive relations from A to B** is the subset $\text{Rel}^{\text{trans}}(A)$ of $\text{Rel}(A, A)$ spanned by the transitive relations.
- 00UV** 2. The **poset of relations from A to B** is the subposet $\text{Rel}^{\text{trans}}(A)$ of $\text{Rel}(A, A)$ spanned by the transitive relations.

00UW Proposition 10.4.1.1.4. Let R and S be relations on A .

- 00UX** 1. *Interaction With Inverses.* If R is transitive, then so is R^\dagger .
- 00UY** 2. *Interaction With Composition.* If R and S are transitive, then $S \diamond R$ **may fail to be transitive**.

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: See [MSE 2096272].⁸

□

00UZ 10.4.2 The Transitive Closure of a Relation

Let R be a relation on A .

⁸*Intuition:* Transitivity for R and S fails to imply that of $S \diamond R$ because the composition operation for relations intertwines R and S in an incompatible way:

- If $a \sim_{S \diamond R} c$ and $c \sim_{S \diamond R} e$, then:
 - There is some $b \in A$ such that:
 - * $a \sim_R b$;
 - * $b \sim_S c$;
 - There is some $d \in A$ such that:
 - * $c \sim_R d$;
 - * $d \sim_S e$.

00V0 Definition 10.4.2.1.1. The **transitive closure** of \sim_R is the relation \sim_R^{trans} ⁹ satisfying the following universal property:¹⁰

(★) Given another transitive relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{trans}} \subset \sim_S$.

00V1 Construction 10.4.2.1.2. Concretely, \sim_R^{trans} is the free non-unital monoid on R in $(\mathbf{Rel}(A, A), \diamond)$ ¹¹, being given by

$$\begin{aligned} R^{\text{trans}} &\stackrel{\text{def}}{=} \prod_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \bigcup_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists some } (x_1, \dots, x_n) \in R^{\times n} \\ \text{such that } a \sim_R x_1 \sim_R \dots \sim_R x_n \sim_R b \end{array} \right\}. \end{aligned}$$

Proof. Clear. □

00V2 Proposition 10.4.2.1.3. Let R be a relation on A .

00V3 1. *Adjointness.* We have an adjunction

$$((-)^{\text{trans}} \dashv \overline{}): \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{trans}}} \\ \perp \\ \xleftarrow{\overline{}} \end{array} \mathbf{Rel}^{\text{trans}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{trans}}(R^{\text{trans}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \mathbf{Obj}(\mathbf{Rel}^{\text{trans}}(A, A))$ and $S \in \mathbf{Obj}(\mathbf{Rel}(A, B))$.

00V4 2. *The Transitive Closure of a Transitive Relation.* If R is transitive, then $R^{\text{trans}} = R$.

00V5 3. *Idempotency.* We have

$$(R^{\text{trans}})^{\text{trans}} = R^{\text{trans}}.$$

⁹Further Notation: Also written R^{trans} .

¹⁰Slogan: The transitive closure of R is the smallest transitive relation containing R .

¹¹Or, equivalently, the free non-unital \mathbb{B}_1 -monoid on R in $(\mathbf{N}_\bullet(\mathbf{Rel}(A, A)), \diamond)$.

00V6 4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A) \\ (-)^{\dagger} \downarrow & & \downarrow (-)^{\dagger} \\ \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A). \end{array}$$

$$(R^{\dagger})^{\text{trans}} = (R^{\text{trans}})^{\dagger},$$

00V7 5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\ (-)^{\text{trans}} \times (-)^{\text{trans}} \downarrow & \text{X} & \downarrow (-)^{\text{trans}} \\ \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A). \end{array}$$

$$(S \diamond R)^{\text{trans}} \stackrel{\text{poss.}}{\neq} S^{\text{trans}} \diamond R^{\text{trans}},$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the transitive closure of a relation, stated in **Definition 10.4.2.1.1**.

Item 2, The Transitive Closure of a Transitive Relation: Clear.

Item 3, Idempotency: This follows from **Item 2**.

Item 4, Interaction With Inverses: We have

$$\begin{aligned} (R^{\dagger})^{\text{trans}} &= \bigcup_{n=1}^{\infty} (R^{\dagger})^{\diamond n} \\ &= \bigcup_{n=1}^{\infty} (R^{\diamond n})^{\dagger} \\ &= \left(\bigcup_{n=1}^{\infty} R^{\diamond n} \right)^{\dagger} \\ &= (R^{\text{trans}})^{\dagger}, \end{aligned}$$

where we have used, respectively:

- **Definition 10.4.2.1.2.**
- **Constructions With Relations**, ?? of ??.
- **Constructions With Relations**, ?? of **Definition 9.2.3.1.2**.

- **Definition 10.4.2.1.2.**

This finishes the proof.

Item 5, Interaction With Composition: This follows from **Item 2** of **Definition 10.4.1.1.4.**

□

00V8 10.5 Equivalence Relations

00V9 10.5.1 Foundations

Let A be a set.

00VA **Definition 10.5.1.1.1.** A relation R is an **equivalence relation** if it is reflexive, symmetric, and transitive.¹²

00VB **Example 10.5.1.1.2.** The **kernel of a function** $f: A \rightarrow B$ is the equivalence relation $\sim_{\text{Ker}(f)}$ on A obtained by declaring $a \sim_{\text{Ker}(f)} b$ iff $f(a) = f(b)$.¹³

00VC **Definition 10.5.1.1.3.** Let A and B be sets.

00VD 1. The **set of equivalence relations from A to B** is the subset $\text{Rel}^{\text{eq}}(A, B)$ of $\text{Rel}(A, B)$ spanned by the equivalence relations.

00VE 2. The **poset of relations from A to B** is the subposet $\text{Rel}^{\text{eq}}(A, B)$ of $\text{Rel}(A, B)$ spanned by the equivalence relations.

00VF 10.5.2 The Equivalence Closure of a Relation

Let R be a relation on A .

00VG **Definition 10.5.2.1.1.** The **equivalence closure**¹⁴ of \sim_R is the relation \sim_R^{eq} ¹⁵ satisfying the following universal property:¹⁶

¹²*Further Terminology:* If instead R is just symmetric and transitive, then it is called a **partial equivalence relation**.

¹³The kernel $\text{Ker}(f): A \dashv A$ of f is the underlying functor of the monad induced by the adjunction $\text{Gr}(f) \dashv f^{-1}: A \rightleftarrows B$ in **Rel** of **Constructions With Relations**, ?? of ??.

¹⁴*Further Terminology:* Also called the **equivalence relation associated to \sim_R** .

¹⁵*Further Notation:* Also written R^{eq} .

¹⁶*Slogan:* The equivalence closure of R is the smallest equivalence relation containing R .

(★) Given another equivalence relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{eq}} \subset \sim_S$.

00VH Construction 10.5.2.1.2. Concretely, \sim_R^{eq} is the equivalence relation on A defined by

$$\begin{aligned} R^{\text{eq}} &\stackrel{\text{def}}{=} \left((R^{\text{refl}})^{\text{symm}} \right)^{\text{trans}} \\ &= \left((R^{\text{symm}})^{\text{trans}} \right)^{\text{refl}} \\ &= \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists } (x_1, \dots, x_n) \in R^{\times n} \text{ satisfying at} \\ \text{least one of the following conditions:} \\ \\ 1. \text{ The following conditions are satisfied:} \\ \\ \quad (a) \text{ We have } a \sim_R x_1 \text{ or } x_1 \sim_R a; \\ \quad (b) \text{ We have } x_i \sim_R x_{i+1} \text{ or } x_{i+1} \sim_R x_i \\ \quad \quad \text{for each } 1 \leq i \leq n-1; \\ \quad (c) \text{ We have } b \sim_R x_n \text{ or } x_n \sim_R b; \\ \\ 2. \text{ We have } a = b. \end{array} \right\}. \end{aligned}$$

Proof. From the universal properties of the reflexive, symmetric, and transitive closures of a relation (**Definitions 10.2.2.1.1, 10.3.2.1.1 and 10.4.2.1.1**), we see that it suffices to prove that:

00VJ 1. The symmetric closure of a reflexive relation is still reflexive.

00VK 2. The transitive closure of a symmetric relation is still symmetric.

which are both clear. □

00VL Proposition 10.5.2.1.3. Let R be a relation on A .

00VM 1. *Adjointness.* We have an adjunction

$$((-)^{\text{eq}} \dashv \overline{}): \mathbf{Rel}(A, B) \begin{array}{c} \xrightarrow{(-)^{\text{eq}}} \\ \perp \\ \xleftarrow{\overline{}} \end{array} \mathbf{Rel}^{\text{eq}}(A, B),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{eq}}(R^{\text{eq}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{eq}}(A, B))$ and $S \in \text{Obj}(\mathbf{Rel}(A, B))$.

00VN 2. *The Equivalence Closure of an Equivalence Relation.* If R is an equivalence relation, then $R^{\text{eq}} = R$.

00VP 3. *Idempotency.* We have

$$(R^{\text{eq}})^{\text{eq}} = R^{\text{eq}}.$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the equivalence closure of a relation, stated in **Definition 10.5.2.1.1**.

Item 2, The Equivalence Closure of an Equivalence Relation: Clear.

Item 3, Idempotency: This follows from **Item 2**. □

00VQ 10.6 Quotients by Equivalence Relations

00VR 10.6.1 Equivalence Classes

Let A be a set, let R be a relation on A , and let $a \in A$.

02B1 **Definition 10.6.1.1.1.** The **equivalence class associated to** a is the set $[a]$ defined by

$$\begin{aligned} [a] &\stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\} \\ &= \{x \in X \mid a \sim_R x\}. \end{aligned} \quad (\text{since } R \text{ is symmetric})$$

02B2 10.6.2 Quotients of Sets by Equivalence Relations

Let A be a set and let R be a relation on A .

02B3 **Definition 10.6.2.1.1.** The **quotient of** X **by** R is the set X/\sim_R defined by

$$X/\sim_R \stackrel{\text{def}}{=} \{[a] \in \mathcal{P}(X) \mid a \in X\}.$$

00VV **Remark 10.6.2.1.2.** The reason we define quotient sets for equivalence relations only is that each of the properties of being an equivalence relation—reflexivity, symmetry, and transitivity—ensures that the equivalence classes $[a]$ of X under R are well-behaved:

- *Reflexivity.* If R is reflexive, then, for each $a \in X$, we have $a \in [a]$.

- *Symmetry*. The equivalence class $[a]$ of an element a of X is defined by

$$[a] \stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\},$$

but we could equally well define

$$[a]' \stackrel{\text{def}}{=} \{x \in X \mid a \sim_R x\}$$

instead. This is not a problem when R is symmetric, as we then have $[a] = [a]'$.¹⁷

- *Transitivity*. If R is transitive, then $[a]$ and $[b]$ are disjoint iff $a \not\sim_R b$, and equal otherwise.

00VW Proposition 10.6.2.1.3. Let $f: X \rightarrow Y$ be a function and let R be a relation on X .

- 02B4** 1. *As a Coequaliser*. We have an isomorphism of sets

$$X/\sim_R^{\text{eq}} \cong \text{CoEq}\left(R \hookrightarrow X \times X \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} \end{array} X\right),$$

where \sim_R^{eq} is the equivalence relation generated by \sim_R .

- 02B5** 2. *As a Pushout*. We have an isomorphism of sets¹⁸

$$X/\sim_R^{\text{eq}} \cong X \amalg_{\text{Eq}(\text{pr}_1, \text{pr}_2)} X, \quad \begin{array}{ccc} X/\sim_R^{\text{eq}} & \longleftarrow & X \\ \uparrow \ulcorner & & \uparrow \\ X & \longleftarrow & \text{Eq}(\text{pr}_1, \text{pr}_2) \end{array}$$

where \sim_R^{eq} is the equivalence relation generated by \sim_R .

¹⁷When categorifying equivalence relations, one finds that $[a]$ and $[a]'$ correspond to presheaves and copresheaves; see Constructions With Categories, ??.

¹⁸Dually, we also have an isomorphism of sets

$$\text{Eq}(\text{pr}_1, \text{pr}_2) \cong X \times_{X/\sim_R^{\text{eq}}} X, \quad \begin{array}{ccc} \text{Eq}(\text{pr}_1, \text{pr}_2) & \longrightarrow & X \\ \downarrow \lrcorner & & \downarrow \\ X & \longrightarrow & X/\sim_R^{\text{eq}} \end{array}$$

02B6 3. *The First Isomorphism Theorem for Sets*. We have an isomorphism of sets^{19,20}

$$X/\sim_{\text{Ker}(f)} \cong \text{Im}(f).$$

00W0 4. *Descending Functions to Quotient Sets, I*. Let R be an equivalence relation on X . The following conditions are equivalent:

02B7 (a) There exists a map

$$\bar{f}: X/\sim_R \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \exists \nearrow \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

02B8 (b) We have $R \subset \text{Ker}(f)$.

02B9 (c) For each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$.

¹⁹*Further Terminology*: The set $X/\sim_{\text{Ker}(f)}$ is often called the **coimage** of f , and denoted by $\text{CoIm}(f)$.

²⁰In a sense this is a result relating the monad in **Rel** induced by f with the comonad in **Rel** induced by f , as the kernel and image

$$\begin{aligned} \text{Ker}(f) &: X \multimap X, \\ \text{Im}(f) &\subset Y \end{aligned}$$

of f are the underlying functors of (respectively) the induced monad and comonad of the adjunction

$$(\text{Gr}(f) \dashv f^{-1}): \begin{array}{ccc} & \text{Gr}(f) & \\ & \downarrow \perp & \\ A & \xrightarrow{\quad} & B \\ & \uparrow \perp & \\ & f^{-1} & \end{array}$$

of **Constructions With Relations**, ?? of ??.

- 00W1 5. *Descending Functions to Quotient Sets, II.* Let R be an equivalence relation on X . If the conditions of **Item 4** hold, then \bar{f} is the *unique* map making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \nearrow \exists! \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

- 00W2 6. *Descending Functions to Quotient Sets, III.* Let R be an equivalence relation on X . We have a bijection

$$\text{Hom}_{\text{Sets}}(X/\sim_R, Y) \cong \text{Hom}_{\text{Sets}}^R(X, Y),$$

natural in $X, Y \in \text{Obj}(\text{Sets})$, given by the assignment $f \mapsto \bar{f}$ of **Items 4** and **5**, where $\text{Hom}_{\text{Sets}}^R(X, Y)$ is the set defined by

$$\text{Hom}_{\text{Sets}}^R(X, Y) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(X, Y) \left| \begin{array}{l} \text{for each } x, y \in X, \\ \text{if } x \sim_R y, \text{ then} \\ f(x) = f(y) \end{array} \right. \right\}.$$

- 02BA 7. *Descending Functions to Quotient Sets, IV.* Let R be an equivalence relation on X . If the conditions of **Item 4** hold, then the following conditions are equivalent:

02BB (a) The map \bar{f} is an injection.

02BC (b) We have $R = \text{Ker}(f)$.

02BD (c) For each $x, y \in X$, we have $x \sim_R y$ iff $f(x) = f(y)$.

- 02BE 8. *Descending Functions to Quotient Sets, V.* Let R be an equivalence relation on X . If the conditions of **Item 4** hold, then the following conditions are equivalent:

02BF (a) The map $f: X \rightarrow Y$ is surjective.

02BG (b) The map $\bar{f}: X/\sim_R \rightarrow Y$ is surjective.

02BH 9. *Descending Functions to Quotient Sets, VI.* Let R be a relation on X and let \sim_R^{eq} be the equivalence relation associated to R . The following conditions are equivalent:

02BJ (a) The map f satisfies the equivalent conditions of **Item 4**:

- There exists a map

$$\bar{f}: X/\sim_R^{\text{eq}} \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \searrow \bar{f} & \uparrow \exists \\ X/\sim_R^{\text{eq}} & & \end{array}$$

commute.

- For each $x, y \in X$, if $x \sim_R^{\text{eq}} y$, then $f(x) = f(y)$.

02BK (b) For each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$.

Proof. **Item 1**, As a Coequaliser: Omitted.

Item 2, As a Pushout: Omitted.

Item 3, The First Isomorphism Theorem for Sets: Clear.

Item 4, Descending Functions to Quotient Sets, I: See [Pro25c].

Item 5, Descending Functions to Quotient Sets, II: See [Pro25d].

Item 6, Descending Functions to Quotient Sets, III: This follows from **Items 5** and **6**.

Item 7, Descending Functions to Quotient Sets, IV: See [Pro25b].

Item 8, Descending Functions to Quotient Sets, V: See [Pro25a].

Item 9, Descending Functions to Quotient Sets, VI: The implication **Item 8a** \implies **Item 8b** is clear.

Conversely, suppose that, for each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$. Spelling out the definition of the equivalence closure of R , we see that the condition $x \sim_R^{\text{eq}} y$ unwinds to the following:

- (★) There exist $(x_1, \dots, x_n) \in R^{\times n}$ satisfying at least one of the following conditions:
 - The following conditions are satisfied:

- * We have $x \sim_R x_1$ or $x_1 \sim_R x$;
- * We have $x_i \sim_R x_{i+1}$ or $x_{i+1} \sim_R x_i$ for each $1 \leq i \leq n - 1$;
- * We have $y \sim_R x_n$ or $x_n \sim_R y$;
- We have $x = y$.

Now, if $x = y$, then $f(x) = f(y)$ trivially; otherwise, we have

$$\begin{aligned}
 f(x) &= f(x_1), \\
 f(x_1) &= f(x_2), \\
 &\vdots \\
 f(x_{n-1}) &= f(x_n), \\
 f(x_n) &= f(y),
 \end{aligned}$$

and $f(x) = f(y)$, as we wanted to show. □

Appendices

A Other Chapters

Preliminaries

1. Introduction
2. A Guide to the Literature

Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

Relations

8. Relations
9. Constructions With Relations
10. Conditions on Relations

Categories

11. Categories
12. Presheaves and the Yoneda Lemma

Monoidal Categories

13. Constructions With Monoidal Categories

Bicategories

14. **Types of Morphisms in Bicat-
egories**

Extra Part

15. **Notes**

References

- [MSE 2096272] **Akiva Weinberger**. *Is composition of two transitive relations transitive? If not, can you give me a counterexample?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2096272> (cit. on p. 9).
- [Pro25a] Proof Wiki Contributors. *Condition For Mapping from Quotient Set To Be A Surjection* — Proof Wiki. 2025. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Surjection (cit. on p. 18).
- [Pro25b] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be An Injection* — Proof Wiki. 2025. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Injection (cit. on p. 18).
- [Pro25c] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be Well-Defined* — Proof Wiki. 2025. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Well-Defined (cit. on p. 18).
- [Pro25d] Proof Wiki Contributors. *Mapping From Quotient Set When Defined Is Unique* — Proof Wiki. 2025. URL: https://proofwiki.org/wiki/Mapping_from_Quotient_Set_when_Defined_is_Unique (cit. on p. 18).