Types of Morphisms in Bicategories

The Clowder Project Authors

July 12, 2025

019H In this chapter, we study special kinds of morphisms in bicategories:

1. *Monomorphisms and Epimorphisms in Bicategories (Sections 13.1 and 13.2).* There is a large number of different notions capturing the idea of a "monomorphism" or of an "epimorphism" in a bicategory.

   Arguably, the notion that best captures these concepts is that of a *pseudomonic morphism* (Definition 13.1.10.1.1) and of a *pseudoepic morphism* (Definition 13.2.10.1.1), although the other notions introduced in Sections 13.1 and 13.2 are also interesting on their own.

Contents

13.1 Monomorphisms in Bicategories 2
13.1.1 Representably Faithful Morphisms 2
13.1.2 Representably Full Morphisms 3
13.1.3 Representably Fully Faithful Morphisms 4
13.1.4 Morphisms Representably Faithful on Cores 6
13.1.5 Morphisms Representably Full on Cores 7
13.1.6 Morphisms Representably Fully Faithful on Cores 8
13.1.7 Representably Essentially Injective Morphisms 9
13.1.8 Representably Conservative Morphisms 10
13.1.9 Strict Monomorphisms 11
13.1.10 Pseudomonic Morphisms 11

13.2 Epimorphisms in Bicategories **16**
13.2.1 Corepresentably Faithful Morphisms 16
13.2.2 Corepresentably Full Morphisms 17
13.2.3 Corepresentably Fully Faithful Morphisms 19
13.2.4 Morphisms Corepresentably Faithful on Cores 21
13.2.5 Morphisms Corepresentably Full on Cores 21
13.2.6 Morphisms Corepresentably Fully Faithful on Cores 22
13.2.7 Corepresentably Essentially Injective Morphisms 24
13.2.8 Corepresentably Conservative Morphisms 24
13.2.9 Strict Epimorphisms 25
13.2.10 Pseudoepic Morphisms 26

A Other Chapters **28**

019J

13.1 Monomorphisms in Bicategories

019K

13.1.1 Representably Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

019L

Definition 13.1.1.1.1 ► Representably Faithful Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably faithful**[1] if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is faithful.

[1] *Further Terminology*: Also called simply a **faithful morphism**, based on Item 1 of Example 13.1.1.1.3.

019M

Remark 13.1.1.1.2 ► Unwinding Definition 13.1.1.1.1

In detail, f is representably faithful if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

019N **Example 13.1.1.1.3 ▶ Examples of Representably Faithful Morphisms**

Here are some examples of representably faithful morphisms.

019P 1. *Representably Faithful Morphisms in* Cats_2. The representably faithful morphisms in Cats_2 are precisely the faithful functors; see Categories, Item 2 of Proposition 11.6.1.1.2.

019Q 2. *Representably Faithful Morphisms in* **Rel**. Every morphism of **Rel** is representably faithful; see Relations, Item 1 of Proposition 8.4.8.1.1.

019R 13.1.2 Representably Full Morphisms

Let $\mathcal{C}$ be a bicategory.

019S **Definition 13.1.2.1.1 ▶ Representably Full Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably full**[1] if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is full.

[1] *Further Terminology:* Also called simply a **full morphism**, based on Item 1 of Example 13.1.2.1.3.

019T **Remark 13.1.2.1.2 ▶ Unwinding Definition 13.1.2.1.1**

In detail, f is representably full if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon f \circ \phi \Longrightarrow f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[from=U, to=D, Rightarrow, "\alpha"] \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[from=U, to=D, Rightarrow, "\alpha"] \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"{name=U}] \arrow[r, bend right, "f\circ\psi"'{name=D}] & B \arrow[from=U, to=D, Rightarrow, "\beta"] \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

019U **Example 13.1.2.1.3 ► Examples of Representably Full Morphisms**

Here are some examples of representably full morphisms.

019V 1. *Representably Full Morphisms in* Cats_2. The representably full morphisms in Cats_2 are precisely the full functors; see Categories, **??** of Proposition 11.6.2.1.2.

019W 2. *Representably Full Morphisms in* **Rel**. The representably full morphisms in **Rel** are characterised in Relations, Item 2 of Proposition 8.4.8.1.1.

019X 13.1.3 Representably Fully Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

019Y **Definition 13.1.3.1.1 ► Representably Fully Faithful Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably fully faithful**[1] if the following equivalent conditions are satisfied:

019Z 1. The 1-morphism f is representably faithful (Definition 13.1.1.1.1) and representably full (Definition 13.1.2.1.1).

01A0

2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is fully faithful.

[1] *Further Terminology:* Also called simply a **fully faithful morphism**, based on Item 1 of Example 13.1.3.1.3.

01A1

Remark 13.1.3.1.2 ▶ Unwinding Representably Fully Faithful Morphisms

In detail, f is representably fully faithful if the conditions in Remark 13.1.1.1.2 and Remark 13.1.2.1.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[r, "f"] & B, \arrow[Rightarrow, from=U, to=D, "\alpha"'] \arrow[Rightarrow, from=U, to=D, "\beta"] \end{tikzcd}$$

if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon f \circ \phi \Longrightarrow f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"{name=U}] \arrow[r, bend right, "f\circ\psi"'{name=D}] & B \arrow[Rightarrow, from=U, to=D, "\beta"'] \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[Rightarrow, from=U, to=D, "\alpha"'] \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[r, "f"] & B \arrow[Rightarrow, from=U, to=D, "\alpha"'] \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"{name=U}] \arrow[r, bend right, "f\circ\psi"'{name=D}] & B \arrow[Rightarrow, from=U, to=D, "\beta"'] \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

01A2 **Example 13.1.3.1.3 ▶ Examples of Representably Fully Faithful Morphisms**

Here are some examples of representably fully faithful morphisms.

01A3 1. *Representably Fully Faithful Morphisms in* Cats_2. The representably fully faithful morphisms in Cats_2 are precisely the fully faithful functors; see Categories, Item 6 of Proposition 11.6.3.1.2.

01A4 2. *Representably Fully Faithful Morphisms in* **Rel**. The representably fully faithful morphisms of **Rel** coincide (Relations, Item 3 of Proposition 8.4.8.1.1) with the representably full morphisms in **Rel**, which are characterised in Relations, Item 2 of Proposition 8.4.8.1.1.

01A5 ## 13.1.4 Morphisms Representably Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

01A6 **Definition 13.1.4.1.1 ▶ Morphisms Representably Faithful on Cores**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably faithful on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

given by postcomposition by f is faithful.

01A7 **Remark 13.1.4.1.2 ▶ Unwinding Definition 13.1.4.1.1**

In detail, f is representably faithful on cores if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

01A8 ## 13.1.5 Morphisms Representably Full on Cores

Let $\mathcal{C}$ be a bicategory.

01A9 **Definition 13.1.5.1.1 ▶ Morphisms Representably Full on Cores**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably full on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

given by postcomposition by f is full.

01AA **Remark 13.1.5.1.2 ▶ Unwinding Definition 13.1.5.1.1**

In detail, f is representably full on cores if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon f \circ \phi \overset{\sim}{\Longrightarrow} f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left=60, "f\circ\phi"{name=U}] \arrow[r, bend right=60, "f\circ\psi"'{name=D}] & B \arrow[Rightarrow, from=U, to=D, "\beta"] \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} X \arrow[r, bend left=60, "\phi"{name=U}] \arrow[r, bend right=60, "\psi"'{name=D}] & A \arrow[Rightarrow, from=U, to=D, "\alpha"] \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left=60, "\phi"{name=U}] \arrow[r, bend right=60, "\psi"'{name=D}] & A \arrow[Rightarrow, from=U, to=D, "\alpha"] \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left=60, "f\circ\phi"{name=U}] \arrow[r, bend right=60, "f\circ\psi"'{name=D}] & B \arrow[Rightarrow, from=U, to=D, "\beta"] \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

01AB

13.1.6 Morphisms Representably Fully Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

01AC

Definition 13.1.6.1.1 ▶ Morphisms Representably Fully Faithful on Cores

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably fully faithful on cores** if the following equivalent conditions are satisfied:

01AD 1. The 1-morphism f is representably faithful on cores (Definition 13.1.5.1.1) and representably full on cores (Definition 13.1.4.1.1).

01AE 2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

given by postcomposition by f is fully faithful.

01AF

Remark 13.1.6.1.2 ▶ Unwinding Definition 13.1.6.1.1

In detail, f is representably fully faithful on cores if the conditions in Remark 13.1.4.1.2 and Remark 13.1.5.1.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[r, "f"] & B, \arrow[from=U, to=D, Rightarrow, "\alpha"'] \arrow[from=U, to=D, Rightarrow, "\beta"] \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon f \circ \phi \overset{\sim}{\Longrightarrow} f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

01AG

13.1.7 Representably Essentially Injective Morphisms

Let $\mathcal{C}$ be a bicategory.

01AH

Definition 13.1.7.1.1 ► Representably Essentially Injective Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably essentially injective** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is essentially injective.

01AJ **Remark 13.1.7.1.2 ▶ Unwinding Definition 13.1.7.1.1**

In detail, f is representably essentially injective if, for each pair of morphisms $\phi, \psi \colon X \rightrightarrows A$ of $\mathcal{C}$, the following condition is satisfied:

($\star$) If $f \circ \phi \cong f \circ \psi$, then $\phi \cong \psi$.

01AK

13.1.8 Representably Conservative Morphisms

Let $\mathcal{C}$ be a bicategory.

01AL **Definition 13.1.8.1.1 ▶ Representably Conservative Morphisms**

A 1-morphism $f \colon A \to B$ of $\mathcal{C}$ is **representably conservative** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_* \colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is conservative.

01AM **Remark 13.1.8.1.2 ▶ Unwinding Definition 13.1.8.1.1**

In detail, f is representably conservative if, for each pair of morphisms $\phi, \psi \colon X \rightrightarrows A$ and each 2-morphism

$$\alpha \colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$, if the 2-morphism

$$\mathrm{id}_f \star \alpha \colon f \circ \phi \Longrightarrow f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\mathrm{id}_f \star \alpha \Downarrow"] & B \end{tikzcd}$$

is a 2-isomorphism, then so is α.

01AN 13.1.9 Strict Monomorphisms

Let $\mathcal{C}$ be a bicategory.

01AP **Definition 13.1.9.1.1 ▶ Strict Monomorphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is a **strict monomorphism** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is injective on objects, i.e. its action on objects

$$f_*\colon \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

is injective.

01AQ **Remark 13.1.9.1.2 ▶ Unwinding Definition 13.1.9.1.1**

In detail, f is a strict monomorphism in $\mathcal{C}$ if, for each diagram in $\mathcal{C}$ of the form

$$X \underset{\psi}{\overset{\phi}{\rightrightarrows}} A \xrightarrow{f} B,$$

if $f \circ \phi = f \circ \psi$, then $\phi = \psi$.

01AR **Example 13.1.9.1.3 ▶ Examples of Strict Monomorphisms**

Here are some examples of strict monomorphisms.

01AS 1. *Strict Monomorphisms in* Cats_2. The strict monomorphisms in Cats_2 are precisely the functors which are injective on objects and injective on morphisms; see Categories, Item 1 of Proposition 11.7.2.1.2.

01AT 2. *Strict Monomorphisms in* **Rel**. The strict monomorphisms in **Rel** are characterised in Relations, Proposition 8.4.7.1.1.

01AU 13.1.10 Pseudomonic Morphisms

Let $\mathcal{C}$ be a bicategory.

01AV

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **pseudomonic** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is pseudomonic.

01AW **Remark 13.1.10.1.2 ▶ Unwinding Definition 13.1.10.1.1**

In detail, a 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is pseudomonic if it satisfies the following conditions:

01AX 1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

01AY 2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon f \circ \phi \overset{\sim}{\Longrightarrow} f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

01AZ

Let $f\colon A \longrightarrow B$ be a 1-morphism of $\mathcal{C}$.

01B0 1. *Characterisations.* The following conditions are equivalent:

01B1 (a) The morphism f is pseudomonic.

01B2 (b) The morphism f is representably full on cores and representably faithful.

01B3 (c) We have an isocomma square of the form

$$A \overset{\text{eq.}}{\cong} A \overset{\leftrightarrow}{\times}_B A, \qquad \begin{tikzcd} A \arrow[r, "\mathrm{id}_A"] \arrow[d, "\mathrm{id}_A"'] & A \arrow[d, "F"] \\ A \arrow[r, "F"'] \arrow[ur, dashed, leftrightarrow] & B \end{tikzcd}$$

in $\mathcal{C}$ up to equivalence.

01B4 2. *Interaction With Cotensors.* If $\mathcal{C}$ has cotensors with $\mathbb{1}$, then the following conditions are equivalent:

(a) The morphism f is pseudomonic.

(b) We have an isocomma square of the form

$$A \overset{\text{eq.}}{\cong} A \overset{\leftrightarrow}{\times}_{\mathbb{1} \pitchfork F} B, \qquad \begin{tikzcd} A \arrow[r, hook] \arrow[d, "F"'] & \mathbb{1} \pitchfork A \arrow[d, "\mathbb{1} \pitchfork F"] \\ B \arrow[r, hook] \arrow[ur, dashed, leftrightarrow] & \mathbb{1} \pitchfork B \end{tikzcd}$$

in $\mathcal{C}$ up to equivalence.

Proof 13.1.10.1.4 ► Proof of Proposition 13.1.10.1.3

Item 1: Characterisations

Omitted.

Item 2: Interaction With Cotensors

Omitted.

01B5 ## 13.2 Epimorphisms in Bicategories

01B6 ### 13.2.1 Corepresentably Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

01B7 **Definition 13.2.1.1.1 ▶ Corepresentably Faithful Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably faithful** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is faithful.

01B8 **Remark 13.2.1.1.2 ▶ Unwinding Definition 13.2.1.1.1**

In detail, f is corepresentably faithful if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

01B9 **Example 13.2.1.1.3 ▶ Examples of Corepresentably Faithful Morphisms**

Here are some examples of corepresentably faithful morphisms.

01BA 1. *Corepresentably Faithful Morphisms in* Cats_2. The corepresentably faithful morphisms in Cats_2 are characterised in Categories, Item 5 of Proposition 11.6.1.1.2.

01BB 2. *Corepresentably Faithful Morphisms in* **Rel**. Every morphism of **Rel** is corepresentably faithful; see Relations, Item 1 of Proposition 8.4.10.1.1.

01BC 13.2.2 Corepresentably Full Morphisms

Let $\mathcal{C}$ be a bicategory.

01BD **Definition 13.2.2.1.1 ▶ Corepresentably Full Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably full** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is full.

01BE **Remark 13.2.2.1.2 ▶ Unwinding Definition 13.2.2.1.1**

In detail, f is corepresentably full if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon \phi \circ f \Longrightarrow \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f\,.$$

01BF

Here are some examples of corepresentably full morphisms.

01BG 1. *Corepresentably Full Morphisms in* Cats_2. The corepresentably full morphisms in Cats_2 are characterised in Categories, Item 7 of Proposition 11.6.2.1.2.

01BH 2. *Corepresentably Full Morphisms in* **Rel**. The corepresentably full morphisms in **Rel** are characterised in Relations, Item 2 of Proposition 8.4.10.1.1.

01BJ 13.2.3 Corepresentably Fully Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

01BK **Definition 13.2.3.1.1 ▶ Corepresentably Fully Faithful Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably fully faithful**[1] if the following equivalent conditions are satisfied:

01BL 1. The 1-morphism f is corepresentably full (Definition 13.2.2.1.1) and corepresentably faithful (Definition 13.2.1.1.1).

01BM 2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is fully faithful.

[1] *Further Terminology:* Corepresentably fully faithful morphisms have also been called **lax epimorphisms** in the literature (e.g. in [Adá+01]), though we will always use the name "corepresentably fully faithful morphism" instead in this work.

01BN **Remark 13.2.3.1.2 ▶ Unwinding Definition 13.2.3.1.1**

In detail, f is corepresentably fully faithful if the conditions in Remark 13.2.1.1.2 and Remark 13.2.2.1.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon \phi \circ f \Longrightarrow \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f\,.$$

01BP **Example 13.2.3.1.3 ▶ Examples of Corepresentably Fully Faithful Morphisms**

Here are some examples of corepresentably fully faithful morphisms.

01BQ 1. *Corepresentably Fully Faithful Morphisms in* Cats_2. The fully faithful epimorphisms in Cats_2 are characterised in Categories, Item 10 of Proposi-

tion 11.6.3.1.2.

01BR 2. *Corepresentably Fully Faithful Morphisms in* **Rel**. The corepresentably fully faithful morphisms of **Rel** coincide (Relations, Item 3 of Proposition 8.4.10.1.1) with the corepresentably full morphisms in **Rel**, which are characterised in Relations, Item 2 of Proposition 8.4.10.1.1.

01BS 13.2.4 Morphisms Corepresentably Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

01BT **Definition 13.2.4.1.1 ▶ Morphisms Corepresentably Faithful on Cores**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably faithful on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(B, X)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(A, X))$$

given by precomposition by f is faithful.

01BU **Remark 13.2.4.1.2 ▶ Unwinding Definition 13.2.4.1.1**

In detail, f is corepresentably faithful on cores if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

01BV 13.2.5 Morphisms Corepresentably Full on Cores

Let $\mathcal{C}$ be a bicategory.

01BW **Definition 13.2.5.1.1 ▶ Morphisms Corepresentably Full on Cores**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably full on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(B,X)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(A,X))$$

given by precomposition by f is full.

01BX **Remark 13.2.5.1.2 ▶ Unwinding Definition 13.2.5.1.1**

In detail, f is corepresentably full on cores if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon \phi \circ f \overset{\sim}{\Longrightarrow} \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left=60, "\phi\circ f"] \arrow[r, bend right=60, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left=60, "\phi"] \arrow[r, bend right=60, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left=60, "\phi"] \arrow[r, bend right=60, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left=60, "\phi\circ f"] \arrow[r, bend right=60, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f\,.$$

01BY

13.2.6 Morphisms Corepresentably Fully Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

01BZ **Definition 13.2.6.1.1 ▶ Morphisms Corepresentably Fully Faithful on Cores**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably fully faithful on cores** if the following equivalent conditions are satisfied:

01C0 1. The 1-morphism f is corepresentably full on cores (Definition 13.2.5.1.1) and corepresentably faithful on cores (Definition 13.2.1.1.1).

01C1 2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(B, X)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(A, X))$$

given by precomposition by f is fully faithful.

01C2 **Remark 13.2.6.1.2 ▶ Unwinding Definition 13.2.6.1.1**

In detail, f is corepresentably fully faithful on cores if the conditions in Remark 13.2.4.1.2 and Remark 13.2.5.1.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon \phi \circ f \overset{\sim}{\Longrightarrow} \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f\,.$$

01C3 **13.2.7 Corepresentably Essentially Injective Morphisms**

Let $\mathcal{C}$ be a bicategory.

01C4 **Definition 13.2.7.1.1 ▶ Corepresentably Essentially Injective Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably essentially injective** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is essentially injective.

01C5 **Remark 13.2.7.1.2 ▶ Unwinding Definition 13.2.7.1.1**

In detail, f is corepresentably essentially injective if, for each pair of morphisms $\phi, \psi\colon B \rightrightarrows X$ of $\mathcal{C}$, the following condition is satisfied:

($\star$) If $\phi \circ f \cong \psi \circ f$, then $\phi \cong \psi$.

01C6 **13.2.8 Corepresentably Conservative Morphisms**

Let $\mathcal{C}$ be a bicategory.

01C7 **Definition 13.2.8.1.1 ▶ Corepresentably Conservative Morphisms**

A 1-morphism $f\colon A \longrightarrow B$ of $\mathcal{C}$ is **corepresentably conservative** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is conservative.

01C8 **Remark 13.2.8.1.2 ▶ Unwinding Definition 13.2.8.1.1**

In detail, f is corepresentably conservative if, for each pair of morphisms $\phi, \psi\colon B \rightrightarrows X$ and each 2-morphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, if the 2-morphism

$$\alpha \star \mathrm{id}_f\colon \phi \circ f \Longrightarrow \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\alpha\star\mathrm{id}_f \Downarrow"] & X \end{tikzcd}$$

is a 2-isomorphism, then so is α.

01C9 ## 13.2.9 Strict Epimorphisms

Let $\mathcal{C}$ be a bicategory.

01CA **Definition 13.2.9.1.1 ▶ Strict Epimorphisms**

A 1-morphism $f\colon A \longrightarrow B$ is a **strict epimorphism in** $\mathcal{C}$ if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is injective on objects, i.e. its action on objects

$$f_*\colon \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(B, X)) \longrightarrow \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(A, X))$$

is injective.

01CB **Remark 13.2.9.1.2 ▶ Unwinding Definition 13.2.9.1.1**

In detail, f is a strict epimorphism if, for each diagram in $\mathcal{C}$ of the form

$$A \xrightarrow{f} B \underset{\psi}{\overset{\phi}{\rightrightarrows}} X,$$

if $\phi \circ f = \psi \circ f$, then $\phi = \psi$.

01CC **Example 13.2.9.1.3 ▶ Examples of Strict Epimorphisms**

Here are some examples of strict epimorphisms.

01CD 1. *Strict Epimorphisms in* Cats_2. The strict epimorphisms in Cats_2 are characterised in Categories, Item 1 of Proposition 11.7.3.1.2.

01CE 2. *Strict Epimorphisms in* **Rel**. The strict epimorphisms in **Rel** are characterised in Relations, Proposition 8.4.9.1.1.

01CF 13.2.10 Pseudoepic Morphisms

Let $\mathcal{C}$ be a bicategory.

01CG **Definition 13.2.10.1.1 ▶ Pseudoepic Morphisms**

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **pseudoepic** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is pseudomonic.

01CH **Remark 13.2.10.1.2 ▶ Unwinding Definition 13.2.10.1.1**

In detail, a 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is pseudoepic if it satisfies the following conditions:

01CJ 1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

01CK 2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon \phi \circ f \overset{\sim}{\Longrightarrow} \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f\,.$$

01CL **Proposition 13.2.10.1.3 ► Properties of Pseudoepic Morphisms**

Let $f\colon A \longrightarrow B$ be a 1-morphism of $\mathcal{C}$.

01CM 1. *Characterisations.* The following conditions are equivalent:

01CN (a) The morphism f is pseudoepic.

01CP (b) The morphism f is corepresentably full on cores and corepresentably faithful.

01CQ (c) We have an isococomma square of the form

$$B \overset{\text{eq.}}{\cong} B \overset{\leftrightarrow}{\coprod}_A B, \qquad \begin{tikzcd} B & B \arrow[l, "\mathrm{id}_B"'] \\ B \arrow[u, "\mathrm{id}_B"] & A \arrow[l, "F"] \arrow[u, "F"'] \arrow[ul, dashed] \end{tikzcd}$$

in $\mathcal{C}$ up to equivalence.

Proof 13.2.10.1.4 ▶ Proof of Proposition 13.2.10.1.3

Item 1: Characterisations

Omitted. ∎

Appendices

A Other Chapters

Preliminaries

1. Introduction
2. A Guide to the Literature

Sets

3. Sets
4. Constructions With Sets
5. Monoidal Structures on the Category of Sets
6. Pointed Sets
7. Tensor Products of Pointed Sets

Relations

8. Relations
9. Constructions With Relations
10. Conditions on Relations

Categories

11. Categories

Monoidal Categories

12. Constructions With Monoidal Categories

Bicategories

13. Types of Morphisms in Bicategories

Extra Part

14. Notes

References

[Adá+01] Jiří Adámek, Robert El Bashir, Manuela Sobral, and Jiří Velebil. "On Functors Which Are Lax Epimorphisms". In: *Theory Appl. Categ.* 8 (2001), pp. 509–521. ISSN: 1201-561X (cit. on p. 19).