Types of Morphisms in Bicategories

The Clowder Project Authors

November 22, 2024

In this chapter, we study special kinds of morphisms in bicategories:

1. *Monomorphisms and Epimorphisms in Bicategories (Sections 1 and 2).* There is a large number of different notions capturing the idea of a "monomorphism" or of an "epimorphism" in a bicategory.

   Arguably, the notion that best captures these concepts is that of a *pseudomonic morphism* (Definition 1.10.1) and of a *pseudoepic morphism* (Definition 2.10.1), although the other notions introduced in Sections 1 and 2 are also interesting on their own.

Contents

1 Monomorphisms in Bicategories 1
1.1 Representably Faithful Morphisms 1
1.2 Representably Full Morphisms 2
1.3 Representably Fully Faithful Morphisms 3
1.4 Morphisms Representably Faithful on Cores 6
1.5 Morphisms Representably Full on Cores 7
1.6 Morphisms Representably Fully Faithful on Cores 8
1.7 Representably Essentially Injective Morphisms 9
1.8 Representably Conservative Morphisms 10
1.9 Strict Monomorphisms 11
1.10 Pseudomonic Morphisms 12

2 Epimorphisms in Bicategories 14
2.1 Corepresentably Faithful Morphisms 14

2.2 Corepresentably Full Morphisms 16
2.3 Corepresentably Fully Faithful Morphisms 17
2.4 Morphisms Corepresentably Faithful on Cores 20
2.5 Morphisms Corepresentably Full on Cores 21
2.6 Morphisms Corepresentably Fully Faithful on Cores 22
2.7 Corepresentably Essentially Injective Morphisms 23
2.8 Corepresentably Conservative Morphisms 24
2.9 Strict Epimorphisms 25
2.10 Pseudoepic Morphisms 26

A Other Chapters 28

1 Monomorphisms in Bicategories

1.1 Representably Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.1.1 ▶ Representably Faithful Morphisms

A 1-morphism $f\colon A \longrightarrow B$ of $\mathcal{C}$ is **representably faithful**[1] if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is faithful.

[1] *Further Terminology*: Also called simply a **faithful morphism**, based on Item 1 of Example 1.1.3.

Remark 1.1.2 ▶ Unwinding Definition 1.1.1

In detail, f is representably faithful if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

Example 1.1.3 ▶ Examples of Representably Faithful Morphisms

Here are some examples of representably faithful morphisms.

1. *Representably Faithful Morphisms in* Cats_2. The representably faithful morphisms in Cats_2 are precisely the faithful functors; see Categories, Item 2 of Proposition 6.1.2.

2. *Representably Faithful Morphisms in* **Rel**. Every morphism of **Rel** is representably faithful; see Relations, Item 1 of Proposition 3.8.1.

1.2 Representably Full Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.2.1 ▶ Representably Full Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably full**[1] if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is full.

[1] *Further Terminology:* Also called simply a **full morphism**, based on Item 1 of Example 1.2.3.

Remark 1.2.2 ▶ Unwinding Definition 1.2.1

In detail, f is representably full if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon f \circ \phi \Longrightarrow f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

Example 1.2.3 ▶ Examples of Representably Full Morphisms

Here are some examples of representably full morphisms.

1. *Representably Full Morphisms in* Cats_2. The representably full morphisms in Cats_2 are precisely the full functors; see Categories, **??** of Proposition 6.2.2.

2. *Representably Full Morphisms in* **Rel**. The representably full morphisms in **Rel** are characterised in Relations, Item 2 of Proposition 3.8.1.

1.3 Representably Fully Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.3.1 ▶ Representably Fully Faithful Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably fully faithful**[1] if the following equivalent conditions are satisfied:

1. The 1-morphism f is representably faithful (Definition 1.1.1) and representably full (Definition 1.2.1).

2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is fully faithful.

[1] *Further Terminology:* Also called simply a **fully faithful morphism**, based on Item 1 of Example 1.3.3.

Remark 1.3.2 ▶ Unwinding Representably Fully Faithful Morphisms

In detail, f is representably fully faithful if the conditions in Remark 1.1.2 and Remark 1.2.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

   if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

   then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon f \circ \phi \Longrightarrow f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

   of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

   of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

   of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

Example 1.3.3 ▶ Examples of Representably Fully Faithful Morphisms

Here are some examples of representably fully faithful morphisms.

1. *Representably Fully Faithful Morphisms in* Cats_2. The representably fully faithful morphisms in Cats_2 are precisely the fully faithful functors; see Categories, Item 6 of Proposition 6.3.2.

2. *Representably Fully Faithful Morphisms in* **Rel**. The representably fully faithful morphisms of **Rel** coincide (Relations, Item 3 of Proposition 3.8.1) with the representably full morphisms in **Rel**, which are characterised in Relations, Item 2 of Proposition 3.8.1.

1.4 Morphisms Representably Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

Definition 1.4.1 ▶ Morphisms Representably Faithful on Cores

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably faithful on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

given by postcomposition by f is faithful.

Remark 1.4.2 ▶ Unwinding Definition 1.4.1

In detail, f is representably faithful on cores if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

1.5 Morphisms Representably Full on Cores

Let $\mathcal{C}$ be a bicategory.

Definition 1.5.1 ▶ Morphisms Representably Full on Cores

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably full on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

given by postcomposition by f is full.

Remark 1.5.2 ▶ Unwinding Definition 1.5.1

In detail, f is representably full on cores if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon f \circ \phi \overset{\sim}{\Longrightarrow} f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

1.6 Morphisms Representably Fully Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

Definition 1.6.1 ▶ Morphisms Representably Fully Faithful on Cores

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably fully faithful on cores** if the following equivalent conditions are satisfied:

1. The 1-morphism f is representably faithful on cores (Definition 1.5.1) and representably full on cores (Definition 1.4.1).
2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

given by postcomposition by f is fully faithful.

Remark 1.6.2 ▶ Unwinding Definition 1.6.1

In detail, f is representably fully faithful on cores if the conditions in Remark 1.4.2 and Remark 1.5.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & A \arrow[r, "f"] & B, \arrow[from=U, to=D, Rightarrow, "\alpha"'] \arrow[from=U, to=D, Rightarrow, "\beta"] \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon f \circ \phi \overset{\sim}{\Longrightarrow} f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"{name=U}] \arrow[r, bend right, "f\circ\psi"'{name=D}] & B \arrow[from=U, to=D, Rightarrow, "\beta"'] \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} \quad = \quad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

1.7 Representably Essentially Injective Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.7.1 ▶ Representably Essentially Injective Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably essentially injective** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is essentially injective.

Remark 1.7.2 ▶ Unwinding Definition 1.7.1

In detail, f is representably essentially injective if, for each pair of morphisms $\phi, \psi\colon X \rightrightarrows A$ of $\mathcal{C}$, the following condition is satisfied:

($\star$) If $f \circ \phi \cong f \circ \psi$, then $\phi \cong \psi$.

1.8 Representably Conservative Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.8.1 ▶ Representably Conservative Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **representably conservative** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is conservative.

Remark 1.8.2 ▶ Unwinding Definition 1.8.1

In detail, f is representably conservative if, for each pair of morphisms $\phi, \psi\colon X \rightrightarrows A$ and each 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

of $\mathcal{C}$, if the 2-morphism

$$\mathrm{id}_f \star \alpha\colon f \circ \phi \Longrightarrow f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\mathrm{id}_f\star\alpha\Downarrow"] & B \end{tikzcd}$$

is a 2-isomorphism, then so is α.

1.9 Strict Monomorphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.9.1 ▶ Strict Monomorphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is a **strict monomorphism** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \to \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is injective on objects, i.e. its action on objects

$$f_*\colon \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(X, A)) \to \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(X, B))$$

is injective.

Remark 1.9.2 ▶ Unwinding Definition 1.9.1

In detail, f is a strict monomorphism in $\mathcal{C}$ if, for each diagram in $\mathcal{C}$ of the form

$$X \underset{\psi}{\overset{\phi}{\rightrightarrows}} A \xrightarrow{f} B,$$

if $f \circ \phi = f \circ \psi$, then $\phi = \psi$.

Example 1.9.3 ▶ Examples of Strict Monomorphisms

Here are some examples of strict monomorphisms.

1. *Strict Monomorphisms in* Cats_2. The strict monomorphisms in Cats_2 are precisely the functors which are injective on objects and injective on morphisms; see Categories, Item 1 of Proposition 7.2.2.

2. *Strict Monomorphisms in* **Rel**. The strict monomorphisms in **Rel** are characterised in Relations, Proposition 3.7.1.

1.10 Pseudomonic Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 1.10.1 ▶ Pseudomonic Morphisms

A 1-morphism $f\colon A \longrightarrow B$ of $\mathcal{C}$ is **pseudomonic** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f_*\colon \mathrm{Hom}_{\mathcal{C}}(X, A) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(X, B)$$

given by postcomposition by f is pseudomonic.

Remark 1.10.2 ▶ Unwinding Definition 1.10.1

In detail, a 1-morphism $f\colon A \longrightarrow B$ of $\mathcal{C}$ is pseudomonic if it satisfies the following conditions:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & A \arrow[r, "f"] & B, \end{tikzcd}$$

   if we have

$$\mathrm{id}_f \star \alpha = \mathrm{id}_f \star \beta,$$

   then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon f \circ \phi \overset{\sim}{\Longrightarrow} f \circ \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

   of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \end{tikzcd}$$

   of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} X \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & A \arrow[r, "f"] & B \end{tikzcd} = \begin{tikzcd} X \arrow[r, bend left, "f\circ\phi"] \arrow[r, bend right, "f\circ\psi"'] \arrow[r, phantom, "\beta\Downarrow"] & B \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \mathrm{id}_f \star \alpha.$$

Proposition 1.10.3 ► Properties of Pseudomonic Morphisms

Let $f\colon A \longrightarrow B$ be a 1-morphism of $\mathcal{C}$.

1. *Characterisations.* The following conditions are equivalent:

   (a) The morphism f is pseudomonic.

   (b) The morphism f is representably full on cores and representably faithful.

   (c) We have an isocomma square of the form

$$A \overset{\text{eq.}}{\cong} A \overset{\leftrightarrow}{\times}_B A, \qquad \begin{tikzcd} A \arrow[r, "\mathrm{id}_A"] \arrow[d, "\mathrm{id}_A"'] & A \arrow[d, "F"] \\ A \arrow[r, "F"'] \arrow[ur, dashed, leftrightarrow] & B \end{tikzcd}$$

   in $\mathcal{C}$ up to equivalence.

2. *Interaction With Cotensors.* If $\mathcal{C}$ has cotensors with $\mathbb{1}$, then the following conditions are equivalent:

   (a) The morphism f is pseudomonic.

   (b) We have an isocomma square of the form

$$A \overset{\text{eq.}}{\cong} A \overset{\leftrightarrow}{\times}_{\mathbb{1} \pitchfork F} B, \qquad \begin{tikzcd} A \arrow[r, hook] \arrow[d, "F"'] & \mathbb{1} \pitchfork A \arrow[d, "\mathbb{1} \pitchfork F"] \\ B \arrow[r, hook] \arrow[ur, dashed, leftrightarrow] & \mathbb{1} \pitchfork B \end{tikzcd}$$

   in $\mathcal{C}$ up to equivalence.

Proof 1.10.4 ▶ Proof of Proposition 1.10.3

Item 1: Characterisations

Omitted.

Item 2: Interaction With Cotensors

Omitted. ∎

2 Epimorphisms in Bicategories

2.1 Corepresentably Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.1.1 ▶ Corepresentably Faithful Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably faithful** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is faithful.

Remark 2.1.2 ▶ Unwinding Definition 2.1.1

In detail, f is corepresentably faithful if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

Example 2.1.3 ▶ Examples of Corepresentably Faithful Morphisms

Here are some examples of corepresentably faithful morphisms.

1. *Corepresentably Faithful Morphisms in* Cats_2. The corepresentably faithful morphisms in Cats_2 are characterised in Categories, Item 5 of Proposition 6.1.2.

2. *Corepresentably Faithful Morphisms in* **Rel**. Every morphism of **Rel** is corepresentably faithful; see Relations, Item 1 of Proposition 3.10.1.

2.2 Corepresentably Full Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.2.1 ▶ Corepresentably Full Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably full** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is full.

Remark 2.2.2 ▶ Unwinding Definition 2.2.1

In detail, f is corepresentably full if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta\colon \phi \circ f \Longrightarrow \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-morphism

$$\alpha\colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f.$$

Example 2.2.3 ► Examples of Corepresentably Full Morphisms

Here are some examples of corepresentably full morphisms.

1. *Corepresentably Full Morphisms in* Cats_2. The corepresentably full morphisms in Cats_2 are characterised in Categories, Item 7 of Proposition 6.2.2.

2. *Corepresentably Full Morphisms in* **Rel**. The corepresentably full morphisms in **Rel** are characterised in Relations, Item 2 of Proposition 3.10.1.

2.3 Corepresentably Fully Faithful Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.3.1 ► Corepresentably Fully Faithful Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably fully faithful**[1] if the following equivalent conditions are satisfied:

1. The 1-morphism f is corepresentably full (Definition 2.2.1) and corepresentably faithful (Definition 2.1.1).

2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is fully faithful.

[1] *Further Terminology:* Corepresentably fully faithful morphisms have also been called **lax epimorphisms** in the literature (e.g. in [Adá+01]), though we will always use the name "corepresentably fully faithful morphism" instead in this work.

Remark 2.3.2 ▶ Unwinding Definition 2.3.1

In detail, f is corepresentably fully faithful if the conditions in Remark 2.1.2 and Remark 2.2.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & X \arrow[from=U, to=D, Rightarrow, "\alpha"'] \arrow[from=U, to=D, Rightarrow, "\beta"] \end{tikzcd},$$

   if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

   then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-morphism

$$\beta \colon \phi \circ f \Longrightarrow \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"{name=U}] \arrow[r, bend right, "\psi\circ f"'{name=D}] & X \arrow[from=U, to=D, Rightarrow, "\beta"'] \end{tikzcd}$$

   of $\mathcal{C}$, there exists a 2-morphism

$$\alpha \colon \phi \Longrightarrow \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & X \arrow[from=U, to=D, Rightarrow, "\alpha"'] \end{tikzcd}$$

   of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"{name=U}] \arrow[r, bend right, "\psi"'{name=D}] & X \arrow[from=U, to=D, Rightarrow, "\alpha"'] \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"{name=U}] \arrow[r, bend right, "\psi\circ f"'{name=D}] & X \arrow[from=U, to=D, Rightarrow, "\beta"'] \end{tikzcd}$$

   of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f.$$

Here are some examples of corepresentably fully faithful morphisms.

1. *Corepresentably Fully Faithful Morphisms in* Cats_2. The fully faithful epimorphisms in Cats_2 are characterised in Categories, Item 10 of Proposition 6.3.2.

2. *Corepresentably Fully Faithful Morphisms in* **Rel**. The corepresentably fully faithful morphisms of **Rel** coincide (Relations, Item 3 of Proposition 3.10.1) with the corepresentably full morphisms in **Rel**, which are characterised in Relations, Item 2 of Proposition 3.10.1.

2.4 Morphisms Corepresentably Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

Definition 2.4.1 ▶ Morphisms Corepresentably Faithful on Cores

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably faithful on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(B, X)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(A, X))$$

given by precomposition by f is faithful.

Remark 2.4.2 ▶ Unwinding Definition 2.4.1

In detail, f is corepresentably faithful on cores if, for all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

2.5 Morphisms Corepresentably Full on Cores

Let $\mathcal{C}$ be a bicategory.

Definition 2.5.1 ▶ Morphisms Corepresentably Full on Cores

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably full on cores** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(B,X)) \to \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(A,X))$$

given by precomposition by f is full.

Remark 2.5.2 ▶ Unwinding Definition 2.5.1

In detail, f is corepresentably full on cores if, for each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon \phi \circ f \overset{\sim}{\Longrightarrow} \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f.$$

2.6 Morphisms Corepresentably Fully Faithful on Cores

Let $\mathcal{C}$ be a bicategory.

Definition 2.6.1 ▶ Morphisms Corepresentably Fully Faithful on Cores

A 1-morphism $f\colon A \longrightarrow B$ of $\mathcal{C}$ is **corepresentably fully faithful on cores** if the following equivalent conditions are satisfied:

1. The 1-morphism f is corepresentably full on cores (Definition 2.5.1) and corepresentably faithful on cores (Definition 2.1.1).
2. For each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(B,X)) \longrightarrow \mathrm{Core}(\mathrm{Hom}_{\mathcal{C}}(A,X))$$

given by precomposition by f is fully faithful.

Remark 2.6.2 ▶ Unwinding Definition 2.6.1

In detail, f is corepresentably fully faithful on cores if the conditions in Remark 2.4.2 and Remark 2.5.2 hold:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

if α and β are 2-isomorphisms and we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon \phi \circ f \overset{\sim}{\Longrightarrow} \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f.$$

2.7 Corepresentably Essentially Injective Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.7.1 ► Corepresentably Essentially Injective Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably essentially injective** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is essentially injective.

Remark 2.7.2 ► Unwinding Definition 2.7.1

In detail, f is corepresentably essentially injective if, for each pair of morphisms $\phi, \psi\colon B \rightrightarrows X$ of $\mathcal{C}$, the following condition is satisfied:

($\star$) If $\phi \circ f \cong \psi \circ f$, then $\phi \cong \psi$.

2.8 Corepresentably Conservative Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.8.1 ▶ Corepresentably Conservative Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **corepresentably conservative** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is conservative.

Remark 2.8.2 ▶ Unwinding Definition 2.8.1

In detail, f is corepresentably conservative if, for each pair of morphisms $\phi, \psi\colon B \rightrightarrows X$ and each 2-morphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

of $\mathcal{C}$, if the 2-morphism

$$\alpha \star \mathrm{id}_f\colon \phi \circ f \Longrightarrow \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\alpha\star\mathrm{id}_f \Downarrow"] & X \end{tikzcd}$$

is a 2-isomorphism, then so is α.

2.9 Strict Epimorphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.9.1 ▶ Strict Epimorphisms

A 1-morphism $f\colon A \to B$ is a **strict epimorphism in** $\mathcal{C}$ if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is injective on objects, i.e. its action on objects

$$f_*\colon \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(B, X)) \to \mathrm{Obj}(\mathrm{Hom}_{\mathcal{C}}(A, X))$$

is injective.

Remark 2.9.2 ▶ Unwinding Definition 2.9.1

In detail, f is a strict epimorphism if, for each diagram in $\mathcal{C}$ of the form

$$A \xrightarrow{f} B \underset{\psi}{\overset{\phi}{\rightrightarrows}} X,$$

if $\phi \circ f = \psi \circ f$, then $\phi = \psi$.

Example 2.9.3 ▶ Examples of Strict Epimorphisms

Here are some examples of strict epimorphisms.

1. *Strict Epimorphisms in* Cats_2. The strict epimorphisms in Cats_2 are characterised in Categories, Item 1 of Proposition 7.3.2.

2. *Strict Epimorphisms in* **Rel**. The strict epimorphisms in **Rel** are characterised in Relations, Proposition 3.9.1.

2.10 Pseudoepic Morphisms

Let $\mathcal{C}$ be a bicategory.

Definition 2.10.1 ▶ Pseudoepic Morphisms

A 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is **pseudoepic** if, for each $X \in \mathrm{Obj}(\mathcal{C})$, the functor

$$f^*\colon \mathrm{Hom}_{\mathcal{C}}(B, X) \to \mathrm{Hom}_{\mathcal{C}}(A, X)$$

given by precomposition by f is pseudomonic.

Remark 2.10.2 ▶ Unwinding Definition 2.10.1

In detail, a 1-morphism $f\colon A \to B$ of $\mathcal{C}$ is pseudoepic if it satisfies the following conditions:

1. For all diagrams in $\mathcal{C}$ of the form

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow\Downarrow\beta"] & X, \end{tikzcd}$$

   if we have

$$\alpha \star \mathrm{id}_f = \beta \star \mathrm{id}_f,$$

   then $\alpha = \beta$.

2. For each $X \in \mathrm{Obj}(\mathcal{C})$ and each 2-isomorphism

$$\beta\colon \phi \circ f \overset{\sim}{\Longrightarrow} \psi \circ f, \qquad \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

   of $\mathcal{C}$, there exists a 2-isomorphism

$$\alpha\colon \phi \overset{\sim}{\Longrightarrow} \psi, \qquad \begin{tikzcd} B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd}$$

   of $\mathcal{C}$ such that we have an equality

$$\begin{tikzcd} A \arrow[r, "f"] & B \arrow[r, bend left, "\phi"] \arrow[r, bend right, "\psi"'] \arrow[r, phantom, "\alpha\Downarrow"] & X \end{tikzcd} = \begin{tikzcd} A \arrow[r, bend left, "\phi\circ f"] \arrow[r, bend right, "\psi\circ f"'] \arrow[r, phantom, "\beta\Downarrow"] & X \end{tikzcd}$$

of pasting diagrams in $\mathcal{C}$, i.e. such that we have

$$\beta = \alpha \star \mathrm{id}_f.$$

Proposition 2.10.3 ► Properties of Pseudoepic Morphisms

Let $f\colon A \longrightarrow B$ be a 1-morphism of $\mathcal{C}$.

1. *Characterisations.* The following conditions are equivalent:

   (a) The morphism f is pseudoepic.

   (b) The morphism f is corepresentably full on cores and corepresentably faithful.

   (c) We have an isococomma square of the form

$$B \overset{\text{eq.}}{\cong} B \overset{\leftrightarrow}{\coprod}_A B, \qquad \begin{tikzcd} B & B \arrow[l, "\mathrm{id}_B"'] \\ B \arrow[u, "\mathrm{id}_B"] & A \arrow[l, "F"] \arrow[u, "F"'] \arrow[ul, dashed] \end{tikzcd}$$

   in $\mathcal{C}$ up to equivalence.

Proof 2.10.4 ► Proof of Proposition 2.10.3

Item 1: Characterisations

Omitted. ∎

Appendices

A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Monoidal Structures on the Category of Sets
4. Pointed Sets
5. Tensor Products of Pointed Sets

Relations

6. Relations
7. Constructions With Relations
8. Equivalence Relations and Apartness Relations

Category Theory

9. Categories

Monoidal Categories

10. Constructions With Monoidal Categories

Bicategories

11. Types of Morphisms in Bicategories

References

[Adá+01] Jiří Adámek, Robert El Bashir, Manuela Sobral, and Jiří Velebil. "On Functors Which Are Lax Epimorphisms". In: *Theory Appl. Categ.* 8 (2001), pp. 509–521. ISSN: 1201-561X.