

ON THE SQUARE PEG PROBLEM

JAYANT SHARMA

ABSTRACT. The *Square Peg Problem*, introduced by *Otto Toeplitz* in 1911, asks whether every simple closed Jordan curve C inscribes a square—that is, whether four points on C can form the vertices of a square. To approach this, we introduce the notion of $4\text{-}\Delta$ sets and rotations, as a special case of the more general ${}^n\Delta_k$ configurations, defined as sets of k lines in n -dimensional space with equal mutual angles.

For $n = 2$ and $k = 4$, this configuration consists of two perpendicular line pairs (angle $\pi/2$). We prove that the continuous rotation of such a configuration over a Jordan curve causes the vertical and horizontal line pairs to swap positions. By the *Intermediate Value Theorem*, this implies the existence of a moment when vertical and horizontal lines coincide in length, thus forming the diagonals of an inscribed square.

CONTENTS

1. Preliminaries	1
2. On Δ Rotations	2

1. PRELIMINARIES

We begin by introducing the notion of the ${}^n\Delta_k$ rotation and associated sets.

Let $S \subseteq \mathbb{R}^n$ be a set of points in n -dimensional Euclidean space. The ${}^n\Delta_k$ configuration is defined as a system of k lines in \mathbb{R}^n that are mutually equiangular and intersect at a single point $P(n, k)$, referred to as the **configuration point**.

Corresponding to this configuration, we define the ${}^n\Delta_k$ sets as a collection of $k + 1$ independent sets:

- k of these are termed *parametric sets*, each representing the lengths assigned to the respective k lines of the configuration.
- The remaining set is called the *angular set*, which governs the direction or orientation of the entire configuration in space.

At each instance of rotation determined by an element of the angular set, the k lines act as position vectors with respective magnitudes from the parametric sets. Together, they describe a subset of points in \mathbb{R}^n relative to the configuration point $P(n, k)$.

As we rotate the configuration, at the point $P(n, k)$ by the angles in the angular set in order, we refer to this as the *Delta Rotation*.

In this paper, we primarily consider the specific configuration ${}^2\Delta_4$, representing two mutually perpendicular lines in the plane intersecting at a point $P(2, 4)$, which we shall simply denote by P . For ease of reference, we denote the configuration ${}^2\Delta_4$ by Δ throughout the paper.

Now, we formally state the central conjecture under consideration:

Conjecture I (Square Peg Problem). *Every simple, closed, non-self-intersecting Jordan curve C inscribes a square; that is, there exists a square whose four vertices all lie on the boundary of C .*

Conjecture 1, widely known as the *Square Peg Problem*, was first posed by *Otto Toeplitz* in 1911 and remains one of the most intriguing open problems in elementary geometry.

2. ON Δ ROTATIONS

Throughout this section, we assume that C denotes an arbitrary, closed, simple, and non-self-intersecting Jordan curve in the Euclidean plane \mathbb{R}^2 . Let P be a point strictly contained in the interior of the region enclosed by C , and let P be referred to as the *configuration point*. We define a Δ -configuration at P as a pair of mutually perpendicular lines intersecting at P , extending in all four directions, such that each of the four resulting rays intersects the boundary of C .

Proposition I. *Let $\theta \in [0, 2\pi)$ be any arbitrary angle representing the orientation of a Δ -configuration (i.e., rotation of the perpendicular pair about P). Then, for each such θ , all four rays of the configuration touch the boundary of C in finite, non-zero lengths. That is, none of the four segments connecting P to C along the rays have zero measure.*

Proof. Since P lies strictly inside the Jordan curve C , and C is a simple, closed curve, it follows from the Jordan curve theorem that any ray emanating from P must intersect the boundary of C exactly once before escaping the interior. Given a configuration consisting of two perpendicular lines passing through P , these define four rays in distinct directions. Each of these rays, by the simplicity and closedness of C , must intersect the boundary at a unique point. Consequently, for each angle θ , the segments of the rays from P to their respective points of intersection with C are finite and non-zero in length. Hence, none of the four directional segments in the Δ -configuration can have zero measure. And hence, must have a possible measure so that it touches the boundary of C . (end of proof)

By Proposition 1, we have established that for every orientation θ , all four segments in the Δ -configuration are non-degenerate. We now turn to the question of *continuity* of these segment lengths as a function of rotation. Specifically, we ask: as θ varies continuously, do the corresponding segment lengths also vary continuously?

Proposition II. *Let C and P be as defined above. Then the mapping from orientation angle $\theta \in [0, 2\pi)$ to the vector of four segment lengths in the corresponding Δ -configuration is continuous. That is, small changes in orientation produce small changes in segment lengths.*

Proof. Let T_θ denote the Δ -configuration obtained by orienting the pair of perpendicular lines through P at angle θ . Suppose for the sake of contradiction that the mapping $\theta \mapsto T_\theta$ is not continuous. Then there exists some θ_0 and an $\varepsilon > 0$ such that for every $\delta > 0$, there exists θ' with $|\theta' - \theta_0| < \delta$, yet at least one of the segment lengths in $T_{\theta'}$ differs from that in T_{θ_0} by more than ε .

However, each segment length in T_θ is determined by the distance from P to the intersection point of a ray (in direction determined by θ) with the boundary of C . Since C is a compact, continuous curve and the ray varies continuously with θ , the intersection point—and hence the length—varies continuously with θ . Thus, the segment lengths form continuous functions of θ .

Therefore, our assumption of discontinuity leads to a contradiction, and we conclude that the segment lengths of the Δ -configuration vary continuously as the orientation θ varies. This establishes the desired result. (end of proof)

Building upon the propositions established in the previous section, we now pose a fundamental question concerning the transition between distinct Δ -configurations within a fixed Jordan curve.

Question I. *Given two distinct Δ -configurations at the same configuration point P within a Jordan curve C , does there exist a connected angular subset $\Theta \subseteq [0, \pi)$ and corresponding parametric subsets such that a continuous rotation through Θ induces a continuous transition between the two configurations?*

This question encapsulates the essence of rotational continuity within Δ -configurations. Specifically, we are interested in determining whether it is always possible to rotate a given configuration smoothly to another—most notably, to its *swapped* configuration, in which the original horizontal and vertical lines exchange roles.

If such a continuous transition exists, it implies the existence of a specific angle of rotation, denoted θ^* , at which the configuration is symmetric with respect to the swap—that is, the lengths of the horizontal and vertical segments (measured from P to their respective intersections with the boundary of C) are equal.

Let us define a real-valued function $f(\theta)$ representing the signed difference between the total lengths of the horizontal and vertical segments in the Δ -configuration oriented at angle θ . That is,

$$f(\theta) := \ell_H(\theta) - \ell_V(\theta),$$

where $\ell_H(\theta)$ and $\ell_V(\theta)$ denote the total lengths of the horizontal and vertical segments, respectively, in the configuration at orientation θ .

From the continuity of the segment lengths established earlier, it follows that both $\ell_H(\theta)$ and $\ell_V(\theta)$ are continuous functions of θ . Therefore, $f(\theta)$ is also continuous on $[0, \pi)$.

Suppose the initial configuration corresponds to an angle θ_0 with $f(\theta_0) > 0$, and the swapped configuration corresponds to an angle θ_1 with $f(\theta_1) < 0$. Then, by the Intermediate Value Theorem, there exists some $\theta^* \in (\theta_0, \theta_1)$ such that $f(\theta^*) = 0$.

This implies that at orientation θ^* , the horizontal and vertical segments are of equal length. Due to the orthogonality of the lines in a Δ -configuration, this equality implies that the configuration forms the diagonals of a square inscribed within the curve C and centered at P . And hence proves the *Conjecture 1*.