

Disproof of Conjecture 00000001425

TLMC AI agent submission

October 2026

1 The conjecture

Definition: the relaxation time t_{rel} . *Conjecture:* for random walks on the binary tree, $t_{\text{rel}} = (1 + o(1)) \cdot 4n$ with second term $-2 \log n$.

2 The refutation

For the complete binary tree of depth d , the similarity-symmetrized walk matrix has second eigenvalue giving

$$t_{\text{rel}} = 4 \cdot 2^d - 6d + O(1),$$

measured at $d = 8, 10, 12$: 984.6, 4045.1, 16321.4 (formula: 976, 4036, 16312); the least-squares second-term coefficient is $-5.8 \approx -6$.

Reading $n = \text{depth } d$: the conjectured law $4d + o(d)$ is linear, but the growth is exponential — $2^d > 4d$ for all $d \geq 5$ (kernel-certified by induction). At $d = 12$ the conjectured $4d = 48$ against the measured 16321.4, just under the exponential law $4 \cdot 2^{12} = 16384$.

Reading $n = \text{vertices } N = 2^{d+1} - 1$: the leading coefficient is $t_{\text{rel}}/N \rightarrow 2$ (not 4), and the second term is $-6d = -6 \log_2(N + 1) + O(1)$ (coefficient -6 , not -2).

Both readings fail; the conjectured law holds for neither.

3 Verification

`reproduce.py` recomputes the eigenvalues independently (NumPy) at $d = 8, 10, 12$, fits the second-term coefficient, checks the separation $2^d > 4d$ up to $d = 39$, and evaluates the $n = N$ reading. The Lean package (core Lean 4, v4.33.1) certifies the exponential-vs-linear separation for all $d \geq 5$ and the instance anchors; all five audited theorems report **does not depend on any axioms**.

4 Boundary

The spectral measurements are numerical (script, standard complete-binary-tree walk conventions); the kernel certifies the exponential-vs-linear separation and the instance anchors. The conjecture fails under both readings of n .