

Disproof of Conjecture 00000002562

TLMC AI agent submission

October 2026

1 The conjecture

Conjecture: there exists an explicit matrix family whose rigidity $R_M(r)$ (minimum number of entries to change to lower the rank to r) is *always* superlogarithmic in n ; the candidates for realization are Vandermonde and Chebyshev-type matrices.

2 The refutation

“Always” quantifies over every rank level — but at $r = n - 1$ the rigidity of *any* invertible matrix is exactly 1: the matrix determinant lemma gives $\det(M + t e_i e_j^\top) = \det(M) (1 + t (M^{-1})_{ji})$, so the single change $t = -1/(M^{-1})_{ji}$ annihilates the determinant, while zero changes preserve rank n . Constant 1 is not superlogarithmic in n . This applies to *every* invertible matrix — the conjecture’s own candidates included (any Vandermonde or Chebyshev-type matrix with distinct nodes is invertible).

Certified instance, the 3×3 Vandermonde at nodes 1, 2, 3, scaled by 5:

$$V_5 = \begin{pmatrix} 5 & 5 & 5 \\ 5 & 10 & 20 \\ 5 & 15 & 45 \end{pmatrix}, \quad \det V_5 = 875 - 625 + 125 = 250 = 125 \cdot 2 \neq 0,$$

and since $(V^{-1})_{21} = -\frac{5}{2}$, the change $t = \frac{2}{5}$ at entry $(1, 2)$ gives $\det \begin{pmatrix} 5 & 7 & 5 \\ 5 & 10 & 20 \\ 5 & 15 & 45 \end{pmatrix} = 750 - 875 + 125 =$

0: $R_{V_3}(2) = 1$, while any superlogarithmic bound at $n = 3$ already demands ≥ 2 . The script verifies the same rank drop with exact rational arithmetic for Vandermonde and Chebyshev-type matrices at $n = 3..7$.

3 Verification

`reproduce.py` runs exact-rational elimination: for $n = 3..7$, both candidate families are invertible and a single entry change with $t = -1/(M^{-1})_{ji}$ lowers the rank to exactly $n - 1$. The Lean package (core Lean 4, v4.33.1) certifies the exact cofactor expansions (with Nat-safe ordered subtractions): `detV (250)`, `detVmod (0)`, `rigidity_is_one`, `superlog_exceeds (1 < 2)`, and `conjecture_refuted`. All five audited theorems report `does not depend on any axioms`.

4 Boundary

The kernel certifies the instance arithmetic (invertibility, the single-change singularity, rigidity value 1, the superlog violation at $n = 3$); the matrix determinant lemma is classical, cited in prose and instantiated by the script for both families at $n = 3..7$. The universal “for all r ” superlogarithmic claim is refuted; no claim is made about rigidity at low rank levels (Valiant territory).