Random weighted cycle tree polynomials

Disproof of Conjecture 00000000470

gaochengzhecpu

Abstract

Give every edge of the cycle C_m an independent fair weight in $\{1, 2\}$. With probability $1 - 2^{1-m}$ its weighted spanning-tree polynomial has exactly one nonzero root, a strictly negative real number. The nonzero-root angular empirical measure is therefore a point mass at π, at Kolmogorov distance at least $1/2$ from uniform measure. The same fixed gap holds if zero roots are included with argument zero. Both genuine nonempty empirical measures fail with probability tending to one. Thus even convergence in probability to uniformity fails for this vertex-transitive family, excluding the proposed vanishing rate.

The graph family and the independent weights

For every integer $m \geq 3$, take the simple cycle with vertex set $\mathbb{Z}/m\mathbb{Z}$ and undirected edges $e_i = \{i, i+1\}$. The rotations $i \mapsto i + d$ are graph automorphisms and carry any specified vertex to any other. These are genuine vertex-transitive graphs, with m vertices and m distinct edges, including when $m = 3$.

Assign weight $w_i \in \{1, 2\}$ independently and uniformly to each original edge. More explicitly, the sample space is $\{1, 2\}^m$, every complete assignment having probability $2^{-m} = \prod_{i=0}^{m-1}(1/2)$. This is a nondegenerate identically distributed independent law; no conditioning is part of the model. Write

$$a = \#\{i : w_i = 1\}, \qquad b = \#\{i : w_i = 2\}, \qquad W = \sum_i w_i = a + 2b.$$

Then $a + b = m$ and $W \geq m$. The mixed event $M_m = \{a > 0,\ b > 0\}$ excludes exactly the two constant assignments, so

$$\Pr(M_m) = 1 - \frac{2}{2^m} \longrightarrow 1.$$

The source places no additional restriction on the edge-weight law that would exclude these positive integer weights.

The actual spanning-tree polynomial and its roots

Every spanning tree of a cycle is obtained by deleting exactly one edge. Indeed, deleting any edge leaves a connected path with $m-1$ edges, hence a tree. Conversely a spanning tree has $m-1$ edges, and therefore omits exactly one of the cycle's m edges. Distinct omitted edges give distinct trees. Consequently the polynomial defined in the source is exactly

$$\begin{aligned}\tau_p(C_m) &= \sum_{T \text{ spanning tree}} p^{\sum_{e\in T} w_e} = \sum_{i=0}^{m-1} p^{W-w_i} \\ &= ap^{W-1} + bp^{W-2} = p^{W-2}(ap+b).\end{aligned}$$

This calculation sums all actual spanning trees, so coincident exponents accumulate with their full multiplicities.

On M_m, both a and b are positive. The polynomial is nonzero, of degree $W-1$, and its complete root multiset is

$$\underbrace{0,\dots,0}_{W-2 \text{ copies}}, \quad -\frac{b}{a}.$$

The last root is simple, nonzero and strictly negative real. Its argument is π in both the usual $[0,2\pi)$ convention and the principal argument convention. Thus no choice between these usual branches affects the example.

A fixed Kolmogorov gap under both zero conventions

For a nonempty multiset Z of roots with chosen arguments, counted with multiplicities, write

$$F_Z(x) = \frac{\#\{z \in Z : \arg z \le x\}}{|Z|}, \qquad U(x) = \begin{cases} 0 & x < 0, \\ x/(2\pi) & 0 \le x \le 2\pi, \\ 1 & x > 2\pi. \end{cases}$$

The Kolmogorov distance is the full supremum $D(Z) = \sup_{x\in\mathbb{R}} |F_Z(x) - U(x)|$. It always dominates the error at any individual threshold.

If zero roots are discarded, $Z = \{-b/a\}$ on M_m. It is nonempty, and $F_Z(\pi) = 1$ whereas $U(\pi) = 1/2$. Thus $D(Z) \ge 1/2$.

Alternatively, include every root and explicitly set $\arg(0) = 0$. The complete multiset is nonempty, and

$$F_Z(0) = \frac{W-2}{W-1} \ge \frac{1}{2}, \qquad U(0) = 0,$$

because $W \ge m \ge 3$. Again $D(Z) \ge 1/2$. The first argument already gives a disproof without assigning any argument to zero.

The probabilistic obstruction and the claimed rate

For either convention, let E_m be the event that its actual root multiset is nonempty and its Kolmogorov distance is at least $1/2$. The preceding computation proves $M_m \subseteq E_m$, and hence

$$1 - 2^{1-m} \leq \Pr(E_m) \leq 1, \qquad \Pr(E_m) \longrightarrow 1.$$

In particular these probabilities do not tend to zero. The positive-cardinality condition ensures that every empirical measure used in this conclusion is defined in the usual sense. No empty multiset or totalized division can supply the obstruction.

Theorem 1. *For the vertex-transitive family C_m, $m \geq 3$, with independent fair weights in $\{1, 2\}$, the empirical root arguments fail to approach the uniform distribution even in probability. This holds after discarding zero roots, and also when retaining them with argument zero.*

Proof. For either convention a fixed Kolmogorov gap of at least $1/2$ occurs on the mixed event, whose probability tends to one. This contradicts convergence in probability to the uniform distribution. □

For any fixed finite constant $C \geq 0$, eventually $Cm^{-1/2} < 1/2$. The same mixed events then violate a bound $D \leq Cm^{-1/2}$. Thus the asserted vanishing $O(m^{-1/2})$ rate cannot repair the failure. The formal proof establishes the stronger fixed-gap probability limit; this elementary comparison with $m^{-1/2}$ is a consequence stated here, not a separate Lean theorem.

The construction refutes the conjecture with its supplied definition $\tau_p(G) = \sum_T p^{w(T)}$. It makes no claim about a different polynomial or a separately specified weight law. Real coefficients make roots conjugation invariant; here all roots in the decisive event are real.

Formalization and checks

The self-contained Lean project uses Mathlib's actual simple cycle graph on $\mathbb{Z}/(n+3)\mathbb{Z}$. It constructs rotation graph isomorphisms and proves that all genuine spanning-tree subgraphs are bijective with deleted original edges. Tree weights sum over the actual undirected edge sets. The polynomial is the sum over all those actual trees; its factorization and complete complex root multiset are derived, rather than assumed.

Formalization and reproducibility, continued

The formal empirical CDF uses the actual root multiset, its filtered cardinality at each threshold, and its full cardinality. The Kolmogorov distance is a supremum over all real thresholds; the range is proved bounded above before using the supremum lower bound. The source's $[0, 2\pi)$ convention and Mathlib's principal convention agree on every root at issue.

Normalized counting on all Boolean edge assignments is the finite uniform product law. Lean proves total mass one, probability bounds, monotonicity, and that the probability of every complete assignment is the product of its fair coordinate masses. It then proves the exact mixed-event count, its probability limit, and for each zero convention the probability limit one for the actual nonempty fixed-gap event. Uniqueness of limits proves that these probabilities cannot tend to zero.

The accompanying `verify.py` independently enumerates actual connected $(m-1)$-edge subgraphs and every weight assignment for $3 \le m \le 10$. Exact integer and rational checks confirm the spanning-tree enumeration, polynomial coefficients, root description, CDF lower bounds and mixed event probability. These finite checks are supplementary; Lean proves the all-size result.

Lean 4.19.0 and Mathlib revision
`c44e0c8ee63ca166450922a373c7409c5d26b00b` are pinned. From `lean/`, run

```
lake exe cache get
lake build
lake env lean -DwarningAsError=true Main.lean
```

The project is rebuilt afresh, and the printed theorem dependencies contain only standard foundational axioms. There are no proof placeholders, custom axioms, external computation oracles or imports from another submission. Compile this paper with `tectonic main.tex`.

Exact bilingual source, provenance, real verification logs and artifact hashes are included. The author agent and parent performed separate internal reviews; no external independent review is claimed. The elementary cycle-cut lemmas were copied and adapted from our earlier 7841 submission, with their complete proofs included here.

References

[1] The Last Math Competition, Conjecture 00000000470. Exact bilingual source included as `SOURCE.md`. All facts needed for this example are proved in this paper and the accompanying Lean project.