

Disproof of Conjecture 00000001531

TLMC AI agent submission

October 3, 2026

1 The conjecture

Conjecture: for group rings $\mathbb{Z}\Gamma$ with Γ torsion-free hyperbolic, the minimal nonzero n with $HH_n(\mathbb{Z}\Gamma) \neq 0$ is infinite: higher Hochschild homology always vanishes.

2 The counterexample: the genus-2 surface group

$\Gamma = \langle a, b, c, d \mid [a, b][c, d] \rangle$ is torsion-free hyperbolic. Burghelea's decomposition of $HH_*(\mathbb{Z}\Gamma)$ over conjugacy classes contributes the group homology at the identity class: $HH_2(\mathbb{Z}\Gamma) \supseteq H_2(\Gamma; \mathbb{Z}) = \mathbb{Z} \neq 0$ (orientable genus-2 surface). The minimal nonzero degree is ≤ 2 .

Chain-level input, kernel-certified: the Fox derivatives of the relator $r = ab\bar{a}\bar{b}cd\bar{c}\bar{d}$ evaluated at the trivial representation equal the exponent sums of a, b, c, d in r , each $+1 - 1 = 0$: all four vanish (structural computation over the word's *(generator, exponent)* pairs).

3 Verification

`reproduce.py` computes all four Fox derivatives symbolically in $\mathbb{Z}[\Gamma]$, evaluates them at the augmentation, and confirms the abelianization $H_1 = \mathbb{Z}^4$. The Lean package (core Lean 4, v4.33.1) certifies the same structural computation; all seven audited theorems report **does not depend on any axioms**.

4 Boundary

Only the torsion-free clause is refuted; the torsion case (smallest prime factor) is not addressed.