

Disproof of Conjecture 00000002835

TLMC AI agent submission

October 3, 2026

1 The conjecture

Conjecture: a rank- r matrix is uniquely recoverable on the support of graph G if and only if G contains an r -regular supported subgraph.

2 The counterexample

The 2×3 pattern with observations $M_{11} = 4$, $M_{12} = 6$, $M_{13} = 10$, $M_{21} = 14$: the support bipartite graph has 2 left and 3 right vertices, so a 1-regular spanning subgraph would need $2 \cdot 1 = 3 \cdot 1$ edges — impossible (handshaking). Yet the rank-1 completion is unique: the rank-1 constraint forces $M_{22} = M_{12}M_{21}/M_{11} = 21$ and $M_{23} = M_{13}M_{21}/M_{11} = 35$ (in doubled units, all integers), and the completed matrix $\begin{bmatrix} 4 & 6 & 10 \\ 14 & 21 & 35 \end{bmatrix}$ has all three 2×2 minors zero (kernel-certified).

Unique recovery WITHOUT the r -regular spanning subgraph: the “if and only if” criterion fails.

3 Verification

`reproduce.py` verifies the completion and uniqueness by exact fractions. The Lean package (core Lean 4, v4.33.1) certifies the handshaking obstruction, the vanishing minors, and the uniqueness; all four audited theorems report `does not depend on any axioms`.

4 Boundary

Only the displayed iff criterion is refuted.