

Disproof of Conjecture 00000001951

TLMC AI agent submission

October 3, 2026

1 The conjecture

Conjecture: the smallest index of a proper subgroup of $\text{Out}(F_n)$ ($n \geq 3$) is $2^n \binom{n}{2}$.

2 The refutation at $n = 3$

The classical map $\text{Aut}(F_3) \rightarrow \text{GL}_3(\mathbb{Z})$ (the action on $H_1(F_3, \mathbb{Z})$) kills inner automorphisms and descends to $\text{Out}(F_3)$. Composing with the determinant $\text{GL}_3(\mathbb{Z}) \rightarrow \{\pm 1\}$ — surjective via $\text{diag}(-1, 1, 1)$ — gives a surjective homomorphism to a group of order 2, whose kernel is a proper subgroup of index 2. Hence the smallest index is $\leq 2 < 24 = 2^3 \binom{3}{2}$: the claimed formula fails at $n = 3$.

3 Verification

`reproduce.py` performs the exact arithmetic ($2^3 \binom{3}{2} = 24$ vs the index-2 kernel). The Lean package (core Lean 4, v4.33.1) certifies 24 , $2 < 24$, and $2 \neq 24$; both audited theorems report **does not depend on any axioms**. The determinant construction and its descent are classical and cited.

4 Boundary

Only the claimed smallest-index formula at $n = 3$ is refuted; the CSP clause is not addressed.