

Non-Orientable Foundations: Geometric Resolution of Set-Theoretic Paradoxes via the Möbius Connection

Adam Kevin Morgan
Independent Researcher

April 16, 2026

Abstract

We observe that every classical set-theoretic paradox (Russell, Cantor, Burali-Forti) is an instance of Lawvere’s fixed point theorem, and that in each case the self-referential operation generating the fixed point is an *involution*—a $\mathbb{Z}_2$ action. The unique $U(1)$ connection on S^1 with $\mathbb{Z}_2$ holonomy is the Möbius connection $\omega = \frac{1}{2}d\theta$. Its holonomy group generates a minimal topos—the category $\mathbb{Z}_2\text{-Set}$ of sets with involution—whose subobject classifier has exactly three truth values: *fixed true*, *fixed false*, and *sheet-dependent* (true on one sheet of the double cover, false on the other). This three-valued logic preserves classical reasoning locally while obstructing the global operations that generate paradoxes.

We demonstrate three consequences: (i) the Burali-Forti paradox is resolved because the collection of all ordinals, while existing as a completed totality, fails to be globally well-ordered; (ii) Cantor’s paradox is resolved because the diagonal set, while well-defined locally, fails to globalize as a section of the subobject classifier; and (iii) the Kunen inconsistency theorem, which prevents nontrivial elementary embeddings $j: V \rightarrow V$ in ZFC, loses its logical supports in the non-orientable setting. These resolutions share a uniform mechanism: non-orientability permits totalities to exist while preventing paradox-generating operations from globalizing.

The three-valued logic of $\mathbb{Z}_2\text{-Set}$ is shown to be a genuine fourth category distinct from classical logic, glut paraconsistency, and gap logic: its third truth value encodes topological orientation-dependence rather than contradiction or indeterminacy. This distinction allows the framework to resolve paradoxes without inheriting the metatheoretical difficulties that paraconsistent foundations have faced.

Given the connection’s existence, the construction is self-grounding at the level of categorical primitives: the Möbius connection generates $\mathbb{Z}_2\text{-Set}$, which is a cartesian closed category to which Lawvere’s theorem applies, and within which the diagonal arguments reproduce the $\mathbb{Z}_2$ involution that determines the connection. The categorical content of the paradoxes is a fixed point of its own generative principle. The deeper derivation of the connection itself from arithmetic structure is the subject of a companion paper.

1 Introduction

The classical paradoxes of naive set theory—Russell’s paradox, Burali-Forti’s paradox of the ordinal of all ordinals, Cantor’s paradox of the set of all sets—motivated the axiomatic foundations of modern mathematics. Zermelo–Fraenkel set theory with the axiom of choice (ZFC) resolves each paradox by restricting the formation of totalities: the axioms of Foundation, Separation, and Replacement collectively ensure that no “set of all sets,” “ordinal of all ordinals,” or “set of all sets not members of themselves” can be formed as a completed object within the theory.

This resolution is effective but *prohibitive* in character. It works by forbidding the constructions that generate paradox without identifying a structural reason why those constructions fail. The

set/proper class distinction—the central device—is imposed by axiomatic fiat rather than derived from a mathematical principle.

A structural explanation already exists, though its geometric implications have not been fully pursued. Lawvere’s fixed point theorem [1] demonstrates that Cantor’s diagonal argument, Russell’s paradox, Gödel’s incompleteness theorem, and the halting problem are instances of a single categorical fact: in any cartesian closed category $\mathcal{C}$ with a point-surjective morphism $A \rightarrow Y^A$, every endomorphism $Y \rightarrow Y$ has a fixed point. The paradoxes arise when this fixed point cannot be consistently accommodated—when the category’s structure is too rigid to contain the object that self-reference demands.

The present paper takes Lawvere’s unification as its starting point and asks: *what is the geometry of the fixed point?* We observe that in every classical paradox, the self-referential operation is an *involution*—an operation of order 2 that maps truth to falsity, membership to non-membership, or a function value to its complement. This $\mathbb{Z}_2$ structure is not incidental; it is the categorical content of negation. The unique $U(1)$ connection on S^1 whose holonomy realizes this $\mathbb{Z}_2$ action is $\omega = \frac{1}{2}d\theta$, the Möbius connection. Its holonomy group generates a topos—the category $\mathbb{Z}_2\text{-Set}$ of sets equipped with a $\mathbb{Z}_2$ involution—whose internal logic is classical locally but admits a third, *sheet-dependent* truth value for propositions that reverse under the involution.

This paper develops the consequences of this observation for set-theoretic foundations. Section 2 establishes the $\mathbb{Z}_2$ structure of diagonal arguments via Lawvere’s theorem. Section 3 derives the Möbius connection as the unique geometric realization of this $\mathbb{Z}_2$ structure and introduces the topos $\mathbb{Z}_2\text{-Set}$. Section 4 establishes that the construction is self-grounding: the connection generates the categorical setting in which Lawvere’s theorem reproduces the connection. Section 5 presents an elementary two-chart construction illustrating the abstract structure in concrete terms. Sections 6–7 resolve the Burali-Forti and Cantor paradoxes. Section 8 situates the three-valued logic of $\mathbb{Z}_2\text{-Set}$ within the taxonomy of classical, paraconsistent, and gap logics, and addresses the metatheoretical concerns of Field [14] and Shapiro [15]. Section 9 dissolves the Kunen barrier. Section 10 discusses implications and open problems.

2 Diagonal Arguments as $\mathbb{Z}_2$ Involutions

2.1 Lawvere’s fixed point theorem

We recall the result that unifies the classical diagonal arguments.

Theorem 2.1 (Lawvere, 1969 [1]). *Let $\mathcal{C}$ be a cartesian closed category. If there exists a point-surjective morphism $\phi: A \rightarrow Y^A$, then every endomorphism $f: Y \rightarrow Y$ has a fixed point: there exists a point $y: 1 \rightarrow Y$ such that $f \circ y = y$.*

The diagonal arguments of set theory arise when the category is **Set** and the endomorphism f has no fixed point (e.g., $f = \neg: \{0, 1\} \rightarrow \{0, 1\}$), implying that the point-surjective morphism ϕ cannot exist. The “paradox” in each case is the attempt to construct ϕ for an f that is fixed-point-free.

2.2 The involutory structure of negation

Proposition 2.2 (Diagonal arguments are $\mathbb{Z}_2$ actions). *In each classical paradox, the endomorphism $f: Y \rightarrow Y$ whose fixed-point-freeness generates the contradiction is an involution: $f^2 = \text{id}_Y$. Equivalently, f is a $\mathbb{Z}_2$ action on Y .*

Proof. We verify this in each case.

Cantor’s diagonal argument. The endomorphism is complementation: $f: \{0, 1\} \rightarrow \{0, 1\}$, $f(b) = 1 - b$. This satisfies $f^2(b) = 1 - (1 - b) = b$, so $f^2 = \text{id}$.

Russell’s paradox. The relevant operation maps the proposition $x \in S$ to $x \notin S$. This is logical negation: $\neg: \Omega \rightarrow \Omega$, where $\Omega = \{\text{true}, \text{false}\}$ is the subobject classifier of **Set**. Negation satisfies $\neg\neg = \text{id}$.

Burali-Forti’s paradox. The contradiction arises from $\Omega \in \Omega$ versus $\Omega \notin \Omega$ —the same membership/non-membership involution applied to the collection of all ordinals.

Gödel’s incompleteness. The Gödel sentence G asserts its own unprovability. The relevant map sends “provable” to “true but unprovable,” which composes with itself to return to “provable.” The self-referential core is the negation in “I am not provable,” which is again $\neg$.

In every case, the fixed-point-free endomorphism is negation or complement—an operation of exact order 2. $\square$

Remark 2.3. The $\mathbb{Z}_2$ structure is not a contingent feature of these particular arguments. In a cartesian closed category, the subobject classifier Ω carries a canonical involution (negation), and Lawvere’s theorem applied with this involution produces exactly the classical paradoxes. The $\mathbb{Z}_2$ action *is* the categorical content of self-referential negation.

3 The Möbius Connection and Its Topos

3.1 Deriving the connection from the $\mathbb{Z}_2$ structure

The diagonal arguments produce a $\mathbb{Z}_2$ involution. We now ask: what is the *geometric* structure that carries a $\mathbb{Z}_2$ action canonically?

Definition 3.1 (The Möbius connection). The **Möbius connection** is the connection 1-form $\omega = \frac{1}{2}d\theta$ on the circle S^1 , viewed as a $U(1)$ -connection. Its holonomy around the generating loop is:

$$\text{hol}(\omega) = \exp\left(i \oint_{S^1} \frac{1}{2} d\theta\right) = e^{i\pi} = -1.$$

This $\mathbb{Z}_2 \subset U(1)$ holonomy generates multiplication by -1 : a quantity transported around the loop returns with its sign reversed.

Proposition 3.2 (Uniqueness of the Möbius connection). *The connection $\omega = \frac{1}{2}d\theta$ is the unique $U(1)$ -connection on S^1 satisfying:*

- (i) *Nontrivial holonomy: $\text{hol}(\omega) \neq 1$.*
- (ii) *Minimal nontrivial cover: the associated covering space has exactly two sheets.*
- (iii) *Field extension: the double cover, equipped with the holonomy-induced identification, carries the structure of a field extension of $\mathbb{R}$.*

Proof. A general $U(1)$ -connection on S^1 has the form $\omega_\alpha = \alpha d\theta$ for $\alpha \in \mathbb{R}$. The holonomy is $e^{2\pi i \alpha}$.

If $\alpha \in \mathbb{Z}$, the holonomy is trivial: condition (i) fails.

If $\alpha = k/n$ with $n > 2$ in lowest terms, the holonomy has order $n > 2$, producing an n -sheeted cover: condition (ii) fails.

If $\alpha \notin \mathbb{Q}$, the holonomy generates a dense subgroup of $U(1)$, and the associated cover does not close: condition (ii) fails.

The remaining case is $\alpha = k/2$ for odd k . All such connections have holonomy $e^{i\pi k} = -1$ and produce the same 2-sheeted double cover. The connections differ by gauge transformation (adding an integer to α), so $\alpha = 1/2$ is the unique representative in the fundamental domain $\alpha \in (0, 1]$.

For condition (iii): the double cover of S^1 is again S^1 , and the deck transformation (the $\mathbb{Z}_2$ action exchanging sheets) provides the conjugation that, together with the algebra structure of the covering space, yields $\mathbb{C}$ as a degree-2 field extension of $\mathbb{R}$. No other connection satisfies all three conditions. $\square$

3.2 The topos of $\mathbb{Z}_2$ -sets

The holonomy group of the Möbius connection generates a natural categorical setting.

Definition 3.3 (The topos $\mathbb{Z}_2\text{-Set}$). The category $\mathbb{Z}_2\text{-Set}$ has as objects pairs (X, σ) where X is a set and $\sigma: X \rightarrow X$ is an involution ($\sigma^2 = \text{id}_X$). Morphisms are equivariant functions: $f: (X, \sigma_X) \rightarrow (Y, \sigma_Y)$ satisfying $f \circ \sigma_X = \sigma_Y \circ f$.

Proposition 3.4 (Standard; see [8]). $\mathbb{Z}_2\text{-Set}$ is an elementary topos. In particular:

- (a) It is cartesian closed.
- (b) It has a subobject classifier Ω .
- (c) It has all finite limits and colimits.

The key structure for our purposes is the subobject classifier.

Proposition 3.5 (Three-valued subobject classifier). The subobject classifier of $\mathbb{Z}_2\text{-Set}$ is $\Omega = (\{\text{true}, \text{false}, \text{swap}\}, \sigma_\Omega)$ where:

$$\begin{aligned}\sigma_\Omega(\text{true}) &= \text{true}, \\ \sigma_\Omega(\text{false}) &= \text{false}, \\ \sigma_\Omega(\text{swap}) &= \text{swap},\end{aligned}$$

with the understanding that **swap** classifies subobjects that are “swapped” by the involution—elements that belong to the subobject on one sheet of the double cover and do not belong on the other.

More precisely, for any $\mathbb{Z}_2$ -set (X, σ) and $\mathbb{Z}_2$ -subset $A \hookrightarrow X$, the characteristic morphism $\chi_A: X \rightarrow \Omega$ assigns:

- $\chi_A(x) = \text{true}$ if $x \in A$ and $\sigma(x) \in A$,
- $\chi_A(x) = \text{false}$ if $x \notin A$ and $\sigma(x) \notin A$,
- $\chi_A(x) = \text{swap}$ if exactly one of x and $\sigma(x)$ belongs to A .

Remark 3.6 (Interpretation of the three truth values). The three values correspond to three relationships between a proposition and the $\mathbb{Z}_2$ involution:

- **true**: The proposition is *invariant*—it holds on both sheets and is unaffected by the holonomy. This is classically true.
- **false**: The proposition fails on both sheets. This is classically false.
- **swap**: The proposition holds on one sheet and fails on the other. Transport around the non-trivial loop reverses its truth value. This is the *boundary* truth value—locally definite, globally orientation-dependent.

Within the fixed-point locus of the involution (elements x with $\sigma(x) = x$), only **true** and **false** arise: classical logic holds in full. The third value appears only for elements that participate in the non-trivial $\mathbb{Z}_2$ action.

4 Self-Grounding

A natural foundational question arises: the derivation in Section 2 begins from Lawvere’s theorem, which requires a cartesian closed category. Where does this category come from? If it must be postulated externally, the non-orientable framework merely shifts the foundational burden from set theory to category theory.

We observe that the categorical setting is generated by the Möbius connection itself, and that Lawvere’s theorem applied within this setting reproduces the connection. The foundational loop closes at the level of categorical primitives given the connection. A further level of closure—deriving the continuous mathematical machinery used to define the connection—is addressed by the companion paper [16]; see Remark 4.4.

Theorem 4.1 (Self-grounding of the Möbius connection). *The following loop of derivations is consistent:*

- (i) *The Möbius connection $\omega = \frac{1}{2}d\theta$ has holonomy group $\mathbb{Z}_2$.*
- (ii) *The holonomy group $\mathbb{Z}_2$ generates the topos $\mathbb{Z}_2\text{-Set}$ (the category of presheaves on the orbit category of $\mathbb{Z}_2$).*
- (iii) *$\mathbb{Z}_2\text{-Set}$ is a cartesian closed category (Proposition 3.4).*
- (iv) *Lawvere’s fixed point theorem (Theorem 2.1) applies within $\mathbb{Z}_2\text{-Set}$.*
- (v) *The subobject classifier of $\mathbb{Z}_2\text{-Set}$ carries a canonical involution (negation), which is a $\mathbb{Z}_2$ action (Proposition 3.5).*
- (vi) *By Proposition 2.2, this $\mathbb{Z}_2$ action is the categorical content of the diagonal arguments.*
- (vii) *By Proposition 3.2, the unique connection with $\mathbb{Z}_2$ holonomy is $\omega = \frac{1}{2}d\theta$.*

The connection generates the topos in which Lawvere’s theorem produces the $\mathbb{Z}_2$ involution that uniquely determines the connection. Given the connection’s existence, the categorical content of the diagonal arguments forms a fixed point of its own generative principle. The independent derivation of the connection from arithmetic structure (the deeper closure) is developed in the companion paper [16]; see Remark 4.4.

Proof. Each step invokes established mathematics: (i) is a standard computation of holonomy; (ii) is the standard construction of a presheaf topos from a group; (iii) is a theorem of topos theory [8]; (iv) is Lawvere’s theorem [1]; (v) is Proposition 3.5; (vi) is Proposition 2.2; (vii) is Proposition 3.2. No step in the loop is novel; each is an application of known results. The observation is that these known results compose into a self-referential loop with a unique fixed point. $\square$

Remark 4.2 (Fixed point, not circular argument). The self-grounding structure is a *fixed point*, not a vicious circle. A vicious circle of the form “ A presupposes B presupposes A ” generates no content. A fixed point of the form “ A generates a setting in which A is the unique solution” is a *consistency condition*: it constrains the structure rather than leaving it undetermined. That the foundation of the framework is itself an instance of the fixed-point phenomenon it describes (Lawvere’s theorem) is a feature of self-referential structures, not a defect.

This situation has precedent. Any foundational system must eventually account for its own expressibility. ZFC is formulated in first-order logic, which is itself studied within ZFC. Category theory is described using categories. The novel feature here is that the self-reference is *mathematically precise*: the connection generates the topos, and Lawvere’s theorem within the topos reproduces the connection. The self-reference is a theorem, not an unexamined presupposition.

Remark 4.3 (Relationship to external foundations). The self-grounding does not eliminate the need to *communicate* the construction using some pre-existing mathematical language. The present paper uses standard mathematical notation and reasoning, which the reader may ground in ZFC,

category theory, or any other framework. The claim is not that the construction requires no language of presentation, but that the *mathematical content* of the construction does not depend on an external foundation: the connection, the topos, and Lawvere’s theorem form a self-consistent package.

Remark 4.4 (Scope of the self-grounding claim). A careful reader may note that Theorem 4.1 assumes the existence of the Möbius connection as a mathematical object. Defining $\omega = \frac{1}{2}d\theta$ rigorously requires the circle S^1 as a manifold, real numbers for the continuous parameter θ , differential forms, integration (to compute holonomy via $\oint$), and $U(1)$ as a Lie group. Each of these presupposes substantial mathematical machinery that standardly requires a set-theoretic or equivalent foundation. The self-grounding established in this paper therefore operates *at the level of categorical primitives given the connection’s existence*: it shows that the topos, its three-valued logic, and the diagonal arguments form a closed fixed-point loop, *not* that the continuous machinery of the connection itself is generated from nothing.

The deeper foundational closure—deriving the number systems $\mathbb{Z}$, $\mathbb{R}$, and $\mathbb{C}$ themselves from the connection rather than presupposing them—is the subject of the companion paper [16]. There, the connection is shown to be the initial object of the category of oriented arithmetic structures, and the number systems are derived from its holonomy and double-cover structure without recourse to set-theoretic construction. Together, the two papers establish self-grounding at both levels: Paper [16] grounds the continuous machinery of the connection, and the present paper grounds the categorical content of the diagonal arguments. Each paper makes its own modular contribution; the full foundational story requires both.

This staged structure is itself a response to foundational scope: claiming total self-grounding in a single paper would either overreach or require compressing both derivations into a single document. The modular approach allows each claim to be evaluated on its own terms while making the composition transparent.

5 Illustration: The Two-Chart Construction

We present a concrete, elementary construction that illustrates the abstract structure of Section 3 in familiar terms. The construction uses standard mathematical language for accessibility; the underlying structure is derived from the categorical framework of the preceding sections, not from the set-theoretic notions used to illustrate it.

Construction 5.1 (Non-orientable colimit of ordered sets). Define two copies of a segment of the natural numbers:

$$\begin{aligned} A &= \{0_A, 1_A, 2_A, 3_A, 4_A, 5_A, 6_A, 7_A, 8_A, 9_A, \dots\}, \\ B &= \{0_B, 1_B, 2_B, 3_B, 4_B, 5_B, 6_B, 7_B, 8_B, 9_B, \dots\}, \end{aligned}$$

each equipped with the standard well-ordering inherited from ω .

Declare an overlap region $U = \{3, 4, 5, 6, 7\}$ in each chart, and define the transition function $\varphi: U_A \rightarrow U_B$ by:

$$\varphi(n_A) = (10 - n)_B \quad \text{for } n \in \{3, 4, 5, 6, 7\}.$$

Explicitly: $3_A \mapsto 7_B$, $4_A \mapsto 6_B$, $5_A \mapsto 5_B$, $6_A \mapsto 4_B$, $7_A \mapsto 3_B$. This is an order-reversing bijection: the discrete analogue of an orientation-reversing transition function.

The **colimit space** $X = A \sqcup B / \sim$, where $a \sim \varphi(a)$ for all $a \in U_A$, identifies the overlap elements. Writing $p = [3_A] = [7_B]$, $q = [4_A] = [6_B]$, $r = [5_A] = [5_B]$, $s = [6_A] = [4_B]$, $t = [7_A] = [3_B]$, the

colimit space consists of elements from each chart below the overlap, the five identified overlap elements, and elements from each chart above the overlap.

Proposition 5.2 (Local well-ordering is preserved). *Each chart’s image in the colimit inherits a well-ordering from its source.*

Chart A ’s image: $0_A < 1_A < 2_A < p < q < r < s < t < 8_A < 9_A < \dots$

Chart B ’s image: $0_B < 1_B < 2_B < t < s < r < q < p < 8_B < 9_B < \dots$

Both are isomorphic to ω as ordered sets.

Proof. The ordering on each chart’s image is the pushforward of the standard ordering on ω , which is a well-ordering. The chart maps are injective on non-overlap portions and agree with the identification on the overlap. $\square$

Theorem 5.3 (Global well-ordering fails). *There is no total order on X that restricts to the inherited well-ordering on both charts simultaneously.*

Proof. Chart A requires $p < q$ (since $p = [3_A]$, $q = [4_A]$, and $3 < 4$ in ω). Chart B requires $q < p$ (since $q = [6_B]$, $p = [7_B]$, and $6 < 7$ in ω). No total order satisfies both. $\square$

Remark 5.4 (Connection to the Möbius strip). Construction 5.1 is the discrete ordered analogue of the Möbius strip. The Möbius strip is formed by gluing two oriented strips with an orientation-reversing transition function. Each strip is locally oriented; the resulting surface is globally non-orientable. Here, each chart is locally well-ordered; the colimit is not globally well-orderable. The obstruction is identical: a $\mathbb{Z}_2$ holonomy (order reversal) around the identification cycle, generated by the Möbius connection of Definition 3.1.

Remark 5.5 (The three truth values, concretely). Consider the proposition “ $p < q$ ” in the colimit space X . In chart A , this proposition evaluates to **true**. In chart B , this proposition evaluates to **false**. The proposition’s truth value is *sheet-dependent*: it is the **swap** value of the subobject classifier (Proposition 3.5). This is the concrete manifestation of the three-valued logic derived abstractly from $\mathbb{Z}_2\text{-Set}$.

6 Resolution of the Burali-Forti Paradox

6.1 The classical paradox

The Burali-Forti paradox [4] proceeds in five steps:

- (BF1) **Collection.** Let Ω be the collection of all ordinals.
- (BF2) **Well-ordering.** Ω is well-ordered by $\in$.
- (BF3) **Transitivity.** Ω is transitive.
- (BF4) **Ordinalhood.** Since Ω is transitive and well-ordered, Ω is an ordinal.
- (BF5) **Contradiction.** $\Omega \in \Omega$, contradicting irreflexivity.

In ZFC, the paradox is blocked at (BF1): Ω is a proper class, not a set.

6.2 The orientability dependence

Lemma 6.1 (Step (BF2) is a global orientability condition). *The claim that Ω is well-ordered by $\in$ requires that the local well-orderings of individual ordinals cohere into a single global well-ordering of the entire collection. This coherence requires a consistent “direction” of succession defined everywhere. In the language of the subobject classifier (Proposition 3.5), it requires that every comparison $\alpha < \beta$ between ordinals evaluates to **true** or **false**—never to **swap**.*

Proof. A well-ordering requires totality: for any $\alpha, \beta \in \Omega$, exactly one of $\alpha < \beta$, $\alpha = \beta$, $\beta < \alpha$ holds. This is a global condition on all pairs. If some comparison evaluates to **swap** (true on one sheet, false on the other), the totality condition fails: the comparison does not have a definite global truth value.

By Theorem 5.3, non-orientable structures necessarily produce such sheet-dependent comparisons. Hence global well-ordering is obstructed. $\square$

6.3 Resolution

Theorem 6.2 (Geometric resolution of the Burali-Forti paradox). *In the topos $\mathbb{Z}_2\text{-Set}$, the collection Ω of all ordinal objects exists as a completed totality. It is transitive (BF3) and locally well-ordered (within any chart, ordinals and their predecessors form a well-ordered set). However, Ω is not globally well-ordered (BF2 fails), and therefore does not satisfy the definition of an ordinal. Step (BF4) does not proceed, and no contradiction arises.*

Proof. Existence. The topos $\mathbb{Z}_2\text{-Set}$ does not impose the set/class distinction that blocks (BF1) in ZFC. Totalities exist as objects of the topos.

Transitivity. Every element of an ordinal is an ordinal. This is a local property depending only on the membership relation within individual ordinals, unaffected by the global involution.

Local well-ordering. Within the fixed-point locus of the involution (elements x with $\sigma(x) = x$), the subobject classifier restricts to $\{\mathbf{true}, \mathbf{false}\}$ and classical logic holds. Any individual ordinal α and its predecessors $\{\beta : \beta < \alpha\}$ live within this classical region.

Global well-ordering fails. By Lemma 6.1, comparisons between ordinals in different charts can evaluate to **swap**, obstructing global totality. The local well-orderings do not cohere.

Ordinalhood fails. Ω is transitive but not globally well-ordered. It does not satisfy the definition of an ordinal. Self-membership never arises. $\square$

Remark 6.3 (Comparison with ZFC). ZFC prevents Ω from existing. The non-orientable framework permits Ω to exist but prevents it from being an ordinal. The former is ontological restriction; the latter is topological obstruction. The non-orientable resolution identifies a *reason* for the failure: global well-ordering is an orientability condition, and the total structure is non-orientable.

7 Resolution of Cantor's Paradox

7.1 The classical paradox

Cantor's paradox [3] proceeds in four steps:

- (C1) **Collection.** Let V be the set of all sets.
- (C2) **Power set.** V has a power set $\mathcal{P}(V)$.
- (C3) **Cantor's theorem.** $|\mathcal{P}(V)| > |V|$.
- (C4) **Contradiction.** $\mathcal{P}(V) \subseteq V$, so $|\mathcal{P}(V)| \leq |V|$.

In ZFC, the paradox is blocked at (C1): V is a proper class.

7.2 Local preservation of diagonalization

Proposition 7.1 (Diagonalization holds locally). *For any object S in $\mathbb{Z}_2\text{-Set}$ contained within the fixed-point locus of the involution (i.e., $\sigma_S = \text{id}_S$), Cantor's theorem holds: $|\mathcal{P}(S)| > |S|$.*

Proof. Within the fixed-point locus, the subobject classifier restricts to $\{\mathbf{true}, \mathbf{false}\}$, the logic is classical, and the standard diagonal argument applies without modification. $\square$

Remark 7.2. Proposition 7.1 ensures that the cardinal hierarchy $\aleph_0 < \aleph_1 < \aleph_2 < \dots$ is intact for ordinary mathematical objects. The modification occurs only at the scale of self-referential totalities.

7.3 Failure of global diagonalization

Theorem 7.3 (Cantor’s theorem does not apply to the total structure). *Let V be the total structure of $\mathbb{Z}_2\text{-Set}$. The diagonal set $D = \{x \in V : x \notin f(x)\}$, for any $f: V \rightarrow \mathcal{P}(V)$, fails to be a well-defined global subobject of V .*

Proof. The diagonal set D is defined by the condition $x \in D \iff x \notin f(x)$. The characteristic morphism of D is the composition:

$$V \xrightarrow{\langle \text{id}, f \rangle} V \times \mathcal{P}(V) \xrightarrow{\text{ev}} \Omega \xrightarrow{\neg} \Omega,$$

where ev is the evaluation (membership test) and $\neg$ is negation.

In **Set** (the classical topos), $\neg$ maps **true** $\mapsto$ **false** and **false** $\mapsto$ **true**. The composition always lands in $\{\text{true}, \text{false}\}$, and D is a well-defined subset.

In $\mathbb{Z}_2\text{-Set}$, the evaluation ev can produce any of the three truth values. Negation acts on the subobject classifier as: $\neg(\text{true}) = \text{false}$, $\neg(\text{false}) = \text{true}$, $\neg(\text{swap}) = \text{swap}$.

The last equation is the critical point: negating a sheet-dependent truth value yields another sheet-dependent truth value. The characteristic morphism $\chi_D = \neg \circ \text{ev} \circ \langle \text{id}, f \rangle$ therefore takes the value **swap** on elements x for which the membership $x \in f(x)$ is sheet-dependent.

A subobject classified by a morphism taking the value **swap** is not a $\mathbb{Z}_2$ -invariant subobject—it is a subobject on one sheet but not the other. It does not exist as a global subobject of V . The diagonal “set” D is a *local section* of the subobject classifier that fails to globalize.

Cantor’s theorem requires D to be a well-defined subobject. In $\mathbb{Z}_2\text{-Set}$ applied to the total structure, D is not well-defined as a global subobject, and the theorem does not apply. $\square$

Corollary 7.4 ($\mathcal{P}(V) \subseteq V$ is consistent). *Since Cantor’s theorem does not apply to the total structure, $\mathcal{P}(V) \subseteq V$ is consistent in $\mathbb{Z}_2\text{-Set}$. The power set of the whole can be contained in the whole.*

Remark 7.5 (Scale-dependent validity). The resolution is sharply delineated by the involution structure:

- For objects in the fixed-point locus ($\sigma = \text{id}$): classical logic, binary truth values, Cantor’s theorem holds.
- For the total structure (nontrivial σ): three-valued logic, diagonal set can be sheet-dependent, Cantor’s theorem need not apply.

The boundary between regimes is the fixed-point locus of the $\mathbb{Z}_2$ action—a mathematically precise criterion, not an ad hoc distinction.

8 The Three-Valued Logic and Paraconsistency

The paradox resolutions of Sections 6–7 depend on a logic that is not classical: the subobject classifier of $\mathbb{Z}_2\text{-Set}$ admits a third truth value **swap** alongside **true** and **false**. This section addresses an obvious question raised by this fact: how does the three-valued logic of $\mathbb{Z}_2\text{-Set}$ relate to the well-studied paraconsistent logics of philosophical and mathematical logic, and does it inherit their foundational difficulties?

We argue that the logic of $\mathbb{Z}_2\text{-Set}$ is a genuine *fourth category* distinct from classical logic, dialetheic (glut) paraconsistency, and gap logic. Its third truth value encodes *topological orientation-dependence* rather than logical contradiction or indeterminacy. This distinction allows the framework to resolve paradoxes without incurring the metatheoretical difficulties that paraconsistent foundations have historically faced.

8.1 The taxonomy of third truth values

Logics admitting a third truth value arise in response to several distinct phenomena. We distinguish the principal types:

Definition 8.1 (Four logical regimes).

- (a) **Classical logic.** Every proposition takes exactly one of the values $\{\mathbf{true}, \mathbf{false}\}$. Truth is globally consistent. The law of excluded middle and the law of non-contradiction both hold.
- (b) **Glut paraconsistency** (e.g., Priest’s LP [12], dialetheism): Some propositions are *both true and false*. The third value represents genuine contradictions tolerated by the logic; excluded middle holds, but non-contradiction fails.
- (c) **Gap logic** (e.g., Kleene’s strong three-valued logic [13], many intuitionistic variants): Some propositions are *neither true nor false*. The third value represents indeterminacy, undefinedness, or absence of truth value; non-contradiction holds, but excluded middle fails.
- (d) **Orientation-dependent logic** (the logic of $\mathbb{Z}_2\text{-Set}$): Every proposition has a definite truth value at every point, but some propositions have truth values that *vary across the base space*. The third value encodes that a proposition is **true** on one sheet of a $\mathbb{Z}_2$ cover and **false** on the other, with no global aggregation. Neither excluded middle nor non-contradiction is violated pointwise; what fails is the assumption that pointwise truth values aggregate into a single global truth value.

Remark 8.2 (The essential distinction). Regimes (b) and (c) modify the *local* behavior of truth: at a given point, a proposition is ambiguous (glut) or absent (gap). Regime (d) preserves classical local behavior and modifies only the *global* aggregation. At every point of $\mathbb{Z}_2\text{-Set}$, a proposition is either **true** or **false**—but the point at which it is **true** on one sheet is the same point at which it is **false** on the other, and the involution σ exchanges these sheets. The third value **swap** is not a local truth value but a *global descriptor* of how a pointwise-classical proposition behaves under the $\mathbb{Z}_2$ action.

Proposition 8.3 ($\mathbb{Z}_2\text{-Set}$ -logic is not paraconsistent in the glut sense). *The logic of $\mathbb{Z}_2\text{-Set}$ does not admit any proposition that is both **true** and **false** at any point. The non-contradiction law $P \wedge \neg P = \mathbf{false}$ holds pointwise for all propositions.*

Proof. At any fixed point x of the base space, the truth value of a proposition $P(x)$ is classical: either $P(x) = \mathbf{true}$ or $P(x) = \mathbf{false}$. The third value **swap** arises only when comparing $P(x)$ to $P(\sigma(x))$, which are *different points* related by the involution. The pointwise conjunction $P(x) \wedge \neg P(x)$ is **false** at every point, regardless of whether P takes the value **swap** globally. $\square$

Proposition 8.4 ($\mathbb{Z}_2\text{-Set}$ -logic is not a gap logic). *Every proposition in $\mathbb{Z}_2\text{-Set}$ has a definite truth value at every point. There are no “truth-value gaps.”*

Proof. The subobject classifier of $\mathbb{Z}_2\text{-Set}$ assigns, for every point x of every $\mathbb{Z}_2$ -set, a truth value in $\{\mathbf{true}, \mathbf{false}\}$. The third global value **swap** arises from the *relationship* between values at x and $\sigma(x)$, not from any point’s lacking a value. At no point is truth left undetermined. $\square$

Corollary 8.5 ($\mathbb{Z}_2$ -**Set**-logic is genuinely a fourth category). *The logic of $\mathbb{Z}_2$ -**Set** differs from classical logic (which has no third value at all), from glut paraconsistent logics (which allow local contradiction), and from gap logics (which allow local indeterminacy). Its third truth value arises from the topology of the base space rather than from any modification of local logical structure.*

8.2 The metatheoretical question

A persistent objection to paraconsistent foundations, advanced by Field [14] and Shapiro [15], concerns the metatheory of a paraconsistent logic. The objection runs as follows.

Let L be a paraconsistent logic that tolerates some contradictions, and let $M(L)$ be the metatheory in which L 's rules, soundness, and justification are articulated. Two cases arise:

- If $M(L)$ is *classical*, then the choice of L over classical logic at the object level requires a classical justification, undermining L 's claim to be foundational.
- If $M(L)$ is *paraconsistent*, then arguments in $M(L)$ justifying L can themselves be both valid and invalid. Shapiro [15] argues that dialetheism cannot adequately express itself without a consistent metatheory, so the paraconsistent option is unstable.

This dilemma does not apply to the logic of $\mathbb{Z}_2$ -**Set**, for a structural reason:

Theorem 8.6 (The metatheory of $\mathbb{Z}_2$ -**Set** is classical, consistently). *The derivation of $\mathbb{Z}_2$ -**Set** and its three-valued subobject classifier (Theorem 4.1) uses classical reasoning throughout. The classical reasoning lives in the fixed-point locus of the involution—the region in which the subobject classifier restricts to $\{\mathbf{true}, \mathbf{false}\}$. This is consistent with, rather than opposed to, the three-valued logic of the total structure: the metatheory is the fixed-point restriction of the object-level logic, which is exactly where classical reasoning is valid within the framework.*

Proof sketch. The self-grounding loop of Theorem 4.1 consists of propositions such as “the Möbius connection has holonomy $\mathbb{Z}_2$,” “ $\mathbb{Z}_2$ -**Set** is cartesian closed,” “Lawvere’s theorem applies.” Each of these is a classical mathematical statement evaluated at no particular point of any non-orientable structure—they are propositions *about* the structure, not *within* it. They live in the standard mathematical metatheory, which is a classical topos (e.g., **Set**, viewed as the trivial $\mathbb{Z}_2$ -**Set** where $\sigma = \text{id}$ everywhere).

Within **Set**, classical logic holds and all the derivations proceed as standard mathematics. The three-valued logic appears only when reasoning *inside* a non-trivial $\mathbb{Z}_2$ -**Set**-object—that is, when evaluating propositions at points where $\sigma \neq \text{id}$.

The framework’s structural claim is precisely that the fixed-point locus of the involution is where classical reasoning is legitimate, and that classical metatheoretical reasoning about $\mathbb{Z}_2$ -**Set** is an instance of such reasoning. The metatheory is classical because it operates in the region where classical logic is structurally warranted, not despite the three-valued object-level logic. $\square$

Remark 8.7 (The recursive structure). The relationship between the metatheory and the object theory is recursive in a well-behaved way:

- Object-level logic: three-valued, governing propositions within non-orientable structures.
- Meta-level logic: classical, governing propositions about the structures themselves.
- Classical logic is the *restriction* of three-valued logic to the fixed-point locus.
- The fixed-point locus is where propositions are $\mathbb{Z}_2$ -invariant—where the involution acts trivially.

The metatheory is not a separate, external logic; it is the classical sublogic of the three-valued logic, restricted to propositions the involution fixes. This dissolves Field’s dilemma: there is no choice between a classical and a paraconsistent metatheory, because the classical metatheory is a derived substructure of the three-valued object theory.

Remark 8.8 (Dissolving Shapiro’s concern). Shapiro’s concern is that a dialetheist cannot articulate dialetheism without presupposing a consistent metatheory, since the articulation invokes notions (proof, truth, negation) that must function consistently in order to state the thesis at all. This concern does not apply to $\mathbb{Z}_2$ -**Set**-logic, because $\mathbb{Z}_2$ -**Set**-logic is not dialetheic: it does not assert any contradictions (Proposition 8.3). The framework makes no contradictory claims anywhere—it claims that truth values can vary across a non-orientable base space, which is a consistent and geometrically precise statement expressible in classical metatheory.

8.3 Relationship to established paraconsistent traditions

While $\mathbb{Z}_2$ -**Set**-logic is not itself paraconsistent, it bears on the philosophical project of paraconsistent logic in a specific way: it provides a geometric grounding for three-valued logical structure that avoids both the glut and gap options. The third truth value is neither a contradiction to be tolerated nor an indeterminacy to be accepted, but a topological feature to be described. Its mathematical content is fully determinate; what it lacks is globalization, not local determinacy.

This reframes a long-standing foundational debate. The motivation for paraconsistent logic has often been the resolution of self-referential paradoxes (Russell, the Liar, the Sorites). The standard paraconsistent move—tolerating contradiction—has foundational costs that Field and Shapiro have identified. The $\mathbb{Z}_2$ -**Set** alternative resolves the same paradoxes while avoiding those costs: the paradox-generating operations produce **swap** truth values that fail to globalize, rather than **true** $\wedge$ **false** contradictions that violate non-contradiction.

Remark 8.9 (Why earlier paraconsistent approaches could not see this). The reason paraconsistent logic has not previously identified the orientation-dependent option is that traditional paraconsistent investigations have proceeded from logical modifications (what axioms to drop or add, what valuations to permit) rather than from geometric structures (what spaces and connections produce logical features as emergent consequences). The $\mathbb{Z}_2$ involution and its three-valued classifier do not appear as natural objects in the logical literature because they arise from a different starting point: a specific geometric object (the Möbius connection) whose topos is computed, and whose logic is then found to be three-valued. The three-valued logic is *discovered* rather than *designed*.

9 The Kunen Barrier

9.1 The classical result

The large cardinal hierarchy [7] measures the consistency strength of axioms asserting elementary embeddings $j: V \rightarrow M$ where M is an inner model. The hierarchy’s natural limit is $j: V \rightarrow V$: a nontrivial elementary embedding of the universe into itself. A cardinal witnessing such an embedding is called a **Reinhardt cardinal**.

Theorem 9.1 (Kunen, 1971 [2]). *Assuming ZFC, there is no nontrivial elementary embedding $j: V \rightarrow V$.*

The proof constructs a critical sequence $\kappa, j(\kappa), j^2(\kappa), \dots$, defines $\lambda = \sup\{j^n(\kappa) : n \in \omega\}$, and uses the Axiom of Choice to well-order V_λ , deriving a combinatorial contradiction from j ’s action on $V_{\lambda+1}$.

9.2 Logical dependencies

The proof depends on three properties that are present in ZFC but not in $\mathbb{Z}_2$ -**Set**:

Proposition 9.2 (Choice obstruction). *In $\mathbb{Z}_2\text{-Set}$, the axiom of choice fails globally. A well-ordering of the total structure requires a globally consistent orientation (selecting which chart’s ordering “wins” on overlaps), which the $\mathbb{Z}_2$ holonomy obstructs.*

Choice holds locally: any object in the fixed-point locus can be well-ordered. But the global well-ordering of V_λ required in Kunen’s proof is unavailable.

Proposition 9.3 (Well-foundedness obstruction). *The critical sequence $\kappa, j(\kappa), j^2(\kappa), \dots$ is the orbit of κ under j . Analysis of this orbit—in particular, the existence and properties of the supremum λ —requires that ordinals are globally well-ordered. By the Burali-Forti resolution (Theorem 6.2), the ordinals in $\mathbb{Z}_2\text{-Set}$ are locally well-ordered but not globally so. The supremum λ , defined as a global ordinal-theoretic operation, is not necessarily well-defined.*

Proposition 9.4 (Logic obstruction). *Kunen’s proof proceeds by contradiction, using excluded middle throughout. In $\mathbb{Z}_2\text{-Set}$, the internal logic is classical for propositions taking values in $\{\text{true}, \text{false}\}$ but not for propositions taking the value **swap**. Key steps of the proof—particularly those involving membership relations at the scale of V_λ —may involve sheet-dependent propositions for which classical reasoning does not apply.*

Theorem 9.5 (The Kunen barrier does not apply in $\mathbb{Z}_2\text{-Set}$). *Each of the three obstructions (Propositions 9.2–9.4) independently blocks Kunen’s proof. Together, they imply that the theorem $\neg \exists j: V \rightarrow V$ is not derivable in the internal logic of $\mathbb{Z}_2\text{-Set}$.*

Remark 9.6 (Strength of the claim). Theorem 9.5 establishes that Kunen’s *proof* does not apply. It does not by itself prove that $j: V \rightarrow V$ is consistent with $\mathbb{Z}_2\text{-Set}$ (which would require constructing a model) or that such embeddings exist in $\mathbb{Z}_2\text{-Set}$ (which would require further argument). We note, however, that a non-orientable structure naturally carries self-maps: the $\mathbb{Z}_2$ involution σ defining any object of $\mathbb{Z}_2\text{-Set}$ is a nontrivial endomorphism. The total structure of $\mathbb{Z}_2\text{-Set}$, equipped with its global involution, already possesses self-referential maps of the type that Kunen’s theorem would prohibit. Whether these can be promoted to full elementary embeddings is an open question (Open Question 10.2).

Remark 9.7 (The geometric novelty). It is well known that Kunen’s theorem depends on the Axiom of Choice, and that dropping AC lifts the barrier: this is the starting point of the investigation of Reinhardt cardinals in ZF. The contribution of the present framework is *not* the observation that AC failure permits $j: V \rightarrow V$ —that observation is standard. The contribution is identifying a *specific geometric origin* for the AC failure.

In the standard ZF-without-Choice treatment, AC is simply absent as an axiom. The question “why might AC fail?” has no principled answer; AC is an axiom that may or may not be assumed. The question “what structure does the universe have if AC fails?” is also open: ZF without AC admits many models, none of them canonical.

In the present framework, AC failure is *forced* by the non-orientable topology of the underlying geometric object. Global well-ordering is obstructed not by the absence of an axiom but by the presence of $\mathbb{Z}_2$ holonomy, which is in turn forced by Lawvere’s theorem applied to any self-referentially rich category (Sections 2–3). The universe of $\mathbb{Z}_2\text{-Set}$ is a canonical object, uniquely determined by the geometric and logical constraints of the construction.

This distinction is substantive. The framework’s Kunen dissolution is not “AC fails, so of course Kunen lifts,” but rather “the geometric structure that forces the three-valued logic also forces AC to fail globally, and the specific pattern of failure is determined by the $\mathbb{Z}_2$ involution rather than being arbitrary.” The same non-orientability that obstructs global well-ordering provides the nontrivial self-map σ , suggesting that self-embedding and AC failure are not independent phenomena but joint consequences of a single geometric structure.

The open question for large cardinal theory, reframed through this lens, is whether the canonical non-orientable structure of $\mathbb{Z}_2\text{-Set}$ is the *right* ZF-without-Choice universe for studying Reinhardt-type phenomena—whether the historically motivated Reinhardt cardinals correspond to the self-maps that non-orientability naturally provides.

10 Discussion

10.1 Uniform mechanism

The three resolutions exhibit a uniform pattern:

Paradox	Operation	Local	Global	Blocked step
Burali-Forti	Well-ordering	Valid	Obstructed	(BF2)
Cantor	Diagonalization	Valid	Obstructed	(C3)
Kunen	Choice + classical logic	Valid	Obstructed	(K4)

In each case: (a) the totality (Ω , V , or the self-embedding target) exists; (b) local structure is fully classical; (c) the paradox-generating operation produces a **swap** truth value and fails to globalize; (d) the paradox dissolves without axiomatic prohibition.

The uniformity follows from the common origin: all three paradoxes are instances of Lawvere’s fixed point theorem (Section 2), all involve $\mathbb{Z}_2$ involutions (Proposition 2.2), and all are resolved by the three-valued subobject classifier that the $\mathbb{Z}_2$ action generates (Proposition 3.5).

10.2 Relationship to ZFC

The non-orientable framework is not a replacement for ZFC but a geometric explanation of why ZFC’s axioms take the form they do. The axioms of Foundation, Separation, and Replacement—which collectively prevent paradoxical totalities—encode, at the axiomatic level, the topological constraints that non-orientability provides geometrically.

Within the fixed-point locus of the $\mathbb{Z}_2$ action, all ZFC axioms hold, classical logic applies, and standard mathematics proceeds without modification. The non-orientable framework extends beyond ZFC only at the scale of self-referential totalities—precisely the scale where ZFC invokes the set/proper class distinction.

10.3 Relationship to other foundational frameworks

- **Homotopy Type Theory (HoTT):** $\mathbb{Z}_2\text{-Set}$ can be viewed as a model of HoTT where the relevant path structure is $\mathbb{Z}_2$ -valued. The three truth values correspond to paths that are trivially true, trivially false, or nontrivial (the “swap” path).
- **Non-well-founded set theories** (Aczel’s AFA [10]): These permit circular membership. The non-orientable framework permits similar self-referential structure but constrains it geometrically (via the involution) rather than axiomatically.
- **Paraconsistent set theories** (Priest’s LP [12] and related): These tolerate controlled contradictions. The non-orientable framework is *not* paraconsistent in this sense (Proposition 8.3)—the **swap** value is not “both true and false” but “true on one sheet, false on the other,” which is an orientation-dependent rather than contradictory property. The framework thereby resolves paradoxes without incurring the metatheoretical difficulties identified by Field [14] and Shapiro [15] (Section 8).

- **Topos theory [8]:** $\mathbb{Z}_2\text{-Set}$ is a topos—one of the simplest nontrivial examples. The present paper applies standard topos-theoretic results to this specific topos and draws foundational consequences. Individual classical topoi (equivalent to **Set**) arise as the fixed-point locus of the involution.

10.4 Open problems

Open Question 10.1 (Consistency). Is the non-orientable framework consistent relative to ZFC? The topos $\mathbb{Z}_2\text{-Set}$ is a well-studied mathematical object whose consistency is not in doubt. The foundational claims of this paper concern the *interpretation* of $\mathbb{Z}_2\text{-Set}$ as a setting for set-theoretic practice. The question is whether this interpretation introduces inconsistencies beyond those already known to be absent in $\mathbb{Z}_2\text{-Set}$ itself.

Open Question 10.2 (Reinhardt cardinals). Does $\mathbb{Z}_2\text{-Set}$ (or a suitable extension) provide a model in which Reinhardt cardinals are consistent? The $\mathbb{Z}_2$ involution of $\mathbb{Z}_2\text{-Set}$ provides a natural nontrivial self-map of the total structure. Whether this self-map can be promoted to a full elementary embedding is the key question.

Open Question 10.3 (The continuum hypothesis). What is the status of the continuum hypothesis in the internal logic of $\mathbb{Z}_2\text{-Set}$? The three-valued subobject classifier introduces structure beyond that of **Set**. Whether this additional structure determines a definite truth value for CH is an open question with direct bearing on the independence results of Gödel [5] and Cohen [6].

Open Question 10.4 (Scope of the $\mathbb{Z}_2$ resolution). The present paper resolves three paradoxes (Burali-Forti, Cantor, Kunen) via a single mechanism (the **swap** truth value obstructing globalization). To what extent does this mechanism apply to other independence results and undecidability phenomena in mathematics? The companion paper [16] identifies compensatory uses of the Axiom of Choice in the construction of the number systems and shows that the Möbius connection provides canonical replacements. A systematic investigation of the relationship between $\mathbb{Z}_2$ holonomy and AC-dependent constructions across mathematics remains open.

Open Question 10.5 (Arithmetic extension and constructive set theory). The topos $\mathbb{Z}_2\text{-Set}$ captures the non-orientability of the Möbius connection but not its full arithmetic structure. The connection’s multiplicative sector—prime-indexed transport cycles and their intersection lattice—carries richer structure than the rank-1 scalar fibers of $\mathbb{Z}_2\text{-Set}$. A natural extension of the present framework would treat sets constructively as intersection patterns of prime transport cycles, with the Möbius inversion formula serving as the inclusion-exclusion principle for this lattice. This arithmetic enrichment connects directly to the L-function classification of the Langlands program, where each L-function corresponds to a coherent response of higher-rank fibers to prime holonomy. This extension is the subject of forthcoming work.

References

- [1] F. W. Lawvere, *Diagonal arguments and cartesian closed categories*, in: Category Theory, Homology Theory and their Applications II, Lecture Notes in Mathematics **92**, Springer, 1969, pp. 134–145.
- [2] K. Kunen, *Elementary embeddings and infinitary combinatorics*, J. Symbolic Logic **36** (1971), 407–413.
- [3] G. Cantor, *Über eine elementare Frage der Mannigfaltigkeitslehre*, Jahresbericht der DMV **1** (1891), 75–78.

- [4] C. Burali-Forti, *Una questione sui numeri transfiniti*, Rendiconti del Circolo Matematico di Palermo **11** (1897), 154–164.
- [5] K. Gödel, *The consistency of the axiom of choice and of the generalized continuum-hypothesis*, Proc. Natl. Acad. Sci. USA **24** (1938), 556–557.
- [6] P. Cohen, *The independence of the continuum hypothesis*, Proc. Natl. Acad. Sci. USA **50** (1963), 1143–1148.
- [7] A. Kanamori, *The Higher Infinite*, 2nd ed., Springer, 2003.
- [8] S. Mac Lane and I. Moerdijk, *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*, Springer, 1992.
- [9] The Univalent Foundations Program, *Homotopy Type Theory: Univalent Foundations of Mathematics*, Institute for Advanced Study, 2013.
- [10] P. Aczel, *Non-Well-Founded Sets*, CSLI Lecture Notes, 1988.
- [11] E. Steinitz, *Algebraische Theorie der Körper*, J. Reine Angew. Math. **137** (1910), 167–309.
- [12] G. Priest, *The logic of paradox*, J. Philosophical Logic **8** (1979), 219–241.
- [13] S. C. Kleene, *Introduction to Metamathematics*, North-Holland, 1952.
- [14] H. Field, *Saving Truth from Paradox*, Oxford University Press, 2008.
- [15] S. Shapiro, *Incompleteness and inconsistency*, Mind **111** (2002), 817–832.
- [16] A. K. Morgan, *The Möbius connection as canonical foundation: orientation, arithmetic, and the Riemann Hypothesis*, in preparation, 2026.