

AMS/MAA | PROBLEM BOOKS

Doorway to Math Olympiads

Arjun Agarwal

The study of mathematics, like
the Nile, begins in minuteness
but ends in magnificence.

Charles Caleb Colton

The essence of mathematics is
not to make simple things
complicated, but to make
complicated things simple.

S. Gudder

© 2023 Arjun Agarwal

Text licensed under CC-by-SA-4.0

This is (still!) an incomplete draft.

Please send corrections, comments, pictures of kittens, etc.

to anulick0@gmail.com

Last updated March 26, 2024.

*To Aparna Thakur Ma'am, Sylvia Marcarhanas Ma'am, Prashant Jain Sir
and Abhay Mahajan Sir, who have taught me all the math I know.*

Contents

Preface	xii	
0.1	Why does this book exist?	xii
0.2	Secret Sauce	xii
0.3	A small history	xiii
How to use this book	xv	
0.4	Prerequisites	xv
0.5	Structure of the book	xv
0.6	Approach	xvi
0.7	Bluntly speaking...	xvi
Part 1 Sales Pitch	1	
1 Introductory Problems	3	
1.1	Introduction	3
	Exercises	3
Part 2 Combinatorics	7	
2 Permutations and Combinations and a pinch of Probability	9	
2.1	Fundamental Principle of Counting	10
2.2	Equivalence	10
2.3	The special sign!	12
2.4	Criminal Lineups	13
2.5	Circles	14
2.6	Team selection	15
2.7	Subsets	16
2.8	Probability	17
2.9	Exercise	17
	Exercises	17
3 Some methods of counting	21	
3.1	Casework	21

3.2	Complementary Counting	22
3.3	Principle of Inclusion and Exclusion	22
3.4	Exercises	25
	Exercises	25
4	A guessing game	29
	Exercises	29
5	Stars, Bars and Hockey Sticks	33
5.1	Stars and Bars	33
5.2	Counting in two ways	34
5.3	Pascal's Triangle	36
5.4	Exercises	38
	Exercises	38
6	Geometrical Combinatorics	41
6.1	Geometric Counting	41
6.2	Geometric Probability	42
	Exercises	43
Part 3 Down the Rabbit Hole		47
7	Power Up Unlocked!	49
7.1	Recursion	49
7.2	Fibonacci Sequence	50
7.3	Solving Liner Recursion	51
7.4	Binet's Formula	52
7.5	But Wait, what if the roots repeat?	53
7.6	Catalan Number's	54
7.7	Incidence Matrices	55
	Exercises	57
8	Graph Theory	61
8.1	Definitions	61
8.2	Paths and Walks	63
8.3	Trees	64
8.4	Real life applications	66
	Exercises	67
9	A New Breed of Counting	71
9.1	Pigeon Hole Principle	72
9.2	Expected Value and Probabilistic Method	74
	Exercises	76
	Exercises	78

Contents	vii
Part 4 Algebra	79
10 Algebraic Manipulations	81
10.1 Binomial Theorem	81
10.2 Common Expansions and Factorization	83
10.3 More Factorization Tricks	84
10.4 Quadratic Equations	86
10.5 Vieta's Formula	88
10.6 The Fundamental Theorem of Algebra	89
10.7 Newton's Sums	91
10.8 Reciprocal Relations	93
10.9 Some Special Polynomials	94
Exercises	95
11 Inequalities	99
11.1 Another note on notation	99
11.2 AM-GM and Muirhead	99
11.3 Non-homogeneous equations	102
11.4 Some advanced inequalities	104
11.5 A note	106
Exercises	106
12 Sequence and Series	109
12.1 Some notes on notation	110
12.2 Arithmetic Progression	110
12.3 Geometric Progression	113
12.4 Induction and Summation	115
12.5 Telescoping	116
12.6 Recurrence Series	121
Exercises	122
Part 5 The Red Pill	125
13 Calculus I: Limits, Continuity and Derivative	127
13.1 The Rate of Change	128
13.2 Some functions	129
13.3 Inverse Trigonometric functions	133
13.4 Derivative	134
13.5 Some more notation	136
13.6 Product and Quotient rules	136
13.7 Chain rule	138
13.8 Limits	141
13.9 Continuity	149
13.10 Application of Differentiation	150

Exercises	152
14 Calculus II: Indefinite Integration	155
14.1 The Fundamental nature of graphs	156
14.2 Integration	158
14.3 Techniques of indefinite Integration: Substitution	158
14.4 Techniques of indefinite Integration: Integration by Parts	185
14.5 Definite integration	189
14.6 King's Rule	190
14.7 Leibnitz Theorem	192
14.8 Taylor Maclaurian Series	193
14.9 Differential Equations	193
14.10 What's more?	195
Exercises	195
15 Ch-14 Linear Algebra	199
15.1 Determinants	200
15.2 Properties of Determinants	203
15.3 Application of Determinants	208
15.4 Crammer's Rule	210
15.5 Types Matrices	212
15.6 Arithmetic of Matrices	213
15.7 The bridge	216
15.8 Some more special matrices	217
15.9 Adjoint and Inverse of Matrices	218
15.10 System of linear Equations using Matrices	220
15.11 Characteristic Equation and Cayley Hamilton Theorem	221
15.12 Netflix and Spotify and Matrices...	222
Exercises	222
16 Inequalities Revisited	225
16.1 Rearrangement Inequalities	225
16.2 Inequalities in Arbitrary Functions	227
16.3 Tangent Line Trick	230
16.4 SEBACS Generalized	231
16.5 Lagrange Multipliers	232
16.6 Sum uses of Calc	234
Exercises	237
Part 6 Number Theory	239
17 Wearing the Crown of Mathematics	241
17.1 Division	242
17.2 Congruence Modulo	246

17.3	Prime Numbers	248
17.4	Number Bases	251
17.5	Divisibility rules	253
17.6	Nature of Factors	254
17.7	Legendre's Theorem	255
17.8	Irrationality	256
	Exercises	256
18	Modular Arithmetic	261
18.1	Modulo Inverse	261
18.2	Fermat's Little Theorem	262
18.3	Euler's Totient Theorem	263
18.4	Wilson's Theorem	264
18.5	Chinese Remainder Theorem	266
	Exercises	268
19	Functional Equations	271
19.1	Some definitions	271
19.2	Basic Functional Equations	272
19.3	Some methods of solving	273
19.4	Cauchy's Functional Equations	275
19.5	Checklist for Functional Equation Solving	277
	Exercises	277
20	Diophantine Equations	281
20.1	Linear Diophantine Equations	281
20.2	Parity Arguments	282
20.3	Algebraic methods	283
20.4	Modular Contradiction Method	285
20.5	Pythagorean Triplets	286
20.6	Infinite Descent	289
20.7	Vieta Jumping	290
	Exercises	291
Part 7	The Number's Awaken	293
21	Bazooka!	295
21.1	Surprisingly, not complex!	295
21.2	Orders	297
21.3	Chicken McNugget Theorem	298
21.4	Pell's Equations	300
21.5	Floor, Ceiling and fractional function	302
21.6	Black Boxes	303
	Exercises	308

22 Constructions	311
22.1 Chinese Remainder Theorem	311
Exercises	312
Part 8 Geometry	313
23 Angle Chasing	315
23.1 Triangle Centers	315
23.2 Cyclic Quadrilaterals	316
Part 9 Extras	317
24 Anti-problems	319
24.1 Anti-Problems	320
Exercise	320
25 Engineering	321
25.1 Abuse of Degree of Freedom	322
Part 10 Appendix	323
A Appendix A: Justifications, Ramifications and Excuses	325
A.1 Introductory Problems	325
B Appendix B: Hints	327
C Appendix C: Selected Solutions	331
D Appendix D: Borrowed Brilliance	333
D.1 Almost every part	333
D.2 Introductory Problems	334
D.3 Permutations and Combinations	335
D.4 Down The Rabbit Hole	335
D.5 Algebra	336
D.6 The Red Pill	336
D.7 Number Theory	337
D.8 The Numbers Awaken	339
E Appendix E: Problem Sources	341

Preface

0.1 Why does this book exist?

As Austin Kleon wisely put it, “Write the book you’d want to read.”

Well, here I am, crafting a math adventure that’s as enjoyable as your favorite Netflix series (okay, maybe not that enjoyable, but close).

I’ve been diving into math contests since grade 3, where “speed math” was my jam. Fast forward to 8th grade, and I stumbled upon Olympiad math, and it was love at first equation. My evenings turned into PRMO paper-solving sessions with friends (who were probably having more fun looking at me cursing the problem, than actually doing the problems). I even found myself deep in calculus on a lazy Sunday just because a tricky problem caught my eye.

However, my math journey, while thrilling, was a bit like a rollercoaster with no seatbelt — wild and unpredictable. I’d study whatever piqued my curiosity, sometimes going way beyond what was needed. That led me down rabbit holes like exploring the mystical nature of π in grade 4 and sneakily applying calculus in a non-calc test in grade 10. Oh, and let’s not forget the countless distractions like sketch videos and attempting problems way beyond my level, only to ponder the meaning of life afterward.

But wait, there’s hope! This book is my attempt at a cure. It’s the guide I desperately wished I had during my math escapades.

0.2 Secret Sauce

If you’ve ever stumbled upon the Wikipedia pages of IMO gold medalists, you might notice a common thread — one or both of their parents likely hold fancy STEM degrees. Now, before you roll your eyes, there’s a simple reason behind this: success in math Olympiads often starts early and requires some serious elbow grease.¹

¹Polgár, L. (1989). Bring Up Genius! Budapest, H. v1.1 2017-07-31

They say it takes a whopping 10,000 hours of practice to become an expert at something.² Most of us didn't even consider real math until later in life. I mean, till grade 7, my idea of math was calculating the total of the menu of the restaurant before the starters arrived! So, hitting that 10,000-hour goal for Olympiad-level math is a bit of a stretch.

But hold on, there's more to it than just clocking in the hours. To improve, we need feedback, genuine feedback. Unfortunately, most of our well-meaning teachers, parents, and pals struggle to tackle even the simplest Olympiad questions. Case in point:

Example 0.1 (IMO 2023, P1). Determine all composite integers $n > 1$ that satisfy the following property: if d_1, d_2, \dots, d_k are all the positive divisors of n with $1 = d_1 < d_2 < \dots < d_k = n$, then d_i divides $d_{i+1} + d_{i+2}$ for every $1 \leq i \leq k - 2$.

They mean well, but let's face it — their feedback can be as skewed as a wonky compass. Only the best teachers are willing to go the extra mile to meet your needs and help you out. But that's often not enough. Why, you ask?

Well, studies show that anesthesiologists get better at their job over time, while radiologists don't improve much from their initial performance. Why? Anesthesiologists get instant feedback, while radiologists have to rack their brains to figure out what went wrong weeks ago.³

The second secret sauce is a valid environment and deliberate practice. We can't just study random topics and solve random problems and expect to shine. We need a structured plan for learning and practice that pushes us just beyond our comfort zone.⁴ Remember, it's like chess players vs. roulette players. After 10 years, a chess player is a chess wizard, while a roulette player ...well, they're just spinning a wheel and praying. We don't want to leave our Olympiad journey to chance, do we?

For the final secret sauce, we turn to the school system. Now, I know the school system has its flaws, but there's one thing it excels at: repetition. Repetition is the secret sauce to improvement. Chess grand masters can play the entire game blindfolded imagining the board and pieces in their head, but if you give them a random arrangement of pieces(which can't occur in a chess

²From Malcom Gladwell's *Outliers: The story of success*

³Goldberg, S. B., Rousmaniere, T., Miller, S. D., Whipple, J., Nielsen, S. L., Hoyt, W. T., & Wampold, B. E. (2016). Do psychotherapists improve with time and experience? A longitudinal analysis of outcomes in a clinical setting. *Journal of Counseling Psychology*, 63(1), 1.

⁴Ericsson, K. A., Krampe, R. T., & Tesch-Römer, C. (1993). The role of deliberate practice in the acquisition of expert performance. *Psychological Review*, 100(3), 363

game) they are just as bad as remembering it as everyone else.⁵ By repeating the same things, we can start seeing the patterns, making our brains better at dealing with them. We need to do the same with problems.

We've got the complete pedagogy of every chapter laid out in Appendix A for all the teachers out there who might be using this text or for the curious learners looking to understand our secret sauce, ingredient by ingredient.

0.3 A small history

This book's journey began back in class 9, during a car trip to my sister's wedding venue. However, it took a little hiatus for two years while I tackled my boards and sailed through 11th grade. But come 12th grade, I had a wild idea—teaching Olympiad math. And as I dove into teaching, my notes slowly evolved into what you're holding now.

As I write this latest version of the preface, the book is gearing up for its first round of updates—more hints, more questions, a dash more theory, and answers to select questions. So, without further ado, let the math adventures begin!

Happy reading,

Warm regards,
Arjun Agarwal

⁵Chase, W. G., & Simon, H. A. (1973). Perception in chess. *Cognitive psychology*, 4(1), 55-81

How to use this book

0.4 Prerequisites

Don't worry, I am not expecting you to perform calculus in your sleep or recite π to a thousand decimal places. But I do have some prerequisites, though, just to make sure you're in the right ballpark.

First off, you should be on friendly terms with basic arithmetic. Yep, that means adding, subtracting, multiplying, and dividing numbers should be old hat for you. I won't judge if you occasionally rely on your trusty calculator, but let's aim for some proficiency here.

Now, about those prime numbers and factoring and HCF and LCM. I'll dive into them later, but it'd be grand if you knew the basics. No sweat if you don't; I've got your back.

But wait, there's more! Beyond crunching numbers, you'll need to use a part of brain. A hefty dose of logical thinking will be used. I am talking creativity, the same thing you use in art and painting(yes, it's used in math as well).

Finally, You need to be ready to swap a bit of your overconfidence for some bona fide thinking prowess. If you say yes, then this book is your doorway to math olympiad.

0.5 Structure of the book

Every chapter has a few pages detailing the techniques and mathematical tools for that chapter. The examples which are not followed by solutions are normally quite simple and meant to reinforce what you just read.

The examples followed by the answer are typically harder and meant to teach a problem solving concept.

The problems at the end of the chapter start easy and move upward. They are meant to challenge you and get you out of the comfort zone while giving you the practice you require.

The hints are numbered and appear in random order in Appendix B, and selected solutions in Appendix C. I have also tried to include the sources of the

problems, so that a diligent reader can find more solutions online (for example on the Art of Problem Solving forums). A full listing of contest acronyms appears in Appendix D.

0.6 Approach

Readers are encouraged to not be bureaucratic in their learning and move around as they see fit, e.g., skipping complicated sections and returning to them later, or moving quickly through familiar material.

However, I recommend not skipping problems out of anger and instead spending time before looking at the solution. However, I also don't recommend obsessing over a problem or being embarrassed of using the solutions or hints.

0.7 Bluntly speaking...

Use Paper

See I care about the environment, but using paper while you read this book scratching notes and solving questions on paper rather than in your mind, is essential.

A few papers won't kill the earth (and you can offset the pollution simply by planting a single tree which will give you 10000 sheets of paper⁶, You'll probably end up using less than that.)

You are not God. You cannot keep everything in your head (and if you think you can, you'll get proven wrong quite quickly), so please do yourself a favor and **use paper**.

⁶Ribble Packaging Ltd. (2022). How much paper comes from one tree?

Part 1

Sales Pitch

1

Introductory Problems

1.1 Introduction

In this chapter, we have gathered 12 logic problems. To solve them, you do not need anything but common sense and the simplest calculation skills. However, that is not to say that the problems are easy, quite to the contrary. You might find yourself struggling after the first 2 or 3. However, the reason we included the particular selection of problems is two-fold. First and foremost, I want to from the start instill that math is more than just a set of formulas—it is a way of thinking in terms of numbers. Secondly, I aim to highlight common errors that often plague problem solvers, but rather than telling you what they are, we have decided to let you make them and find out. With that lets begin.

The ♣ is placed before extensions to problems which require a bit more math than the original problem

Solve till you reach at least [69★]. This exercise is worth [92★].

Exercises

- (1) [2★] A caterpillar crawls up a pole 75 inches high, starting from the ground. Each day it crawls up 5 inches, and each night it slides down 4 inches. When will it first reach the top of the pole? **Hints:** [31](#)
- (2) [2★] In a certain year, there were exactly four Fridays and exactly four Mondays in January. On what day of the week did the 20th of January fall that year? **Hints:** [49](#)
- (3) [3★](PROMYS 2023) How many boxes are crossed by a diagonal in a rectangular table formed by 199×991 small squares? ♣[5★] Try generalizing for a diagonal passing through $m \times n$ grid. **Hints:** [67](#)

- (4) [3★] Cross out 10 digits from the number 1234512345123451234512345 so that the remaining number is as large as possible. **Hints:** 18

- (5) [2★](Arvind Gupta) Two friends sit in the mess of IIT-Kanpur. One of them has tea and the other has Khus Soda(A south-east Asian grass which is genetically quite close to lemon grass. Consumed with soda during summer due to its ability to hold electrolytes.)

Our glasses are of 100 ml. Each is filled to 90ml with its respective drink. They both decide to make their friendship stronger by mixing their drinks. The first friend fills the tea glass with khus soda. The glass is then stirred vigorously without loss. Then 10ml of the mixture is added to the khus soda(returning both the glasses to 90ml). Now does the tea have more soda or does the soda have more tea? What if we did the same twice? ♣ [2★] What would happen if we kept doing this again and again for a very long period of time? ♣[9★] Can we generate a function, $f(n)$ for the percentage of khus soda in tea if we have done the process n times? **Hints:** 47 5 2

- (6) [3★]Distribute 127 one dollar bills among 7 wallets so that any integer sum from 1 through 127 dollars can be paid without opening the wallets. **Hints:** 12

- (7) (David Morrin) [9★]You visit a remote desert island inhabited by one hundred very friendly dragons, all of whom have green eyes. They haven't seen a human for many centuries and are very excited about your visit. They show you around their island and tell you all about their dragon way of life (dragons can talk, of course). They seem to be quite normal, as far as dragons go, but then you find out something rather odd. They have a rule on the island that states that if a dragon ever finds out that he/she has green eyes, then at precisely midnight at the end of the day of this discovery, he/she must relinquish all dragon powers and transform into a long-tailed sparrow. However, there are no mirrors on the island, and the dragons never talk about eye color, so they have been living in blissful ignorance throughout the ages. Upon your departure, all the dragons get together to see you off, and in a tearful farewell you thank them for being such hospitable dragons. You then decide to tell them something that they all already know (for each can see the colors of the eyes of all the other dragons): You tell them all that at least one of them has green eyes. Then you leave, not thinking of the consequences (if any). Assuming that the dragons are (of course) infallibly logical, what happens? **Hints:** 22

- (8) (David Morrin) [5★]A rubber band with an initial length of 10 cm has one end attached to a wall. At $t = 0$, the other end is pulled away from the wall at a constant speed of 10 cm/s. (Assume that the rubber band stretches uniformly.) At the same time, an ant located at the end not attached to the wall begins to crawl toward the wall, with a constant speed of 5 cm/s

relative to the band. Will the ant reach the wall? If so, how much time will it take? ♣[5★] What if the rubber band has length L , speed u , and the ant crawls with speed v ; what is the limit for which it will not reach the end? **Hints:** 52 69

- (9) (TED Ed) [5★] You are given twelve coins, eleven of which have the same weight, and one of which has a weight different from the others (either heavier or lighter; you do not know). You have a balance scale. What is the minimum number of weighings required in order to guarantee that you can determine which coin has the different weight and also whether it is heavier or lighter than the rest? ♣ [9★] You are given N coins, $N - 1$ of which have the same weight, and one of which has a weight different from the others (either heavier or lighter; you do not know). You are allowed W weighings on a balance scale. What is the maximum value of N , as a function of W , for which you are guaranteed to be able to determine which coin has the different weight and also whether it is heavy or light? **Hints:** 63 30 10 11

- (10) (Even Chen) [3★] Determine, with proof, the smallest positive integer c such that for any positive integer n , the decimal representation of the number $c^n + 2014$ has digits all less than 5. **Hints:** 68

- (11) (Arthur Engel) [5★] The numbers $1, 2, \dots, 10$ are written on a board. Every minute, one can select three numbers a, b, c on the board, erase them, and write $\sqrt{a^2 + b^2 + c^2}$ in their place. This process continues until no more numbers can be erased. What is the largest possible number that can remain on the board at this point? ♣ [2★] What about if we started with $1, 2, 3, \dots, n$? **Hints:** 41

- (12) (Putnam 2017, A1) [9★] Find the smallest set S of positive integers such that

- (a) $2 \in S$
- (b) $n \in S$ whenever $n^2 \in S$
- (c) $(n + 5)^2 \in S$ whenever $n \in S$

Hints: 29 38

- (13) (Iceland 2009) [9★] A number of persons seat at a round table. It is known that there are 7 women who have a woman to their right and 12 women who have a man to their right. We know that 3 out of each 4 men have a woman to their right. How many people are seated at the table?

Hints: 65

Part 2

Combinatorics

2

Permutations and Combinations and a pinch of Probability

I'll begin this book with what we first learnt math as, counting.

Example 2.1. Cricket t-shirts have 2-digit numbers on them using six possible digits: 0, 1, 2, 3, 4, 5 How many different t-shirts can be formed?

While we can write down all the possible t-shirt numbers and count them, we can do something faster and more beautiful. The first digit can be any of the 6 given digits, and so can the second. As they are completely independent, we can simply multiply them.

So we have $6 \times 6 = 36$ such T-Shirts.

Example 2.2. How many 4 digit numbers can be formed using the digits 1, 2, 3 and 4 (Without repetition).

This question is similar as while we can simply try to write all the numbers and then just count, but it is smarter to do something else.

This means we have 4 choices for the thousands digit. As one number is now used, we only have 3 choices for the hundreds digit. As two numbers are now used, we have only 2 choices for the tens digit. With all other numbers used, the last digit will be used as the one's digit, leaving us with only 1 choice. As all these choices are independent, we say that we can make $4 \times 3 \times 2 \times 1 = 24$

Let's explore this new type of counting a bit more.

2.1 Fundamental Principle of Counting

Example 2.3. A person can travel from city A to city B via Road (Car/Bus/Bike), Train (Express/Mail) or Flight (Economy/Business). In how many ways can a person go from city A to city B ?

Pretty standard, right? We can simply count the ways as Number of Road ways + Number of trains + Number of Flights = $3 + 2 + 2 = 7$. Note that we decided to add here.

Example 2.4. A person can travel from city A to city B via Road (Car/Bus/Bike), Train (Express/Mail) or Flight (Economy/Business). He can further travel from city B to city C via Road (Car/Bus/Bike) and Train (Shatabdi/Express/Mail). In how many ways can a person go from city A to city C via city B ?

Number of Ways to travel from City A to City B are 7. The ways to travel from City B to City C are 6 by similar logic.

But as which way we choose after reaching City B doesn't depends on which way we used to actually get there, we can multiply them to get the total ways as $6 \times 7 = 42$.

The main idea I want you to take here is when do we add and when do we multiply. Here is an exercise for you to check if you got the concept.

Example 2.5. The Hermetian alphabet consists of only three letters: A, B, and C. A word in this language is an arbitrary sequence of no more than four letters. How many words does the Hermetian language contain?

2.2 Equivalence

The definition of what it means to be equal is at heart of counting. For example, lets say I ask a 7 year old how many apples do I have if my mother gave me 2 apples and my father gave me 3 apples and I already had 2 apples to begin with.

We expect the child to count on their fingers and report the answer as 7 apples. But here is my point, I asked the child to count apples, why did the child count their fingers?

The answer is simple, because apples and fingers are equivalent. But that makes no sense.

Apples	Fingers
Apples are a product of the <i>Malus domestica</i> genus, characterized by a spherical or slightly oblate shape. The smooth, hydrophobic epidermis is rich in cutin and epicuticular waxes, forming a protective shield for the succulent parenchyma. The parenchyma contains water-soluble polysaccharides like pectin, providing a balance of tartness from malic acid and sweetness from saccharose and fructose.	The finger is a manual digit with bilateral symmetry and flexible articulation. It comprises metacarpal and phalangeal elements, with a keratinocyte-rich epidermis forming a protective barrier. Precision grip is facilitated by a complex musculature network, and distal phalanges house the nail apparatus. Vascularization sustains metabolic demands, and mechanoreceptors enable tactile perception crucial for haptic interaction.

or as a less rigorous individual would put it, apple and fingers are different, bruh!

So how did we draw an equivalence?

We said that, mathematically, the additive property of both of them is the same. This means, with respect to this property, sheep and apple and pear are all equal because their addition follows the same set of rules. Although, we don't compare apple to pizza as while we can cleanly give someone $\frac{3}{8}$ of a pizza, doing so with the apple is much harder. So if two problems are mathematically equivalent, we can take advantage and do the calculations on the easier to solve the easier problem. Here is an example:

Example 2.6. There are 10 lamps in a hall. Each one of them can be switched on independently. The number of ways in which the hall can be illuminated is:

Example 2.7. Ramin language is a very unique language which has 10 symbols and words are a set of more than one symbol. One word has no repetitions of a symbol and the order of symbols doesn't matter. How many words are in Ramin?

These both questions ask us the number of way we can choose some subset of 10 objects with the size (also called cardinality) of the subset is not 0. However, the first question is much easier to understand as the lamp question as its obvious there that each lamp is either on or off, so we have 2^{10} configurations. As the hall is illuminated, all lamps are off condition is to be removed, so our answer to both the questions is $2^{10} - 1$.

We'll explore this idea further in this chapter as well as in a lot more detail as double counting(in chapter 5) and bijections(in chapter 7). Let's now formalize what we learnt till here.

2.3 The special sign!

Normally when you see a ! sign in a book, it refers to the author trying to be funny. But if it comes after a number it means as follows:

Definition 2.8 (Factorial). $n! =$ Number of way to arrange n distinct objects in a straight line.

This means that by the definition of factorial the example about number of 4 digit numbers which can be made with 1, 2, 3, 4 without repetition can be answered simply as $4! = 24$.

Theorem 2.9. $n! = n \times (n - 1) \times \dots \times 3 \times 2 \times 1 \implies n! = n \times (n - 1)!$

Theorem 2.10 (Vacuous Truth). $0! = 1$

What is vacuous truth? A truth which is true as it being false would make life difficult. This is not the actual definition of the word, but I am avoiding the glaring landmine called mathematical philosophy. The reason for including this is you'll find a lot of bad proofs of the same which challenge the definition of factorial. Any proof of this is ill motivated and wrong despite what it may seem.

The most common one is $1! = 1 = 1 * 0! \implies 0! = 1$ which is false as $0! = 0 * (-1)! = 0$ is contradictory. This is defined much better by the real analysis definition of this, aka the famous Γ (Gamma) function. However, we shall not discuss this in this book as real analysis is outside the scope of olympiads.

Let's now come back to the topic at hand.

Example 2.11. For a three digit number $\overline{ABC} = A! + B! + C!$. Find $A+B+c$.

Solution. I had asked this question to Aparna ma'am when I was in 8th grade. After trying to solve it algebraically, I gave up. Ma'am solved it very quickly by trial and error. It felt like cheating for a long time, till I realized that a lot of math, especially at research level, is reducing the noise and then doing trial and error. I like to call it 'The art of Trial and Error'.

For this question, we need to remember the values of first 7 – 8 factorials. That will happen with practice and repetition. We need to note that our number can obviously not have any number greater than 6 as it is a three digit number. Better yet, we can remove 6 from the pool as adding anything to 720 will automatically give us a 7, 8, 9 in the number which is not possible.

This forces us to have a 5 in the number as if we don't have a $5! = 120$, we can't have a three digit number. This is true as $4! + 4! + 4! = 72$ which is not a three digit number.

$0! =$	1
$1! =$	1
$2! =$	2
$3! =$	6
$4! =$	24
$5! =$	120
$6! =$	720
$7! =$	5040
$8! =$	40320
$9! =$	362880

Here, we will try to determine the number of 5's in \overline{ABC} . If all three are 5, we have $555 = 360$ which is false.

If we have 2 5's, the number will also have a 2 as $5! + 5! = 240$. This implies $5! + 5! + 2! = 242 = 255$ which is false.

This means we have only one 5. This also means we have a 1 in \overline{ABC} . Next, we can do trial error on 0, 4.

$1! + 5! + 0! = 122$	false
$1! + 5! + 1! = 122$	false
$1! + 5! + 2! = 123$	false
$1! + 5! + 3! = 127$	false
$1! + 5! + 4! = 145$	true

Thus, the $\overline{ABC} = 145$. Thus, $A + B + C = 1 + 4 + 5 = 10$. \square

2.4 Criminal Lineups

If we have 5 suspects lined up, we can arrange them in $5! = 120$ ways. But if 2 of them are wearing squid game masks them? Now they are identical and hence interchangeable. The ways to arrange them now will be halved as they both being switched doesn't create a new permutation.

Now what if the remaining three of them wear Joker masks. We'll have to divide the permutations by $3! = 6$ as all of them are identical. This can obviously be generalized to:

Theorem 2.12. *When arranging a total of a objects, where there are groups of identical objects denoted by k, l, m, n , etc., the number of distinct arrangements is given by dividing the factorial of the total number of objects ($a!$) by the product of the factorials of the counts of each group of identical objects ($k!, l!, m!, n!$, etc.).*

Let's try to understand this more clearly through an example.

Example 2.13. How many ways can the letters of the word 'AGARWAL' be arranged?

Solution. My surname has 3 A's and 4 distinct letters. Let's consider the A's distinct as A_1, A_2, A_3 . Then we have, $7!$ arrangements. However, the A's are not distinct. Thus, the arrangement of A's doesn't matter. We have $3!$ arrangement of A's. Thus, we can divide it from the total to get the actual number of total arrangements.

Thus, the answer is $\frac{7!}{3!}$

□

The above question comes regularly in collage entrance exams. A better question type which also can be found on collage entrances is:

Example 2.14. Using all the letters M, O, P, R , and X , we can form five-letter "words". If these "words" are arranged in alphabetical order, then what position does the "word" $PRMOX$ occupy?

Solution. We can first look at all words starting with letter M which is alphabetically the first.

There are $4!$ such words. Similarly for O brings the total to 48

In words staring with P we first look at words which start with PM which we have $3!$ of. Then PO .

This brings the total count to $48 + 12 = 60$

$PRMOx$ is the 61st word as after PR the rest of the letters are in alphabetically order and hence, it is the very next word.

□

We can, without much avail, try to make these questions look more difficult by having repeat letters. But as you'll find upon solving, the question is still rather trivial.

Example 2.15. If letters of the word 'JUGNU' are arrranged to form all possible words, what is the rank of 'JUGNU' in the list alphabetically?

2.5 Circles

Example 2.16. A stadium is to have the flags of 12 different teams arranged around the ground in a circle. In how many ways can it be done?

Solution. We need to notice that the arrangements need to be unique upto rotation but not to flipping. This means that if two arrangements are such that we can rotate them to obtain the other, they are the same. Flipping means that if we move the flags about the diagonal. Obviously, once the

flags are put, we can rotate them by changing our viewpoint, but we can realistically not flip them.

Let's now move to the solving. If rotation was not possible, we would have $12!$ arrangements. But as rotation is possible, in all the 12 arrangements, only 1 will be unique.

This means we have $\frac{12!}{12} = 11!$ arrangements. □

What would happen if we could flip the stadium as well? For the sake of imagination, let's say you have 12 gemstones. You want to string them into a necklace. What is the number of ways to do so?

We obviously would have only $11!$ ways if flipping was not possible. But as it is possible, the two flipped permutations are now considered the same.

This means we have $\frac{11!}{2}$ ways to make the necklace.

Let's generalize the two problems we just solved:

Theorem 2.17. *The number of ways of arranging n objects in a circle where rotations of the same arrangement are not considered distinct is $(n - 1)!$*

Theorem 2.18. *The number of ways of arranging n objects in a circle where rotations of the same arrangement are not considered distinct and reflections of the same arrangement are not considered distinct is $\frac{(n-1)!}{2}$*

Circular counting happens to be much more complicated than normal once we start having repeated elements. We will solve such questions using casework in the next chapter and then generalize a formula using group actions later.

2.6 Team selection

Your school is having an inter-class cricket tournament. From every class of 30 we need to choose 11 players. How many ways can we do it?

For the first players we have 30 choices, then 29 and so on. But that's not all. The order in which players are chosen doesn't matter as they are a team in the end. So we need to divide it in the end by $11!$. So the number of possible teams will be $\frac{30 \times 29 \times 28 \dots 22 \times 21 \times 20}{11!} = 54,627,300$. Generalizing:

Theorem 2.19. *Number of ways of choosing k objects from n , where order doesn't matter is $\binom{n}{k} = \frac{n!}{k!(n-k)!}$*

You might remember that we have already reached this theorem in the examples we solved right at the start. We are just happening to formalize this here.

Also notice that $\binom{n}{k} = \binom{n}{k-n}$ as ways of choosing k things to be selected is the same as choosing $n - k$ things to not be selected.

We will also add, subtract and do a bunch of things with it later.

2.7 Subsets

A set is a collection of things. A subset is a smaller collection of things all of which are part of the set it is subset of.

Example 2.20. If a set has n distinct elements in it, How many subsets of that set exist?

Solution. Every element is either in the subset or not in it. Hence we have two possibilities for every element. Hence we can say 2^n subsets exist. \square

Theorem 2.21 (Subset Theorem). *The number of subsets of a set of size n is 2^n .*

Another theorem which we happened to derive in the introduction itself. But here is a much cooler property we can prove using this:

Example 2.22. Prove that:

$$\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n} = 2^n$$

It will be much more instructive if you find the proof yourself. As a hint, this is equivalent to the subset theorem.

Finally, note that we have considered the empty set(the one with zero elements) and the full set(the set with all the elements) to be a subset of the set. Please check if the question is considering the same. If not subtract them from 2^n . A lot of times the empty or full set are not included but the same is not explicitly mentioned, always keep in mind:**Reading the question explains the question.**

Also note that no formula exists for sets with some repeating elements. We'll solve such questions using beggars theorem(aka Stars and Bars), you'll learn more about it later.

Example 2.23. (AMC 10 2008) Two subsets of the set $S = \{a, b, c, d, e\}$ are to be chosen so that their union is S and their intersection contains exactly two elements. In how many ways can this be done, assuming that the order in which the subsets are chosen does not matter?

Solution. Let the subsets be A and B , hence $A \cup B = S$.

We are basically looking to divide S into three sets. The elements which only lie in A , the elements which only lie in B and the elements which lie in $A \cap B$

As 2 elements lie in $A \cap B$, we have $\binom{5}{2}$ ways to determine them.

The other three elements need to be divided to A and B , therefore by using the subset theorem we have 2^3 ways to do so.

Thus, the total ways to do so are $2^3 \cdot \binom{5}{2} = 80$. But, wait, we are not done yet. We need to divide by 2 as we have over counted the case where A and B have just interchanged. So the answer is $\frac{80}{2} = \boxed{40}$ \square

2.8 Probability

Probability is basically the chance of something occurring. While probability and its theories are their own branch, which we will explore later in more detail, with a more focused perspective, The only thing we need to know(which you probably already do) is

Theorem 2.24. *Probability* = $\frac{\text{Number of desired outcomes}}{\text{Total number of outcomes}}$

Let's try to understand that with an example:

Example 2.25. (AIME 2000) A deck of forty cards consists of four 1's, four 2's,..., and four 10's. A matching pair (two cards with the same number) is removed from the deck. Given that these cards are not returned to the deck, let $\frac{m}{n}$ be the probability that two randomly selected cards also form a pair, where m and n are relatively prime positive integers. Find $m + n$

Solution. Without loss of generality, Let the removed pair be of 1.

The first card can either be 1 or not be 1 (obviously). Based on this the probability second card forming a pair is either $\frac{1}{37}$ or $\frac{3}{37}$.

The probability of first card being 1 is $\frac{2}{38}$ and that of it not being 1 is obviously $\frac{36}{38}$

Hence, the probability of getting a pair is $\frac{2}{38} \cdot \frac{1}{37} + \frac{36}{38} \cdot \frac{3}{37} = \frac{2+108}{38 \cdot 37} = \frac{55}{19 \cdot 37} = \frac{55}{703}$

Thus, $m + n = 55 + 703 = 758$ \square

And we shall end this chapter here. This chapter dealt with the common definitions and techniques in counting. We also explored equivalence. We worked on when to multiply and divide in great detail as well as bookmarked things we shall explore more later. Now let's solve some problems.

2.9 Exercise

Solve at least questions worth [52★]. This exercise has a total of [68★].

Exercises

- (AMC 10 2019) [2★] A child builds towers using identically shaped cubes of different colors. How many different towers with a height of 8 cubes can the child build with 2 red cubes, 3 blue cubes, and 4 green cubes? (One cube will be left out.)? **Hints:** [23](#)

- (2) (AMC 10 2006) [2★] A license plate in a certain state consists of 4 digits, not necessarily distinct, and 2 letters, also not necessarily distinct. These six characters may appear in any order, except that the two letters must appear next to each other. How many distinct license plates are possible?

Hints: 66

- (3) (AMC 10 2017) [2★] At a gathering of 30 people, there are 20 people who all know each other and 10 people who know no one. People who know each other hug, and people who do not know each other shake hands. How many handshakes occur within the group?

- (4) (AMC 10 2004) [2★] Henry's Hamburger Haven serves its hamburgers with the following condiments: ketchup, mustard, mayonnaise, tomato, lettuce, pickles, cheese, and onions. A customer can choose one, two, or three meat patties and any collection of condiments. How many different kinds of hamburgers can be ordered?

- (5) (AMC 12 2022) [5★] What is the number of ways the numbers from 1 to 14 can be split into 7 pairs such that for each pair, the greater number is at least 2 times the smaller number? Hints: 20

- (6) (AMC 12 2003) [5★] How many 15-letter arrangements of 5 A's, 5 B's, and 5 C's have no A's in the first 5 letters, no B's in the next 5 letters, and no C's in the last 5 letters? Hints: 62

- (7) (AMC 10 2021) [2★] A deck of cards has only red cards and black cards. The probability of a randomly chosen card being red is $1/3$. When 4 black cards are added to the deck, the probability of choosing red becomes $1/4$. How many cards were in the deck originally?

- (8) (AMC 10 2006) [2★] Bob and Alice each have a bag that contains one ball of each of the colors blue, green, orange, red, and violet. Alice randomly selects one ball from her bag and puts it into Bob's bag. Bob then randomly selects one ball from his bag and puts it into Alice's bag. What is the probability that after this process the contents of the two bags are the same?

- (9) (AMC 10 2020) [5★] Ms. Carr asks her students to read any 5 of the 10 books on a reading list. Harold randomly selects 5 books from this list, and Betty does the same. What is the probability that there are exactly 2 books that they both select? Hints: 17

- (10) (AMC 12 2021) [9★] Two fair dice, each with at least 6 faces are rolled. On each face of each dice is printed a distinct integer from 1 to the number of faces on that die, inclusive. The probability of rolling a sum of 7 is $3/4$ of the probability of rolling a sum of 10, and the probability of rolling a

sum of 12 is $1/12$. What is the least possible number of faces on the two dice combined? **Hints:** [25](#) [50](#)

- (11) (AMC 10 2009) [3★] Two cubical dice each have removable numbers 1 through 6. The twelve numbers on the two dice are removed, put into a bag, then drawn one at a time and randomly reattached to the faces of the cubes, one number to each face. The dice are then rolled and the numbers on the two top faces are added. What is the probability that the sum is 7?
Hints: [14](#)

- (12) (AMC 12 2019) [3★] The numbers 1, 2, ..., 9 are randomly placed into the 9 squares of a 3 by 3 grid. Each square gets one number, and each of the numbers is used once. What is the probability that the sum of the numbers in each row and each column is odd?

Hints: [24](#)

- (13) (AMC 12 2003) [2★] Let S be the set of permutations of the sequence 1, 2, 3, 4, 5 for which the first term is not 1. A permutation is chosen randomly from S . The probability that the second term is 2, in lowest terms, is a/b . What is $a + b$?

- (14) (AMC 10 2018) [2★] A box contains 5 chips, numbered 1, 2, 3, 4, and 5. Chips are drawn randomly one at a time without replacement until the sum of the values drawn exceeds 4. What is the probability that 3 draws are required? **Hints:** [27](#)

- (15) (AMC 10 2021) [3★] Each of the 20 balls is tossed independently and at random into one of the 5 bins. Let p be the probability that some bin ends up with 3 balls, another with 5 balls, and the other three with 4 balls each. Let q be the probability that every bin ends up with 4 balls. What is p/q ? **Hints:** [32](#)

- (16) (ISRO Interview) [3★] A bag contains 2007 red balls and 2007 black balls. We remove two balls at a time repeatedly and

- (i) discard them if they are of the same color,
- (ii) discard the black ball and return to the bag the red ball if they are of different colors.

What is the probability that this process will terminate with one red ball in the bag? **Hints:** [34](#)

- (17) [5★] Two evenly matched teams play in the world series, a best of seven competition in which the competition stops as soon as one team has won four games. Is the world series more likely to end in six or seven games?

- (18) You toss n coins, and you win if you turn up an even number of heads. Otherwise, Bob Hough takes your lunch money.
- (a) [5★] Show that your odds of winning are 50% if all the coins are fair coins.
- (b) [3★] Better yet, show that your odds of winning are 50% if at least one of the coins is fair.
- (19) (AMC 10 2018) [3★] Three young brother-sister pairs from different families need to take a trip in a van. These six children will occupy the second and third rows in the van, each of which has three seats. To avoid disruptions, siblings may not sit right next to each other in the same row, and no child may sit directly in front of his or her sibling. How many seating arrangements are possible for this trip?
- (20) Slips of paper with the numbers from 1 to 99 are placed in a hat. Five numbers are randomly drawn out of the hat one at a time (without replacement). What is the probability that the numbers are chosen in increasing order?

3

Some methods of counting

The last chapter dealt with mostly constructive counting. There we directly using permutations and combinations try to count exactly what we are asked to count. Straightforward, right?.

However, life is not that simple. Sometimes it is easier to count what we are not asked to count and simply subtract(complimentary counting) or divide our problem into smaller cases and add them(casework). Let me illustrate:

3.1 Casework

Example 3.1 (Motivating Example). For how many pairs of consecutive integers lying in 1000, 1001, 1002, . . . 1998, 1999, 2000, does their addition happen without any carry over?

Solution. We need to notice that carry over happens if the sum of two digits at a place are above 10. This means we can easily limit out the numbers which are like 1a85 or 1a89 as there successor has a 9 right below the 8. Keeping our observations in mind, we can see that: 1999 meets the criteria

1a99 meets the criteria if and only if $a = 0, 1, 2, 3, 4$

1ab9 meets the criteria if and only if $a, b = 0, 1, 2, 3, 4$

1abc meets the criteria if and only if $a, b, c = 0, 1, 2, 3, 4$

That will lead to $1 + 5 + 5^2 + 5^3 = 156$ such pairs existing. \square

When we break a question down into multiple cases and then solve them and add it, it is called casework.

This technique is the messiest of the bunch and PnC as branch has developed to try to avoid it. Much of our further chapters will be about developing stronger techniques to avoid casework.

3.2 Complementary Counting

Example 3.2 (Motivating Example). How many three digit numbers exist such that the number contains at least one 0 or 5?

Solution. Instead of looking for the digits which with the condition, it is easier in this case to look for digits which don't. If a three digit number doesn't have 0 or 5, we have 8 choices for every digit. Thus, we have $8^3 = 512$ such digits.

Thus, as there are 900 total three digit numbers and 512 are not suitable so we have $900 - 512 = 388$ \square

This concept is known as complementary counting, counting the thing opposite to asked and then subtracting from the total.

It is the hardest to spot in the wild but also the most versatile. I normally encourage you to at least give it a thought if the phrase 'at least' makes an appearance.

3.3 Principle of Inclusion and Exclusion

Example 3.3 (Motivating Example). (AIME) Many states use a sequence of three letters followed by a sequence of three digits as their standard license-plate pattern. Given that each three-letter three-digit arrangement is equally likely, the probability that such a license plate will contain at least one palindrome (a three-letter arrangement or a three-digit arrangement that reads the same left-to-right as it does right-to-left) is m/n , where m and n are relatively prime positive integers. Find $m + n$.

Solution. A three character palindrome is of the form "♥♦♥". Note that ♥ = ♠ is also valid.

This means that the number of letter palindromes is $26 * 26 = 26^2$ and number of number palindromes is $10 * 10 = 10^2$. So total plates with at least one palindrome is the sum of plates where number is the palindrome and the plates where letter is the palindrome. However, we'll subtract number of plates where both are palindrome as they were counted twice. So we can hence get the total plates with at least one palindrome as $26^2 * 10^3 + 10^2 * 26^3 - 26^2 * 10^2$. As the question asks probability, lets divide this by total number of plates which is $26^3 * 10^3$

$$\begin{aligned}
 & \frac{26^2 * 10^3 + 10^2 * 26^3 - 26^2 * 10^2}{26^3 * 10^3} \\
 &= \frac{26^2 * 10^2(10 + 26 - 1)}{26^2 * 10^2(26 * 10)} \\
 &= \frac{35}{260} = \frac{7}{52}
 \end{aligned}$$

Hence the answer is $7 + 52 = 59$

□

This is called the principal of inclusion and exclusion or PIE for short. This is used in questions where two or more conditions are to be satisfied. Here, we would like introduce a general theorem for the same but will revise two definitions before doing the same:

Definition 3.4. $A \cup B$ refers to the set of all elements which occur in A and B, with duplicates(elements common to both sets) are only written once. This is known as union and read as A union B.

Definition 3.5. $A \cap B$ refers to the set of the common elements which occur in both A and B. This is known as intersection and read as A intersection B.

Now you are ready for the PIE theorem:

Theorem 3.6. $A \cup B = A + B - A \cap B$

Which is what we just intuitively used in the above example.

We also have PIE for three sets which is as follows:

Theorem 3.7.

$$\begin{aligned}
 A \cup B \cup C &= A + B + C \\
 &\quad - A \cap B - B \cap C - C \cap A \\
 &\quad + A \cap B \cap C
 \end{aligned}$$

This is easier to understand using a Venn diagram. If you draw it and label each circle as a set, you'll intuitively get why this is true.

This also extends to union of 4 or more sets. While we can't draw such Venn Diagrams easily, we can prove that the pattern holds true for larger number of sets using induction.

Theorem 3.8. If $(A_i)_{1 \leq i \leq n}$ are finite sets, then:

$$\begin{aligned} \left| \bigcup_{i=1}^n A_i \right| &= \sum_{i=1}^n |A_i| \\ &\quad - \sum_{i < j} |A_i \cap A_j| \\ &\quad + \sum_{i < j < k} |A_i \cap A_j \cap A_k| \\ &\quad - \dots \\ &\quad + (-1)^{n-1} |A_1 \cap \dots \cap A_n| \end{aligned}$$

Proof. (B) We can see this obviously works if we have a single set A .

(S) Let's assume this is true for some n number of sets. We will prove that then it will be true for $n+1$ number of sets as well.

Let's consider an element $k \in A_{n+1}$ which is also part of $A_1 \cup A_2 \cup \dots \cup A_n$. This means we need to prove that we still count it only once.

Without loss of generality, assume that k is a part of $A_1, A_2, A_3 \dots A_l$.

This means the number of times we count k is originally:

$$\binom{l}{1} - \binom{l}{2} + \binom{l}{3} - \dots = 1$$

This will be 1 as we have assumed the statement true for n .

This number of times we will count k when A_{n+1} is present is:

$$\binom{l+1}{1} - \binom{l+1}{2} + \binom{l+1}{3} - \dots$$

Here we make the observation:

$$\begin{aligned} \binom{n}{k} + \binom{n}{k-1} &= \frac{n!}{k!(n-k)!} + \frac{n!}{(k-1)!(n-k+1)!} \\ &= \frac{n!}{(k-1)!(n-k)!} \left(\frac{1}{k} + \frac{1}{n-k+1} \right) \\ &= \frac{n!}{(k-1)!(n-k)!} \left(\frac{n+1}{(k)(n-k+1)} \right) \\ &= \frac{(n+1)!}{k!(n-k+1)!} \\ &= \binom{n+1}{k} \end{aligned}$$

This is known as Pascal's identity. We will discuss it in detail later.

But for this proof, we can use it to break the present sum.

$$\begin{aligned}
 \binom{l+1}{1} - \binom{l+1}{2} + \binom{l+1}{3} - \dots &= \left(\binom{l}{1} + \binom{l}{0} \right) - \left(\binom{l}{2} + \binom{l}{1} \right) \\
 &\quad + \left(\binom{l}{3} + \binom{l}{2} \right) \dots \\
 &= \binom{l}{0} + \left(\binom{l}{1} - \binom{l}{2} + \dots \right) \\
 &\quad - \left(\binom{l}{1} - \binom{l}{2} + \dots \right) \\
 &= 1 + 1 - 1 = 1
 \end{aligned}$$

Thus, by induction, the above is true. \square

Please feel free to forget this proof as more than 3 subsets is rare in math contests. Programming although does often walk into larger cases. But anyways, A rare specimen of more than 3 subsets from ARML is included in the problems.

This brings us at the end of fundamental principles of counting. Here are the arithmetic rules we used:

Theorem 3.9. Multiplication rule: *If two choices are independent, we can multiply them.*

Addition rule: *If we can break a problem into multiple sub problems after a certain choice, we can add the results.*

Subtraction Rule: *If we can count the complement of what we are asked, then we can subtract it from the total.*

Division Rule: *We can divide to account for multiple countings of repeated objects.*

3.4 Exercises

Solve at least questions worth [50★]. This exercise has a total of [71★].

Exercises

- (1) (AMC 12 2014) [2★] A fancy bed and breakfast inn has 5 rooms, each with a distinctive color-coded decor. One day 5 friends arrive to spend the night. There are no other guests that night. The friends can room in any combination they wish, but with no more than 2 friends per room. In how many ways can the innkeeper assign the guests to the rooms? **Hints:** 48.

- (2) (AMC 10 2021) [3★] A farmer's rectangular field is partitioned into 2 by 2 grid of 4 rectangles. In each section the farmer will plant one crop: corn, wheat, soybeans, or potatoes. The farmer does not want to grow corn and wheat in any two sections that share a border, and the farmer does not want to grow soybeans and potatoes in any two sections that share a border. Given these restrictions, in how many ways can the farmer choose crops to plant in each of the four sections of the field?
- (3) (AMC 12 2021) [3★] Azar and Carl play a game of tic-tac-toe. Suppose the players make their moves at random, rather than trying to follow a rational strategy, and that Carl wins the game when he places his third O. How many ways can the board look after the game is over? **Hints:** 58
- (4) (AMC 10 2004) [2★] Coin A is flipped three times and coin B is flipped four times. What is the probability that the number of heads obtained from flipping the two fair coins is the same?
- (5) (AMC 10 2014) [3★] Three fair six-sided dice are rolled. What is the probability that the values shown on two of the dice sum to the value shown on the remaining die?
- (6) (AMC 10 2015) [2★] How many rearrangements of $abcd$ are there in which no two adjacent letters are also adjacent letters in the alphabet? For example, no such rearrangements could include either ab or ba . **Hints:** 44
- (7) [3★] How many three-digit numbers are composed of three distinct digits such the tens digit is the average of the other two? **Hints:** 51 54
- (8) (AMC 10 2020) [9★] There are 10 people standing equally spaced around a circle. Each person knows exactly 3 of the other 9 people: the 2 people standing next to her or him, as well as the person directly across the circle. How many ways are there for the 10 people to split up into 5 pairs so that the members of each pair know each other? **Hints:** 45 42
- (9) (AMC 10 2020) [9★] Jason rolls three fair standard six-sided dice. Then he looks at the rolls and chooses a subset of the dice (possibly empty, possibly all three dice) to reroll. After rerolling, he wins if and only if the sum of the numbers face up on the three dice is exactly 7. Jason always plays to optimize his chances of winning. What is the probability that he chooses to reroll exactly two of the dice? **Hints:** 61
- (10) (AMC 12 2021) [9★] Each of the 12 edges of a cube is labeled 0 or 1. Two labeling are considered different even if one can be obtained from the other by a sequence of one or more rotations and/or reflections. For how many such labeling is the sum of the labels on the edges of each of the 6 faces of the cube equal to 2? **Hints:** 43

- (11) (AMC 10 2018) [2★] How many subsets of $2, 3, 4, 5, 6, 7, 8, 9$ contain at least one prime number?
- (12) [3★] Let (a, b, c, d) be an ordered quadruple of not necessarily distinct integers, each one of them in the set $0, 1, 2, 3$. For how many such quadruples is it true that $a * d - b * c$ is odd?
- (13) (AMC 10) [3★] In how many ways can the sequence $1, 2, 3, 4, 5$ be rearranged so that no three consecutive terms are increasing and no three consecutive terms are decreasing?
- (14) [9★] Each unit square of a 3-by-3 unit-square grid is to be colored either blue or red. For each square, either color is equally likely to be used. The probability of obtaining a grid that does not have a 2-by-2 red square is m/n , where m and n are relatively prime positive. Find $m + n$ **Hints:** [35](#) [60](#)
- (15) (2015 ARML) [9★] Six people of different heights are getting in line to buy donuts. Compute the number of ways they can arrange themselves in line such that no three consecutive people are in increasing order of height, from front to back. [The rare specimen I promised!] **Hints:** [55](#)

4

A guessing game

We stop your usual broadcast to present a guessing game for you.

The reason why most Olympiad givers find PnC hard is as they cannot decide which strategy to use. While the other strategies we are yet to study have a signature framing which you can notice with practice, the most difficult to discern and the most fundamental methods of counting by far are: constructive counting(ch-2), complementary counting(ch-3) and casework(ch-3). In the next few questions you have to first guess what to use. Then solve the question and check if your guess was correct.

I have not provided points or hints for these questions as they are more like a test then an exercise.

Also note that the questions were randomized by a computer, so don't look for patterns. You'll not find them.

Exercises

- (1) (IOQM 2021) Find the number of maps $f : \{1, 2, 3\} \rightarrow \{1, 2, 3, 4, 5\}$ such that $f(i) \leq f(j)$ whenever $i < j$.
- (2) Consider all 3 element subsets of $1, 2, 3 \dots 298, 299, 300$ What number of the subsets have the sum of elements divisible by 3?
- (3) How many three term increasing GPs can be obtained from the GP $1, 2, 2^2, \dots 2^n$
- (4) In how many ways can five cards labeled A, B, C, D, E be rearranged so that no card is more than one away from its original position?(Not moving the cards is also a valid arrangement)

- (5) (ARML 2011) The six sides of a convex hexagon $A_1A_2A_3A_4A_5A_6$ are colored red. Each of the diagonals of the hexagon is colored either red or blue. If N is the number of coloring's such that every triangle $A_iA_jA_k$, where $1 \leq i < j < k \leq 6$, has at least one red side, find the sum of the digits of N .
- (6) (AIME 2010) Let N be the number of ways to write 2010 in the form $2010 = a_3 \cdot 10^3 + a_2 \cdot 10^2 + a_1 \cdot 10 + a_0$, where the a_i 's are integers, and $0 \leq a_i \leq 99$. An example of such a representation is $1 \cdot 10^3 + 3 \cdot 10^2 + 67 \cdot 10^1 + 40 \cdot 10^0$. Find N .
- (7) (Purple Comet 2013) How many four-digit positive integers have exactly one digit equal to 1 and exactly one digit equal to 3?
- (8) (Pascal 2005) A number is called special if each of its digits is less than the digit to the left. For example, 5420 is special. How many special numbers are there between 200 and 700?
- (9) How many three element subsets of 1, 2, 3 . . . 18, 19, 20 have there product divisible by 4?
- (10) A cube is constructed from 4 white unit cubes and 4 blue unit cubes. How many different ways are there to construct the $2 * 2 * 2$ cube using these smaller cubes? (Two constructions are considered the same if one can be rotated to match the other.)
- (11) (IOQM 2020) Three couples sit for a photograph in 2 rows of three people each such that no couple is sitting in the same row next to each other or in the same column one behind the other. How many arrangements are possible?
- (12) Let A denote a subset of $1, 11, 21, \dots, 531, 541, 551$, where any two elements of A do not add up to 552. What is the maximum number of elements in A ?
- (13) (AIME 2005) A game uses a deck of n dierent cards, where n is an integer and $n \geq 6$. The number of possible sets of 6 cards that can be drawn from the deck is 6 times the number of possible sets of 3 cards that can be drawn. Find n .
- (14) (Fermat 2007) What is the number of 3-digit positive integers a such that both a and $2a$ have only even digits?
- (15) Find how many committees(of any number of people more than 1) with a chairman can be chosen from a set of n persons.

- (16) (AIME 2014) Let the set $S = \{P_1, P_2, \dots, P_{12}\}$ consist of the twelve vertices of a regular 12-gon. A subset Q of S is called "communal" if there is a circle such that all points of Q are inside the circle, and all points of S not in Q are outside of the circle. How many communal subsets are there? (Note that the empty set is a communal subset.)
- (17) (Cayley 2009) How many integers n are there with the property that the product of the digits of n is 0, where $5000 \leq n \leq 6000$?
- (18) How many 4 digits exist such that the thousands digit is 1 and the number has exactly two equal digits?
- (19) (Fermat 2010) A gumball machine that randomly dispenses one gumball at a time contains 13 red, 5 blue, 1 white, and 9 green gumballs. What is the least number of gumballs that Wally must buy to guarantee that he receives 3 gumballs of the same colour?
- (20) (2004 AIME) An integer is called snakelike if its decimal representation $a_1a_2a_3\dots a_k$ satisfies $a_i < a_{i+1}$ if i is odd and $a_i > a_{i+1}$ if i is even. How many snakelike integers between 1000 and 9999 have four distinct digits?
- (21) (AIME 2014) An urn contains 4 green balls and 6 blue balls. A second urn contains 16 green balls and N blue balls. A single ball is drawn at random from each urn. The probability that both balls are of the same color is 0.58. Find N .
- (22) (1992 AIME) A positive integer is called ascending if, in its decimal representation, there are at least two digits and each digit is less than any digit to its right. How many ascending positive integers are there?
- (23) (AIME 2018) Find the number of permutations of 1, 2, 3, 4, 5, 6 such that for each k with $1 \leq k \leq 5$ at least one of the first k terms of the permutation is greater than k .
- (24) (IOQM 2021) In how many ways can four married couples sit in a merry-go-round with identical seats such that men and women occupy alternate seats and no husband sits next to his wife?
- (25) Call a positive integer an uphill integer if every digit is strictly greater than the previous digit. For example, 1357, 89, and 5 are all uphill integers, but 32, 1240, and 466 are not. How many uphill integers are divisible by 15?
- (26) For a particular peculiar pair of dice, the probabilities of rolling 1, 2, 3, 4, 5, 6, on each die are in the ratio 1 : 2 : 3 : 4 : 5 : 6. What is the probability of rolling a total of 7 on the two dice?

- (27) (IOQM 2023) Unconventional dice are to be designed such that the six faces are marked with numbers from 1 to 6 with 1 and 2 appearing on opposite faces. Further, each face is colored either red or yellow with opposite faces always of the same color. Two dice are considered to have the same design if one of them can be rotated to obtain a dice that has the same numbers and colors on the corresponding faces as the other one. Find the number of distinct dice that can be designed.
- (28) A restaurant has six appetizers, five main courses, and four deserts to choose from its menu. How many possible dinners are there if a main course is required but appetizers and deserts are not?
- (29) (AIME 2010) Dave arrives at an airport which has twelve gates arranged in a straight line with exactly 100 feet between adjacent gates. His departure gate is assigned at random. After waiting at that gate, Dave is told the departure gate has been changed to a different gate, again at random. Let the probability that Dave walks 400 feet or less to the new gate be a fraction $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.
- (30) (Cayley 2010) What is the number of 3-digit positive integers that have exactly one even digit?

5

Stars, Bars and Hockey Sticks

While most problems can be solved using the basic methods, sometimes it's better to remember a certain fact. This chapter covers some common equivalence as well as some common combinatorial sums. We will start with Stars and Bars or Beggar's Theorem which is one of the most famous equivalences ever.

5.1 Stars and Bars

Example 5.1 (Motivating Example). In how many ways can 10 chocolates be divided among 3 children (The children are distinguishable, the chocolates aren't)?

Solution. This question is basically number of ordered tuples (a, b, c) such that $a + b + c = 10$ given that $a, b, c \geq 0$.

This means we can re frame the question as ways to insert two identical bars among ten identical stars. Note that this is equivalent as the stars will be divided into three parts which will sum to 10. As the problem is same, so is the solution, hence:

We are looking for the permutations of this configuration which will be:
 $\frac{(10+2)!}{2! \cdot 10!}$ Which is equal to 66. □

Note that it is also possible to get the solution with casework in this particular question. However, doing so will become increasingly impractical as the number of children and chocolates will increase.

We can generalize the above idea as:

Theorem 5.2 (Stars and Bars). *we can say the number ways to put n similar objects in k distinguishable bins is equivalent to permuting n stars and $k - 1$ bars which is equal to $\frac{(n+k-1)!}{n!*(k-1)!} = \binom{n+k-1}{n}$*

The Stars and Bars has various uses. The vanilla use of it comes up routinely in collage entrances. It can be made a little more spicy by doing a small change.

Example 5.3. Alice has 24 apples. In how many ways can she share them with Becky and Chris so that each of the three people has at least two apples?

Solution. As everyone needs to have 2 apples, let's start by giving everyone 2 apples to begin with. This means we have to divide 18 apples with 3 people. This is the classical stars and bars. So the answer is $\binom{20}{2}$. \square

This question can be boosted a bit more if every apple limit? For example let's say Alice must have at least 3 apples, Becky at least 2 and Chris at least 1. The solving(and in this case the answer) remains the same.

We can get it a bit more spicy.

Example 5.4. Find the number of positive integer quadruples (a, b, c, d) that satisfy $a + b + c + d < 24$.

Solution. Let $a + b + c + d = 24 - e \iff a + b + c + d + e = 24$ where e is a positive integer. This means a, b, c, d, e are all positive integers. Note that 0 is neither positive nor negative. So all of them are at least 1.

This means we are back to a Stars and Bars problem. Give all of them 1 to start with and then inserting in the theorem we get $\binom{23}{4}$. \square

To lead us to the next part, I'll need you to notice something:

We could use casework on this question. So we want to solve $a+b+c+d = k$ for $k = 3, \dots, 23$. Using Stars and bars, we get the solution to each of the case by giving a, b, c, d 1 each to begin with. So using stars and bars, $\binom{k-4+3}{3} = \binom{k-1}{3}$ for some value of k . This means $\binom{3}{3} + \dots + \binom{22}{3}$ is the answer.

We already know the answer is $\binom{23}{4}$ and as they are equal, we get: $\binom{3}{3} + \dots + \binom{22}{3} = \binom{23}{4}$

5.2 Counting in two ways

Let's talk about a last chapters once. While playing the guessing game, you may have found yourself having two or more ways to solve the same question both, hopefully, leading to the same answer. But as one method is easier to

compute than the other, we should use that. But that doesn't prevent us from thinking about it.

Given a configuration, we should get the same answer from any which methods. This means we can count in two ways.

This idea is what we use to create some combinatorial identities. We calculate the same thing generalized thing in two ways and equate them to get an identity, which we can use elsewhere as it is true in general.

This is called counting in two ways. Let's see it in action:

Example 5.5 (Motivating Example). How many councils with at least 1 member and at most n members be made from a pool of n people?

Solution. We obviously know that the answer is $2^n - 1$ from the subset theorem.

However we can also write this as $\binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n}$

We can, hence say, $\binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} = 2^n - 1$

$$\therefore \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} + 1 = 2^n$$

as we know $\binom{n}{0} = 1$, we can say: $\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} = 2^n$

□

What we just derived is called the Binomial identity.

Theorem 5.6. $\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} = 2^n$

Let's talk about the name of the identity for a while.

Binomial in math refers to polynomial with 2 terms. For example $x + 2$ or $3x + 7$, in general $ax + b$.

If you have already studied some algebras, you may know that $(a+b)^2 = a^2 + 2ab + b^2$ and $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$. These are normally derived by opening the brackets and multiplying(FOIL).

But how do we find $(a+b)^4$ or worse $(a+b)^{10}$.

We can notice that all the terms of the expansion of $(a+b)^k$ are $a^m b^n$ where $m + n = k$. I smell some combinatorics...

Theorem 5.7 (Binomial Theorem).

$$(a+b)^n = \binom{n}{0}a^n b^0 + \binom{n}{1}a^{n-1} b^1 + \dots + \binom{n}{n-1}a^1 b^{n-1} + \binom{n}{n}a^0 b^n = \sum_{m=0}^k \binom{k}{m} \cdot a^m \cdot b^{k-m}$$

Proof. As expected, The Binomial Theorem has a nice combinatorial proof:

We can write $(a+b)^k = \underbrace{(a+b) \cdot (a+b) \cdot (a+b) \cdots \cdot (a+b)}_k$. Repeatedly

using the distributive property, we see that for a term $a^m b^{k-m}$, we must choose m of the k terms to contribute an a to the term, and then each of the other $k - m$ terms of the product must contribute a b . Thus, the coefficient

of $a^m b^{k-m}$ is the number of ways to choose m objects from objects k , or $\binom{k}{m}$. Extending this to all possible values of m from 0 to k , we see that $(a+b)^k = \sum_{m=0}^k \binom{k}{m} \cdot a^m \cdot b^{k-m}$, as claimed. \square

This is also the reason $\binom{n}{k}$ is called a binomial coefficient.

Why is the binomial theorem useful here? We can use the binomial theorem to expand $(1+1)^n = \binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{n-1} + \binom{n}{n} = 2^n$

There are a lot more uses of binomial theorem as well. Unfortunately, we shall not cover them in this book.

Let's look at some more identities.

Example 5.8 (Motivating Example). Suppose a committee consists of m men and n women. In how many ways can a subcommittee of r members be formed?

Solution. We obviously know that the answer is $\binom{m+n}{r}$.

However we can also do some case work. Let's say a committee has 0 men and r women. Then 1 man and $r-1$ women. And so on. We can write this as $\binom{m}{0} * \binom{n}{r-0} + \binom{m}{1} * \binom{n}{r-1} + \dots + \binom{m}{r-1} * \binom{n}{1} + \binom{m}{r} * \binom{n}{0}$

We can, hence say,

$$\binom{m}{0} * \binom{n}{r-0} + \binom{m}{1} * \binom{n}{r-1} + \dots + \binom{m}{r-1} * \binom{n}{1} + \binom{m}{r} * \binom{n}{0} = \binom{m+n}{r}$$

\square

This is called Vandermonde's Identity.

Theorem 5.9. $\binom{m}{0} * \binom{n}{r-0} + \binom{m}{1} * \binom{n}{r-1} + \dots + \binom{m}{r-1} * \binom{n}{1} + \binom{m}{r} * \binom{n}{0} = \binom{m+n}{r}$

5.3 Pascal's Triangle

The triangle somewhere on the top of the page is called Pascal's triangle.

It has a lot of fun and amazing properties (try to find them, you'll be surprised).

We are going to exploit two of them right now. First being, Every term in subsequent line is made by adding the two above it. As it turned out, every ancient civilization did reach this triangle by doing just that. The second one, the reason Blaise Pascal gets to have his name on it. This is exceptionally useful as this leads us to Pascal's Identity....

Theorem 5.10. $\binom{n}{k} + \binom{n}{k+1} = \binom{n+1}{k+1}$

Remember, we saw this while proving principle of inclusion exclusion. There we proved it algebraically. We also 'proved' it in a sort of hand wavy




Figure 5.0. Pascal's Triangle: A wonder of the mathematical world

$$\begin{array}{c}
 \binom{0}{0} \\
 \binom{1}{0} \quad \binom{1}{1} \\
 \binom{2}{0} \quad \binom{2}{1} \quad \binom{2}{2} \\
 \binom{3}{0} \quad \binom{3}{1} \quad \binom{3}{2} \quad \binom{3}{3} \\
 \binom{4}{0} \quad \binom{4}{1} \quad \binom{4}{2} \quad \binom{4}{3} \quad \binom{4}{4} \\
 \binom{5}{0} \quad \binom{5}{1} \quad \binom{5}{2} \quad \binom{5}{3} \quad \binom{5}{4} \quad \binom{5}{5}
 \end{array}$$

Figure 5.0. As it turns out, we can write everything as a binomial coefficient here.

manner above using pascal's triangle. Here is an another proof, combinatorial this time, for it.

Proof. Let's consider choosing a team of k lawyers from a pool of $n - 1$ junior lawyers and 1 Harvey Specter.

The answer is $\binom{n}{k}$. Using casework, We can either have Harvey on the team or not on it. If Harvey is on the team, we have $\binom{n-1}{k-1}$ ways to choose rest of the lawyers. If we don't have Harvey on it, we have $\binom{n-1}{k}$ ways to choose the team.

$$\text{This means } \binom{n-1}{k-1} + \binom{n-1}{k} = \binom{n}{k}. \quad \square$$

Finally, here is the hockey stick identity(try looking at a diagonal in the triangle, that's where the name comes from). We started this section at this identity, and I'll let you prove it.

$$\textbf{Theorem 5.11. } \binom{k}{k} + \binom{k+1}{k} + \cdots + \binom{n}{k} = \binom{n+1}{k+1}$$

Now here is the first thing, you should remember these identities. They can be memorized quite simply by writing them on a piece of paper and taping them to a wall next to where you sleep. Every morning look at it just after you wake up, and every night just before you sleep. You'll have them memorized in less then a week.

The another way to remember(the one which I remember), is by simply solving the questions and deriving every identity you forget, no turning the pages back. That will get it done in less than two hours flat.

5.4 Exercises

Solve at least questions worth [60★]. This exercise has a total of [75★].

Exercises

- (1) [2★] In how many ways can one get 10 upon rolling 7 dice?
- (2) [2★] How many 4 digit numbers have a sum of 9?
- (3) [3★] How many ordered pairs (a, b, c, d) where $a \leq b \leq c \leq d \leq 5$ and $a, b, c, d \in \mathbb{N}$? **Hints:** [13](#)
- (4) (AIME 2000) [9★] Given that

$$\frac{1}{2!17!} + \frac{1}{3!16!} + \frac{1}{4!15!} + \frac{1}{5!14!} + \frac{1}{6!13!} + \frac{1}{7!12!} + \frac{1}{8!11!} + \frac{1}{9!10!} = \frac{N}{1!18!}$$

find the greatest integer that is less than $\frac{N}{100}$. **Hints:** [46](#)

- (5) (AIME 2001) [3★] A fair die is rolled four times. The probability that each of the final three rolls is at least as large as the roll preceding it may be expressed in the form m/n where m and n are relatively prime positive integers. Find $m + n$.
- (6) (AMC 8 2018) [3★] From a regular octagon, a triangle is formed by connecting three randomly chosen vertices of the octagon. What is the probability that at least one of the sides of the triangle is also a side of the octagon? **Hints:** [33](#)
- (7) (AIME 2020) [2★] A club consisting of 11 men and 12 women needs to choose a committee from among its members so that the number of women on the committee is one more than the number of men on the committee. The committee could have as few as 1 member or as many as 23 members. Let N be the number of such committees that can be formed. If $N = \binom{a}{b}$, find $a + b$ **Hints:** [39](#)
- (8) (AIME 2015) [3★] Consider all 1000-element subsets of the set $1, 2, 3, \dots, 2015$. From each such subset choose the least element. The arithmetic mean of all of these least elements is p/q , where p and q are relatively prime positive integers. Find $p + q$. **Hints:** [19](#)
- (9) [2★] For how many positive integers x_1, x_2, \dots, x_{10} do we have $x_1 + x_2 + \dots + x_{10} = 50$?
- (10) (AIME 2011) [5★] Ed has five identical green marbles, and a large supply of identical red marbles. He arranges the green marbles and some of the red ones in a row and finds that the number of marbles whose right hand neighbor is the same color as themselves is equal to the number of marbles whose right hand neighbor is the other color. An example of such an arrangement is *GGRRRGGRG*. Let m be the maximum number of red marbles for which such an arrangement is possible, and let N be the number of ways he can arrange the $m + 5$ marbles to satisfy the requirement. Find the remainder when N is divided by 1000. **Hints:** [16](#) [1](#)
- (11) (AIME 2011) [9★] Six men and some number of women stand in a line in random order. Let p be the probability that a group of at least four men stand together in the line, given that every man stands next to at least one other man. Find the least number of women in the line such that p does not exceed 1 percent. **Hints:** [7](#) [64](#)
- (12) [2★] Let n be a positive integer. In how many ways can one write a sum of at least two positive integers that add up to n ?
- (13) (AIME 2013) [5★] Melinda has three empty boxes and 12 textbooks, three of which are mathematics textbooks. One box will hold any three of her textbooks, one will hold any four of her textbooks, and one will hold

any five of her textbooks. If Melinda packs her textbooks into these boxes in random order, the probability that all three mathematics textbooks end up in the same box can be written as $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$. **Hints:** 57 37

- (14) (AMC 10 2016) [3★] For some particular value of N , when $(a + b + c + d + 1)^N$ is expanded and like terms are combined, the resulting expression contains exactly 1001 terms that include all four variables a, b, c , and d , each to some positive power. What is N ?
- (15) (AMC 12 2021) [9★] A choir director must select a group of singers from among his 6 tenors and 8 basses. The only requirements are that the difference between the number of tenors and basses must be a multiple of 4, and the group must have at least one singer. Let N be the number of different groups that could be selected. What is the remainder when N is divided by 100? **Hints:** 36
- (16) (IMO 1981/2) [9★] Let $1 \leq r \leq n$ and consider all subsets of r elements of the set $\{1, 2, \dots, n\}$. Each of these subsets has a smallest member. Let $F(n, r)$ denote the arithmetic mean of these smallest numbers; prove that

$$F(n, r) = \frac{n+1}{r+1}.$$

Hints: 4

- (17) [5★] Prove that

$$\sum_{k=0}^n k \binom{n}{k}^2 = n \binom{2n-1}{n-1}$$

- (18) Given a positive integer n , what is the largest k such that the numbers $1, 2, \dots, n$ can be put into k boxes such that the sum of the numbers in each box is the same?

6

Geometrical Combinatorics

I will not write an introduction here as my thoughts differ on both the sections.

While geometrical counting is a sticker to make simple questions feel difficult, geometrical probability is a very powerful technique which is used in research as well.

However, I have decided to cover them in the same chapter as they both start with the prefix 'geometric'.

6.1 Geometric Counting

Example 6.1 (Motivating Example). How many rectangles of any and all sizes can be formed in a rectangular grid of size $m * n$?

Solution. This can simply be solved by saying that a rectangle is formed when we choose two vertical and two horizontal lines.

In an $m * n$ grid, we have $m + 1$ verticals and $n + 1$ horizontals.

Hence, we can say the number of rectangles is: $\binom{m+1}{2} * \binom{n+1}{2}$

□

This is all geometric counting is. We have a geometric figure and have to count something about it. It is the cheapest trick in the question writers tool box, if a question seems too simple, stick it on the top of a geometric figure and suddenly half the test takers will not attempt it.

We don't want to be those people.

6.2 Geometric Probability

Geometric probability is a way to calculate probability by measuring the number of outcomes geometrically, in terms of length, area, or volume. Why would we do that? Because sometimes, it is easier to solve for the area than the actual probability. We have a chocolate in India called 'Melody' which has a distinct chocolaty and addictive taste. Its tagline was: "Khud khao, khud jaan jao(Try it yourself to find it for yourself)" The same applies to geometric probability. Let's try it out:

Example 6.2 (Motivating Example). (AMC 10 2017) Chloe chooses a real number uniformly at random from the interval $[0, 2017]$. Independently, Laurent chooses a real number uniformly at random from the interval $[0, 4034]$. What is the probability that Laurent's number is greater than Chloe's number? (Assume they cannot be equal)

Solution. I have not provided a figure to motivate you to draw it.

Let's call Chloe's number as x and Laurent's number as y , all their choices can be represented as a rectangle which is 2017 units on the x axis and 4034 units on the y axis.

We are looking for cases where Laurent's number is greater than Chloe's, or $y > x$

This is a line from $(0, 0)$ to $(2017, 2017)$. We can now find the area of the rectangle above this line divided by the total area of the rectangle which will lead to the answer:

$$\begin{aligned} & \frac{(2017+4034)*2017/2}{2017*4034} \\ &= \frac{2017*1.5}{2017*2} \\ &= \frac{3}{4} \end{aligned}$$
□

Geometric probability can be useful when the number of possible outcomes is infinite and we can easily make a diagram to represent the outcomes.

Another common type of question is:


Example 6.3. A coin of radius r is thrown randomly on a floor tiled with squares of side l . Two players bet that the coin will land on exactly one or more than one square respectively. What relation should l and r satisfy for the game to be fair?

Solution. The coin lies inside the square if the center of the coin is at least r distance away from the boundary of the square.

Thus, thus the bet turns to choosing a random point inside a square. If it is at a distance $d \geq r$ then player one wins or else player two wins.

This means for the game to be fair:

$$\frac{(l-2r)^2}{l^2} = \frac{1}{2}$$



$$\begin{aligned}
 &\iff 2(l - 2r)^2 = l^2 \\
 &\iff 2l^2 - 8lr + 8r^2 = l^2 \\
 &\iff l^2 - 8lr + 8r^2 = 0 \\
 \therefore l &= \frac{8r \pm \sqrt{64r^2 - 32r^2}}{2} \\
 &\iff l = \frac{8r \pm 4r\sqrt{2}}{2} \\
 &\iff l = 4r \pm 2r\sqrt{2}
 \end{aligned}$$


We reject the minus case as then l will be less than r which is not possible.
 $\therefore l = 4r + 2r\sqrt{2}$ □

In conclusion, Geometric probability is the exact converse of geometric counting. There we used to convert a geometrical problem into a counting problem, here we convert a counting problem to a geometry problem. Also it was useless and unnecessary, while this is useful and beautiful.

Solve at least questions worth [45★]. This exercise has a total of [60★].

Exercises

- (1) (AMC 10 2004) [2★] The 5×5 grid shown contains a collection of squares with sizes from 1×1 to 5×5 . How many of these squares contain the black center square?



Hints: [56](#)

- (2) (Extension to 1) [3★] How many rectangles formed by the grid lines in this 5×5 grid contain the black center square?
- (3) (AMC 10 2007) [3★] A set of 25 square blocks is arranged into a 5×5 square. How many different combinations of 3 blocks can be selected from that set so that no two are in the same row or column?
- (4) (AMC 10 2021) [3★] How many ways are there to place 3 indistinguishable red chips, 3 indistinguishable blue chips, and 3 indistinguishable green chips in the squares of a 3 by 3 grid so that no two chips of the same color are directly adjacent to each other, either vertically or horizontally? **Hints:** [40](#)
- (5) (AIME 2006) [5★] There is an unlimited supply of congruent equilateral triangles made of colored paper. Each triangle is a solid color with the same color on both sides of the paper. A large equilateral triangle is constructed from four of these paper triangles. Two large triangles are considered distinguishable if it is not possible to place one on the other, using translations, rotations, and/or reflections, so that their corresponding small triangles are of the same color. Given that there are six different colors of triangles from which to choose, how many distinguishable large equilateral triangles may be formed? **Hints:** [83](#)
- (6) (AMC 10 2019) [5★] Real numbers between 0 and 1, inclusive, are chosen in the following manner. A fair coin is flipped. If it lands heads, then it is flipped again and the chosen number is 0 if the second flip is heads, and 1 if the second flip is tails. On the other hand, if the first coin flip is tails, then the number is chosen uniformly at random from the closed interval $[0, 1]$. Two random numbers x and y are chosen independently in this manner. What is the probability that $|x - y| > \frac{1}{2}$? **Hints:** [9](#)

- (7) (AIME 2004) [3★] A circle of radius 1 is randomly placed in a 15×36 rectangle ABCD so that the circle lies completely within the rectangle. Given that the probability that the circle will not touch diagonal AC is m/n , where m and n are relatively prime positive integers. Find $m + n$.
- (8) (AIME 1989) [2★] 10 points are marked on a circle. How many distinct convex polygons can be made using a subset of them as vertices?
- (9) [3★] There are 10 points on the circumference of a circle. How many points of intersection of line segments with ends at these points will be inside the circle? **Hints:** 6
- (10) [2★] There are 6 points on the x axis and 9 points on the y axis. How many points of intersection of these points lie in the first quadrant?
- (11) [2★] There are 10 lines in a plane, with no three being concurrent and no two being parallel. How many triangles can be formed using these points of intersection?
- (12) [2★] Sheldon is standing at $(0,0,0)$ and can move one unit along either the x, y or z axis. IN how many ways can he walk up to $(3,3,3)$?
- (13) [9★] Choose n points randomly from a circle, What is the probability that all the points are in one semicircle? **Hints:** 70 53
- (14) (Amazon Interview) [9★] A stick is broken in two random places. What are the odds that the three pieces can form the sides of a triangle? **Hints:** 21
- (15) (OMCC 2003) [9★] A square board with side-length of 8 cm is divided into 64 squares with side-length of 1 cm each. Each square can be painted black or white. Find the total number of ways to color the board so that every square with side-length of 2 cm formed with 4 small squares with a common vertex has two black squares and two white squares. **Hints:** 59 15

Part 3

Down the Rabbit Hole

7

Power Up Unlocked!

In this chapter we'll expand on some things we explored in the previous part, namely recursion and incidence matrices. Recursion is the mathematical version of the principle "We can't do the same thing again and again and expect different results".

Incidence matrices is another way to perform double counting to solve much more complicated questions.

This chapter is the start of our PnC journey away from short answer contests towards a more involved, more rigorous PnC.

7.1 Recursion

Example 7.1 (Motivating Example). A tower of n circular discs of different sizes is stacked on one of the 3 given pegs in decreasing size from the bottom. The task is to transfer the entire tower to another peg by sequence of moves under the following conditions:


- (i) each moves carries exactly one disc, and
- (ii) no disc can be placed on top of a smaller one.

It is said that there is temple in Kashi where in its basement are priests playing this game with 64 discs. They started playing at the dawn of time and when they finish the universe will die. Considering it takes them 1 sec to move a disc, how long do we have?

Solution. 64 discs seem quite a lot to work with. Let's look at a Small case. If they had 1 disc, it would take them 1 move.

If they had 2 discs, it would take 3 moves.

If they had 3 discs, it would take 7 moves, as illustrated below:



We can see a pattern here. For some n , let M_n be the moves required. Then $M_{n+1} = 2M_n + 1$. Let's prove it now.

Lets consider it took some M_n moves to move n discs.

So to move $n+1$ discs we first move the top n discs to the second pole in M_n moves. We then move the largest disc to third pole in 1 move. We then finally move the n discs to the third pole in M_n moves.

This means it takes a total of $2M_n + 1$ moves to move $n+1$ discs. This is called the recursive formula as values of every term depends on the previous terms.

We can try to find the explicit formula of M_n for any n by observing that $M_1 = 1 = 2^1 - 1$

Which we can put into the recursive formula to get the form $M_n = 2^n - 1$

For $n = 64$, we will get such a big number that even in seconds that is 42 times the scientific life of universe. \square

While the temple story is great, its fictional. It was created by a friend of Edward Lucas, the mathematician behind this game. There are some more versions of this problem which will make an appearance in the exercises.

7.2 Fibonacci Sequence

Any discussion on recurrence is incomplete without a discussion of the famous Fibonacci Sequence.

1, 1, 2, 3, 5, 8, 13...

is a sequence at the heart of math and nature. We can see it in plants, animals to in the great pyramids to temples of Greece.

What makes this sequence so special?

It is the simplest form of recursion defined by $F_n = F_{n-1} + F_{n-2}$.

But it gets better!

Example 7.2. (IMO/3 1981) Determine all (m, n) where m and n are integers satisfying $(n^2 - mn - m^2)^2 = 1$.

Solution. We need to first see that $(n^2 - mn - m^2)^2 = 1 \implies n^2 - mn - m^2 = \pm 1$

We can see that $(m, n) = (1, 1)$ is a solution, so is $(m, n) = (1, 2)$ and so is $(m, n) = (2, 3)$.

FIBONACCI!

We'll just need to prove this now.

Claim: If and only if (m, n) is a solution than so is $(n, m + n)$.

Let $(n^2 - mn - m^2)^2 = 1$ then:

$$\begin{aligned} & ((m+n)^2 - (m+n)n - n^2)^2 \\ &= (m^2 + 2mn + n^2 - mn - n^2 - n^2)^2 \\ &= (m^2 + mn - n^2)^2 \\ &= (-m^2 + mn - n^2)^2 \\ &= (n^2 - mn + m^2)^2 \\ &= 1 \end{aligned}$$

This proves the if. To prove the only if, we'll argue as follows.

Let's assume, to the contrary, that there is some $(n, m + n)$ which is a solution but (m, n) is not a solution.

This means $(m^2 + mn - n^2)^2 = 1$ and $(n^2 - mn + m^2)^2 = (-m^2 + mn - n^2)^2 = (m^2 + mn - n^2)^2 \neq 1$

This is absurd!

Hence, contradiction. Thus, our initial claim must be false and there exists no such $(n, m + n)$

With this we have proven our claim. □

We'll see Fibonacci occur in other places as well.

For our further purposes, we ask ourselves how do we find the 500th Fibonacci number without needing to find the other 499? Or basically what is the closed form of Fibonacci numbers?

7.3 Solving Liner Recursion

A linear recursion (where we the power/exponent of previous terms is 1) can be solved to get it as a difference of some geometric progressions. However, determining them is a bit more involved.

Definition 7.3 (Characteristic polynomial). Let $x_1, x_2, x_3 \dots$ be a series following the recurrence

$$x_k = c_{k-1}x_{k-1} + c_{k-2}x_{k-2} + c_{k-3}x_{k-3} \dots$$

where c refers to the coefficients of the terms in the recurrence relation. Then, the characteristic polynomial of the recurrence is:

$$x^k = c_{k-1}x^{k-1} + c_{k-2}x^{k-2} + c_{k-3}x^{k-3} \dots$$

When we cancel the common powers, we will be left with the simplified characteristic polynomial.

For example, for the Fibonacci numbers, we have:

$$\begin{aligned} F_n &= F_{n-1} + F_{n-2} \\ \implies x^n &= x^{n-1} + x^{n-2} \\ \implies x^2 &= x^1 + 1 \end{aligned}$$

This is the characteristic polynomial. What is so special about this?

Theorem 7.4 (Expanded form). *If the roots of the characteristic polynomial P are $r_1, r_2, r_3 \dots$ then there exist $a_1, a_2, a_3 \dots$:*

$$x_n = a_1 r_1^n + a_2 r_2^n \dots$$

Here, if given the initial values of x_n we can find $a_1, a_2 \dots$

This theorem is proven by (strong) induction. You can give it a try but I am not discussing it as it is quite long and tedious and unnecessary. What I will discuss is how to use this to find the n^{th} Fibonacci number.

7.4 Binet's Formula

We can replace all F_k with x_k . Thus,

$$x_k = x_{k-1} + x_{k-2},$$

implying

$$x^2 - x - 1 = 0.$$

The solutions to this equation are

$$x_{1,2} = \frac{1 \pm \sqrt{1 + 4 \cdot (-1) \cdot (-1)}}{2} = \frac{-1 \pm \sqrt{5}}{2}.$$

Thus, the formula must be of the form

$$F_n = a_1 \left(\frac{-1 + \sqrt{5}}{2} \right)^n + a_2 \left(\frac{-1 - \sqrt{5}}{2} \right)^n,$$

for constants a_1 and a_2 . To find these two constants, we simply plug in $n = 0$ and $n = 1$ and get the desired result.

If you solve for a_1 and a_2 , you'll reach the result: $a_1 = a_2 = \frac{1}{\sqrt{5}}$.

Putting that in the expression we get:

$$F_n = \frac{\left(\frac{-1 + \sqrt{5}}{2} \right)^n + \left(\frac{-1 - \sqrt{5}}{2} \right)^n}{\sqrt{5}},$$

This is known as the Binet Formula for the n^{th} Fibonacci number. While you'll not get to use this formula a lot, but isn't amazing that despite being filled with so many $\sqrt{5}$ which are irrational, we will always get an positive integer.

Think about it...

7.5 But Wait, what if the roots repeat?

Nice question, let's look at an example from INMO 1996.

Example 7.5. The sequence $\{a_n\}_{n \in \mathbb{N}}$ is defined by $a_1 = 1, a_2 = 2$, and $a_{n+2} = 2a_{n+1} - a_n + 2$ for $n \geq 1$. Prove that for any m , $a_m a_{m+1}$ is also a term of the sequence.

Proof. This example illustrates two concepts. First is what to do if we have a constant in the recurrence.

We are given $a_{n+2} = 2a_{n+1} - a_n + 2$, which means $a_{n+3} = 2a_{n+2} - a_{n+1} + 2$. Let's subtract them.

$$\begin{aligned} a_{n+3} - a_{n+2} &= 2a_{n+2} - 3a_{n+1} + a_n \\ \iff a_{n+3} &= 3a_{n+2} - 3a_{n+1} + a_n \end{aligned}$$

We can solve this recurrence. Let's make a characteristic pronominal,

$$\begin{aligned} x^3 &= 3x^2 - 3x + 1 \\ \iff x^3 - 3x^2 + 3x - 1 &= 0 \\ \iff (x - 1)^3 &= 0 \end{aligned}$$

Thus, we have all three roots equal to 1. In this case, our characteristic polynomial is something like this.

$$a_n = c_1 1^n + (n-1)c_2 1^n + (n-2)c_3 1^n$$

$a_n = c_1 + (n-1)c_2 + (n-2)(n-1)c_3$ We did the process three times as the multiplicity(times the root appears or the power of the factor giving the root) of the root is 3.

How do we figure out the coefficients? Using the values of a we already know.

$$\begin{aligned} a_1 &= c_1 = 1 \\ a_2 &= c_1 + c_2 = 1 + c_2 = 2 \iff c_2 = 1 \end{aligned}$$

But what about c_3 ? We use the final part of the question, the recurrence with the constant.

$$\begin{aligned} a_3 &= c_1 + 2c_2 + 2c_3 = 3 + 2c_3 \text{ As } a_3 = 2a_2 - a_1 + 2 \\ \therefore 3 + 2c_3 &= 4 - 1 + 2 \\ \iff c_3 &= 1 \end{aligned}$$

$$\begin{aligned} \text{Thus, } a_n &= 1 + n - 1 + (n-1)(n-2) = n + n^2 - 3n + 2 \\ &= n^2 - 2n + 2 = (n-1)^2 + 1 \end{aligned}$$

The last form will help us prove the required faster. Now, we'll prove that for any m , $a_m a_{m+1}$ is also a term of the equation.

$$\begin{aligned}
 & a_m a_{m+1} \\
 &= ((m-1)^2 + 1)(m^2 + 1) \\
 &= (m^2 - 2m + 2)(m^2 + 1) \\
 &= m^4 - 2m^3 + 3m^2 - 2m + 2
 \end{aligned}$$

We need this to be equal to $(k-1)^2 + 1$ for some positive integer k .

So are looking for, $(k-1)^2 = m^4 - 2m^3 + 3m^2 - 2m + 1 = (m^2 - m + 1)^2$
 $\implies k = \pm(m^2 - m + 2)$, which is positive for one case of the sign.

□

While the proof itself used a very smart algebraic solving, the main meet was the way to deal with double roots. It's proof also follows from strong induction, and will not be covered for the sake of brevity.

7.6 Catalan Number's

Example 7.6 (Motivating Example). Five boys and five girls want to have ice cream. The ice cream costs 5 rupees. All the boys have a single 10 rupee note. All the girls have a single 5 rupee note. The cashier starts with no change. In how many ways can the boys and girls line themselves so that the cashier ends up giving everyone change?

This questions phrasing makes it very hard to think of what to do first. The question is basically asking us the number of ways to arrange the boys and girls so that the number of girls at any point is more than the number of boys.

Proof. Let's rephrase the question as: The number of lattice paths from $(0, 0)$ to $(5, 5)$ using only unit up and right steps, such that the path stays in the region $x \geq y$. We'll try to solve it for the general case from $(0, 0)$ to (n, n) . These are formally referred to as Dyck paths. Convince yourself that both problems are the same things.

Now let's find the total number of such paths. The total number of lattice paths from $(0, 0)$ to (n, n) without the $x \geq y$ restriction is equal is $\binom{2n}{n}$ using basic PnC.

Using the principle of complementary counting, Let us count the number of paths that go into the $x < y$ region. They shall be called the bad paths. We'll subtract them later.

Suppose that P is a bad path. Since P goes into the region $x < y$, it must hit the line $y = x + 1$ at some point. Let X be the first point on the path P that lies on the line $y = x + 1$

Now, reflect the portion of path P up to X about the line $y = x + 1$, Keeping the latter portion of P the same. This gives us a new path P' . It is trivial to observe that every bad path will have one and only one reflection.




Figure 7.0. One such path.




Figure 7.0. An example

The number of bad paths is therefore equal to the number of lattice paths from $(-1, 1)$ to (n, n) using only unit up and right steps. Notice, that there is no restriction anymore, so we can compute the number of such paths simply as: $\binom{2n}{n+1}$.

So the number of Dyck paths from $(0, 0)$ to (n, n) are:

$$\binom{2n}{n} - \binom{2n}{n+1} = \binom{2n}{n} - \frac{n}{n+1} \binom{2n}{n} = \frac{1}{n+1} \binom{2n}{n}$$

□

The sequence of such numbers is called the Catalan numbers after the French-Belgian mathematician Eugène Charles Catalan. They come up a lot. The sequence goes like: 1, 2, 5, 14, 42, The n^{th} term is (as we derived above):

Theorem 7.7. $C_n = \frac{1}{n+1} \binom{2n}{n}$

7.7 Incidence Matrices

We used the concept of counting in two ways while deriving the identities. In this chapter, we will now learn about incidence matrices and use them to set

up the double counting. While the incidence matrix is a very powerful tool, and can solve a lot of questions, we'll see even stronger methods later.

Example 7.8. In a certain committee, each member belongs to exactly three subcommittees, and each subcommittee has exactly three members. Prove that the number of members equals to the number of subcommittees.

Here's how we usually set up the incidence matrix. In our incidence matrix, each row represents a member, and each column represents a subcommittee. An entry is 1 if the member corresponding to its row belongs to the subcommittee corresponding to its column; otherwise, the entry is 0. Of course, the roles of rows and columns may be interchanged without any loss of generality. If we had 5 members, and 5 subcommittee's, we'd have a matrix looking something like:

$$\begin{pmatrix} 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 \end{pmatrix}$$

Suppose that there are n subcommittees and m members. Then the incidence matrix is a $m \times n$ matrix. The given conditions tell us that each row contains 3 ones, so there are $3m$ ones in total.

On the other hand, each column contains 3 ones, so there are $3n$ ones in total. Equating the two counts, we see that $3m = 3n$, so $m = n$, which is what we wanted to prove.

The methodology of incidence matrix is based on a simple idea, **the sum of elements taken one row at a time is equal to sum of elements taken one column at a time**.

However, this approach is often not enough. Oftentimes, we are given some restriction that applies to every pair of organizations (or individuals). For example, it may be that every two organizations share exactly one common member. In this case, counting the number of 1's as we did above does not incorporate all the given information, and thus would likely be unsuccessful. Fortunately, such problems can usually be approached by counting pairs of 1's. Specifically, we are interested in the number of pairs of 1's that lie on the same column (or row).

Example 7.9. (IMC 2002) Two hundred students participated in a mathematical contest. They had six problems to solve. It is known that each problem was correctly solved by at least 120 participants. Prove that there must be two participants such that every problem was solved by at least one of these two students.

Proof. Let's assume, to the contrary, that no two students managed to together solve all the problems. That means, for any set of two students, one problem must be unsolved by both.

This prompts us to count the pairs of students with their unsolved problem. Let us consider the incidence matrix of this configuration. We have six rows, each representing a problem, and 200 columns, each representing a student. In light of the above remark, we make an entry of the matrix 1 if the student corresponding to the column did not solve the problem corresponding to the row, and make the entry 0 otherwise. The setup is illustrated below:

$$\begin{pmatrix} 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & \dots & 0 & 0 \end{pmatrix}$$

Let T denote the set of pairs of 1 in the same row. We'll be considering the cardinality(fancy way to say number of elements) of the set T .

Counting by column: As there exists a problem unsolved by any two students, we should have at least a pair of ones in one of the rows if we choose any two columns. Basically, $|T| \geq \binom{200}{2}$.

Counting by row: As every problem was solved by at least 120 students, there are at most 80 ones in every row. So we have $\binom{80}{2}$ pairs of one per row. That means, $|T| \leq 6\binom{80}{2}$

We are done as the above implies: $\binom{200}{2} \leq |T| \leq 6\binom{80}{2}$, which is false as $\binom{200}{2} \geq 6\binom{80}{2}$. Hence, our initial assumption was false. Thus, the converse is true.

Hence, proved. □

Exercises

- (1) How many ways are there to tile an 10 by 2 board with 1 by 2 dominoes such that each domino covers exactly two squares and no domino overlaps?
- (2) (BMO 2013) Isaac is planning a nine-day holiday. Every day he will go surfing, or water skiing, or he will rest. On any given day he does just one of these three things. He never does different water-sports on consecutive days. How many schedules are possible for the holiday?
- (3) (AIME 2015) There are $2^{10} = 1024$ possible 10-letter strings in which each letter is either an A or a B . Find the number of such strings that do not have more than 3 adjacent letters that are identical.

- (4) Given a regular $2n$ -gon, how many ways are there to pair vertices and draw line segments between those vertices such that no two line segments intersect?
- (5) (IMO/6 1979) Let A and E be opposite vertices of an octagon. A frog starts at vertex A . From any vertex except E it jumps to one of the two adjacent vertices. When it reaches E it stops. Let a_n be the number of distinct paths of exactly n jumps ending at E . Prove that:


$$a_{2n-1} = 0, \quad a_{2n} = \frac{(2 + \sqrt{2})^{n-1} - (2 - \sqrt{2})^{n-1}}{\sqrt{2}}.$$

- (6) (AIME 2006) A collection of 8 cubes consists of one cube with edge-length k for each integer $k, 1 \leq k \leq 8$. A tower is to be built using all 8 cubes according to the rules:
 - Any cube may be the bottom cube in the tower.
 - The cube immediately on top of a cube with edge-length k must have edge-length at most $k + 2$.
 Let T be the number of different towers than can be constructed. What is the remainder when T is divided by 1000?

- (7) (ISI 2023) Three left brackets and three right brackets have to be arranged in such a way that if the brackets are serially counted from the left, then the number of right brackets counted is always less than or equal to the number of left brackets counted. In how many ways can this be done? ♣
 What about for n left brackets and n right brackets.
- (8) Find the number of triangulation's of a convex $(n+2)$ gon into n triangles by $n-1$ diagonals that do not intersect their interiors.
- (9) (USAMO 1996) An n -term sequence (x_1, x_2, \dots, x_n) in which each term is either 0 or 1 is called a binary sequence of length n . Let a_n be the number of binary sequences of length n containing no three consecutive terms equal to 0, 1, 0 in that order. Let b_n be the number of binary sequences of length n that contain no four consecutive terms equal to 0, 0, 1, 1 or 1, 1, 0, 0 in that order. Prove that $b_{n+1} = 2a_n$ for all positive integers n .
- (10) (IOQM 2023) Given a 2×2 tile and seven dominoes (2×1 tile), find the number of ways of tiling (that is, cover without leaving gaps and without overlapping of any two tiles) a 2×7 rectangle using some of these tiles.
- (11) (China 1993) A group of 10 people went to a bookstore. It is known that:
1. Everyone bought exactly 3 books
 2. For every two persons, there is at least one book that both of them bought.

What is the least number of people that could have bought the book purchased by the greatest number of people?

- (12) (China 1992) Sixteen students took part in a math competition where every problem was a multiple choice question with four choices. After the contest, it is found that any two students had at most one answer in common. Determine the maximum number of questions
- (13) (China 1995) Twenty-one people took a test with 15 true and false questions. It is known that for every two people, there is at least one question that both have answered correctly. Determine the minimum possible number of people that could have correctly answered the question that most number of people are correct on.
- (14) (China 1996) Eight singers participate in an art festival where m songs are performed. Each song is performed by 4 singers, and each pair of singers performs together in the same number of songs. Find the smallest m for which this is possible.
- (15) (AIME 2018) The wheel shown below consists of two circles and five spokes, with a label at each point where a spoke meets a circle. A bug walks along the wheel, starting at point A . At every step of the process, the bug walks from one labeled point to an adjacent labeled point. Along the inner circle the bug only walks in a counterclockwise direction, and along the outer circle the bug only walks in a clockwise direction. For example, the bug could travel along the path $AJABCHCHIJA$, which has 10 steps. Let n be the number of paths with 15 steps that begin and end at point A . Find the remainder when n is divided by 1000.



- (16) How many n -digit numbers whose digits are in the set $\{2, 3, 7, 9\}$ are divisible by 3?

8

Graph Theory

Now in normal language, Graph refers to a paper with a grid of squares. In more formal language we define it as follows:

A graph is a mathematical object we use to think about networks. It consists of a bunch of points, called vertices, and lines joining pairs of vertices, called edges.

While it is not 'explicitly' required in Olympiads, it makes easy work of a lot of problems which would otherwise be a lot harder. Also it's an active area of research so familiarity with it would not hurt.

8.1 Definitions

As I mentioned above, graphs are usually networks. We usually let G denote the graph, V the set of vertices and E the set of edges, and write $G = (V, E)$. We say two vertices are adjacent or are neighbors if they are joined by an edge, and nonadjacent otherwise. We say an edge and a vertex are incident if the vertex is one of the endpoints of the edge. The degree of a vertex v is the number of edges incident with v .

Example 8.1. What is least number of edges a graph on n vertices can have? What is the most number of edges a graph on n vertices can have?

The least number will be 0, as that will just be some points on a plane, with no edges joining them. That's called an independent set or empty graph.




Figure 8.0. Some graphs

The most will be $\binom{n}{2}$ where every node is attached to every other node. This is called a clique graph or a complete graph.

Believe it or not, with so little theory, we are now ready for our very first theorem.

Theorem 8.2. *Let $G = (V, E)$ be a graph. Then:*

$$\sum_{v \in V} \deg v = 2|E|$$

The proof is quite trivial. The LHS counts the degree of every vertices, which will be twice the number of edges as every vertices has 2 edges, which will both add it to the total. This is called the Handshake lemma as we can let the vertices be the number of people in a corporate boardroom and the edges be the handshake between them. If we ask every person how many hands he shook(the degree of his vertices) and add it up, it is quite elementary that we'll get twice the number of handshakes.

Example 8.3. Is there a family of graphs such that every vertex has degree 0? 1? 2? 1 or 2? ♣ Given integers p and q , is there a graph such that every vertex either has degree p or q

Example 8.4. Let a vertex be even if it has even degree, and odd if it has odd degree. Which of the following doesn't exist:

- (1) Graph with only even vertices
- (2) Graph with only odd vertices
- (3) Graph with exactly one even vertex
- (4) Graph with exactly one odd vertex

8.2 Paths and Walks

A walk in a graph G is a sequence of vertices $v_0 - v_1 - v_2 - \cdots - v_n$ such that each vertex v_i is adjacent to the vertex v_{i-1} before it and the vertex v_{i+1} after it. We call L the length of the walk.

A path in a graph G is called a walk if all the vertices are different. Basically, every walk is a path, but every path is not a walk.

This leads us to ask: Given a walk between two vertices in a graph, how do we obtain a path between them? Is there always a walk between two vertices in a graph?

The answer is trivial, if a walk has cycles where we start at v_n and after exploring some more vertices end at v_n , we can just remove them and have a path.

The answer to the second one is simple: no. For example in an independent set, we'll obviously not have a walk between any two points.

This leads us to a few more definitions:

Definition 8.5. A cycle in a graph G is a sequence of vertices $v_0 - v_1 - v_2 - \cdots - v_n = v_0$ such that each vertex v_i is adjacent to the vertex v_{i-1} before it and the vertex v_{i+1} after it. We call L the length of the cycle.

Definition 8.6. A graph is disconnected if it can be divided into two parts with no edges between the parts, otherwise it is connected.

Definition 8.7. A connected component of a graph is a connected subgraph which is as large as possible.

Definition 8.8. Let G be a connected graph. A cut vertex in G is a vertex whose removal disconnects the graph. A cut edge in G is an edge whose removal disconnects the graph




Figure 8.0. e is the cut edge and u, v the cut vertices

And now a theorem

Theorem 8.9. *If a graph has a cut edge, it has a cut vertex.*

However the converse of this is not true. If it has a cut vertex, it may or may not have a cut edge.

Proof. The theorem is trivial for a *graph* with less than 3 vertices.

Let G have more than 3 vertices, and the cut edge $e = uv$. The other vertices will have a path to either u or v without using e but not both as that would violate the assumption that e is the cut edge.

Here if one of u or v is removed, the graph would get disconcerted. If and only if $\deg u$ or $\deg v$ is zero, it will not be a cut vertex(which can happen for only one of them).

The converse of this theorem is however not true. A counterexample will suffice: \square




Figure 8.0. The Counter-example

Example 8.10. Let G be a graph with $n \geq 2$ vertices such that every vertex has degree at least $\frac{n-1}{2}$. Show that G is connected.

8.3 Trees

Example 8.11. What is the smallest number of edges a connected graph of n vertices can have?

If you try to draw connected graphs with the fewest number of edges possible, you'll start to notice that they have a particular structure. They should, if you look closely, look like trees.

Definition 8.12. A graph is acyclic if it contains no cycles. We call an acyclic graph a forest, and a connected acyclic graph a tree. A leaf is a vertex in a tree with degree one.



Figure 8.0. Some trees

All trees are forests, but all forests are not trees.

You would have noticed that the number of edges in a tree of n vertices is $n - 1$. You should also notice that every edge is a cut edge. Another thing that is elementary to note is that adding any new edge will create a cyclic path. A less obvious fact, however, is:

Theorem 8.13. *A forest with n vertices and k components contains $n - k$ edges*

Proof. Forest is called forest as every component will be a tree(as in real life). So we have k trees which in total have n vertices. Let the individual trees have $n_1, n_2, n_3 \dots n_k$ vertices and $n_1 + n_2 + n_3 \dots + n_k = n$, so the number of edges will be $n_1 - 1 + n_2 - 1 + n_3 - 1 \dots n_k - 1 = n - k$

□

Let's now talk about one very last topic, and the very reason why we studied trees.

Definition 8.14. A spanning tree of a graph G is a subgraph of G that is a tree containing all the vertices of G .

The fun fact is:

Theorem 8.15. *A graph is connected if and only if it has a spanning tree.*

Proof. As the statement has if and only if, we'll need to prove the theorem as well as its converse.

Let's first prove that if G has a spanning tree it is connected. Suppose G has a spanning tree. Then G is connected, since for every pair of vertices u, v there is a path from u to v , namely the path in the spanning tree.

Now we'll prove that every connected graph has a spanning tree.

Suppose G is connected. Let e_1, e_2, \dots, e_m be the edges of G in some order. If we go through the edges of G in order and remove edge e_i which is in a cycle then we end up with a graph G' . It is trivial that G' is connected (if we remove e_i then since e_i is in a cycle the remaining graph is still connected). If we break all such cycles, we'll be left with T which will be the spanning tree of graph G .

□

Example 8.16. How many spanning trees does a tree have? How many spanning trees does a cycle have? How many spanning trees does a complete graph have?

8.4 Real life applications

I had promised in the start that graph theory has a variety of real life applications. We'll go through a few here.

One application of paths in graphs is finding driving directions. Let's say I want to go from Cambridge University to Mathematical Bridge. What is the fastest way to get there? We can formulate this problem as a graph theory




Figure 8.0. The google map

problem, where $G = (V, E)$ is a graph whose vertices V are intersection points between roads in Cambridge, and edges are road segments. Then we can think of the university and Math Bridge as two vertices u and v in my graph G , and my questions become the following: How can we find whether there is a path between u and v ? How can we find a shortest path from u to v ?

We can do this in many ways, the most basic being the depth-first search and the breadth-first search algorithm.

8.4.1 Depth-first search. The idea of depth first search is: Start at u and ‘keep walking’, i.e. walk as far as possible searching for v , and if you hit a dead end backtrack.

The algorithm is as follows: Start at u and move to any of its neighbours. Let this be v' . Now from v' move to another neighbour other than u . The algorithm stops on three conditions:

- (1) If it reaches v , it then returns the path as the output.

- (2) If it reaches a v' with no unvisited neighbours. It will retract its path and move till the next fork where it will make a new choice and try again.
- (3) If it reaches $v' = u$ and has no unvisted neighbours, it will declare that we have no path from u to v

Depth-first search allows us to find a path from u to v , or verify that no such path exists. But the path it finds has no guarantee of being the shortest path. In sad cases, it may generate a path from Cambridge to Kings Collage and then to the bridge.

8.4.2 Breadth-first search. The idea od Breadth first search is: Start at u , keep track of the vertices ‘closest’ to u (u ’s neighbors) and see if they’re v . If not, see if any of vertices ‘second-closest’ to u (neighbors of u ’s neighbors) are v . Eventually we’ll find v .

This method is superior to depth search in the aspect that it will find the path with least number of edges. However, it also assumes that there is a path from u to v , which if untrue, it will not detect and go till the system crashes(or it reaches a point where all vertices are neighbour less and terminates and declares no path).

However, they both don’t give us the fastest path as we are taking all edges as equal. But roads are of different lengths and we know that longer roads take more time to travel. According to both of or methods, we’ll end up considering Saudi’s Highway 10(256 km) as just as short as Scotland’s Ebenezer Place(2.46 meters).

To capture this, I will need the concept of weighted edges. A weighted graph is a graph where each edge is assigned a number, w called its weight. For an edge e we let $\omega(e)$ denote its weight.

8.4.3 Dijkstra’s Algorithm. The idea of Dijkstra is Start at u , for each vertex v' keep track of an estimated (weighted) distance from u to v . Visit the vertices in order of how ‘close’ they are to u and see if they’re v . The more nitty gritty of this is left to the diligent reader.

However, despite being quite efficient, google maps doesn’t work on Dijkstra. It instead uses A^* algorithm. It basically makes the estimation step more accurate. Google also uses bi-directional search to make it quicker(looking for path from u to v as well as v to u simultaneously hoping that they meet somewhere in the middle) along with a lot of trade secret pruning, shortcuts and caching. But that is more computer science and less math.

Exercises

- (1) Prove that for any graph with n vertices and m edges, has the lowest $\deg v \leq \frac{2m}{n}$ and the highest $\deg v \geq \frac{2m}{n}$.

- (2) . Is it possible to build a house with exactly eight rooms, each with three doors, and such that exactly three of the house's doors lead outside?
- (3) The complement of a graph G is the graph G' obtained by including an edge if and only if it was not present in the original graph. Prove that between G and G' , one and only one is connected.
- (4) Show that at any party, there are always at least two people with exactly the same number of friends at the party.
- (5) (Martin Gardner) My wife and I recently attended a party at which there were four other married couples. Various handshakes took place. No one shook hands with oneself, nor with one's spouse, and no one shook hands with the same person more than once. After all the handshakes were over, I asked each person, including my wife, how many hands he (or she) had shaken. To my surprise each gave a different answer. How many hands did my wife shake?
- (6) (IMOSL 2001) Define a k -clique to be a set of k people such that every pair of them are acquainted with each other. At a certain party, every pair of 3-cliques has at least one person in common, and there are no 5-cliques. Prove that there are two or fewer people at the party whose departure leaves no 3-clique remaining
- (7) (Italy 2007) Let n be a positive odd integer. There are n computers and exactly one cable joining each pair of computers. You are to color the computers and cables such that no two computers have the same color, no two cables joined to a common computer have the same color, and no computer is assigned the same color as any cable joined to it. Prove that this can be done using n colors.
- (8) USAMO 2007, edited An animal with n cells is a connected figure consisting of n equal-sized cells which are square(a n –tile polyomino). A dinosaur is an animal with at least 2023 cells. It is said to be primitive if its cells cannot be partitioned into two or more dinosaurs. Find the maximum number of cells in a primitive dinosaur
- (9) Show that the Petersen graph has 2000 spanning trees
- (10) (Cayley's Formula) A labelled tree of n vertices is a tree of n vertices where all vertices are given distinct labels from 1 to n . Prove that there are n^{n-2} labelled trees of n vertices.
- (11) (Veblen's Theorem) Prove that the edges of a graph can be partitioned into cycles if and only if each vertex has even degree.




Figure 8.0. Petersen Graph

- (12) (Euler's Formula) Prove that given graph G be a connected planar graph with V vertices, E edges and F faces (number of parts it divides the plane into, note: the outside of graph is also a face). Then $F + V - E = 2$
- (13) (IrMO 1989) . Each of the n members of a club is given a different item of information. They are allowed to share the information, but, for security reasons, only in the following way: A pair may communicate by telephone. During a telephone call only one member may speak. The member who speaks may tell the other member all the information he or she knows. Determine the minimal number of phone calls that are required to convey all the information to each other.
- (14) (IrMO 1994) If a square is partitioned into n convex polygons, determine the maximum number of edges present in the resulting figure.
- (15) (IMO 2019, edited) A social network has 2023 users, some pairs of which are friends (friendship is symmetric). If A, B, C are three users such that AB are friends and AC are friends but BC is not, then the administrator may perform the following operation: change the friendships such that BC are friends, but AB and AC are no longer friends. Initially, 1011 users have 1012 friends and 1012 users have 1011 friends. Prove that the administrator can make a sequence of operations such that all users have at most 1 friend.
- (16) (Moser's Circle Problem) Given n points on the circumference of pizza, what is the maximum number of parts the circle is separated by the chords connecting all the n points to each other? (Note: Don't use Engineer's induction)

9

A New Breed of Counting

We are going to encounter a new breed of Combinatorics now onward where we are not so interested in counting in the ways to do something but in proving that there exists some way of doing something or more often, there exists no way of doing something. We'll now look at tiling's and coloring and expected values and betting and games and a lot, lot more. While this section begins our foray into the more robust real life uses of Combi, it is still relevant to Olympiads. A beginning example could be:

Example 9.1. (RMO 2023) Consider a set of 16 points arranged in 4×4 square grid formation. Prove that if any 7 of these points are coloured blue, then there exists an isosceles right-angled triangle whose vertices are all blue.

Proof. Probably the hardest problem on the test.


Fun story, I actually gave the test when this question came. Overconfident in my combi, I messed it up pretty bad as I created a non-existent counter example. I'll leave the details to your imagination.

The solution proceeds by observing the possible isosceles triangles.

We need to see that either three points lie on the vertices of a square or 2 on the vertices of the surrounding three squares and 1 on the center.

We therefore have at most 2 points on the outside squares, which is $2 * 3 = 6$. But as 7 points are colored, either a square has more than 2 vertices colored or the center square has a colored vertex.

Thus, the stated claim is true. □



H

Let's get started with the most complex simple thing!

9.1 Pigeon Hole Principle

This PHP while very powerful, seems stupid when heard/read for the first time:

Theorem 9.2 (Pigeon hole principle). *Consider a flock of pigeons nestled in a set of n pigeonholes. If there are n pigeons, then it is possible for all of the pigeons to rest happily in separate pigeonholes. However, if at least one more pigeon arrives, making a total of more than n pigeons, then at least one of the pigeonholes, inevitably, will end up with more than one pigeon.*

In particular, if k pigeons are put into n holes, there are at least $\lceil \frac{k}{n} \rceil$ in one of the holes.

While PHP may seem pointless, even offensively funny to some, a correct choice of pigeons and holes can simplify quite a lot of problems. It was first used by Dirichlet and is therefore also called the Dirichlet principle. Here is an IMO problem to show PHP's power:

Example 9.3. (IMO/1 1972) Prove that from a set of ten distinct two-digit numbers (in the decimal system), it is possible to select two disjoint subsets whose members have the same sum?

Proof. Let the number of subsets be the pigeons and the possible sums be the holes.

We have $2^{10} - 2 = 1022$ subsets and $90 + 91 + 92 + \dots + 99 - 10 + 1$ possible sums. As $100 * 10 = 1000 \leq 1022$, the pigeons are far less than the holes. Hence, two are sure to share the hole and hence, we will always have two disjoint subsets whose members have the same sum. \square

You have already used PHP before in a much simpler form to answer riddles like: "You have 14 brown socks, 14 blue socks and 14 black socks in

your sock drawer. How many socks must you remove (without looking to be sure) to have a matched pair?"

We didn't give it a name then as the uses were obvious. As they become more complex, a name was given to the idea.

PHP, while simple, can be extended to real analysis and other branches of mathematics in the following form:

Theorem 9.4 (Infinite PHP). *Given an infinite pigeons, if they are put into finite pigeon holes, there is at least one pigeon hole with infinite pigeons.*

We can use this to prove something of the form:

Example 9.5. A 100×100 board is divided into unit squares. In every square there is an arrow that points up, down, left or right. The board square is surrounded by a wall, except for the right side of the top right corner square. An insect is placed in one of the squares. Each second, the insect moves one unit in the direction of the arrow in its square. When the insect moves, the arrow of the square it was in moves 90 degrees clockwise. If the indicated movement cannot be done, the insect does not move that second, but the arrow in its squares does move. Is it possible that the insect never leaves the board?

Proof. Let's assume to the contrary that there is some arrangement such that the insect is trapped.

In this case, it will make infinite moves. As there are only 100^2 pigeon holes, the insect visits some square infinite time.

As the arrows keep changing, it also visits the neighbouring squares infinite times.

By the same logic, it visits all the squares infinite times. Hence, it visits the top right square infinite times, which would mean it is not trapped.

This is a contradiction, hence our initial assumption is false. There exists no such arrangement such that the insect is trapped.

Hence, proved. □

Before we move ahead, here is a classic PHP question for you to try which we can also do using Extremal principle(later in this chapter).

Example 9.6. There are 17 points inside an equilateral triangle with side lengths 1. Prove there are at least two points within distance $\frac{1}{4}$ of each other.
Hint: Try dividing the triangle in equal parts.

9.2 Expected Value and Probabilistic Method

The probabilistic method refers to proving the existence of some element in a set by showing the probability of its existence is non-zero.

Cutting out the jagron, let's say I have bag of candy. How do you prove that I have a Kit-Kat in the bag? You either keep on removing chocolates till you get a Kit-Kat, or through some tricks you prove that the probability of drawing a Kit-Kat is not zero.

Another allegory would be thinking of it like rolling a dice. If you roll a regular six-sided dice, there's a chance of getting any number from 1 to 6. The probabilistic method is a bit like saying, "Hey, there's a chance, maybe a small one, but it's not zero, that I'll get a 6." So, you can be pretty confident that a 6 exists on that dice, even if you haven't seen it yet.

This was explained thrice to provide the motivation to define the following.

Definition 9.7 (Expected Value). For a random variable X , $\mathbb{E}(x) = \sum x_i \cdot P(x_i)$ where x_i are the possible values of X and $P(x_i)$ is the probability of the value being x_i

For example for a dice, let X be the number on the die. Then $\mathbb{E}(x) = \frac{1}{6}1 + \frac{1}{6}2 + \dots + \frac{1}{6}6 = 3.5$.

What this tells us is that if we roll a dice, we expect the value to be 3.5. While 3.5 is not a number on the dice, over a lot of throws this is the average of roll of a dice.

We now define the most important theorem of the probabilistic method

Theorem 9.8 (Linearity of Expectations). $\mathbb{E}(X_1 + X_2 + X_3 + \dots + X_n) = \mathbb{E}(x_1) + \mathbb{E}(x_2) + \dots + \mathbb{E}(x_n)$

This is obvious when the variables are independent. However, the beauty of this theorem is based on the fact that that even if they are not independent, this still stands. The proof has been omitted as it can be made using a incidence matrices and then summing up. You will end up with a ugly summation which will mean the same as above. Google it if you are curious! Let's see a classic example

Example 9.9 (HMMT 2006). At a nursery, 2006 babies sit in a circle. Suddenly, each baby randomly pokes either the baby to its left or to its right. What is the expected value of the number of unpoked babies?

Solution. Let me introduce you to the most classy way of using the probabilistic method. We define a variable P_n for the n^{th} baby in the circle where

$P = 0$ if the baby is poked and $P = 1$ if it is unpoked.

The probability of a baby being unpoked is $\frac{1}{4}$. Which means expected value of P_n is $\frac{1}{4}$ which means using Linearity of Expectations(LOE) we can say that $\mathbb{E}(\text{Number of unpoked babies}) = \mathbb{E}(P_1 + P_2 + \dots + P_{2006}) = \mathbb{E}(P_1) + \dots + \mathbb{E}(P_{2006}) = \frac{2006}{4}$

□

We can also have questions which while can be done using the probabilistic method, also have simpler solutions.

Example 9.10. (AMC 10 2021) Five balls are arranged around a circle. Chris chooses two adjacent balls at random and interchanges them. Then Silva does the same, with her choice of adjacent balls to interchange being independent of Chris's. What is the expected number of balls that occupy their original positions after these two successive transpositions?

Solution. We can also do this by the definition of expected value. Without loss of generality, let's say Chris interchanged the first ball with the second ball.

Silva can now interchange either the same two ($\frac{1}{5}$ probability) which leaves us with 5 balls in the original position.

Silva can interchange one of the balls Chris switched with another neighbour ($\frac{2}{5}$ probability) which leaves 2 balls in the original position.

Silva can interchange two entirely new balls ($\frac{2}{5}$ probability) which leaves 1 ball in the original position.

This means the expected value is $\frac{1}{5} * 5 + \frac{2}{5} * 2 + \frac{2}{5} * 1 = 1 + 0.8 + 0.4 = 2.2$
If we want to use probabilistic method, we can define a variable P for every ball which is 0 if the ball is not on its original position and 1 if it is.

So the Expected value of P is the sum of the probability of the ball never being switched and it being switched twice.

$$\therefore \mathbb{E}(P) = \left(\frac{3}{5}\right)^2 + \frac{2}{5} \cdot \frac{1}{5} = \frac{11}{25}$$

As we have 5 balls, Using LOE, the expected number of balls in the original position are $\frac{11}{25} * 5 = \frac{11}{5} = 2.2$ □

To the contrary of the last example, here is one which cannot be solved neatly without LOE.

Example 9.11. (Putnam 2014, A4) Suppose X is a random variable that takes on only non-negative integer values, with $\mathbb{E}(X) = 1$, $\mathbb{E}(X^2) = 2$, and $\mathbb{E}(X^3) = 5$. Determine the smallest possible value of the probability of the event $X = 0$.

Proof. We need to realize that given expected value of x, x^2 and x^3 , we can find the expected value of any polynomial with degree 3 or less courtesy

linearity of expectations.

This means we know the expected value of $f(x) = (x - 1)(x - 2)(x - 3) = x^3 - 6x^2 + 11x - 6$ which is 0 for 1, 2, 3.

$$\mathbb{E}(f(x)) = 5 - 6 * 2 + 11 * 1 - 6 = 5 - 12 + 11 - 6 = -2$$

We need to now notice that $f(0) = -6$ and $f(x) > 0$ for all $x > 3$. So as the probability of $x > 3$ increases, probability of $x = 0$ also increases.

For the minimum case, let probability of $x = 0$ be p and probability of $x = 1, 2, 3$ be $1 - p$.

$$\text{This means } -6p = -2 \iff p = \frac{1}{3}.$$

We can construct probability for 1, 2, 3 to satisfy the equations. That is left for you to try yourself. \square

Before moving onto some other applications of this method, I'll give a brief view of how it ends up used in active research. Also I'll introduce a powerful graph theory theorem at the same time, and show off another way of setting the variable. One stone, three pigeons

Theorem 9.12 (Turán's Theorem). *For $G = (V, E)$. Let an independent set be a set of vertices such that no two are adjacent. G contains an independent set of size at least $\sum_{v \in V} \frac{1}{d(v)+1}$*

Proof. We'll permute all the vertices in a line. Let's define a variable S as the number of vertices such that a vertex in the permutation lies before all vertices adjacent to it. The expected value of this is obviously the expected size of an independent set.

Let another variable P be 1 if all neighbours of a vertex are after it in the permutation, and 0 otherwise. Using LOE, we can say that $\mathbb{E}(S) = \sum_{v \in V} \mathbb{E}(P_v)$.

We need to note that for any vertex v , it had $d(v)$ adjacent vertices. Only one of them, among the vertex and its neighbours, has all its neighbours after it. This means that the probability of P_v being 1 is $\frac{1}{d(v)+1}$ which means

$$\mathbb{E}(P_v) = \frac{1}{d(v)+1}$$

$$\text{Thus, expected size of independent set } \mathbb{E}(S) = \sum_{v \in V} \frac{1}{d(v)+1} \quad \square$$

We will also prove the same using Extremal principal later.

Exercises

- (1) (AMC 10 2006) A player pays 5 dollars to play a game. A die is rolled. If the number on the die is odd, the game is lost. If the number on the die is even, the die is rolled again. In this case the player wins if the second number matches the first and loses otherwise. How much should the player win if the game is fair? (In a fair game the probability of winning times the amount won is what the player should pay.)

- (2) (AMC 12) A school has 100 students and 5 teachers. In the first period, each student is taking one class, and each teacher is teaching one class. The enrollments in the classes are 50, 20, 20, 5, and 5. Let t be the average value obtained if a teacher is picked at random and the number of students in their class is noted. Let s be the average value obtained if a student was picked at random and the number of students in their class, including the student, is noted. What is $t - s$?
- (3) (OMM 1998) The sides and diagonals of a regular octagon are colored black or red. Show that there are at least 7 monochromatic triangles with vertices in the vertices of the octagon.
- (4) Let F_n be the n^{th} Fibonacci numbers. Show that for some $n \geq 1$, F_n ends with 2023 zeros.
- (5) Let S be a subset of $1, 2, 3, \dots, 2n$ with $n + 1$ elements. Show that there are two elements in S which are relatively prime.
- (6) (continuing of the above question) Show that there are two elements in S , one divisible by the other
- (7) At a party, certain pairs of individuals have shaken hands. Prove that there exist two persons who have shaken the same number of hands
- (8) Given 7 lines on the plane, prove that two of them form an angle less than 26°
- (9) A chess grandmaster has 77 days to prepare for a tournament. He wants to play at least one game per day, but not more than 132 games in total. Prove that there is a sequence of successive days on which he plays exactly 21 games in total.
- (10) (Canada 2004, edited) Let T be the set of all positive integer divisors of 2023^{100} . What is the largest possible number of elements that a subset S of T can have if no element of S is an integer multiple of any other element of S ?
- (11) Show that there is some n for which 111...111 (with n ones) is divisible by 2023
- (12) (IMO/4 1964) Seventeen people correspond by mail with one another - each one with all the rest. In their letters only three different topics are discussed. Each pair of correspondents deals with only one of these topics. Prove that there are at least three people who write to each other about the same topic.
- (13) Prove that the decimal representation of any irrational number has at least two digits appearing infinitely often

- (14) (IMO/4 1985) Given a set M of 1985 distinct positive integers, none of which has a prime divisor greater than 23, prove that M contains a subset of 4 elements whose product is the 4th power of an integer.
- (15) Prove that among any $2m+1$ distinct integers of absolute value less than or equal to $2m - 1$, there are three whose sum is zero.(Although can be solved using PHP, there is a much simpler way out)
- (16) (Putnam/A2 2002) Given any five points on a sphere, show that some four of them must lie on a closed hemisphere.
- (17) Given nine points inside the unit square, prove that some three of them form a triangle whose area does not exceed $\frac{1}{8}$.

Exercises

- (1) (IMO/4 2000) A magician has one hundred cards numbered 1 to 100. He puts them into three boxes, a red one, a white one and a blue one, so that each box contains at least one card. A member of the audience selects two of the three boxes, chooses one card from each and announces the sum of the numbers on the chosen cards. Given this sum, the magician identifies the box from which no card has been chosen. How many ways are there to put all the cards into the boxes so that this trick always works?
- (2) (Japan 2009) Let N be a positive integer. Suppose that a collection of integers was written in a blackboard so that the following properties hold:
- each written number k satisfies $1 \leq k \leq N$;
 - each k with $1 \leq k \leq N$ was written at least once;
 - the sum of all written numbers is even.
- Show that it is possible to label some numbers with ♡ and the rest with ♠ so that the sum of all numbers with ♡ is the same as the sum of all numbers with ♠
- (3) (Ukraine 1999) Two players alternately write integers on a blackboard as follows: the first player writes a_1 arbitrarily, then the second player writes a_2 arbitrarily, and thereafter a player writes a number that is equal to the sum of the two preceding numbers. The player after whose move the obtained sequence contains terms such that $a_i - a_j$ and $a_{i+1} - a_{j+1}$ ($i \neq j$) are divisible by 1999, wins the game. Which of the players has a winning strategy?

Part 4

Algebra

10

Algebraic Manipulations

You are already expected to know basic algebra, I think it will be embarrassing for both of us if we need to discuss $x - 3 = 5 \iff x = 8$

Hence, I hope that you not will wonder what x, y or z has to do with math. We'll study about manipulation in this chapter. Manipulation normally is a bad thing, its the act of saying or doing things to control or influence (a person or situation) cleverly or unscrupulously, for your gain. We should not be manipulative and be away from such people.

However, when we refer to manipulation in math, we refer to playing around with an algebraic equation to make it more favorable or easy to solve. We'll study some methods of manipulation in this chapter.

10.1 Binomial Theorem

If you have already studied algebraic identities, you may know that $(a+b)^2 = a^2 + 2ab + b^2$ and $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$. These are normally derived by opening the brackets and multiplying(FOIL). But how do we find $(a+b)^4$ or worse $(a+b)^{10}$.

We can notice that all the terms of the expansion of $(a+b)^k$ are $a^m b^n$ where $m + n = k$. Does this give you a feel of PnC?

Theorem 10.1 (Binomial Theorem). $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$

Proof. As expected, The Binomial Theorem has a nice combinatorial proof: We can write $(a+b)^k = \underbrace{(a+b) \cdot (a+b) \cdot (a+b) \cdots \cdot (a+b)}_k$. Repeatedly

using the distributive property, we see that for a term $a^m b^{k-m}$, we must choose m of the k terms to contribute an a to the term, and then each of the other $k - m$ terms of the product must contribute a b . Thus, the coefficient of $a^m b^{k-m}$ is the number of ways to choose m objects from objects k , or $\binom{k}{m}$. Extending this to all possible values of m from 0 to k , we see that $(a+b)^k = \sum_{m=0}^k \binom{k}{m} \cdot a^m \cdot b^{k-m}$, as claimed. \square

We also need to note that:

Theorem 10.2 (Binomial Approximation). $(1+x)^n = 1 + nx$ for $x \ll 1$

$$\text{Proof. } (1+x)^n \\ = 1 + \binom{n}{1}x + \binom{n}{2}x^2 + \dots$$

As $x \ll 1$, then x^2 will be insignificant in comparison to 1.

$$(1+x)^n = 1 + nx \quad \square$$

Let's try a problem:

Example 10.3. (JEE Adv 2023) Let a and b be two nonzero real numbers. If the coefficient of x^5 in the expansion of $(ax^2 + \frac{70}{27bx})^4$ and the coefficient of x^{-5} in $(ax - \frac{1}{bx^2})^7$ are equal, then what is the value of $2b$?

Solution. Coefficient of x^5 in $(ax^2 + \frac{70}{27bx})^4$ will be $\binom{4}{k} a^k (\frac{70}{27b})^{4-k}$. We also know $(x^2)^k \cdot \frac{1}{x^k} = x^5$.

Hence, we can figure $k = 3$. Thus, the coefficient is $\binom{4}{3} a^3 (\frac{70}{27b})$.

Lets now do the same for the coefficient of x^{-5} in $(ax - \frac{1}{bx^2})^7$. The coefficient is: $\binom{7}{L} a^L (\frac{-1}{b})^{7-L}$ and we know that $x^L - (\frac{1}{x^2})^L = x^{-5}$.

It's trivial that $L = 3$. Thus, the coefficient is $\binom{7}{3} a^3 (\frac{-1}{b})^{7-3}$.

Equating them: $\binom{7}{3} a^3 (\frac{-1}{b})^{7-3} = \binom{4}{3} a^3 (\frac{70}{27b})$

$$\iff \binom{7}{3} (\frac{1}{b})^4 = \binom{4}{3} (\frac{70}{27b})$$

$$\iff \frac{7!}{4! \cdot 3!} = \frac{4!}{3!} \left(\frac{70}{27} \right)$$

$$\iff b^3 = \frac{7 \cdot 6 \cdot 5 \cdot 4!}{4! \cdot 3!} \cdot \frac{27}{70}$$

$$\iff b^3 = \frac{27}{8}$$

$$\iff b = \frac{3}{2}$$

$$\iff 2b = 3 \quad \square$$

Here is an example for you to try.

Example 10.4. (JEE adv 2013) The coefficient of three consecutive terms of $(1+x)^{n+5}$ are in the ratio 5 : 10 : 14 then what is the value of n ?

10.2 Common Expansions and Factorization

Below are some expansions/factorization which occur quite often. While you can easily prove them using binomial theorem or by expansion. I think you may already know most of them! $(x + y)^2 = x^2 + 2xy + y^2$

$$(x - y)^2 = x^2 - 2xy + y^2$$

$$(x + y)^2 = (x - y)^2 + 4xy$$

$$x^2 - y^2 = (x - y)(x + y)$$

$$(x + y + z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$$

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$ This knowledge makes us quite powerful. Let's use that power now.

Example 10.5. Simplify $a^2 + b^2 + c^2 - ab - bc - ca$

Solution. $a^2 + b^2 + c^2 - ab - bc - ca$

$$= \frac{1}{2}(2a^2 + 2b^2 + 2c^2 - 2ab - 2bc - 2ca)$$

$$= \frac{1}{2}(a^2 - 2ab + b^2 + b^2 - 2bc + c^2 + c^2 - 2ca + a^2)$$

$$= \frac{1}{2}((a - b)^2 + (b - c)^2 + (c - a)^2) \quad \square$$

Now that we have unlocked the power of algebraic expansion, we can use them to clean some dirty calculations:

Example 10.6. Calculate $\frac{(2020^2 - 20100)(20100^2 - 100^2)(2000^2 + 200100)}{2010^6 - 10^6}$

Solution. KEEP YOUR CALCULATOR DOWN. With our new found powers, the question will shrivel before your eyes.

$$\frac{(2020^2 - 20100)(20100^2 - 100^2)(2000^2 + 200100)}{2010^6 - 10^6}$$

$$= \frac{10^2(202^2 - 201)10^4(201^2 - 1)10^2(200^2 + 201)}{10^6(201^6 - 1)}$$

$$= \frac{10^8(202^2 - 201)(201^2 - 1)(200^2 + 201)}{10^6(201^6 - 1)}$$

Notice that the question has most of the terms somewhat close to 201. Let

$$a = 201 \therefore \frac{10^8(202^2 - 201)(201^2 - 1)(200^2 + 201)}{10^6(201^6 - 1)}$$

$$= \frac{10^2((a+1)^2 - a)(a^2 - 1)((a-1)^2 + a)}{a^6 - 1}$$

$$= \frac{10^2(a^2 + 1 + a)(a^2 - 1)(a^2 + 1 - a)}{a^6 - 1}$$

$$= \frac{10^2(a^2 + 1)^2 - a^2(a^2 - 1)}{a^6 - 1}$$

$$= \frac{10^2(a^4 + a^2 + 1)(a^2 - 1)}{a^6 - 1}$$

$$= \frac{10^2(a^6 - 1)}{a^6 - 1}$$

$$= 10^2 = 100 \quad \square$$

Now you try:

Example 10.7. Calculate $\sqrt{(500)(501)(502)(503) + 1}$

10.3 More Factorization Tricks

While we can solve linear equations with ease, and we'll learn how to solve a quadratic (You may already know that); solving higher power equations is not that nice. Hence, we try to break them into smaller, more nicer equations. Some methods of the same are shown here. Up first: A generalization of an identity.

Theorem 10.8. $x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \dots + xy^{n-2} + y^{n-1})$

NOTE: This happens for all natural values of n and the sign in the second bracket is all positive.

Theorem 10.9. $x^{2n+1} + y^{2n+1} = (x + y)(x^{2n} - x^{2n-1}y + \dots - xy^{2n-1} + y^{2n})$

NOTE: This only happens for odd powers, and the sign in second bracket alternates.

We can prove the first simply by expanding. The second simply follows by $y \rightarrow -y$.

$$\begin{aligned} \text{Proof. } & (x - y)(x^{n-1} + x^{n-2}y + \dots + xy^{n-2} + y^{n-1}) \\ &= x^n + \cancel{x^{n-1}y} + \dots + \cancel{x^2y^{n-2}} + \cancel{xy^{n-1}} - \cancel{x^{n-1}y} - \cancel{x^{n-2}y^2} - \dots - \cancel{xy^{n-1}} - y^n \\ &= x^n - y^n \quad \square \end{aligned}$$

We will now talk about my favorite factorization trick, or more accurately, Simon's Favorite Factoring Trick (SFFT).

The Simon here refers to Simon Rubinstein-Salzedo, the director of Euler's Circle, a very prestigious math program. The legend goes that he was one of the earliest people posting on AOPS. Richard Rusczyk hired him as a teaching assistant after he graduated high school. During this time, he solved a question using the method we are gonna talk about. He couldn't remember its name, and was going to say "Using my favorite factoring trick." but instead said, "Using Simon's favorite factoring trick."

Richard liked the name so much that he referred to it as Simon's Favorite Factoring Trick in the critically acclaimed Art of Problem Solving books. And the name stuck.

Here is what he says about this:

It is very strange to me that my greatest claim to fame in life is a single forum post I made when I was 18 years old. I think I've done much better things in my life, especially running Euler Circle and trying to revolutionize gifted mathematics education, giving strong students an opportunity to get a dignified and challenging education that no one else is willing to offer them. However, it does seem that people find my factoring trick memorable with my name on it, so it seems to have done a small part in improving students' problem-solving abilities; I'm glad about that. The really weird part, though, is that my factoring trick has made me something of a celebrity among math contest kids: they sometimes ask me for my autograph when I run math circles or show up at competitions. I certainly don't deserve that

So that's the story. Now let's come to the actual trick.

Theorem 10.10 (SFFT). $xy + kx + ly = C \Rightarrow (x + l)(y + k) = C + kl$

Line of Thought. $xy + kx + ly = C$

$$\begin{aligned} &\iff x(y + k) + ly = C \\ &\iff x(y + k) + ly + kl = C + kl \\ &\iff x(y + k) + l(y + k) = C + kl \\ &\iff (x + l)(y + k) = C + kl \end{aligned}$$

□

This may seem harmless enough, but it can take down all sorts of questions. Case in point:

Example 10.11. (AIME 1987) m, n are integers such that $m^2 + 3m^2n^2 = 30n^2 + 517$. Find $3m^2n^2$.

Solution. Moving things around and Dividing by three will give us the Simon form in m^2, n^2

$$\begin{aligned} m^2n^2 + \frac{m^2}{3} - 10n^2 &= \frac{517}{3} \\ \iff (m^2 - 10)(n^2 + 1/3) &= \frac{507}{3} \\ \iff (m^2 - 10)(3n^2 + 1) &= 507 \end{aligned}$$

This clearly limits our values as (m, n) are integers and $507 = 1 * 507 = 3 * 169 = 13 * 39$. This forces us to $3n^2 + 1 = 13 \iff n = 2$ and therefore $m = 7$. The question wants us to return $3m^2n^2 = 3 * 2^2 * 7^2 = 3 * 4 * 49 = 588$ □

A commonly occurring but hard to notice factorization is the Sophie Germain identity.

Theorem 10.12 (Sophie Germain identity). $x^4 + 4y^4 = (x^2 + 2y^2 - 2xy)(x^2 + 2y^2 + 2xy)$

This is a result of a process called completing the square.

Line of Thought. We know that $a^2 + 2ab + b^2 = (a + b)^2$. Keeping that in mind,

$$\begin{aligned} x^4 + 4y^4 &= (x^2)^2 + (2y^2)^2 \\ &= (x^2)^2 + (2y^2)^2 + 2(x^2)(2y^2) - 2(x^2)(2y^2) \\ \text{We are doing this to use the identity of } (a+b)^2 \\ &= (x^2 + 2y^2)^2 - (2xy)^2 \\ &= (x^2 + 2y^2 - 2xy)(x^2 + 2y^2 + 2xy) \end{aligned} \quad \square$$

The method of thinking here is more useful rather than the identity. We will also use it again in this chapter to unlock an even more powerful theorem. Here is a token use of Sophi

Example 10.13. Prove that $x^4 + 1$ is composite for all integer values of x such that $|x| > 1$.

Proof. What would other wise be a headache with number theory techniques becomes a mild embarrassment using Sophie Germain.

$$\begin{aligned} x^4 + 1 &= x^4 + 4 * 1^4 \\ &= (x^2 + 2 * 1^2 - 2x)(x^2 + 2 * 1^2 + 2x) \\ &= (x^2 + 2 - 2x)(x^2 + 2 + 2x) \end{aligned}$$

The only way for $x^4 + 1$ to be prime is if one of these is equal to 1. That occurs if and only if either $x^2 - 2x + 2 = 1 \iff x^2 - 2x + 1 = 0 \iff (x-1)^2 = 0 \iff x = 1$ or $x^2 + 2x + 2 = 1 \iff x^2 + 2x + 1 = 0 \iff (x+1)^2 = 0 \iff x = -1$ which are both not possible as $|x| > 1$. Hence, proved. \square

The final factorization(which we have already seen in an example) we should keep in mind is(also a result of completing the square):

Theorem 10.14. $x^4 + x^2 + 1 = (x^2 + x + 1)(x^2 - x + 1)$

While there are a lot more factorization which can be used, these are the most common(and almost everything else is derived from these or can be found only on expansion). You'll see there power as the chapter progresses.

10.4 Quadratic Equations

An equation of the form $ax^2 + bx + c = 0$ is known as a quadratic. It has many real life uses and comes up more than one expects. We solved one just above, but not all of them are that friendly. We have three common methods to solve quadratics.

10.4.1 Splitting the middle term. If $a = 1$ and you can find two numbers, α and β such that $\alpha + \beta = b$ and $\alpha \cdot \beta = c$, then:

$$x^2 + bx + c = 0 \Rightarrow x^2 + \alpha x + \beta x + \alpha\beta = 0$$

$$\Rightarrow x(x + \alpha) + \beta(x + \alpha) = 0$$

$$\Rightarrow (x + \beta)(x + \alpha) = 0$$

$$\Rightarrow x = -\beta \text{ or } -\alpha$$

If $a \neq 1$ then either we can divide the equation by a and do the above or look for α and β such that $\alpha + \beta = b$ and $\alpha \cdot \beta = ac$ and then factorize.

Example 10.15. $20x^2 + 51x + 22 = 0$

$$20x^2 + 40x + 11x + 22 = 0$$

$$20x(x + 2) + 11(x + 2) = 0$$

$$(x + 2)(20x + 11) = 0$$

10.4.2 Completing the Square.

Example 10.16. $2x^2 + 7x - 4 = 0$ Here we can see that factorization is quite difficult. So we'll complete the square.

$$x^2 + \frac{7}{2}x - 2 = 0$$

We know that $(x + a)^2 = x^2 + 2ax + a^2$ lets create that over here.

$$x^2 + 2\frac{7}{4}x + \frac{49}{16} - \frac{49}{16} - 2 = 0$$

$$(x + \frac{7}{4})^2 = \frac{49}{16} + 2$$

$$(x + \frac{7}{4})^2 = \frac{49+32}{16} = \frac{81}{16}$$

$$x + \frac{7}{4} = \frac{\pm 9}{4}$$

$$x = \frac{2}{4} \text{ or } \frac{-16}{4}$$

$$x = \frac{1}{2} \text{ or } -4$$

As you might have noticed, this is literally SFFT with $x = y$. But for the times when we don't wish to use even the smallest part of our brain:

10.4.3 Quadratic formula. Using completing the square on the general equation, $ax^2 + bx + c = 0$

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0$$

$$x^2 + 2\frac{b}{2a}x + \frac{b^2}{4a^2} - \frac{b^2}{4a^2} + \frac{c}{a} = 0$$

$$(x + \frac{b}{2a})^2 = \frac{b^2}{4a^2} - \frac{c}{a}$$

$$(x + \frac{b}{2a})^2 = \frac{b^2 - 4ac}{4a^2}$$

$$x + \frac{b}{2a} = \frac{\pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

This is the quadratic formula for every single equation. While a bit messy, it solves the complicated equations the other methods can't.

Theorem 10.17. For $ax^2 + bx + c = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

The part inside the square root, $b^2 - 4ac$ is known as the discriminant and decides the nature of roots.

If the discriminant is positive, we have 2 real roots; if it is zero, one real root and if it is negative no real roots(the roots are then imaginary. We'll talk about them later).

If the discriminant a square number and b and $2a$ are rational, then the roots are rational. If it is not a square number, and b and $2a$ are rational, the roots are irrational.

10.5 Vieta's Formula

The last section was us finding the roots using the coefficients, now we'll use the roots to find the coefficients.

Let α and β be the roots of $ax^2 + bx + c = 0$, then:

$$\begin{aligned} \alpha + \beta &= \frac{-b + \sqrt{b^2 - 4ac}}{2a} + \frac{-b - \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-2b}{2a} \\ &= \frac{b}{a} \\ \text{and also, } \alpha \cdot \beta &= \frac{-b + \sqrt{b^2 - 4ac}}{2a} \cdot \frac{-b - \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{b^2 - b^2 + 4ac}{4a^2} \\ &= \frac{c}{a} \end{aligned}$$

These are known as Vieta's formula. Another derivation of them happens simply using the splitting of middle term. It's quite trivial. Just divide the equation by a and the rest is by definition of roots.

Theorem 10.18. For an equation, $ax^2 + bx + c = 0$

Sum of roots is $\frac{-b}{a}$ and product of roots is $\frac{c}{a}$

However, Vieta can be generalized for higher degree polynomials as well.

Theorem 10.19. For $ax^n + bx^{n-1} + cx^{n-2} \dots = 0$ and the roots being

$r_1, r_2, r_3 \dots r_n$,

$$\sum_{k=1}^n r_k = \frac{-b}{a}$$

$$\sum_{k,l=1}^n = \frac{c}{a}$$

$$\sum_{k,l,m}^n = \frac{-d}{a}$$

\vdots

Note: We are basically taking sum of terms first one at a time, then two at a time(all possible pair of two terms) and then three at a time(all possible triplets of three terms) and so on, till we reach the product.

The proof will require the help of one additional theorem...

10.6 The Fundamental Theorem of Algebra

Theorem 10.20. *For any polynomial $ax^n + bx^{n-1} + cx^{n-2} \dots = 0$ and the roots being $r_1, r_2, r_3 \dots r_n$, we can say*

$$\begin{aligned} ax^n + bx^{n-1} + cx^{n-2} \dots &= 0 \\ \Rightarrow a(x - r_1)(x - r_2)(x - r_3) \dots (x - r_n) &= 0 \end{aligned}$$

The above follows by the definition of being a root. We can now prove the generalized Vieta.

Proof. Let $P(x) = a_n(x - r_1)(x - r_2) \dots (x - r_n)$

Now for some x^k we'll need to choose some k of the n factors to give us the x and $n - k$ to give us the coefficient. The thing is that we can do this in all the ways $n - k$ can be chosen from n .

So the total coefficient will end being $-1^{n-k}a_n$ times the sum of the product of $n - k$ roots in every which way (let's call it S_{n-k}). So we can say $-1^{n-k}aS_{n-k}$ is the coefficient of x^k which we'll denote as c_k .

This leads to $S_{n-k} = -1^{n-k} \frac{c_k}{a} \iff S_k = -1^k \frac{c_{n-k}}{a}$. Which is the generalized Vieta.

□

Another theorem which is a consequence of the same is:

Theorem 10.21. *When $P(x)$ is divided by $x - r$ the remainder is $P(r)$*

Proof. Let $x - a$ divide $P(x)$ and there exists a quotient $Q(x)$ and remainder $R(x)$ such that

$$P(x) = (x - a)Q(x) + R(x)$$

We need to notice that $\deg R(x) < \deg(x - a) = 1$ as if had equal or higher degree, we could divide it further.

Now as $\deg(x - a)$ is 1, $\deg R(x)$ is 0, or it is a constant. Let this constant be r . We may substitute this into our original equation and rearrange to yield

$$r = P(x) - (x - a)Q(x).$$

When $x = a$, this equation becomes $r = P(a)$. Hence, the remainder upon dividing $P(x)$ by $x - a$ is equal to $P(a)$. □

This short theorem might seem rather unremarkable but has a variety of uses. Case in point:

Example 10.22. Given the cubic $f(x) = x^3 + x + 1$, $g(x)$ is another cubic whose roots are the square of the roots of $f(x)$. Given $g(0) = -1$, find $g(9)$

Solution. Let $f(x) = (x - a)(x - b)(x - c)$ making a, b, c roots $f(x)$.

Therefore, $g(x) = k(x - a^2)(x - b^2)(x - c^2)$

Taking $x = 0$,

$$g(0) = -ka^2b^2c^2 = -1 \iff ka^2b^2c^2 = 1$$

We know $abc = 1$ using vieta on $f(x)$.

$$\therefore k * 1 = 1 \iff k = 1$$

$$\therefore g(9) = (9 - a^2)(9 - b^2)(9 - c^2)$$

$$= (3 + a)(3 - a)(3 + b)(3 - b)(3 + c)(3 - c)$$

We can consider $f(3) = (3 - a)(3 - b)(3 - c)$ and $-f(-3) = -1(-3 - a)(-3 - b)(-3 - c) = -1^3(-3 - a)(-3 - b)(-3 - c) = (3 + a)(3 + b)(3 + c)$

$$\therefore g(9) = -f(3) * f(-3) = 899 \quad \square$$

However, there is even a shorter way out...

Alternate solution. As the squares of roots of $f(x)$ are the roots of $g(x)$, it is of the form $g(x) = kf(\sqrt{x})$.

So we have: $x^{3/2} + x^{1/2} + 1 = 0$

$$\iff x^{1/2}(x + 1) = -1$$

$$\iff x(x^2 + 2x + 1) = 1$$

$$\iff x^3 + 2x^2 + x - 1 = 0$$

Hence, $g(x) = k(x^3 + 2x^2 + x - 1)$. As $g(0) = -1$, $k = 1$.

Therefore, $g(x) = x^3 + 2x^2 + x - 1$. We can now simply compute $g(9) = 899 \quad \square$

Another theorem, which provides a valid but tedious and messy way to find rational roots is:

Theorem 10.23. Given an integer polynomial $P(x)$ with leading coefficient a_n and constant term a_0 , if $P(x)$ has a rational root $r = p/q$ in lowest terms, then p divides a_0 and q divides a_n

While not ideal, this theorem is sometimes the only weapon of attack we have against more complicated polynomials. Once we find a single root, we can use the factor theorem and divide the polynomial into simpler parts. Below is a proof for the same:

Proof. Let $\frac{p}{q}$ be in its simplest form and a rational root of $P(x) = a_nx^n + a_{n-1}x^{n-1} + \dots + a_0$, where every a_r is an integer;

Since $\frac{p}{q}$ is a root of $P(x)$,

$$0 = a_n \left(\frac{p}{q}\right)^n + a_{n-1} \left(\frac{p}{q}\right)^{n-1} + \dots + a_1 \left(\frac{p}{q}\right) + a_0.$$

Multiplying by q^n yields

$$0 = a_n p^n + a_{n-1} p^{n-1} q + \cdots + a_1 p * q^{n-1} + a_0 q^n.$$

We need to notice every term other than $a_0 q^n$ is divisible by p and so is zero. Which means $a_0 q^n$ is also divisible by p . However, as p and q have no common factors(as $\frac{p}{q}$ is in its simplest form), $\therefore a_0$ is divisible by p . Simlerly, a_n is divisible by q . \square

We'll give it a whirl with this token example:

Example 10.24. Find all rational roots of $6x^3 + x^2 - 19x + 6$

Solution. The possible values of p are $-6, -3, -2, -1, 1, 2, 3, 6$ and possible values of q are $1, 2, 3, 6$ (if both are negative, it cancels and becomes both positive)

Hence, $\frac{p}{q}$ has possible values, $-6, -3, -2, -1, \frac{-3}{2}, \frac{-1}{2}, \frac{-2}{3}, \frac{-1}{3}, \frac{-1}{6}, \frac{1}{6}, \frac{1}{3}, \frac{2}{3}, \frac{1}{2}, \frac{3}{2}, 1, 2, 3, 6$
Of all the eighteen values, we'll check the simplest ones first. -2 is a root.
Therefore, our polynomial is divisible by $x + 2$ which will leave a a quadratic which we can factorize quite easily.

$$\begin{aligned}\therefore 6x^3 + x^2 - 19x + 6 &= (6x^2 - 11x + 3)(x + 2) \\ &= (2x - 3)(3x - 1)(x + 2)\end{aligned}$$

Hence the rational roots are: $\frac{3}{2}, \frac{1}{3}$ and 2 . \square

10.7 Newton's Sums

Next, we'll discuss Newton Sums. We can add roots being multiplied to other roots using Vieta. What about squares of all the roots? What about the cubes? This solves questions which are seemingly complex using Vieta's very easily.

Theorem 10.25. For $P(x) = a_n x^n + a_{n-a} x^{n-a} + \cdots + a_1 x + a_0$ with roots $r_1, r_2, r_3, \dots, r_n$,

Let: $S_1 = r_1 + r_2 + \dots + r_n$

$$S_2 = r_1^2 + r_2^2 + \dots + r_n^2$$

\vdots

$$S_k = r_1^k + r_2^k + \dots + r_n^k$$

then the following will hold true:

$$a_n S_1 + a_{n-1} = 0$$

$$a_n S_2 + a_{n-1} S_1 + 2a_{n-2} = 0$$

\vdots

Basically what the theorem says is:

- (1) Start with a S_k value and multiply by it by the leftmost polynomial coefficient.
- (2) Then, multiply S_{k-1} by the polynomial's coefficient right after it.
- (3) Continue doing so and summing the products until a_{k-i} becomes 0 in which case we simply add the last term and stop
- (4) Set your final sum of terms to be equal to 0

Proof. Let $\alpha, \beta, \gamma, \dots, \omega$ be the roots of a given polynomial $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$. Then, we have that

$$P(\alpha) = P(\beta) = P(\gamma) = \dots = P(\omega) = 0$$

Thus,

$$\begin{cases} a_n \alpha^n + a_{n-1} \alpha^{n-1} + \dots + a_0 = 0 \\ a_n \beta^n + a_{n-1} \beta^{n-1} + \dots + a_0 = 0 \\ \vdots \\ a_n \omega^n + a_{n-1} \omega^{n-1} + \dots + a_0 = 0 \end{cases}$$

Multiplying each equation by $\alpha^{k-n}, \beta^{k-n}, \dots, \omega^{k-n}$, respectively,

$$\begin{cases} a_n \alpha^{n+k-n} + a_{n-1} \alpha^{n-1+k-n} + \dots + a_0 \alpha^{k-n} = 0 \\ a_n \beta^{n+k-n} + a_{n-1} \beta^{n-1+k-n} + \dots + a_0 \beta^{k-n} = 0 \\ \vdots \\ a_n \omega^{n+k-n} + a_{n-1} \omega^{n-1+k-n} + \dots + a_0 \omega^{k-n} = 0 \end{cases}$$

$$\begin{cases} a_n \alpha^k + a_{n-1} \alpha^{k-1} + \dots + a_0 \alpha^{k-n} = 0 \\ a_n \beta^k + a_{n-1} \beta^{k-1} + \dots + a_0 \beta^{k-n} = 0 \\ \vdots \\ a_n \omega^k + a_{n-1} \omega^{k-1} + \dots + a_0 \omega^{k-n} = 0 \end{cases}$$

now taking the sum,

$$\begin{aligned} & a_n \underbrace{(\alpha^k + \beta^k + \dots + \omega^k)}_{P_k} + a_{n-1} \underbrace{(\alpha^{k-1} + \beta^{k-1} + \dots + \omega^{k-1})}_{P_{k-1}} + \\ & + a_0 \underbrace{(\alpha^{k-n} + \beta^{k-n} + \dots + \omega^{k-n})}_{P_{k-n}} = 0 \end{aligned}$$

Therefore,

$$a_n P_k + a_{n-1} P_{k-1} + a_{n-2} P_{k-2} + \dots + a_0 P_{k-n} = 0.$$

□

This is a monsterous proof. No wonder Newton got it during the black plague.
Let's now use the theorem as a example:

Example 10.26. (USAMO 1973) Determine all the roots, real or complex, of the system of simultaneous equations $x + y + z = 3$, $x^2 + y^2 + z^2 = 3$, $x^3 + y^3 + z^3 = 3$.

Solution. Let x, y, z be the roots of some cubic polynomial $f(t) = at^3 + bt^2 + ct + d$.

Let $x + y + z = S_1 = 3$, $x^2 + y^2 + z^2 = S_2 = 3$ and $x^3 + y^3 + z^3 = S_3 = 3$

Then using newton sums: $3a + b = 3$; $3a + 3b + 2c = 0$; $3a + 3b + 3c + 3d = 0$
as $b = -3a$ and $a + b + c + d = 0$, $\therefore 3a + 3b + 2c = 0 \iff 2b + 2c = 0 \iff b + c = 0 \iff b = -c$

$$\therefore a = -d$$

$$\therefore f(t) = at^3 - 3at^2 + 3at - a$$

Now we need to find roots of this equation:

$$at^3 - 3at^2 + 3at - a = 0$$

$$\iff t^3 - 3t^2 + 3t - 1 = 0 \iff (t - 1)^3 = 0$$

$$\iff t = 1$$

Thus, the only possible solution is $x = y = z = 1$

□

10.8 Reciprocal Relations

The given relations tend to come up time to time in speed solving competitions(where you should remember them) and admission tests. There only other use occurs in the simplification of even symmetric polynomials(see below). So it is recommended to memorize them by simply deriving them yourself. Do it the next day as well.

You will now not forget them for rest of your days.

Theorem 10.27. If $x + \frac{1}{x} = a$ then:

$$x^2 + \frac{1}{x^2} = a^2 - 2$$

$$x^3 + \frac{1}{x^3} = a^3 - 3a$$

$$x^4 + \frac{1}{x^4} = (a^2 - 2)^2 - 2$$

Theorem 10.28. If $x - \frac{1}{x} = a$ then:

$$x^2 + \frac{1}{x^2} = a^2 + 2$$

$$x^3 - \frac{1}{x^3} = a^3 + 3a$$

$$x^4 + \frac{1}{x^4} = (a^2 + 2)^2 + 2$$

10.9 Some Special Polynomials

We are going to talk about a few special polynomials now. These tend to appear a lot and we should be alert when they do as they are like Unicorn sightings, scary at first but wonderful on retrospection.

10.9.1 Symmetric Polynomial.

Definition 10.29. A polynomial $P(x) = a_nx^n + a_{n-1}x^{n-1} \cdots + a_1x + a_0$ is symmetric if and only if:

$$\begin{aligned} a_n &= a_0 \\ a_{n-1} &= a_1 \\ &\vdots \\ a_{n-k} &= a_k \end{aligned}$$

If it is of an even degree, it can be solved with the following algorithm:

Theorem 10.30 (Algorithm). (1) *Divide by $x^{n/2}$*

- (2) *Group x^k with $\frac{1}{x^k}$*
- (3) *Make the substitution $t = x + \frac{1}{x}$*
- (4) *Solve the simplified polynomial.*

Let's solve an example for more clarity.

Example 10.31. The real number x satisfies the equation $x + \frac{1}{x} = \sqrt{5}$. What is the value of $x^{11} - 7x^7 + x^3$?

Solution. Before you complain that the given equation is neither symmetric nor of even degree, take note that it is neither symmetric nor of even degrees. We can see that $a_0 = a_1 = a_2 = 0$ and hence I can keep $a_{14} = a_{13} = a_{12} = 0$. Making it both even and symmetric.

$$\begin{aligned} &\therefore x^{11} - 7x^7 + x^3 \\ &= 0x^{14} + x^{11} - 7x^7 + x^3 + 0 \text{ Now let's divide by } x^7 \\ &= x^4 - 7 + \frac{1}{x^4} \text{ using the reciprocal relations,} \\ &= (\sqrt{5}^2 - 2)^2 - 2 - 7 \\ &= 0 \end{aligned}$$

□

10.9.2 Reciprocal Roots.

Theorem 10.32. If the roots of polynomial $P(x) = a_nx^n + a_{n-1}x^{n-1} \cdots + a_1x + a_0$ are r_1, r_2, \dots, r_n then the roots of $a_0x^n + a_1x^{n-1} \cdots + a_{n-1}x + a_n$ are

$$\frac{1}{r_1}, \frac{1}{r_2}, \dots, \frac{1}{r_n}$$

No proof is required as it is quite trivial to see using Vieta.

10.9.3 Adding to all roots.

Theorem 10.33. If the roots of polynomial $P(x) = a_nx^n + a_{n-1}x^{n-1} \cdots + a_1x + a_0$ are r_1, r_2, \dots, r_n then the roots of $a_n(x - k)^n + a_{n-1}(x - k)^{n-1} \cdots + a_1(x - k) + a_0$ are $r_1 + k, r_2 + k, \dots, r_n + k$

This doesn't require a proof as it follows from the definition of root.
Now let's get started with some exercises.

Exercises

- (1) (AIME) Solve for positive value of x:

$$\frac{1}{x^2-10x-29} + \frac{1}{x^2-10x-45} - \frac{2}{x^2-10x-69} = 0$$

- (2) (AIME) Let (a, b, c) be the real solution of the system of equations $x^3 - xyz = 2, y^3 - xyz = 6, z^3 - xyz = 20$. The greatest possible value of $a^3 + b^3 + c^3$ can be written in the form $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$

- (3) (AIME) Find $x^2 + y^2$ if x and y are positive integers such that

$$\begin{aligned} xy + x + y &= 71, \\ x^2y + xy^2 &= 880. \end{aligned}$$

- (4) (AIME) The equation $2^{333x-2} + 2^{111x+2} = 2^{222x+1} + 1$ has three real roots.

Given that their sum is m/n where m and n are relatively prime positive integers, find $m + n$.

- (5) (AIME) What is the product of the real roots of the equation $x^2 + 18x + 30 = 2\sqrt{x^2 + 18x + 45}$?

- (6) If the roots of the quadratic $3x^2 + 6x - 5$ are r and s, find $r^3 + s^3$.

- (7) If r, s, and t are roots of the cubic $x^3 - 6x - 5 = 0$, find $r^3 + s^3 + t^3$?

- (8) (AIME) Suppose that the roots of $x^3 + 3x^2 + 4x - 11 = 0$ are a, b, and c, and that the roots of $x^3 + rx^2 + sx + t = 0$ are $a+b, b+c$, and $c+a$. Find t.?

- (9) Solve for real roots of $10000x^4 - 5000x^3 + 825x^2 - 50x + 1 = 0$

- (10) Find the products of the solutions for:

$$\sqrt{5|x| + 8} = \sqrt{x^2 - 16}$$

- (11) (AIME) The only real root of $8x^3 - 3x^2 - 3x - 1 = 0$ can be written as $\frac{\sqrt[3]{a} + \sqrt[3]{b} + 1}{c}$ in simplest form. What is $a + b + c$
- (12) (AMC 12) A rectangular floor measures a by b feet, where a and b are positive integers with $b > a$. An artist paints a rectangle on the floor with the sides of the rectangle parallel to the sides of the floor. The unpainted part of the floor forms a border of width 1 foot around the painted rectangle and occupies half of the area of the entire floor. How many possibilities are there for the ordered pair (a, b) ?
- (13) (AMC 12) For certain real numbers a , b , and c , the polynomial
- $$g(x) = x^3 + ax^2 + x + 10$$
- has three distinct roots, and each root of $g(x)$ is also a root of the polynomial
- $$f(x) = x^4 + x^3 + bx^2 + 100x + c.$$
- What is $f(1)$?
- (14) (IOQM 2023) A positive integer m has the property that m^2 is expressible in the form $4n^2 - 5n + 16$ where n is an integer (of any sign). Find the maximum possible value of $|m - n|$.
- (15) (AIME) The polynomial $P(x)$ is cubic. What is the largest value of k for which the polynomials $Q_1(x) = x^2 + (k - 29)x - k$ and $Q_2(x) = 2x^2 + (2k - 43)x + k$ are both factors of $P(x)$?
- (16) (CAMO 1998) For what real values of k do $1988x^2 + kx + 8891$ and $8891x^2 + kx + 1988$ have a common zero?
- (17) (AIME 2018) A real number a is chosen randomly and uniformly from the interval $[-20, 18]$. The probability that the roots of the polynomial $x^4 + 2ax^3 + (2a - 2)x^2 + (-4a + 3)x - 2$ are all real can be written in the form $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.
- (18) (AIME 2001) Find the sum of the roots, real and non-real, of the equation $x^{2001} + (\frac{1}{2} - x)^{2001} = 0$, given that there are no multiple roots.
- (19) (AIME 1996) Suppose that the roots of $x^3 + 3x^2 + 4x - 11 = 0$ are a , b , and c , and that the roots of $x^3 + rx^2 + sx + t = 0$ are $a + b$, $b + c$, and $c + a$. Find t .
- (20) (AIME 2005) The equation $2^{333x-2} + 2^{111x+2} = 2^{222x+1} + 1$ has three real roots. Given that their sum is m/n where m and n are relatively prime positive integers, find $m + n$.

- (21) (AIME 1983) Suppose that the sum of the squares of two complex numbers x and y is 7 and the sum of the cubes is 10. What is the largest real value that $x + y$ can have?
- (22) (AIME 2015) Let x and y be real numbers satisfying $x^4y^5 + y^4x^5 = 810$ and $x^3y^6 + y^3x^6 = 945$. Evaluate $2x^3 + (xy)^3 + 2y^3$.
- (23) (PRMO 2019) Let $f(x) = x^2 + ax + b$. If for all nonzero real x :
- $$f\left(x + \frac{1}{x}\right) = f(x) + \frac{1}{x}$$
- If the roots of $f(x)$ are integers, what is the value of $a^2 + b^2$?
- (24) $x + y + z = 1$
 $x^2 + y^2 + z^2 = 2$
 $x^3 + y^3 + z^3$
Find $x^4 + y^4 + z^4$?
- (25) (INMO 1991) How many ordered triples (x, y, z) of real numbers satisfy the system of equations:
 $x^2 + y^2 + z^2 = 9$
 $x^4 + y^4 + z^4 = 33$
 $xyz = -4$
- (26) (AIME 2016) Let $P(x)$ be a nonzero polynomial such that $(x-1)P(x+1) = (x+2)P(x)$ for every real x , and $(P(2))^2 = P(3)$. Then $P\left(\frac{7}{2}\right) = \frac{m}{n}$, where m and n are relatively prime positive integers. Find $m+n$.
- (27) (AIME 1983) What is the product of the real roots of the equation $x^2 + 18x + 30 = 2\sqrt{x^2 + 18x + 45}$?
- (28) (USAMO 1984) In the polynomial $x^4 - 18x^3 + kx^2 + 200x - 1984 = 0$, the product of 2 of its roots is -32 . Find k .
- (29) (RMO 2007) Let $P(x) = x^2 + \frac{x}{2} + b$ and $Q(x) = x^2 + cx + d$ be two polynomials with real coefficients such that $P(x)Q(x) = Q(P(x))$ for all real x . Find all real roots of $P(Q(x)) = 0$.
- (30) (PRMO 2018) Let $P(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n$ be a polynomial in which a_i is non-negative integer for each $i \in 0, 1, 2, 3, \dots, n$. If $P(1) = 4$ and $P(5) = 136$, what is the value of $P(3)$?
- (31) (PRMO 2018) Integers a, b, c satisfy $a + b - c = 1$ and $a^2 + b^2 - c^2 = -1$. What is the sum of possible values of $a^2 + b^2 + c^2$?
- (32) (CMI 2019) Find all roots of

$$\frac{8^x + 27^x}{12^x + 18^x} = \frac{7}{6}$$

- (33) (AIME 1986) The polynomial $1 - x + x^2 - x^3 + \cdots + x^{16} - x^{17}$ may be written in the form $a_0 + a_1y + a_2y^2 + \cdots + a_{16}y^{16} + a_{17}y^{17}$, where $y = x + 1$ and the a_i 's are constants. Find the value of a_2 .

11

Inequalities

Till now we were dealing with the cases where $f(x) = g(y)$. The thing to note is that they were mostly equal. This is normally not true in real life. Say you have a budget of 10,000 dollars for your wedding. Our expenditure needs to be less than or equal to 10,000 dollars. I don't think people will hold it against you for saving money.

This chapter doesn't use = much. Instead we'll have two sides which will not be equal.

11.1 Another note on notation

We will make use of two new notations, \sum_{cyc} and \sum_{sym} which mean to cycle through all the n values and go through all the $n!$ values respectively. For example:

$$\sum_{\text{cyc}} a^2 = a^2 + b^2 + c^2 + d^2$$

$$\sum_{\text{cyc}} ab = ab + bc + ca + ad$$

$$\sum_{\text{sym}} a^2 = 6(a^2 + b^2 + c^2 + d^2)$$

$$\sum_{\text{sym}} ab = ab + ac + ad + bc + bd + ad$$

11.2 AM-GM and Muirhead

Theorem 11.1 (AM-GM).

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

The equality holds, if and only if, $x_1 = x_2 = x_3 \cdots = x_n$

Proof. AM-GM can be proven in various ways. However, the simplest and most logical one is presented here.

We'll refer to the inequality as I_n with n being the number of terms.

Let's first prove it for $n = 2$.

Let $x_1 + k = x_2$. Proof by induction:

$$(B) \text{ for } k = 1, \left(\frac{x_1+x_1+1}{2}\right)^2$$

$$= (x_1 + \frac{1}{2})^2$$

$$= x_1^2 + x_1 + \frac{1}{4} > x_1(x_1 + 1) = x_1^2 + x_1$$

(S) Let's assume this is true for some k leading to $x_1^2 + kx_1 + \frac{k^2}{4} > x_1(x_1 + k)$, then:

$$x_1(x_1 + k + 1)$$

$$= x_1^2 + kx_1 + x_1$$

$$= x_1(x_1 + k) + x_1$$

$$< x_1^2 + kx_1 + \frac{k}{4} + x_1$$

$$< x_1^2 + (k + 1)x_1 + \frac{(k+1)^2}{4}$$

Hence, proved.

Now with this information, let's do something unique. Assuming, I_n holds, let's prove that I_{2n} also holds.

Let x_1, x_2, \dots, x_{2n} be any list of nonnegative reals. Then, because the two lists x_1, x_2, \dots, x_n and $x_{n+1}, x_{n+2}, \dots, x_{2n}$, each have n variables,

$$\frac{x_1 + x_2 + \cdots + x_n}{n} \geq \sqrt[n]{x_1 x_2 \cdots x_n}$$

and

$$\frac{x_{n+1} + x_{n+2} + \cdots + x_{2n}}{n} \geq \sqrt[n]{x_{n+1} x_{n+2} \cdots x_{2n}}.$$

Adding these two inequalities together and dividing by 2 yields

$$\frac{x_1 + x_2 + \cdots + x_{2n}}{2n} \geq \frac{\sqrt[n]{x_1 x_2 \cdots x_n} + \sqrt[n]{x_{n+1} x_{n+2} \cdots x_{2n}}}{2}.$$

Using I_2 , on $\sqrt[n]{x_1 x_2 \cdots x_n}$ and $\sqrt[n]{x_{n+1} x_{n+2} \cdots x_{2n}}$ we get

$$\frac{\sqrt[n]{x_1 x_2 \cdots x_n} + \sqrt[n]{x_{n+1} x_{n+2} \cdots x_{2n}}}{2} \geq \sqrt[2n]{x_1 x_2 \cdots x_{2n}}.$$

Plugging this in proves this for I_{2n}

Finally, we'll perform the third and final (S) this time we'll assume that I_n holds and prove that so does I_{n-1} , causing all the dominos to topple.

Letting $x_n = \frac{x_1+x_2+\cdots+x_{n-1}}{n-1}$, we have that

$$\frac{x_1 + x_2 + \cdots + x_{n-1} + \frac{x_1+x_2+\cdots+x_{n-1}}{n-1}}{n} \geq \sqrt[n]{x_1 x_2 \cdots x_{n-1} \left(\frac{x_1 + x_2 + \cdots + x_{n-1}}{n-1} \right)}.$$

Because we assumed AM-GM in n variables, equality holds if and only if $x_1 = x_2 = \dots = x_{n-1} = \frac{x_1+x_2+\dots+x_{n-1}}{n-1}$. However, note that the last equality is implied if all the numbers of x_1, x_2, \dots, x_{n-1} are the same; thus, equality holds if and only if $x_1 = x_2 = \dots = x_{n-1}$.

We first simplify the lefthand side. Multiplying both sides of the fraction by $n-1$ and combining like terms, we get that

$$\frac{x_1 + x_2 + \dots + x_{n-1} + \frac{x_1+x_2+\dots+x_{n-1}}{n-1}}{n} = \frac{nx_1 + nx_2 + \dots + nx_{n-1}}{n(n-1)} = \frac{x_1 + x_2 + \dots + x_{n-1}}{n-1}$$

Plugging this into the earlier inequality yields

$$\frac{x_1 + x_2 + \dots + x_{n-1}}{n-1} \geq \sqrt[n]{x_1 x_2 \dots x_{n-1} \left(\frac{x_1 + x_2 + \dots + x_{n-1}}{n-1} \right)}.$$

Raising both sides to the n th power yields

$$\left(\frac{x_1 + x_2 + \dots + x_{n-1}}{n-1} \right)^n \geq x_1 x_2 \dots x_{n-1} \left(\frac{x_1 + x_2 + \dots + x_{n-1}}{n-1} \right).$$

From here, we divide by $\frac{x_1+x_2+\dots+x_{n-1}}{n-1}$ and take the $(n-1)$ th root to get that

$$\frac{x_1 + x_2 + \dots + x_{n-1}}{n-1} \geq \sqrt[n-1]{x_1 x_2 \dots x_{n-1}}.$$

This is I_{n-1} .

With the most advanced thing we have used and will probably use in induction, we can finally say:

Hence, proved. □

This theorem is the king of inequalities. A lot of people believe it to be one of the most important facts in maths. It stands for 'arithmetic mean, geometric mean'. While it is quite a well known inequality, its proof is less known. If you ever find yourself having crush on a math lover, just present this proof from memory. Thank me later.

Now back to math, We can use AM-GM to get the following things:

$$a^2 + b^2 \geq 2ab \text{ and } a^3 + b^3 + c^3 \geq 3abc$$

You can sum such inequalities to solve simple questions.

Example 11.2. For $a, b, c > 0$ prove that $a^2 + b^2 + c^2 \geq ab + bc + ca$

Proof. As we just discussed above, $a^2 + b^2 \geq 2ab$, $b^2 + c^2 \geq 2bc$, $c^2 + a^2 \geq 2ac$. Adding them gives us:

$$2a^2 + 2b^2 + 2c^2 \geq 2ab + 2bc + 2ca$$

$$\therefore a^2 + b^2 + c^2 \geq ab + bc + ca$$

□

Example 11.3. For $a, b, c > 0$ prove that $a^4 + b^4 + c^4 \geq a^2bc + ab^2c + abc^2$

Proof. $a^4 + a^4 + b^4 + c^4 \geq 4a^2bc$

$$a^4 + b^4 + b^4 + c^4 \geq 4ab^2c$$

$$a^4 + b^4 + c^4 + c^4 \geq 4abc^2$$

Adding these gives us the inequality. \square

In most symmetric inequality, both sides will be equal when we set all variables equal.

Moreover, we often compare expressions which are the same degree, or homogeneous. For example when we write $a^2 + b^2 + c^2 \geq ab + bc + ca$, both sides are degree 2. (Notice that the AM-GM inequality itself has the same property!)

The reason for this is x^5 and x^3 are not comparable for generic $x > 0$, since the behaviors when x is very small and x is very large are different. So a non-homogeneous inequality like $a^2 + b^2 + c^2 \geq a^3 + b^3 + c^3$ will definitely not be true in general, since the behaviors if $a = b = c = 0.01$ and $a = b = c = 100$ will be different. You may have already picked up some intuition: more “mixed” terms are smaller. For example, for degree 3, the polynomial $a^3 + b^3 + c^3$ is biggest and $3abc$ is the smallest. Roughly, the more “mixed” polynomials are the smaller.

This intuition can be formalized as Muirhead inequality

Definition 11.4. If $x_1 + x_2 + \dots + x_k \geq y_1 + y_2 + \dots + y_k$ for all $0 \leq k \leq n$ then $x_1, x_2, \dots, x_n \succ y_1, y_2, \dots, y_n$

The \succ symbol in speaking changes to majorizes. Using this definaion, we can say

Theorem 11.5 (Muirhead).

$$\sum_{sym} a_1^{x_1} a_2^{x_2} \dots a_n^{x_n} \geq \sum_{sym} a_1^{y_1} a_2^{y_2} \dots a_n^{y_n}$$

If $x_1, x_2, x_3 \dots x_n \succ y_1, y_2, y_3 \dots y_n$

Since, $(5, 0, 0) \succ (3, 1, 1) \succ (2, 2, 1)$

$$a^5 + a^5 + b^5 + b^5 + c^5 + c^5 \geq a^3bc + a^3bc + ab^3c + ab^3c + abc^3 + abc^3 \geq a^2b^2c + a^2b^2 + ab^2c^2 + ab^2c^2 + a^2bc^2 + a^2bc^2$$

Note: It is symmetric, not cyclic. Our only tool for cyclic is still AM-GM.

11.3 Non-homogeneous equations

11.3.1 Even's Substitution. If an inequality has the condition $abc = 1$, one can also sometimes use the substitution $(a, b, c) = (x/y, y/z, z/x)$ which

will transform it into a homogeneous inequality automatically. I call it the Even's substitution as I first learnt it from Evan Chan's handouts(Its not an official name.).

Example 11.6. Prove that if $abc = 1$ then $a^2 + b^2 + c^2 \geq a + b + c$

Proof. Using Evan's substitution, the problem statement transforms to $\frac{x^2y^4+y^2z^4+z^2x^4}{x^2y^2z^2} \geq \frac{xy^2+yz^2+zx^2}{xyz}$
 $\iff x^2y^4+y^2z^4+z^2x^4 \geq x^2y^3z+xy^2z^3+x^3yz^2$
Using, $4x^2y^4+y^2z^4+z^2x^4 \geq 6x^2y^3z$, and adding its cycles, we get the inequality. \square

11.3.2 Ravi's Substitution. Sometimes, an inequality will refer to the a, b, c as the sides of a triangle. In that case, one can replace $(a, b, c) = (y+z, z+x, x+y)$ where $x, y, z > 0$ are real numbers using the properties of in-circle.

This is colloquially known as the Ravi substitution, after Ravi Vakil, a well




Figure 11.0. Ravi's Substitution

known math professor at Stanford and an IMO gold medalist(and 4 time Putnam fellow)

Example 11.7. Find the smallest constant k such that $k > \frac{a^2+b^2+c^2}{ab+bc+ca}$ where a, b, c are the sides of a triangle.

Proof. Using Ravi's Substitution,

$$\begin{aligned} & \frac{a^2+b^2+c^2}{ab+bc+ca} \\ &= \frac{(x+y)^2+(y+z)^2+(z+x)^2}{(x+y)(y+z)+(y+z)(z+x)+(x+z)(x+y)} \\ &= \frac{2(x^2+y^2+z^2+xy+yz+zx)}{x^2+y^2+z^2+3(xy+yz+zx)} \end{aligned}$$

Hence we are looking for k such that:

$$k(x^2+y^2+z^2)+3k(xy+yz+zx) > 2(x^2+y^2+z^2)+2(xy+yz+zx)$$

$$(3k - 2)(xy + yz + zx) > (2 - k)(x^2 + y^2 + z^2)$$

We already know that $x^2 + y^2 + z^2 > xy + yz + zx$; Hence,

$$(3k - 2)(x^2 + y^2 + z^2) > (2 - k)(x^2 + y^2 + z^2)$$

$$3k - 2 > 2 - k$$

$$4k > 4$$

$$k > 1$$

Thus, minimum value of k is 1. \square

11.3.3 Schur's Inequality. It's canny how the three tools for non homogeneity all are named after someone but for completely different reasons. This one was discovered by Issai Schur, the Russian mathematician and is hence named after him.

Theorem 11.8 (Schur's Inequality). *For all non-negative $a, b, c \in \mathbb{R}$ and $r > 0$:*

$$a^r(a - b)(a - c) + b^r(b - a)(b - c) + c^r(c - a)(c - b) \geq 0$$

The four equality cases occur when $a = b = c$ or when two of a, b, c are equal and the third is 0.

Proof. Let $a \geq b \geq c$.

$$\begin{aligned} & \therefore a^r(a - b)(a - c) + b^r(b - a)(b - c) \\ &= a^r(a - b)(a - c) - b^r(a - b)(b - c) \\ &= (a - b)(a^r(a - c) - b^r(b - c)) \end{aligned}$$

Clearly, $a^r \geq b^r \geq 0$, and $a - c \geq b - c \geq 0$.

Thus, $(a - b)(a^r(a - c) - b^r(b - c)) \geq 0$

$$\therefore a^r(a - b)(a - c) + b^r(b - a)(b - c) \geq 0.$$

As, $c^r(c - a)(c - b) \geq 0$

$$\therefore a^r(a - b)(a - c) + b^r(b - a)(b - c) + c^r(c - a)(c - b) \geq 0. \quad \square$$

11.4 Some advanced inequalities

11.4.1 Extended AM-GM.

Theorem 11.9 (Weighted Power Mean). *For $a_1, a_2, a_3 \dots a_n$ positive reals and $w_1, w_2, w_3, \dots w_n$ where $w_1 + w_2 + w_3 + \dots + w_n = 1$. For any real number R , we'll define the weighted power mean as:*

$$P(x) = \begin{cases} (w_1 a_1^r + w_2 a_2^r + \dots + w_n a_n^r)^{1/r} & r \neq 0 \\ a_1^{w_1} a_2^{w_2} \dots a_n^{w_n} & r = 0 \end{cases}$$

Then, if $k > l$ then $P(k) \geq P(l)$. Equality occurs when $a_1 = a_2 = \dots = a_n$

Theorem 11.10 (Simplified Weighted Mean). *The r^{th} power mean refers to:*

$$p(x) = \begin{cases} \left(\frac{a_1^r + a_2^r + \dots + a_n^r}{n}\right)^{1/r} & r \neq 0 \\ \sqrt[n]{a_1 a_2 \dots a_n} & r = 0 \end{cases}$$

Basically, we have set $w_1 = w_2 = \dots = w_n = \frac{1}{n}$. The inequality still holds true, if $k > l$ then $P(k) \geq P(l)$. Equality occurs when $a_1 = a_2 = \dots = a_n$

Using the Simplified Weighted Mean, and setting r to $(2, 1, 0, -1)$ respectively will give us:

Theorem 11.11 (RMS-AM-GM-HM).

$$\begin{aligned} \sqrt{\frac{a_1^2 + a_2^2 + \dots + a_n^2}{n}} &\geq \frac{a_1 + a_2 + \dots + a_n}{n} \\ &\geq \sqrt[n]{a_1 \cdot a_2 \cdot \dots \cdot a_n} \\ &\geq \frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n}} \end{aligned}$$

I'll not prove any of these. While RMS-AM-GM-HM can be proven using the same mechanism we used to prove AM-GM, the weighted form will use some math which we are yet to explore. That technique appears in chapter 15

11.4.2 Cauchy-Schwarz(aka SEBACS). The following theorem is taught under many names. Cauchy had originally proposed it, it was generalized by Schwarz and Bunyakovsky, given an Olympiad friendly form by Arthur Engel and popularized by Titu Andreescu and Nairi Sedrakyan. I propose it being called SEBACS Inequality.

Theorem 11.12 (Original Form). *for any list of reals a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n ,*

$$(a_1^2 + a_2^2 + \dots + a_n^2)(b_1^2 + b_2^2 + \dots + b_n^2) \geq (a_1 b_1 + a_2 b_2 + \dots + a_n b_n)^2,$$

Theorem 11.13 (Modern Form). *for any list of reals a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n :*

$$\frac{a_1^2}{b_1} + \frac{a_2^2}{b_2} + \dots + \frac{a_n^2}{b_n} \geq \frac{(a_1 + a_2 + \dots + a_n)^2}{b_1 + b_2 + \dots + b_n}.$$

Proving any of these will require us to use vectors which we'll learn later (in geometry). With them it will become quite trivial and we'll prove it then. The non-geometrical proof is found in chapter 15

For now, let's try an example:

Example 11.14. for positive a, b, c , prove that $\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2}$,

Proof. By SEBACS Inequality:

$$[\sqrt{(b+c)^2} + \sqrt{(c+a)^2} + \sqrt{(a+b)^2}] \left(\sqrt{\frac{1}{b+c}}^2 + \sqrt{\frac{1}{c+a}}^2 + \sqrt{\frac{1}{a+b}}^2 \right) \geq (\sqrt{\frac{b+c}{b+c}} + \sqrt{\frac{a+c}{a+c}} + \sqrt{\frac{b+a}{b+a}})^2,$$

Upon expanding this gives us:

$$2 \left(\frac{a+b+c}{b+c} + \frac{a+b+c}{c+a} + \frac{a+b+c}{a+b} \right) \geq 9$$

$$\Leftrightarrow \frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2}$$

□

11.5 A note

A lot of more advanced inequalities need calculus for complete exploration. Hence, we've decided to explore them in greater detail in Chapter-15. Inequalities tend to not occur in lower levels and occur at an advanced state at higher level. Hence, we have been unable to include a lot of PYQs here.

Exercises

- (1) Prove that $a^5 + b^5 + c^5 \geq a^3bc + b^3ca + c^3ab \geq abc(ab + bc + ca)$.
- (2) (PRMO) What is the largest positive integer n such that $\frac{a^2}{\frac{a}{29} + \frac{c}{31}} + \frac{b^2}{\frac{c}{29} + \frac{a}{31}} + \frac{c^2}{\frac{a}{29} + \frac{b}{31}} \geq n(a + b + c)$
- (3) For a, b, c the sides of some triangle, prove the inequality $\frac{1}{b+c-a} + \frac{1}{c+a-b} + \frac{1}{a+b-c} \geq \frac{1}{a} + \frac{1}{b} + \frac{1}{c}$
- (4) (Candian MO) For positive reals a, b, c ,

$$\frac{a^3}{bc} + \frac{b^3}{ac} + \frac{c^3}{ab} \geq a + b + c$$

- (5) Prove that for any non-negative a, b, c :

$$\frac{a}{b+c} + \frac{b}{a+c} + \frac{c}{a+b} \geq \frac{3}{2}$$

- (6) Prove that:

$$\sqrt{a_1^2 + a_2^2 + \dots} + \sqrt{b_1^2 + b_2^2 + \dots} \geq \sqrt{(a_1 + b_1)^2 + (a_2 + b_2)^2 + \dots}$$

(7) If $a, b, c > 0$, prove that:

$$a^3b + b^3c + c^3a \geq abc(a + b + c)$$

(8) (IMO 1995) Let a, b, c be positive real numbers such that $abc = 1$. Prove that

$$\frac{1}{a^3(b+c)} + \frac{1}{b^3(c+a)} + \frac{1}{c^3(a+b)} \geq \frac{3}{2}.$$

(9) If $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 1$, then $(a+1)(b+1)(c+1) \geq 64$

(10) . If $abcd = 1$, then $a^4b + b^4c + c^4d + d^4a \geq a + b + c + d$.

(11) Prove that:

$$\frac{a}{b+c} + \frac{b}{a+c} + \frac{c}{a+b} \geq \frac{3}{2}$$

for any non-negative a, b, c

(12) If $a, b, c > 0$ prove that

$$abc(a+b+c) \geq a^3b + b^3c + c^3a$$

(13) For $a, b, c > 0$, prove that:

$$abc \geq (a+b-c)(b+c-a)(c+a-b)$$

(14) Let a, b, c, d be non-negative real numbers such that $a + b + c + d = 1$.

Find the minimum value of $a^2 + b^2 + c^2 + d^2$

(15) (AMC 10) Let A, M , and C be nonnegative integers such that $A + M + C = 10$. What is the maximum value of $A \cdot M \cdot C + A \cdot M + M \cdot C + C \cdot A$?

(16) (AMC 10) Let a, b , and c be real numbers such that

$$a + b + c = 2, \text{ and}$$

$$a^2 + b^2 + c^2 = 12$$

What is the difference between the maximum and minimum possible values of c ?

(17) (MPFG) Find the least real number K such that for all real numbers x and y , we have $(1 + 20x^2)(1 + 19y^2) \geq Kxy$

(18) Prove that

$$\frac{x}{x+y} + \frac{y}{y+z} + \frac{z}{z+x} \leq 2$$

12

Sequence and Series

I normally like to start this chapter with a very well known story. Let's go to a classroom in somewhere near 1785. The math teacher is a man who is not interested in teaching and wants to chat with the female school workers. So he decides that he must keep you all engaged and quiet while he tries to get a life partner. He writes on the board: $1 + 2 + 3 + \dots + 100$ Do not speak until complete

Saying that he leaves. While everyone else is diligently adding, one of the kid stands up after a minute and walks outside the class to find the teacher. "The answer is 5050" he announces.

"It's wrong!" The teacher replies, as he doesn't really know the correct answer. He just figures that a 8 year old couldn't add this quickly.

The child replies, "Sir, its correct."

"Are you the teacher or am I?"

"With all due respect, sir it's correct."

"See, you could by no means add so quickly. You are just guessing."

"Sir, I didn't add much. I multiplied."

"What?"

" $1 + 100 = 101$, $2 + 99 = 101$ and so on, $50 + 51 = 101$. We have 50 such pairs. So the answer must be $50 * 101 = 5050$." The teacher patted the child, impressed. "Well that's something."

The child was the great mathematician Carl Friedrich Gauss. The teacher, although more interested in love than math, took notice of Gauss's talents and worked actively with the university to nurture him.

12.1 Some notes on notation

Definition 12.1. A sequence is an ordered list of numbers. A series is the sum of that list.

This chapter will also be the first time we'll use the summation notation. Why use it instead of just some terms, ... and more terms? Here is a quote from Calculus II for Dummies.

Mathematicians just love sigma notation for two reasons. First, it provides a convenient way to express a long or even infinite series. But even more important, it looks really cool and scary, which frightens non-mathematicians into revering mathematicians and paying them more money.

Definition 12.2. In the summation notation or sigma notation:

$$\sum_{a=1}^b f(a) = f(1) + f(2) + \cdots + f(b-1) + f(b)$$

By the definition itself, we can state two very useful properties of sigma notation.

Theorem 12.3. *Distributively*

$$\sum k f(x) = f \sum k(x)$$

Commutativity

$$\sum [f(x) + g(x)] = \sum f(x) + \sum g(x)$$

And that's all I wish to tell you about notation. Let's come back to math now.

12.2 Arithmetic Progression

An arithmetic progression is the more a sequence of numbers such that the difference between any two consecutive terms is constant. This constant is called the common difference of the sequence. 8, 11, 14, 17 is an AP
More formally:

Definition 12.4. The sequence a_1, a_2, \dots, a_n is an arithmetic progression if and only if $a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1}$

Now Lets consider $a_1, a_2, a_3 \dots a_n$ an AP with common difference d .

Theorem 12.5. *The n^{th} theorem of an Arithmetic Sequence is:*

$$a_n = a + (n - 1)d$$

This follows from the definition.

Theorem 12.6. *Number of terms in an AP:*

$$n = \frac{a_n - a_1}{2} + 2$$

This follows from the n^{th} term formula.

Theorem 12.7. *Average of n terms of AP:*

$$\frac{a_1 + a_n}{2}$$

This also follows from the definition of AP.

Theorem 12.8. *Sum of n terms of an AP:*

$$\frac{a_1 + a_n}{2} \cdot n$$

$$= \frac{a_1 + a_1 + (n - 1)d}{2} \cdot n$$

$$= \frac{2a_1 + (n - 1)d}{2} \cdot n$$

This follows from the definition of average.

With these trivial observations in your toolkit, you can now make some remarkable observations.

Let's consider the AP $1, 2, 3, \dots, n$ with $a_1 = 1$ and $d = 1$. Then

Theorem 12.9.

$$\sum_{k=1}^n k = \frac{n + 1}{2} \cdot n = \frac{n(n + 1)}{2}$$




Figure 12.0. A proof without words for triangle numbers

The numbers which can be written in the form of $\frac{n(n+1)}{2}$ are called triangular numbers. This is because a given number of balls, we can arrange them into a perfect triangle with a base of n .

The theorem above can also be shown without words. You can also realize that this is exactly what Gauss did. Now let's consider the AP $2, 4, 8, 16 \dots 2n$

Theorem 12.10.

$$\sum_{k=1}^n 2k = 2 \sum_{k=1}^n k = 2 * \frac{n(n+1)}{2} = n(n+1)$$

And finally, we'll consider the AP $1, 3, 5, 7 \dots 2n - 1$. Its sum is quite beautiful:

Theorem 12.11.

$$\sum_{k=1}^n 2k - 1 = \sum_{k=1}^n 2k - \sum_{k=1}^n 1 = n(n+1) - n = n(n+1-1) = n^2$$

This can also be basically proven without words as:

Now using whatever we just learnt, let's obliterate some questions.

Example 12.12. (AIME 2012) The terms of an arithmetic sequence add to 715. The first term of the sequence is increased by 1, the second term is increased by 3, the third term is increased by 5, and in general, the k th term is increased by the k th odd positive integer. The terms of the new sequence add to 836. Find the sum of the first, last, and middle terms of the original sequence.

Solution. Let's first realize that the question is asking the sum of first, last and the middle term of the AP. We also know that the middle term is the average of first and last term. So basically, we are looking for three times the




Figure 12.0. Counting the number of squares in the alternating black and white gives us the proof

middle term.

Now let's find it. Let the sequence of $a_1, a_2, a_3 \dots a_n$:

$$\sum_{i=1}^n a_i = 715 \text{ and adding the odd numbers}$$

$$\sum_{i=1}^n a_i + 2n - 1 = 835$$

$$\iff \sum_{i=1}^n a_i + \sum_{i=1}^n 2n - 1 = 835$$

$$\iff 715 + n^2 = 835$$

$$\iff n = 11$$

Hence, we have 11 terms in the sequence. As the sum is 715, the middle term(or the average) is 65. Hence, our answer is $65 * 3 = 195$

□

Example 12.13. (AIME 2005, edited)For each positive integer k , let S_k denote the increasing arithmetic sequence of integers whose first term is 1 and whose common difference is k . For example, S_3 is the sequence 1, 4, 7, 10, For how many values of k does S_k contain the term 2023?

12.3 Geometric Progression

A lot of math jokes begin with: An infinite amount of mathematicians walk into a bar.

This can have one continuation for every mathematician in the bar. The most classic one is: The first one asks for a beer. The second one asks for half a beer. The third one asks for a quarter of a beer. The fourth one asks for an eighth of a beer... . The bartender pours two beers, saying "You guys ought to know your limits".

Let's try to understand what just happened. A geometric progression, is a

sequence of numbers such that the ratio between any two consecutive terms is constant. This constant is called the common ratio of the sequence. For example, 1, 2, 4, 8 is a geometric sequence with common ratio 2 and 100, -50, 25, -25/2 is a geometric sequence with common ratio $-1/2$; however, 1, 3, 9, -27 and -3, 1, 5, 9, ... are not geometric sequences, as the ratio between consecutive terms varies. More formally,

Definition 12.14. The sequence a_1, a_2, \dots, a_n is a geometric progression if and only if $a_2/a_1 = a_3/a_2 = \dots = a_n/a_{n-1}$

Let $g_1, g_2, g_3 \dots g_n$ be a GP with common ratio r .

Theorem 12.15. The n^{th} term in a GP is:

$$\begin{aligned} g_n &= g_1 \cdot r^{n-1} \\ g_n &= g_m \cdot r^{n-m} \end{aligned}$$

This follows from the definition.

Theorem 12.16. Sum of first n terms of a GP is:

$$g_1 \cdot \frac{r^n - 1}{r - 1}$$

NOTE: $\frac{r^n - 1}{r - 1}$ may be replaced with $\frac{1 - r^n}{1 - r}$ whenever convenient.

Proof. This is one of the most beautiful proof in all of mathematics. Let $S_n = a_1 + a_2 + \dots + a_n$

This can be re-written as: $S_n = a_1 + a_1r + \dots + a_1r^{n-1}$ Multiplying both sides by r , $S_n r = a_1r + a_1r^2 + \dots + a_1r^n$ Subtracting the original equation from this equation,

$$\begin{aligned} S_n r - S_n &= a_1r^n - a_1 \\ \iff S_n(r - 1) &= a_1(r^n - 1) \\ \iff S_n &= \frac{a_1(r^n - 1)}{r - 1} \end{aligned}$$

□

Quite cool, but here is something even more cool.

Theorem 12.17. Sum of infinite terms of a converging GP:

$$\frac{g_1}{1 - r}$$

NOTE: This only holds for $-1 \geq r \geq 1$, as other series sum keeps increasing to either ∞ or $-\infty$

Proof. The proof is quite similar to above. With the only difference being that $n \rightarrow \infty$. Let $S_n = a_1 + a_2 + \dots$

This can be re-written as: $S_n = a_1 + a_1r + \dots$ Multiplying both sides by r , $S_n r = a_1r + a_1r^2 + \dots$ Subtracting this equation from the original equation,

$$S_n - S_n r = a_1$$

$$\iff S_n(1 - r) = a_1$$

$$\iff S_n = \frac{a_1}{1-r}$$

□

Now we can truly understand the joke at the start of the chapter.

The first mathematician asked for 1 beer, The second for half, the third for a quarter. This is a GP with $a = 1$ and common ratio $r = \frac{1}{2}$. So we can easily add it using the above formula:

$$\frac{1}{1-\frac{1}{2}} = \frac{1}{\frac{1}{2}} = 2$$

Now let's blast some questions:

Example 12.18. (AIME 2002) Two distinct, real, infinite geometric series each have a sum of 1 and have the same second term. The third term of one of the series is $1/8$, and the second term of both series can be written in the form $\frac{\sqrt{m}-n}{p}$, where m , n , and p are positive integers and m is not divisible by the square of any prime. Find $100m + 10n + p$.

Solution. Let the second term of both the series be x . Then the first term of the first series is $8x^2$ and the common ratio $\frac{1}{8x}$. As it's infinite sum is 1,

$$1 = \frac{8x^2}{1-\frac{1}{8x}}$$

$$\iff 64x^3 - 8x + 1 = 0$$

$$\iff (4x-1)(16x^2+4x-1) = 0 \iff x = \frac{1}{4}, \frac{\sqrt{5}-1}{8}, \frac{-\sqrt{5}-1}{8}$$

As x is of the form $\frac{\sqrt{m}-n}{p}$

$$\therefore x = \frac{\sqrt{5}-1}{8}$$

$$\therefore m = 5, n = 1, p = 8$$

Hence, $100m + 10n + p = 518$

□

Example 12.19. The sum of the first 2011 terms of a geometric sequence is 200. The sum of the first 4022 terms is 380. Find the sum of the first 6033 terms.

12.4 Induction and Summation

While many of the summations are quite trivial to derive, some of them cannot be derived. Only proven. These are typically observed and then proved using a technique called induction.

Almost everyone has once had fun arranging dominoes(or jenga bricks) in a row and starting a wave. Push the first domino and it topples the second, the second will topple the third and so forth until all the dominoes are toppled. Now let us change the set of dominoes into an infinite series of propositions: $P_1, P_2, P_3 \dots$. Assume that we can prove that:

(B): that P_1 of the series is true;

(S): If P_k is true, than P_{k+1} is also true.

Then, in fact, we will have proven that all the propositions in the series are true. We just pushed the first domino and proved that every dominoes is close enough that it will topple with the falling of its predecessor.

This is a description of the method of mathematical induction (MMI). Theorem (B) is called base of induction, and theorem (S) is the inductive step.

This is much better understood by actually seeing it being used:

Theorem 12.20.

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

Proof. (B): For $n=1$, $\frac{1*2*(2*1+1)}{6} = 1^2 = 1$. Hence, it is true for $n = 1$

(S): Let $\sum_{k=1}^m k^2 = \frac{m(m+1)(2m+1)}{6}$ be true for some m . Then:

$$\begin{aligned} & \sum_{k=1}^{m+1} k^2 \\ &= (m+1)^2 + \sum_{k=1}^m k^2 \\ &= (m+1)^2 + \frac{m(m+1)(2m+1)}{6} \\ &= \frac{6(m+1)^2 + m(m+1)(2m+1)}{6} \\ &= \frac{(m+1)(2m^2+m+6m+6)}{6} \\ &= \frac{(m+1)(2m^2+7m+6)}{6} \\ &= \frac{(m+1)(2m+3)(m+2)}{6} \\ &= \frac{(m+1)(m+1+1)(2(m+1)+1)}{6} \end{aligned}$$

As now both (B) and (S) have been proved, our theorem is also true. \square

We'll leave one as exercise for you. It's almost the same, but with cube.

Theorem 12.21.

$$\sum_{k=1}^n k^3 = \left(\frac{k(k+1)}{2}\right)^2$$

12.5 Telescoping

Telescoping is a method of writing a summation out in such a way that a lot of it gets canceled. While naturally such series don't occur a lot, they do appear

in human generated numbers. A very common use is in the dividend discount model. You may have heard that the stock prices rise and plummet on the basis of demand and supply. How and why of this is guarded by mathematical algorithms. The Dividend discount model prices a stock based on the amount of money it will pay to the holder in dividends discounted to present value. Let's break that down. A stock is basically a part, a stake in the company you have. Basically, companies want you to hold their stocks. So they pay yearly dividends to the holders. Basically, it's your part of the pie for owning a part of the company. It's value is not decided by a person but an algorithm which based on market patterns predicts what the stock will pay you in dividends over perpetuity(a very long period of time, technically ∞ , however taken as 5-10 year period for the purpose of economic calculations).

However, you might have heard people say that 'inflation is rising'. Your parents or grandparents probably would have said that in their days the things were cheaper. So whatever the stock is expected to pay you in the future, will be affected by inflation. So we adjust it for the same.

If a company pays us the same amount and the rate of inflation is constant, we'll have a simple infinite GP. However, stocks with constant dividends are less liked, hence companies increase the dividends from time to time. This leads to an infinite summation which normally gives us the value of the stock. For example: ITC, Indian Tobacco Company, had a dividend of 5.25 INR in 2012 and it has increased annually by an average 2.85%. The inflation in Indian market has increased by an average of 6.02% in the same time period, ouch!. So based on this data, its price should be $5.25 + 5.25 * (1 + \frac{2.85}{100}) * (1 - \frac{6.02}{100}) + \dots = \sum_{n=1}^{\infty} 5.25 * ((\frac{2.85}{100}) * (1 - \frac{6.02}{100}))^{n-1}$
 $= \sum_{n=1}^{\infty} 5.25 * 0.9683^{(n-1)}$

This is now a summation we can compute. While this is not telescopic, note that the numbers were averages. Exact dividend growth and inflation are different every year which will lead to a telescope, but we'll not demonstrate that here.

Solving this, we get 165.61 as the price which is shockingly close to its price in 2012 mid year when dividends were paid(167).

The actual price is found using estimates for inflation and other data which are useful but messy functions. However, as these numbers are based on human activity or selected by humans, they tend to lead to telescoping.

Telescoping is something that is also a very common question pattern in Olympiads. We basically ask you to compute some strange summation which can be broken down into subtraction of a few functions which happen to cancel out leaving very little computation to be done. Just like a telescope.

Example 12.22 (Motivating Example). (Stanford 2011) Evaluate the sum:

$$\sum_{k=1}^{\infty} \frac{7k+32}{k(k+2)} \cdot \frac{3^k}{4^k}$$

Solution. Decomposing the denominator:

$$\frac{7k+32}{k(k+2)} = \frac{16}{k} - \frac{9}{k+2}$$

$$\text{As } 9 = 16 * \frac{3^2}{4^2}$$

$$\begin{aligned} \therefore \sum_{k=1}^{\infty} \frac{7k+32}{k(k+2)} \cdot \frac{3^k}{4^k} &= \frac{16}{k} * \frac{3k}{4^k} - \frac{9}{k+2} * \frac{3^k}{4^k} \\ &= \frac{16}{k} * \frac{3^k}{4^k} - \frac{16}{k+2} * \frac{3^{k+2}}{4^{k+2}} \end{aligned}$$

We'll only need to solve for $k = 1, 2$ as rest will get canceled. Thus, the answer is: $16 * \frac{3}{4} + 8 * \frac{9}{16} = 12 + \frac{9}{2} = \frac{33}{2}$ \square

In the motivating example, in order to cancel things out, we decomposed the denominator to reach a solution. The thing is that this can be done with every factorizable denominator and is of great value in integral calculus.

A, B, C need to be figured out separately, which you can do in two ways. I'll

Table 12.0. Rational Functions and Their Decompositions

General Form	Decomposition
$\frac{px+q}{(x-a)(x-b)}$	$\frac{A}{x-a} + \frac{B}{x-b}$
$\frac{px+q}{(x-a)(x-a)^2}$	$\frac{A}{x-a} + \frac{B}{(x-a)^2}$
$\frac{px^2+qx+r}{(x-a)(x-b)(x-c)}$	$\frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$
$\frac{px^2+qx+r}{(x-a)^2(x-c)}$	$\frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{C}{x-b}$
$\frac{px^2+qx+r}{(x-a)(x^2+bx+c)}$	$\frac{A}{x-a} + \frac{Bx+C}{x^2+bx+c}$

demonstrate them:

Example 12.23.

$$\sum_{n=4}^{\infty} \frac{n}{n^3 - 6n^2 + 11n - 6}$$

Solution. We can factorize $n^3 - 6n^2 + 11n - 6$ as $(n-1)(n-2)(n-3)$.

Therefore we can decompose $\frac{n}{n^3 - 6n^2 + 11n - 6} = \frac{A}{(n-1)} + \frac{B}{(n-2)} + \frac{C}{(n-3)}$

$$\therefore A(n-2)(n-3) + B(n-1)(n-3) + C(n-1)(n-2) = n$$

Here the paths diverge. We'll start with Method I, which is more widely published, but much slower.

$$\iff A(n^2 - 5n + 6) + B(n^2 - 4n + 3) + C(n^2 - 3n + 2) = n$$

$$\iff (A + B + C)n^2 - (5A + 4B + 3C)n + (6A + 3B + 2C) = 0n^2 + n + 0$$

$$\iff A + B + C = 0; 5A + 4B + 3C = -1; 6A + 3B + 2C = 0$$

$\iff 2A+B = -1; 4A+B = 0$ This follows by repeated subtraction of the first equation

$$\iff A = \frac{1}{2}$$

$$\therefore A = \frac{1}{2}; B = -2; C = \frac{3}{2}$$

If we move back to the fork in the road, we can now use method II, which is less commonly known but much faster.

As we have, $A(n-2)(n-3) + B(n-1)(n-3) + C(n-1)(n-2) = n$, If

$n-1=0 \iff n=1$ then: $A(1-2)(1-3)+B(1-1)(1-3)+C(1-1)(1-2)=1$

$$\iff A(-1)(-2) = 1$$

$$\iff A = \frac{1}{2}$$

We can now take $n-2=0 \iff n=2$ and then $n-3=0 \iff n=3$ giving us $B=-2$ and $C=\frac{3}{2}$. It's so simple that you can probably do it in your head.

Now let's destroy the question:

$$\begin{aligned} & \sum_{n=4}^{\infty} \frac{n}{n^3 - 6n^2 + 11n - 6} \\ &= \sum_{n=4}^{\infty} \frac{1}{2(n-1)} - \frac{2}{n-2} + \frac{3}{2(n-3)} \\ &= \sum_{n=4}^{\infty} \frac{1}{2(n-1)} - \frac{4}{2(n-2)} + \frac{3}{2(n-3)} \\ &= \frac{1}{2} \sum_{n=4}^{\infty} \frac{1}{(n-1)} - \frac{4}{(n-2)} + \frac{3}{(n-3)} \\ &= \frac{1}{2} \sum_{n=4}^{\infty} \frac{1}{(n-1)} - \frac{4}{(n-2)} + \frac{3}{(n-3)} \end{aligned}$$

To find the summation notice that the diagonals of the table below are summing to zero. I've crossed one out to make it more clear. The rest terms also get canceled other than the three values in the right corner, but they have

$\frac{1}{(n-1)}$	$\frac{-4}{(n-2)}$	$\frac{3}{(n-3)}$
$\frac{1}{3}$	$\frac{-4}{2}$	$\frac{3}{1}$
$\frac{1}{4}$	$\frac{-4}{3}$	$\frac{3}{2}$
$\frac{1}{5}$	$\frac{-4}{4}$	$\frac{3}{3}$
$\frac{1}{6}$	$\frac{-4}{5}$	$\frac{3}{4}$
\vdots	\vdots	\vdots

not been crossed to make sure the pattern is evident.

Hence we can now say that:

$$\begin{aligned} & \frac{1}{2} \sum_{n=4}^{\infty} \frac{1}{(n-1)} - \frac{4}{(n-2)} + \frac{3}{(n-3)} \\ &= \frac{1}{2} \cdot \left(\frac{-4}{2} + 3 + \frac{3}{2} \right) \end{aligned}$$

$$= \frac{1}{2} \cdot \frac{5}{2}$$

$$= \frac{5}{4}$$

□

That was quite beautiful! Let's do another question.

Example 12.24. (USAMTS 1999, edited) $\sqrt{1 + \frac{1}{1^2} + \frac{1}{2^2}} + \sqrt{1 + \frac{1}{2^2} + \frac{1}{3^2}} + \sqrt{1 + \frac{1}{3^2} + \frac{1}{4^2}} + \cdots + \sqrt{1 + \frac{1}{2023^2} + \frac{1}{2024^2}}$

Solution. The question is basically:

$$\sum_{n=1}^{2023} \sqrt{1 + \frac{1}{n^2} + \frac{1}{(n+1)^2}}$$

The square root seems to be the most irritating part of the question. Let's see if we can somehow get rid. What will happen if we try to simplify the stuff inside the radical?

$$\begin{aligned} & 1 + \frac{1}{n^2} + \frac{1}{(n+1)^2} \\ &= \frac{(n(n+1))^2 + (n+1)^2 + n^2}{(n(n+1))^2} \\ &= \frac{n^4 + 2n^3 + 3n^2 + 2n + 1}{(n(n+1))^2} \\ &= \frac{n^4 + n^3 + n^2 + n^2 + n + n^2 + n + 1}{(n(n+1))^2} \\ &= \frac{n^2(n^2 + n + 1) + n(n^2 + n + 1) + 1(n^2 + n + 1)}{(n(n+1))^2} \\ &= \frac{n^2 + n + 1}{n(n+1)}^2 \end{aligned}$$

Voila! The square root has been defeated. Now all is left is to solve whatever remains have been left.

$$\begin{aligned} & \sum_{n=1}^{2023} \sqrt{1 + \frac{1}{n^2} + \frac{1}{(n+1)^2}} \\ &= \sum_{n=1}^{2023} \sqrt{\frac{n^2 + n + 1}{n(n+1)}} \\ &= \sum_{n=1}^{2023} \frac{n^2 + n + 1}{n(n+1)} \\ &= \sum_{n=1}^{2023} \frac{n(n+1) + 1}{n(n+1)} \\ &= \sum_{n=1}^{2023} 1 + \frac{1}{n(n+1)} \\ &= \sum_{n=1}^{2023} 1 + \sum_{n=1}^{2023} \frac{1}{n(n+1)} \\ &= 2023 + \sum_{n=1}^{2023} \frac{1}{n(n+1)} \\ &= 2023 + \sum_{n=1}^{2023} \frac{1}{n} - \frac{1}{n+1} \\ &= 2023 + \frac{1}{1} - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \frac{1}{3} - \cdots - \frac{1}{2022} + \frac{1}{2023} - \frac{1}{2024} \\ &= 2023 + 1 - \frac{1}{2024} \\ &= 2023 + \frac{2023}{2024} \end{aligned}$$

□




Figure 12.0. The rabbit graph, source: Murderous Math: The Key to Universe

12.6 Recurrence Series

Let's for our final story, go to 12th century. We are in Pisa. The construction for a white bell tower has just began. The church despite scientific advice, has decided to go for it in the land available. The subsoil seems a bit unsteady, but the architect is confident.

A mathematician has bought a pair of young male and female rabbits from the animal seller as his Christmas present to himself. He leaves them in his backyard. In January, they are all grown up. By February, they give birth to another pair which happens to be off opposite genders.

In March, the original pair gives birth to two more baby rabbits and the first children are of reproductive age. In April, we get two new pairs of rabbits and the second born are of reproductive age. We can notice that the number of pairs of rabbits in a given month follow the sequence are:

1, 1, 2, 3, 5, 8, ...

The pattern is evident. The mathematician's name was Fibonacci and this sequence is called the Fibonacci sequence after him.

What happened of the rabbits? They were plentiful in Pisa(couldn't override the town as they neither live or breed for eternity) and most either lived happily chewing on grass near the white tower. It is believed that their descendants are still there, chewing on the grass, unaware about their contribution. But back to math, a series which depends on its previous terms for its further terms is called a recursive series. We studied about them from a combinatorial perspective in chapter 7. Most of it still applies, and I recommend that you go through it if you haven't already.

Exercises

- (1) (AIME) Find the eighth term of the sequence 1440, 1716, 1848, ... whose terms are formed by multiplying the corresponding terms of two arithmetic sequences.
- (2) (AIME) Two geometric sequences a_1, a_2, a_3, \dots and b_1, b_2, b_3, \dots have the same common ratio, with $a_1 = 27$, $b_1 = 99$, and $a_{15} = b_{11}$. Find a_9 .
- (3) (AIME) For $-1 < r < 1$, let $S(r)$ denote the sum of the geometric series

$$12 + 12r + 12r^2 + 12r^3 + \dots$$

Let a between -1 and 1 satisfy $S(a)S(-a) = 2016$. Find $S(a) + S(-a)$

- (4) (AIME) An infinite geometric series has sum 2005. A new series, obtained by squaring each term of the original series, has 10 times the sum of the original series. The common ratio of the original series is $\frac{m}{n}$ where m and n are relatively prime integers. Find $m + n$.
- (5) (AIME) Call a 3-digit number geometric if it has 3 distinct digits which, when read from left to right, form a geometric sequence. Find the difference between the largest and smallest geometric numbers.
- (6) (AIME) The sum of an infinite geometric series is a positive number S , and the second term in the series is 1. What is the smallest possible value of S ?
- (7) (AIME) In an increasing sequence of four positive integers, the first three terms form an arithmetic progression, the last three terms form a geometric progression, and the first and fourth terms differ by 30. Find the sum of the four terms.
- (8) (AIME) A sequence of positive integers with $a_1 = 1$ and $a_9 + a_{10} = 646$ is formed so that the first three terms are in geometric progression, the second, third, and fourth terms are in arithmetic progression, and, in general, for all $n \geq 1$, the terms $a_{2n-1}, a_{2n}, a_{2n+1}$ are in geometric progression, and the terms a_{2n}, a_{2n+1} , and a_{2n+2} are in arithmetic progression. Let a_n be the greatest term in this sequence that is less than 1000. Find $n + a_n$.

- (9) (AIME) Consider the sequence defined by $a_k = \frac{1}{k^2 + k}$ for $k \geq 1$. Given that $a_m + a_{m+1} + \dots + a_{n-1} = \frac{1}{29}$, for positive integers m and n with $m < n$, find $m + n$.

- (10) (AIME) Given that

$$x_1 = 211,$$

$$x_2 = 375,$$

$$x_3 = 420,$$

$$x_4 = 523, \text{ and}$$

$$x_n = x_{n-1} - x_{n-2} + x_{n-3} - x_{n-4} \text{ when } n \geq 5,$$

find the value of $x_{531} + x_{753} + x_{975}$.

- (11) (AIME) The sequence $\{a_n\}$ is defined by

$$a_0 = 1, a_1 = 1, \text{ and } a_n = a_{n-1} + \frac{a_{n-1}^2}{a_{n-2}} \text{ for } n \geq 2.$$

The sequence $\{b_n\}$ is defined by

$$b_0 = 1, b_1 = 3, \text{ and } b_n = b_{n-1} + \frac{b_{n-1}^2}{b_{n-2}} \text{ for } n \geq 2.$$

Find $\frac{b_{32}}{a_{32}}$.

- (12) (AIME 2004) Consider the sequence defined by $a_k = \frac{1}{k^2 + k}$ for $k \geq 1$.

Given that $a_m + a_{m+1} + \cdots + a_{n-1} = \frac{1}{29}$, for positive integers m and n with $m < n$, find $m + n$.

- (13) (AIME 1989) If the integer k is added to each of the numbers 36, 300, and 596 one obtains the squares of three consecutive terms of an arithmetic series. Find k .

- (14) Compute this sum:

$$\left(\frac{1}{2^2} + \frac{1}{3^2} + \dots\right) + \left(\frac{1}{2^3} + \frac{1}{3^3} + \dots\right) + \dots$$

- (15) (Putnam 2013,edited) For positive integers n , let the numbers $c(n)$ be determined by the rules: $c(1) = 1$, $c(2n) = c(n)$, and $c(2n+1) = (-1)^n c(n)$. Find the value of:

$$\sum_{n=1}^{2023} c(n)c(n+2)$$

- (16) If a, b, c form an arithmetic progression, and $a = x^2 + xy + y^2$

$$b = x^2 + xz + z^2$$

$c = y^2 + yz + z^2$ where $x + y + z = 0$, prove that x, y , and z also form an arithmetic progression.

- (17) (IOQM 2023) The sequence a_n with $n \geq 0$ is defined by $a_0 = 1, a_1 = -4$ and $a_{n+2} = -4a_{n+1} - 7a_n$ for $n \geq 0$. Find $a_{50}^2 - a_{49}a_{51}$

(18)

- (19) (AMC 12) [5★] Three balls are randomly and independently tossed into bins numbered with the positive integers so that for each ball, the probability that it is tossed into bin i is 2^{-i} for $i = 1, 2, 3, \dots$. More than one ball is allowed in each bin. The probability that the balls end up evenly spaced in distinct bins is p/q , where p and q are relatively prime positive integers. (For example, the balls are evenly spaced if they are tossed into bins 3, 17, and 10.) What is $p + q$?

Part 5

The Red Pill

13

Calculus I: Limits, Continuity and Derivative

Calculus, the mathematics of change, and change, mysterious. Some things grow imperceptibly... others zoom... hair grows slowly and suddenly cut... temperatures rise and fall... smoke curls through the air... planets wheel through space... and time, time never stops... - Cartoon Guide to Calculus

Imagine you're sitting in a car with your family, driving along a winding mountain road. The scenery outside is breathtaking, with lush green forests, towering cliffs, and a deep blue sky. As you cruise along, you can't help but notice how the world around you is constantly transforming.

Now, let's say you're curious about how fast things are changing. You're particularly interested in how quickly your car is ascending the mountain. You pull out your trusty stopwatch and start measuring the time it takes to climb each mile of the road. You also jot down your car's speed at each mile marker. As you collect this data, something remarkable happens. You realize that the speed of your car is not constant—it varies with each mile. Sometimes you're going uphill at a slow pace, and other times you're speeding down a hill, faster than you thought possible.

What's even more intriguing is that you notice a pattern: the steeper the incline, the slower your car seems to go. Conversely, when the road levels out or slopes downward, your car speeds up. This variation in speed as you move along the road is what we call...

13.1 The Rate of Change

In a still shot of a baseball game, the ball is clear, not moving at the instant of the photo. A good camera can take many such photos each with the ball clear and stopped.

So if in any instant the ball is stopped, how is it truly moving? This is



Figure 13.0. Is the ball moving? How do you know?

courtesy, Champaign News-Gazette and photographer Robin Scholtz, 2003

known as Zeno's paradox. The answer is rather simple. The ball, like every other moving object, has an unseen invisible quantity called velocity which is dictating its speed and direction.

While we can define velocity as rate of change of distance(displacement is more physically accurate as it also encompasses the direction), what about changing velocity?

As a baseball is thrown in the air, its velocity decreases as it slows down before it stops at its highest point. It then comes down, speeding up along the way. However, here is a small issue. The speedometer of the car was showing you a speed all the time without ever actually having memory of what time it is and how much distance we had travelled.

How did that work? The speedometer was showing us the speed in the last 0.01 sec or even smaller. That was the cars instantaneous speed.

If this feels sort of strange, it should. This is all kind of paradoxical, as we are trying to look for the speed of something in a snapshot, which is not possible as we cannot divide by zero.

If the $f(x)$ is our distance function and time is along the x axis, we can rep-




Figure 13.0. The graph at heart of calculus

resent velocity between time x_0 and $x_0 + \Delta x$ as the slope of the line between $f(x_0) = P$ and $f(x_0 + \Delta x) = Q$.

As $\Delta x \rightarrow 0$, $f(x_0 + \Delta x) = Q \rightarrow P$, which makes the secant between P and Q a tangent at P of which we can find the slope off.

This is essentially what your speedometer does, it measures the slope of the the tangent of your distance graph.

This is essentially what calculus is all about.

13.2 Some functions

We will end up using some functions again and again in calculus. These functions using in combination with each other make up most of the things we see in physics, which is the major use of calculus. I will take this moment to remind you that a function is a relation from a domain to co-domain.

13.2.1 Constant Function. A simple function which returns the same value even in plugging out different values. Domain is $x \in \mathbb{R}$, while the co-domain is a single Real number.




Figure 13.0. The constant function

13.2.2 Power Function. These are the functions with formulas x, x^2, x^3, \dots, x^n , where n is a positive integer. When n is even, these functions all have bowl-shaped graphs as $(-x)^{2n} = x^{2n}$. If n is odd, then $(-x)^{2n-1} = -(x^{2n-1})$, causing the graphs bend downward on the left.




Figure 13.0. Power Function

13.2.3 Polynomials. Polynomials are basically made by multiplying powers by constants and adding them. You already know from algebra that a polynomial with degree n has at most n roots or n points where the graph cuts zero. We can also notice that every polynomial either goes to ∞ or $-\infty$, so every time it cuts at zero more than once, it will also have a turning. So basically, it has at most $n - 1$ turnings. The turnings will become use full in a minute.

13.2.4 Negative powers. We can also write negative powers using the fact $x^{-n} = \frac{1}{x^n}$. Their graph also varies according to the parity of n .

13.2.5 Fractional Powers. We can also plot $x^{\frac{1}{n}} = \sqrt[n]{x}$. Also without any surprise, you might have already understood that it also depends on parity.

13.2.6 Exponential functions. Power functions: We get large pretty quickly
Exponential functions: Hold my cup...

Exponential functions are defined as $f(x) = a^x$ where a is constant. Physicists like using e as the base where e is Euler's number and is equal to $2.718\dots$

Like π it is irrational, however unlike π , its definition is not geometric. It was arrived upon by Jacob Bernoulli while pondering the given question




Figure 13.0. Negative Powers




Figure 13.0. Fractional Powers

Example 13.1. An account starts with 1 dollar and pays 100 percent interest per year. If the interest is credited once, at the end of the year, the value of the account at year-end will be 2 dollars. What happens if the interest is computed and credited more frequently during the year?

Using our knowledge of sequence and series, we could compute that if the interest was compounded n times a year, we would get $(1 + \frac{1}{n})^n$. If we keep making n larger and larger, till it approaches ∞ , we'll be left with e . How exactly? We'll talk about that in just a minute

We can also say that $e^r = a$ will have a unique solution for r as long as $a \neq 0$. This is used so often that we have a notation for it: $\ln(a) = r$ where \ln refers to the natural logarithm. Its properties are discussed in greater detail in the logarithm's chapter. However, using only the fact that it exists we can say $f(x) = e^{rx}$.

This graph grows exponentially for $r > 0$, is constant for $r = 0$ and decays exponentially for $r < 0$.

13.2.7 Trigonometric Functions. If you are unfamiliar with trigonometry, I recommend reading through that chapter before continuing this .

Here we have a circle of radius 1 and make two perpendicular lines through the center. A radius is made which goes from the center to some point on the circumference P making an angle θ with the horizontal, which is measured




Figure 13.0. Meet the unit circle

in radians. Radians are another way to measure angles which is the norm in physics. It basically is the arc length of the arc subtended by your angle divided by the radius. Physicist's like it better as it doesn't use an abstract number 360 as the reference for the angles instead uses an physical quantity. But back to the diagram, The perpendicular distance of this from the y axis is called $\cos \theta$ and the perpendicular distance from the x axis is called $\sin(\theta)$. We define $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$

A trigonometric function refers to $f(x) = \sin(x)$; $f(x) = \cos(x)$; $f(x) = \tan(x)$ or a combination of them with coefficients where the domain is $x \in \mathbb{R}$ and the co-domain for $\sin(x)$ and $\cos(x)$ is $\{-1, 1\}$. The co-domain is all real numbers for $\tan(x)$.




Figure 13.0. Trigonometric Functions

13.2.8 Composing Functions. A composing function is defined as $f(x) = h(g(x))$ basically, a combination of two functions. NOTE: While sometimes, $h(g(x)) = g(h(x))$ this is often untrue. We should not interchange the functions unless we are absolutely sure we can.

13.2.9 Inverting Functions. If composing function $f(x) = h(g(x)) = C$ where C is a constant then $h(x)$ is the inverse of $g(x)$ and vice versa. The inverse of a function $k(x)$ is sometimes denoted as $k^{-1}(x)$. You may feel that an inverse composition may be switched around but consider this: $g(x) = x^2, h(x) = \sqrt{x}$ then while $f(x) = g(h(x))$ has a domain of all positive real numbers, $f(x) = h(g(x))$ has a domain of all real numbers; making them fundamentally different.

13.3 Inverse Trigonometric functions

This difference is even more profound with trigonometric functions (whose inverse are written as $\sin^{-1}, \cos^{-1}, \tan^{-1}$ and have domains of $(-1, 1); (-1, 1); \theta \in \mathbb{R}$ respectively.)

We need to note that while $\sin(x)$ keeps on going up and down, it is strictly increasing in the range $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$. Fun fact: A lot of people refer to \sin^{-1} as arcsin. This is because it basically corresponds to the length of the arc the radii is subtending given the perpendicular distance from the vertical.

I expect you to find the co-domain of \arccos by yourself now.




Figure 13.0. The return of the unit circle

Example 13.2. Find the co-domain of \arccos

Let's talk about \arctan . The co-domain of this is pleasantly surprising as it has a massive domain.




Figure 13.0. The Unit Circle Awakens

13.4 Derivative

As derivative is the slope of a function $f(x)$ at a particular point, we can also plot the derivative as a function $f'(x)$ as we move that particular point.

Theorem 13.3 (Differentiation formula).

$$f'(x) = \frac{f(x+h) - f(x)}{h}, h \rightarrow 0$$

This is just formalization of the definition of derivative.
Using this definition, we can quite easily figure some more things.

Theorem 13.4 (Two truths of differentiation). $(f(x) + g(x))' = f'(x) + g'(x)$
 $\therefore (Cf(x))' = Cf'(x)$ where C is a constant

While this seems both trivial and extraordinary, here is the proof

$$\begin{aligned} & \text{Proof. } (f(x) + g(x))' \\ &= \frac{f(x+h) + g(x+h) - f(x) - g(x)}{h} \\ &= \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \\ &= f'(x) + g'(x) \end{aligned}$$

□

We will also prove some standard derivatives which you should remember(just the identity, the proof is obvious).

Theorem 13.5 (Power rule of differentiation). If $f(x) = x^n$,
 $f'(x) = nx^{n-1}$

Proof. $f'(x) = \frac{x+h^n-x^n}{h}$
 $= \frac{x^n+nx^{n-1}h+\dots+nxh^{n-1}+h^n-x^n}{h} = nx^{n-1} + \binom{n}{2}x^{n-2}h + \dots + nxh^{n-2} + h^{n-1}$ Using $h \rightarrow 0$
 $= nx^{n-1}$ \square

With this much, we are now qualified enough to find the derivative of all polynomials.

Theorem 13.6. If $f(x) = \sin(x)$, then:

$$f'(x) = \cos(x)$$

If $f(x) = \cos(x)$, then:

$$f'(x) = -\sin(x)$$

Proof. $f'(x) = \frac{\sin(x+h)-\sin(x)}{h}$ By the trigonometric property for $\sin(\alpha + \beta)$
 $= \frac{\sin(x)\cos(h)+\cos(x)\sin(h)-\sin(x)}{h}$
 $= \cos(x)\frac{\sin(h)}{h} + \sin(x)\frac{\cos(h)-1}{h}$ We can notice that $\sin(h) = h$ for small values of h as the perpendicular distance to horizontal and the arc length are almost equal.

We also need to notice that $\cos(h)-1$ is almost equal to 0 as the perpendicular distance to vertical is almost equal to the radius of the unit circle which is 1.
 $\therefore \cos(x)\frac{\sin(h)}{h} + \sin(x)\frac{\cos(h)-1}{h}$
 $= \cos(x)\frac{h}{h} + \sin(x)\frac{0}{h} = \cos(x)$ \square

I'll leave the full proof for the derivative of $\cos(x)$ to you but the simple, non trigonometric, proof without words is in noticing that $\cos(x)$ is exactly like $\sin(x)$ shifted left by $\frac{\pi}{2}$.

We'll come to $\tan(x)$ in a minute.




Figure 13.0. A proof without words

Theorem 13.7. If $f(x) = e^x$:

$$f'(x) = e^x$$

$$\therefore f(x) = a^x = f'(x) \iff a = e.$$

This is what makes e so special.

Proof. Let $f(x) = a^x = f'(x)$ $f'(x) = \frac{a^{x+h}-a^x}{h} = a^x$

$$\iff a^x \frac{a^h - 1}{h} = a^x$$

$$\iff \frac{a^h - 1}{h} = 1$$

$\iff a^h = 1 + h$ Remember the definition of e ? This is where that definition becomes significant (we'll prove it, wait for it). As $e = (1 + \frac{1}{n})^n, n \rightarrow \infty$, we can set $h \rightarrow \frac{1}{n} \rightarrow 0$

$$\therefore e = (1 + h)^{\frac{1}{h}}$$

$$\therefore e^h = 1 + h$$

Which implies, $a = e$

Hence proved. \square

One of my most favorite proofs.

13.5 Some more notation

Calculus has had a dark history. Newton and Leibniz contested on who had invented it. The thing is that they both assumed that the other had copied it. Historical records show that the discovery was 1. independent, 2. already discovered approximately 2000 years ago by the Greek, Chinese, Arabic and Indian mathematicians 3. already in use in different forms by modern mathematicians and physicists like Galileo and Kepler

So in classic European fashion, they took something that was already there, gave it a name and then argued over who 'discovered' it.

What they did do was give us some sort of a better notation system. Till now we were using the Newton's notation where derivatives are shown as $f'(x)$ but we'll use Leibniz notation a bit more in the future as it simplifies a lot of things which Newton would complicate.

In Leibniz notation, the derivative is taken of an equation rather than of a function. For example:

$$y = x^2$$

$\therefore \frac{dy}{dx} = 2x$ Here, $\frac{dy}{dx}$ implies the derivative. The d here is a version of Δ , which is what we use to show big change, but d shows microscopic changes. Basically, this is just the original meaning of differentiation.

We also need to notice that we just multiplied the entire equation by $\frac{d}{dx}$ and solved the derivatives and got the result.

This makes the two truths feel like an elementary fact.

13.6 Product and Quotient rules

Theorem 13.8 (Product rule). $(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$

Proof. We'll take $f(x+h) = f(x) + \Delta(f)$ where $h, \Delta(f) \rightarrow 0$

$$\begin{aligned} \therefore (f(x)g(x))' &= \frac{f(x+h)*g(x+h)-f(x)g(x)}{h} \\ &= \frac{(f(x)+\Delta(f))*(g(x)+\Delta(g))-f(x)g(x)}{h} \\ &= \frac{f(x)g(x)+g(x)\Delta(f)+f(x)\Delta(g)+\Delta(f)\Delta(g)-f(x)g(x)}{h} \\ &= g(x)\frac{\Delta(f)}{h} + f(x)\frac{\Delta(g)}{h} + \frac{\Delta(f)\Delta(g)}{h} \\ \text{Using the fact that } \Delta(f), \Delta(g) &\rightarrow 0 \text{ and } f(x+h) - f(x) = \Delta(f) \text{ and } g(x+h) - g(x) = \Delta(g) \\ g(x)\frac{\Delta(f)}{h} + f(x)\frac{\Delta(g)}{h} + \frac{\Delta(f)\Delta(g)}{h} &= g(x)\frac{f(x+h)-f(x)}{h} + f(x)\frac{g(x+h)-g(x)}{h} \\ &= g(x)f'(x) + g'(x)f(x) \end{aligned}$$

□

Also I'd like you to note that, by the same proof, if we want to differentiate more than two functions we can go for: $(fg h \dots)' = f'gh \dots + fg'h \dots + fgh' \dots$

Theorem 13.9 (Reciprocal rule). $(\frac{1}{f(x)})' = \frac{-f'(x)}{(f(x))^2}$

$$\begin{aligned} \text{Proof. } (\frac{1}{f(x)})' &= \frac{\frac{1}{f(x+h)} - \frac{1}{f(x)}}{h} \\ &= \frac{\frac{f(x) - f(x+h)}{f(x+h)f(x)}}{h} \\ &= \frac{-f'(x)}{f(x+h)(f(x))} \text{ Using } h \rightarrow 0 \\ &= \frac{-f'(x)}{(f(x))^2} \end{aligned}$$

□

Combining the product and reciprocal rule gives us:

Theorem 13.10 (Quotient Rule). $(\frac{f(x)}{g(x)})' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$

We are now powerful enough to solve a few more differential questions:

Example 13.11. For $f(x) = \tan(x)$, compute:
 $f'(x)$

Solution. We know that the formula for $\tan(\alpha + \beta)$ is quite messy. However, we also know that $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$ as well as the derivatives for $\sin(\theta)$ and $\cos(\theta)$, this seems like an use the quotient rule.

$$\begin{aligned} \therefore f'(x) &= \frac{(\sin(x))'\cos(x) - (\cos(x))'\sin(x)}{\cos^2(x)} \\ &= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} \end{aligned}$$

We know that $\sin^2(x) + \cos^2(x) = 1$ using the unit circle,

$$\begin{aligned} & \therefore \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} \\ &= \frac{1}{\cos^2(x)} \\ &= \sec^2(x) \end{aligned}$$

The final step is by the definition that $\frac{1}{\sin} = \csc$; $\frac{1}{\cos} = \sec$; $\frac{1}{\tan} = \cot$ \square

I will leave it to you as exercise to differentiate the rest of the trigonometric functions by yourself.

Theorem 13.12. $x^{-n} = -nx^{(-n+1)}$

While, this is exactly what the power rule states. It's proof is different from that of the power rule as we can't use the binomial expansion here. However, it's trivial as we are just using the reciprocal rule and the power rule together. I expect that you'll take it upon yourself to prove it once for practice.

13.7 Chain rule

While we can differentiate a lot of things, we still fail to differentiate things like e^{2x} or $\sin(\cos(x))$.

Here is where the chain rule comes.

Theorem 13.13. $f(g(x))' = g'(x)f'(g(x))$

This probably looks worse than it actually is. It's proof will appear in a minute. But till then let's do a question to understand this better.

Example 13.14. $\frac{d}{dx}e^{2x}$

Solution. The chain rule basically states that for the derivative of $f(g(x))$ we will first take $f'(x)$ and plug $g(x)$ in place of x . We'll then multiply that by $g'(x)$

In this case, $f(x) = e^x \implies f'(x) = e^x$ and $g(x) = 2x \implies g'(x) = 2$, therefore:

$$(e^{2x})' = 2e^{2x}$$
 \square

Example 13.15. $\frac{d}{d\theta}\sin(\cos(\theta))$

I believe this one will be a cake walk for you to do.

Now let's talk about inverses. The chain rule can solve for the inverse given we know the derivative of the original function.

Theorem 13.16 (Inverse Rule). $(f^{-1}(x))' = \frac{1}{f'(f^{-1}(x))}$

$$\begin{aligned} \text{Proof. } & x = f(f^{-1}(x)) \\ \iff & \frac{dx}{dx} = \frac{d}{dx} f(f^{-1}(x)) \\ \iff & 1 = f'^{-1}(x) \cdot f'(f^{-1}(x)) \\ \iff & (f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))} \end{aligned}$$

□

If you use this formula on $x^{\frac{1}{n}}$, you'll get what we get by the power rule. If you think that's cool, hold my cup:

Theorem 13.17. $\frac{d}{dx} \ln(x) = \frac{1}{x}$

This looks wild. How did this even happen?

Proof. As $f(x) = \ln(x)$ is the inverse of $g(x) = e^x \implies g'(x) = e^x$, we can use the inverse rule.

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}} = \frac{1}{x}$$

□

This obviously doesn't happen with other functio...

Theorem 13.18. $\frac{d}{dx} \arcsin(x) = \frac{1}{\sqrt{1-x^2}}$

Proof. As $f(x) = \arcsin(x)$ is the inverse of $g(x) = \sin(x) \implies g'(x) = \cos(x)$, we can use the inverse rule.

$$\frac{d}{dx} \arcsin(x) = \frac{1}{\cos(\arcsin(x))}$$

As $\sin^2(x) + \cos^2(x) = 1$, therefore $\cos(x) = \sqrt{1 - \sin^2(x)} = \sqrt{1 - x^2}$

Plugging in $x = \arcsin(x)$ we get: $\cos(x) = \sqrt{1 - x^2}$, therefore:

$$\frac{1}{\cos(\arcsin(x))} = \frac{1}{\sqrt{1-x^2}}$$

□

So maybe this the only trig funct...

Theorem 13.19. $\frac{d}{dx} \arctan(x) = \frac{1}{1+x^2}$

Proof. As $f(x) = \arctan(x)$ is the inverse of $g(x) = \tan(x) \implies g'(x) = \sec^2(x)$, we can use the inverse rule.

$$\frac{d}{dx} \arctan(x) = \frac{1}{\sec^2(\arctan(x))}$$

As $\sec^2(x) - \tan^2(x) = 1$ (Just write both in terms of sin and cos and it will become obvious), therefore $\sec^2(x) = \tan^2(x) + 1$

Plugging in $x = \arctan(x)$ we get: $\sec^2(\arctan(x)) = \tan^2(\arctan(x)) + 1 = x^2 + 1$, therefore:

$$\frac{1}{\sec^2(\arctan(x))} = \frac{1}{1+x^2}$$

□

Also, as you might have already guessed. The same happens to every trigonometric function. I leave the proof up to you.

The strange part is that nobody has an intuitive explanation why this happens. It just does through proven formulas and we don't question it.

But wait, we haven't proven the chain rule. So let's do it.

Let's go on a small tangent first. Remember that functions are just arrows from the domain to co-domain?

So we can write functions in a parallel view like this. Here I ask you a simple




Figure 13.0. Parallel view of function

question, by what scale has h been increased due to the function? $\frac{\Delta f}{h}$ is the obvious answer.

But what happens when $h \rightarrow 0$, things start to breakdown as they get smaller, don't they?

Let's talk about what we mean to be small. Smallness is relative, a mouse is small in comparison to an elephant. A flea is small in comparison to the mouse. The flea is beneath notice in comparison to the elephant.

In terms of math, elephant refers to macro numbers like $x, f(x)$ while they can be zero in some cases, they are mostly not.

The increment h is the mouse. Which while very small, is not beyond notice. However, anything which when divided by h is approaches zero can be considered a flea. This makes $h^2, h^3, h^{\frac{4}{3}}$ fleas as $h \rightarrow 0$.

We can say that $\frac{\text{flea}}{h} = \text{mouse}$ (as it is approaching 0, not reaching it)

We can also say that $h * \text{mouse} = \text{flea}$

This all may seem interesting but where are we going with this?

Remember the secant and tangent which we used to create the differentiation formula? We are finally going to return to it.

We know that $\frac{\Delta f}{h} = f'(x)$ as $h \rightarrow 0$

Then we can say $\frac{\Delta f}{h} - f'(x) = 0$ as $h \rightarrow 0$

But as h approaches 0, not reaches it, $\frac{\Delta f}{h} - f'(x) = \text{mouse}$

Multiplying by h gives us: $\Delta f = hf'(x) + \text{flea}$

Graphically this is nothing but the fact that as P and Q come closer, the secant and tangent have the difference of the most minuscule flea.

This in terms of our parallel view of functions we can now say that the scaling factor was $f'(x)h + \text{flea}$ where the flea can be ignored. This is monumental as we only need the value of x to actually find by what the function becomes. This proves the chain rule almost instantly.




Figure 13.0. The chain rule, in parallel form

Proof. We can say that $\Delta u = u'(x)h$ and therefore:

$$\Delta v = v'(u(x))\Delta u$$

$$= v'(u(x))u'(x)h$$

$$\therefore v(u(x))' = \frac{\Delta v}{h} = v'(u(x))u'(x)h$$

And we are done. □

While all this is sure neat, but what is it actually used for? You'll see it in a minute.

13.8 Limits

All this time we have been taking derivatives of functions assuming that they exist. What about when they don't? What about times where a point is not in the domain of $f(x)$ but is in the domain of $f'(x)$? What about all that?

Imagine a blind pirate captain named Goldeyes who lands on an island with a mysterious heavy object hidden in a hole. He faces a dilemma: should he try to retrieve this object for his ship's treasure or leave it behind? To help him decide, Goldeyes turns to his two most trusted companions for their opinions. Here are the three possible outcomes:

Disagreement: Both companions provide conflicting opinions—one claims it's

just a common rock, while the other insists it's the valuable treasure of John Timbers. In this scenario, Goldeyes is left uncertain about the true nature of the object. Consequently, he decides to leave it on the island, as he cannot confidently determine its identity.

Partial Information: One of his companions confidently identifies the object as John Timbers' treasure, but the other companion is unsure and cannot confirm. Again, Goldeyes is faced with uncertainty, and he opts to leave the object behind because he lacks a clear understanding of what it truly is.

Consensus: In the final case, both companions unanimously agree that the object is the same, whether they both claim it's a rock or John Timbers' gold. Goldeyes, despite being blind and unable to verify the object himself, trusts the consensus of his companions and takes action accordingly. If they both say it's gold, he will take it as treasure(even if it is rock); if they both say it's a rock, he will disregard it as a worthless item(even if it is treasure).

How does this have anything to do with calculus? In a minute it will become clear.

Till now we were dealing with continuous functions. A continuous graph refers to functions whose graph can be drawn without lifting the pen. However, we need to understand that some graphs are discontinuous at a number of points.

Let's consider the function $f(x) = \frac{x^2 - 9}{x - 3}$ which has a domain of $\mathbb{R} - 3$ as $\frac{0}{0}$ is undefined. This also makes it discontinuous as the graph has a hole at $x = 3$. However, for rest of the real numbers, $f(x) = x + 3$. So if we look slightly on the left of $x = 3$ we can get a number close to 6 and just on the right of $x = 3$ we have a number close to 6 as well. So we can say that: $\lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3} = 6$. We need to understand that like Captain Goldeye's we still have zero idea on what is in the hole, but based on what two people are telling us we are making out mind on what is in the hole. This is called limit of a function $f(x)$ for $x = a$.

The limit for $f(x)$ at $x = \alpha$ is said to exist if and only if $\lim_{x \rightarrow \alpha^-} f(x) = \lim_{x \rightarrow \alpha^+} f(x)$ and then $\lim_{x \rightarrow \alpha^-} f(x) = \lim_{x \rightarrow \alpha^+} f(x) = \lim_{x \rightarrow \alpha} f(x)$

However, this rule has an exception. If both the left hand limit(LHL) and the right hand limit(RHL) are going to ∞ or $-\infty$ the limit is known as infinite limit. This limit doesn't exist as infinite is very large. Let me explain.

For example, if a man from Paris and a woman from Bordeaux leave for Russia, do they meet? Unless they are characters in a rom-com, the event seem unlikely as Russia is quite large.

The same happens here. Infinity is quite large, how do we know that both the LHL and RHL meet?

Theorem 13.20 (Fundamental theorem of Limits). *If $\lim_{x \rightarrow \alpha} f(x) = l$ and $\lim_{x \rightarrow \alpha} g(x) = k$ then:*

$$\lim_{x \rightarrow \alpha} f(x) \pm g(x) = l \pm m$$

$$\lim_{x \rightarrow \alpha} f(x) \cdot g(x) = l \cdot m$$

$$\lim_{x \rightarrow \alpha} \frac{f(x)}{g(x)} = \frac{l}{m}$$

$$\lim_{x \rightarrow \alpha} n f(x) = n \lim_{x \rightarrow \alpha} f(x) = nl$$

$$\lim_{x \rightarrow \alpha} g(f(x)) = g(\lim_{x \rightarrow \alpha} f(x)) = g(l) \text{ provided } g \text{ is continuous at } g(x) = m$$

We can now do some simple questions pertaining to limits.

Example 13.21. Let $f(x) = \begin{cases} \frac{1}{2} & \text{for } x > 1 \\ 0 & \text{for } x = 1 \\ \frac{1}{2} & \text{for } x < 1 \end{cases}$

What is $\lim_{x \rightarrow 1} f(x)$?

I recommend you to think about this question and make a prediction about this answer. Do not look at the solution. Done.

The answer is not 0. Notice that the function is discontinuous at $x = 1$. We have a hole in the graph and what did Captain Goldeyes teach us? We will trust the LHL and RHL even if what they say is untrue. Here slightly less than 1 would give us $\frac{1}{2}$ and so would slightly greater than 1. Hence, the limit is $\frac{1}{2}$.

It is only in continuous functions(or continuous regions of discontinuous functions) where we can plug in whatever the number is approaching to and expect the limit to be the same. For a point of discontinuity, we will look at the LHL and RHL.

While the above examples was 150% unnecessary and just me trying to be a little tricky and cheeky as this limit has no real significance, we should ask where limits are used?

In certain time functions which occur in physics, the discontinuous points lead to indeterminate forms. These are seven things which are indeterminable in math. However, time is continuous so we are certain that the function needs to have a value. This is what limits allows us to do. The indeterminate forms are $\frac{0}{0}, \frac{\infty}{\infty}, 0 \times \infty, \infty - \infty, \infty^0, 0^0, 1^\infty$. Through algebraic manipulations, all these forms may be inter-converted.

Here are also some forms which a lot of people mistake for indeterminate but are actually not: $\infty + \infty = \infty, \infty \times \infty = \infty, \frac{\alpha}{\infty} = 0$ if α is finite.

Also before someone tries to argue $\infty + \infty = \infty \iff 2 = 1$ remember, $\frac{\infty}{\infty}$ is indeterminate.

Also note: $\frac{\infty}{a}, \frac{a}{0}$ are also indeterminate for $a \in \mathbb{R}$. They are not part of the seven forms as they have a variable as part of their formulation.

Example 13.22. We know that for $\lim_{x \rightarrow 0} \frac{x^2}{x}$ no limit exists, as LHL \neq RHL. Does $\lim_{x \rightarrow 0} \frac{\lfloor x^2 \rfloor}{x}$ exist (here $\lfloor a \rfloor$ refers to the greatest integer less than or equal to a)?

Again I encourage you to think about this. Before we see the solution. While this is also like the previous is 150% unnecessary and just me trying to be a little tricky and cheeky, it touches on some critical concepts.

If you answered that the limit doesn't exist, you are wrong. The greatest integer function (GIF) causes a subtle but monumental change. Let's consider the LHL for maybe $x = -0.001$, $\frac{\lfloor -0.001^2 \rfloor}{-0.001} = \frac{\lfloor 0.000001 \rfloor}{-0.001} = \frac{0}{-0.001} = 0$. If we do the same for RHL we can notice that the GIF has caused LHL = RHL = 0 and hence the limit is equal to 0.

We can resolve indeterminacy in a few ways. We can Simplify, factorize or rationalize the function like we did with $f(x) = \frac{x^2-9}{x-3}$. We can use certain standard limits which we will learn about in a moment. Or we can make some approximations and solve it, like we did with $f(x) = \frac{\sin(x)}{x}$. The most intuitive and simple one is using approximations. You'll see it destroy questions other methods took time around. The only thing is that you'll have to hold your curiosity as their proof is not gonna appear for quite some time, it is in another chapter. But for now, you'll have to trust me. If you do, then here are some common approximations:

Theorem 13.23 (Taylor-Maclaurin Series). $e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$
 $a^x = 1 + \frac{x \ln(a)}{1!} + \frac{x^2 \ln^2(a)}{2!} + \frac{x^3 \ln^3(a)}{3!} + \dots \quad \ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots \text{ for } -1 < x \leq 1$
 $\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$
 $\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$
 $\tan(x) = x + \frac{x^3}{3} + \frac{2x^5}{15} + \dots$
In general, $f(x) = f(0) + \frac{xf'(0)}{1!} + \frac{x^2f''(0)}{2!} + \frac{x^3f'''(0)}{3!} + \dots$

As a even further cheat, we normally use less than the first three terms in the approximations if $x \rightarrow 0$, mostly a single term does the trick. I call this the Brahmastra over the mythological weapon in Indian mythology which can destroy anything and everything the universe.

Let's explore the concepts more through some problems.

Example 13.24.

$$\lim_{x \rightarrow 3} \frac{x^2 - 2x - 3}{x^2 - 4x + 3}$$

$$\begin{aligned} \text{Solution. } & \frac{x^2 - 2x - 3}{x^2 - 4x + 3} \\ &= \frac{(x-3)(x+1)}{(x-3)(x-1)} \\ &= \frac{x+1}{x-1} \end{aligned}$$

$\therefore \lim_{x \rightarrow 3} \frac{x^2 - 2x - 3}{x^2 - 4x + 3} = \lim_{x \rightarrow 3} \frac{3+1}{3-1}$ as the new function is continuous at $x = 3$

$$= \frac{4}{2} = 2$$

□

Example 13.25.

$$\lim_{x \rightarrow 2+\sqrt{3}} \frac{x^4 - 7x^3 + 14x^2 - 7x + 1}{x^2 - 4x + 1}$$

$$\begin{aligned} \text{Solution. } & \frac{x^4 - 7x^3 + 14x^2 - 7x + 1}{x^2 - 4x + 1} = \frac{(x^2 - 3x + 1)(x^2 - 4x + 1)}{x^2 - 4x + 1} \\ &= x^2 - 3x + 1 \\ \therefore & \lim_{x \rightarrow 2+\sqrt{3}} \frac{x^4 - 7x^3 + 14x^2 - 7x + 1}{x^2 - 4x + 1} \\ &= \lim_{x \rightarrow 2+\sqrt{3}} x^2 - 3x + 1 \text{ as } x^2 - 3x + 1 \text{ is continuous at } x = 2 + \sqrt{3}, \text{ we can} \\ &\text{plug it in to get the answer as } 2 + \sqrt{3} \end{aligned}$$

□

So those two were quite standard. Let's do something more fun!

Example 13.26.

$$\lim_{x \rightarrow 1} \frac{3 - \sqrt{8x + 1}}{5 - \sqrt{24x + 1}}$$

Solution. We can solve this in two ways. While one is more tedious other is just chad. We'll do the tedious one first.

To prevent the need to type the limit again and again, assume it's presence throughout the solve.

$$\begin{aligned} & \frac{3 - \sqrt{8x + 1}}{5 - \sqrt{24x + 1}} \\ &= \frac{3 - \sqrt{8x + 1}}{5 - \sqrt{24x + 1}} \cdot \frac{3 + \sqrt{8x + 1}}{3 + \sqrt{8x + 1}} \cdot \frac{5 + \sqrt{24x + 1}}{5 + \sqrt{24x + 1}} \end{aligned}$$

Notice that the last fraction doesn't have any indeterminacy, and is continuous and can hence be separated from the limit using the properties of limits. Therefore we can write it as:

$$\begin{aligned} & \frac{5}{3} \cdot \frac{9 - 8x - 1}{25 - 24x - 1} \\ &= \frac{5}{3} \cdot \frac{8(1-x)}{24(x+1)} \\ &= \frac{5}{3} * \frac{1}{3} = \frac{5}{9} \end{aligned}$$

Now let's do it using the chad method.

We want to use approximations, hence we need the limit to be tending to 0. We'll substitute $x = 1 + t$ and $t \rightarrow 0$,

$$\lim_{t \rightarrow 0} \frac{3 - \sqrt{9 + 8t}}{5 - \sqrt{25 + 24t}}$$

While doesn't seem anything too special, remember the binomial approximation from chapter-9? Assume the limit to be present from here onward

$$\begin{aligned}
&= \frac{3-3(1+\frac{8}{9}t)^{\frac{1}{2}}}{5-5(1+\frac{24}{25}t)^{\frac{1}{2}}} \\
&= \frac{3-3(1+\frac{4t}{9})}{5-5(1+\frac{12t}{25})} \\
&= \frac{3(\frac{4t}{9})}{5(\frac{12t}{25})} \\
&= \frac{3}{5} \cdot \frac{4}{9} \cdot \frac{25}{12} \\
&= \frac{5}{9}
\end{aligned}$$

□

You may have felt that the chad method wasn't really that good...

Example 13.27.

$$\lim_{x \rightarrow 1} \frac{x}{\sqrt{1+x} - \sqrt{1-x}}$$

Solution. Using binomial approximation,

$$\begin{aligned}
\frac{x}{\sqrt{1+x} - \sqrt{1-x}} &= \frac{x}{\sqrt{1+\frac{x}{2}} - \sqrt{1+\frac{1}{x}}} \\
&= \frac{2\cancel{x}}{\cancel{x}} \\
&= 2
\end{aligned}$$

□

I leave trying out the rigorous method to you. Let's now try a problem where the rigorous method will become so complex that I don't think it will be worth the time or effort.

Example 13.28.

$$\lim_{x \rightarrow 2} \frac{3 - \sqrt[3]{x^2 + 5x + 13}}{4 - \sqrt{x^2 + 3x + 6}}$$

Proof. Let $x = 2 + t$ and $t \rightarrow 0$,

$$\begin{aligned}
&\frac{3 - \sqrt[3]{x^2 + 5x + 13}}{4 - \sqrt{x^2 + 3x + 6}} \\
&= \frac{3 - \sqrt[3]{(t+2)^2 + 5(t+2) + 13}}{4 - \sqrt{(t+2)^2 + 3(t+2) + 6}} \\
&= \frac{3 - \sqrt[3]{(t^2 + 4t + 4) + 5t + 10 + 13}}{4 - \sqrt{(t^2 + 4t + 4) + 3t + 6 + 6}} \\
&= \frac{3 - \sqrt[3]{(t^2 + 9t + 27)}}{4 - \sqrt{(t^2 + 7t + 16)}}
\end{aligned}$$

Remember the entire mice and flea discussion? Can we neglect the fleas(t^2) as after all $t \rightarrow 0$?

$$\begin{aligned}
&= \frac{3 - \sqrt[3]{(9t+27)}}{4 - \sqrt{(7t+16)}} \\
&= \frac{3 - 3(1+\frac{t}{3})^{\frac{1}{3}}}{4 - 4(1+\frac{7t}{16})^{\frac{1}{2}}} \\
&= \frac{3(1-1-\frac{t}{3})}{4(1-1-\frac{7t}{32})} \\
&= \frac{3}{4} \cdot \frac{t}{9} \cdot \frac{32}{7t} \\
&= \frac{8}{21}
\end{aligned}$$

□

We need to make a small note here. We cannot just neglect powers which don't exist. For example in $\lim_{x \rightarrow 0} \frac{(x^2+x+1)-(x+1)}{x}$ we cannot neglect x^2 in comparison to x as the x itself is getting canceled out. Remember, size is relative. We need someone in comparison to actually make a neglection. Now let's talk about some standard limits and why people who try to memorize them are ignorant fools.

Theorem 13.29 (Standard results). (1) $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = \lim_{x \rightarrow 0} \frac{\tan(x)}{x} = 1$

$$(2) \lim_{x \rightarrow 0} \frac{\arcsin(x)}{x} = \lim_{x \rightarrow 0} \frac{\arctan(x)}{x} = 1$$

$$(3) \lim_{x \rightarrow \infty} (1 + \frac{1}{x})^x = \lim_{x \rightarrow 0} (1 + x)^{\frac{1}{x}} = e$$

$$(4) \lim_{x \rightarrow \infty} (1 + \frac{a}{x})^x = \lim_{x \rightarrow 0} (1 + ax)^{\frac{1}{x}} = e^a$$

$$(5) \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$$

$$(6) \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln(a) \text{ for } a > 0$$

$$(7) \lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1$$

$$(8) \lim_{x \rightarrow \alpha} \frac{x^n - \alpha^n}{x - \alpha} = n\alpha^{n-1}$$

Proof. The first and second were explained earlier using unit circle. If someone want's they can use the Taylor series but that's overkill.

The third one is a specific case($a = 1$) of the fourth which we'll prove. Assuming limit to be wherever necessary,

$$\text{Let } \lim_{x \rightarrow 0} (1 + ax)^{\frac{1}{x}} = y$$

$$\iff \frac{1}{x} \ln(1 + ax) = \ln(y)$$

$\iff \frac{ax}{x} = \ln(y)$ This is using the Brahmastra.

$$\iff \ln(y) = a$$

$$\iff y = e^a$$

The infinity form is just obtained by taking $x = \frac{1}{n}$.

The fifth is a mild embarrassment using Brahmastra, $\frac{e^x - 1}{x} = \frac{x+1-1}{x} = \frac{x}{x} = 1$

The sixth limit will use the same methodology. $\frac{a^x - 1}{x} = \frac{e^{x \ln(a)} - 1}{x} = \frac{x \ln a + 1 - 1}{x} = \ln(a)$

The seventh limit just shouts Brahmastra,

$$\frac{\ln(1+x)}{x} = \frac{x}{x} = 1$$

The eight limit will however have to wait a minute. \square

Taking the logarithm of the limit(as we did for the forth one) is quite a common technique to deal with limits with variable exponents and I refer to it as Vayuasta, as a reference to the Indian mythological weapon which gives the user complete control of the sky and air. And third and final weapon for

attacking limits is L'Hopital's rule, which I call Agniasthra as a reference to, you guessed it, Indian mythology. As Agniasthra gives the user immense fire power, so does L'Hopital.

Theorem 13.30 (L'Hopital's Rule).

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow x_0} \frac{f'(x)}{g'(x)}$$

If and only if, $f(x)$ and $g(x)$ are differentiable on all points other than x_0 with $g'(x_0) \neq 0$ and $\frac{f(x)}{g(x)}$ is indeterminate which means $f(x)$ or $g(x)$ are both either 0 or ∞

Proof. We know that $\Delta f = hf'(x) \iff f(x+h) - f(x) = hf'(x) \iff f(x+h) = f(x) + hf'(x)$

Now we'll compute the limit for $x \rightarrow x_0$ where $f(x_0) = g(x_0) = 0$

$$\begin{aligned} & \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} \\ &= \lim_{x \rightarrow x_0, h \rightarrow 0} \frac{f(x+h)}{g(x+h)} \\ &= \lim_{x \rightarrow x_0, h \rightarrow 0} \frac{hf'(x) + f(x_0)}{hg'(x) + g(x_0)} \\ &= \lim_{x \rightarrow x_0, h \rightarrow 0} \frac{\cancel{hf'(x)}}{\cancel{hg'(x)}} \\ &= \lim_{x \rightarrow x_0} \frac{f'(x)}{g'(x)} \end{aligned}$$

Hence, proved. □

This makes proving the eighth result a piece of cake.

$$\frac{x^n - a^n}{x - a} = \frac{nx^{n-1}}{1} = nx^{n-1}$$

All these standard results are hence quite easy to get and keeping them in the brain wastes the space which may be used elsewhere. Let's do some examples.

Example 13.31.

$$\lim_{x \rightarrow 0} \frac{3^x - 1}{2^x - 1}$$

Solution. While this question is destroyed in shreds by Brahmastra, I'll use the Agniasthra for instructive purposes.

$(a^x)' = (e^{x \ln(a)})'$ we can use the chain rule here, to get:

$$\begin{aligned} &= \ln(a)e^{x \ln(a)} \\ &= \ln(a)a^x \end{aligned}$$

Using this in our limit gives:

$$\begin{aligned} & \lim_{x \rightarrow 0} \frac{3^x - 1}{2^x - 1} \\ &= \lim_{x \rightarrow 0} \frac{\ln(3)3^x}{\ln(2)2^x} \text{ as the function is now continuous, we can just plug in } x = 0 \\ &= \frac{\ln(3)}{\ln(2)} = \log_2(3) \end{aligned} \quad \square$$

Example 13.32.

$$\lim_{x \rightarrow 0} \frac{1 - \cos 3x}{x^2}$$

Solution. For the first time, we'll have to use the Brahmastra upto two places.

$$\begin{aligned} & \frac{1 - \cos 3x}{x^2} \\ &= \frac{1 - (1 - \frac{(3x)^2}{2!})}{x^2} \\ &= \frac{\cancel{1} - \cancel{1}}{\cancel{x^2}} \\ &= \frac{9}{2} \end{aligned}$$

□

13.9 Continuity

As we have already defined, continuity refers to the fact a graph(or a region of it) can be drawn without lifting the pencil. However, more formally:

Definition 13.33. A function $f(x)$ is continuous at $x = a$ if and only if:
 $f(a^-) = f(a^+) = f(a)$

Basically the right hand function must agree with the left hand function as well as the instantaneous value. Unlike with limits, here $f(a)$ actually matters.

Example 13.34. Let $f(x) = \begin{cases} (1+x)^{\frac{1}{x}} & \text{for } x > 0 \\ a & \text{for } x = 0 \\ \frac{1-\cos(x)}{bx^2} & \text{for } x < 0 \end{cases}$

If $f(x)$ is continuous at $x = 0$, find $\lfloor a + b \rfloor$

Solution. Using the defination of continuity, we can say:

$$(1+x)^{\frac{1}{x}} = a = \frac{1-\cos(x)}{bx^2} \text{ for } x \rightarrow 0$$

You might recall that the limit of $(1+x)^{\frac{1}{x}} = e$ for $x \rightarrow 0$

That means $a = e$. Let's now find b .

$$\begin{aligned} & \lim_{x \rightarrow 0} \frac{1-\cos(x)}{bx^2} = e \\ \Leftrightarrow & \frac{1}{b} \lim_{x \rightarrow 0} \frac{1-\cos(x)}{x^2} = e \\ \Leftrightarrow & b = \frac{1}{e} \lim_{x \rightarrow 0} \frac{1-\cos(x)}{x^2} \\ \Leftrightarrow & b = \frac{1}{e} \lim_{x \rightarrow 0} \frac{1-(1-\frac{x^2}{2})}{x^2} \\ \Leftrightarrow & b = \frac{1}{e} \cdot \frac{1}{2} \end{aligned}$$

Using Brahmastra, obviously Therefore, $\lfloor a + b \rfloor = \lfloor e + \frac{1}{2e} \rfloor = \lfloor 2.90\dots \rfloor = 2$ □

13.10 Application of Differentiation

We are finally nearing the end of our first foray into calculus. We are now going to return from where we had started, real life.

Example 13.35. A 15m ladder is propped up against a window of 12m. It starts moving away from the wall at 1m/s. At what speed is it falling?

Solution. A classic physics problems which can be done using two more physics approaches. However, we'll study the mathematical approach here.

Let the height be $x = 12$ and the run(distance from the base of wall) be y , we can then say:

$x^2 + y^2 = 15^2 \iff y = 9$ We want to know the rate of change of x in terms of time. So we can multiply by $\frac{d}{dt}$ to get:

$$\frac{d(x^2)}{dt} + \frac{d(y^2)}{dt} = \frac{d(15^2)}{dt}$$

$$\iff 2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\iff 2 * 12 \frac{dx}{dt} + 2 * 9 * 1 = 0$$

$$\iff \frac{dx}{dt} = -\frac{3}{4}$$

Which represents that the ladder is moving downward at the speed of $\frac{3}{4}$ m/s. \square

Another use of derivatives is in optimization. Remember, I had promised that we'll talk about the turnings of the graph? The most optimized point for a function where it reaches its maxima or minima will be the cusp of a turning. We can notice that the slope of the tangent is 0 at the points of maxima and minima. Hence, the extreme points are the roots of $f'(x)$.

But its not as simple, the converse is not always true, zero slope may just be a point of inflection(a point higher or lower than its neighbours but not the maxima or minima)

We need to be able to tell whether tell if a point is a maxima or a minima, so here is what we need to notice. The slope starts as positive for a maxima and then becomes negative. The opposite is true for minima. This means that we can take the rate of change of the slope and if it is less than 0, the point is a maxima. If it is more than 0, the point is a minima.

This is the second derivative test. But what if we don't wish to take two derivatives?

The we can notice that the sign of the slope changes at the points. So we can notice the value of $f'(x)$ between two roots to guess if its a minima or maxima.

We don't need to do this for every interval as the sign just alternates.

You'll understand this better through a problem:




Figure 13.0. The turnings have become significant

Example 13.36. An open topped box is to be constructed by removing equal squares from each corner of a 3 metre by 8 metre rectangular sheet of aluminium and folding up the sides. Find the volume of the largest such box.

Solution. Let the side of the square removed be x . We can notice that the volume would be $f(x) = x(3 - 2x)(8 - 2x) = 4x^3 - 22x^2 + 24x$ meter cubed. We can differentiate it to get the equation: $f'(x) = 12x^2 - 44x + 24$

Setting this as zero has two solutions, $x = \frac{2}{3}, 3$

While we know the answer is $\frac{2}{3}$ as one side of the shape is 3. Let's confirm it is the maxima using both the methods.

Single Derivative method: We can notice that $f'(0) = 24 > 0$ which means that $f'(x)$ is positive till $x = \frac{2}{3}$, then negative till $x = 2$, and then positive forever.

We can use this to say $x = \frac{2}{3}$ is the point of maxima and $x = 3$ the point of minima(realistically it is $x = \frac{3}{2}$) Double derivative method: We can compute that $f''(x) = 24x - 44$ which is less than zero for $x = \frac{2}{3}$ and greater than zero for $x = 3$ making them the maxima and minima repectivly.

With the tests out of the way, we can now find the largest possible volume by just plugging in $x = \frac{2}{3}$ to get that the maximum volume would be: $\frac{200}{3}$ meter cubed. \square

And finally, we talk about video games. Most video games need to make trigonometric calculations in an instant especially online FPS and fighting games. While we could have it compute the actual values, that would be mind numbingly slow and make the gameplay worse. For the quick tactile feel we need, we use approximations. However, these approximations also need to be accurate or else the game will feel unrealistic.

How do we achieve this? Remember the mouse-flea discussion and its form we had created during the proof of L'-Hopital?

$f(x+h) = f(x) + hf'(x)$ may seem harmless enough but if we choose a value of $f(x)$ we already know and then move recursively ahead we can get increasingly accurate answers.

Example 13.37. Calculate $\sqrt{73}$

Solution. Let's take $f(x) = \sqrt{x}$ and $x = 8^2 = 64$,

$$\therefore h = 73 - 64 = 9 \text{ and } f'(x) = \frac{1}{2\sqrt{x}}$$

Then we can say that,

$$\begin{aligned}\sqrt{73} &= f(73) = f(64+9) = f(64) + 9f'(64) = 8 + \frac{9}{16} \\ &= 8.5625\end{aligned}$$

The actual value of $\sqrt{73} = 8.54400$ which is within 1% of what we found using the formula only once.

A computer would now compute 8.5625^2 and use that for an even more refined approximation. Most games do it 2-3 times, while even some calculators also use this but do it 5-10 times. \square

And with that this chapter is concluded.

Exercises

- (1) What is the co domain of $f(X) = \arctan(x) + \frac{1}{2} \arcsin(x)$?

(2) Evaluate

$$\lim_{x \rightarrow -\frac{1}{3}} \frac{1}{x} [\frac{-1}{x}]$$

where $[x]$ represents the Greatest integer less than or equal to x

(3) Evaluate:

$$\lim_{n \rightarrow \infty} \frac{5^{n+1} + 3^n - 2^{2n}}{5^n + 3^{2n+3} + 2^n}$$

(4) The value of the given limit is:

$$\lim_{x \rightarrow 0} \frac{\cos(\sin(x)) - \cos(x)}{x^4}$$

(5) If

$$\lim_{n \rightarrow \infty} \frac{1^2 n + 2^2(n-1) + 3^2(n-2) + \dots + n^2 1}{1^3 + 2^3 + 3^3 + \dots + n^3} = \frac{a}{b}$$

then find the value of $a^3 + b^3$

(6)

$$\lim_{x \rightarrow 0} \left(\sum_{r=1}^n r^{\csc^2(x)} \right)^{\sin^2(x)}$$

(7)

$$\lim_{n \rightarrow \infty} [(1 + \frac{1}{n})^n - (1 - \frac{1}{n})]^{-n}$$

(8) If a_1 is the greatest value of $f(x)$, where $f(x) = \frac{1}{2 + [\sin(x)]}$ and $a_{n+1} = a_n + \frac{(-1)^{n+2}}{n+1}$, then $\lim_{n \rightarrow \infty} a_n = ?$ (here $[x]$ is the greatest integer function)

(9) Calculate

$$\lim_{x \rightarrow 0} \left\{ \left[\frac{100x}{\sin(x)} \right] + \left[\frac{99 \sin(x)}{x} \right] \right\}$$

(10) If a_n and b_n are positive integers and $a_n + \sqrt{2}b_n = (2 + \sqrt{2})^n$ then calculate $\lim_{n \rightarrow \infty} (\frac{a_n}{b_n}) = ?$ (11) If $f(x) = \begin{cases} k\sqrt{x+1}, & \text{if } x \leq 3 \\ 2 + mx, & \text{if } x > 3 \end{cases}$ is differentiable at $x = 3$, then find the value if m and k is:(12) (JEE Mains 2018) Let $S = t \in R : f(x) = |x - \pi| \cdot (e^{|x|} - 1) \sin(x)$ is not differentiable then the set S is equal to:(13) (JEE Mains 2021) The number of points at which the function $f(x) = |2x+1| - 3|x+2| + |x^2+x-2|$ with $x \in \mathbb{R}$ is not differentiable, is:

(14) Let $f(x) = e^{x-1} - ax^2 + b$ and $g(x) = \begin{cases} e^{x-1} & \text{if } x \leq 1 \\ x^2 + 1 & \text{if } x > 1 \end{cases}$ then find the value of a and b such that $f(x) \times g(x)$ is differentiable at $x = 1$

(15) (JEE Mains 2020) For a function f defined on $(-\frac{1}{3}, \frac{1}{3})$ by $f(x) = \begin{cases} \frac{1}{x} \ln \frac{1+3x}{1-2x} & \text{when } x \neq 0 \\ k, & \text{when } x = 0 \end{cases}$ is continuous, then k is equal to

(16) (AIEEE 2011) The value of p and q for which the function:

$$f(x) = \begin{cases} \frac{\sin(p+1)x + \sin(x)}{x}, & x < 0 \\ q, & x = 0 \\ \frac{\sqrt{x^2+x} - \sqrt{x}}{x^2}, & x > 0 \end{cases}$$

is continuous for all $x \in \mathbb{R}$ are

(17) (JEE Mains 2020) If $y(a) = \sqrt{2\left(\frac{\tan(x) + \cot(x)}{1 + \sin^2(a)} + \frac{1}{\sin^2(a)}\right)}$ where $a \in (\frac{3\pi}{4}, \pi)$, then $\frac{dy}{dx}$ at $y = \frac{5\pi}{6}$ is:

(18) (IIT 2006) If $f''(x) = -f(x)$ and $g(x) = f'(x)$ and $F(x) = (f(\frac{x}{2}))^2 + (g(\frac{x}{2}))^2$ and given that $F(5) = 5$, then $F(10)$ is

(19) (IIT Adv 2014) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ be respectively given by $f(x) = |x| + 1$ and $g(x) = x^2 + 1$. Defined $h : \mathbb{R} \rightarrow \mathbb{R}$ by $h(x) = \begin{cases} \max\{f(x), g(x)\}, & x \leq 0 \\ \min\{f(x), g(x)\}, & x > 0 \end{cases}$ The number of points at which $h(x)$ is not differentiable is:

14

Calculus II: Indefinite Integration

In Calculus I, we delved into the fundamental concepts of limits, derivatives, continuity, and tried to understand how things change. We learned how to find slopes of curves, calculate instantaneous rates of change, and understand the concept of a limit - and got somewhat comfortable with the idea that as we let things get infinitely close, we can figure out what's happening at a particular point. That was the beginning of our journey into understanding the mathematical underpinnings of change.

Now, as we move into Calculus II, we're going to take this journey a step further. We're going to explore the opposite process: instead of finding rates of change, we're going to learn how to work backward and find the cumulative effect of change over a range, and this is where integration comes into play. But why is this important?

Well, calculus isn't just an abstract exercise in mathematical manipulation. It's the language of the universe. From understanding how objects move under the influence of forces to predicting the spread of diseases in a population, calculus is the tool we use to describe and make sense of the real world. Think of it as both a powerful microscope and telescope that allows us to zoom in and out on nature's processes.

Integration helps us calculate areas, accumulate quantities, and understand the net effect of change over time. Whether you're an engineer designing a bridge, a physicist modeling the motion of planets, an economist analyzing markets, or a biologist studying population growth, you'll find integration to be a fundamental tool.


So, let's get started on this adventure together. Calculus II is where we dig deeper into the mathematical tools that shape our understanding of the universe. It's going to be exciting, challenging, and, I hope, a lot of fun. Welcome to the world of Calculus II!

14.1 The Fundamental nature of graphs

We begin this chapter with a theorem which while of no practical use, is of great importance to the foundations of Calculus.

Theorem 14.1 (Extreme value theorem). *For a function $f(x)$ continuous between $[c, d]$ attains some maximum value $f(a) = M$ such that $f(a) = M \geq f(X)$ for all $x, a \in [c, d]$*

We also claim that a $f(b) = m$ exists such that $f(b) = m \leq f(X)$ for all $x, b \in [c, d]$ using the fact that $-f(x)$ is also continuous and it's maxima also exists.



COULD BE IN THE INTERIOR OR AT ONE OF THE ENDS!

Figure 14.0. Extreme Value Theorem

I have not included the complete rigorous proof of this. However, I'll try to give you an intuitive understanding of the same.

Let's assume that our theorem is false, that we don't have $f(a) = M \geq f(x)$ for $x, a \in [c, d]$. This means that for some values of $x \in [c, d]$, $f(x) = \pm\infty$. Let one of those points be g . Therefore we can say, $\lim_{x \rightarrow g} f(a)$ is infinite limit at $x = g$ making all its neighbours(an infinite of them) incomparable to each other. However, if we have an infinite limit, which we know is not continuous, this will violate the assumption that $f(x)$ is continuous. Hence, our assumption is false and we there exists a $f(a) = M \geq f(x)$ for $x, a \in [c, d]$. Using the extreme value theorem we can get:

Theorem 14.2 (Rolle's Theorem). *If $f(x)$ is continuous in $[c, d]$ and differentiable in (c, d) , and $f(c) = f(d) = 0$, then there is atleast one point $a \in (c, d)$ such that $f'(a) = 0$*

While all this may sound like strange jargon, but it basically says that between any two roots of $f(x)$ we have one point of maxima, minima or inflection. While it is intuitively correct, as we did in Calculus I, here is the more formal version:

Proof. If f is the constant function $f(x) = 0$, then the result is obvious. If f is not constant, then it has non-zero values. Therefore, it attains either a maximum $M > 0$ or a minimum $m < 0$ at some point a , by the Extreme Value Theorem. As a is not one of the endpoints because $M, m \neq 0$, and hence, $f'(a) = 0$

□

And this finally gives rise to another theorem:

Theorem 14.3 (Mean Value Theorem). *If $f(x)$ is continuous in $[c, d]$ and differentiable in (c, d) , then there is a point $a \in (c, d)$ such that $f'(a) = \frac{f(d)-f(c)}{d-c}$*

Proof. Let's define a new function by changing the axis of $f(x)$ by making the chord from c to d the y axis. This can be done simply by subtracting the chord from the function.

$g(x) = f(x) - \frac{f(d)-f(c)}{d-c}(x - c) - f(c)$ Then using Rolle's theorem we can claim that there exists $g'(a) = 0$ where $a \in [c, d]$

$$\text{as } g'(x) = f'(x) - \frac{f(d)-f(c)}{d-c}$$

$$\therefore g'(a) = f'(a) - \frac{f(d)-f(c)}{d-c}$$

$$\iff 0 = f'(a) - \frac{f(d)-f(c)}{d-c}$$

$$\iff f'(a) = \frac{f(d)-f(c)}{d-c}$$

Hence, proved.

□

You should notice that these theorems just say that a point fulfilling some conditions exist. They don't find such points or reveal anything about their nature.

Such theorems are called non-constructive.

14.2 Integration

Let's again think about our car example. If we know the speed of the car and the time it travelled at that speed we can easily find its distance by simply multiplying the numbers.

What if we knew its speed in the first interval and then its speed in second interval and so on? You would say that the distance traveled is the multiplication of the velocity by the length of the interval. More mathematically, you would say it is $\sum_{i=1} v_i t_i$ where the sequence v_1, v_2, \dots is the collection of speeds in the intervals t_1, t_2, \dots .

What if the velocity function had some closed form as a function $f(t)$ of time? Then one particular speed would only be for an infinitely small interval which we can write in Leibniz notation as dt . Hence, the distance traveled can be written as $\sum f(t)dt$. However, \sum is used when we put integer values, but time is continuous and we need all real values between the integers.

We write this using a new notation known as the integral.

$\int_{t_{\text{initial}}}^{t_{\text{final}}} f(t)dt$ Here, you may wonder that as addition is easier than division, why did we not do integral first and then derivatives?

Let's try to solve the integral. We can through the picture attached notice that the integral is just the area under the graph. If I consider a small interval h and consider that the area under the graph changes as our range of integration change(which is obvious), by some function $A(x)$ which is also continuous if $f(x)$ is continuous.

We can say that: $A(t+h) - A(t) = f(t) \cdot h$ as $h \rightarrow 0$

$\therefore f(t) = \frac{A(t+h) - A(t)}{h}, h \rightarrow 0$ which is the definition of differentiation.

$\therefore f(t) = A'(t) \iff f'(t) = A(t) \iff f'(t) = \int f(t)dt$

This relates integration with differentiation which gives us the fundamental theorem of calculus(not the fundamental equation, that was the mouse-flea theorem)

Theorem 14.4 (Fundamental Theorem of Calculus). $\int f'(x)dx = f(x)$

However, integration tends to be a bit tougher(an understatement)

14.3 Techniques of indefinite Integration: Substitution

When we don't put limits on the integration and are only interested in the function describing the area $A(x)$ we call it an indefinite integration. We generally find the anti derivative and add a constant C to it.

A lot of people planning to give MCQ exams skip it as it is tricky and then just differentiate the four/five options. Examiners therefore nowadays put in some random values as the limits of the integration to force the candidate to




Figure 14.0. The Integral

compute the exact integral.

However, all of this is not as difficult if we are clear with Calc I.

Theorem 14.5. (1) $\int x^n dx = \frac{x^{n+1}}{n+1} + C$

$$(2) \int \frac{1}{x} dx = \frac{-1}{x} + C$$

$$(3) \int \frac{1}{\sqrt{x}} dx = 2\sqrt{x} + C$$

$$(4) \int \frac{1}{x} dx = \ln x + C$$

$$(5) \int e^x dx = e^x + C$$

$$(6) \int a^x dx = \frac{a^x}{\ln a}$$

$$(7) \int \sin(x) dx = -\cos(x) + C$$

$$(8) \int \cos(x) dx = \sin(x) + C$$

$$(9) \int \sec^2(x) dx = \tan(x) + C$$

$$(10) \int \csc^2(x) dx = -\cot(x) + C$$

$$(11) \int \sec(x) \tan(x) dx = \sec(x) + C$$

$$(12) \int \csc(x) \cot(x) dx = -\csc(x) + C$$

$$(13) \int \frac{1}{\sqrt{1-x^2}} dx = \arcsin(x) + C$$

$$(14) \int \frac{1}{1+x^2} dx = \arctan(x) + C$$

$$(15) \int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1}(x) + C$$

This all seems like a revision of Calc-I doesn't it?
Let's now look at some real integrals

Theorem 14.6. (1) $\int \tan(x) dx = -\ln |\cos(x)| + C = \ln \sec(x) + C$

$$(2) \int \cot(x) dx = \ln |\sin(x)| + C = -\ln |\csc(x)| + C$$

$$\begin{aligned} (3) \int \sec(x) dx &= \ln |\sec(x) + \tan(x)| + C \\ &= -\ln |\sec(x) - \tan(x)| + C \\ &= \ln |\tan(\frac{\pi}{4} + \frac{x}{2})| + C \end{aligned}$$

$$\begin{aligned} (4) \int \csc(x) dx &= \ln |\csc(x) - \cot(x)| + C \\ &= -\ln |\csc(x) + \cot(x)| + C \\ &= \ln |\tan(\frac{x}{2})| + C \end{aligned}$$

How did these appear? We'll see in just a minute. Just remember them for now, we will derive them in a while. Till then here are some more facts about integrals:

Theorem 14.7 (Basic Facts about integrals). $\int f_1(x) \pm f_2(x) dx = \int f_1(x) dx \pm \int f_2(x) dx$
 $\therefore \int Kf(x) dx = K \int f(x) dx$

Let's do some examples now.

Example 14.8. If $f''(x) = 10$ and $f(1) = 6$ and $f(1) = 4$ then find $f(-1)$

Solution. As $f''(x) = 10$

$$\therefore f'(x) = 10x + C \text{ by integrating on both sides}$$

$$\therefore f'(1) = 10 + C = 6 \iff C = -4$$

$$\therefore f'(x) = 10x - 4$$

$$\therefore f(x) = 5x^2 - 4x + c$$

$$\therefore f(1) = 5 - 4 + c = 4 \iff c = 3$$

$$\therefore f(x) = 5x^2 - 4x + 3$$

$$\therefore f(-1) = 5 + 4 + 3 = 12$$

□

This question was just the start. Remember Brahmastra? Let's derive it for $\arctan(x)$

Proof. We know that $\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \dots$ using sum of infinite GP.
 Integrating both sides will give us:

$$\int \frac{1}{1+x^2} dx = \arctan(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

□

A lot of the approximations will get proved by this method, the others will have to wait a while.

Example 14.9. Evaluate:

$$\int \frac{1}{\sin^2(x) \cos^2(x)} dx$$

Solution. Using the fact $\sin^2(x) + \cos^2(x) = 1$,

$$\begin{aligned} & \int \frac{1}{\sin^2(x) \cos^2(x)} dx \\ &= \int \frac{\sin^2(x) + \cos^2(x)}{\sin^2(x) \cos^2(x)} dx = \int \frac{\sin^2(x)}{\sin^2(x) \cos^2(x)} dx + \int \frac{\cos^2(x)}{\sin^2(x) \cos^2(x)} dx = \int \frac{1}{\cos^2(x)} dx + \\ & \int \frac{1}{\sin^2(x)} dx = \int \sec^2(x) dx + \int \csc^2(x) dx = \tan(x) - \cot(x) \end{aligned}$$

□

That was quite fun, wasn't it? Like a swimming pool dive, scary till we jumped and fun after a while. Here is an even easier example for you to try.

Example 14.10. Evaluate:

$$\int \tan^2(x) dx$$

Let's now look at more integrals:

Example 14.11. Evaluate: $\int \frac{1}{1+\cos(2x)} dx$

Solution. Using the 'remember for life' facts from the trigonometry chapter.

$$\begin{aligned} & \int \frac{1}{1+\cos(2x)} dx \\ &= \int \frac{1}{2\cos^2(x)} dx \\ &= \frac{1}{2} \int \sec^2(x) dx \\ &= \frac{\tan(x)}{2} \end{aligned}$$

□

Example 14.12. Evaluate: $\int \frac{\cos(x)-\cos(2x)}{1-\cos(x)} dx$

Solution. Using another of the 'remember for life' facts from trigonometry:

$$\begin{aligned} & \int \frac{\cos(x)-\cos(2x)}{1-\cos(x)} dx \\ &= \int \frac{\cos(x)-2\cos^2(x)+1}{1-\cos(x)} dx \\ &= \int \frac{2\cos^2(x)-\cos(x)-1}{\cos(x)-1} dx \\ &= \int \frac{(2\cos(x)+1)(\cos(x)-1)}{\cos(x)-1} dx \\ &= \int 2\cos(x) + 1 dx \\ &= 2\sin(x) + x \end{aligned}$$

□

Example 14.13. Evaluate:

$$\int \frac{x^2+\cos^2(x)}{1+x^2} \csc^2(x) dx$$

Solution. Algebra and Trig in one question seem bad. What if we can cancel at least one of them? Also, isn't $\csc(x) = \frac{1}{\sin(x)}$?

$$\begin{aligned} & \int \frac{x^2+\cos^2(x)}{1+x^2} \csc^2(x) dx \\ &= \int \frac{x^2+1-\sin^2(x)}{1+x^2} \csc^2(x) dx \\ &= \int \frac{x^2+1}{1+x^2} \csc^2(x) dx - \int \frac{\sin^2(x)}{1+x^2} \csc^2(x) dx \\ &= \int \csc^2(x) dx - \int \frac{1}{1+x^2} dx \\ &= -\cot(x) - \arctan(x) \end{aligned}$$

□

And with this out of the way, we are ready for one the most scary integral which can be solved using these formulas:

Example 14.14. Evaluate: $\int \frac{\sin^6(x)+\cos^6(x)}{\sin^2(x)\cos^2(x)} dx$

Solution. Every time we look at anything with trig, we should remember the formulas for life.

$$\begin{aligned}
 & \int \frac{\sin^6(x) + \cos^6(x)}{\sin^2(x) \cos^2(x)} dx \\
 &= \int \frac{1 - 3\sin^2(x)\cos^2(x)}{\sin^2(x) \cos^2(x)} dx \\
 &= \int \frac{1}{\sin^2(x) \cos^2(x)} dx - \int 3dx \text{ Here is a cardinal rule of integration, we only} \\
 &\text{respect or fear an integral once. The next time its a tamed beast, we proceed} \\
 &\text{to use it.} \\
 &= \tan(x) - \cot(x) - 3x \quad \square
 \end{aligned}$$

Before we move to our first integration technique, here is an observation you would have made:

Theorem 14.15 (Integration of function on a linear equation). *If $\int f(x)dx = F(x) + C$, then $\int f(ax + b)dx = \frac{1}{a}F(ax + b) + c$*

We'll prove this in a while, but let's take it for a test ride.

Example 14.16. Evaluate: $\int \frac{2x+7}{\sqrt{3x-5-\sqrt{x-2}}} dx$

Solution. Remember, we can rationalize the radical?

$$\begin{aligned}
 \int \frac{2x+7}{\sqrt{3x-5-\sqrt{x-2}}} dx &= \int \frac{(2x+7)(\sqrt{3x-5+\sqrt{x-2}})}{(\sqrt{3x-5-\sqrt{x-2}})(\sqrt{3x-5+\sqrt{x-2}})} dx = \int \frac{(2x+7)(\sqrt{3x-5+\sqrt{x-2}})}{2x+7} dx \\
 &= \int \sqrt{3x-5} + \sqrt{x-2} dx = \frac{2}{9}(3x-5)^{\frac{3}{2}} + \frac{2}{3}(x-2)^{\frac{3}{2}} \quad \square
 \end{aligned}$$

And the final question before we see the more powerful methods,

Example 14.17. Evaluate: $\int \sin^4\left(\frac{x}{4}\right) + \cos^4\left(\frac{x}{4}\right) dx$

Solution. It is uncanny how useful the trigonometry formulas have been. We'll use the formula for life of $\sin^2(\theta)$

$$\begin{aligned}
 & \int \sin^4\left(\frac{x}{4}\right) + \cos^4\left(\frac{x}{4}\right) dx \\
 &= \int 1 - 2\sin^2\left(\frac{x}{4}\right)\cos^2\left(\frac{x}{4}\right) dx \\
 &= \int 1 - \frac{4}{2}\sin^2\left(\frac{x}{4}\right)\cos^2\left(\frac{x}{4}\right) dx \\
 &= \int 1 - \frac{1}{2}\sin^2\left(\frac{x}{2}\right) dx \\
 &= \int \frac{3}{4} - \frac{\cos(x)}{4} dx \text{ Using what we just discussed} \\
 &= \frac{3x}{4} + \frac{\sin(x)}{4} \quad \square
 \end{aligned}$$

And now let's learn a new technique.

14.3.1 Integration by Substitution.

Theorem 14.18 (Integration by Substitution). *For $I = \int f(g(x)) \cdot g'(x)dx$, We can put $g(x) = t$ and $g'(x)dx = dt$,*
 $\therefore I = \int f(t)dt$

This theorem is trivial to prove as it does nothing new, just does it in a way cooler fashion. Let's just see it in action to understand

Example 14.19 (Motivating example). Evaluate: $\int x^2 e^{x^3} dx$

$$\begin{aligned} \text{Solution. } & \text{Let } t = x^3, \\ & \therefore dt = 3x^2 dx \therefore \int x^2 e^{x^3} dx \\ &= \int \frac{1}{3} 3x^2 e^{x^3} dx \\ &= \frac{1}{3} \int e^t dt \\ &= \frac{1}{3} e^t \\ &= \frac{e^{x^3}}{3} \end{aligned}$$

□

Not that this is the first and last time we'll actually show that we multiply and divide to get the coefficient. Next question onward, we just directly bring the reciprocal of the coefficient out of the integral.

We can also use it to prove the integration of a function of a linear equation:

Proof. Let $\int f(x)dx = F(x) + C$.

Then to evaluate $\int f(ax + b)dx$, let $ax + b = t$ and therefore, $adx = dt$, therefore:

$$\begin{aligned} & \int f(ax + b)dx \\ &= \frac{1}{a} \int f(t)dt \\ &= \frac{1}{a} F(t) + C \\ &= \frac{1}{a} F(ax + b) + C \end{aligned}$$

□

We can also prove the integral of $\tan(x)$, $\csc(x)$, $\sec(x)$, $\cot(x)$ using this technique.

Proof. Let's prove for $\tan(x)$ first. We know that $\tan(x) = \frac{\sin(x)}{\cos(x)}$ and that the derivative of $\cos(x)$ is $-\sin(x)$, therefore:

$$\begin{aligned} & \int \tan(x)dx \\ &= \int \frac{\sin(x)}{\cos(x)} dx \\ &= -\int \frac{1}{t} dt \text{ We have put } t = \cos(x) \\ &= -\ln t \\ &= \ln \sec(x) \end{aligned}$$

□

I leave the proof for the integral of $\cot(x)$ for you to do, its just the same but the substitution changes slightly.

Proof. Let's now prove for $\csc(x)$. We have two ways of doing this, the other one appears later.

$$\begin{aligned} & \int \csc(x) dx \\ &= \int \csc(x) \frac{\csc(x) + \cot(x)}{\csc(x) + \cot(x)} dx \\ &= \int \frac{\csc^2(x) + \csc(x) \cot(x)}{\csc(x) + \cot(x)} dx \end{aligned}$$

We will now put $t = \csc(x) + \cot(x)$ and therefore $dt = (\csc^2(x) + \csc(x) \cot(x))dx$

$$\begin{aligned} & \therefore \int \frac{\csc^2(x) + \csc(x) \cot(x)}{\csc(x) + \cot(x)} dx \\ &= \int \frac{1}{t} dt \\ &= \ln t \\ &= \ln \csc(x) + \cot(x) \quad \square \end{aligned}$$

The proof for the integral of $\sec(x)$ is quite similar and should be attempted by you before moving on.

Done? Write it in the margins or on a paper and let's continue with more questions.

Example 14.20. Evaluate: $\int \frac{\sin(\sqrt{x})}{\sqrt{x}} dx$

This is quite simple. Let $t = \sqrt{x}$, you ~~can~~ should solve from here onward by taking $dt = \frac{1}{2\sqrt{x}}$

Example 14.21. Evaluate: $\int \frac{1}{x^2 \sin \frac{1}{x}} dx$

Solution. Here the sine is only there to shift your focus away from the fact that $(\frac{1}{x})' = \frac{-1}{x^2}$.

$$\begin{aligned} & \therefore \int \frac{1}{x^2 \sin \frac{1}{x}} dx \\ &= \int \frac{1}{x^2} \csc \frac{1}{x} dx \\ &= - \int \csc t dt \\ &= - \ln |\csc(t) - \cot(t)| \\ &= \ln |\csc(t) + \cot(t)| \\ &= \ln |\tan(\frac{t}{2})| \text{ Remember, } \int \csc(x) dx \text{ has three forms?} \quad \square \end{aligned}$$

This all was quite mild, right? Let's taste the real chilli

Example 14.22. Evaluate $\int \frac{1+\ln x}{x^x} dx$

Solution. The x^x seems like the most scary part. Let's take that as t . We have differentiated this before and know that $(x^x)' = x^x(1 + \ln x)$

With that in mind, we can say let $t = x^x$ and $dt = x^x(1 + \ln x)dx \iff \frac{dt}{x^x} = 1 + \ln x dx$

$$\begin{aligned}\therefore \int \frac{1+\ln x}{x^x} dx \\ &= \int \frac{1}{x^x \cdot x^x} dt \\ &= \int \frac{1}{t^2} dt \\ &= \frac{-1}{t} \\ &= \frac{-1}{x^x}\end{aligned}$$

□

Integration becomes hard for only one reason, that we need to do some hit and trial to get an answer.

In this method, we need to try a few things before we can be sure about what to substitute.

The only way to achieve speed and intuition in the game of hit and trial is to play the game many times. But sometimes, sometimes a little wishful thinking also helps out. Here are two examples to illustrate my point:

Example 14.23. Evaluate $\int \frac{\sec^4(x)}{\sqrt{\tan(x)}} dx$

Solution. How great would it be if we had $\sec^2(x)$ instead of $\sec^4(x)$. Everything else as $\tan x$ and a single $\sec^2(x)$... Let's make that true:

$$\begin{aligned}\int \frac{\sec^4(x)}{\sqrt{\tan(x)}} dx \\ &= \int \frac{\sec^2(x) \cdot \sec^2 x}{\sqrt{\tan(x)}} dx \\ &= \int \frac{\sec^2 x (1 + \tan^2(x))}{\sqrt{\tan(x)}} dx \text{ Let } \tan(x) = t \text{ and } dt = \sec^2(x) dx \\ \therefore \int \frac{\sec^2 x (1 + \tan^2(x))}{\sqrt{\tan(x)}} dx \\ &= \int \frac{1+t^2}{\sqrt{t}} dt \\ &= 2t^{\frac{1}{2}} + \frac{2}{5}t^{\frac{5}{2}} = 2\tan(x)^{\frac{1}{2}} + \frac{2}{5}(\tan(x))^{\frac{5}{2}}\end{aligned}$$

□

Example 14.24. Evaluate: $\int \frac{1}{e^x+1} dx$

Solution. How much better would it be only if we had e^x as the numerator? Why not make it happen?

$$\begin{aligned}\int \frac{1}{e^x+1} dx \\ &= \int \frac{1+e^x-e^x}{e^x+1} dx \\ &= \int 1 dx - \int \frac{e^x}{e^x+1} dx \text{ And now we can take } e^x + 1 = t \text{ and } dt = e^x dx, \\ \therefore \int 1 dx - \int \frac{e^x}{e^x+1} dx \\ &= x - \int \frac{1}{t} dt \\ &= x - \ln t \\ &= x - \ln e^x + 1\end{aligned}$$

□

And whenever this fails, we can use our strategy of just making the longest term equal t

Example 14.25. Evaluate $\int \frac{(x+\sqrt{1+x^2})^{2023}}{\sqrt{1+x^2}} dx$

Solution. The longest term is $x + \sqrt{1+x^2}$ so let $t = x + \sqrt{1+x^2}$ and on differentiating we get $dt = (1 + \frac{2x}{2\sqrt{1+x^2}})dx \iff dt = \frac{x+\sqrt{1+x^2}}{\sqrt{1+x^2}}$

This essentially solves the question.

$$\begin{aligned} & \therefore \int \frac{(x+\sqrt{1+x^2})^{2023}}{\sqrt{1+x^2}} dx \\ &= \int \frac{(x+\sqrt{1+x^2})^{2022} \cdot (x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx \\ &= \int t^{2022} dt \\ &= \frac{t^{2023}}{2023} \\ &= \frac{x+\sqrt{1+x^2}^{2023}}{2023} \end{aligned}$$

□

This technique didn't use much tri...

Example 14.26. Evaluate $\int (\sec(x) + \tan(x))^{\frac{1}{2}} \sec^2(x) dx$

Proof. We can solve this in many ways, but looking at $\sec^2(x)dx$ I want to substitute $\tan(x) = t$. But the problem is that it converts the integral to $\int \sqrt{1+t^2+t} dt$ which is still quite hard to solve.

What if we let $\sec(x) + \tan(x) = t$ which would mean $\sec^2(x) - \tan^2(x) = 1 \iff (\sec(x) - \tan(x))(\sec(x) + \tan(x)) = 1 \iff \sec(x) - \tan(x) = \frac{1}{t}$.

Using these two equations, we can say $2\tan(x) = t - \frac{1}{t}$

$\therefore 2\sec^2(x)dx = (1 + \frac{1}{t^2})dt$, now we can say:

$$\begin{aligned} & \int (\sec(x) + \tan(x))^{\frac{1}{2}} \sec^2(x) dx \\ &= \frac{1}{2} \int (t)^{\frac{1}{2}} (1 + \frac{1}{t^2}) dt \\ &= \frac{1}{2} \int (t)^{\frac{1}{2}} + (t)^{-\frac{3}{2}} dt \\ &= \frac{1}{2} (\frac{2}{3} t^{\frac{3}{2}} - 2t^{-\frac{1}{2}}) = \frac{1}{3} t^{\frac{3}{2}} - t^{-\frac{1}{2}} = \frac{(\sec(x)+\tan(x))^{\frac{3}{2}}}{3} - (\sec(x) + \tan(x))^{\frac{-1}{2}} \end{aligned}$$

□

However, till now we had elements in the function which we substituted as t . Sometimes, the strategy needs to change:

Example 14.27. Evaluate $\int \frac{\sqrt{x}}{\sqrt{1-x^3}} dx$

Solution. We can try substituting $x^3, 1-x^3, \sqrt{x}, \sqrt{1-x^3}$ all with no avail. What we need to realize is that we know the integral of $\frac{1}{\sqrt{1-x^2}}$. What if we could somehow convert x^3 to x^2 . Let's be wishful and assume $t^2 = x^3 \iff$

$t = x^{\frac{3}{2}}$ and therefore $\frac{3}{2}\sqrt{x}dx = dt$, and we are done.

$$\begin{aligned} & \therefore \int \frac{\sqrt{x}}{\sqrt{1-x^3}} dx \\ &= \frac{2}{3} \int \frac{1}{\sqrt{1-t^2}} dt \\ &= \frac{2}{3} \arcsin(t) \\ &= \frac{2}{3} \arcsin(x^{\frac{3}{2}}) \end{aligned}$$

□

Before we move to the next technique in integration, here is another common configuration which occurs in integration by substitution.

Theorem 14.28. For integral in the form of: $\int \frac{1}{x(x^n+1)} dx$; $\int \frac{1}{x^n(x^n+1)^{\frac{1}{n}}} dx$; $n \in \mathbb{N}$ We can take x^n common and pull it out of the bracket and substitute $1+x^{-n} = t$

The theorem may seem a bit confusing, here is an example to show how its used.

Example 14.29 (Motivating Example). (Jee Mains 2015) $\int \frac{1}{x^2(1+x^4)^{\frac{3}{4}}} dx = ?$

Proof. We'll take x^4 common from the bracket.

$$\begin{aligned} & \int \frac{1}{x^2(1+x^4)^{\frac{3}{4}}} dx \\ &= \int \frac{1}{x^2(x^4(\frac{1}{x^4}+1))^{\frac{3}{4}}} dx \\ &= \int \frac{1}{x^5((\frac{1}{x^4}+1))^{\frac{3}{4}}} dx \end{aligned}$$

Note: I have shown the entire taking common and then applying the exponent, here just for instructive purposes. I don't intend to show that again.

Now we take $t = 1 + x^{-4}$ and therefore $dt = \frac{-4}{x^5} dx$, therefore:

$$\begin{aligned} & \int \frac{1}{x^5((\frac{1}{x^4}+1))^{\frac{3}{4}}} dx \\ &= \frac{-1}{4} \int \frac{1}{t^{\frac{3}{4}}} dt \\ &= \frac{-1}{4} \cdot 4 \cdot t^{\frac{1}{4}} \\ &= -1 + \frac{1}{x^4} \end{aligned}$$

□

As you could see, this algorithm made quick work of a seemingly scary question.

Here is one for you to try

Example 14.30. $\int \frac{1}{x(x^5+1)} dx$

And before we move to another method, here is a lengthy(albeit not that hard) example

Example 14.31. Given that $f(0) = f'(0) = 0$ and $f''(x) = \sec^4(x) + \sec^2(x)\tan^2(x) + 4$, find $f(x)$

Solution. It's obvious that we need to integrate $f''(x)$ twice and get the C for those values. So lets do it:

$$\begin{aligned} f'(x) &= \int \sec^4(x) + \sec^2(x)\tan^2(x) + 4dx \\ &= \int \sec^2(x)(\sec^2(x) + \tan^2(x))dx + 4x + C \\ &= \int \sec^2(x)(1+2\tan^2(x))dx + 4x + C \text{ we obviously take } t = \tan(x) \text{ and therefore } dt = \sec^2(x)dx \\ &= \int 1 + 2t^2dt + 4x + C \\ &= t + \frac{2t^3}{3} + 4x + C \\ &= \tan(x) + \frac{2\tan^3(x)}{3} + 4x + C \text{ As } f'(0) = 0, \text{ we can easily say } C = 0. \text{ Hence,} \\ f'(x) &= \tan(x) + \frac{2\tan^3(x)}{3} + 4x, \text{ which we'll have to integrate again(screaming inside).} \end{aligned}$$

$$\begin{aligned} &\int \tan(x) + \frac{2\tan^3(x)}{3} + 4xdx \\ &= \ln|\sec(x)| + \frac{2}{3} \int \tan^3(x)dx + 2x^2 + C \\ &= \ln|\sec(x)| + \frac{2}{3} \int \tan(x) \cdot \tan^2(x)dx + 2x^2 + C \\ &= \ln|\sec(x)| + \frac{2}{3} \int \tan(x) \cdot (1 - \sec^2(x))dx + 2x^2 + C \\ &= \ln|\sec(x)| + \frac{2}{3} (\int \tan(x)dx - \int \tan(x) \sec^2(x)dx) + 2x^2 + C \\ &= \ln|\sec(x)| + \frac{2}{3} (\ln|\sec(x)| - \frac{\tan^2(x)}{2}) + 2x^2 + C \\ &= \frac{5}{3} \ln|\sec(x)| - \frac{\tan^2(x)}{3} + 2x^2 + C \\ \text{As } f(0) &= 0, C = 0 \\ \therefore f(x) &= \frac{5}{3} \ln|\sec(x)| - \frac{\tan^2(x)}{3} + 2x^2 \end{aligned}$$

□

14.3.2 Standard Integrations. Now let's refine this substitution to solve certain questions at extreme speeds.

Theorem 14.32. $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \arctan(\frac{x}{a})$

Proof. I'll offer two proofs for this. One uses a substitution trick and other is vanilla integration. Let's do the vanilla one before we add the chocolate sauce.

$$\begin{aligned} \int \frac{1}{a^2+x^2} dx &= \int \frac{1}{a^2(1+\frac{x^2}{a^2})} dx \\ &= \frac{1}{a^2} \int \frac{1}{1+\frac{x^2}{a^2}} dx = \frac{1}{a^2} \cdot a \cdot \arctan \frac{x}{a} = \frac{1}{a} \arctan \frac{x}{a} \text{ The integral got solved using} \\ &\text{the integration of function over linear equation.} \end{aligned}$$

Now let's add the chocolate sauce. As $\tan(\theta)$ can take any real value, we can

claim that $x = a \tan(\theta) \iff \theta = \arctan(\frac{x}{a})$ and therefore $dx = a \sec^2(\theta)d\theta$

$$\begin{aligned} & \therefore \int \frac{1}{a^2+x^2} dx \\ &= \int \frac{a \sec^2(\theta)}{a^2+a^2 \tan^2(\theta)} d\theta \\ &= \frac{1}{a} \int \frac{\sec^2(\theta)}{1+\tan^2(\theta)} d\theta \\ &= \frac{1}{a} \int \frac{\cancel{\sec^2(\theta)}}{\cancel{\sec^2(\theta)}} d\theta \\ &= \frac{\theta}{a} \\ &= \frac{1}{a} \arctan\left(\frac{x}{a}\right) \end{aligned}$$

□

The second proof may seem unnecessary complex, but we'll see its power in a while.

Theorem 14.33. $\int \frac{1}{x^2-a^2} dx = \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right|$ Multiplying by -1 on both sides will give us:

$$\int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \ln \left| \frac{x+a}{x-a} \right|$$

Proof. Unfortunately, the chocolate sauce method is useless here as we have no part which even resembles a trig function.

So let's do the vanilla solving here:

$$\begin{aligned} & \int \frac{1}{x^2-a^2} dx \\ &= \int \frac{1}{(x-a)(x+a)} dx \\ &= \int \frac{1}{2a} \frac{(x+a)-(x-a)}{(x-a)(x+a)} dx \\ &= \frac{1}{2a} \int \frac{(x+a)-(x-a)}{(x-a)(x+a)} dx \\ &= \frac{1}{2a} \left(\int \frac{1}{x-a} dx - \int \frac{1}{x+a} dx \right) \\ &= \frac{1}{2a} (\ln|x-a| - \ln|x+a|) \\ &= \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| \text{ The other one simply comes by multiplying } -1 \text{ on both sides.} \end{aligned}$$

□

Theorem 14.34. $\int \frac{1}{\sqrt{x^2-a^2}} dx = \ln|x + \sqrt{x^2-a^2}|$

Proof. And suddenly the vanilla methods don't seem to work. Let $x = a \sec(\theta)$ and therefore $dx = a \sec(\theta) \tan(\theta)d\theta$, therefore:

$$\begin{aligned} & \int \frac{1}{\sqrt{x^2-a^2}} dx = \int \frac{a \sec(\theta) \tan(\theta)}{\sqrt{a^2(\sec^2(\theta)-1)}} d\theta \\ &= \int \frac{a \sec(\theta) \tan(\theta)}{a \sqrt{\tan^2(\theta)}} d\theta \\ &= \int \frac{\sec(\theta) \tan(\theta)}{\tan^2(\theta)} d\theta \\ &= \int \sec(\theta) d\theta \\ &= \ln|\sec \theta + \tan \theta| = \ln\left|\frac{x}{a} + \frac{\sqrt{x^2-a^2}}{a}\right| = \ln|x + \sqrt{x^2-a^2}| - \ln a = \ln|x + \sqrt{x^2-a^2}| \end{aligned}$$

In the last step, $\ln a$ is just absorbed into the integration constant.

□

Theorem 14.35. $\int \frac{1}{\sqrt{x^2+a^2}} dx = \ln|x + \sqrt{x^2+a^2}|$

This proof is left to you. Just take $x = a \tan(\theta)$ and the proof follows from that.

Theorem 14.36. $\int \frac{1}{\sqrt{a^2 - x^2}} dx = \arcsin\left(\frac{x}{a}\right)$

Proof. Here we can take $x = a \sin(\theta) \iff \theta = \arcsin\left(\frac{x}{a}\right)$ as if $x^2 > a^2 \iff |\frac{x}{a}| > 1$, we'll get a negative inside the radical, which is not possible(in this plane at least). Therefore $dx = a \cos(\theta)d\theta$

, using this:

$$\begin{aligned} & \int \frac{1}{\sqrt{a^2 - x^2}} dx \\ &= \int \frac{a \cos(\theta)}{\sqrt{a^2 - a^2 \sin^2(\theta)}} d\theta \\ &= \int \frac{a \cos(\theta)}{\sqrt{a^2(1 - \sin^2(\theta))}} d\theta \\ &= \int \frac{a \cos(\theta)}{a \cos(\theta)} d\theta \\ &= \theta \\ &= \arcsin\left(\frac{x}{a}\right) \quad \square \end{aligned}$$

While all these derivations are not particularly tricky, I strongly recommend that you remember these theorems. A simple way to do so, as I have said before, is to write them on a page and look at it for three days straight just after waking up and just before going to sleep.

Now let's do some examples:

Example 14.37. Evaluate: $\int \frac{x^2}{x^6 + a^6} dx$

Solution. The x^2 in the numerator seems unnecessary. We can simply remove it if we take $t = x^3$ and that seems promising as it allows us to make the denominator one of the functions we just studied. Therefore, $dt = 3x^2 dx$, Using this:

$$\begin{aligned} & \int \frac{x^2}{x^6 + a^6} dx \\ &= \frac{1}{3} \int \frac{1}{t^2 + (a^3)^2} dt \\ &= \frac{1}{3} \cdot \frac{1}{a^3} \arctan \frac{t}{a^3} \\ &= \frac{1}{3a^3} \arctan \frac{x^3}{a^3} \quad \square \end{aligned}$$

Remember, we had learnt that every quadratic can be written as perfect square \pm some constant?

We'll use it now

Theorem 14.38. For integral of the type: $\int \frac{1}{ax^2 + bx + c} dx$; $\int \frac{1}{\sqrt{ax^2 + bx + c}} dx$ We should express $ax^2 + bx + c$ in the form of a perfect square \pm some constant

and then apply the standard results.

Let's see an example:

Example 14.39. Evaluate: $\int \frac{1}{2x^2 - 2x + 3} dx$

$$\begin{aligned} \text{Solution. } \int \frac{1}{2x^2 - 2x + 3} dx &= \frac{1}{2} \int \frac{1}{x^2 - x + \frac{3}{2}} dx = \frac{1}{2} \int \frac{1}{x^2 - x + \frac{1}{4} - \frac{1}{4} + \frac{6}{4}} dx = \frac{1}{2} \int \frac{1}{(x - \frac{1}{2})^2 + \frac{5}{4}} dx = \\ &\frac{1}{2} \int \frac{1}{(x - \frac{1}{2})^2 + \frac{5}{4}} dx = \frac{1}{2} \cdot \frac{2}{\sqrt{5}} \arctan(\frac{2x-1}{\sqrt{5}}) = \frac{1}{\sqrt{5}} \arctan(\frac{2x-1}{\sqrt{5}}) \quad \square \end{aligned}$$

And here is one for you to try:

Example 14.40. Evaluate: $\int \frac{1}{\sqrt{x^2 - 4x + 6}} dx$

Now let's look at another configuration, but this time through a motivating example:

Example 14.41 (Motivation Example). Evaluate: $\int \frac{6x+7}{\sqrt{3x^2+7x+65}} dx$

Solution. The numerator looks awful lot like the differentiation of the polynomial inside the radical. Let $t = 3x^2 + 7x + 65$ and therefore $dt = (6x + 7)dx$. Apple is red, question is dead.

$$\begin{aligned} &\int \frac{6x+7}{\sqrt{3x^2+7x+65}} dx \\ &= \int \frac{1}{\sqrt{t}} dt \\ &= 2\sqrt{t} \\ &= 2\sqrt{3x^2 + 7x + 65} \quad \square \end{aligned}$$

The only reason this question was in this section is because a lot of people try to use standard results here, where normal substitution is enough. Here is the general form of this configuration:

Theorem 14.42. For integral of the type: $\int \frac{px+q}{ax^2+bx+c} dx$; $\int \frac{px+q}{\sqrt{ax^2+bx+c}} dx$ We should express $px + q$ in the form of the derivative of the denominator times a constant \pm some other constant and then apply the standard results.

Let's do a more juicy example

Example 14.43. Evaluate: $\int \frac{2x+1}{\sqrt{2x^2+x-1}} dx$

Solution. This gets very, very messy.

We know that the derivative of $2x^2 + x - 1$ is $4x + 1$. We can write $2x + 1$ as $\frac{1}{2}(4x + 1) + \frac{1}{2}$. Therefore:

$$\begin{aligned} & \int \frac{2x+1}{\sqrt{2x^2+x-1}} dx \\ &= \frac{1}{2} \int \frac{4x+1}{\sqrt{2x^2+x-1}} dx + \frac{1}{2} \int \frac{1}{\sqrt{2x^2+x-1}} dx = \sqrt{2x^2+x-1} + \frac{1}{2\sqrt{2}} \int \frac{1}{\sqrt{x^2+\frac{x}{2}-\frac{1}{2}}} dx = \\ & \sqrt{2x^2+x-1} + \frac{1}{2\sqrt{2}} \int \frac{1}{\sqrt{x^2+\frac{x}{2}+\frac{1}{16}-\frac{1}{16}-\frac{8}{16}}} dx = \sqrt{2x^2+x-1} + \frac{1}{2\sqrt{2}} \int \frac{1}{\sqrt{(x+\frac{1}{4})^2-\frac{9}{16}}} dx = \\ & \sqrt{2x^2+x-1} + \frac{1}{2\sqrt{2}} \ln x + \frac{1}{4} - \sqrt{(x+\frac{1}{4})^2-\frac{9}{16}} \\ &= \sqrt{2x^2+x-1} + \frac{1}{2\sqrt{2}} \ln x + \frac{1}{4} - \sqrt{(x+\frac{1}{4})^2-(\frac{3}{4})^2} \\ &= \sqrt{2x^2+x-1} + \frac{1}{2\sqrt{2}} \ln x + \frac{1}{4} - \sqrt{x^2+\frac{x}{2}-\frac{1}{2}} \end{aligned}$$

□

Not all integration's lead to pretty results, at least this didn't. But not all such are this messy

Example 14.44. Evaluate $\int \frac{3x-2}{x^2+4x+3} dx$

Solution. As the derivative of $x^2 + 4x + 3$ is $2x + 4$, let's write $3x - 2 = \frac{3}{2}(2x + 4) - 8$. We should also see that $x^2 + 4x + 3 = (x + 2)^2 - 1$, therefore:

$$\begin{aligned} & \int \frac{3x-2}{x^2+4x+3} dx \\ &= \frac{3}{2} \int \frac{2x+4}{x^2+4x+3} dx - 8 \int \frac{1}{(x+2)^2-1} dx \\ &= \frac{3}{2} \ln x^2 + 4x + 3 - 8 \frac{1}{2} \ln \frac{x+2-1}{x+2+1} \\ &= \frac{3}{2} \ln x^2 + 4x + 3 - 4 \ln \frac{x+1}{x+3} \end{aligned}$$

□

Sometimes we need to use two techniques together.

Example 14.45. Evaluate: $\int \frac{e^x}{\sqrt{e^{2x}+7e^x+1}} dx$

Proof. This one is quite simple, in comparison to what we had been doing till now. Simply let $t = e^x$ and therefore $dt = e^x dx$. And now its the questions funeral

$$\begin{aligned} & \int \frac{e^x}{\sqrt{e^{2x}+7e^x+1}} dx \\ &= \int \frac{1}{\sqrt{t^2+7t+1}} dt \\ &= \int \frac{1}{\sqrt{t^2+7t+\frac{49}{4}-\frac{49}{4}+1}} dt \\ &= \int \frac{1}{\sqrt{(t+\frac{7}{2})^2-\frac{45}{4}}} dt \\ &= \ln |t + \frac{7}{2} + \sqrt{t^2 + 7t + 1}| \\ &= \ln |e^x + \frac{7}{2} + \sqrt{e^{2x} + 7e^x + 1}| \end{aligned}$$

□

Here is one question which at the start seems hopeless but then crumbles to dust.

Example 14.46. Evaluate $\int \sqrt{\sec(x) - 1} dx$

Proof. The question would become so easy if we had a $\tan(x)$ or maybe $\sec^2(x)$ somewhere. Let's make them appear:

$$\begin{aligned} & \int \sqrt{\sec(x) - 1} dx \\ &= \int \sqrt{\sec(x) - 1} \cdot \sqrt{\frac{\sec(x)+1}{\sec(x)+1}} dx \\ &= \int \sqrt{\frac{\sec^2(x)-1}{\sec(x)+1}} dx \\ &= \int \frac{\tan(x)}{\sqrt{\sec(x)+1}} dx \end{aligned}$$

The $\sec(x)+1$ inside the radical seems nasty. Even if we take it as t , the integral seems messy. Let's do away with the polynomial and the radical in a single motion by $t^2 = \sec(x) + 1 \iff \sec(x) = t^2 - 1 \iff t = \sqrt{\sec(x) + 1}$, therefore $2tdt = \sec(x)\tan(x)dx \iff \tan(x)dx = \frac{2tdt}{\sec(x)} \iff \tan(x)dx = \frac{2tdt}{t^2-1}$, using this

$$\begin{aligned} & \therefore \int \frac{\tan(x)}{\sqrt{\sec(x)+1}} dx \\ &= \int \frac{2t}{t(t^2-1)} dt \\ &= \int \frac{2}{(t^2-1)} dt \\ &= 2 \int \frac{1}{(t^2-1)} dt \\ &= 2 \ln \frac{t-1}{t+1} \\ &= 2 \ln \frac{\sqrt{\sec(x)+1}-1}{\sqrt{\sec(x)+1}+1} \end{aligned}$$

□

We can also solve this using sine and cosine. I leave it to you to try and find that solution.

With that before we move on to the next method, here the final boss fight question:

Example 14.47. Evaluate: $\int \sqrt{\frac{\sin(x-a)}{\sin(x+a)}} dx$

Solution. We'll use the trigonometric formula for life that $\sin(A+B)\sin(A-B) = \cos^2(A) - \cos^2(B)$:

$$\begin{aligned} & \int \sqrt{\frac{\sin(x-a)}{\sin(x+a)}} dx \\ &= \int \frac{\sin(x-a)}{\sqrt{\cos^2(x)-\cos^2(a)}} dx \\ &= \int \frac{\sin(x)\cos(a)-\cos(x)\sin(a)}{\sqrt{\cos^2(x)-\cos^2(a)}} dx \\ &= \int \frac{\sin(x)\cos(a)}{\sqrt{\cos^2(x)-\cos^2(a)}} dx - \int \frac{\cos(x)\sin(a)}{\sqrt{\cos^2(x)-\cos^2(a)}} dx \end{aligned}$$

$$\begin{aligned}
&= \cos(a) \int \frac{\sin(x)}{\sqrt{\cos^2(x) - \cos^2(a)}} dx - \sin(a) \int \frac{\cos(x)}{\sqrt{\cos^2(x) - \cos^2(a)}} dx \\
&= -\cos(a) \int \frac{1}{\sqrt{t^2 - \cos^2(a)}} dt - \sin(a) \int \frac{\cos(x)}{\sqrt{1 - \sin^2(x) - 1 + \sin^2(a)}} dx \\
&= -\cos(a) \int \frac{1}{\sqrt{t^2 - \cos^2(a)}} dt - \sin(a) \int \frac{\cos(x)}{\sqrt{\sin^2(a) - \sin^2(x)}} dx \\
&= -\cos(a) \int \frac{1}{\sqrt{t^2 - \cos^2(a)}} dt - \sin(a) \int \frac{1}{\sqrt{\sin^2(a) - T^2}} dT \\
&= -\cos(a) \ln t - \sqrt{t^2 - \cos^2(a)} - \sin(a) \arcsin\left(\frac{T}{\cos(a)}\right) \\
&= -\cos(a) \ln \cos(x) - \sqrt{\cos^2(x) - \cos^2(a)} - \sin(a) \arcsin\left(\frac{\sin(x)}{\sin(a)}\right)
\end{aligned}$$

□

14.3.3 Trigonometric Substitution. We have already used it above, but here is the full thing. Sometimes it is better if we substitute some trigonometric form as t as that would make the question easier. Here are some forms which should be substituted by trigonometric functions

Theorem 14.48 (Trigonometric Substitution). (1) $a^2 - x^2$ or $\sqrt{a^2 - x^2} \rightarrow x = a \sin(\theta)$ or $a \cos(\theta)$

(2) $a^2 + x^2$ or $\sqrt{a^2 + x^2} \rightarrow x = a \tan(\theta)$ or $a \cot(\theta)$

(3) $x^2 - a^2$ or $\sqrt{x^2 - a^2} \rightarrow x = a \sec(\theta)$ or $a \csc(\theta)$

(4) $\sqrt{a+x}$, $\sqrt{a-x}$, $\sqrt{\frac{a+x}{a-x}}$ or $\sqrt{\frac{a-x}{a+x}}$ $\rightarrow x = a \cos(2\theta)$

(5) $\sqrt{\frac{x-a}{b-x}}$ or $\sqrt{(x-a)(b-x)}$ $\rightarrow x = a \cos^2 2\theta + b \sin^2(\theta)$

This doesn't require any proof as it is just using substitution a different way. Let's now use it to make short work of things that were considerably harder with standard integrals.

Example 14.49. Evaluate $\int \sqrt{\frac{x-1}{2-x}} dx$

Solution. Let's substitute $x = \cos^2(\theta) + 2 \sin^2(\theta) \iff x = 1 + \sin^2(\theta) \iff \theta = \arcsin(\sqrt{x-1})$ therefore $dx = \sin(2\theta)d\theta$. Using this:

$$\begin{aligned}
&\int \sqrt{\frac{x-1}{2-x}} dx \\
&= \int \sqrt{\frac{1+\sin^2(\theta)-1}{2-(1+\sin^2(\theta))}} d\theta \\
&= \int \sqrt{\frac{\sin^2(\theta)}{1-\sin^2(\theta)}} \sin(2\theta) d\theta \\
&= \int \sqrt{\frac{\sin^2(\theta)}{\cos^2(\theta)}} \sin(2\theta) d\theta \\
&= \int \tan(\theta) 2 \sin(\theta) \cos(\theta) d\theta
\end{aligned}$$

$$\begin{aligned}
 &= \int 2 \sin^2(\theta) d\theta \\
 &= \int 1 - \cos(2\theta) d\theta = \theta - \frac{\sin(2\theta)}{2} = \arcsin(\sqrt{x-1}) \frac{\sin(2 \arcsin(\sqrt{x-1}))}{2} = \arcsin(\sqrt{x-1}) \frac{2(\sqrt{x-1})}{\arcsin(\sqrt{x-1}) \sqrt{(2-x)(x-1)}} = \arccos(\sqrt{2-x}) \sqrt{(2-x)(x-1)} \quad \square
 \end{aligned}$$

We could also do this by substitution by rationalizing, but with trig, it almost solves itself.

Here are some more useful integrals which can be derived using trigonometric substitution. The proof is quite basic and hence, I expect that you'll be able to do that yourself.

Theorem 14.50. (1) $\int \sqrt{a^2 - x^2} dx = \frac{1}{2}x\sqrt{a^2 - x^2} + \frac{1}{2}a^2 \arcsin \frac{x}{a}$

(2) $\int \sqrt{a^2 + x^2} dx = \frac{1}{2}x\sqrt{a^2 + x^2} + \frac{1}{2}a^2 \ln x + \sqrt{x^2 + a^2}$

(3) $\int \sqrt{x^2 - a^2} dx = \frac{1}{2}x\sqrt{x^2 - a^2} - \frac{1}{2}a^2 \ln x + \sqrt{x^2 - a^2}$

14.3.4 Integration Trigonometric Functions. While we were seeing trig functions since the start of this chapter, we'll be studying about only their integration in this subsection.

First recall that $\sin^2(x) = \frac{1-\cos(2x)}{2}$ and $\cos^2(x) = \frac{1+\cos(2x)}{2}$ from the formula's for life table. We'll be using them almost in all the questions in this section.

Theorem 14.51. $\int \cos^2(\theta) d\theta = \frac{\theta}{2} + \frac{\sin(2x)}{4}$

$$\int \sin^2(\theta) d\theta = \frac{\theta}{2} + \frac{\cos(2x)}{4}$$

I think you are more than capable now in proving both of these.

Example 14.52. Evaluate $\int \cos^4(x) dx$

Solution.

$$\begin{aligned}
 &\int \cos^4(x) dx \\
 &= \int \left(\frac{1+\cos(2x)}{2}\right)^2 dx \\
 &= \int \frac{1+2\cos(2x)+\cos^2(2x)}{4} dx \\
 &= \frac{1}{4} \int 1 + 2\cos(2x) + \cos^2(4x) dx \\
 &= \frac{1}{4} \int 1 + 2\cos(2x) + \cos^2(4x) dx \\
 &= \frac{1}{4} \int 1 + 2\cos(2x) + \frac{1+\cos(4x)}{2} dx \\
 &= \frac{1}{4} \int \frac{3}{2} + 2\cos(2x) + \frac{\cos(4x)}{2} dx \\
 &= \frac{1}{4} \left(\frac{3x}{2} + \sin(2x) \frac{\sin(x)}{8} \right)
 \end{aligned}$$

□

And here is one for you to solve:

Example 14.53. Evaluate $\int \sin^4(x)dx$

And on while we are on the topic of powers of sine and cosine:

Example 14.54. Evaluate $\int \sin^3(x)dx$

Solution. Remember the formula's for life which had $\sin^3(x)$?

$$\begin{aligned} & \int \sin^3(x)dx \\ &= \int \frac{3\cos(x)-\cos(3x)}{4} dx \\ &= \frac{1}{4} \int 3\cos(x) - \cos(3x) dx \\ &= \frac{1}{4} \left(3\sin(x) - \frac{\sin(3x)}{3} \right) \end{aligned}$$

□

You know the drill by now...

Example 14.55. Evaluate $\int \cos^3(x)dx$

Now let's use another trig formula to break a complicated looking question

Example 14.56. Evaluate: $\int \cos(x) \cos(2x) \cos(3x)dx$

Solution. Remember the product sum formulas? We will use them here.

$$\begin{aligned} & \int \cos(x) \cos(2x) \cos(3x)dx \\ &= \frac{1}{2} \int (\cos(3x) + \cos(x)) \cos(3x) dx \\ &= \frac{1}{2} \int \cos^2(3x) + \cos(x) \cos(3x) dx \\ &= \frac{1}{4} \int 1 + \cos(6x) + \cos(4x) + \cos(2x) dx \\ &= \frac{1}{4} \left(x + \frac{\sin(6x)}{6} + \frac{\sin(4x)}{4} + \frac{\sin(2x)}{2} \right) \end{aligned}$$

□

This works for sine and cosine but we don't have anything of this sort tan then what do we do if:

Example 14.57. Evaluate: $\int \tan(x) \tan(2x) \tan(3x)dx$

Solution. While we don't know any product sum formulas for tan, we do know the formulation for $\tan(A + B)$ and we can clearly see $x + 2x = 3x$

$$\begin{aligned} & \therefore \tan(x + 2x) = \tan(3x) \\ & \iff \frac{\tan(x) + \tan(2x)}{1 - \tan(x)\tan(2x)} = \tan(3x) \\ & \iff \tan(x) + \tan(2x) = \tan(3x) - \tan(x)\tan(2x)\tan(3x) \\ & \iff \tan(x)\tan(2x)\tan(3x) = \tan(3x) - \tan(2x) - \tan(x) \end{aligned}$$

And this converts our integral to:

$$\begin{aligned} & \int \tan(x) \tan(2x) \tan(3x) dx \\ &= \int \tan(3x) - \tan(2x) - \tan(x) dx \\ &= \frac{\ln |\sec(3x)|}{3} - \frac{\ln |\sec(2x)|}{2} - \ln |\sec(x)| \end{aligned}$$

□

Let's now make things a bit more crunchy!

Example 14.58. Evaluate $\int \frac{1}{\cos(x-a)\cos(x-b)} dx$

Solution. We can try using the product sum but that will be to no avail. What else can we do?

If we had $\sin((x-a)-(x-b))$ as the numerator, the question would simplify a lot. Notice that $\sin((x-a)-(x-b)) = \sin(b-a)$ which is a constant.

$$\begin{aligned} & \therefore \int \frac{1}{\cos(x-a)\cos(x-b)} dx \\ &= \frac{1}{\sin(b-a)} \int \frac{\sin((x-a)-(x-b))}{\cos(x-a)\cos(x-b)} dx \\ &= \frac{1}{\sin(b-a)} \int \frac{\sin(x-a)\cos(x-b)-\cos(x-a)\sin(x-b)}{\cos(x-a)\cos(x-b)} dx \\ &= \frac{1}{\sin(b-a)} \int \tan(x-a) - \tan(x-b) dx \\ &= \frac{1}{\sin(b-a)} (\ln |\sec(x-a)| - \ln |\sec(x-b)|) \end{aligned}$$

□

This example is the reason why a lot of people hate integral calculus. How were you supposed to think about multiplying by $\sin(b-a)$ of all thing?

The answer is practice. There is no way to make sure you see these patterns then to practice them.

Let's now talk about another common configuration:

Theorem 14.59. If we wish to find $\int \frac{\cos(x)+\sin(x)}{f(\sin(2x))} dx$ we take $\cos(x)-\sin(x) = t$

If we wish to find $\int \frac{\cos(x)-\sin(x)}{f(\sin(2x))} dx$ we take $\cos(x)+\sin(x) = t$

Let's just do an example to understand what this substitution wishes to say:

Example 14.60 (Motivating Example). Evaluate: $\int \frac{\cos(x)-\sin(x)}{\sqrt{\sin(2x)}} dx$

Solution. We'll put $\cos(x)+\sin(x) = t$ and therefore, $(\cos(x)-\sin(x))dx = dt$ and $t^2 = \cos^2(x)+\sin^2(x)+2\sin(x)\cos(x) = 1+\sin(2x) \iff \sin(2x) = t^2-1$; using all of this,

$$\int \frac{\cos(x)-\sin(x)}{\sqrt{\sin(2x)}} dx$$

$$\begin{aligned}
 &= \int \frac{1}{\sqrt{t^2-1}} dt \\
 &= \ln t + \sqrt{t^2 - 1} \\
 &= \ln \cos(x) + \sin(x) + \sqrt{\sin(2x)}
 \end{aligned}$$

□

But sometimes the form is not so direct...

Example 14.61. Evaluate: $\int \frac{1}{1+\cot(x)} dx$

Proof. $\int \frac{1}{1+\cot(x)} dx$

$$\begin{aligned}
 &= \int \frac{\sin(x)}{\sin(x)+\cos(x)} dx \text{ We see } \sin(x) + \cos(x) \text{ in the denominator which makes us} \\
 &\quad \text{want to substitute } t = \sin(x) + \cos(x), \text{ therefore } dt = (\cos(x) - \sin(x))dx. \\
 \therefore \int \frac{\sin(x)}{\sin(x)+\cos(x)} dx &= \frac{1}{2} \int \frac{\sin(x)+\cos(x)+\sin(x)-\cos(x)}{\sin(x)+\cos(x)} dx = \frac{1}{2}(x + \int \frac{\sin(x)-\cos(x)}{\sin(x)+\cos(x)} dx) = \frac{1}{2}(x + - \int \frac{1}{t} dt) \\
 &= \frac{1}{2}(x + - \ln t) \\
 &= \frac{1}{2}(x + - \ln \sin(x) + \cos(x))
 \end{aligned}$$

□

Example 14.62. Evaluate $\int \frac{\sin(x)}{\cos(3x)} dx$

Solution. This question has a very arbitrary multiplication, which you may not like...

$$\begin{aligned}
 &\int \frac{\sin(x)}{\cos(3x)} dx \\
 &= \int \frac{\cos(x) \sin(x)}{\cos(x) \cos(3x)} dx \\
 &= \frac{1}{2} \int \frac{\sin(2x)}{\cos(x) \cos(3x)} dx \\
 &= \frac{1}{2} \int \frac{\sin(3x-x)}{\cos(x) \cos(3x)} dx \\
 &= \frac{1}{2} \int \frac{\sin(3x) \cos(x) - \sin(x) \cos(3x)}{\cos(x) \cos(3x)} dx \\
 &= \frac{1}{2} \int \tan(3x) - \tan(x) dx \\
 &= \frac{1}{2} \left(\frac{\ln |\sec(3x)|}{3} - \ln |\sec(x)| \right) \text{ Here the } \cos(x) \text{ multiplication was motivated by} \\
 &\quad \text{the fact that generally trig integrals with } x, 2x, 3x \text{ all can be simplified.}
 \end{aligned}$$

Let's do one simple question to close this subsection:

Example 14.63. Evaluate $\int \frac{\sqrt{\tan(x)}}{\sin(x) \cos(x)} dx$

Solution. We'll use the great rule: Whenever I see $\tan(x)$, I want $\sec^2(x)$.

$$\begin{aligned}
 &\int \frac{\sqrt{\tan(x)}}{\sin(x) \cos(x)} dx \\
 &= \int \frac{\sqrt{\tan(x)}}{\frac{\sin(x)}{\cos(x)} \cos^2(x)} dx
 \end{aligned}$$

$$\begin{aligned}
 &= \int \frac{\sec^2(x) \sqrt{\tan(x)}}{\tan(x)} dx \\
 &= \int \frac{1}{\sqrt{t}} dt \\
 &= 2\sqrt{t} \\
 &= 2\sqrt{\tan(x)}
 \end{aligned}$$

□

14.3.5 Integration by Partial Fractions. We have studied the exercise of breaking a function into partial fractions in sequence and series. That's exactly what we do before applying the standard forms and getting an answer. Let's just do a token example before doing some real fun with it:

Example 14.64. Evaluate:

$$\int \frac{x^2}{(x-1)(x-2)(x-3)}$$

Solution. Through partial fractions we know that for some A, B, C , we have $A(x-2)(x-3) + B(x-1)(x-3) + C(x-1)(x-2) = x^2$, setting $x = 1, 2, 3$ we get $A = \frac{1}{2}$; $B = -4$; $C = \frac{9}{2}$

. The integral transforms into:

$$\begin{aligned}
 &\int \frac{x^2}{(x-1)(x-2)(x-3)} \\
 &= \frac{1}{2(x-1)} - \frac{4}{x-2} + \frac{9}{2(x-3)} dx \\
 &= \frac{1}{2} \ln(|x-1|) - 4 \ln|x-2| + \frac{9}{2} \ln|x-3|
 \end{aligned}$$

□

Sometimes the partial fraction is disguised as trigonometry...

Example 14.65. Evaluate: $\int \frac{1}{\sin(x)+\sin(2x)} dx$

$$\begin{aligned}
 &\text{Solution. } \int \frac{1}{\sin(x)+\sin(2x)} dx \\
 &= \int \frac{1}{\sin(x)+2\sin(x)\cos(x)} dx \\
 &= \int \frac{1}{\sin(x)(1+2\cos(x))} dx \\
 &= \int \frac{\sin(x)}{\sin^2(x)(1+2\cos(x))} dx \\
 &= \int \frac{\sin(x)}{(1-\cos^2(x))(1+2\cos(x))} dx \\
 &= \int \frac{1}{(1-t^2)(1+2t)} dt \\
 &= \int \frac{1}{(1-t)(1+t)(1+2t)} dt \\
 &= \frac{\ln|1-t|}{6} + \frac{\ln|1+t|}{4} + \frac{4\ln|1+2t|}{3}
 \end{aligned}$$

□

... other times as exponentials.

Example 14.66. Evaluate: $\int \frac{1}{(e^x-1)^2} dx$

$$\begin{aligned}
 & \text{Solution. } \int \frac{1}{(e^x - 1)^2} dx \\
 &= \int \frac{e^x}{e^x(e^x - 1)^2} dx \\
 &= \int \frac{1}{t(t-1)^2} dt \\
 &= \int \frac{1}{t} dt - \int \frac{1}{t-1} dt + \int \frac{1}{(t-1)^2} dt \\
 &= \ln t - \ln t - 1 - \frac{1}{(t-1)} \\
 &= x - \ln e^x - 1 - \frac{1}{e^x - 1}
 \end{aligned}$$

□

As you can see partial fractions are normally quite simple. However, there is one particular configuration where they are a bit harder to see:

Theorem 14.67. For $\int \frac{x^2 \pm a}{x^4 + kx^2 + a^2}$ where k is a constant, divide the numerator and denominator by x^2 and then take $x \mp \frac{a}{x} = t$

Let's try a motivating example to see what the theorem wants to say:

Example 14.68 (Motivating Example). Evaluate $\int \frac{x^2 - 1}{x^4 + x^2 + 1} dx$

Solution. The given question fits the configuration. This leads us to dividing numerator and denominator by x^2 .

$$\begin{aligned}
 & \therefore \int \frac{x^2 - 1}{x^4 + x^2 + 1} dx \\
 &= \int \frac{1 - \frac{1}{x^2}}{x^2 + 1 + \frac{1}{x^2}} dx \text{ Now let } x + \frac{1}{x} = t, \text{ therefore } 1 - \frac{1}{x^2} dx = dt \text{ and } x^2 + \frac{1}{x^2} + 2 = t^2 \\
 & \therefore \int \frac{1 - \frac{1}{x^2}}{x^2 + 1 + \frac{1}{x^2}} dx = \int \frac{1}{t^2 - 1} dt \\
 &= \frac{1}{2} \ln \frac{t-1}{t+1} \\
 &= \frac{1}{2} \ln \frac{x + \frac{1}{x} - 1}{x + \frac{1}{x} + 1} \\
 &= \frac{1}{2} \ln \frac{x^2 + 1 - x}{x^2 + 1 + x}
 \end{aligned}$$

□

Now let's go through some final configurations before we are done with integration by substitution

14.3.6 Some non-intuitive substitutions.

Theorem 14.69. $\int \frac{1}{(px+q)\sqrt{ax+b}} dx; \int \frac{1}{(px^2+qx+r)\sqrt{ax+b}} dx$ Substitute $ax+b \rightarrow t^2$
 $\int \frac{1}{(px+1)\sqrt{ax^2+bx+c}} \text{ substitute } px+q = \frac{1}{t}$
 $\int \frac{1}{(px^2+q)\sqrt{ax^2+b}} \text{ substitute } x = \frac{1}{t}$

Let's just see one example of each before moving on.

Example 14.70 (Motivating Example). Evaluate $\int \frac{1}{(x-4)\sqrt{x+5}} dx$

Solution. Take $x + 5 = t^2$ and therefore $dx = 2tdt$. This transforms the question into:

$$\begin{aligned} & \int \frac{1}{(x-4)\sqrt{x+5}} dx \\ &= \int \frac{2t}{(t^2-9)t} dt \\ &= \int \frac{2}{(t^2-9)} dt \\ &= 2 \int \frac{1}{(t^2-9)} dt \\ &= 2 \frac{1}{6} \ln \frac{t-3}{t+3} \\ &= \frac{1}{3} \ln \frac{\sqrt{x+5}-3}{\sqrt{x+5}+3} \end{aligned}$$

□

Example 14.71 (Motivating Example). Evaluate $\int \frac{1}{(x^2+3x+2)\sqrt{x+4}} dx$

Proof. Take $x + 4 = t \iff x = t - 4$ and therefore $dx = 2tdt$. This transforms the question into:

$$\begin{aligned} & \int \frac{1}{(x^2+3x+2)\sqrt{x+4}} dx \\ &= \int \frac{2t}{((t-4)^2+3(t-4)+2)t} dt \\ &= 2 \int \frac{1}{t^2-8t+16+3t-12+2} dt \\ &= 2 \int \frac{1}{t^2-5t+6} dt \\ &= 2 \int \frac{1}{(t-2)(t-3)} dt \\ &= 2 \int \frac{(t-2)-(t-3)}{(t-2)(t-3)} dt \\ &= 2(\ln t - 3 - \ln t - 2) \\ &= 2 \ln \frac{x+1}{x+2} \end{aligned}$$

□

Example 14.72 (Motivating example). Evaluate $\int \frac{1}{(x-2)\sqrt{x^2-4}} dx$

Solution. Let $x - 2 = \frac{1}{t} \iff x = \frac{2t+1}{t}$ and $dx = \frac{-1}{t^2} dt$, now the equation evolves to:

$$\begin{aligned} & \int \frac{1}{(x-2)\sqrt{x^2-4}} dx \\ &= \int \frac{\frac{-1}{t^2}}{\frac{1}{t}\sqrt{((\frac{2t+1}{t})^2-4}} dt \\ &= \int \frac{\frac{-1}{t}}{\sqrt{(\frac{4t^2+4t+1}{t^2})-4}} dt \\ &= \int \frac{\frac{-1}{t}}{\sqrt{(\frac{4t^2+4t-3}{t^2})}} dt \\ &= - \int \frac{1}{\sqrt{4t^2+4t-3}} dt \\ &= -2 \frac{\sqrt{4t+1}}{4} = -\frac{\sqrt{4t+1}}{2} = -\frac{\sqrt{\frac{4}{x-2}+1}}{2} = -\frac{\sqrt{\frac{x+2}{x-2}}}{2} \end{aligned}$$

□

Example 14.73. Evaluate $\int \frac{1}{(1+x^2)\sqrt{1-x^2}} dx$

Solution. We'll take $x = \frac{1}{t} \iff t = \frac{1}{x}$ and therefore $dx = -\frac{1}{t^2} dt$, the equation converts to:

$$\begin{aligned} & \int \frac{1}{(1+x^2)\sqrt{1-x^2}} dt \\ &= \int \frac{\frac{-1}{t^2}}{(1+\frac{1}{t^2})\sqrt{1-\frac{1}{t^2}}} dt \\ &= - \int \frac{t}{t^2+1\sqrt{t^2-1}} dt \end{aligned}$$

We substitute for the second time, $t^2-1=y^2 \iff y=\sqrt{t^2-1}$ and therefore $2tdt=2ydy \iff tdt=ydy$, hence we can say:

$$\begin{aligned} &= - \int \frac{t}{t^2+1\sqrt{t^2-1}} dt \\ &= - \int \frac{y}{(y^2+2)y} dy \\ &= - \int \frac{1}{(y^2+2)} dy \\ &= -\frac{1}{\sqrt{2}} \arctan\left(\frac{y}{\sqrt{2}}\right) \\ &= -\frac{1}{\sqrt{2}} \arctan\left(\frac{\sqrt{t^2-1}}{\sqrt{2}}\right) \\ &= -\frac{1}{\sqrt{2}} \arctan\left(\frac{\sqrt{(\frac{1}{x})^2-1}}{\sqrt{2}}\right) \end{aligned}$$

□

A very famous, and somewhat sneaky substitution is the universal trigonometric substitution or its misnomer Weierstrass substitution (was discovered by Lagrange, Weierstrass had no relation whatsoever)

Theorem 14.74 (Universal Trigonometric Substitution(UTS)). For $\int \frac{1}{a \sin(x)+b \cos(x)+c} dx$ we will substitute $t = \tan(\frac{x}{2})$ and therefore $\sin x = \frac{2t}{1+t^2}$, $\cos x = \frac{1-t^2}{1+t^2}$, and $dx = \frac{2}{1+t^2} dt$

This may seem somewhat complicated and unnecessary but it finishes highly trigonometric questions with relative ease, and without arbitrary multiplication. Let's see the second proof for the integral of $\sec(x)$ and $\csc(x)$

Proof. Using UTS $\int \sec(x) dx$

$$\begin{aligned} &= \int \frac{1}{\cos(x)} dx \\ &= \int \frac{1+t^2}{1-t^2} \frac{2}{1+t^2} dt \\ &= 2 \int \frac{1}{1-t^2} dt \\ &= 2 \frac{1}{2} \ln \left| \frac{1+t}{1-t} \right| \\ &= \ln \left| \tan\left(\frac{x}{2} + \frac{\pi}{4}\right) \right| \text{ The last step came from the formula for life for } \tan(a+b) = \frac{\tan(a)+\tan(b)}{1-\tan(a)\tan(b)} \end{aligned}$$

□

I imagine that you have already started to scribble in the margins to prove the same for $\csc(x)$.

The UTS is like life with cheat codes as it can solve any trigonometric integral(provided that you are able to solve the algebraic integral), while it is best used with the configuration prescribed, it can solve the other cases as well. Here is token example:

Example 14.75. Evaluate $\int \frac{1}{5+4\cos(x)} dx$

Solution. Using UTS,

$$\int \frac{1}{5+4\cos(x)} dx = \int \frac{1}{5+4\frac{1-t^2}{1+t^2}} \frac{2}{1+t^2} dt = \int \frac{1+t^2}{5+5t^2+4-4t^2} \frac{2}{1+t^2} dt = 2 \int \frac{1}{t^2+9} dt = \frac{2}{3} \arctan\left(\frac{t}{3}\right) = \frac{2}{3} \arctan\left(\frac{\tan\left(\frac{x}{2}\right)}{3}\right) \quad \square$$

The next two substitutions are seen very less, however, I have decided to include them in order to be more than sure that all substitution tricks/methods were present.

Theorem 14.76. For $\int \frac{1}{a\cos^2(x)+b\sin^2(x)+c\sin(x)\cos(x)} dx$ divide the numerator and denominator by $\cos^2(x)$ in order to take $\tan(x) = t$ and then solve.

Theorem 14.77. For $\int \frac{p\cos(x)+q\sin(x)+r}{a\cos(x)+b\sin(x)+c} dx$ we will try to express the numerator N as $N = \alpha D + \beta D' + \gamma$ where D is the denominator function.

We will do one token example each for both of them:

Example 14.78. Evaluate: $\int \frac{1}{5\cos^2(x)-4\cos(x)\sin(x)-2\sin^2(x)} dx$

$$\begin{aligned} & \text{Solution. } \int \frac{1}{5\cos^2(x)-4\cos(x)\sin(x)-2\sin^2(x)} dx \\ &= \int \frac{\sec^2(x)}{5-4\tan(x)-2\tan^2(x)} dx \\ &= \int \frac{1}{5-4t-2t^2} dt \\ &= \frac{-1}{2} \int \frac{1}{t^2+2t-\frac{5}{2}} dt \\ &= \frac{-1}{2} \int \frac{1}{(t+1)^2-\frac{7}{2}} dt \\ &= \frac{-1}{2} \frac{1}{2\sqrt{\frac{7}{2}}} \ln \frac{t+1-\sqrt{\frac{7}{2}}}{t+1+\sqrt{\frac{7}{2}}} = \frac{-1}{2\sqrt{14}} \ln \frac{t+1-\sqrt{\frac{7}{2}}}{t+1+\sqrt{\frac{7}{2}}} \end{aligned} \quad \square$$

Not a pretty expression... Let's hope the next one is not as awful

Example 14.79. Evaluate: $\int \frac{2\sin(x)-\cos(x)+1}{7\sin(x)+2\cos(x)+2} dx$

Solution. Let $2\sin(x) - \cos(x) + 1 = (7\sin(x) + 2\cos(x) + 2)a + (7\cos(x) - 2\sin(x))b + c$

We can by comparing the coefficients say that: $7a - 2b = 2$; $2a + 7b = -1$; $2a + c = 1$

Solving these will give us $a = \frac{12}{53}$; $b = -\frac{1}{53}$; $c = \frac{29}{53}$

Using this we get:

$$\begin{aligned} & \int \frac{2\sin(x)-\cos(x)+1}{7\sin(x)+2\cos(x)+2} dx \\ &= \int \frac{\frac{12}{53}(7\sin(x)+2\cos(x)+2) - \frac{11}{53}(7\cos(x)-2\sin(x)) + \frac{29}{53}}{7\sin(x)+2\cos(x)+2} dx \\ &= \int \frac{\frac{12}{53}dx}{7\sin(x)+2\cos(x)+2} - \int \frac{\frac{11}{53}(7\cos(x)-2\sin(x))}{7\sin(x)+2\cos(x)+2} dx + \int \frac{\frac{29}{53}}{7\sin(x)+2\cos(x)+2} dx \\ &= \frac{12x}{53} - \frac{11}{53} \ln|7\sin(x) + 2\cos(x) + 2| + \frac{29}{53} \int \frac{1}{7\sin(x)+2\cos(x)+2} dx \\ &= \frac{12x}{53} - \frac{11}{53} \ln|7\sin(x) + 2\cos(x) + 2| + \frac{29}{53} \int \frac{1}{7t+1} dt \\ &= \frac{12x}{53} - \frac{11}{53} \ln|7\sin(x) + 2\cos(x) + 2| + \frac{29}{53} \frac{\ln|7t+2|}{7} \\ &= \frac{12x}{53} - \frac{11}{53} \ln|7\sin(x) + 2\cos(x) + 2| + \frac{29}{53} \frac{\ln|7t+2|}{7} \end{aligned}$$

□

I knew a man who said every integral is beautiful. He died of a heart attack after seeing this one.

14.4 Techniques of indefinite Integration: Integration by Parts

Using substitutions and the many, many types of it; we could integrate huge swath of functions. However, we still fail to answer a very simple $\int \ln x dx$ or a more complicated $\int x^3 e^x dx$. What now?

Theorem 14.80 (Integration by Parts). $\int u dv = uv - \int v du$

This may seem strange, but here is it in action (we'll prove it in a while):

Example 14.81 (Motivating Example). (a) $\int xe^{3x} dx = ?$

(b) $\int x \sin(x) dx = ?$

Solution. (a) Let $u = x$ and $e^{3x} dx = dv$, therefore $du = 1 dx$ and $v = \frac{e^{3x}}{3}$, using this:

$$\begin{aligned} \int xe^{3x} dx &= x \frac{e^{3x}}{3} - \int \frac{e^{3x}}{3} dx \\ &= \frac{xe^{3x}}{3} - \frac{e^{3x}}{9} \end{aligned}$$

If you got this, I ask you to try (b) out before you read through my solution.

Done? Here's what I did

(b) $u = x$ and $dv = \sin(x)dx$, therefore $du = dx$ and $v = -\cos(x)$. Using these, we can say:

$$\begin{aligned} \int x \sin(x)dx &= -x \cos(x) - \int -\cos(x)dx \\ &= -x \cos(x) + \int \cos(x)dx \\ &= -x \cos(x) + \sin(x) \end{aligned}$$

□

You might have got it by now. Now here is the proof for why this must work:

Proof. We know from the product rule of differentiation:

$d(uv) = u dv + v du$, Integrating on both sides gives us

$\Leftrightarrow uv = \int u dv + \int v du$, which rearranges to

$\Leftrightarrow \int u dv = uv - \int v du$

□

Sometimes, however you'll need to do this process twice or thrice because of the nature of $\int v du$ in which case the chances of things going wrong increases many fold(the sign of the integral keeps on changing). So I'll share the secret method I use instead, it is faster, better, safer and easier. The only problem is that in a subjective examination, the paper checker may not allow this. So in those cases be a little careful.

Theorem 14.82 (The DI method). *Divide the function $f(x)g(x)$ into two parts, a D say $f(x)$ and I say $g(x)$. The D is easy to differentiate and the I is easy to integrate. We write this in this in columns like this:*

We do this till we hit one of the three stops:

s	D	I
+	$f(x)$	$g(x)$
-	$f'(x)$	$\int g(x)$
+	$f''(x)$	$\int \int g(x)$
⋮	⋮	⋮

- (1) *The product of the D and I of the same row becomes 0 in which case we just cross multiple(D of one row to the I of the row below it and add it to the solution keeping the sign in mind)*

- (2) *The product of the D and I of the same row become a function we can integrate, we cross multiply for all rows above them and then add the integration of that function.*
- (3) *The product of the D and I of a row become the same function we were integrating (with negative sign probably). Then we cross multiply till that row and then divide the whole thing by two.*

This may seem more complicated than it actually is. Here is the DI method in action:

Example 14.83 (Motivating Example). $\int x^2 \cos(2x) dx = ?$

Solution. We take $D = x^2$ and $I = \cos(2x)$ We have hit the first stop and

$$\begin{array}{c|cc} s & D & I \\ + & x^2 & \cos(2x) \\ - & 2x & \frac{1}{2} \sin(2x) \\ + & 2 & \frac{-1}{4} \cos(2x) \\ - & 0 & \frac{-1}{8} \sin(2x) \end{array}$$

therefore, the integral is:

$$x^2 \cdot \frac{1}{2} \sin(2x) - 2x \cdot \frac{-1}{4} \cos(2x) + 2 \cdot \frac{-1}{8} \sin(2x) \\ = \frac{x^2 \sin(2x)}{2} + \frac{x \cos(2x)}{2} - \frac{\sin(2x)}{4}$$

□

I dare you to try to solve it with the usual method, without either taking too long or making some stupid error. The DI method on the other hand makes quick work of all this...

Example 14.84 (Motivating Example). $\int \arcsin(x) dx = ?$

Solution. We take $D = \arcsin(x)$ and $I = 1$ We see the second stop here and

$$\begin{array}{c|cc} s & D & I \\ + & \arcsin(x) & 1 \\ - & \frac{1}{1+x^2} & x \end{array}$$

write the answer simply as:

$$x \arcsin(x) - \int \frac{x}{1+x^2} dx \\ = x \arcsin(x) - \frac{1}{2} \ln 1 + x^2 dx$$

□

And the final example:

Example 14.85 (Motivating Example). $\int e^{-x} \cos(x) dx = ?$

Solution. We take $D = e^{-x}$ and $I = \cos(x)$ And we have hit the third stop.

$$\begin{array}{c|cc} s & D & I \\ + & e^{-x} & \cos(x) \\ - & -e^{-x} & \sin(x) \\ + & +e^{-x} & -\cos(x) \end{array}$$

We can now write the answer as:

$$\begin{aligned} & \frac{1}{2}(e^{-x} \sin(x) - (-e^{-x})(-\cos(x))) \\ &= \frac{1}{2}(e^{-x} \sin(x) - (e^{-x})(\cos(x))) \\ &= \frac{e^{-x}}{2}(\sin(x) - \cos(x)) \end{aligned} \quad \square$$

If you have understood this technique I expect the below example be a child's play:

Example 14.86. (a) $\int \ln x dx$
 (b) $\int \sec^3(x) dx$
 (c) $\int e^{2x} \sin(3x) dx$

Here are the final few configurations for indefinite integration:

Theorem 14.87. (1) $\int e^{ax} \cos(bx) dx = \frac{e^{ax}}{a^2+b^2}(a \cos(bx) + b \sin(bx))$

$$(2) \int e^{ax} \sin(bx) dx = \frac{e^{ax}}{a^2+b^2}(a \sin(bx) - b \cos(bx))$$

$$(3) \int e^x(f(x) + f'(x)) dx = e^x f(x)$$

$$(4) xf'(x) + f(x) dx = xf(x)$$

I normally despise doing this, but I want to leave these proofs as exercise for you to solve. As they are standard functions, you can obviously use google but I really wish you don't need to do that.

At this point you may feel that every integral is solvable and maybe will try to solve for the indefinite integral of any and every function you encounter(I used to do this a lot), but we can't integrate everything. Many integrals such as $\int e^{x^2} dx$ are famously hard(known as the error

function). While we can compute integrals for $\ln x$, $\sin(x)$, $\cos(x)$, $x \sin(x)$, $\sin^2(x)$ we fail to compute of $\int \ln \ln x dx$, $\int \frac{\sin(x)}{x} dx$, $\int \frac{1}{\ln x} dx$, $\int \frac{1}{xe^x} dx$, $\int \sin(x^2) dx$, $\int \cos(x^2) dx$ and many many more functions. The why and how for each of these functions is a topic I encourage you to explore on your own as that touches. However, It will not be covered in this book as a lot of the required math us way beyond the scope of any single book.

14.5 Definite integration

In indefinite integration, we've become masters at finding anti derivatives, or what you might call the 'net change' in a quantity. We've been able to calculate functions that, when differentiated, give us the original function we started with. This is a powerful tool for understanding how quantities change over an interval or across a curve

Now, with definite integration, we're going to focus on a more specific question: How can we calculate the 'net change' or the 'accumulated total' of a quantity over a given interval? Think of it as going from knowing how a car's speed changes at every instant (indefinite integration) to finding out how far the car has traveled during a specific time frame (definite integration)

It is basically the area under the curve of the graph as we had proven using the fundamental theorem of calculus.

Reiterating,

Theorem 14.88 (Fundamental Theorem of Calculus). *If $\int f(x)dx = F(x)$ then $\int_a^b f(x)dx = F(b) - F(a)$*

With only this, we can convert almost every indefinite integration problem into a definite integration problem by slapping on two limits.

However, that will not be discussed here as that is just an answer procurement method. Actual definite integration refers to integration of an otherwise impossible to integrate function over some limit.

Theorem 14.89 (Some properties of the integral). (1) $\int_a^b f(x)dx = \int_a^b f(t)dt$

$$(2) \int_a^b f(x)dx = - \int_b^a f(x)dx$$

$$(3) \int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx \text{ for } c \in \mathbb{R}$$

All of these seem to be trivial. So here is another simple(but powerful) manipulation:

14.6 King's Rule

Theorem 14.90 (King's Rule). $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

The rule is just an elementary manipulation of the variable using the 1st basic property of the integral.

Let's take it for a spin:

Example 14.91. $\int_0^{\frac{\pi}{2}} \frac{\cos(x)}{\cos(x)+\sin(x)} dx$

Solution. Using the King's rule:

$$\begin{aligned} I &= \int_0^{\frac{\pi}{2}} \frac{\cos(x)}{\cos(x)+\sin(x)} dx \\ \iff I &= \int_0^{\frac{\pi}{2}} \frac{\cos(\frac{\pi}{2}-x)}{\cos(\frac{\pi}{2}-x)+\sin(\frac{\pi}{2}-x)} dx \\ \iff I &= \int_0^{\frac{\pi}{2}} \frac{\sin(x)}{\sin(x)+\cos(x)} dx \end{aligned}$$

Adding this to the original equation gives us:

$$\begin{aligned} 2I &= \int_0^{\frac{\pi}{2}} \frac{\cos(x)}{\cos(x)+\sin(x)} dx + \int_0^{\frac{\pi}{2}} \frac{\sin(x)}{\sin(x)+\cos(x)} dx \\ \iff 2I &= \int_0^{\frac{\pi}{2}} \frac{\sin(x)+\cos(x)}{\sin(x)+\cos(x)} dx \\ \iff 2I &= \int_0^{\frac{\pi}{2}} dx \\ \iff 2I &= \frac{\pi}{2} \\ \iff I &= \frac{\pi}{4} \end{aligned}$$

□

That made quick work of something we would otherwise have to integrate, get a bad answer and then substitute.

Here is another example for you:

Example 14.92. $\int_0^{\frac{\pi}{2}} \frac{\sqrt{\tan(x)}}{\sqrt{\cot(x)}+\sqrt{\tan(x)}} dx$

And if you feel this is too easy to actually come in a paper...

Example 14.93. (JEE Mains 2015) $\int_2^4 \frac{\ln x^2}{\ln(x^2)+\ln(36-12x+x^2)} dx$

Solution. Using the kings rule:

$$\begin{aligned} I &= \int_2^4 \frac{\ln x^2}{\ln x^2+\ln 36-12x+x^2} dx \\ \iff I &= \int_2^4 \frac{\ln(6-x)^2}{\ln(6-x)^2+\ln 36-12(6-x)+(6-x)^2} dx \end{aligned}$$

$$\iff I = \int_2^4 \frac{\ln 36 - 12x + x^2}{\ln 36 - 12x + x^2 + \ln 36 - 72 + 12x + 36 - 12x + x^2} dx \iff I = \int_2^4 \frac{\ln 36 - 12x + x^2}{\ln 36 - 12x + x^2 + \ln x^2} dx$$

Adding to the first equation:

$$2I = \int_2^4 \frac{\ln 36 - 12x + x^2 + \ln x^2}{\ln 36 - 12x + x^2 + \ln x^2} dx$$

$$2I = 2$$

$$I = 1$$

□

And once in a while, the problem writer tries to be sneaky

Example 14.94. $\int_0^\pi \frac{1}{1+2^{\tan(x)}}$

Till now we were having functions we could integrate and kings rule just saved time, this time we a function we can't integrate but like a king, the kings rule doesn't care.

Solution. Using the Kings rule:

$$I = \int_0^\pi \frac{1}{1+2^{\tan(x)}}$$

$$\iff I = \int_0^\pi \frac{1}{1+2^{\tan(\pi-x)}} \iff I = \int_0^\pi \frac{1}{1+2^{-\tan(x)}} \iff I = \int_0^\pi \frac{2^{\tan(x)}}{1+2^{\tan(x)}}$$

Adding to the first equation:

$$2I = \int_0^\pi \frac{2^{\tan(x)}}{1+2^{\tan(x)}} + \int_0^\pi \frac{1}{1+2^{\tan(x)}}$$

$$\iff 2I = \pi$$

$$\iff I = \frac{\pi}{2}$$

□

and once in a while the question is a bit cheeky

Example 14.95. $I = \int_1^{2019} (1-x)(2-x)(3-x)\dots(2019-x)dx$

Solution. The answer is obviously zero over integers. But as integration is over real numbers, we'll have to get 0 some other way.

Using the kings rule:

$$I = \int_1^{2019} (1-x)(2-x)(3-x)\dots(2019-x)dx$$

$$\iff I = \int_1^{2019} (1-2020+x)(2-2020+x)(3-2020+x)\dots(2019-2020+x)dx$$

$$\iff I = \int_1^{2019} (x-2019)(x-2018)(x-2017)\dots(x-1)dx$$

$$\iff -1^{2019} I = -I = \int_1^{2019} (1-x)(2-x)(3-x)\dots(2019-x)dx$$

$$\therefore -I = I$$

$$\therefore I = 0$$

□

and sometimes the question maker will try make it seem complex:

Example 14.96. If [...] stands for the greatest Integer function. Calculate

$$\int_4^{10} \frac{[x^2]}{[x^2 - 28x + 196] + [x^2]} dx$$

Solution. As we cannot directly integrate the greatest integer function this seems scary. But the king's rule disagrees:

$$\begin{aligned} I &= \int_4^{10} \frac{[x^2]}{[x^2 - 28x + 196] + [x^2]} dx \\ \iff I &= \int_4^{10} \frac{[x^2]}{[(14-x)^2] + [x^2]} dx \\ \iff I &= \int_4^{10} \frac{[(14-x)^2]}{[(14-x)^2] + [x^2]} dx \end{aligned}$$

Adding to the first equation gives:

$$\begin{aligned} 2I &= \int_4^{10} dx \\ \iff 2I &= 6 \\ \iff I &= 3 \end{aligned}$$

□

And to end kings rule, here is a beautiful question of that topic:

Example 14.97. If $f(x)$ is continuous and symmetric about $x = 2$ and $\int_1^3 f(x)dx = 2$, then find $\int_1^3 xf(x)dx$

Solution. We need to realize that $f(4-x) = f(x)$ using king's rule on the integral.

Using kings rule on the second integral now:

$$\begin{aligned} I &= \int_1^3 xf(x)dx \\ \iff I &= \int_1^3 (4-x)f(4-x)dx \iff I = \int_1^3 (4-x)f(x)dx \iff I = \\ &\int_1^3 4f(x)dx - \int_1^3 xf(x)dx \\ \iff I &= 4 \int_1^3 f(x)dx - I \\ \iff 2I &= 8 \\ \iff I &= 4 \end{aligned}$$

□

14.7 Leibnitz Theorem

Theorem 14.98. If $F(x) = \int_{g(x)}^{h(x)} \phi(t)f(t)dt$,
 $F'(x) = \phi(x)(f(h(x))h'(x) - f(g(x))g'(x))$

This seems messy. It basically converts a function defined as an integral with variable limits into a simpler function.

The proof is using the definition of integration.

It's use cases are few and far between so we'll take a question from IIT 2005 for the example:

Example 14.99. (IIT 2005) If $\int_{\sin(x)}^1 t^2 f(t)dt = 1 - \sin(x)$ for $x \in (0, \frac{\pi}{2})$ then $f(\frac{1}{\sqrt{3}})$ is?

Solution. Using Leibnitz Theorem:

$$\begin{aligned} F'(x) &= -\sin^2(x)f(\sin(x)) \cdot \cos(x) = -\cos(x) \iff f(\sin(x)) = \frac{1}{\sin^2(x)} \\ \iff f(x) &= \frac{1}{x} \end{aligned}$$
□

14.8 Taylor Maclaurian Series

We had used them in Calc-I where I had promised a proof later. I will honour my promise now.

Proof. We know that $f(x) = f(0) + \int_0^x f'(t)dt$ using the fundamental theorem.

$$\therefore f(x) = f(0) + \int_0^x f'(x-t)dt \text{ using King's rule}$$

$$\therefore f(x) = f(0) + xf'(0) + \int_0^x tf'(x-t)dt \text{ using Integration by parts}$$

$\therefore f(x) = f(0) + xf'(0) + \frac{1}{2}f''(0)x^2 + \frac{1}{2} \int_0^x t^2 f'''(x-t)dt$ Reiterating this again and again will give us

$$f(x) = f(0) + \frac{xf'(0)}{1!} + \frac{x^2 f''(0)}{2!} + \frac{x^3 f'''(0)}{3!} + \dots$$

And with almost 30 pages of learning we have finally proved Taylor Maclaurian Series. **QED** □

14.9 Differential Equations

Example 14.100 (Motivating Example). The population of the city of Yogyakarta increases at a rate proportional to the number of its inhabitants present at any time t . If the population of Yogyakarta was 30,000 in 1970 and 35,000 in 1980, what will be the population of Yogyakarta in 1990?

We need to realize that as number of people in a place increase, so does the rate of increase of population. That is if there are more people, more of them reproduce, leading to more population and the cycle continues(not really, nature has a carrying capacity and above which population doesn't grow, but below that our analysis is mostly true)

So we can say that the rate of change of population is proportional to the population. That means $\frac{dy}{dt} = Ky$ where K is a constant.

Such equations are called differential equations as they have a differential term. This particular one is quite elementary to solve:

$$\frac{dy}{dt} = Ky$$

$$\iff \frac{dy}{Ky} = dt \text{ Now we integrate on both the sides}$$

$$\iff \int \frac{dy}{Ky} = \int dt$$

$$\iff \frac{1}{K} \ln y = t + C$$

The variables are K and C and we have two equations and two variables, so they can be solved. I leave the solving to you.

Such equations are common place in physics as the rate of a lot of things in nature is based on its quantity which itself is changing. For example:

The rate at which the nuclei of a substance decay is proportional to the amount (more precisely, the number of nuclei) $A(x)$ of the substance remaining at time t . This leads to a differential equation $\frac{dA}{dt} = kA$. Another is the cooling or warming of objects in air. Suppose T represents the temperature of a body at time t , T_m the temperature of the surrounding medium. The rate at which the temperature of a body changes is proportional to the difference between the temperature of the body and the temperature of the surrounding medium. This leads to the differential equation $\frac{dT}{dt} = k(T - T_m)$.

While the above differential equations were elementary to solve, linear differential equations tend to lead to some strange solutions.

Theorem 14.101. *For a linear differential equation of the form $\frac{dy}{dx} + Py = Q$ for P, Q being either constants or functions of x . The solution is $y \cdot IF = \int Q \cdot IF dy + C$ where $IF = e^{\int P dx}$ where IF stands for the integrating factor*

It is recommended to remember this as it is as the proof is a bit non-intuitive

Proof. Multiplying the equation by $g(x)$ such that the RHS becomes the first derivative of $yg(x)$

$$\begin{aligned} & \therefore \frac{dy}{dx} + Py = Q \\ & \iff g(x) \frac{dy}{dx} + Pg(x)y = Qg(x) \\ & \therefore g(x) \frac{dy}{dx} + Pg(x)y = \frac{d}{dx}(yg(x)) \\ & \iff g(x) \frac{dy}{dx} + Pg(x)y = \frac{dy}{dx}g(x) + yg'(x) \\ & \iff Pg(x)y = yg'(x) \\ & \iff P = \frac{g'(x)}{g(x)} \\ & \implies \int P dx = \int \frac{g'(x)}{g(x)} dx \\ & \iff \int P dx = \ln g(x) \\ & \iff g(x) = e^{\int P dx} \end{aligned}$$

We can now use this value of $g(x)$ in the equation to get:

$$\begin{aligned} & (e^{\int P dx}) \frac{dy}{dx} + Py(e^{\int P dx}) = Q(e^{\int P dx}) \\ & \iff \frac{d}{dx}(ye^{\int P dx}) = Qe^{\int P dx} \text{ And now we can integrate on both sides to get:} \\ & ye^{\int P dx} = \int Qe^{\int P dx} dx + C \quad \square \end{aligned}$$

We'll not do any exercises or examples of this as it is only included to complete the topic, I have not seen it being used in any olympiad ever.

14.10 What's more?

This chapter was one the longest in this book. It took me more than 20 hours in total to write. However, this is not even 1% of calculus. While this is more than what you generally study in high school, we have not talked about hyperbolic functions which predict how a lot of things behave. Including but not limited to hanging chains and rotating wheels. While I mentioned in passing



Figure 14.0. A hanging chain experiences a different force everywhere, as the distance from support increases. The shape it forms is called a Catenary. It uses calc which we didn't study

some functions without proper integrals, we didn't study about their approximate integrals. We also worked only on the real plane, what about imaginary plane(where $i = \sqrt{-1}$ lies) which occur a lot in electricity and quantum mechanics? Also our work was mostly contained with two variables. What about 3D graphs? What about even higher dimensions? While we learnt to find area under lines, but what about the area under a curved cloth? We didn't study about what if instead of an equation we have an inequality? While we did explore differential equations, what about when they are squared? And cubed? While you'll learn some of these things later in this book itself, a lot while interesting is beyond the scope of this book. While I encourage you to dive head on into calculus and have provided the resources in the appendix D, I'll like to take a moment to remark on the fact that so much of the world around us is explained on the basis on two operations differentiation and integration. While no one knows who truly discovered them in the ancient civilizations, here is to Zeno, Archimedes, Cavalieri, Torricelli, Galileo, Newton, Leibniz, Taylor, Maclaurin, Abel, Euler, Lagrange and many more whose work we will not study in any part of this book for giving us calculus. Now to the exercises!

Exercises

(1) Compute: $\int \frac{x^4 + 5x - 1}{\sqrt{x}} dx$

- (2) Compute: $\int \frac{x^4}{x^2+1} dx$
- (3) Compute: $\int \sqrt{1 - \sin(2x)} dx$ when
 (a) $x \in (\frac{\pi}{4}, \frac{5\pi}{4})$
 (b) $x \in (\frac{-3\pi}{4}, \frac{\pi}{4})$
- (4) Compute $\int \frac{2x+3}{x+1} dx$
- (5) Evaluate: $\int (x - \tan(x)) \tan^2(x) dx$
- (6) (Jee Mains 2021) If $f(x) = \int \frac{5x^8 + 7x^6}{(x^2+1+2x^7)^2} dx, x \geq 0, f(0) = 0$ and $f(1) = \frac{1}{k}$,
 the value of k is:
- (7) Evaluate: $\int \frac{1}{\sqrt{(x-1)(x-2)}} dx$
- (8) Compute: $\int \sqrt{\frac{x-2}{x+2}} dx$
- (9) Compute: $\int \frac{\tan(x)}{\sqrt{a+b\tan^2(x)}} dx$
- (10) Compute $\int \sqrt{x^2 + 2x + 5} dx$
- (11) Compute: $\int \frac{\sin(2x)}{\sin(5x)\sin(3x)} dx$
- (12) Compute: $\int \sin^2(x) \cos^2(x) dx$
- (13) Compute: $\int \sin^4(x) \cos^5(x) dx$
- (14) Compute: $\int \tan^2(x) \sec^4(x) dx$
- (15) Evaluate: $\int \frac{1}{\sin(x)\cos^2(x)} dx$
- (16) Evaluate: $\int \frac{\cos(\theta)+\sin(\theta)}{\sqrt{5+\cos(2\theta)}} d\theta$
- (17) (AIEEE 2008) The value of $\sqrt{2} \int \frac{\sin(x)dx}{\sin(x-\frac{\pi}{4})}$ is:
- (18) Evaluate: $\int \frac{\cos(x)-\sin(x)+1-x}{e^x \sin(x)+x} dx$
- (19) Evaluate: $\int \frac{\sin(x)}{\sin(4x)} dx$
- (20) Evaluate $\int \frac{2x^2}{1+x^4} dx$
- (21) Evaluate $\int \frac{\sqrt{x+1}}{(x+2)\sqrt{x+3}} dx$
- (22) Evaluate $\int \frac{1+\sin(x)}{\sin(x)(1+\cos(x))} dx$
- (23) Compute $\int \frac{2\sin(x)+\cos(x)}{7\sin(x)-5\cos(x)} dx$

- (24) Compute $\int \frac{1}{2+\sin^2(x)} dx$
- (25) (JEE Mains 2013) If $\int f(x)dx = g(x)$, then $\int x^5 \cdot f(x^3)dx$ is equal to
- (26) Evaluate $\int e^x \frac{2+\sin(2x)}{1+\cos(2x)}$
- (27) If $\int_{-1}^{-4} f(x)dx = 4$ and $\int_2^{-4} e - f(x)dx = 7$ then $\int_1^2 f(x)dx = ?$
- (28) (IIT 1995) Evaluate $\int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}}$
- (29) Evaluate $\int_{-\pi}^{\pi} \frac{\cos^2(x)}{1+2^x}$
- (30) Suppose there is continuous function such that $f(x) + f(-x) = x^2$, what is $\int_{-1}^1 f(x)dx$ equal to?
- (31) Evaluate $\int \frac{\ln(1+x)}{1+x^2} dx$

15

Ch-14 Linear Algebra

Let me address the fundamental question on the face of it: what is Linear Algebra, and why should we bother learning it? You see, mathematics is not just a set of abstract rules and theorems; it's a language that allows us to describe and understand the world around us. Linear Algebra is a crucial part of this mathematical language, and it serves as a bridge between various mathematical concepts and the real-world problems we aim to solve.

Now, linear algebra may seem like a bit of a departure from your previous mathematical endeavors, and I want to assure you that it's quite different, yet fundamentally tied to the mathematics you've encountered before. You see, on the face of it linear algebra serves as a bridge between algebra and geometry, and at a greater depth it achieves a lot more.

At its core, Linear Algebra deals with matrices, which are essentially a different way of representing data than the vectors we'll explore later on. Think of a matrix as a structured arrangement of numbers, much like an Excel spreadsheet. You might remember incidence matrices from Power Overwhelming chapter. These numbers hold hidden patterns and relationships that we'll uncover throughout this chapter.

Now, I must offer a small apology in advance. This chapter, and indeed much of Linear Algebra, may seem abstract at first. We'll delve into concepts that may not seem immediately applicable to the real world. But rest assured, as we progress, we'll unlock the practical applications of these abstract notions. Think of this abstract treatment as laying the groundwork, building a solid foundation for the magnificent mathematical structures we'll explore later. Much like constructing a sturdy building, we need a strong foundation to support the practical applications of Linear Algebra that we'll discuss in due

time.

I want to also extend my apologies once again, before we embark on this journey through Linear Algebra. While our goal is to explore the beauty and utility of matrices, determinants, and their applications, I must acknowledge that I have not provided the rigorous proofs for every statement and concept we encounter.

You see, Linear Algebra can be quite intricate, and proving every assertion can often be a lengthy and intricate process, especially when we're dealing with general matrices. Consider, for instance, taking a random matrix, multiplying it by another, and then attempting to factorize the result. This task, though fundamental and basic algebra, can indeed become quite tedious. I however, encourage you to do so. You won't need any more knowledge than what we covered in basic algebraic manipulations

With that out of the way, I must remind you that Linear algebra is not just about abstract problems and equations. Linear Algebra is the muscle behind modern technology. It powers your smartphone apps and games, helps Netflix recommend your next binge-worthy series, and even guides spacecraft to explore the cosmos. Heck, it's even there when you make a simple Google search. So, if you ever wondered how your world works, this is a fantastic place to start.

As for the how and why of these real-world applications, they'll come, I promise! You see, linear algebra is like a Swiss Army knife for scientists, engineers, and data scientists. We can use it to solve systems of equations, analyze networks, optimize processes, and even make sense of vast amounts of data in fields like machine learning and statistics. So, while we may start in the abstract, by the end of this journey, you'll understand why linear algebra is so essential in the modern world.

15.1 Determinants

Let's for now assume matrix to be a row and column of numbers. We'll talk more about them later.

The determinant is only defined for a matrix with equal columns and rows, or a square matrix. The number of rows or columns is called the order. We'll define it as follows:

Definition 15.1. Determinant of matrix of order 1:

$$|a| = a$$

Definition 15.2. Determinant of matrix of order 2:

$$\begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} = a_1 \cdot b_2 - b_1 \cdot a_2$$

We can remember this definition by looking at it as cross multiplying on then diagonals and then subtracting.

Here are a few examples of taking the determinant:

Example 15.3 (Motivating Example). $\begin{vmatrix} 4 & 3 \\ 1 & -3 \end{vmatrix}$

Solution. This quite simple.

$$\begin{aligned} 4 * (-3) - 3 * 1 \\ = -12 - 3 \\ = -15 \end{aligned}$$

□

Example 15.4. If $\begin{vmatrix} e^x & \sin(x) \\ \cos(x) & \ln 1 + x \end{vmatrix} = A + Bx + Cx^2 + \dots$ What is the value of A ?

Solution. Before we pull out the expansions from calculus, notice that taking $x = 0$ will make the determinant A and that is what is asked from us.

$$\begin{aligned} \therefore A &= \begin{vmatrix} e^0 & \sin(0) \\ \cos(0) & \ln 1 + 0 \end{vmatrix} \\ &= \begin{vmatrix} 1 & 0 \\ 1 & 0 \end{vmatrix} \\ &= 0 \end{aligned}$$

□

Now we enter the actually confusing area. For any general matrix we will here onward write the term at the intersection of the i^{th} column and j^{th} row as a_{ij} . Using this notation we define the determinant of order 3 as:

Definition 15.5. Determinant of order 3 $\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$
 $= a_{11}(a_{22} \cdot a_{33} - a_{23} \cdot a_{32}) - a_{12}(a_{21} \cdot a_{33} - a_{23} \cdot a_{31}) + a_{13}(a_{21} \cdot a_{32} - a_{22} \cdot a_{31})$

We can remember this by basically removing the row and column of a number, taking the determinant of the rest of the matrix(now of order 2) and then multiply them. Then we take the next term in the column or row, repeat the process but alternating the sign. We can do this for any row or column to get the determinant(the definition is on the first row but we can choose the one which leads to the shortest calculations).

Let's try an example:

Example 15.6 (Motivating example). $\begin{vmatrix} 3 & 3 & 2 \\ 5 & 4 & 7 \\ 5 & 7 & 6 \end{vmatrix}$

Solution. Using the first row seems promising as it has relatively small numbers.

$$\begin{aligned} \begin{vmatrix} 3 & 3 & 2 \\ 5 & 4 & 7 \\ 5 & 7 & 6 \end{vmatrix} &= 3(4*6 - 7*7) - 3(5*6 - 7*5) + 2*(5*7 - 4*5) \\ &= 3*(-25) - 3(-5) + 2(15) \\ &= -75 + 15 + 30 \\ &= 30 \end{aligned}$$

□

Now we'll define two new terms called minor and cofactor.

Definition 15.7. The minor M_{ij} of an element a_{ij} in a matrix $\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ is the determinant obtained by deleting the i^{th} row and j^{th} column.

Definition 15.8. The co-factor C_{ij} of an element a_{ij} is given by:
 $C_{ij} = (-1)^{i+j} M_{ij}$

Let's do an example to get comfortable with the definition.

Example 15.9 (Motivating example). Find the co-factors of all the elements of : $\begin{bmatrix} 1 & -5 & -1 \\ 5 & 0 & 3 \\ -3 & 7 & 9 \end{bmatrix}$

Solution. For $a_{11} = 1$, the $C_{11} = (-1)^{1+1} M_{11}$

$$= (-1)^2 \begin{vmatrix} 0 & 3 \\ 7 & 9 \end{vmatrix}$$

$$= -21$$

For $a_{12} = -5$, the $C_{12} = (-1)^{1+2} M_{12}$

$$= (-1)^3 \begin{vmatrix} 5 & 3 \\ -3 & 9 \end{vmatrix}$$

$$= -54$$

For $a_{13} = -1$, the $C_{13} = (-1)^{1+3} M_{13}$

$$= (-1)^4 \begin{vmatrix} 5 & 0 \\ -3 & 7 \end{vmatrix}$$

$$= 38$$

And we can do the same with the rest of the matrix. I think, you will be able to do that by yourself.

□

What we need to notice is that we will always have some matrix

Definition 15.10. The adjacent of some matrix A $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ Is defined as $\text{adj}(A) = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}$

This leads us to another definition of the determinant:

Definition 15.11. $\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \cdot C_{11} + a_{12} \cdot C_{12} + a_{13} \cdot C_{13}$ Or in words we can say: The sum of product of elements of any row or column of A with the corresponding row or column of $\text{adj}(A)$ is equal to the determinant of A $\det(A)$.

A classic matrix we should remember the determinant to is:

$$\text{Example 15.12. } \begin{vmatrix} 1 & z & -y \\ -z & 1 & x \\ y & -x & 1 \end{vmatrix}$$

Solution. Expanding along the first row gives us:

$$\begin{aligned} & (1+x^2) - z(-z-xy) + (-y)(zx-y) \\ &= 1+x^2+z^2+xyz-xyz+y^2 \\ &= 1+x^2+y^2+z^2 \end{aligned}$$

□

Another thing which is quite interesting, although useless, is:

Theorem 15.13. *The sum of product of any row or column of matrix A with the corresponding any other row or column (other than the corresponding) of $\text{adj}(A)$ is zero.*

15.2 Properties of Determinants

Theorem 15.14. *Value of determinant remains unchanged on interchanging rows and columns.*

$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{vmatrix}$ The flipping of rows and columns in called transposing.

Theorem 15.15. If any two rows (columns) of a determinant are inter-

changed, the value of determinant changes sign.

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = - \begin{vmatrix} a_{31} & a_{32} & a_{33} \\ a_{21} & a_{22} & a_{23} \\ a_{11} & a_{12} & a_{13} \end{vmatrix}$$

Theorem 15.16. If any two rows (columns) are identical then value of determinant is zero

We can use this properties to solve problems which are otherwise harder to solve:

Example 15.17. Find x if

$$\begin{vmatrix} 1 & 1 & 0 \\ (x^2 - 5x + 7) & 1 & 0 \\ 2 & 1 & 1 \end{vmatrix} = 0$$

Solution. This is quite simple as we can notice that two terms of the first and second row are equal. Hence if the third are also equal, the determinant will become 0.

$$\therefore x^2 - 5x + 7 = 1$$

$$\therefore x^2 - 5x + 6 = 0$$

$$\therefore x = 2, 3$$

□

Here is another theorem, this one is actually quite use full, in both questions and in real world.

Theorem 15.18. If all the elements of any row or column be multiplied by a number K then value of determinant is multiplied by K

$$K \times \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} Ka_{11} & Ka_{12} & Ka_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} Ka_{11} & a_{12} & a_{13} \\ Ka_{21} & a_{22} & a_{23} \\ Ka_{31} & a_{32} & a_{33} \end{vmatrix}$$

Let's annihilate a JEE advance question to prove the power of this:

Example 15.19. If $b_{ij} = 2^{i+j}a_{ij}$ where a_{ij} and b_{ij} are elements of 3×3 determinants Δ_1 and Δ_2 respectively, the find Δ_2 if $\Delta_1 = 2$

Solution. Let the determinant Δ_1 be of the matrix:

$$\begin{aligned} & \left| \begin{array}{ccc} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{array} \right|, \text{ therefore} \\ & \Delta_2 = \left| \begin{array}{ccc} 2^2 a_{11} & 2^3 a_{12} & 2^4 a_{13} \\ 2^3 a_{21} & 2^4 a_{22} 2^5 & a_{23} \\ 2^4 a_{31} & 2^5 a_{32} & 2^6 a_{33} \end{array} \right| \\ & = 2 \left| \begin{array}{ccc} 2^2 a_{11} & 2^3 a_{12} & 2^4 a_{13} \\ 2^2 a_{21} & 2^3 a_{22} 2^4 & a_{23} \\ 2^4 a_{31} & 2^5 a_{32} & 2^6 a_{33} \end{array} \right| \\ & = 2 * 2^2 \left| \begin{array}{ccc} 2^2 a_{11} & 2^3 a_{12} & 2^4 a_{13} \\ 2^2 a_{21} & 2^3 a_{22} 2^4 & a_{23} \\ 2^2 a_{31} & 2^3 a_{32} & 2^4 a_{33} \end{array} \right| \\ & = 2^3 * 2^2 * 2^3 * 2^4 \left| \begin{array}{ccc} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{array} \right| \\ & = 2^{12} \cdot \Delta_1 \\ & = 2^{13} \end{aligned}$$

□

Theorem 15.20. If each element of any row (or column) is expressed as sum of two (or more) terms then the determinant can be expressed as the sum of two (or more) determinants.

$$\left| \begin{array}{ccc} a_{11} + x & a_{12} + y & a_{13} + z \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{array} \right| = \left| \begin{array}{ccc} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{array} \right| + \left| \begin{array}{ccc} x & y & z \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{array} \right|$$

This theorem is more commonly used to combine matrices than to split them. Here is an example which will also brush up your sequence and series:

$$\text{Example 15.21. If } \Delta_r = \left| \begin{array}{ccc} r+x & n(n+1) & n^2+n^3 \\ 2^r & 4(2^n-1) & n^2+n+1 \\ 3^r & 3(3^n-1) & 2n+1 \end{array} \right| \text{ then } \sum_{r=1}^n \Delta_r = ?$$

Solution. We can notice that the second column and the third column are same in all the matrices. Therefore we can simply combine over the first column.

$$\text{As } \sum_{r=1}^n r = \frac{r(r+1)}{2}$$

$$\sum_{r=1}^n 2^r = \frac{2(2^n-1)}{2-1}$$

$$\sum_{r=1}^n 3^r = \frac{3(3^n+1)}{3-1}$$

$$\text{Therefore, } \sum_{r=1}^n \Delta_r = \left| \begin{array}{ccc} \frac{r(r+1)}{2} & n(n+1) & n^2+n^3 \\ \frac{2(2^n-1)}{2-1} & 4(2^n-1) & n^2+n+1 \\ \frac{3(3^n+1)}{3-1} & 3(3^n-1) & 2n+1 \end{array} \right|$$

$$= \frac{1}{2} \begin{vmatrix} r(r+1) & n(n+1) & n^2 + n^3 \\ 4(2^n - 1) & 4(2^n - 1) & n^2 + n + 1 \\ 3(3^n + 1) & 3(3^n - 1) & 2n + 1 \end{vmatrix}$$

$$= 0$$

As column 1 is equal to column 2.

□

Theorem 15.22. *The value of determinants is not altered by adding or subtracting the multiple of any row or column in other row or column.*

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11} + Ka_{31} & a_{12} & a_{13} \\ a_{21} + Ka_{32} & a_{22} & a_{23} \\ a_{31} + Ka_{33} & a_{32} & a_{33} \end{vmatrix}$$

What makes this theorem even more powerful is the fact we can do 2 operations at a time.

What I mean to say is that:

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11} - a_{12} & a_{12} - a_{13} & a_{13} \\ a_{21} - a_{22} & a_{22} - a_{23} & a_{23} \\ a_{31} - a_{32} & a_{32} - a_{33} & a_{33} \end{vmatrix}$$

This allows us to solve a lot of problems at extreme speeds as long as we can find the correct manipulation.

For example:

$$\begin{vmatrix} 1 & a & b+c \\ 1 & b & c+a \\ 1 & c & a+b \end{vmatrix}$$

can be calculated in an instant by simply sum-

ming up the second and third column and then taking $a + b + c$ as common.
Let's do something a tad more difficult:

Example 15.23. Evaluate:

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$$

Solution. While expanding the determinant along the first row is quite trivial, let's use properties to make a joke of this.

Subtracting the first column from the second and third from the second:

$$\begin{vmatrix} 0 & 0 & 1 \\ a-b & b-c & c \\ a^2-b^2 & b^2-c^2 & c^2 \end{vmatrix}$$

We could now expand with even greater ease, but wait there's more:

$$(a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 1 & 1 & c \\ a+b & b+c & c^2 \end{vmatrix}$$

Which simplifies to $(a-b)(b-c)(c-a)$ and we are done.

□

I recommend remembering this result as a standard form.
Here is an very similar example for you to try:

$$\text{Example 15.24. } \begin{vmatrix} 1 & a & bc \\ 1 & b & ac \\ 1 & c & ab \end{vmatrix}$$

As simple way to hide the three ones looks like:

$$\text{Example 15.25. } \begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$$

Solution. We notice that the sum of all three rows is equal.
Therefore, we add along the rows to get:

$$\begin{aligned} & \begin{vmatrix} a+b+c & b & c \\ a+b+c & c & a \\ a+b+c & a & b \end{vmatrix} \\ &= (a+b+c) \begin{vmatrix} 1 & b & c \\ 1 & c & a \\ 1 & a & b \end{vmatrix} \end{aligned}$$

And the three one's present themselves. \square

It's a common approach to try to get the three one's to simplify the determinant.

$$\text{Example 15.26. } \begin{vmatrix} b^2c^2 & bc & b+c \\ c^2a^2 & ca & c+a \\ a^2b^2 & ab & a+b \end{vmatrix}$$

Solution. We'll multiply the first row by a , second row by b and the third row by c to get:

$$\begin{aligned} & \frac{1}{abc} \begin{vmatrix} ab^2c^2 & abc & a(b+c) \\ bc^2a^2 & bca & b(c+a) \\ ca^2b^2 & cab & c(a+b) \end{vmatrix} \\ &= \frac{(abc)^2}{abc} \begin{vmatrix} bc & 1 & ab+ac \\ ca & 1 & bc+ab \\ ab & 1 & ac+bc \end{vmatrix} \end{aligned}$$

This may seem good enough as we already have the three one's but we are gonna utterly humiliate the question by adding column 3 to column 1.

$$\begin{array}{ccc|c} ab+bc+ca & 1 & ab+ac \\ abc & ab+bc+ca & 1 & bc+ab \\ ab+bc+ca & 1 & ac+bc \end{array}$$

$$= abc(ab + bc + ca) \begin{vmatrix} 1 & 1 & ab + ac \\ 1 & 1 & bc + ab \\ 1 & 1 & ac + bc \end{vmatrix}$$

$$= 0$$

And we are done. \square

15.3 Application of Determinants

We are now done with determinants. Before looking at Matrices, let's talk about some applications of them.

Theorem 15.27 (Shoelace formula). *Area of a triangle with the vertices at coordinates $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ = magnitude of $\frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$*

We'll see a more simplified and generalized form of this in coordinate geometry, from which this is derived. But the determinant form has its own use, specifically in the form:

Theorem 15.28 (Condition of Colinearity). *If $\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = 0$ then the points $(x_1, y_1); (x_2, y_2); (x_3, y_3)$ are colinear.*

This is true as the area of a line is zero.

This tends to occur in questions in a very typical pattern:

Example 15.29. If the points $(at_1^2, 2at_1), (at_2^2, 2at_2), (a, 0)$ are colinear. If $(t_1 \neq t_2)$, then find the minimum value of $t_1^2 + 9t_2^2$.

Solution. Using the condition of Colinearity, $\begin{vmatrix} at_1^2 & 2at_1 & 1 \\ at_2^2 & 2at_2 & 1 \\ a & 0 & 1 \end{vmatrix} = 0$

$$\begin{aligned} &\Leftrightarrow \begin{vmatrix} at_1^2 - a & 2at_1 & 0 \\ at_2^2 - a & 2at_2 & 0 \\ a & 0 & 1 \end{vmatrix} = 0 \\ &\Leftrightarrow (at_1^2 - a)(2at_2) = (2t_2^2 - a)(2at_1) \\ &\Leftrightarrow 2a^2t_2t_1^2 - 2a^2t_2 = 2a^2t_1t_2^2 - 2a^2t_1 \\ &\Leftrightarrow t_2t_1^2 - t_2 = t_1t_2^2 - t_1 \\ &\Leftrightarrow t_2 - t_1 = t_2t_1(t_1 - t_2) \end{aligned}$$

$$\Leftrightarrow t_1 t_2 = -1$$

At this point we can use AM-GM inequality to say that:

$$\frac{t_1^2 + 9t_2^2}{2} \geq \sqrt{9t_1^2 t_2^2}$$

$$\Leftrightarrow t_1^2 + 9t_2^2 \geq 2\sqrt{9}$$

$$\Leftrightarrow t_1^2 + 9t_2^2 \geq 6$$

Hence, the minimum value of $t_1^2 + 9t_2^2$ is 6. \square

Here is an example for you to try. This one uses the AM-HM inequality.

Example 15.30. If the points $(a, 0), (x, y), (0, b)$ are colinear where $a, b, x, y > 0$ then find the minimum value of $\frac{a}{x} + \frac{b}{y}$?

Theorem 15.31 (Concurrency of Lines). *Three lines $a_1x + b_1y + c_1 = 0$; $a_2x + b_2y + c_2 = 0$; $a_3x + b_3y + c_3 = 0$ are concurrent if and only if they are non-*

parallel and $\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = 0$

This can also be proven using coordinate geometry.

Let's use this to solve a pretty difficult question from JEE advance, which has appeared multiple times in mains as well:

Example 15.32. If the lines $ax + y + 1 = 0$, $x + by + 1 = 0$ and $x + y + c = 0$ where a, b, c being distinct and different from unity, are concurrent, then the value of $\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c}$ is

Solution. Using the concurrency of lines condition:

$$\begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0$$

$$\Leftrightarrow \begin{vmatrix} a-1 & 1-b & 0 \\ 0 & b-1 & 1-c \\ 1 & 1 & c \end{vmatrix} = 0 \text{ Expanding along the first column,}$$

$$(a-1)[(b-1)c - (1-c)] + (1-b)(1-c) = 0$$

$$\Leftrightarrow (a-1)(b-1)(c) - (a-1)(1-c) + (1-b)(1-c) = 0$$

This seems like a good time to divide by $(a-1)(b-1)(c-1)$

$$\frac{c}{c-1} + \frac{1}{b-1} + \frac{1}{a-1} = 0$$

$$\Leftrightarrow \frac{c-1+1}{c-1} + \frac{1}{b-1} + \frac{1}{a-1} = 0$$

$$\Leftrightarrow \frac{1}{c-1} + \frac{1}{b-1} + \frac{1}{a-1} - 1 = 0$$

$$\Leftrightarrow \frac{1}{c-1} + \frac{1}{b-1} + \frac{1}{a-1} = 1$$

\square

15.4 Crammer's Rule

Have you ever wondered how computers can solve 7-8 simultaneous linear equations in seconds. The answer I am not looking for is 'They are computers, duh!'

The answer is using Matrices and Determinants. The matrix version of this is going to come up in a minute, but the determinant version is known as Crammer's Rule over Gabriel Crammer who generalized this for n variables and n equations. We will only get to use it for 2, 3 variables and 2, 3 equations as we don't want to compute determinants of order 4 by hand. Those versions were originally found by Colin Maclaurin of the Taylor-Maclaurin Series. But he had already stuck his name elsewhere, so he let Crammer have it(also after all Crammer was the one who generalized it)

Theorem 15.33 (Crammer's Rule). *For the system of liner equations: $a_1x + b_1y + c_1z = d_1$; $a_2x + b_2y + c_2z = d_2$; $a_3x + b_3y + c_3z = d_3$, the solution is:*

$$x = \frac{\Delta_x}{\Delta}, y = \frac{\Delta_y}{\Delta}, z = \frac{\Delta_z}{\Delta}$$

$$\text{where } \Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}, \Delta_x = \begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}, \Delta_y = \begin{vmatrix} a_1 & d_1 & c_1 \\ a_2 & d_2 & c_2 \\ a_3 & d_3 & c_3 \end{vmatrix}, \Delta_z = \begin{vmatrix} a_1 & b_1 & d_1 \\ a_2 & b_2 & d_2 \\ a_3 & b_3 & d_3 \end{vmatrix}$$

The simplest proof follows from opening the determinants and comparing them to the equation's substitutions.

I recommend you trying this out on random systems of equations. But before yo do that here is small life saver:

Definition 15.34. If system has solution (unique or many) than it is called consistent, otherwise it is called inconsistent.

If while using the Crammer's rule $\Delta = \Delta_x = \Delta_y = \Delta_z = 0$ then the system has infinite solutions, and if $\Delta = 0$ but even one of the rest of the determinants is non-zero, then the system has no solutions.

If $\Delta \neq 0$ then it has one unique solution. We need to make sure that all coefficients are not zero in any of the $\Delta = 0$ cases.

This definition may seem like the normal math speak for be careful, but we actually use it quite a lot in questions as well. Here is an example for us to nuke from IIT 2004.

Example 15.35. (IIT 2004) Find k such that, $2xy + 2z = 2, x2y + z = 4, x + y + kz = 4$ has no solution

Solution. For the equation to have no solution, $\Delta = 0$

$$\therefore \begin{vmatrix} 2 & -1 & 2 \\ 1 & -2 & 1 \\ 1 & 1 & k \end{vmatrix} = 0$$

Opening the determinant along the third row,

$$3 - 3k = 0$$

$$\iff k = 1$$

Technically, we need to check if $\Delta_x, \Delta_y, \Delta_z$ are zero, but knowing that this is an exam with an answer, we can be assured that there is only one solution which is $k = 1$ \square

However, we have yet to consider the case where the system is homogeneous.

Definition 15.36. If the constant terms in the system of equations (i.e. d_1, d_2, d_3) are all zero, then system is called homogeneous system of equations

Here are some more definitions pertaining to homogeneous systems:

Theorem 15.37. (1) Homogeneous system is always consistent (as $(0, 0, 0)$ always satisfies it). (2) $(0, 0, 0)$ is also called trivial solution. (3) Homogeneous system has infinite non-trivial (i.e. non-zero) solutions if and only if $\Delta = 0$

We can use this questions such as:

Example 15.38. (IIT 2000) If the system of equations $xKyz = 0$

$$Kxyz = 0$$

$x + yz = 0$ Has a non zero solution then $K =$

Solution. We basically want $\begin{vmatrix} 1 & -k & -1 \\ k & -1 & -1 \\ 1 & 1 & -1 \end{vmatrix} = 0$

$$\iff -(k+1) - (-1)(1+k) + (-1)(-1+k^2) = 0$$

$$\iff -(k+1) + (1+k) + (1-k^2) = 0$$

$$\iff -k - 1 + 1 + k + 1 - k^2 = 0$$

$$\iff k^2 = 1$$

$$\iff k = \pm 1$$

\square

15.5 Types Matrices

If you know determinants, you already know most of matrix.

This section is just me telling you all the names you need to know in order to solve ahead.

Definition 15.39. Matrix an arrangement of $m \times n$ elements in ‘ m ’ rows and

‘ n ’ columns.

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

Unlike determinants, it is possible that $m \neq n$.

Now let's discuss some special types of matrices.

Definition 15.40. Row Matrix is a matrix having only one row. $[a \ b \ c \ \dots]$

Definition 15.41. Column Matrix is a matrix having only one column.

$$\begin{bmatrix} a \\ b \\ c \\ \vdots \end{bmatrix}$$

Definition 15.42. Null matrix or zero is a matrix with all elements 0.

$$\begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix}$$

Definition 15.43. Square matrix is a matrix where $m = n$

In a square matrix, we define the following as well:

Definition 15.44. (1) a_{ii} are called the diagonal elements

(2) a_{ij} and a_{ji} are the conjugate elements

(3) $\sum_{i=1}^n a_{ii}$ is called the trace of the matrix

While the square matrix is in itself very spacial, we define some more special matrix within it as well.

Definition 15.45. A Triangular Matrix is a square matrix with elements on only one side of the diagonal. For example:

$$\begin{bmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

is an lower triangular matrix while:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix}$$

is an upper triangular matrix.

We need to note that the determinant of a triangular matrix is the product of its diagonal elements. That is $a_{11} \cdot a_{22} \cdot a_{33}$ in the above examples.

Definition 15.46. A diagonal matrix is a square matrix with all non-diagonal elements being 0. For example:

$$\begin{bmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{bmatrix}$$

is a diagonal matrix. Its determinant is also the product of the diagonal elements.

Definition 15.47. A scalar matrix is a diagonal matrix with all elements on the diagonal equal. For example:

$$\begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$$

Is a scalar matrix. It's determinant is a^3

Definition 15.48. The Identity matrix is a scalar matrix with $a = 1$. The scalar matrix of order three is:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

That concludes the first round of definitions.

15.6 Arithmetic of Matrices

Two matrices are said to be equal if they have the same number of rows and column(order of matrix) and the corresponding terms are equal in the rows and columns are equal.

We can only add matrices of same order. The addition of matrix A and B is just creating a new matrix C where: $c_{ij} = a_{ij} + b_{ij}$

Let's solve an example to understand.

Example 15.49 (Motivating Example). $\begin{bmatrix} 3 & 9 & 10 & 3 \\ 4 & 4 & 7 & 5 \end{bmatrix} + \begin{bmatrix} 10 & 6 & 8 & 10 \\ 3 & 6 & 2 & 3 \end{bmatrix}$

Solution. The sum is $\begin{bmatrix} 3+10 & 9+6 & 10+8 & 3+10 \\ 3+4 & 4+6 & 7+2 & 5+3 \end{bmatrix}$
 $= \begin{bmatrix} 13 & 15 & 18 & 13 \\ 7 & 10 & 9 & 8 \end{bmatrix}$

□

The multiplication of a matrix by a constant follows as:

Definition 15.50. $K \cdot \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$
 $= \begin{bmatrix} Ka_{11} & Ka_{12} & Ka_{13} \\ Ka_{21} & Ka_{22} & Ka_{23} \\ Ka_{31} & Ka_{32} & Ka_{33} \end{bmatrix}$

We need to realize that this is different from the determinant multiplication by scalar as that only was multiplied to one row or column, while here K is multiplied to every element. We will now discuss the most important part of matrices, matrix multiplication to matrix.

Definition 15.51. We define matrix multiplication as $A_{m \times n} \times B_{n \times p} = C_{m \times p}$ where A, B and C are matrices.

A is called the pre-multiplier while B is the post multiplier. two Matrices can only be multiplied if number of columns of pre-multiplier is equal to number of rows of post multiplier.

How do we actually do the multiplication? We multiply the rows of the pre multiplier to the columns of the post multiplier.

For example $\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \times \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix} = \begin{bmatrix} a_{11} \cdot b_{11} + a_{12} \cdot b_{21} & a_{11} \cdot b_{12} + a_{12} \cdot b_{22} \\ a_{21} \cdot b_{11} + a_{22} \cdot b_{21} & a_{21} \cdot b_{12} + a_{22} \cdot b_{22} \\ a_{31} \cdot b_{11} + a_{32} \cdot b_{21} & a_{31} \cdot b_{12} + a_{32} \cdot b_{22} \end{bmatrix}$

Not the prettiest thing, but I hope you can somewhat understand what's going on. I have given two very simple examples for you to solve to check if you have understood the concept.

Example 15.52 (Motivating Example). $\begin{bmatrix} 2 & 3 & 4 \\ 1 & 2 & 3 \end{bmatrix} \times \begin{bmatrix} 1 & 0 & 1 & 2 \\ 2 & 1 & 1 & 2 \\ 0 & 1 & -1 & 3 \end{bmatrix}$

Example 15.53 (Motivating Example). $\begin{bmatrix} 1 & 0 & 2 & 3 \end{bmatrix} \times \begin{bmatrix} 1 & 0 & 1 & 2 \\ 2 & 1 & 1 & 2 \\ 0 & 1 & -1 & 3 \end{bmatrix}$

Here are some properties of matrix multiplication:

Theorem 15.54. Here A, B, C all represent Matrices whose product is defined

- (1) It is not commutative. In general $AB \neq BA$.
This leads to a peculiar form of $(A + B)^2 = A^2 + AB + BA + B^2$
- (2) It is associative. $(A \times B) \times C = A \times (B \times C)$
- (3) It distributes over addition. $A \times (B+C) = A \times B + A \times C$ and $(B+C) \times A = B \times A + C \times A$
- (4) The identity matrix I which can multiply A will show the following
 $I \times A = A \times I = A$
- (5) Any matrix multiplied with null matrix gives a null matrix. However the the converse is not true. If $A \times B$ is null matrix then it is not necessary that either A or B will be null matrix.
- (6) Laws of exponents hold as the are:

$$A^m \times A^n = A^{m+n}$$

$$A^n = A^{n-1} \times A$$

$$A^{mn} = A^{mn}$$

A very common question which comes using these properties is:

Example 15.55. If $A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ then A^{2023} is equal to:

Solution. I'll first show you the solution which a lot of books have. Which while great for competitive speed papers, is not acceptable in written example.

Notice that:

$$A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \times \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

This establishes a pattern, which means $A^{2023} = \begin{bmatrix} 1 & 0 \\ 2023 & 1 \end{bmatrix}$

You can see that this is a very bad explanation. The actual way to do it is to prove that $A^n = \begin{bmatrix} 1 & 0 \\ n & 1 \end{bmatrix}$ using Induction. The first one was engineers induction(see Power Overwhelming), explains quite well why such questions occur in engineering entrance exams the most.

□

15.7 The bridge

We will now start the process of connecting Matrices with determinants. We already know that we can find determinant of a square matrix. Using that we can claim the following:

Theorem 15.56. *If A and B are two square matrices of same order then $|A \times B| = |A| \times |B|$*

Also if we remember the multiplication of matrices and determinants by a constant and more specifically their one difference, we can also say:

Theorem 15.57. *If A_n is a square matrix of order n and K is a constant then: $|KA_n| = K^n|A_n|$*

You may also remember the transposing of determinant. We'll define it formally here.

Definition 15.58. Matrix obtained by interchanging rows and columns is called transpose of matrix, for some matrix A it is denoted by A^T . $A =$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$\therefore A^T = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{bmatrix}$$

Using the definition we can also notice the following facts:

Theorem 15.59. (1) $(A^T)^T = A$

(2) $(A + B)^T = A^T + B^T$

(3) $(KA)^T = K(A^T)$ where K is a constant

(4) $(AB)^T = B^T A^T$

(5) $(ABC)^T = C^T B^T A^T$

(6) $(A^n)^T = (A^T)^n$

15.8 Some more special matrices

Now that we have defined transpose, we can define some more special matrices on the basis of it.

Definition 15.60. Symmetric matrix: If $(A_n)^T = A_n$ for a square matrix A_n then it is called a symmetric matrix. Basically, $a_{ij} = a_{ji}$

Definition 15.61. Skew Symmetric Matrix: If $(A_n)^T = -A_n$ for a square matrix A_n then it is called a skew symmetric matrix. Basically, $a_{ij} = -a_{ji}$

With this much, we can start proving almost surprising facts:

Example 15.62. Prove that for any square matrix A , $\frac{1}{2}(A + A^T)$ is symmetric matrix and $\frac{1}{2}(A - A^T)$ is skew symmetric matrix.

Proof. We just take the transpose.

$$\begin{aligned} & \left(\frac{1}{2}(A + A^T)\right)^T \\ &= \frac{1}{2}(A^T + A) \text{ which makes it symmetric.} \\ & \left(\frac{1}{2}(A - A^T)\right)^T \\ &= \frac{1}{2}(A - A^T) \\ &= \frac{-1}{2}(A^T - A) \text{ which makes it skew-symmetric.} \end{aligned}$$

□

This also leads to a surprising fact: Every square matrix A can be represented as a sum of symmetric and skew symmetric matrix. Here is an example for you to solve

Example 15.63. If A and B are symmetric matrices of same order then prove that $AB - BA$ is skew symmetric matrix.

Definition 15.64. A square matrix is orthogonal if $AA^T = I$

We can note that the determinant of A must be ± 1

Expanding the multiplication gives us: the sum of squares of elements in any row or column is one and the pairwise product and sum of two rows or columns is zero.

The last one allows us to detect Orthogonal matrices in the wild. For example in this Question from IIT.

Example 15.65. (IIT 2005) Find $P^T Q^{2005} P$, where $P = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ \frac{-1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$, $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $Q = PAP^T$

We need to only notice that P is an orthogonal matrix. After that, the question basically solves itself.

Definition 15.66. A square matrix is called idempotent if $A^2 = A$. Clearly, A^n will also be equal to A for all $n \geq 2$

At this time I think it is also important for you to know that $(A + I)^n$ can be opened like a binomial expansion. This can only be done for $(A + I)$ not for $(A + B)$. This is because $A \times I = I \times A = A$

And here I ask you to ponder before reading the solution:

Example 15.67. If A is an idempotent matrix then $(I + A)^n =$

Solution. Expanding using binomial theorem, $I^n + \binom{n}{1} I^{n-1} A + \cdots + A^n = I + (2^n - 1)A$ Using the fact that $I^n = I$ and $A \times I = A$ and $\sum_{n=1}^n \binom{n}{0} = 2^n - 1$. The last one as you may recall comes from combinatorics.

□

And we end this section with a few miscellaneous definitions.

Definition 15.68. A square matrix is called involutory if $A^2 = I$

Definition 15.69. A square matrix is called nilpotent matrix of order m if: $A^m = 0$ and $A^{m-1} \neq 0$

Currently, there is no way to check whether matrix is nilpotent or not, other than checking the powers manually.

Definition 15.70. A matrix is called singular if its determinant is zero, otherwise it is called non-singular.

15.9 Adjoint and Inverse of Matrices

Remember the co-factor matrix we had studied earlier? We'll use it in a minute to find inverse of a matrix, the one thing for which we have literally studied matrix for.

But first let's talk about adjoint of a matrix.

Definition 15.71. For any square matrix, its adjoint is defined as transpose of its cofactor matrix.

Using whatever we know about the co-factor matrix and about transpositions, we'll get at the following properties of adjoint matrices:

Theorem 15.72. For square matrices A and B of order n , we have:

$$(1) \quad |\text{adj}A| = |A|^{n-1}$$

$$(2) \quad \text{adj}(\text{adj}A) = |A|^{n-2} A$$

$$(3) \quad \text{adj}(A^T) = (\text{adj}A)^T$$

$$(4) \quad \text{adj}(KA) = K^{n-1}(\text{adj}A)$$

$$(5) \quad \text{adj}(A^n) = (\text{adj}A)^n$$

$$(6) \quad \text{adj}(AB) = (\text{adj}B)(\text{adj}A)$$

Here is the reason why we learnt about the adjoint:

Definition 15.73. Square matrix B_n is called inverse matrix of A_n if: $AB = BA = I$

Clearly, if B is inverse of A then A is also inverse of B . Formula for $A^{-1} = \frac{1}{|A|} \times \text{adj}(A)$

Clearly, this makes singular matrices have no inverses.

At this point I recommend you taking the inverse of a random 3×3 matrix. While, we have done all the operations and transformations previously, doing it only once will give you some amount of confidence in what to do. Here are some properties of the inverse:

Theorem 15.74. (1) $|A^{-1}| = \frac{1}{|A|}$

$$(2) \quad (A^T)^{-1} = (A^{-1})^T$$

$$(3) \ adj(A^{-1}) = (adjA)^{-1} = \frac{A}{|A|}$$

$$(4) \ (AB)^{-1} = B^{-1}A^{-1}$$

These properties wreck questions such as:

Example 15.75. (JEE Mains 2014) If A is a 3×3 non-singular matrix such that $AA^T = A^TA$ and $B = A^1A^T$ then $BB^T =$

Solution. This question is perfect as it uses a good number of the properties we discussed.

$$\begin{aligned} BB^T &= (A^{-1}A^T)(A^{-1}A^T)^T \\ &= A^{-1}A^T(A^T)^T(A^{-1})^T \\ &= A^{-1}(A^T(A^T)^T)(A^{-1})^T \\ &= A^{-1}(A^TA)(A^T)^{-1} \\ &= A^{-1}I(A^T)^{-1} \\ &= IA^{-1}(A^T)^{-1} \\ &= I^2 \\ &= I \end{aligned}$$

□

15.10 System of linear Equations using Matrices

Now we finally see the back end of Crammer's Rule.

But before that let's go on a little tangent.

Suppose we have three matrices A, X, B where A and X are multiply-able and A is invertible(has an inverse, or is non-singular)

If $AX = B$ then can we just divide both sides by A ? Obviously not. We can't truly divide matrices. But we can instead premultiply both the sides by A^{-1} to get $A^{-1}AX = A^{-1}B$ which simplifies to $IX = A^{-1}B \iff X = A^{-1}B$

We need to note this is not the same as dividing as the order of the multiplication matters. If we, in confusion, take $X = BA^{-1}$, we will get a wrong answer. But how is all this related? We need to notice that for a system of equations:

$$\begin{aligned} a_1x + b_1y + c_1z &= d_1 \\ a_2x + b_2y + c_2z &= d_2 \\ a_3x + b_3y + c_3z &= d_3 \end{aligned}$$

Is in all ways and forms equivalent to $\begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix} \times \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix}$

This is exactly the case we just discussed above. While we can use inverse to

find the system of solutions and in school exams you have to do that(Don't for the love of god use Crammer's rule here, you will lose marks). However, competitively, Crammer's rule is much quicker form of doing the same.

Also to be complete here are the solution conditions for the inverse matrix form.

Theorem 15.76. Let's denote $\begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$ as A and $\begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix}$ as B If $|A| \neq 0$,

we have one unique solution.

If $|A| = 0$ and $(adj A) \times B = 0$ then infinitely many solutions provided that all coefficients are not 0

If $|A| = 0$ and $(adj A) \times B \neq 0$ then no solutions exist.

15.11 Characteristic Equation and Cayley Hamilton Theorem

Definition 15.77. If A is any square matrix, then $|AxI| = 0$ is called its characteristic equation. Roots of Characteristic Equation are called as Eigen values or Characteristic roots

For example the characteristic equation $A = \begin{bmatrix} 8 & 6 \\ 5 & 9 \end{bmatrix}$ is:

$$|A - xI| = 0$$

$$\Leftrightarrow \begin{vmatrix} 8-x & 6 \\ 5 & 9-x \end{vmatrix} = 0$$

$$\Leftrightarrow (8-x)(9-x) - 30 = 0$$

$$\Leftrightarrow (72 - 17x + x^2) - 30 = 0$$

$$\Leftrightarrow x^2 - 17x + 42 = 0$$
 is the characteristic equation of A.

Like recursion, this is called the characteristic equation as it is satisfied by only and only A. In this case it means, $A^2 - 17A + 42I = 0$

The fact that every matrix satisfies it's characteristic equation is known as Cayley-Hamiltonian theorem. We can observe from our solving and using vieta, the sum of its eigen values is equal to the trace and the product of the eigen values is equal to the determinant.

Also using the properties of A^{-1} we can say that if λ is an eigen value of A then $\frac{1}{\lambda}$ is an eigen value of A^{-1}

All this is interesting, but what is the use?

Glad you asked:

Example 15.78. If $A = \begin{bmatrix} 1 & 0 & 2 \\ 1 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$ then find k such that $A^3 - kA^2 + 7A + 2I = 0$

Solution. Let's find the characteristic equation.

$$\begin{aligned} & \begin{bmatrix} 1-x & 0 & 2 \\ 0 & 2-x & 1 \\ 2 & 0 & 3-x \end{bmatrix} = 0 \\ \iff & (2-x)(3-x)(1-x) - 4(2-x) = 0 \\ \iff & -x^3 + 6x^2 - 7x - 2 = 0 \\ \implies & A^3 - 6A^2 + 7A + 2I = 0 \end{aligned}$$

Thus, $k = 6$

□

15.12 Netflix and Spotify and Matrices...

Okay, so you know when Netflix is like, 'Hey, watch this show!' or when Spotify suggests your next favorite jam? Well, behind the scenes, there's some linear algebra happening. Netflix has a huge matrix where rows are people and columns are movies or shows. Each cell is like a 'how much they like it' score.

Using inverses and multiplication, They break this table into two smaller tables – one for people's taste (call it 'U' for users. One is linked with every account) and one for awesomeness of a show (we'll call it 'V' for value, one is linked with every movie).

If Netflix wants to find the right show for the right user so they multiply your U with the shows V and takes the determinant. The higher the value, the better the fit is. This might seem simple in speaking but under the hood this is a very complicated algorithm which whose nitty and gritty are better suited for a computing book. Also a lot of it is proprietary, or top secret, so it's not possible for us to know everything but this is the gist of it.

This is exactly how websites are ranked by assigning a matrix to the search quarry and a matrix to the site and then multiplying and taking determinant. We also use it in geometry and physics as we'll see later

Exercises

$$(1) \begin{vmatrix} \sin(2x) & 1 - \cos(2x) & 2\sin(x) \\ \cos(x) & \sin(x) & 1 \\ \sin(x) & \cos(x) & 1 \end{vmatrix}$$

- (2) (JEE Mains 2020) Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two 3×3 real matrices such that $b_{ij} = 3^{i+j-2}a_{ji}$, where $i, j = 1, 2, 3$. If the determinant of B is

81, then the determinant of A is:

(3) If $\Delta_r = \begin{vmatrix} 4 & 612 & 915 \\ 101r^2 & 2r & 3r \\ r & \frac{1}{r} & \frac{1}{r^2} \end{vmatrix}$ then the value of $\lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{r=1}^n \Delta_r$

(4) If $s = (a+b+c)$, then the value of $\begin{vmatrix} s+c & a & b \\ c & s+a & b \\ c & a & s+b \end{vmatrix}$ is (in terms of s):

(5) (IIT 2011) The Real roots of $\begin{vmatrix} \sin(x) & \cos(x) & \cos(x) \\ \cos(x) & \sin(x) & \cos(x) \\ \cos(x) & \cos(x) & \sin(x) \end{vmatrix}$ in the interval $\frac{-\pi}{4} \leq x \leq \frac{\pi}{4}$ is (are):

(6) (JEE Mains 2020) Let $a - 2b + c = 1$, If $f(x) = \begin{vmatrix} x+a & x+2 & x+1 \\ x+b & x+3 & x+2 \\ x+c & x+4 & x+3 \end{vmatrix}$,

then find the algebraic form of $f(x)$

(7) Find values of p, q such that $x+y+z = 6; 2x+5y+pz = q; x+2y+3z = 14$

(a) has unique solution

(b) has infinitely many solutions

(8) If 't' is real and $\lambda = \frac{t^2-3t+4}{t^2+3t+4}$ then find number of solution of $3xy+4z=3$
 $x+2y+3z=2$
 $6x+5y+\lambda z=3$

(9) (JEE Mains 2020) The system of equation $3x+4y+5z=\mu, x+2y+3z=1, 4x+4y+4z=\delta$ is inconsistent, then (δ, μ) can be

(10) For matrices A, B , If $AB = A$ and $BA = B$ then $B^2 =$

(11) If A and B are square matrices of order 3 such that $|A| = -1, |B| = 3$, then the determinant of $2A^3B^2$ is equal to:

(12) If P is a 3×3 matrix such that $P^T = 2P + I$ then prove that $P + I = 0$

(13) Let A be the set of all 3×3 matrices which are symmetric with entries 0 or 1. If there are five 1's and four 0's, then number of matrices in A is:

(14) If $A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & -2 & 1 \\ 4 & 2 & 1 \end{bmatrix}$, then find K such that $A^3 - kA - 40I = 0$

(15) (Poh-Shen Loh) Calculate the determinant of

$$\begin{vmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 5 & 3 & 8 & 1 & 9 & 9 \\ 6 & 5 & 1 & 1 & 6 & 6 & 4 \\ 1 & 1 & 3 & 3 & 8 & 5 & 6 \\ 3 & 2 & 7 & 8 & 9 & 9 & 8 \end{vmatrix}$$

16

Inequalities Revisited

We finally will now learn about some more advanced methods of solving inequalities and summations. The last 90 to 100 pages, while essential in their own right, were mostly so that we can do this.

16.1 Rearrangement Inequalities

Theorem 16.1 (Rearrangement Inequality). *If $a_1 \geq \dots \geq a_n$ and $b_1 \geq \dots \geq b_n$ then for any permutation (rearrangement) c_1, \dots, c_n of b_1, \dots, b_n ,*

$$a_1b_1 + \dots + a_nb_1 \leq a_1c_1 + \dots + a_nc_n \leq a_1b_1 + \dots + a_nb_n$$

Proof. The proof of the Rearrangement Inequality can be handled with proof by contradiction. We will prove the maximization first, the minimization will follow from that.

Let us first consider the case where $n = 2$. We can take $a_1 \geq a_2$ and $b_1 \geq b_2$.

$$\begin{aligned} & \therefore (a_1 - a_2)(b_1 - b_2) \geq 0 \\ \iff & a_1b_1 + a_2b_2 - a_1b_2 - a_2b_1 \geq 0 \\ \iff & a_1b_1 + a_2b_2 \geq a_1b_2 + a_2b_1 \end{aligned}$$

Now for the general case. Let $a_1 \geq a_2 \geq \dots \geq a_n$ and $b_1 \geq b_2 \geq \dots \geq b_n$; and let's to the contrary assume that in the grouping maximizing the sum, a_m is not paired with b_m . We'll instead assume that a_m is paired with b_l and b_m is paired with a_l .

Hence we are claiming, $a_m b_l + a_l b_m \geq a_m b_m + a_l b_l$ which is untrue as we showed above.

The minimization equality can be very easily proved by noting that if we

have the set $\{-b_1, -b_2, \dots, -b_n\}$, ordered in increasing order (which makes $b_1 \geq b_2 \geq b_3 \dots \geq b_n$) and the set $\{a_1, a_2, \dots\}$, ordered in decreasing order, then the maximum sum is just $-a_1b_n - a_2b_{n-1} + \dots$. Whose negative is $a_1b_n + \dots + a_nb_1$ which will be the minimum possible value.

□

A more refined form of the rearrangement inequality is

Theorem 16.2 (Chebyshev's Inequality). *if $a_1 \geq a_2 \geq \dots \geq a_n$ and $b_1 \geq b_2 \geq \dots \geq b_n$ then the following inequality holds:*

$$n(\sum_{i=1}^n a_i b_i) \geq (\sum_{i=1}^n a_i)(\sum_{i=1}^n b_i).$$

On the other hand, if $a_1 \geq a_2 \geq \dots \geq a_n$ and $b_n \geq b_{n-1} \geq \dots \geq b_1$ then:

$$n(\sum_{i=1}^n a_i b_i) \leq (\sum_{i=1}^n a_i)(\sum_{i=1}^n b_i).$$

Proof. The proof is simple. We know that $\sum_{i=1}^n a_i b_i$ is maximal.

$$\therefore \sum_{i=1}^n a_i b_i \geq a_1 b_1 + a_2 b_2 + \dots + a_n b_n$$

$$\sum_{i=1}^n a_i b_i \geq a_1 b_2 + a_2 b_3 + \dots + a_n b_1$$

⋮

$$\sum_{i=1}^n a_i b_i \geq a_1 b_n + a_2 b_1 + \dots + a_n b_{n-1}$$

Adding them will give us the inequality.

□

These two inequalities make quick work of even IMO problems. Case in point:

Example 16.3. (IMO 1975) Let x_i, y_i ($i = 1, 2, \dots, n$) be real numbers such that

$$x_1 \geq x_2 \geq \dots \geq x_n \text{ and } y_1 \geq y_2 \geq \dots \geq y_n.$$

Prove that, if z_1, z_2, \dots, z_n is any permutation of y_1, y_2, \dots, y_n , then

$$\sum_{i=1}^n (x_i - y_i)^2 \leq \sum_{i=1}^n (x_i - z_i)^2.$$

Proof. To the untrained eye, this seems quite bad. But let's expand.

$$\sum_{i=1}^n (x_i - y_i)^2 \leq \sum_{i=1}^n (x_i - z_i)^2.$$

$$\iff \sum_{i=1}^n (x_i^2 + y_i^2 - 2x_i y_i) \leq \sum_{i=1}^n (x_i^2 + z_i^2 - 2x_i z_i) \text{ as } y_1^2 + \dots + y_n^2 = z_1^2 + \dots + z_n^2 \text{ due to the fact that } z \text{ is just a rearrangement of } y.$$

$$\iff \sum_{i=1}^n -2x_i y_i \leq \sum_{i=1}^n -2x_i z_i$$

$$\iff \sum_{i=1}^n x_i y_i \geq \sum_{i=1}^n x_i z_i$$

which is just the rearrangement inequality. And we are done

□

16.2 Inequalities in Arbitrary Functions

Till now our inequalities dealt in well understood functions like square, cube, average, reciprocal etc. But what about when the inequality deals in a function like $\sin(x)$ or $\ln x$ or maybe $\sin(\ln x)$, what do we do now?

Before we look at the actual inequalities, we'll define two new terms.

Definition 16.4. A convex function is a continuous function whose value at the midpoint of every interval in its domain does not exceed the arithmetic mean of its values at the ends of the interval.

Definition 16.5. A concave function is the opposite of a convex function, i.e. a function f is concave if and only if $-f$ is convex.

Basically, A concave function is a continuous function whose value at the midpoint of every interval in its domain exceeds the arithmetic mean of its values at the ends of the interval.

A simple way to remember is that happy face is convex, while a sad face is concave. We can basically remember that happy face is convex and sad face




Figure 16.0. Convex and Concave graphs

is concave.

For a function, whose graph is harder to plot, we define a function as convex if $f''(x) \geq 0$ and concave if $f''(0) \leq 0$. This allows us to talk about two more powerful inequalities

Theorem 16.6 (Jensen Inequality). *Let F be a convex function of one real variable. Let $x_1, \dots, x_n \in \mathbb{R}$ and let $a_1, \dots, a_n \geq 0$ satisfy $a_1 + \dots + a_n = 1$. Then*

$$F(a_1x_1 + \dots + a_nx_n) \leq a_1F(x_1) + \dots + a_nF(x_n)$$

If F is a concave function, the inequality reverses.

Proof. We only prove the case where F is concave. The proof for the other case follows from the fact that $-F$ is convex.

Let $\bar{x} = \sum_{i=1}^n a_i x_i$. As F is concave, its slope F' is decreasing. We consider two cases. If $x_i \leq \bar{x}$, then

$$\int_{x_i}^{\bar{x}} F'(t) dt \geq \int_{x_i}^{\bar{x}} F'(\bar{x}) dt.$$

If $x_i > \bar{x}$, then

$$\int_{\bar{x}}^{x_i} F'(t) dt \leq \int_{\bar{x}}^{x_i} F'(\bar{x}) dt.$$

By the fundamental theorem of calculus, we have

$$\int_{x_i}^{\bar{x}} F'(t) dt = F(\bar{x}) - F(x_i).$$

Evaluating the integrals, each of the last two inequalities implies the same result:

$$F(\bar{x}) - F(x_i) \geq F'(\bar{x})(\bar{x} - x_i)$$

so this is true for all x_i . Then we have

$$\begin{aligned} & F(\bar{x}) - F(x_i) \geq F'(\bar{x})(\bar{x} - x_i) \\ \implies & a_i F(\bar{x}) - a_i F(x_i) \geq F'(\bar{x})(a_i \bar{x} - a_i x_i) \quad \text{as } a_i > 0 \\ \implies & F(\bar{x}) - \sum_{i=1}^n a_i F(x_i) \geq F'(\bar{x}) \left(\bar{x} - \sum_{i=1}^n a_i x_i \right) \quad \text{as } \sum_{i=1}^n a_i = 1 \\ \implies & F(\bar{x}) \geq \sum_{i=1}^n a_i F(x_i) \quad \text{as } \bar{x} = \sum_{i=1}^n a_i x_i \end{aligned}$$

as desired. \square

We need to note that the weighted AM-GM follows from this.
Also we can rewrite Jensen as $\frac{f(a_1) + \dots + f(a_n)}{n} \geq f\left(\frac{a_1 + \dots + a_n}{n}\right)$ for f is convex and reverse if f is concave.

Theorem 16.7 (Karamata's Inequality). *If f is convex and (x_n) majorizes (y_n) then:*

$$f(x_1) + \dots + f(x_n) \geq f(y_1) + \dots + f(y_n)$$

The reverse inequality hold true when f is concave.

Proof. We will use the fact that $\frac{f(x)-f(y)}{x-y}$ or the slope of graph is decreasing when it is convex. Also using the fact that addition is commutative, we can, without loss of generality assume that $x_1 \geq x_2 \geq \dots \geq x_n$ and $y_1 \geq y_2 \geq \dots \geq y_n$.

We can use that to say: $c_{i+1} = \frac{f(x_{i+1}) - f(y_{i+1})}{x_{i+1} - y_{i+1}} \leq \frac{f(x_i) - f(y_i)}{x_i - y_i} = c_i$
 We will also define $A_i = x_1 + \dots + x_i$ and $B_i = y_1 + \dots + y_i$

$$\begin{aligned} \sum_{i=1}^n (f(x_i) - f(y_i)) &= \sum_{i=1}^n c_i(x_i - y_i) \\ &= \sum_{i=1}^n c_i \left(\underbrace{A_i - A_{i-1}}_{=x_i} - \underbrace{(B_i - B_{i-1})}_{=y_i} \right) \\ &= \sum_{i=1}^n c_i(A_i - B_i) - \sum_{i=1}^n c_i(A_{i-1} - B_{i-1}) \\ &= c_n \underbrace{(A_n - B_n)}_{=0} + \sum_{i=1}^{n-1} \underbrace{(c_i - c_{i+1})(A_i - B_i)}_{\geq 0} - c_1 \underbrace{(A_0 - B_0)}_{=0} \\ &\geq 0, \end{aligned}$$

The proof for concave is similar. □

Note that both the proofs are only given for completeness. You can clearly get away with not memorizing them as long as you know the actual inequality. And here is a question from the IMO shortlist to show how powerful inequalities become:

Example 16.8. (IMOSL 2009) Given $a + b + c = \frac{1}{a} + \frac{1}{b} + \frac{1}{c}$, Prove that $\frac{1}{(2a+b+c)^2} + \frac{1}{(a+2b+c)^2} + \frac{1}{(a+b+2c)^2} \leq \frac{3}{16}$

Proof. Using the fact that $a + b + c = \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \iff \frac{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}{a+b+c} = 1$

$$\frac{1}{(2a+b+c)^2} + \frac{1}{(a+2b+c)^2} + \frac{1}{(a+b+2c)^2} \leq \frac{3}{16} \cdot \frac{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}{a+b+c}$$

As the inequality is homogeneous, we can set $a + b + c = 3$

This leaves us with:

$$\sum_{cyc} \frac{1}{(3+a)^2} \leq \sum_{cyc} \frac{1}{16a}$$

$$\iff 0 \leq \sum_{cyc} \frac{1}{16a} - \frac{1}{(3+a)^2}$$

We define $f(x) = \frac{1}{16x} - \frac{1}{(3+x)^2}$, which is convex.

Then using Jensen, $\frac{1}{16a} - \frac{1}{(3+a)^2} + \frac{1}{16b} - \frac{1}{(3+b)^2} + \frac{1}{16c} - \frac{1}{(3+c)^2} \geq 3 \left(\frac{1}{16} - \frac{1}{(3+\frac{a+b+c}{3})^2} \right)$

As $a + b + c = 3$,

$$\sum_{cyc} \frac{1}{16a} - \frac{1}{(3+a)^2} \geq 3 \left(\frac{1}{16} - \frac{1}{4^2} \right)$$

Which gives us

$$\sum_{cyc} \frac{1}{16a} - \frac{1}{(3+a)^2} \geq 0$$

Which proves the inequality. \square

16.3 Tangent Line Trick

For some homogeneous inequality, if we can prove that the function is above the tangent line at the equality value for the sum being 1, we can prove the inequality by summation.

This explanation seems more complicated than the trick actually is. Here is a question to try it out

Example 16.9. (USAMO 2003) Let a, b, c be positive real numbers. Prove that

$$\frac{(2a + b + c)^2}{2a^2 + (b + c)^2} + \frac{(2b + c + a)^2}{2b^2 + (c + a)^2} + \frac{(2c + a + b)^2}{2c^2 + (a + b)^2} \leq 8.$$

Proof. Since, the inequality is homogeneous we may assume that $a+b+c = 1$ and $0 < a, b, c < 1$.

The first time on the LHS is the inequality will be:

$$f(a) = \frac{(a+1)^2}{2a^2 + (1-a)^2} = \frac{a^2 + 2a + 1}{3a^2 - 2a + 1}$$

Note that equality holds when $a = b = c = 1/3$.

We can differentiate to git that the tangent has the equation of the form $y = \frac{12x+4}{3}$.

So we claim that

$$f(a) = \frac{a^2 + 2a + 1}{3a^2 - 2a + 1} \leq \frac{12a + 4}{3} \text{ for } 0 < a < 1$$

Upon clearing the denominators, it is equivalent to:

$$36a^3 - 15a^2 - 2a + 1 \geq 0$$

Note that since the curve and the line intersect at $1/3$, $3a - 1$ would be a factor.

$$36a^3 - 15a^2 - 2a + 1 = (3a - 1)^2(4a + 1) \geq 0 \text{ for } 0 < a < 1$$

Adding the similar inequalities for b and c gives:

$$f(a) + f(b) + f(c) \leq \frac{12(a + b + c) + 12}{3} = 8$$

\square

While I recommend trying Jensen and Karamarta before pulling out the Tangent Trick. However, when nothing else works, kill with calc.

16.4 SEBACS Generalized

Theorem 16.10 (Holder's Inequality). *If $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n, z_1, z_2, \dots, z_n$ are non-negative real numbers and $\lambda_a, \lambda_b, \dots, \lambda_z$ are non-negative reals with sum of 1, then*

$$(a_1 + \dots + a_n)^{\lambda_a} (b_1 + \dots + b_n)^{\lambda_b} \cdots (z_1 + \dots + z_n)^{\lambda_z} \geq a_1^{\lambda_a} b_1^{\lambda_b} \cdots z_1^{\lambda_z} + \dots + a_n^{\lambda_a} b_n^{\lambda_b} \cdots z_n^{\lambda_z}.$$

Note that with two sequences \mathbf{a} and \mathbf{b} , and $\lambda_a = \lambda_b = 1/2$, this is SEBACS inequality.

Proof. This is one the most easy proof in this book. We need to notice that $(a_1 + \dots + a_n), (b_1 + \dots + b_n), \dots, (z_1 + \dots + z_n)$ are homogeneous, allowing us to take $a_1 + \dots + a_n = b_1 + \dots + b_n = \dots = z_1 + \dots + z_n = 1$, converting the inequality to:

$$1 \geq a_1^{\lambda_a} b_1^{\lambda_b} \cdots z_1^{\lambda_z} + \dots + a_n^{\lambda_a} b_n^{\lambda_b} \cdots z_n^{\lambda_z}$$

Which is trivially true as $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n, z_1, z_2, \dots, z_n$ are non-negative real numbers. \square

However, sadly SEBACS tends to be used more often. Also I am yet to see any inequality which can be done by Holder but not using Jensen or Karmata. However, here is an example(which can be done using Jensen, and I encourage you to do that)

Example 16.11. Let a, b, c be positive real numbers. Prove that $\frac{a}{\sqrt{a^2+8bc}} + \frac{b}{\sqrt{b^2+8ca}} + \frac{c}{\sqrt{c^2+8ab}} \geq 1$.

Proof. Jensen does this using $abc = 1$ and then the question dies in silence. However, for Holder, we'll need to 'observe' that:

$$a(a^2 + 8bc) \cdot \left(\frac{a}{\sqrt{a^2+8bc}}\right)^2 = a^3$$

Which we can use to say, using Holder:

$$\sum_{cyc} a(a^2 + 8bc) \cdot \sum_{cyc} \left(\frac{a}{\sqrt{a^2+8bc}}\right)^2 \geq \sum_{cyc} a^3$$

$$\iff \sum_{cyc} \left(a(a^2 + 8bc)\right)^{\frac{1}{3}} \cdot \sum_{cyc} \left(\left(\frac{a}{\sqrt{a^2+8bc}}\right)\right)^{\frac{2}{3}} \geq a + b + c$$

Which will change our to prove to:

$$(a+b+c)^3 \geq \sum_{cyc} a(a^2 + 8bc)$$

$$\iff a^3 + b^3 + c^3 + 3(a+b)(b+c)(c+a) \geq a^3 + b^3 + c^2 + 24abc$$

$$\iff 3(a+b)(b+c)(c+a) \geq 24abc$$

$$\iff (a+b)(b+c)(c+a) \geq 8abc$$

Notice that $a+b \geq 2\sqrt{ab}$ and similar for the others, leading to the multiplication:

$$(a+b)(b+c)(a+c) \geq 2\sqrt{ab} \cdot 2\sqrt{bc} \cdot 2\sqrt{ac}$$

$$\iff (a+b)(b+c)(a+c) \geq 8\sqrt{a^2b^2c^2}$$

$$\iff (a+b)(b+c)(a+c) \geq 8abc$$

\square

16.5 Lagrange Multipliers

This is the most powerful trick/methodtheorem about inequalities. This is the reason why I can solve almost any inequality in an instant. Is this almost cheating? YES. Is it allowed if you write it up correctly? YES!

Lagrange multiplier finds the extreme of multi variable functions given a constraint.

Example 16.12 (Motivating Example). Maximize $f(x, y) = 2x + y$ for $x^2 + y^2 = 1$.

Solution. Before we go further, I know that this example is doable with basic calculus as well. However, let's try to explore a new technique.

We'll use a concept called gradients(∇f), vectors(lines) which are perpendicular to the tangent of a graph.

This is useful as the extremal value will occur at the point where $f(x, y) = 2x + y$ is perpendicular to $g(x, y) = x^2 + y^2 = 1$

While in this case the lines are straight, sometimes we end up with a much




Figure 16.0. The point of maxima and minima

more curvy line, however the observation holds.

Another key observation to make is that if two graphs are tangent, the gradient lines at point of tangency are parallel. We can therefore say that, $\nabla f(x, y) = \lambda \nabla g(x, y)$ as they are parallel.

We now need to calculate the gradient. The gradient can be written as a col-

umn matrix $\begin{bmatrix} \frac{\partial f(x,y)}{\partial x} \\ \frac{\partial f(x,y)}{\partial y} \end{bmatrix}$ where $\frac{\partial}{\partial x}$ means the derivative of $f(x, y)$ considering

everything else other than x as a constant.

Why does this definition work? That should keep you curious till the Calc III course in college.

But with this in hand, we can solve our question:

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix} = \lambda \begin{bmatrix} 2x \\ 2y \end{bmatrix}$$

Using the definition of equal matrices, $2\lambda x = 2; 2\lambda y = 1$

Which gives us $x = 2y$ which means $4y^2 + y^2 = 1$

$$\Leftrightarrow y = \pm \frac{1}{\sqrt{5}}$$

Which means the maxima of $2x + y = 5y$ is $\sqrt{5}$ and minima is $-\sqrt{5}$

□

This may seem excessive, and frankly overkill for the given question, but it becomes increasingly more powerful as we look at Olympiad questions. We basically partial differentiate both the functions and declare their ratio as λ . This along with the original equation leads to the maxima or minima, which solves the inequality.

Be very careful while using it as it will attract the wrath of the grader as it is an undergraduate method. However, if we are careful, it is also free marks. This is at the end to discourage you from only using Lagrange (like I did for quite a long time) and use the actual inequalities.

Everytime we have an inequality question, we can do it by the others as well. Just that Lagrange is easier and quicker most of the times. Here is an example from MOP, to put a close on inequalities

Example 16.13. (MOP 2012) Given that $a + b + c + d = 4$, for positive reals a, b, c, d prove that

$$\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + \frac{1}{d^2} \geq a^2 + b^2 + c^2 + d^2$$

Proof. While this can be done using Tangent line, Jensen and more, we'll use Lagrange.

We'll rewrite the inequality as: $\frac{1}{a^2} - a^2 + \frac{1}{b^2} - b^2 + \frac{1}{c^2} - c^2 + \frac{1}{d^2} - d^2 \geq 0$

$$\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \lambda \begin{bmatrix} \frac{-2}{a^3} - 2a \\ \frac{-2}{b^3} - 2b \\ \frac{-2}{c^3} - 2c \\ \frac{-2}{d^3} - 2d \end{bmatrix}$$

We can notice that $a = b = c = d = \frac{4}{4} = 1$ at the minima which is 0. This makes the inequality true

NOTE: If you are rigorous, you may notice that we don't know if 0 is the maxima or the minima. In this case we can test it by having two variables (and rest as 1s) and proving that it is always above 0. This while still not the correct way, is satisfactory (as this is almost similar to the single derivative test).

The more accurate methods of determining this will come in undergraduate studies. \square

16.6 Sum uses of Calc

Example 16.14 (Motivating Example). For a polynomial $p(x)$ with roots $r_1, r_2, r_3 \dots r_n$ find:

$$\sum_{i=1}^n \frac{1}{x - r_i}$$

Solution. While we can do this using vieta, here is the most chad way of going about it.

We know that $p(x) = a(x - r_1)(x - r_2) \dots (x - r_n)$

We can convert it to sum by taking $\log p(x) = \log a + \log x - r_1 + \dots + \log x - r_n$

We can convert it into reciprocals by diffrentiation:

$$\frac{p'(x)}{p(x)} = \frac{1}{x - r_1} + \frac{1}{x - r_2} + \dots + \frac{1}{x - r_n}$$

This solves the question. \square

As we can see, discrete summations can also be done using analytic techniques(calc). However, the fun part is almost all discrete sums are solvable by these methods.

However, we'll make a few assumptions about the question before solving them. While this may seem wrong, here is an assumption we have been making for quite some time:

$$1 - 1 + 1 - 1 + \dots$$

Some of you may compute this as $(1 - 1) + (1 - 1) + \dots = 0$ or you may compute this as $1 + (-1 + 1) + (-1 + 1) \dots = 1$ or in the correct way as an infinite GP $\frac{1}{1 - (-1)} = \frac{1}{2}$.

The other two are wrong as the given series is conditionally convergent (converges in only a few cases, and diverges in other, definition wise has the absolute sum of terms diverging) which makes grouping inherently wrong. The Riemann Series theorem (which you'll learn more about in real analysis, not in this book) proves that this is not allowed. However, we have been rearranging and grouping sums for quite a long time without running into any such issues. This is because we are assuming that the series is absolutely convergent (converges all the time). This allows grouping, order changes and all the other techniques.

We'll now with the knowledge about our ignorance, proceed to ignore it once more as questions are questions as the series is absolutely convergent.

You will study more about conditional convergence in real analysis and discrete analysis courses.

Definition 16.15. A power series (centered at 0) is an infinite series of the form

$$\sum_{n=0}^{\infty} a_n z^n = a_0 + a_1 z + a_2 z^2 + \dots$$

where $a_i \in \mathbb{C}$ for all $i \geq 0$.

While it is hard to see whether every such series converges, however here is a simple way to check:

Definition 16.16.

$$R = (\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}})^{-1}$$

Here \limsup refers to the maximum possible value of the function at the limit. Basically, for a function like \sin this refers to 1 at ∞ as otherwise it's an oscillating limit.

Here R is the radius of convergence. The series converges(normally) if




Figure 16.0. \limsup and \liminf in a graph

$z : |z| < R$ and diverges to ∞ or $-\infty$ if $z : |z| > R$.

We'll not use this again. It's useless in most cases. As I said before, we normally assume a series is absolutely convergent.

We should also remember the Taylor series expansions, as we are going to use them.

Example 16.17.

$$\sum_{n=0}^{\infty} \frac{1}{(2n)!!}$$

where $n!! = n(n - 2)(n - 4) \dots$ is the double factorial.

Solution. $\sum_{n=0}^{\infty} \frac{1}{(2n)!!}$

$$\sum_{n=0}^{\infty} \frac{1}{(2n)(2n-2)(2n-4)\dots(2)}$$

$$\sum_{n=0}^{\infty} \frac{1}{2^n(n)(n-1)(n-2)\dots(1)}$$

$$\sum_{n=0}^{\infty} \frac{1}{2^n n!}$$

$$\sum_{n=0}^{\infty} \frac{1}{2^n}$$

Which is the Taylor series for $e^{\frac{1}{2}} = \sqrt{e}$.

□

Here is also another fun fact:

Theorem 16.18 (Differentiating and integrating under the power series).
For 'well behaved' function, we can say if:

$$F(x) = \sum_{n=0}^{\infty} a_n x^n$$

then:

$$F'(x) = \sum_{n=1}^{\infty} (n-1)a_n x^{n-1}$$

and:

$$\int F(x) dx = \sum_{n=0}^{\infty} a_n \frac{x^{n+1}}{n+1}$$

Basically, as long as our functions are well behaved, we differentiate and integrate them term by term.

This can also be used to say, for a well behaved function:

$$\text{int} \sum f(x) = \sum \int f(x)$$

We will use these properties to solve the following example:

Example 16.19. (AMSP) Prove that

$$\sum_{r=1}^n \frac{1}{r} \binom{n}{r} = \sum_{r=1}^n \frac{2^r - 1}{r}$$

for $n \in \mathbb{Z}^+$

Proof. We will use the rewrite $\frac{1}{r} = \int_0^1 x^{r-1} dx$, using this on the LHS gives us:

$$\sum_{r=1}^n \binom{n}{r} \int_0^1 x^{r-1} dx \\ \int_0^1 \sum_{r=1}^n \binom{n}{r} x^{r-1} dx$$

This looks an awful lot like the binomial theorem with some modifications.

$$\int_0^1 \frac{(1+x)^n - 1}{x} dx$$

This can be solved by integration by substitution taking $u = x + 1$

$$\int_1^2 \frac{u^n - 1}{u-1} du$$

Which using the sum of GP turns to:

$$\int_1^2 \sum_{r=1}^n u^{r-1} dr$$

$$\sum_{r=1}^n \int_1^2 u^{r-1} dr$$

$$\sum_{r=1}^n \frac{2^r - 1}{r}$$

Which proves the above. □

Exercises

(1) Let a, b, c, d be positive real numbers with $a + b + c + d = 1$. Prove that:
 $6(a^3 + b^3 + c^3 + d^3) \geq a^2 + b^2 + c^2 + d^2 + \frac{1}{8}$

(2) (USAMO 1978) Given that a, b, c, d, e are real numbers such that $a + b + c + d + e = 8$
 $a^2 + b^2 + c^2 + d^2 + e^2 = 16$ determine the minimum value of e .

(3) (Canada 1997) Prove that for non negative real numbers a, b, c with $a + b + c = 1$ we have
 $a^2b + b^2c + c^2a \leq \frac{4}{27}$

(4) (USAMO 2011) If $a^2 + b^2 + c^2 + (a + b + c)^2 \leq 4$, then

$$\frac{1 + ab}{(a + b)^2} + \frac{1 + bc}{(b + c)^2} + \frac{1 + ac}{(a + c)^2} \geq 3$$

(5) Let a, b, c be positive reals satisfying $a + b + c = \sqrt[3]{a} + \sqrt[3]{b} + \sqrt[3]{c}$. Prove that $a^a b^b c^c \geq 1$

(6) Let a, b, c be the sides of a triangle. Prove that:

$$\frac{a}{b+c-a} + \frac{a}{b+c-a} + \frac{a}{b+c-a} \geq 3$$

- (7) Let a, b, c be real numbers such that $0 \leq a, b, c \leq 1$, then prove
$$\frac{a}{1+bc} + \frac{a}{1+bc} + \frac{c}{1+ab} \leq 2$$
- (8) (USAMO 2004) Let a , b , and c be positive real numbers. Prove that
$$(a^5 - a^2 + 3)(b^5 - b^2 + 3)(c^5 - c^2 + 3) \geq (a + b + c)^3.$$

Part 6

Number Theory

17

Wearing the Crown of Mathematics

We can simply, but reasonably correctly, claim that there are three fundamental building blocks of mathematics: numbers, geometric figures and logical reasoning. Just as geometric objects and mathematical statements can be thought of as systematic collections of points and words, so can integers be broken into their indivisible components known as prime numbers. We have already discussed the logical reasoning part in form of algebra and combinatorics, let's talk about numbers now.

Welcome to Number Theory considered by many as the “crown of mathematics”

Example 17.1. Prove that $1^n + 2^n + 3^n + \cdots + (n - 1)^n$ is divisible by n for all odd n .

A human calculator(or a machine one for that matter) can compute to maybe $n = 50$, but that is not even 1% of ∞ for which the proposition must hold.

In contrast, using number theory we can prove this claim to be true within seconds, using only two lines and almost no calculations. How? You'll understand by the end of the chapter. While this question is quite easy, I reckon that some of you know how to solve it, number theory is not always that easy.

Example 17.2. (IMO 1988, P6) Let a and b be positive integers such that $ab + 1$ divides $a^2 + b^2$. Show that $\frac{a^2+b^2}{ab+1}$ is the square of an integer.

Even though this may look overall like some sort of a “simple division exercise,” it is a lot worse. This is believed to be the hardest IMO question of all time. The problem was solved perfectly by only 12 students at the IMO, and very few partial scores were given: it was “all or nothing.” But this is not even a taste of the complexity and beauty of number theory. Any work about number theory would be incomplete if we don’t mention perhaps the greatest math question.

Theorem 17.3 (Fermat’s Last Theorem). $a^n + b^n \neq c^n$ for $n > 2$

This theorem may seem innocent enough. But its proof was a mystery for more than 350 years. The theorem was written in the margin of a book by Piree de Fermat in 1637. He claimed to have “a truly marvelous proof of this proposition which this margin is too narrow to contain”. He never wrote the proof down, and the answer to this was lost to time.

It was proven by Andrew Wiles in 1995. Anyone who even remotely knows about Wiles’s actual proof will tell you that it is impossible to describe it in a few easy-to-understand sentences(it is 100+ pages long and mathematically dense). In fact, it is well beyond the scope of this book or of any highschool or undergraduate facing textbook and becomes accessible only in an advanced graduate course on modular forms. Do I wish to write a book on it one day? YES. Will that day be today? NO.

The amazing thing is that most of this advance math wasn’t available to Fermat in 1637, and we have reason to believe that he really had a proof. Fermat scribbled a lot of mathematical claims in the margins, and all of them were eventually proven using 17th century math. This was the only exception. It was Fermat’s last theorem to be left unproven. While Wiles’s proof is an achievement in its own regard, we still don’t know how Fermat would have proven this.

Fermat’s mathematical knowledge was similar to that of Olympiad mathematics. Which raises the question: Could you be the high school mathematician to find the elementary proof which has evaded the community for centuries? Who knows? But before we are even capable of approaching such a problem, let’s actually learn the tools we would need.

17.1 Division

I expect you to be familiar with division in the form:
You must also know that:

Quotient → 015
 Divisor → 32 487
 Dividend → 48
 32
 167
 160
 Remainder → 7

Figure 17.0. The long division

Theorem 17.4. $a = qd + r$ such that $0 \leq r < |d|$
Where q, r are uniquely determined for a, b

Do we notice any concepts whose appearance seems striking or “unjustified”? The absolute value $|d|$ jumps out at us: why do we need it? Well, if we divide by a negative d , we still want to arrive at a non-negative remainder r , but the inequalities $0 \leq r < d < 0$ won’t make much sense! This is remedied by the absolute value $|d| > 0$. For example, dividing 13 by -5 yields $13 = (-2) \cdot (-5) + 3$: the remainder 3 satisfies $0 \leq 3 < |-5|$

Definition 17.5. If $a = qd$ for some non-zero integers a, d , and q , we say that d and q divide a , or that d and q are divisors or factors of a , and write $d|a$ and $q|a$. Further, we say that $d|0$ since $0 = d0$

For instance, $5|10$, but $5 \nmid 13$. Note that the bar in ” $d | a$ ” is vertical, not slanted; while $5/10$ stands for the usual fraction $\frac{1}{2}$, the expression ” 5 divides 10 ” means something different, so be careful! Also, another trivial observation: if $d|a$, it is not necessarily true that $d \leq a$; indeed, $7|(-14)$ but $7 > -14$. To avoid annoying minus signs, we conclude in general that $d|a$ implies $|d| \leq |a|$ (unless $a = 0$).

We also need to note that division is just repeated addition or subtraction(whatever approaches 0). For example for $a = 16, d = 6$, we can find the remainder by $16 \rightarrow 16 - 6 \rightarrow 10 - 6 \rightarrow 4$. However, for $a = -19, d = 5$, we

find the remainder by $-19 \rightarrow -19 + 5 \rightarrow -14 + 5 \rightarrow -9 + 5 \rightarrow -4 + 5 \rightarrow 1$. We stopped the moment the condition $0 \leq r < |d|$ became true. This is called the Euclidean Algorithm for division.

We'll use this algorithm to prove our theorem.

Proof. Proof of existence: Set r to be the smallest non-negative such difference, say, $r = a - qd$ for some q . Since we start with a positive a and decrease it by d each time, we will eventually plunge under 0: the difference right before this plunge is our r . Rewriting, we have $a = qd + r$ with $r \geq 0$. Thus, we have constructed our quotient q and remainder $r \geq 0$; we only need to verify that $r < d$.

To the contrary, suppose $r \geq d$. Say we had 12 apples to give to 5 people. After giving everyone one apple, we are left with 7. If we kept them as remainder, we are greedy. We give the people one more each and have 2 as remainder. In general, we can subtract another d from r and get an even smaller non-negative difference:

$$a - qd = r \rightarrow a - (q + 1)d = r - d \geq 0,$$

which contradicts our choice for $a - qd$ as the smallest such difference. We conclude $r < d$. Hence we have constructed the desired quotient q and remainder r . We can symmetrically prove this for a or d being negative.

Proof of uniqueness: Let's to the contrary assume that q and r are not unique for every a and d .

We suppose that in addition to one quotient q and remainder r , there is another non equal quotient q_1 and remainder r_1 : $a = qd + r = q_1d + r_1$

$$\iff r - r_1 = d(q - q_1)$$

$$\iff d|(r - r_1)$$

$$\iff |r - r_1| \geq d$$

Which is untrue for all $r \neq r_1$ by the fact that $0 \leq r < |d|$

Hence, by contradiction, $r = r_1$ and $q = q_1$.

□

Note that this is often how we prove existence and uniqueness of something.

While this may seem excessive proving an obvious theorem, this is foundation for a lot more complex proves and without certainty of its truth, everything built on its foundation is shaky at best.

We will also take this moment to define two new functions:

Definition 17.6 (Greatest Common Divisor(GCD)). If we take F_a to be the set of all factors of a and F_b to be the set of all factors of b $\gcd(a, b) = \max(F_a \cap F_b)$

Definition 17.7 (Least Common Multiple(LCM)). If we take M_a to be the set of all multiples of a and M_b to be the set of all multiples of b $\text{lcm}(a, b) = \min(F_a \cap F_b)$

We can basically find the GCD using the Euclidean algorithm as follows:
For two natural a, b such that $a > b$, to find

Theorem 17.8. $\gcd(a, b)$ we use the division algorithm repeatedly:

$$a = bq_1 + r_1$$

$$b = r_1q_2 + r_2$$

$$r_1 = r_2q_3 + r_3$$

\vdots

$$r_{n-2} = r_{n-1}q_n + r_n$$

$$r_{n-1} = r_nq_{n+1}.$$

Then we have $\gcd(a, b) = \gcd(b, r_1) = \gcd(r_1, r_2) = \dots = \gcd(r_{n-1}, r_n) = r_n$

This can be proven simply by using the definition of GCD and the Euclidean algorithm.

Another thing to notice is that $\text{lcm}(a, b) \cdot \gcd(a, b) = ab$. Its proof follows from $a = kx$ and $b = ky$ where x, y are coprime(or have $\gcd(x, y) = 1$)

We can also use these functions to prove a theorem which we'll use a lot later.

Theorem 17.9 (Bezout's Theorem). For $a, b \in \mathbb{N}$, there exist $x, y \in \mathbb{Z}$ such that $ax + by = \gcd(a, b)$

Proof. The proof is simply by running the Euclidean Algorithm backwards.

$$\gcd(a, b) = r_{n-2} - r_{n-1}q_n$$

$$= r_{n-2} - (r_{n-3} - r_{n-2}q_{n-1})q_n$$

$$= r_{n-2}(1 + q_nq_{n-1}) - r_{n-3}(q_n)$$

\vdots

$$= ax + by.$$

□

This result must not be that fun. But a rather surprising, and beautiful, result is:

Theorem 17.10. $\gcd(a^m - 1, a^n - 1) = a^{\gcd(m, n)} - 1$

Proof. We can prove this by simply using the algebraic fact that $a^k - 1 | a^m - 1$ if $k|m$, which follows from the sum of GP formula or the general difference

expansion (whichever you like).

The GCD will happen for the largest k such that $k|m, n \iff k = \gcd(m, n)$
Hence, $\gcd(a^m - 1, a^n - 1) = a^{\gcd(m, n)} - 1$.

□

17.2 Congruence Modulo

Before today when we divided two numbers, our interest used to lie in the quotient. But sometimes, remainders become much, much more useful. Once something becomes useful, we give it some respect. Hence, instead of using remainder as if it is leftover soup, we start calling it congruence modulo or modulo.

Definition 17.11. $r \equiv a \pmod{d}$ means that a leaves remainder m when divided by d .

This definition of modulo leads us to some of its fundamental properties. These are all trivial to see and I expect that you will be able to prove them quite easily.

Theorem 17.12. (1) **Reflexivity:** $a \equiv a \pmod{n}$

(2) **Symmetry:** $a \equiv b \pmod{n}$ if and only if $b \equiv a \pmod{n}$

(3) **Transitivity:** If $a \equiv b \pmod{n}$ and $b \equiv c \pmod{n}$, then $a \equiv c \pmod{n}$

These three define the nature of the modulo function, which allows us to state:

Theorem 17.13. (1) **Compatibility with Translation:** $a + k \equiv b + k \pmod{n}$ for any integer k

(2) **Compatibility with Scaling:** $ka \equiv kb \pmod{n}$ for any integer k

(3) We can also state it as: $ka \equiv kb \pmod{kn}$ for any integer k

(4) **Compatibility with Exponentiation:** $a^k \equiv b^k \pmod{n}$ for any non-negative integer k

(5) **Compatibility with Addition:** $a_1 + a_2 \equiv b_1 + b_2 \pmod{n}$

- (6) **Compatibility with Subtraction:** $a_1 - a_2 \equiv b_1 - b_2 \pmod{n}$
- (7) **Compatibility with Multiplication:** $a_1 \cdot a_2 \equiv b_1 \cdot b_2 \pmod{n}$
- (8) **Compatibility with Polynomial Evaluation:** $p(a) \equiv p(b) \pmod{n}$, for any polynomial $p(x)$ with integer coefficients

Compatibility with exponentiation is used in questions a lot. Here are two examples:

Example 17.14. (a) Find the remainders of 2^{1998} divided by 3 and 3^{1998} when divided by 2.

(b) Find the remainders of 5^{2007} divided by 7 and of 17^{1701} divided by 11.

Solution. For (a), we can easily notice that $2 \equiv -1 \pmod{3}$, therefore $2^{1998} \equiv -1^{1998} \pmod{3} = 1 \pmod{3}$

The second one is even worse as $3 \equiv 1 \pmod{2}$, which makes $3^{1998} \equiv 1^{1998} \pmod{2} = 1 \pmod{2}$

For (b), we need to notice that $5^3 = 125 \equiv -1 \pmod{7}$ which solves the question as $\frac{2007}{3} = 669$. This makes $5^{2007} = 125^{669} \equiv -1^{669} \pmod{7} \equiv -1 \pmod{7} \equiv 6 \pmod{7}$

The second one is more like it. Consider $\pmod{11}$ to be applicable throughout the solve and $=$ and \equiv be interchangeable:

$$\begin{aligned} 17^{1701} &= 6^{1701} \\ &= 6 * 36^{850} = 6 * 3^{850} = 6 * 9^{425} \\ &= 54 * 9^{424} = -1 * 81^{212} = -1 * 7^{212} = -1 * 49^{106} \\ &= -1 * 5^{106} = -1 * 25^{53} = -1 * 3^{53} \\ &= -3 * 3^{52} = -3 * 9^{26} = -3 * 81^{13} \\ &= -3 * 4^{13} = -12 * 4^{12} = -1 * 16^6 \\ &= -1 * 5^6 = -1 * 25^3 = -1 * 3^3 = 26 = 4 \end{aligned}$$

Hence the remainder is 4. □

The part 2 of (b) is the common type of modulo you can expect to occur often. The other were designed for elegant solutions.

We can also use the compatibility of exponentiation to get the following result: the units digit, tens digit or any place per say, of any power repeats in cycles, this can be proven by taking d as 10, then 100 and so on.

Here is an example for you to try.

Example 17.15. (AIME 2010) Find the remainder when $9 \times 99 \times 999 \times \dots \times \underbrace{999 \dots 9}_{999 \text{ 9's}}$ is divided by 1000.

As we close this section, with our new found knowledge, we are now capable enough to prove the first example.

Proof. $1^n + 2^n + \cdots + (n-1)^n$

We take \pmod{n} to get $1^n + 2^n + \cdots + (-2)^n + (-1)^n$

as n is odd, $1^n - 1^n + 2^n - 2^n + \cdots \equiv 0 \pmod{n}$, which means that $1^n + 2^n + \cdots + (n-1)^n$ is divisible by n .

□

We just proved something about numbers, with powers and sums, without computing a single one. That's quite impressive, isn't it?

We'll return to Congruence modulo in the next chapter.

17.3 Prime Numbers

Example 17.16. There are 100 light bulbs on the wall, all of them initially lit up. Following specific rules:

- (1) The first bulb overheats and breaks.
- (2) A math God decides that bulbs with even numbers (except for 2) will burst.
- (3) A physics God decides that bulbs divisible by 3 (except for 3) will explode.
- (4) A chemistry God decides that bulbs divisible by 5 (except for 5) will turn into glass shards.
- (5) A biology God decides that bulbs divisible by 7 (except for 7) will shatter.

After all these divine interventions, how many lights remain illuminated? (Note: Once a bulb is affected, it cannot be restored by any other divine power.)

You'll be sad to know that we don't have any way to solve this without actually writing the numbers down. As it turns out, after crossing the broken bulbs out, we get:

What we'll notice is that all the bulbs which remain have exactly two divisors or factors: 1 and the number itself. These numbers are called the prime numbers. The crossed out numbers are called the composite numbers

Theorem 17.17 (Fundamental Theorem of Arithmetic). *Every natural number $n \neq 1$ may be prime factorized in one and only one way as $p_1^{e_1} p_2^{e_2} \cdots p_k^{e_k}$ where all p are prime and all $e \in \mathbb{N}$.*




Figure 17.0. The bulbs which remain lit. The diagram is called the Sieve of Eratosthenes

This is another theorem which is quite simple and known to every schoolchild. The only problem, schools don't teach the proof, which is essential for deeper understanding. I have discussed the proof below.

Proof. We divide it into two parts. Showing the existence of a prime factorization and then proving it's uniqueness.

Proof of existence: (B) Prime factorization trivially exists for the first few numbers. Let's consider till $n = 4$ where 2, 3 are prime and $4 = 2^2$.

(S) Let's assume that all numbers less than k can be prime factorized. If k is prime, the number has a trivial prime factorization, $k = k^1$. If k is composite, which means $k = pq$ where p and q are less than k and have a prime factorization. We multiply the prime factorization to get the prime factorization of k .

Proof of uniqueness: Let's to the contrary assume that:

$$p_1^{e_1} p_2^{e_2} \dots p_k^{e_k} = q_1^{f_1} q_2^{f_2} \dots q_l^{f_l} \text{ where terms of both } p, q \text{ are distinct primes.}$$

We can say that as p_1 divides the LHS, it also divides the RHS. This means some q is divisible by p_1 , let that term be q_1 . However, as p_1 and q_1 are primes, $p_1 = q_1$. This leads to $p_i = q_i$ for $i \leq k = l$. This also leads to the exponents being equal, which makes the prime factorization exactly the same, contradicting the initial assumption.

Thus, the assumption was false. This means that every n has one and only one prime factorization of the form $p_1^{e_1} p_2^{e_2} \dots p_k^{e_k}$

□

This leads to the observation that in order to check whether a number n is prime, we need to check all the primes that are less than or equal to \sqrt{n} as any factor of n less than \sqrt{n} has its corresponding factor greater than \sqrt{n} and the same is true vice versa.

Example 17.18. How many prime numbers are there? Prove it!

Note that we have 25 primes less than 100. But only 21 from 100 to 200. It drops to 16 in the range 200 to 300. So a question arises, are their finite primes?

NO, and we'll prove that.

Proof. Let's, to the contrary, assume that we have finite primes. We can list them as p_1, p_2, \dots . However, $(p_1 * p_2 * \dots) + 1$ is not divisible by any of them. If it is a prime, our list is incomplete. If it is not, its prime factors have been left out from the list. In either case, we have a contradiction. Hence, our initial assumption was false. Thus, we have infinite primes. \square

This is one of the most beautiful proofs in math, considered to be the part of "**THE BOOK**".

We normally denote primes with p . While we can prove that there are infinite primes, it is an open question if there are infinite primes of the form $p + 2$ called Euclidian Primes, infinite primes of the form $2p + 1$ called Germain Primes or infinite primes of the form $2^p - 1$ called Mersenne Primes. While proving these is much harder, I do recommend thinking about it when you get time.

Another common question is if we have some formula to find the n^{th} prime. Unfortunately, no such formula exists which is accurate, the best one is called Prime Number Theorem. It uses the famous Riemann Zeta function and a lot of math beyond this book to get the following result:

Theorem 17.19 (Prime Number Theorem). *Number of primes less than $n \approx \frac{n}{\ln n}$ and consequently, the n^{th} prime $\approx n \ln n$*

They both are inaccurate for small n and become increasingly accurate as $n \rightarrow \infty$

NOTE: The prime number theorem is included just for your information, it will never be used in any competition, in any way whatsoever. So you can totally read and forget it.

17.4 Number Bases

To understand the notion of number bases, we look at our own number system. We use the decimal(dec), or base 10, number system. To help explain what this means, consider the number 2746. This number can be rewritten as $2746_{10} = 2 \cdot 10^3 + 7 \cdot 10^2 + 4 \cdot 10^1 + 6 \cdot 10^0$.

Note that each number in 2746 is actually just a placeholder which shows how many of a certain power of 10 there are. The first digit to the left of the decimal place (recall that the decimal place is to the right of the 6, i.e. 2746.0) tells us that there are six 10^0 's, the second digit tells us there are four 10^1 's, the third digit tells us there are seven 10^2 's, and the fourth digit tells us there are two 10^3 's.

Base-10 uses digits 0-9. Usually, the base, or radix, of a number is denoted as a subscript written at the right end of the number. We don't really do that with base 10 as it is the one we use all day. If we did, we would write the above example as 2746_{10} , where 10 is the radix.

The next natural question is: how do we convert a number from another base into base 10? For example, what does 4201_5 mean? Just like base 10, the first digit to the left of the decimal place tells us how many 5^0 's we have, the second tells us how many 5^1 's we have, and so forth. Therefore:

$$\begin{aligned} 4201_5 &= (4 \cdot 5^3 + 2 \cdot 5^2 + 0 \cdot 5^1 + 1 \cdot 5^0)_{10} \\ &= 4 \cdot 125 + 2 \cdot 25 + 1 = 551_{10} \end{aligned}$$

From here, we can generalize. Let $x = (a_n a_{n-1} \cdots a_1 a_0)_b$ be an $n+1$ -digit number in base b . In our example (2746_{10}) $a_3 = 2, a_2 = 7, a_1 = 4$ and $a_0 = 6$. We convert this to base 10 as follows:

$$\begin{aligned} x &= (a_n a_{n-1} \cdots a_1 a_0)_b \\ &= (b^n \cdot a_n + b^{n-1} \cdot a_{n-1} + \cdots + b \cdot a_1 + a_0)_{10} \end{aligned}$$

However, it turns out that converting from base 10 to other bases is far harder for us than converting from other bases to base 10. This shouldn't be a surprise, though. We work in base 10 all the time so we are naturally less comfortable with other bases. Nonetheless, it is important to understand how to convert from base 10 into other bases.

Example 17.20. Convert $1000_{10} = n_7$, Find the value of n .

Basically we are looking for a solution to
 $1000 = a_0 + 7a_1 + 49a_2 + 343a_3 + 2401a_4 + \cdots$

where all the a_i are digits from 0 to 6. Obviously, all the a_i from a_4 and up are 0 since otherwise they will add in a number greater than 1000, and all the terms in the sum are non negative. Then, we wish to find the largest a_3 such that $343a_3$ does not exceed 1000. Thus, $a_3 = 2$ since $2a_3 = 686$ and $3a_3 = 1029$. This leaves us with

$$1000 = a_0 + 7a_1 + 49a_2 + 343(2)$$

$$\iff 314 = a_0 + 7a_1 + 49a_2.$$

Using similar reasoning, we find that $a_2 = 6$, leaving us with $20 = a_0 + 7a_1$.

We use the same procedure twice more to get that $a_1 = 2$ and $a_0 = 6$.

Finally, we have that $1000_{10} = 2626_7$.

An alternative version is to find the "digits" a_0, a_1, \dots starting with a_0 . Note that a_0 is just the remainder of division of 1000 by 7. So, to find it, all we need to do is to carry out one division with remainder. We have $1000 : 7 = 142(R6)$. How do we find a_1 , now? It turns out that all we need to do is to find the remainder of the division of the quotient 142 by 7: $142 : 7 = 20(R2)$, so $a_1 = 2$. Now, $20 : 7 = 2(R6)$, so $a_3 = 6$. Finally, $2 : 7 = 0(R2)$, so $a_4 = 2$. We may continue to divide beyond this point, of course, but it is clear that we will just get $0 : 7 = 0(R0)$ during each step.

Note that both versions of this method use computations in base 10.

It's often a good idea to double check by converting your answer back into base 10, since this conversion is easier to do. We know that $2626_7 = 343 \cdot 2 + 6 \cdot 49 + 2 \cdot 7 + 6 = 1000$, so we can rest assured we got the right answer.

While it may seem strange to talk about bases despite using 10 for years without issues, we need to realize that alternate bases are all around us.

Computers use base 2 or binary(bin), which is extremely efficient as it can be represented as two states in a circuit, on and off.

This makes them very fast at running calculations using semi-conductors in the silicon chip. The exact how of this is under the perview of physics and chemistry, not math and is not discussed here.

Base 8 or Octal(oct) is used in music theory due to the fact that we have eight notes. However, this property is also used by transponders in airplanes to communicate messages to the ground.

Base 16 or Hexadecimal(hex) is used in a lot of coding language from C to HTML. The color codes (which look something like #321adf are hexadecimal where we are using 0 – 9 and a – f) are also hexadecimal.

Base 60 or Sexagesimal(sex) is the reason why we have 60 seconds in a minute. 60 minutes in an hour. 360 degrees in an angle. ≈ 360 days in an year, 180 latitudes and 360 longitudes. It is not 'Illuminati' only the ancient Babylonians who invented all of this and they used base 60 which we continue with. And these are only a few examples. Computer science uses a lot more bases in many ways unique to them. They have a modified ternary with the digits -1, 0, 1, and use that in many codes. However, the best use in my opinion is it allows us to have two christmas or Halloween in an year, whichever you like. How? By simply noticing $\text{oct}(31) = \text{dec}(25)$

Example 17.21. $\overline{abc}_7 = \overline{cba}_9$. What is a, b, c in base 10? (here \overline{abc} represents a three digit number with digits a, b, c in the given order)

Solution. We can come to a common base in order to solve the question. Let it be 10 for simplicity

$$\overline{abc}_7 = \overline{cba}_9$$

$$\iff 49a + 7b + c = 81c + 9b + a$$

$$\iff 48a = 80c + 2b$$

$$\iff 24a - 40c = b$$

Here, we need to note that a, b, c are 0 – 6 as they are digits of a base 7 number. We also need to note that $8|b \iff b = 0$.

$$\therefore 3a - 5c = 0$$

$\iff 3a = 5c$ using the bounds,

$$\therefore a = 5, b = 0, c = 3$$

□

Here is another one for you:

Example 17.22. $\overline{xyz}_9 = \overline{zyx}_6$. Find $x + y + z$.

17.5 Divisibility rules

Using modulo, we can create some divisibility rules to check if a number N is divisible by some natural number n .

Here are some common divisibility rules:

Table 17.0. Divisibility Rules

Divisor	Rule
2	Last digit is even
3	Sum of digits is divisible by 3
4	Last 2 digits divisible by 4
5	Last digit is 0 or 5
6	Divisible by 2 and 3
7	Take the last digit, double it, and subtract from the rest If the result is divisible by 7, then the number is divisible by 7
8	Last 3 digits are divisible by 8
9	Sum of digits is divisible by 9
10	Last digit is 0
11	Calculate the sum of odd digits (O) and even digits (E) If $ O - E $ is divisible by 11, then the number is divisible by 11
12	Divisible by 3 and 4
15	Divisible by 3 and 5

Theorem 17.23. *NOTE: The divisibility test for $p * q$ is the combined test of p and q if they are co prime(that is $\gcd p, q = 1$)*

Proof. The proof for $n = 2, 4, 5, 8, 10$ is kinda obvious. If we write the digit as $10^k a_k + 10^{k-1} a_{k-1} + \dots + 10^3 a_3 + 10^2 a_2 + 10 a_1 + a_0$ we can notice that, taking $(\text{mod } 2)$ we get $a_0 \pmod{2}$, which should be 0 if the number is divisible by 2. Hence, a_0 should be even then. I expect that you will be able to prove it for 4 and 8 using this technique.

The proof for $n = 3, 9, 11$ is similar. If the $N = 10^k a_k + 10^{k-1} a_{k-1} + \dots + 10^3 a_3 + 10^2 a_2 + 10 a_1 + a_0$, then on taking $(\text{mod } 9)$ we get $a_k + \dots + a_1 + a_0 \pmod{9}$, which should be 0 if the number is divisible by 9. This means $a_k + \dots + a_1 + a_0$ is divisible by 9. The proof for 3 is almost exactly the same. The one for 11 takes $10 \pmod{11} = -1$ and the proof follows.

The proof for $n = 6, 12, 15$ follows from the fact that if $6|N$, then $N = 6 * k = 2 * 3 * k$ which means, N is divisible by 2 and 3. This only works if n can be broken into two co-prime factors(if a, b have $\gcd a, b = 1$, they are co-prime). The proof for $n = 7$ is a bit more involved. We prove this in reverse, we assume $N = 10a + b$ and $7|a - 2b$ and prove that $7|N$

This can be done by converting the $7|$ to $(\text{mod } 7)$ and noticing that $a \equiv 2b \pmod{7}$ which limits $a \equiv 0, 2, 4, 6 \pmod{7}$ for $a \equiv 0, 1, 2, 3 \pmod{7}$ respectively.

This makes $N \equiv 10*0+0, 10*2+1, 10*4+2, 10*6+3 \pmod{7} \equiv 0, 21, 42, 63 \pmod{7} = 0 \pmod{7}$ proving the rule.

□

Here is a question which can be solved quite easily if you have understood how the rules were derived.

Example 17.24. Find and prove a divisibility rule in base 7 arithmetic that is analogous to the rule (in ordinary base 10 arithmetic) for divisibility by 9. See if you can find other divisibility rules in base 7 arithmetic that are similar to rules for base 10

17.6 Nature of Factors

Example 17.25. The number of factors of a number n if it can be written as: $p_1^{e_1} * p_2^{e_2} * \dots * p_n^{e_n}$ where $p_1, p_2 \dots p_n$ are prime and $e_1, e_2 \dots e_n \in \mathbb{Z}^+$ is:

All factors of n will have the same prime factors. Essentially, for every prime p_n of prime factorization of n has a power $0 \geq k_n \geq e_n$ in its factors. Hence, every k_n will have $e_n + 1$ possible values. Thus the total factors are: $(e_1 + 1) * (e_2 + 1) * \dots * (e_n + 1)$

Example 17.26. The sum of the divisors of some natural number is:

$\sigma_{d|n} d = (1+p_1+p_1^2+\cdots+p_1^{q_1})(1+p_2+p_2^2+\cdots+p_2^{q_2}) \cdots (1+p_k+p_k^2+\cdots+p_k^{q_k})$. We can justify this claim using the fact that there are $(q_1+1)(q_2+1)(q_3+1) \cdots (q_k+1)$ products formed by taking one number from each sum, which is the number of divisors of n . Clearly all possible products are divisors of n . Furthermore, all of those products are unique since each positive integer has a unique prime factorization. Since all of these products are added together, we can conclude this gives us the sum of the divisors.

Example 17.27. There are 100 light switches on the wall, all turned off. A hundred toddlers come by. The first toddler flips every switch. Then the second toddler flips just switches 2, 4, 6, 8, Then the third toddler flips switches 3, 6, 9, 12, This pattern continues until finally the 100th toddler flips just switch number 100. How many lights are turned on at the end?

What we need to realize is that for every factor of a number, we have another factor complementing it. This will cause all the bulbs to remain closed as the switch is flipped even number of times. However, this is not the case as the complement can be equal. If this occurs for some factor k , then the number is k^2 . This basically means that all squares have odd number of factors while non-squares have even number of factors.

This means only the bulbs 1, 4, 9, 16, 25, 36, 49, 64, 81, 100 will be turned on.

17.7 Legendre's Theorem

Example 17.28. How many zeros are at the end of $100!$?

Obviously, we are not expecting you to compute $100!$ and count the zeros. What we need to do is that every zero represents the number being divisible by some power of ten. Basically if a number has 2 zeros at the end, then it is divisible by $10^2 = 100$.

$10^n = 2^n 5^n$, and as power of 2 is anyhow more than power of 5 in the prime factorization of $100!$, it is simple that for every power of 5, we'll have one zero at the end.

How do we count those? Every number divisible by 5 will contribute to the exponent once. Every number divisible by 25 will contribute twice.

This leads to the number of zeros being $20 + 4 = 24$

We can write this in the form:

Theorem 17.29 (Legendre's Theorem).

$$v_p(n!) = \sum_{i=1}^{\infty} \left\lfloor \frac{n}{p^i} \right\rfloor = \frac{n - S_p(n)}{p - 1}$$

where p is a prime and $v_p(n!)$ is the exponent of p in the prime factorization of $n!$ and $S_p(n)$ is the sum of the digits of n when written in base p

17.8 Irrationality

Example 17.30. Prove that \sqrt{p} is irrational for any prime p .

Proof. Let's to the contrary assume that \sqrt{p} is rational and equal to $\frac{m}{n}$ in its lowest form

$$\begin{aligned} \therefore p^2 &= \frac{m^2}{n^2} \\ \iff p^2 n^2 &= m^2 \\ \therefore p^2 | m^2 &\iff p | m \iff m = pm' \\ \therefore p^2 &= \frac{p^2 m'^2}{n^2} \\ \iff n^2 &= m'^2 \iff n = m' \end{aligned}$$

This causes $\frac{m}{n} = \frac{pm'}{m'}$ which is a contradiction as m and n should be co-prime for the fraction to be in the lowest form.

Hence, the assumption is false and \sqrt{p} cannot be written as a rational number, or is irrational.

□

This proof is a common place in all good math textbooks. I decided to include at the end of the chapter just before the exercise starts.

Exercises

- (1) (AMC 10) A positive integer divisor of $12!$ is chosen at random. The probability that the divisor chosen is a perfect square can be expressed as $\frac{m}{n}$, where m and n are relatively prime positive integers. What is $m + n$?
- (2) (AMC 10) How many positive even multiples of 3 less than 2020 are perfect squares?
- (3) (AMC 10) How many positive integer divisors of 202^3 are perfect squares or perfect cubes (or both)?
- (4) (AMC 12) What is the sum of the exponents of the prime factors of the square root of the largest perfect square that divides $12!$?
- (5) (AIME 2012) Find the number of positive integers with three not necessarily distinct digits, abc , with $a \neq 0$ and $c \neq 0$ such that both abc and cba are multiples of 4.

- (6) (AMC 10) The 25 integers from -10 to 14 , inclusive, can be arranged to form a 5-by-5 square in which the sum of the numbers in each row, the sum of the numbers in each column, and the sum of the numbers along each of the main diagonals are all the same. What is the value of this common sum?
- (7) (AMC 12) How many odd positive 3-digit integers are divisible by 3 but do not contain the digit 3?
- (8) (AIME 2015) There is a prime number p such that $16p + 1$ is the cube of a positive integer. Find p .
- (9) (AMC 10) Joey and Chloe and their daughter Zoe all have the same birthday. Joey is 1 year older than Chloe, and Zoe is exactly 1 year old today. Today is the first of the 9 birthdays on which Chloe's age will be an integral multiple of Zoe's age. What will be the sum of the two digits of Joey's age the next time his age is a multiple of Zoe's age?
- (10) (AMC 10) Let $N = 34 \cdot 34 \cdot 63 \cdot 270$. What is the ratio of the sum of the odd divisors of N to the sum of the even divisors of N ?
- (11) (AMC 12) In multiplying two positive integers a and b , Ron reversed the digits of the two-digit number a . His erroneous product was 161. What is the correct value of the product of a and b ?
- (12) Positive integers a , b , and 2023, with $a < b < 2023$, form a geometric sequence with an integer ratio. What is a ?
- (13) (AMC 10) Suppose that m and n are positive integers such that $75m = n^3$. What is the minimum possible value of $m + n$?
- (14) (AIME 2005) Find the number of positive integers that are divisors of at least one of 10^{10} , 15^7 , 18^{11} .
- (15) (AMC 12) For each positive integer $n > 1$, let $P(n)$ denote the greatest prime factor of n . For what value of positive integer n is it true that $P(n) = \sqrt{n}$ and $P(n + 48) = \sqrt{n + 48}$?
- (16) (AIME 1995) Let $n = 2^{31}3^{19}$. How many positive integer divisors of n^2 are less than n but do not divide n ?
- (17) (AIME 1990) Let n be the smallest positive integer that is a multiple of 75 and has exactly 75 positive integral divisors, including 1 and itself. Find $\frac{n}{75}$.
- (18) (AMC 10) Let n denote the smallest positive integer that is divisible by both 4 and 9, and whose base-10 representation consists of only 4's and 9's, with at least one of each. What are the last four digits of n ?

- (19) (AIME 2006) Let N be the number of consecutive 0's at the right end of the decimal representation of the product $1!2!3!4!\cdots 99!100!$. Find the remainder when N is divided by 1000
- (20) (IMO 1959) Prove that the fraction $\frac{21n+4}{14n+3}$ is irreducible for every natural number n .
- (21) (AIME 1986) What is the largest positive integer n for which $n^3 + 100$ is divisible by $n + 10$?
- (22) How many consecutive numbers do you need to guarantee that their product is divisible by 30? by 120?
- (23) (AMC 12) What is the largest integer that is a divisor of $(n+1)(n+3)(n+5)(n+7)(n+9)$ for all positive even integers n ?
- (24) (AMC 12) The number $21! = 51,090,942,171,709,440,000$ has over 60,000 positive integer divisors. One of them is chosen at random. What is the probability that it is odd?
- (25) (AIME 1985) The numbers in the sequence 101, 104, 109, 116, ... are of the form $a_n = 100 + n^2$, where $n = 1, 2, 3, \dots$. For each n , let d_n be the greatest common divisor of a_n and a_{n+1} . Find the maximum value of d_n as n ranges through the positive integers.
- (26) (AMC 10) How many ordered pairs (a, b) of positive integers satisfy the equation $a \cdot b + 63 = 20 \cdot \text{lcm}(a, b) + 12 \cdot \gcd(a, b)$,
- $$a \cdot b + 63 = 20 \cdot \text{lcm}(a, b) + 12 \cdot \gcd(a, b),$$
- (27) (AMC 12) There are 10 horses, named Horse 1, Horse 2, ..., Horse 10. They get their names from how many minutes it takes them to run one lap around a circular race track: Horse k runs one lap in exactly k minutes. At time 0 all the horses are together at the starting point on the track. The horses start running in the same direction, and they keep running around the circular track at their constant speeds. The least time $S > 0$, in minutes, at which all 10 horses will again simultaneously be at the starting point is $S = 2520$. Let $T > 0$ be the least time, in minutes, such that at least 5 of the horses are again at the starting point. What is the sum of the digits of T ?
- (28) (AMC 12) Let n be the smallest positive integer such that n is divisible by 20, n^2 is a perfect cube, and n^3 is a perfect square. What is the number of digits of n ?

- (29) (AMC 12) How many positive integers n are there such that n is a multiple of 5, and the least common multiple of 5! and n equals 5 times the greatest common divisor of 10! and n ?
- (30) Find the smallest natural number n such that $n!$ is divisible by 990? What about 560?
- (31) Sabrina multiplied two 2-digit numbers together on the blackboard. Then she changed all numbers to letters (different digits changed to different letters, equal digits to equal letters. She got : $\overline{AB} \cdot \overline{CD} = \overline{EEFF}$. Prove that Sabrina made a mistake somewhere.
- (32) What is the last digit of 2023^{2023} ?
- (33) Find the remainder when $2222^{5555} + 5555^{2222}$ is divided by 7
- (34) Suppose that a, b and c are integers such that $a^2 + b^2 = c^2$. Prove that at least one of a, b and c is divisible by 3.
- (35) . A base-10 3 digit number n is selected at random. What is the probability that the base-9 and base-11 representation of n are both 3-digit numbers?
- (36) The first 2007 positive integers are written in base-3. How many of these numbers are base-3 palindromes?
- (37) $121212121212_3 =$ what number in base 9?
- (38) The increasing sequence $1, 3, 4, 9, 10, 12, 13, \dots$ consists of all positive integers which are powers of 3 or sum of distinct powers of 3. Find the 100^{th} term of this sequence.
- (39) Using a weighing balance, on which weights can be placed on both sides, what is the minimum number of weights one needs to be able to measure all the integral weights between 1 and 1000?
- (40) How many trailing zeros does the number $(2^{16})!$ have in base-2 representation?
- (41) n had 10 positive divisors. $2n$ has 15 positive divisors. $3n$ has 20 positive divisors. How many positive divisors does $4n$ have?
- (42) (IMO 2023, 1) Determine all composite integers $n > 1$ that satisfy the following property: if d_1, d_2, \dots, d_k are all the positive divisors of n with $1 = d_1 < d_2 < \dots < d_k = n$, then d_i divides $d_{i+1} + d_{i+2}$ for every $1 \leq i \leq k - 2$.

- (43) (Putnam 2000, 2) Prove that for the expression

$$\frac{\gcd(m, n)}{n} \binom{n}{m}$$

is an integer for $n \geq m \geq 1$.

- (44) Prove that $a + b + c + d$ is not prime given, $ab = cd$ and that $a, b, c, d \in \mathbb{Z}$

- (45) (India 2017) Let a, b, c, d be pairwise distinct positive integers such that:

$$\frac{a}{a+b} + \frac{b}{b+c} + \frac{c}{c+d} + \frac{d}{d+a}$$

is an integer. Prove that $a + b + c + d$ is not prime.

18

Modular Arithmetic

This chapter we will talk more about Congruence Modulo. It is used in number theory for variety of purposes, from telling time to literally securing passwords, through cryptography.

While our uses of them will be elementary but we'll form the basis of checking card numbers, pins and internet encryption. Also, they help in telling the time.

So with no waste of time, let's begin.

18.1 Modulo Inverse

We start with a strange claim, If $\gcd(a, p) = 1$ then $\{0, a, 2a, 3a, \dots, (p-1)a\} \pmod{p}$ is pairwise distinct.

We can simply prove it by assuming, to the contrary, that two elements ai and aj are equal modulus p .

$$\begin{aligned}\therefore ai &\equiv aj \pmod{p} \iff a(i - j) = 0 \pmod{p} \\ &\iff p|a \text{ or } p|(i - j)\end{aligned}$$

The first case is not possible as $\gcd a, p = 1$, $\therefore p|(i - j)$ which is false as $|i - j| < p$ as $i, j \in \{0, 1, 2, \dots, p-1\}$

Hence, the assumption is false and hence, no two elements must be equal.

This through pigeonhole principle(remember?), means that the sets $\{0, a, 2a, 3a, \dots, (p-1)a\} \equiv \{0, 1, 2, \dots, p-1\} \pmod{p}$

Theorem 18.1 (Equal Sets lemma). $\{0, a, 2a, 3a, \dots, (p-1)a\} \equiv \{0, 1, 2, \dots, p-1\} \pmod{p}$ for $\gcd(a, p) = 1$

We need to note that the sets are equal in the fact that they have the same elements. The position of the elements is clearly different.

Using the equal sets lemma, we can say for any integer $0 < b < p$, we can find an integer x such that $ax \equiv b \pmod{p}$. In particular, if $b = 1$, then $ax \equiv 1 \pmod{p}$. What this means is if $\gcd(a, p) = 1$, then there always exists a multiple of a which is 1 modulo p . This allows us to define:

Definition 18.2. We say that the inverse of a number a modulo m , refers to $b \leq p$ such that $ab \equiv 1 \pmod{m}$.

b is commonly denoted as $a^{-1} \pmod{m}$

All of the work above was done to reach this point as inverses are very useful because they finally enable us to divide under the modulo.

Theorem 18.3. If $b \not\equiv 0 \pmod{p}$, then

$$\frac{a}{b} \equiv a \cdot b^{-1} \pmod{p}$$

18.2 Fermat's Little Theorem

Theorem 18.4 (Fermat's Little Theorem). *If p is prime and does not divide a , then $a^p \equiv a \pmod{p}$, which can also be written as: $a^{p-1} \equiv 1 \pmod{p}$*

Fermat's little theorem is, as we already know, the more benevolent short form of FLT. The other one is Fermat's Last Theorem.

The proof for this FLT is much shorter and sweeter(the last theorem proof is about 130 pages and filled with very complex math).

Proof. The most straightforward way to prove this theorem is by applying the induction principle. We fix p as a prime number. (B) $1^p \equiv 1 \pmod{p}$, is obviously true.

(S) Suppose the statement $a^p \equiv a \pmod{p}$ is true. Then, by the binomial theorem,

$$(a+1)^p = a^p + \binom{p}{1}a^{p-1} + \binom{p}{2}a^{p-2} + \cdots + \binom{p}{p-1}a + 1.$$

Note that p divides into any binomial coefficient of the form $\binom{p}{k}$ for $1 \leq k \leq p-1$. This follows by the definition of the binomial coefficient as $\binom{p}{k} = \frac{p!}{k!(p-k)!}$; since p is prime, then p divides the numerator, but not the denominator.

Taken \pmod{p} , all of the middle terms disappear, and we end up with $(a+1)^p \equiv a^p + 1 \pmod{p}$. Since we also know that $a^p \equiv a \pmod{p}$, then $(a+1)^p \equiv a + 1 \pmod{p}$, as desired \square

18.3 Euler's Totient Theorem

45 years after death of Pierre De Fermat, influenced by his work, Leonhard Euler proved the following, which is also called Euler's generalization or the Fermat-Euler theorem. I however, have gone for the old name which is...

Theorem 18.5 (Euler Totient Function). *Given the general prime factorization of $n = p_1^{e_1} p_2^{e_2} \cdots p_m^{e_m}$, one can compute $\phi(n)$ using the formula*

$$\phi(n) = n \left(1 - \frac{1}{p_1}\right) \left(1 - \frac{1}{p_2}\right) \cdots \left(1 - \frac{1}{p_m}\right).$$

ϕn represents the number of integers in the range $\{1, 2, 3, \dots, n\}$ which are relatively prime to n

Theorem 18.6 (Euler's Totient Theorem). *If a is an integer and m is a positive integer relatively prime to a , then $a^{\phi(m)} \equiv 1 \pmod{m}$.*

Let's now prove both these things,

Proof. To derive the formula, let us first define the prime factorization of n as $n = \prod_{i=1}^m p_i^{e_i} = p_1^{e_1} p_2^{e_2} \cdots p_m^{e_m}$ where the p_i are distinct prime numbers. Now, we can use a PIE argument to count the number of numbers less than or equal to n that are relatively prime to it.

First, let's count the complement of what we want (i.e. all the numbers less than or equal to n that share a common factor with it). There are $\frac{n}{p_1}$ positive integers less than or equal to n that are divisible by p_1 . If we do the same for each p_i and add these up, we get

$$\frac{n}{p_1} + \frac{n}{p_2} + \cdots + \frac{n}{p_m} = \sum_{i=1}^m \frac{n}{p_i}.$$

But we are obviously over counting. We then subtract out those divisible by two of the p_i . There are $\sum_{1 \leq i_1 < i_2 \leq m} \frac{n}{p_{i_1} p_{i_2}}$ such numbers. We continue with this PIE argument to figure out that the number of elements in the complement of what we want is

$$\sum_{1 \leq i \leq m} \frac{n}{p_i} - \sum_{1 \leq i_1 < i_2 \leq m} \frac{n}{p_{i_1} p_{i_2}} + \cdots + (-1)^{m+1} \frac{n}{p_1 p_2 \cdots p_m}.$$

This sum represents the number of numbers less than n sharing a common factor with n , so

$$\phi(n) = n - \left(\sum_{1 \leq i \leq m} \frac{n}{p_i} - \sum_{1 \leq i_1 < i_2 \leq m} \frac{n}{p_{i_1} p_{i_2}} + \cdots + (-1)^{m+1} \frac{n}{p_1 p_2 \cdots p_m} \right)$$

$$\begin{aligned}\phi(n) &= n \left(1 - \sum_{1 \leq i \leq m} \frac{1}{p_i} + \sum_{1 \leq i_1 < i_2 \leq m} \frac{1}{p_{i_1} p_{i_2}} - \cdots + (-1)^m \frac{1}{p_1 p_2 \cdots p_m} \right) \\ \phi(n) &= n \left(1 - \frac{1}{p_1} \right) \left(1 - \frac{1}{p_2} \right) \cdots \left(1 - \frac{1}{p_m} \right).\end{aligned}$$

□

And next we shall prove Euler's Totient Theorem.

Proof. Consider the set of numbers $A = \{n_1, n_2, \dots, n_{\phi(m)}\} \pmod{m}$ such that the elements of the set are the numbers relatively prime to m . We can prove that this set is the same as the set $B = \{an_1, an_2, \dots, an_{\phi(m)}\} \pmod{m}$ where $\gcd(a, m) = 1$ as all elements of B are relatively prime to m and distinct, which by the pigeonhole principle means that B has the same elements as A . In other words, each element of B is congruent to one of A . This means that $n_1 n_2 \dots n_{\phi(m)} \equiv an_1 \cdot an_2 \dots an_{\phi(m)} \pmod{m} \implies a^{\phi(m)} \cdot (n_1 n_2 \dots n_{\phi(m)}) \equiv n_1 n_2 \dots n_{\phi(m)} \pmod{m} \implies a^{\phi(m)} \equiv 1 \pmod{m}$ as desired. Note that dividing by $n_1 n_2 \dots n_{\phi(m)}$ is allowed since it is relatively prime to m and therefore has an inverse. □

18.4 Wilson's Theorem

A few years later, French mathematician John Wilson read both Fermat and Euler's work and gave us:

Theorem 18.7 (Wilson's Theorem). *if integer $p > 1$, then $(p-1)! + 1$ is divisible by p if and only if p is prime. Essentially, $(p-1)! \equiv -1 \pmod{p}$ for prime p .*

Let's now prove this as well.

Proof. Let's define a polynomial with the roots $1, 2, 3, \dots, p-1$:

$$g(x) = (x-1)(x-2)\dots(x-(p-1))$$

Also we can define $h(x)$ such that:

$$h(x) = x^{p-1} - 1$$

$$\therefore h(x) \pmod{p} = \{1, 2, 3, \dots, p-1\}$$

This is a consequence of Fermat's Little Theorem.

Subtracting $g(x)$ from $h(x)$ will give us:

$$f(x) = x^{p-1} - 1 - (x-1)(x-2)\dots(x-(p-1))$$

On taking mod p, we will still have roots 1, 2, 3, ..., p-1. This means, all terms of $f(x)$ are divisible by p, Which means the constant term is also divisible by p

$$\therefore (p-1)! + 1 \equiv 0 \pmod{p}$$

$$\therefore (p-1)! \equiv -1 \pmod{p}$$

□

This triad makes up the fundamental theorems of Modulo.
We can use them independently as in:

Example 18.8. For how many integer values of i , $1 \leq i \leq 1000$, does there exist an integer j , $1 \leq j \leq 1000$, such that i is a divisor of $2^j - 1$

Example 18.9. How many prime numbers p are there such that $29^p + 1$ is a multiple of p

Example 18.10. (ARML 2002) Let $a \in \mathbb{N}$ such that

$$1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{22} + \frac{1}{23} = \frac{a}{23!}$$

Find $a \pmod{13}$

Before you read the given solution, I advise you to try to guess which theorem goes to which question.

Solution. The first one, we can notice that i can't be even as $2^j - 1$ is odd for every j . For an odd i , we can be certain that $2^{\phi(i)} - 1$ is divisible by i using the Euler Totient theorem. Therefore, we have 500 such i from 0 – 100. The second one follows from $29^p + 1 \equiv 29 + 1 \equiv 30 \pmod{p}$ using FLT. As p is a divisor of $29^p + 1$, therefore $p|30$. As p is prime, we can say $p = 2, 3, 5$ The third one can be solved by isolating $a = 23!(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{22} + \frac{1}{23}) = 23! + \frac{23!}{2} + \frac{23!}{3} + \dots + \frac{23!}{22} + \frac{23!}{23}$, at this stage we can notice that all are divisible by 13 other than $\frac{23!}{13}$. This reduces the question to $\frac{23!}{13} \equiv 12! * (13 * 14 * 15 * \dots * 23) \equiv 12! * 10! = 12 * \frac{11!}{11} \equiv 12 * \frac{1}{11} \equiv 12 * 11^{-1} \pmod{13}$
We just used Wilson twice. We'll now finally use inverse of 11 which is 6 as $66 \equiv 1 \pmod{13}$. This means,
 $a \pmod{13} \equiv 12 * 6 \equiv 72 \equiv 7$

□

We'll also see them used together in some while.

18.5 Chinese Remainder Theorem

Theorem 18.11 (Chinese Remainder Theorem). *If a positive number x satisfies the system of congruence's:*

$$x \equiv a_1 \pmod{n_1}$$

$$x \equiv a_2 \pmod{n_2}$$

$$\vdots$$

$$x \equiv a_k \pmod{n_k}$$

where all n_i are relatively prime, then $x \equiv A \pmod{N}$ where $N = n_1 \cdot n_2 \cdot n_3 \dots n_k$

After a bit of a dry stretch, we have a theorem which is so obvious that it entails no proof(You can easily prove it using induction if you are not convinced). However, It only tells you the upper bound of solution, you then will either need to make a Diophantine(we'll learn about them later) or do hit and trial to get the solution.

While this theorem is a bit weak right now, we'll see its true power in the constructions chapter.

The uses we see in this chapter looks like follows

Example 18.12. (AIME 2012) For a positive integer p , define the positive integer n to be p -safe if n differs in absolute value by more than 2 from all multiples of p . For example, the set of 10-safe numbers is $\{3, 4, 5, 6, 7, 13, 14, 15, 16, 17, 23, \dots\}$. Find the number of positive integers less than or equal to 10,000 which are simultaneously 7-safe, 11-safe, and 13-safe

Solution. We can notice that our number must be $3, 4 \pmod{7}, 3, 4, 5, 6, 7, 8, 9 \pmod{11}, 3, 4, 5, 6, 7, 8, 9, 10, 11 \pmod{13}$. Which means that the number is defined for $(\pmod{1001})$. We also need to notice that we'll have an unique solution for unique remainder we have in any of the cases.

This leads to 96 possible values per 1001 integers. Hence we have 960 such numbers in $1 - 10,010$. We just check for $10001 - 10010$ to get that even 100006 and 100007 fulfill the condition. Therefore, we have $960 - 2 = 958$ such numbers. \square

Example 18.13. Consider a number line consisting of all positive integers greater than 7. A hole punch traverses the number line, starting from 7 and working its way up. It checks each positive integer n and punches it if and only if $\binom{n}{7}$ is divisible by 12. As the hole punch checks more and more numbers, the fraction of checked numbers that are punched approaches a limiting number R . Find R

Solution. Using CRT in reverse, $0 \pmod{12}$ can be broken down to $0 \pmod{3}$ and $0 \pmod{4}$. As

$$\binom{n}{7} = \frac{n(n-1)(n-2)(n-3)(n-4)(n-5)(n-6)}{2^4 3^2 5 7}$$

The mod 3 condition forces us to look for 7 consecutive numbers divisible by 27 due to the 3^2 in the denominator. This can obviously only occur if and only if one of the numbers is divisible by 9. This means $n \equiv 0, 1, 2, 3, 4, 5, 6 \pmod{9}$

The mod 4 condition forces us to look for 7 consecutive numbers divisible by 2^6 due to the 2^4 in the denominator. If n is even, this is trivially true.

If n is odd, we have to either have $(n-1), (n-3), (n-5)$ divisible by 2. We also either have $4|n-3$ or $4|(n-1), (n-5)$

This means that in the first case, $16|n-3$ or in the second case $8|(n-1), (n-5)$, As everything in this condition perfectly divides 16, we can convert all the condition for 16 and say $n \equiv 0, 1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 13, 14 \pmod{16}$

Here we again use CRT to say that of every $9 * 16 = 144$ values of n , only $7 * 13 = 91$ satisfy the condition.

Thus, $R = \frac{91}{144}$ □

And just as a small taste of construction till we reach the actual chapter:

Example 18.14. Call a lattice point “visible” if the greatest common divisor of its coordinates is 1. Prove that there exists a 100×100 square on the board none of whose points are visible. Generalize it for $n \times n$ square.

Proof. Without loss of generality, we can let one of the corners be (a, b) As we are only proving the existence of something, we don’t really care if a simpler case exists. We only care if a case exists. Here we can let $a \equiv 0 \pmod{p_1 * p_2 * \dots * p_{100}}$ where p_i is prime.

This means $b \equiv 0 \pmod{p_1}, b+1 \equiv 0 \pmod{p_2}, \dots, b+99 \equiv 0 \pmod{p_{100}}$

We can now declare $a+1 = 0 \pmod{p_{101} * \dots * p_{200}}$

This means $b \equiv 0 \pmod{p_{101}}, b+1 \equiv 0 \pmod{p_{102}}, \dots, b+99 \equiv 0 \pmod{p_{200}}$

So on and so forth. We know that every term $b, b+1, \dots, b+99$ exists using CRT as p_i are prime.

Hence, such a square defiantly exists.

The generalization is left to you to solve. □

The square we made may or may not be the only one, that is a whole differently and much more difficult question. However, it defiantly exists. Such proves are called constructions, and we’ll see more of them, and more techniques to solve them, later.

Exercises

- (1) (AMC 12) Let S be a subset of $\{1, 2, 3, \dots, 30\}$ with the property that no pair of distinct elements in S has a sum divisible by 5. What is the largest possible size of S ?
- (2) (AMC 12) Let $k = 2008^2 + 2^{2008}$. What is the units digit of $k^2 + 2^k$?
- (3) (AIME) Let $a_n = 6^n + 8^n$. Determine the remainder upon dividing a_{83} by 49.
- (4) (AMC 10) What is the hundreds digit of 2011^{2011} ?
- (5) (AIME) The positive integers N and N^2 both end in the same sequence of four digits $abcd$ when written in base 10, where digit a is not zero. Find the three-digit number abc .
- (6) (AMC 10) An integer N is selected at random in the range $1 \leq N \leq 2020$. What is the probability that the remainder when N^{16} is divided by 5 is 1?
- (7) (PUMAC) If p, q, r are primes with $pqr = 7(p+q+r)$, find $p+q+r$
- (8) What are the last two digits of the integer 17^{198} ?
- (9) (AMC 10) Let $a_1, a_2, \dots, a_{2018}$ be a strictly increasing sequence of positive integers such that

$$a_1 + a_2 + \dots + a_{2018} = 2018^{2018}.$$

What is the remainder when $a_1^3 + a_2^3 + \dots + a_{2018}^3$ is divided by 6?

- (10) How many of the first 2018 numbers in the sequence 101, 1001, 10001, 100001, ... are divisible by 101?
- (11) (AMC 10) What is the remainder when $3^0 + 3^1 + 3^2 + \dots + 3^{2009}$ is divided by 8?
- (12) (AMC 10) What is the greatest power of 2 that is a factor of $10^{1002} - 4^{501}$?
- (13) (AMC 10) Let n be a 5-digit number, and let q and r be the quotient and the remainder, respectively, when n is divided by 100. For how many values of n is $q+r$ divisible by 11?
- (14) (AMC 10) Let $N = 123456789101112\dots4344$ be the 79-digit number that is formed by writing the integers from 1 to 44 in order, one after the other. What is the remainder when N is divided by 45?
- (15) (AMC 10) A palindrome between 1000 and 10,000 is chosen at random. What is the probability that it is divisible by 7?

- (16) (Purple Comet 2013) There is a pile of eggs. Joan counted the eggs, but her count was off by 1 in the ones place. Tom counted the eggs, but his count was off by 1 in the tens place. Raoul counted the eggs, but his count was off by 1 in the hundreds place. Sasha, Jose, Peter, and Morris all counted the eggs and got the correct count. When these seven people added their counts together, the sum was 3162. How many eggs were in the pile?
- (17) (Mandelbrot 2008) Determine the smallest positive integer m such that $m^2 + 7m + 89$ is a multiple of 77.
- (18) Find some positive multiple of 21 has 241 as its final three digits.
- (19) Show that if n is an integer greater than 1, then n does not divide $2^n - 1$.
- (20) . Let n be a positive integer such that $n + 1$ is divisible by 24. Prove that the sum of the divisors of n is divisible by 24
- (21) Do there exist 2004 consecutive integers such that each is divisible by a perfect cube bigger than 1?
- (22) . Find all primes of the form $n^4 + 4$.
- (23) Prove that $3^n - 2^n$ is not divisible by n .

19

Functional Equations

Functional equations are equations which are defined on functions rather than in variables. These require surprising amount of intuition and guessing to solve (and then also sometimes come out wrong). However, this should not deter us from learning about them.

We will start of small and finally explore what are called 'monsters'. However, you may need to read some parts more than once to understand what is going on. So I request you to not move on until you are clear about a piece of text.

19.1 Some definitions

Let $f : A \rightarrow B$ be a function. The set A is called the **domain**, and B the **co-domain**. A couple definitions which will be useful:

Definition 19.1. A function $f : A \rightarrow B$ is **injective** if $f(x) = f(y) \iff x = y$. (Sometimes also called **one-to-one**.)

Definition 19.2. A function $: A \rightarrow B$ is **surjective** if for all $b \in B$, there is some $x \in A$ such that $f(x) = b$. (Sometimes also called **onto**.)

Definition 19.3. A function is **bijective** if it is both injective and surjective.

In this chapter, by solve over K , we will mean find all functions $f : K \rightarrow K$ such that the equation holds for all inputs in K .

For example:

There's a function from living humans to $\mathbb{Z}_{\geq 0}$ by taking every human to their

age in years (rounded to the nearest integer). This function is not injective, because for example there are many people with age 20. This function is also not surjective: no one has age 10000. There's also a function taking every Indian citizen to their Aadhar card number, which we view as a function from citizens to $\mathbb{Z}_{\geq 0}$. This is also not surjective (no one has card number equal to 00...3), but it is injective (no two people have the same number).

This example should help remembering the functions.

19.2 Basic Functional Equations

Example 19.4 (Motivation Example). (USAMO 2016) Find all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that for all real numbers x and y ,

$$(f(x) + xy) \cdot f(x - 3y) + (f(y) + xy) \cdot f(3x - y) = (f(x + y))^2.$$

One of the most tricky motivating example of this book. But this one question will explore most of the possible configurations of this chapter.

We can get some insight by first trying out random standard values. Like, let $y = 0$, will give us:

$$\begin{aligned} f(x) * f(x) + f(0) * f(3x) &= f(x)^2 \\ f(0) * f(3x) &= 0 \end{aligned}$$

This means either $f(3x) = 0 \implies f(x) = 0$ or $f(0) = 0$. As we have got one solution, we'll now try to see where $f(0) = 0$ takes us. Let $x = 0$

$$\begin{aligned} f(0) * f(-3y) + f(y) * f(-y) &= f(y)^2 \text{ as } f(0) = 0 \\ \iff f(y)f(-y) &= f(y)^2 \text{ which clearly shouts } f(y) = f(-y) \end{aligned}$$

At this stage my intuition blurs that $f(x) = x^2$ which works for the given equation.

However, we can show that that is the case by simply taking $x - 3y = 3x - y \iff x = -y$ to cancel out the unsymmetrical terms in the question.

$$f(2x)(f(x) - x^2 + f(-x) - x^2) = f(0)^2$$

We use $f(0) = 0$ and $f(x) = f(-x)$

$$(f(x) - x^2)f(4x) = 0 \iff f(x) = x^2 \text{ or } f(4x) = 0$$

This solves the question. However, we need to also prove that no other function satisfies this equation.

We do this by contradiction,

Assume to the contrary that $f(x)$ is such that $f(a) = 0$ for $a \neq 0$ and $f(b) = b^2$.

We'll let $x - 3y = a$ and $3x - y = b$

$$(f(x) + xy)a^2 = f(x + y)^2$$

As b is arbitrarily large as $f(b) = f(2b) = f(4b) = \dots = 0$ as we have seen above, we can claim $x, y > 0$ and therefore $f(x) + xy \geq xy > 0$ and hence, $f(x + y)^2 > 0$ which means $f(x + y) = (x + y)^2$

$$\begin{aligned}
 (x+y)^4 &= (f(x) + xy)(x - 3y)^2 \\
 \iff (x+y)^4 &\leq (x^2 + xy)(x - 3y)^2 \\
 \iff (x+y)^4 &< (x+y)^2(x - 3y)^2 \\
 \iff (x+y)^4 &< (x+y)^4
 \end{aligned}$$

Which is a contradiction and hence no such function exists.

Therefore the only such functions are $f(x) = 0$ or $f(x) = x^2$

The last step of proving is called the pointwise trap and not doing it is the reason why despite being correct a lot of people lose marks in functional equations.

While we used a few tricks here, we'll learn a few more now.

19.3 Some methods of solving

19.3.1 Forced cancellations.

We'll explore this using the given example:

Example 19.5 (Motivating Example). Find all $f : \mathbb{R} \rightarrow \mathbb{R}$ such that:

$$f(x^2 + y) = f(x^{27} + 2y) + f(x^4)$$

for $x, y \in \mathbb{R}$

Solution. A little thinking can convince us that $f(x) = 0$, however how do we go about proving it?

Let's set y such that: $x^2 + y = x^{27} + 2y$

$$\therefore y = x^2 - x^{27}$$

Which will mean, $f(x^4) = 0$, hence $f(x) = 0$ for all positive reals.

Now let's extend it to reals. Let $y = 0$,

$$f(x^2) = f(x^{27}) + f(x^4)$$

$$\iff 0 = f(x^{27}) + 0$$

$\iff f(x^{27}) = 0$ making it 0 for negative reals as well. □

19.3.2 The FFF trick!

Normally, $f(f(x))$ does more bad than good. But sometimes, we can see this as an opportunity and put another f . Let's see it in action:

Example 19.6. Find all strictly increasing functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that $f(f(x)) = x + 2$ for all integers x

Solution. Let's add a f on both the sides.

$$\begin{aligned}
 f(f(f(x))) &= f(x + 2) \\
 \iff f(x) + 2 &= f(x + 2) \\
 \iff f(x) &= x + k
 \end{aligned}$$

We put this in the original equation to solve for k

$$\begin{aligned}
 f(f(x)) &= x + 2 \\
 \iff f(x+k) &= x+2 \\
 \iff x+2k &= x+2 \\
 \iff k &= 1
 \end{aligned}$$

Thus, $f(x) = x + 1$

□

19.3.3 Symmetry. If the equation, or some parts of it are symmetric in terms of x and y , we can interchange them for insight.

Example 19.7. Find all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$xf(x) + y^2 + f(xy) = f(x+y)^2 - f(x)f(y)$$

for all real numbers x and y

Proof. Let's swap for x and y ,

$$yf(y) + x^2 + f(xy) = f(x+y)^2 - f(x)f(y)$$

What we need to notice is that the RHS is exactly the same. So subtracting the original equation from the new one:

$$xf(x) - x^2 = yf(y) - y^2$$

It is rather obvious that the solution is $f(x) = x$

□

Here I would also like to include a exam focused note: that functional equations over real numbers tend to have simpler solutions like $c, x, kx, kx + c, x^n, kx^n, kx^n + c$ and one can literally try all of them out to solve a complex one and just prove that it is the solution. Moreover, $n > 3$ is so rare that it is an abnormality in the space time continuum. Also in the worst case scenario, we may have a polynomial, so that one thing.

Is it correct or the ethical way to solve the problem? NO. Will Cauchy come in your dream and haunt you? Probably. But will it score marks? Defiantly. While, this trick is not applicable to functional equations over integers or whole numbers as you have divisibility, and other things, most of which work because integers are "discrete". However, strange functions, like ones that depend on mods or something are rare enough that you can just use the base forms here as well and still get the answer most of the time.

Example 19.8. (USAMO 2019) Let \mathbb{N} be the set of positive integers. A function $f : \mathbb{N} \rightarrow \mathbb{N}$ satisfies the equation

$$\underbrace{f(f(\dots f(n) \dots))}_{f(n) \text{ times}} = \frac{n^2}{f(f(n))}$$

for all positive integers n . Given this information, determine all possible values of $f(1000)$.

Proof. Assume $f(a) = f(b)$. Then

$$f^{f(a)}(a) = f^{f(b)}(b)$$

since f is applied the same number of times. Thus we get

$$a^2 = f^{f(a)}(a) = f^{f(b)}(b) = b^2$$

so $a = b$. Thus, $f(a) = f(b) \iff a = b$, so f is injective.

Let $n = 1$. Then we get $f^{f(1)}(1) = 1$ and $f^{f(1)} = 1$. Now let $f(1) = k$. We get

$$k^2 = f^{f(k)}(k) = f(f(k)) = f(k)f(1) = k$$

so $k = 1$, therefore $f(1) = 1$.

Now we use **Induction**

Claim: $f(n) = n$ for all odd numbers n . (B) We have that $f(1) = 1$, so our base case is done.

(S) Now assume it is true for $n = 1, 3, \dots, 2k - 1$, and let $n = 2k + 1$. If $f(f(n)) < n$, then $f(f(n)) = m$ for some odd number $m < n$. But then by injectivity, $f(n) = m$, so $n = m$, which is a contradiction. Similarly, if $f^{f(n)}(n) = m < n$, then $n = m$, which is another contradiction. Thus, we must have $f(f(n)) = f^{f(n)}(n) = n$. So our induction is complete.

From injectivity, we know $f(1000)$ can't be odd. Notice that in the above proof, n being odd was important, and if it was even, we couldn't conclude anything. This makes it seem like $f(1000)$ could be any even number, and we only need to find one such function. Now consider the function

$$f(n) = \begin{cases} 2k & \text{if } n = 1000 \\ 1000 & \text{if } n = 2k \\ n & \text{if } n \neq 2k, 1000 \end{cases}$$

This function was chosen since we want $f(f(n)) = n$, which would make the function f satisfy the equation. Thus, the even numbers are all possible values of $f(1000)$. \square

This is a complicated question, so I recommend reading the proof once more. It had taken me way longer than it should have to understand the proof.

19.4 Cauchy's Functional Equations

Cauchy's functional equations refers to certain set of functional equations, all which are plays on $f(x) + f(y) = f(x + y)$. Using them can simplify a lot of questions. I recommend memorizing them so as to not needing to derive them whenever they come up/

Theorem 19.9 (Cauchy's first functional equation). *For $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that:*

$$f(x) + f(y) = f(x + y)$$

where $f(x)$ is continuous if and only if $f(x) = kx$

The proof of the same is trivial via induction. I expect that you'll be able to do this.

Theorem 19.10 (Cauchy's second functional equation). *For $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that:*

$$f(x \cdot y) = f(x) + f(y)$$

where $f(x)$ is continuous if and only if $f(x) = k \ln x$

Proof. We know that every $x = a^u$ for a unique $u \in \mathbb{R}$ for every x, a . This allows us to make a substitution:

$$f(a^u \cdot a^v) = f(a^u) + f(a^v)$$

This transforms into the first Cauchy functional equation by defining $f(a^x) = g(x)$. This solves to give us $f(x) = k \ln x$ \square

Theorem 19.11 (Cauchy's third functional equation). *For $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that:*

$$f(x + y) = f(x) \cdot f(y)$$

where $f(x)$ is continuous and non-zero if and only if $f(x) = a^x$ where $a = f(1)$

Proof. We can take log on both sides.

$$\log f(x + y) = \log f(x) + \log f(y)$$

Taking $\log f(x) = g(x)$, this transforms into the first Cauchy. We solve to get $f(x) = a^x$ where $a = f(1)$ \square

Theorem 19.12 (Cauchy's fourth functional equation). *For $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that:*

$$f(x \cdot y) = f(x) \cdot f(y)$$

where $f(x)$ is continuous and non zero if and only if $f(x) = x^k$

Proof. This time taking log transforms to the second Cauchy. Which leads to $f(x) = x^k$ \square

19.5 Checklist for Functional Equation Solving

At the beginning of a problem:

- Figure out what the answer is using the common solutions list.
- Plug in $x = y = 0, x = 0$ into the equation
- Plug in things that make lots of terms cancel.
- Look to see if the FFF trick can be used.
- Look to check if it is injective or surjective and if that can tell us something
- Look for symmetry (or breaks in that)
- Look for opportunities to use induction
- Look for ways to simplify it to Cauchy
- Make sure you don't fall in the pointwise trap
- Check if the solution actually works

Let's start the exercises now.

Exercises

- (1) (USAJMO 2015) Find all functions $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that

$$f(x) + f(t) = f(y) + f(z)$$

for all rational numbers $x < y < z < t$ that form an arithmetic progression.
(\mathbb{Q} is the set of all rational numbers.)

- (2) (USAMO 2002) Let \mathbb{R} be the set of real numbers. Determine all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x^2 - y^2) = xf(x) - yf(y)$$

for all pairs of real numbers x and y .

- (3) Find all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfying:

$$f\left(\frac{x^2}{2} + y\right) = f(f(x) - y) + 4f(x)y$$

- (4) (IMO 2010) Find all function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that for all $x, y \in \mathbb{R}$ the following equality holds

$$f(\lfloor x \rfloor y) = f(x) \lfloor f(y) \rfloor$$

- (5) (AHSME 1998) Let $f(x)$ be a function with the two properties:

- (a) for any two real numbers x and y , $f(x + y) = x + f(y)$, and
(b) $f(0) = 2$.

What is the value of $f(1998)$?

- (6) (AIME 1994) The function f has the property that, for each real number x , $f(x) + f(x-1) = x^2$. If $f(19) = 94$, what is the remainder when $f(94)$ is divided by 1000?

- (7) (BMO 1997) A non-negative integer $f(n)$ is assigned to each positive integer n in such a way that the following conditions are satisfied:

- (a) $f(mn) = f(m) + f(n)$, for all positive integers m , and n ;
- (b) $f(n) = 0$ whenever the units digit of n (in base 10) is a 3; and
- (c) $f(10) = 0$.

Prove that $f(n) = 0$, for all positive integers N

- (8) (Putnam 1999, A1) Find polynomials $f(x)$, $g(x)$, and $h(x)$, if they exist, such that, for all x :

$$|f(x)| - |g(x)| + h(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0 \end{cases}$$

- (9) (Russia 1988) The functions $f(x)$ and $g(x)$ are defined on the real axis so that they satisfy the following condition: for any real numbers x and y , $f(x + g(y)) = 2x + y + 5$. Find an explicit expression for the function $g(x + f(y))$.

- (10) (IMO 1977) Let $f(n)$ be a function $f : \mathbb{N}^+ \rightarrow \mathbb{N}^+$. Prove that if

$$f(n+1) > f(f(n))$$

for each positive integer n , then $f(n) = n$.

- (11) (Russia 1991) Does there exist a function $F : \mathbb{N} \rightarrow \mathbb{N}$ such that for any natural number x , $F(F(F(\dots F(x)\dots))) = x+1$? Here F is applied $F(x)$ times

- (12) (Putnam 1992, A1 modified) Find all $f : \mathbb{N} \rightarrow \mathbb{N}$ such that:

- (a) $f(f(n)) = n$, for all integers n ;
- (b) $f(f(n+2) + 2) = n$ for all integers n ;
- (c) $f(0) = 1$.

- (13) (Putnam 1971 B2) Let $F(x)$ be a real valued function defined for all real x except for $x = 0$ and $x = 1$ and satisfying the functional equation $F(x) + F(1 - \frac{1}{x}) = 1+x$. Find all functions $F(x)$ satisfying these conditions.

- (14) (a) (IMO 1987, 4) Prove that there is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ which satisfies the functional equation $f(f(n)) = n + 1987$.

- (b) Is there an $f : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $f(f(n)) = n^2$?

- (c) Is there a $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $g(g(x)) = -x$? Is there a continuous g ?

- (15) (IMO 1988, 3) A function f defined on the positive integers (and taking

$$f(1) = 1,$$

$$f(3) = 3$$

positive integers values) is given by:

$$f(2n) = f(n)$$

for

$$f(4n + 1) = 2f(2n + 1) - f(n)$$

$$f(4n + 3) = 3f(2n + 1) - 2f(n),$$

all positive integers n . Determine with proof the number of positive integers ≤ 1988 for which $f(n) = n$.

20

Diophantine Equations

Equations in which we are looking for only integer solutions are called Diophantine equations.

While this book deals with the most elementary cases of Diophantine, remember that some of the most difficult unsolved problems are Diophantine equations. There have been many advanced techniques developed in modern number theory to solve such equations, for instance elliptic curves. These are normally part of **graduate courses**, which I am not and neither claim to be an expert on.

Unlike most of the chapters, We'll explicitly look at it from an Olympiad perspective to make sure that we don't go too ahead of ourselves. Also, Diophantine equations is best learnt through problems.

20.1 Linear Diophantine Equations

Example 20.1 (Motivating Example). Find all integer solutions to $5x + 6y = 12$

This question is quite simple. We can obviously see that one solution is $(0, 2)$, we can now find rest of the solutions by adding 6 to the 0 and subtracting 5 the same number of times from 2. We can also subtract 6 and add 5. This gives us that all solutions are of the form $6k, 2 - 5k$ where $k \in \mathbb{Z}$. We can check this is true by plugging this in the equation $5(6k) + 6(2 - 5k) = 30k + 12 - 30k = 12$.

Let's now generalize,

Theorem 20.2 (Linear Diophantine). *Given $ax + by = c$ over integers, if $\gcd(a, b)|c$, and if one solution is (x', y') then all the solutions are in the form $(x' - bk, y' + ak)$ where $k \in \mathbb{Z}$*

Proof. The GCD condition comes from Bezout's identity(remember?). The rest of it can simply be shown to be true using substitution.

$$a(x' - bk) + b(y' + ak) = ax' + by' - abk + abk = ax' + by' = c$$

This is the entirety of solutions as a linear equation always has only one solution, so the minute we set y, x is defined by default. We are basically setting one of the variables and the other just follows. \square

This theorem, while use full still requires us to provide the initial solution. While in simpler equations like the one above, it is easy to do so, what about something more complex like:

Example 20.3. $125x + 8y = 279$

While this one is still somewhat doable with hit and trial, the fastest way out is taking modulus 8

$$5x = 7 \pmod{8}$$

We now need to only check 7 one digit values for x , where in this case it is obviously 3. This gives us a solution $(3, 12)$. And that generalizes to $(3 - 8k, 12 + 125k)$.

This method is especially use full while working with Diophantine equations arising from CRT(like this one which was a CRT on 1000).

20.2 Parity Arguments

Sometimes we can negate the need to solve a Diophantine by showing that no solution actually exists. Mathematicians don't like to lose, so when they can't do something they just prove it's impossible to do it. Some may call that stubbornness or pride. Mathematicians may call it "certainty".

One way of doing so is through parity arguments.

Example 20.4 (Motivating Example). Can one form a "magic square" out of the first 36 prime numbers? A "magic square" here means a 6×6 array of boxes, with a number in each box, and such that the sum of the numbers along any row, column, or diagonal is constant.

Notice that the question doesn't ask us to find the square. Just to prove if it exists or not.

We notice that The only even prime is 2 and it cannot be part of only three lines(row, column or diagonal).

This means that the sum of those lines is odd as they have 5 odd numbers and 1 even number. However, the sum of the other lines is even as they have 6 odd numbers.

As an odd number can't be equal to an even number, no such magic square exists.

Here is another such question for you to try.

Example 20.5. Let k be an even number. Is it possible to write 1 as the sum of the reciprocals of k odd integers?

20.3 Algebraic methods

We can use algebraic tricks to break Diophantine equations by either factorization or using inequalities. This is mainly the reason why we study number theory after algebra.

Example 20.6 (Motivating Example). (AIME 2000) A point whose coordinates are both integers is called a lattice point. How many lattice points lie on the hyperbola $x^2 - y^2 = 2000^2$?

Solution. We first need to notice that both x and y are even. We can do this by either taking $\pmod{4}$ or by using parity. Thus, we can let $x = 2m, y = 2n$. Now we factorize, $x^2 - y^2 = 2000^2 \iff 4m^2 - 4n^2 = 2000^2 \iff m^2 - n^2 = 1000^2 \iff (m - n)(m + n) = 2^6 5^6$

It is obvious that for different ways of splitting the factors will lead to different solutions. We will double this number as we also need to consider negative values of m, n . Thus, there are $7 * 7 * 2 = 98$ lattice points.

□

Example 20.7. (BMO 2005) The integer n is positive. There are exactly 2005 ordered pairs (x, y) of positive integers satisfying:

$$\frac{1}{x} + \frac{1}{y} = \frac{1}{n}$$

Prove that n is a perfect square.

Proof. We can convert the equation to $nx + ny = xy$ which looks an awful lot like SFFT.

$$xy - nx - ny + n^2 = n^2$$

$$(x - n)(y - n) = n^2$$

This has 2005 solutions. This means n^2 has 2005 factors. If the prime factorization of $n = p_1^{e_1} p_2^{e_2} \dots$, then:

$$(2e_1 + 1)(2e_2 + 1) \cdots = 2005 = 5 * 401$$

Therefore, without loss of generality $e_1 = 2$ and $e_2 = 200$. This makes n have $(2 + 1)(200 + 1) = 3 * 201 = 603$ factors. As 603 is odd, we know that n is square is an odd numbers. \square

Finally here is an INMO question for you to solve.

Example 20.8. (INMO ??) Determine all non negative integral pairs (x, y) for which $(xy - 7)^2 = x^2 + y^2$

Now let's try some questions which will use inequalities to solve the Diophantine equation.

Example 20.9. (Russia) Find all (x, y) such that $x, y \in \mathbb{N}$ $x^3 - y^3 = xy + 61$.

Solution. We need to notice that $x^3 - y^3 = (x - y)(x^2 + xy + y^2) = xy + 61$ means that $x - y > 0 \iff x > y$.

This means $(x^2 + xy + y^2) \leq xy + 61 \iff x^2 + y^2 \leq 61$.

This inequality can be solved by plugging in values of x and then checking for y . This gives us the following possible solutions $(x, y) = (7, 3)(7, 2)(7, 1)(6, 5)(6, 4)(6, 3)(6, 2)$. We can plug in all the cases to notice that only solution is $(6, 5)$

\square

Example 20.10. Find all pairs (x, y) of integers such that $x^3 + y^3 = (x + y)^2$ and $x \neq |y|$

Solution. This is slightly more involved than the last one.

$$(x + y)(x^2 - xy + y^2) - (x + y)^2 = 0$$

$$\iff (x + y)(x^2 + y^2 - xy - x - y) = 0$$

As $x \neq |y|$, $x + y \neq 0$

$$\therefore x^2 + y^2 - xy - x - y = 0$$

$$\iff x^2 + y^2 - 2xy = x + y - xy$$

$$\iff 1 - (x - y)^2 = xy - x - y + 1$$

$$\iff 1 - (x - y)^2 = (x - 1)(y - 1)$$

$$\iff 1 = (x - 1)(y - 1) + (x - y)^2$$

$$\iff 1 \geq (x - 1)(y - 1)$$

This limits our values of $(x, y) = (0, 0)(1, n)(n, 1)(2, 2)$ where $n \in \mathbb{N}$. As, $(0, 0)$ and $(2, 2)$ are rejected, We'll resolve for n now.

$$1 = (1 - 1)(n - 1) + (n - 1)^2$$

$$\iff \pm 1 = n - 1$$

$$\iff n = 0, 2$$

This gives us the four solutions $(1, 2)(2, 1)(0, 1)(1, 0)$. \square

We can also use other inequalities in this process. Here is one which will use AM-HM or SEBACS.

Example 20.11. (Putnam 2005, B2) Find all series k_1, k_2, \dots, k_n such that:
 $k_1 + \dots + k_n = 5n - 4$ and $\frac{1}{k_1} + \dots + \frac{1}{k_n} = 1$

20.4 Modular Contradiction Method

As I discussed before, mathematicians really hate equations they can't solve. Here is another way to flat out with certainty declare that an equation is not solvable.

Example 20.12 (Motivating Example). (RMO 2017) Show that the equation

$$a^3 + (a+1)^3 + \dots + (a+6)^3 = b^4 + (b+1)^4$$

has no solution in integers a, b

This question makes it rather obvious that taking $\pmod{7}$ is a good idea by literally giving us 7 consecutive digits.

We make a table:

This makes it clear that the LHS is $\equiv 0 \pmod{7}$ while the RHS can only be

a	$a^3 \pmod{7}$	$a^4 \pmod{7}$
0	0	0
1	1	1
2	1	2
3	-1	4
4	1	4
5	-1	2
6	-1	1

$1, 3, 6 \pmod{7}$ and hence, by contradiction, the given equation has no solutions.

While making such a table is not much hassle, most questions don't make it obvious which modulus to take. So here is a small list of modulo to consider:

Theorem 20.13. (1) $a^2 \equiv 0, 1 \pmod{3}$

(2) $a^2 \equiv 0, 1 \pmod{4}$

- (3) $a^2 \equiv 0, \pm 1 \pmod{5}$
- (4) $(\text{Odd integer})^2 \equiv 1 \pmod{8}$
- (5) $a^3 \equiv 0, \pm 1 \pmod{7}$
- (6) $a^3 \equiv 0, \pm 1 \pmod{9}$

We can also use FLT here in the following form:

Theorem 20.14. $a^{\frac{p-1}{2}} \equiv 0, \pm 1 \pmod{p}$ for some prime p . Basically for an exponent, if its double plus 1 is prime, we should consider that as a choice the base of our modulo.

We will see another example before we end this section:

Example 20.15. (USAJMO 2013) Are there integers a and b such that a^5b+3 and ab^5+3 are both perfect cubes of integers?

Proof. Either one of a, b is divisible by 3 or not. If a is, we'll have $a^5b+3 \equiv 6 \pmod{9}$ which is not possible for a cube. If b is then, $ab^5+3 \equiv 6 \pmod{9}$ and we run into the same issue.

If they are not, then let's to the contrary assume, $a^5b+3 \equiv \pm 1 \pmod{9} \iff a^5b \equiv 5, 7 \pmod{9}$ and similarly $ab^5+3 \equiv \pm 1 \pmod{9} \iff ab^5 \equiv 5, 7 \pmod{9}$ but as $a^5b \cdot ab^5 = (ab)^6 \equiv 4, 7, 8 \pmod{9}$ which is a contradiction as $x^6 \equiv 0, 1 \pmod{9}$. \square

20.5 Pythagorean Triplets

A Pythagorean triplet is refers to a possible sets of sides of a right angle triangle using the Pythagoras theorem. A primitive Pythagorean triplet is defined as follows.

Definition 20.16. The solution to

$$a^2 + b^2 = c^2$$

where $a, b, c \in \mathbb{N}$ and $\gcd(a, b, c) = 1$ is called a primitive Pythagorean triplet.

We define all primitive Pythagorean triplets using algebra as follows:

Theorem 20.17 (Triplet Formula). *For a primitive Pythagorean triplet where $a, b < c$ and $a, b, c, m, n \in \mathbb{N}$:*

$$\begin{aligned} a &= 2mn \\ b &= m^2 - n^2 \\ c &= m^2 + n^2 \end{aligned}$$

Here is a very simple use of this:

Example 20.18. Prove that for any three primes a, b, c , $a^2 + b^2 \neq c^2$

Proof. While the actual proof is not that hard (parity), the Pythagorean triplets make it an embarrassment.

Without loss of generality, let $a = 2mn$, which if $a = 2$ if a is prime which means $mn = 1 \iff m = 1$ and $n = 1$

Thus, $b = c = 0$ which is not prime.

Hence, $a^2 + b^2 \neq c^2$ for a, b, c being primes. □

Here is much better example.

Example 20.19. (Korea 1993) An integer which is the area of a right-angled triangle with integer sides is called Pythagorean. Prove that for every positive integer $k > 12$ there exists a Pythagorean number p such that $k < p < 2k$.

|

Proof. By the triplet formula, we can say that the area is $\frac{2mn(m^2-n^2)}{2} = mn(m^2-n^2) = mn(m-n)(m+n)$

This is especially great as we can let $n = 1$. We can now notice that for $m = 3$ we have the Pythagorean number 24 which is satisfying for k from 13 – 23. We can take $m = 4$ which is satisfying for k from 31 – 59 using the Pythagorean number 60. $m = 5$ satisfies 61 – 119 and so on. Note 60, 120... are yet to be satisfied.

However, the gap remains for 24 – 29. We'll look at 30 in a minute. Here we can take $n = 2$ and then see that for $m = 3$, we'll have it satisfied for 23 – 29. The only un-satisfied values are at the edges like 30, 60, 120.... We will solve all of them by a simple maneuver. If we scale a primitive Pythagorean triplet by n (increase every term n times), the area increases by n^2 . We have a Pythagorean number 6 where $m = 2$ and $n = 1$. We can now simply scale it by first 3 to get 54 for 30. Then by 4 for 60 and so on.

We can algebraically prove every step, but that's left as some work for you to do. □

Remember Fermat's Last Theorem? While we can't understand the proof of the same, we can still use it.

Theorem 20.20 (Fermat's Last Theorem). $a^n + b^n \neq c^n$ for $n > 2$

Here is an embarrassment of a question.

Example 20.21. (India) If x, y, z and $n > 1$ are all natural numbers with $x^n + y^n = z^n$. Prove that $x, y, z > n$

The question is so bad that whoever wrote it should lose marks for making it.

Anyways, using Fermat's last theorem we know that $n = 2$. After which, this question is burned using simple Pythagoras.

A slightly better use would be:

Example 20.22. Prove that $\sqrt[n]{2}$ is irrational for natural numbers $n > 1$.

Proof. We had proven it for $n = 2$ previously and will not do it here.

For $n > 2$, Let's to the contrary assume that $\sqrt[n]{2} = \frac{p}{q}$ where $p, q \in \mathbb{Z}$.

$$\begin{aligned}\therefore 2 &= \frac{p^n}{q^n} \\ \iff q^n + q^n &= p^n\end{aligned}$$

Which is untrue using Fermat's last theorem. Hence, contradiction. Thus, $\sqrt[n]{2}$ is irrational for natural numbers $n > 1$. \square

Another use, which is not straightaway bad is:

Example 20.23. (Romania) Prove that if n is odd, a, b, c are non-zero integers and $a^{3n} + b^{3n} + 3(abc)^n = c^{3n}$, then $a = b = -c$

Proof. We can rewrite the equation as: $(a^n)^3 + (b^n)^3 + ((-c)^n)^3 + 3a^n b^n (-c)^n = 0$

We can now use $x^3 + y^3 + z^3 - 3xyz = \frac{1}{2}(x+y+z)((x-y)^2 + (y-z)^2 + (z-x)^2)$,
 $\therefore \frac{1}{2}(a^n + b^n - c^n)((a^n - b^n)^2 + (b^n + c^n)^2 + (a^n + c^n)^2) = 0$

$$\therefore a^n + b^n - c^n = 0 \iff a^n + b^n = c^n \text{ or } (a^n - b^n)^2 + (b^n + c^n)^2 + (a^n + c^n)^2 = 0$$

We reject the first one using Fermat's last theorem. The second one is destroyed using the fact $x^2 + y^2 + z^2 = 0 \iff x = y = z = 0$

$$a^n = b^2; b^n = -c^n; c^n = -a^n$$

$$\therefore a = b = -c \quad \square$$

20.6 Infinite Descent

Example 20.24 (Motivating Example). Find all the solutions to $x^2 + y^2 = 3z^2$

Proof. Trying to put things seems like we have no solution other than $(0, 0, 0)$. We can try to solve it by taking modulus 3.

$$\{0, 1\} + \{0, 1\} \equiv 0 \pmod{3}$$

Hence, $x, y \equiv 0 \pmod{3}$. This also forces $z \equiv 0 \pmod{3}$. Let $x = 3a, y = 3b, z = 3c$

$$\begin{aligned} \therefore (3a)^2 + (3b)^2 &= 3(3c)^2 \\ \iff 9a^2 + 9b^2 &= 27c^2 \\ \iff a^2 + b^2 &= 3c^2 \end{aligned}$$

This is exactly the equation we have. So for any answer (x, y, z) then $(\frac{x}{3^n}, \frac{y}{3^n}, \frac{z}{3^n})$ are solutions for $n \in \mathbb{Z}$.

Let for some solution (k, l, m) be the solution which $k+l+m$ is minimized. But $\frac{k}{3} + \frac{l}{3} + \frac{m}{3}$ is smaller. This means that the only possible value is $k+l+m=0$. This forces the solution $x = 0, y = 0, z = 0$. \square

This is called proof by infinite descent, as the solution keeps on descending to infinity. This method was another contribution of Fermat. He claimed, in a letter to Carcavi, that he had a proof using infinite descent to 10 propositions. Only 1 of them has been found yet. All others can be proven using modern techniques, but it really forces us to ask what did Fermat know that we don't?

The most interesting thing is that Fermat claimed that Fermat's Last Theorem could be solved using infinite descent. We are yet to find out how.

Example 20.25. Prove that no non-zero solution exists for $x^4 + y^4 = z^4$ without using Fermat's last theorem.

Proof. This is obviously the $n = 4$ case of Fermat's last theorem. We want to prove it.

We will rewrite the equation as $(x^2)^2 + (y^2)^2 = z'^2$ here $z' = z^2$

Using Pythagorean triplets, $x^2 = 2pq, y^2 = p^2 - q^2; z' = p^2 + q^2$

Using Pythagorean triplets, $q = 2ab; y = a^2 - b^2; p = a^2 + b^2$

This means $x^2 = 2pq = 4ab(a^2 + b^2)$

This means that if $p|a$ or $p|b$ then $p \nmid (a^2 + b^2)$. This means $ab, (a^2 + b^2)$ are co-prime. Which due to it all being equal to x^2 means, ab and $a^2 + b^2$ are square numbers.

But as a, b are co-prime as they are Pythagorean triplets, therefore a, b are both square numbers.

Let $a=A^2, b=B^2$ and $a^2 + b^2 = P^2$

Hence, $P^2 = (A^2)^2 + (B^2)^2 \iff P^2 = A^4 + B^4$

And we are done by infinite descent. \square

This may seem inconsequential but we just proved Fermat's last theorem for all n in the form $n = 4k$ with $k \in \mathbb{N}$. If there exists a simple proof to Fermat, it probably uses infinite descent.

Here is an example for you to solve before we move ahead,

Example 20.26. Solve over integers $x^4 + y^4 + z^4 + t^4 = 2020xyzt$

20.7 Vieta Jumping

As we close this chapter, we'll talk about the legendary IMO 1998, problem 6.

Example 20.27. Let a and b be positive integers such that $ab + 1$ divides $a^2 + b^2$. Show that $\frac{a^2+b^2}{ab+1}$ is the square of an integer.

We will finally solve it.

Proof. Lets assume to the contrary that $\frac{a^2+b^2}{ab+1} = k$ where $\sqrt{k} \notin \mathbb{Z}$. Also without loss of generality, assume that $a \geq b$

$$\therefore a^2 + b^2 = kab + k \iff a^2 - kba + (b^2 - k) = 0$$

This is a quadratic with a root a . Let another root be x . We need to note a few things about x . Using Vieta, $x = kb - a$ which is an positive integer. Also $ax = b^2 - k$ which, as k is not a square means that $x \neq 0$. We will also show that $x \neq a$. As if that was true, $k^2b^2 = 4(b^2 - k)$ and $a = \frac{kb}{2}$ which would mean

$$\frac{\frac{k^2b^2}{4} + b^2}{\frac{kb^2}{2} + 1} = k$$

$$\iff \frac{k^2b^2}{4} + b^2 = \frac{k^2b^2}{2} + k$$

$$\iff b^2 - k = \frac{k^2b^2}{2}$$

Which contradicts the first point. Which means $k = b^2$ which contradicts the fact that k is not a square.

With this, lets without loss of generality assume that $a < x$

As $a \geq b$

$$\iff a^2 \geq b^2$$

$$\iff a^2 > b^2 - k$$

$$\iff a > \frac{b^2 - k}{a}$$

$$\iff a > x$$

Which is a contradiction. This contradiction remains even is $a > x$ as we can

do the same with x as x satisfies the original equation. If $b \geq a$ then we can do the whole process for b . This leaves only one assumption, that $\sqrt{k} \notin \mathbb{Z}$ which is false. Which means k is perfect square.

Hence, proved. □

This method is called Vieta jumping. We basically converted the equation to a quadratic and then used Vieta to jump from one root to the next. This method is best understood by more exploration so here is an example for you before you head into the exercises.

Example 20.28. Let a and b be positive integers such that ab divides $a^2 + b^2 + 1$. Show that:

$$\frac{a^2 + b^2 + 1}{ab} = 3$$

Exercises

- (1) (AMC 12 2005) Let A, M , and C be digits with

$$(100A + 10M + C)(A + M + C) = 2005.$$

What is A ?

- (2) (Purple Comet MS 2011) Find the prime number p such that $71p + 1$ is a perfect square
- (3) (PUMAC 2013) If p, q , and r are primes with $pqr = 7(p + q + r)$, find $p + q + r$
- (4) (AMC 10 2008) How many right triangles have integer leg lengths a and b and a hypotenuse of length $b + 1$, where $b < 100$?
- (5) (Purple Comet HS 2004) Find n such that $n - 76$ and $n + 76$ are both cubes of positive integers.
- (6) (AIME 2013) Positive integers a and b satisfy the condition

$$\log_2(\log_{2^a}(\log_{2^b}(2^{1000}))) = 0.$$

Find the sum of all possible values of $a + b$.

- (7) . (AwesomeMath Test A) Find all pairs of integers (x, y) that satisfy the equation

$$2(x^2 + y^2) + x + y = 5xy.$$

- (8) Find all pairs of integers (x, y) that satisfy the equation

$$x^2 - y! = 2001.$$

- (9) (Hong Kong TST 2002) Prove that if a, b, c, d are integers such that

$$(3a + 5b)(7b + 11c)(13c + 17d)(19d + 23a) = 2001^{2001}$$

then a is even.

- (10) (APMO 2017) We call a 5-tuple of integers arrangeable if its elements can be labeled a, b, c, d, e in some order so that $a - b + c - d + e = 29$. Determine all 2017-tuples of integers $n_1, n_2, \dots, n_{2017}$ such that if we place them in a circle in clockwise order, then any 5-tuple of numbers in consecutive positions on the circle is arrangeable.

- (11) (IMO 2003, P2) Find all $(a, b) \in \mathbb{N}^2$, such that $\frac{a^2}{2ab^2 - b^3 + 1}$ is a positive integer.

- (12) (IMO 2007, P5) Let a and b be positive integers. Show that if $4ab - 1$ divides $(4a^2 - 1)^2$, then $a = b$.

- (13) Find all positive integers m and n for which $1! + 2! + 3! + \dots + n! = m^2$

- (14) (INMO 1988) Find all $(x, y, n) \in \mathbb{N}$ such that $\gcd(x, n+1) = 1$ and $x^n + 1 = y^{n+1}$

- (15) Find the number of triples of (x, y, z) of positive integers satisfying

$$(x + y + z)^2 = 2018xyz$$

Part 7

The Number's Awaken

21

Bazooka!

This chapter talks about some more interesting and complicated concepts of Number Theory. We start with simpler concepts and then move towards more complicated concepts. The chapter will start with extensions of older concepts like Modulo and primes. We'll then explore the nature of functions related to number theory like Euler Totient and then define some more functions. We'll extend on the topic of Diophantine by talking about Pell's equations ($x^2 - dy^2 = 1$ where $d \in \mathbb{Z}^+$). We will then finally talk about some black boxes which like Fermat's Last theorem are very difficult to prove but can be cited to solve problems.

We'll also get to meet some more friendly theorems which couldn't be included elsewhere.

21.1 Surprisingly, not complex!

Not long ago we had learnt that $\sqrt{-1} = i$ where i is imaginary. What if I tell you that we can take some modulo and get not only a real but an integer value for i ? Even better what if I claim, $i \equiv 2, 3 \pmod{5}$?

This all seems rather strange. The upper claim is true as $i^2 = -1 \equiv 3^2 = 9 \equiv 2^2 = 4 \pmod{5}$. This however, doesn't occur with all numbers. The next one where we can seem something like this is 13, and then 17 and then 29. What is the pattern?

All the numbers are primes, but so are many which don't satisfy the given condition like 7, 11, 23. What about the fact that they are $1 \pmod{4}$. That seems to separate all of them. But is it true? If yes, how do we go about

proving it?

Theorem 21.1 (Fermat Christmas Theorem). *There exists an x with $x^2 \equiv -1 \pmod{p}$ if and only if p is a prime and $p \equiv 1 \pmod{4}$*

Proof. Whenever there is an frankly unbelievable result, Fermat is standing by.

Let's prove this in two parts, first that $x^2 \equiv -1 \pmod{p} \implies p \equiv 1 \pmod{4}$ for primes p and second that $x^2 \equiv -1 \pmod{p} \iff p \equiv 1 \pmod{4}$ for primes p .

For the first part, $x^2 \equiv -1 \pmod{p}$

$$\iff (x^2)^{\frac{p-1}{2}} \equiv -1^{\frac{p-1}{2}} \pmod{p}$$

$$\iff x^{p-1} \equiv -1^{\frac{p-1}{2}} \pmod{p}$$

$$\iff 1 \equiv -1^{\frac{p-1}{2}} \pmod{p}$$

$$\implies \frac{p-1}{2} \text{ is even or } p-1 = 4k \iff p \equiv 1 \pmod{4}.$$

The x^{p-1} was from Fermat's little theorem.

We'll now prove the second part. We will prove the existence using basic construction.

$$x = \left(\frac{p-1}{2}\right)!$$

$$\therefore x^2 = \left(\frac{p-1}{2} \cdot \frac{p-3}{2} \cdots 1\right) \cdot \left(\frac{p-1}{2} \cdot \frac{p-3}{2} \cdots 1\right)$$

$$\equiv \left(\frac{p-1}{2} \cdot \frac{p-3}{2} \cdots 1\right) \cdot \left(-\frac{p+1}{2} \cdot -\frac{p+3}{2} \cdots -(p-1)\right) \pmod{p}$$

$$\equiv -1^{\frac{p-1}{2}} ((p-1) \cdot (p-2) \cdots 2 \cdot 1) \pmod{p}$$

$$\equiv -1^{\frac{p-1}{2}} (p-1)! \pmod{p}$$

Which will be equal to one as $p = 4k + 1$ and $(p-1)! = 1 \pmod{p}$ from Wilson's theorem.

Thus, $x^2 \equiv -1 \pmod{p} \iff p \equiv 1 \pmod{4}$ for primes p . Hence, proved. □

Surly we'll use it for a question now? NO. The most surprising fact is that almost no question ever ends up using this and I have still included it.(Don't worry, we'll use it later, It was a joke)

However, here is a token example.

Example 21.2. Prove that there are no positive integers x, k and $n \geq 2$ such that $x^2 + 1 = k(2^n - 1)$

Proof. We need to notice that $2^n - 1 \equiv 3 \pmod{4}$. Which means some prime $p \equiv 3 \pmod{4}$ divides it.

Hence, $x^2 + 1 \equiv 0 \pmod{p}$ which is false by Fermat's Christmas Theorem. □

21.2 Orders

Now here is a much more simple and useful topic.

Definition 21.3. Let p be a prime and a not divisible by p . Then the order of a modulo p is defined to be the smallest positive integer n such that $a^n \equiv 1 \pmod{p}$. We denote it as $\text{ord}_p a$.

Here are somethings we need to notice, and prove.

Theorem 21.4. For any $a, p \in \mathbb{N}$ where p is prime and $a \not\equiv 0 \pmod{p}$, we have $\text{ord}_p a = n|(p - 1)$.

Proof. We know from Fermat's little theorem that $a^{p-1} \equiv 1 \pmod{p}$ which means that if $a^n \equiv 1 \pmod{p}$ and n is smallest possible value to do so. This means $a^n - 1|a^{p-1} - 1$ which we can also write as $\frac{a^{p-1}-1}{a^n-1} = k$, this looks extremely like an GP, doesn't it?

$1 + a^n + a^{2n} + \dots + a^{p-1-n} = k$, also to note is the fact that all exponents are of the type an and therefore the next term of GP is $p - 1$ which is equal to an for some a . This means, $n|(p - 1)$ which means $\text{ord}_p a = n|(p - 1)$. \square

This also implies that the value of order of a modulo p is less than or equal to $p - 1$. The equal to $p - 1$ has a special name.

Definition 21.5. Let p be a prime. Then there exists an integer g , called a primitive root, such that $\text{ord}_p g = p - 1$.

We also need to note that:

Theorem 21.6. $g^{\frac{p-1}{2}} \equiv -1 \pmod{p}$ for g being the primitive root of p and $p > 2$

Proof. $g^{p-1} \equiv 1 \pmod{p} \iff g^{\frac{p-1}{2}} \equiv \pm 1 \pmod{p}$, but since $\frac{p-1}{2} < p - 1$, If $g^{\frac{p-1}{2}} \equiv 1 \pmod{p}$ than it will be the primitive root, which is not contradictory. Hence, $g^{\frac{p-1}{2}} \equiv -1 \pmod{p}$. \square

With this out of the way, we can finally solve some questions:

Example 21.7. Find all n such that $n|2^n - 1$

Proof. We can let p be the smallest prime factor of n . Which means $2^n \equiv 1 \pmod{p} \iff \text{ord}_p 2 | n$. This means $\text{ord}_p 2 | p - 1$.

This means $\text{ord}_p 2 | \gcd(p - 1, n)$. As p is the smallest factor of n , it clearly means that $\gcd(p - 1, n) = 1$. This means $1 \geq \text{ord}_p 2$ which forces $\text{ord}_p 2 = 1$.

As no natural number is less than 1, we know that $n = \text{ord}_p 2 = 1$ which means the only n satisfying the following is 1. \square

Another similar thing to try is:

Example 21.8. Prove that every prime divisor of $2^p - 1$ is greater than p

Proof. This is somewhat easier to do. Let q be the smallest prime factor of $2^p - 1$.

That means $2^p \equiv 1 \pmod{q}$. This means $\text{ord}_q 2 | p$.

This means that $\text{ord}_q 2 = 1$ or p . If it is equal to 1, $2^1 - 1 \equiv 0 \pmod{q}$ which is absurd as no prime is less than equal to 1. Then $\text{ord}_q 2 = p$ which means $p | q - 1$ which allows us to say $p \leq q - 1 < q$. Hence, the smallest prime factor is greater than p . \square

21.3 Chicken McNugget Theorem

After all these heavy topics, let's talk about a rather light one.

Famous problem writer, Henri Picciotto was dining at McDonalds with his son when he noticed that nuggets were sold in packs of 6, 9 and 20. He taught about what is the largest number of nuggets he could order which could not be packed in these boxes. It is said that he worked it all out on a napkin.

The problem then appeared in Games Magazine in 1987.

Let's see if we can find the solution.

We are looking for the largest N such that $6x + 9y + 20z = N$ has no solution. Let's first find all elements $M \in m$ which can't be written as $6x + 9y = M$ and then get rid of all such that $M - 20k \notin m$.

What we need to notice that all multiples of 3 are $\notin m$. What we need to notice is that we can subtract 20 to make anything a multiple.

So what we are looking for are numbers from which when we subtract either 20 if they are 1 modulo 3 and 40 if they are 2 modulo 3. The subtracted number is a multiple of 3 and we need to prove that it can't be partitioned into 6, 9.

We can see (and will formally prove in a minute) that all multiples of 3 can be partitioned into 6, 9 other than 3 itself. Hence, the answer is $3 + 40 = 43$.

Theorem 21.9 (Chicken McNugget Theorem). *For any two relatively prime positive integers m, n , the greatest integer that cannot be written in the form*

$am + bn$ for nonnegative integers a, b is $mn - m - n$.

A consequence of the theorem is that there are exactly $\frac{(m-1)(n-1)}{2}$ positive integers which cannot be expressed in the form $am + bn$

Proof. We know from Bezout's that infinite integer solutions to $am + bn = N$ exist every time we have $\gcd(m, n)|N$. This is true for all N as $\gcd(m, n) = 1$ as they are coprime.

So we always have integer solutions. But the condition of never having positive integer solution is true if and only if either one of the coefficients is all way negative.

Using linear Diophantine, let a solution be (x, y) and then the set of solutions is $(x - kn, y + km)$. Let x be positive and y be negative, and (x, y) be the case where y is the greatest(ie closest to zero). In this case $|y| \geq n$ Adding to y to make it positive should lead to x becoming negative.

For maximum value of N we can let $y = -1$ and $x = n - 1$ as it will follow the above condition.

$$N = m(n - 1) - n$$

$\iff N = mn - m - n$ Hence, proved. □

Now you may feel that 6, 9 are not co-prime, but we can take 3 common. Then we are left with 2, 3 which are coprime. Using the Chicken Mcnugget theorem we can say that $6 - 3 - 2 = 1$ is the largest number not partitionable. This means all other multiple of 3 other than 3 can be made by adding 6, 9. Hence, 43 is the answer.

Let's look at a very basic question:

Example 21.10. (AMC 10B 2015)The town of Hamlet has 3 people for each horse, 4 sheep for each cow, and 3 ducks for each person. What is the largest number which could not possibly be the total number of people, horses, sheep, cows, and ducks in Hamlet?

Solution. $P = 3H, S = 4C, D = 3P$ from the question.

Thus the total number of living organisms is $P + H + S + C + D = 3H + H + 4C + C + 9H = 13H + 5C$

which has no solution for $13 * 5 - 5 - 13 = 47$ living organisms. □

So far so good, let's do something better

Example 21.11 (AIME II 2019). Find the sum of all positive integers n such that, given an unlimited supply of stamps of denominations 5, n , and $n + 1$ cents, 91 cents is the greatest postage that cannot be formed.

Solution. By the Chicken McNugget theorem, the least possible value of n such that 91 cents cannot be formed satisfies $5n - (5 + n) = 91 \implies n = 24$,

so n must be at least 24.

For a value of n to work, we must not only be unable to form the value 91, but we must also be able to form the values 92 through 96, as with these five values, we can form any value greater than 96 by using additional 5 cent stamps.

Notice that we must form the value 96 without forming the value 91. If we use any 5 cent stamps when forming 96, we could simply remove one to get 91. This means that we must obtain the value 96 using only stamps of denominations n and $n + 1$.

Recalling that $n \geq 24$, we can easily figure out the working $(n, n+1)$ pairs that can be used to obtain 96, as we can use at most $\frac{96}{24} = 4$ stamps without going over. The potential sets are $(24, 25), (31, 32), (32, 33), (47, 48), (48, 49), (95, 96)$, and $(96, 97)$.

The last two obviously do not work, since they are too large to form the values 92 through 94, and by a little testing, only $(24, 25)$ and $(47, 48)$ can form the necessary values, so $n \in \{24, 47\}$. $24 + 47 = 071$. □

And finally a question from Indian TST

Example 21.12. (India) On the real number line, paint red all points that correspond to integers of the form $81x + 100y$, where x and y are positive integers. Paint the remaining integer points blue. Find a point P on the line such that, for every integer point T , the reflection of T with respect to P is an integer point of a different colour than T .

Solution. It is easy to notice that the answer is $P = \frac{81*100 - 81 - 100}{2}$. Let's prove it.

Let's assume that this is not the correct answer. Let's assume, to the contrary, that there exists a k such that both are red, that is:

$$81x + 100y = P - k$$

$$81a + 100b = P + k$$

Adding the equations and letting $x + a = k$ and $y + b = l$, we get:

$81k + 100l = 2P = 81 * 100 - 81 - 100$ which is false from Chicken McNugget Theorem.

Hence, Contradiction.

Thus, no such k exists. Hence, $P = \frac{81*100 - 81 - 100}{2} = \frac{7919}{2} = 3959.5$ is the required point. □

21.4 Pell's Equations

I didn't want to include this, it had only been seen in TST's and that too in very rare cases. But then some examiner decided to ask it in AMC (It's the

example at the end of the section) and here we are.

Definition 21.13. An equation of the form $x^2 - dy^2 = 1$ where d is square free is called Pell's Equation.

d needs to be square free as otherwise there will be no integer solutions by difference of squares.

Another thing to note is:

Theorem 21.14. *If a Pell's equation has one solution, then it has infinitely many.*

We will prove this by actually generating the infinite solutions.
We begin by defining some terms.

Definition 21.15. Characteristic complex(z) of a Pell's equation is $x + y\sqrt{d}$ and its conjugate is \bar{z} which is $x - y\sqrt{d}$.

The norm is defined as $N(z) = z\bar{z} = x^2 - dy^2$

This is quite similar to complex numbers we dealt with in algebra. This is because Pell's equation is studied in a field of maths aptly named Algebraic Number Theory where algebraic structures are used to study the properties of numbers.

We'll however not talk about it much.

We can notice that $N(ab) = N(a)N(b)$ which is trivial to prove using basic algebra (and also follows from the complex plane), therefore we can say that if $N(z) = 1$ then $N(z^k) = 1$

Which means that if we can find a single solution of a pell's equation, then we can find the characteristic complex of that solution and hence the rest of the infinite solutions.

Let me illustrate with an example:

Example 21.16. (AMC 12A 2022) A *triangular number* is a positive integer that can be expressed in the form $t_n = 1 + 2 + 3 + \dots + n$, for some positive integer n . The three smallest triangular numbers that are also perfect squares are $t_1 = 1 = 1^2$, $t_8 = 36 = 6^2$, and $t_{49} = 1225 = 35^2$. What is the sum of the digits of the fourth smallest triangular number that is also a perfect square?

Solution. The question is looking for solutions to:

$$\frac{n(n+1)}{2} = k^2$$

$$\begin{aligned} &\iff n^2 + n = 2k^2 \\ &\iff 4n^2 + 4n = 8k^2 \\ &\iff 4n^2 + 4n + 1 - 8k^2 = 1 \\ &\iff (2n+1)^2 - 2(2k)^2 = 1 \end{aligned}$$

This is a Pell's equation we already know a solution to which is $(k, n) = (1, 1)$.

Which means $z = 3 + 2\sqrt{2}$

$$z^2 = 9 + 8 + 12\sqrt{2} = 17 + 12\sqrt{2} \implies (k, n) = (8, 6)$$

$$z^4 = 289 + 288 + 408\sqrt{2} = 577 + 408\sqrt{2} \implies (k, n) = (288, 204)$$

$$204^2 = 41616$$

Thus, the answer is $4 + 1 + 6 + 1 + 6 = 18$

□

21.5 Floor, Ceiling and fractional function

Before we talk about the black boxes, let's explore the final topic which we would reasonably understand.

Definition 21.17. The greatest integer function(GIF) function represents greatest integer less than or equal to x where $x \in \mathbb{R}$. We represent it using $\lfloor x \rfloor$. It is also called the floor function.

The ceiling function represents the smallest integer more than or equal to x where $x \in \mathbb{R}$. We represent it using $\lceil x \rceil$.

The fractional function represents the part of x which cannot be represented as integer. That is $x - \lfloor x \rfloor$. It is represented as $\{x\}$

The ceiling function is simply the GIF function plus 1. Hence, we don't use it normally. In this chapter, $[x]$ represents the greatest integer function. With the formal introduction out of the way, let's have some fun now.

Example 21.18. (PRMO 2017 , edited)Find the maximum value of x such that $\{x\}, [x], x$ form a geometric progression.

Solution. We can simply use the fact $x = [x] + \{x\}$ to solve this question.

$$\begin{aligned} &\iff [x]^2 = [x]\{x\} + \{x\}^2 \\ &\iff \{x\}^2 + [x]\{x\} - [x]^2 \\ &\iff 0 \leq \{x\} = \frac{-[x] + \sqrt{[x]^2 + 4[x]^2}}{2} < 1 \\ &\iff 0 \leq [x](-1 + \sqrt{5}) < 2 \\ &\iff 0 \leq [x] < \frac{2}{\sqrt{5}-1} = \frac{(\sqrt{5}+1)}{2} = 1.61\dots \\ &\therefore [x] = 1 \end{aligned}$$

$$\therefore \{x\} = \frac{\sqrt{5}-1}{2}$$

$$\therefore x = \frac{\sqrt{5}+1}{2}$$

□

This was good but let's take it a notch higher

Example 21.19. Find all x such that $\frac{1}{[x]} + \frac{1}{[2x]} = \{x\} + \frac{1}{3}$

Proof. Here the key fact to note is that $[2x] = 2x$ or $[2x] = 2x + 1$ depending on the fractional part.

Hence, we can divide the question into two cases.

$$\frac{1}{3} \leq \{x\} + \frac{1}{3} = \frac{3}{2[x]} < \frac{5}{6}$$

The RHL is due to the fact that if the fractional part is 0.5 or more than we can't have $[2x] = 2[x]$.

$$\begin{aligned} 3 &\geq \frac{2[x]}{3} > \frac{6}{5} \\ \iff 4.5 &\geq [x] > 1.8 \\ \therefore [x] &= 2, 3, 4 \end{aligned}$$

Now onto the second case.

$$\begin{aligned} \frac{5}{6} &\leq \{x\} + \frac{1}{3} = \frac{1}{[x]} + \frac{1}{2[x]+1} < \frac{4}{3} \\ \iff \frac{5}{6} &\leq \frac{3[x]+1}{2[x]^2+[x]} < \frac{4}{3} \\ \iff 5 &\leq \frac{18[x]+6}{2[x]^2+[x]} < 8 \\ \iff 10[x]^2+5[x] &\leq 18[x]+6 < 16[x]^2+8[x] \\ \iff 10[x]^2-13[x] &\leq 6 < 16[x]^2-10[x] \\ \iff 1 &< [x] \leq 1.6\dots \end{aligned}$$

Which means there is no possible $[x]$.

Thus, we only have three values of x which occur at $[x] = 2, 3, 4$

□

21.6 Black Boxes

We are finally in the section where I'll state some very powerful results without proof. More times than not, they can be cited in a math contest without much worry. This section can also be skipped as you are just memorizing a theorem without proof. The choice is yours. We'll follow every theorem by an example and use the theorem to solve it.

Theorem 21.20 (Schur's Theorem). *If a polynomial $P(x)$ with integer coefficients is non constant, then the set of prime factors of $P(x)$ for all $P(x) \in \mathbb{N}$ is infinite.*

Another contribution by the great Issai Schur. We'll use it on this US-AMO 3

Example 21.21. (USAMO/3 2006) For integral m , let $p(m)$ be the greatest prime divisor of m . By convention, we set $p(\pm 1) = 1$ and $p(0) = \infty$. Find all polynomials f with integer coefficients such that the sequence $\{p(f(n^2)) - 2n\}_{n \geq 0}$ is bounded above. (In particular, this requires $f(n^2) \neq 0$ for $n \geq 0$.)

Proof. Suppose that f satisfies the condition. We suppose that f is irreducible, otherwise look at all irreducible factors. Let $g(x) = f(n^2)$

Assume that $p|g(k)$. We can assume $0 \leq k \leq \frac{p}{2}$, since obviously $g(p \pm k) \equiv g(k) \pmod{p}$.

Then, $p \leq 2k + r$, for some r . So we have either $k = \frac{p}{2}, k = \frac{p-1}{2}, \dots, k = \frac{p-r}{2}$. We know that infinitely many prime number divide the series of $g(n)$ due to Schur. Therefore for some $0 \leq j \leq r$ there are infinitely many primes p for which $p|g(\frac{p-j}{2})$. This implies that $p|g(-\frac{j}{2})$ for infinitely many p , so $g(-\frac{j}{2}) = 0$, thus $2x + j|g(x^2)$, and also $2x - j|g(x^2)$, so $4x^2 - j^2|g(x)$, so $4x - j^2|f(x)$. As f is irreducible, $f = 4x - j^2$.

□

While Schur only played a minor role in this example, Our next theorem will destroy it's following example.

Theorem 21.22 (Anti-Schur). *If a polynomial $P(x)$ with integer coefficients is non-constant and non-reducible, then there exists a prime p such that $P(x)$ has no roots modulo p .*

Now let's hunt down an IMO 6.

Example 21.23. (IMO/6 2003) Let p be a prime number. Prove that there exists a prime number q such that for every integer n , the number $n^p - p$ is not divisible by q .

Proof. The given polynomial is integer and irreducible. Which means by anti-schur, there exists some prime q which gives us $n^p - p \not\equiv 0 \pmod{q}$ which means it is not divisible.

And we are done.

□

This felt almost like cheating, didn't it?

Theorem 21.24 (Kobayashi's theorem). *Let $t \in \mathbb{Z}$ and a set of number a_1, a_2, \dots*

If the set of prime factors of a_1, a_2, \dots is finite; then the set of prime factors of $a_1 + t, a_2 + t, \dots$ is infinite.

I find this theorem quite unsettling.

The unsettling part is who really proved it. It says a high school student named Hiroshi Kobayashi from Ebina Highschool. But that school doesn't exist. No other work by that person either. Just one citation to some Olympiad handout. Makes you wonder...who really did prove this result?

Example 21.25. (STEMS 2021) Determine all non-constant monic polynomials $P(x)$ with integer coefficients such that no prime $p > 10^{100}$ divides any number of the form $P(2^n)$

Proof. For $P(x) = x^d$, Let $P'(x) = x^d + a$ where a represents rest of the equation.

From Kobayashi, $P'(x)$ has infinite prime factors and hence for some n , $P'(2^n)$ will have a factor greater than 10^{100} .

Therefore, $P(x) = x^d$ are the only solutions. □

The next theorem is very useful, especially if you want to date a math lover. Pronounce the name right, and they'll love you forever.

Definition 21.26. Zsigmondy Set refers to the set of n such that for a_n of the series a_1, a_2, \dots all prime p such that $p|a_n$, $p|a_k$ for some $k < n$. It is denoted as $\mathcal{Z}\{a_n\}$

Read the definition twice to actually internalize what it means.

Theorem 21.27 (Zsigmondy Theorem). *If a and b are relatively prime, then $\mathcal{Z}\{a^n - b^n\} \subseteq \{1, 2, 6\}$. In particular:*

- $1 \in \mathcal{Z}\{a^n - b^n\} \iff a - b = 1$
- $2 \in \mathcal{Z}\{a^n - b^n\} \iff a + b = 2^n, n \in \mathbb{N}$
- $6 \in \mathcal{Z}\{a^n - b^n\} \iff a = 2, b = 1$

Similarly, $\mathcal{Z}\{a^n + b^n\} = \emptyset$, with the exception of $2^3 + 1^3$

The last line in itself is a weaker form of Zsigmondy which is used a lot in Olympiads.

Also an elementary proof to the last part of Zsigmondy exists, I however leave finding it out to you.

Example 21.28. Find all triplets (a, n, k) of positive integers such that

$$a^n - 1 = \frac{a^k - 1}{2^k}.$$

Solution. Using Zsigmondy, there is some prime p which divides $a^k - 1$ but doesn't divide $a^n - 1$

This means $p|2^k(a^n - 1)$ which means $p = 2$

This is obviously not true as if the RHS is divisible by 2 which means a is odd which makes LHS even which contradicts the initial assumption.

Hence, we'll have to look at the exception case of Zsigmondy. First we'll note that $a = 1$ which trivially works.

The second is k being in \mathcal{Z} . We'll check it on a case by case basis.

Let $k = 1$, then $n < k$ but that is not possible so this doesn't work.

Let $k = 2$, then $n = 1$ which means:

$$a - 1 = \frac{a^2 - 1}{4}$$

$$\iff 4 = a + 1$$

$$\iff a = 3$$

Then another solution is $a = 3, n = 1, k = 2$

Finally, $k = 6$ for which $a = 2$ by Zsigmondy. But that violates $\frac{a^k - 1}{2^k}$ and hence no such solution exists.

Thus, the only solutions are $(a, n, k) = (1, n, k)$ and $(3, 1, 2)$ \square

Another very useful theorem is:

Theorem 21.29 (Bertrand's postulate). *If $n > 3$, then there's some prime p such that $n < p < 2n$.*

This has an elementary proof, which was found by the great Paul Erdos, it's just tedious. I have decided to not include the proof but greatly encourage you to explore it with the help of internet. Now let's use the theorem.

Example 21.30. Find all n such that

$$n! + (n+1)! + (n+2)!$$

is a square number.

$$\begin{aligned} \text{Solution. } & n! + (n+1)! + (n+2)! \\ &= n!(1+n+1+(n+1)(n+2)) \\ &= n!(n+2)(n+2) \\ &= n!(n+2)^2 \end{aligned}$$

This means we are looking for $n!$ which is a square number.

There is some p such that $\frac{n}{2} < p < n$ from Bertrand's theorem.

For $n!$ to be a square number, we need to have $2p < n$, but:

$$n < 2p < 2n$$

This is a contradiction. This means no such n exist.

But as we have used Bertrand, hence we need to check for $n < 3$

As $0! = 1! = 1^2 = 1$ and $2! = 2 \neq k^2, k \in \mathbb{N}$, it means which means that the only solution are $n = 0, 1$. \square

Let's look at another question

Example 21.31. Prove that $1, \dots, 2n$ can be partitioned into n pairs such that the sum of the numbers in each pair is prime

Proof. We need to note that $2n < p < 4n$ from Bertrand.

This means $p = 2n + m$ for some m . This means $\{m, \dots, 2n\}$ will recursively all get paired as $m + 1 + 2n - 1 = m + 2n = p$

We can now repeat the same for $m - 1$ and so on recursively.

Hence, proved. \square

Finally a theorem which is a glimpse of the next chapter.

Theorem 21.32 (Dirichlet's Theorem). *There are infinite primes of the form $a + nd$ given $\gcd(a, d) = 1$.*

This may either seem obvious or surprising, in any case it is a black box which requires a lot of advanced math to prove. But let's have some fun.

Example 21.33. Show that there are infinitely many positive numbers n that cannot be written as $3ab + a + b$ for any $a, b \in \mathbb{N}$.

Proof. We need to prove that there are infinite such n that $3ab + a + b = n$ This means $3n + 1 = 9ab + 3a + 3b + 1$ has no solutions This means $3n + 1 = (3a + 1)(3b + 1)$ has no solutions.

This will be true if $3n + 1$ is prime. There are infinite such n due to Dirichlet's Theorem. \square

And as we end the chapter, I'll revisit the first black box we saw:

Theorem 21.34 (Prime Number Theorem). *Number of primes less than $n \approx \frac{n}{\ln n}$ and consequently, the n^{th} prime $\approx n \ln n$*

But this form is less fun. A little transformation will give us a much stronger form.

Theorem 21.35 (Prime Number Theorem's Bertrand's form). *For every $\varepsilon > 0$ we have some natural number $n_\varepsilon \in \mathbb{N}$ such that for all $n > n_\varepsilon$, we have some prime p such that $n < p < (\varepsilon + 1)n$*

The proof, while not elementary, can be done using limits (the approx sign is just limit of n tending to ∞).

Example 21.36. Prove that exist infinity prime number which began (from left to right) with 9.

Proof. From the prime number theorem, it follows that for every $\varepsilon > 0$, there exists a natural number N_ε such that for all $n > N_\varepsilon$, there is a prime between n and $(1 + \varepsilon)n$. Applying this to $\varepsilon = \frac{1}{9}$ and $n = 9 \cdot 10^k$ for $k > \log_{10} \frac{N_1}{9}$, we are done. \square

Exercises

(1) (IMO 2005, 4) Determine all positive integers relatively prime to all the terms of the infinite sequence

$$a_n = 2^n + 3^n + 6^n - 1, \quad n \geq 1.$$

(2) (HMMT 2014) Determine all positive integers $1 \leq m \leq 50$ for which there exists an integer n for which m divides $n^{n+1} + 1$.

(3) (IMOSL 1991) Find the largest integer k for which 1991^k divides:

$$1990^{1991^{1992}} + 1992^{1991^{1990}}$$

(4) (AIME 1994) Ninety-four bricks, each measuring $4'' \times 10'' \times 19''$, are to be stacked one on top of another to form a tower 94 bricks tall. Each brick can be oriented so it contributes $4''$ or $10''$ or $19''$ to the total height of the tower. How many different tower heights can be achieved using all ninety-four of the bricks?

- (5) (AIME II 2019) Find the sum of all positive integers n such that, given an unlimited supply of stamps of denominations 5, n , and $n + 1$ cents, 91 cents is the greatest postage that cannot be formed.
- (6) (RMO 2018) Find all natural numbers n such that $1 + [\sqrt{2n}]$ divides $2n$.
- (7) Find $n \in \mathbb{N}$ satisfied: $n^2 + 3^n$ is a square number
- (8) (Brazil 2018) Let $a_0 = a > 1$ and $a_{n+1} = 2^{a_n} + 1$. So that the set of prime divisors of $\{a\}_{n \geq 0}$ is infinite.
- (9) (Hong Kong TST 2018) Find infinitely many positive integers m such that for each m , the number $\frac{2^{m-1} - 1}{8191m}$ is an integer.

22

Constructions

We have dealt with some constructions already. But this time we deal with them formally.

Construction refers to showing that infinite solutions of an equation exist by finding one solution and then using it to generate a family of solutions.

It is somewhat like infinite descent but in reverse.

Sometimes a question is of the form "Does there exist..." these problems become impossible if we assume from the get go that no such answer exists as more often than not there exists. If all constructions fail, then we can assume non existence and then prove the same.

All this will become more clear as we solve some questions.

22.1 Chinese Remainder Theorem

We have already seen this theorem and will now use it for more powerful purposes.

Example 22.1. (IMO 1989) Prove that for each positive integer n there exist n consecutive positive integers none of which is an integral power of a prime number.

Proof. We are basically looking to prove that there are n consecutive positive integers which all have two prime factors or more.

Using the CRT there exists some x such that:

$$x + 1 \equiv 0 \pmod{p_1 q_1} \iff x \equiv -1 \pmod{p_1 q_1}$$

$$x + 2 \equiv 0 \pmod{p_2 q_2} \iff x \equiv -2 \pmod{p_2 q_2}$$

$$\therefore x + n \equiv 0 \pmod{p_n q_n} \iff x \equiv -n \pmod{p_n q_n}$$

For two series of different primes p_1, \dots, p_n and q_1, \dots, q_n . This means $x + 1, x + 2, x + 3, \dots, x + n$ are n such integers where none are integral power of any prime.

□

Here is another way to use CRT:

Example 22.2. (Math Prize 2010) Prove that for every positive integer n , there exists integers a and b such that $4a^2 + 9b^2 - 1$ is divisible by n .

Proof. We can prove this by simply showing $4a^2 + 9b^2 - 1$ is divisible by p^k for some a and b .

$$\therefore 4a^2 + 9b^2 \equiv 1 \pmod{p^k}$$

$$\therefore 4a^2 = 1 \pmod{p^k} \iff a \equiv \frac{1}{2} \pmod{p^k} \text{ and } b \equiv 0 \pmod{} \quad \square$$

Exercises

- (1) (USAMO/1 2008) Prove that for each positive integer n , there are pairwise relatively prime integers k_0, \dots, k_n , all strictly greater than 1, such that $k_0 k_1 \cdots k_n - 1$ is the product of two consecutive integers.

Part 8

Geometry

23

Angle Chasing

Geometry and I have a relation which the less spoken about, the better. We don't get along. While I have a better appreciation for geometry now, I still am somewhat fearful of it. The main reason of my problems with Geometry is fundamentally due to the fact that I memorized theorems early on without understanding(or reading the proof) in many cases.

This is something you need to avoid from the get go.

This chapter is about one of the most critical skills for a geometer. Using data about some angle to find others by chasing. We'll learn geometry through problems rather than through theorems,

A lot of angle chasing is easier to actually solve then to follow from the solution, so maybe ignore my answers and try to solve everything yourself...

23.1 Triangle Centers

We can define the center of triangle in a lot of ways, this leads to there being many centers of the same triangle each one correct based on some criterion. We'll talk about 4 such centers now.

Definition 23.1. • The orthocenter of ABC , usually denoted by H , is the intersection of the perpendiculars (or altitudes) from A to BC , B to CA , and C to AB . The triangle formed by the feet of these altitudes is called the orthic triangle.




Figure 23.0. The four primitive centers

- The centroid, usually denoted by G , is the intersection the medians, which are the lines joining each vertex to the midpoint of the opposite side. The triangle formed by the midpoints is called the medial triangle.
- The incenter, usually denoted by I , is the intersection of the angle bisectors of the angles of ABC . It is also the center of a circle (the incircle) tangent to all three sides. The radius of the incircle is called the inradius.
- The circumcenter, usually denoted by O , is the center of the unique circle (the circumcircle) passing through the vertices of ABC . The radius of this circumcircle is called the circumradius.

23.2 Cyclic Quadrilaterals

Part 9

Extras

24

Anti-problems

One of the first chapters of this book contained questions which had complex solutions but easy to understand formulations. As I close this book, this chapter is the anti-thesis. The questions will have complex formulations but very simple solutions.

However, these are not birthday cakes. The solutions tend to be conceptual and hard to notice. Such problems were earlier called 'Jewish Problems' as Russian Anti-Semitic professors to keep out the Jews and other 'undesirables' from their institutes. While you will surely find some problems fell easy, that is mostly as we have explored such ideas before, in those days, such access was rare. In addition, the students were given these problems one after another until they failed one of them, at which point they were given a failing mark. They were basically like "Olympiad-style" questions, except that you were expected to answer them with far less time than in an actual Olympiad, and work them out orally. Sure you could probably work a few out with paper and pencil given the time, but perhaps you wouldn't always be given a paper and pencil. Often the answer itself would be something trivial or even intuitive, but the trick was rigorously proving your answer beyond the shadow of a doubt. Basically you could be asked to stop at any time and explain the reasoning to your interviewer, and perhaps even asked to take a different line of reasoning. Also, more than the problems being unsolvable - they were also generally designed to be ambiguous in such a way that the examiner could pick whichever interpretation made the candidate's solution invalid.

The problem with the "Jewish questions" was not that they were difficult. It was that they were being administered unfairly.

It is not wrong for an university to ask these sorts of questions on their entrance exams, if they are looking to select for talented applicants who can come up with their own solutions and/or have already learned concepts more advanced than just high school common core.

The problem with the "Jewish questions" was that not all applicants would be asked them. Most applicants would get the fairly standard "solve this quadratic equation" type of questions. But when the examiner would see somebody from an "undesirable caste" of people (e.g. a Jewish student), they would swap the typical questions with one of these problems, which they would almost surely be unable to solve in the time given.

And the entire point of doing this was plausible deniability: The solutions to these problems, once you already know the right way to do it, when written out, do not take up any more space than a solution to a more standard problem would. Therefore, the examiner/institution could feasibly claim impartiality, since to an uninitiated inspector both types of questions seem "roughly similar in difficulty". The institution can pretend to an outside observer that all students are being tested "fairly", since the correct solutions to both kinds of problems look "similar"; while at the same time immediately offing any "undesirable" applicants using a method that is hard to prove (or even spot in the first place) if you are yourself not an expert in mathematical problem solving and testing.

But I will not bore you with history. Here are the problems:

24.1 Anti-Problems

Exercise

(1)

25

Engineering

We are normally shocked at the marvelous pay top engineers earn. It almost feels like they know something that the normal population doesn't. Is it math? no. If you are reading this chapter towards the end, you probably are better at math than most engineers.

Is it coding? Not really. A lot of coders just copy and hack code from google. Is it that they are smarter than the rest of the population? Based on my conversations and IQ tests, Negitive.

So why? because they have learnt mastered the art of engineering. Engineering is finding the answer to a problem without solving it. It's the art of problem solving minus the solving part, we don't do that here. It is finding the answer with less effort than it would take us if we did the problem "properly".

While engineering is a useful skill, it is by no means replacement for learning math. Remember the high ball engineers we talked about can still code if google stopped working. In fact, it will be them who will get it to start working again. They can solve the problem, but for efficiency use such methods.

It shouldn't be satisfying to get an answer with engineering. Engineering a problem is like taking out a loan, that you need to repay with learning the actual solution to the problem. THIS CHAPTER IS NOT FOR PEOPLE WHO ARE YET TO SOLVE THE REST OF THE BOOK.

That is a disclaimer, as if misunderstood this chapter can do more harm than good, and don't say I didn't warn you.

25.1 Abuse of Degree of Freedom

Example 25.1. (PMO 2019) In triangle ABC , D and E are points on sides AB and AC respectively, such that BE is perpendicular to CD . Let X be a point inside the triangle such that $\angle XBC = \angle EBA$ and $\angle XCB = \angle DCA$. If $\angle A = 54^\circ$, what is the measure of $\angle EXD$?

While I assume you can already solve this geometrically, the engineering solution comes by noticing that the question is general, hence if D was placed right on A , and therefore $BE \perp AC$ and therefore X is placed right on B making this a simple angle sum.




Figure 25.0. The question diagram as we engineer it out of existence

Example 25.2. Let S be any point within a rectangle $ABCD$. On AS , a square $ASRF$ is described and E is a point on AB such that $AE = AD$. The value of $((\frac{AF+ES}{FD+AS})^{\frac{1}{2}} + 3)^{\frac{1}{2}}$

This question dies a sad death by simply letting $E = S$

Part 10

Appendix

A

Appendix

Appendix A: Justifications, Ramifications and Excuses

This chapter is me trying to explain why I made whatever choices I have made. While this is an appendix, so most of the readers are expected to skip over this, I recommend the teachers using the book to read through. You may resonate with some of the ideas here and hopefully get something new for your classroom.

A.1 Introductory Problems

B

Appendix

Appendix B: Hints

- (1) As we know m now, can we simply use stars and bars with one red marble always between 2 green.
- (2) Try to compute the fraction to a few more terms and look for patterns.
- (3) Case work onto number of same color triangles on the outside.
- (4) We have already solved a case of it as an AIME 2015 problem, won't the same technique work in general?
- (5) Think about what the end case can be? At what case will the transfers stop mattering, and the entire thing will become invariant?
- (6) While some lines will intersect outside, won't every four points give us a pair of two lines intersecting inside?
- (7) In what ways can the men be arranged in? Use the groups of men as bars and the women as stars
- (8) Focus on the color of the center triangle
- (9) Case work onto the possible coin flips results
- (10) After the first weighing we are sure of some coins. Let's use them to our advantage now.
- (11) How did you solve the case with 12 coins and 3 weighing? The same method is applicable in general.
- (12) Think Binary
- (13) Let $a = 1 + x$, $b = a + y$ and so on. Does this look like a Stars and Bars now?
- (14) Is rolling in anyway different from just pulling two numbers from the bag?

- (15) If we get a BB or WW in a column, the other is determined uniquely
- (16) m is limited by the fact that number of marbles whose right hand neighbor is the other color is maximum 10. How many red marbles will we need for this to lead to a valid arrangement?
- (17) Think about why we really don't care which books Harold selects.
- (18) What is the largest number you can form from an arbitrary set of digits? For example, 5986772. Why did you take the approach you did?
- (19) $a_1 + 2a_2 + 3a_3 + \dots + na_n = (a_1 + a_2 + \dots + a_n) + (a_2 + a_3 + \dots) + \dots + (a_n)$
- (20) The integers from 8 through 14 must be in different pairs, and 7 must pair with 14.. Why is this true? And why does this solve this question?
- (21) Take the sides as $x, y - x, n - y$ where n is the length of the stick.
- (22) Start with 1 dragon. Than 2. Than 3... A pattern becomes obvious.
- (23) Just build a 9 cube tower and ignore the last block.
- (24) In what ways can we get an odd sum? Can you arrange E and O in such a way that this is true? How many rearrangements does your arrangement have?
- (25) In how many ways can we add up to 7? What does that tell us about ways to add up to 10? What does that tell us about one of the dice faces?
- (26) Multiply both sides by 2 and use $\binom{n}{k} = \binom{n}{n-k}$
- (27) Just write all the possible draws till we exceed 4 and you'll be done in no time.
- (28) Add terms to reach an identity which we know the summation to
- (29) Can we combine the b and c?
- (30) Let's start by weighing any four coins, against other four (setting aside the remaining four).
- (31) Why will it slide down after reaching the top of the pole?
- (32) Assume that the balls and bins are both distinguishable and then this is just question 2, but a bit more involved.
- (33) Can we convert the choosing of 3 points as partitioning of remaining 5 points into 3?
- (34) We can draw BB, BR, RB or RR. What happens in each case?
- (35) Complimentary counting seems quite fine
- (36) Casework on the tenors – basses combined with Vandermonde will work.
- (37) Simplify before you compute! Take 9! common, multiply numerator and denominator by 3!4!5!
- (38) 1 and positive integers divisible by 5 will not be in the set. Why? What about all the other integers?

- (39) We'll use Vandermonde identity, now try to solve it.
- (40) The cases seem to be less enough that maybe just counting for one colour followed by some multiplication is good sauce.
- (41) What is the value of the sum of squares of the numbers written on the board as we continue doing this?
- (42) Try casework about the number of diameters used
- (43) Try using casework on the fact that the opposite edges have same number or not.
- (44) With such a strict condition and only $4!$ maximum cases, writing by hand seems quite promising.
- (45) Experiment with smaller cases. Maybe you'll notice something
- (46) Multiply both sides by $19!$
- (47) Just compute the fraction, now think why it makes sense intuitively as well.
- (48) In how many ways can the friends be put into rooms? Like we can have one friend per room, 2 in one room and rest in single room, 2 sharing a room, another 2 sharing a room and one person living in a single room
- (49) Try to determine which weekdays we had five off in that month then.
- (50) Let n be the ways to get 12 and then try solving the equation.
- (51) Write by hand, the condition is quite stringent
- (52) Think in terms of percentage of the journey.
- (53) We have n possible reference points
- (54) We can notice that if abc is such a number then cba is also such number
- (55) Use Complementary counting and the PIE
- (56) Just count!
- (57) Casework onto the box in which the three math textbooks are in
- (58) Make the winning combinations and then try to arrange the 3 X's in any non winning way.
- (59) Try fixing a column
- (60) You'll also need to do some casework on the number of red squares
- (61) Think about the cases when he'll roll 0, 1, 2 or 3 dice
- (62) If we have x B's in the first 5 letters and $5 - x$ C's, what will happen? Does this solve the question?
- (63) The answer is 3. Try walking backward now.
- (64) Use Pascal to simplify and then open the binomial
- (65) Think in the terms of people on the left!

- (66) First choose the numbers and letters and then simply permute them.
- (67) What is the condition for the diagonal intersecting a lattice point? How many times will that occur?
- (68) Think in terms of the unit digit. We can quickly rule out a lot of numbers.
- (69) It will always reach the end (no matter the values as long as $v \neq 0$). Now try to walk backwards from this...
- (70) Choose one point as reference which is one of the diameter points of the semi-circle, we can lie on either side of the diameter.

C

Appendix

Appendix C: Selected Solutions

D

Appendix

Appendix D: Borrowed Brilliance

As I have mentioned many times, this book is ~~steals~~ borrows brilliance from a lot of people who are frankly much more talented and accomplished than me. It is impotent that we pay tribute to them somewhere.

D.1 Almost every part

- (1) *Art of Problem Solving* website and forums: A lot of the theory, questions and their solutions were sourced from AOPS. If I even just mention the people whose work I directly used, this manuscript would get doubled. I am indebted beyond measure to the entire community.
- (2) *Brilliant.org* website: The website hosts a lot of free pages for different theorems and their exposition. I have taken a lot of inspiration from there along with questions. They could have easily kept these for the paid subscribers, but we can clearly see that they care more about math than money.
- (3) *Math Stack Exchange* forums: It is the reason this book was written in this lifetime. I am a very confused person, if the community there hadn't clarified the dumbest of my doubts, we wouldn't have this book. Also a lot of question and answers were sourced from here.
- (4) *Latex Stack Exchange* forums: It is the reason why this looks like a book and not a jumbled mess of words. It pains me that I can't feasibly mention everyone who helped by their username.

- (5) *Overleaf* latex editor: I didn't install Linux and then a latex editor and compiler and pdf viewer and what not to my computer. I used Overleaf as it was more organized, less cumbersome and easier to share and edit. All the die-hard Linux lovers are gonna lose their marbles. Guys, I don't know how to do the installation and I will not learn until I absolutely need to.
- (6) *Math problem book* template by MAA: Used to give the book the 'book feel'.
- (7) *Random Hints* code by Evan Chan: The reason why we have the elegant random hints. It was the easiest one to use within overleaf.

D.2 Introductory Problems

- (1) *Mathematical Circles: The Russian Experience* book by Dmitrii Vladimirovich Fomin, Ilia Itenberg, and S. Genkin: A classic book with a number of good questions. A must read for all primary school teachers.
- (2) *The USSR Olympiad problem book* Book by D. O. Shkliarskii: This book has a lot of interesting questions. Which require nothing more than the mind to solve. A clear reason why the Russian Olympiads are so fabled
- (3) *Friendship over Tea* video and problem by Arvind Gupta: I had decided while writing this book that I'll include Arvind Gupta sir at least once. He is such a legend. The way he has brought science and design education into the poorest parts of the world using toys from trash is commendable. Check him out, you'll not regret it. His Ted Talk was ranked as 2nd in the 5 favorite education talks.
- (4) *The Green Eyed Dragon and Other Mathematical Monsters* by David Morrin: A very fun book. I have long said to people that once a child turns of listening age, read a single problem from this every night and they'll develop the most wonderful sense of math and logic.
- (5) Counterfeit Coin Riddle video by Jennifer Lu(Ted-Ed): Ted-Ed creates these beautiful animation videos explaining science concepts. They have entire series on world mythology, problems and basics of coding. In hostel, sometimes late at night, a bunch of me and my friends used to sit and see Ted-ed all night. We are all quite successful in math Olympiads, and ones watching something else all night, not quite...
- (6) The OTIS Excerpts book by Evan Chen: Although it is borderline promotional material for his paid course, the book is quite educational.

D.3 Permutations and Combinations

- (1) *Mastering AMC 10/12* by Sohil Rathee: A lot of the AMC 10 and 12 questions were sourced from the book. This is especially true for the part 1 of combinatorics. Sohil Rathi is an angel for going through the long history of AMC and painstakingly choosing the questions, and then putting it out for free.
- (2) *Murderous Math: The Perfect Sausage and other fundamental formulas* book by Kjartan Poskitt: The murderous math series was some of first the non-academic math I had studied. A major influence on how I present math.
- (3) *Introduction to Counting and Probability* handout by David Altizio: This handout in 2013-14 Math League gave me the idea and quite a few of the problems for The Guessing Game.
- (4) *Permutation and Combination for IOQM 2023* lecture series by Abhay Mahajan(Vedantu Olympiad School): Some of the best lectures on combinatorics I have watched. No doubt would be much more popular if they were in English, unfortunately, they are in Hindi, so only fraction of you can enjoy them.
- (5) *Yale Putnam Handouts* by Pat Devlin: The long list handouts made by Pat for the Yale Putnam students was of enormous use in sourcing some of the more difficult problems.
- (6) *Stanford Putnam Handouts* by Ravi Vakil: A legend of the math community, Ravi Vakil's handouts have been used all through the book, including in these for the 9 star problems.

D.4 Down The Rabbit Hole

- (1) *Recursion in AIME* handout by Dylan Yu et al.: This handout, which is part of the Euclid's Orchid handouts, was a source for lot of the recursion theory and problems.
- (2) *Recurrence Relations* lecture by Prashant Jain: A great introductory video with a lot of classic examples. Sadly, its in Hindi, making it less accessible.
- (3) *Reccurence for INMO basics* lecture by Abhay Mahaajan: Abhay sir's question picking genius shines here. Every question is slightly harder and before you realize we are from AMC to IMO.
- (4) *IOQM 2022 practice sessions PnC* lecture by Prashant Jain: Prashant sir had introduced me to Catlan numbers here. The example over there is from Prashant sir's class.

- (5) *Bijections* by Yufei Zhao: The Catlan numbers were expanded upon using Yufai's handout.
- (6) *Counting in Two Ways* handout by Yufei Zhao: The idea of incidence matrix was completely taken from here.
- (7) *Introduction to Graph Theory* handout and powerpoint by Irene Lo: This 2019 Berkeley Math Circle handout was the basis for majority of the graph theory chapter.
- (8) *Olympiad Graph Theory* handout by Adam Kelly: This handout was mainly used for the questions in the graph theory chapter.

D.5 Algebra

- (1) *Polynomials in AIME* handout by Dylan Lu et al.: Another great handout in the Euclid's orchid collection. Worth its weight in gold.
- (2) *Sequences and Series in the AMC and AIME* handout by Dylan Lu et al.: Also from the Euclid's orchid collection. Dylan and his gang have really great material on algebra.
- (3) *Series and Sequences* by David Altizio: Used mainly as a question source.
- (4) *A Brief Introduction to Olympiad Inequalities* handout by Evan Chen: This was the main reference for both of the inequalities chapters
- (5) *Inequalities* handout by Ananth Shyamal, Divya Shyamal, Kevin Yang, and Reece Yang: This was a handout for the Iowa City Math Circle. If any of the readers is in Iowa, I recommend visiting these peeps.
- (6) *Inequalities* handout by Dimitar Grantcharov: The handout made for Berkeley Math Circle while lacking in theory has a bunch of great questions.

D.6 The Red Pill

- (1) *This Is the Calculus They Won't Teach You* video by A Well Rested Dog: This YouTube video made during the first Summer of Math Exposition talks about the history of calculus and was referenced as source if historical context.
- (2) *The Cartoon Guide to Calculus* book by Larry Gonick: This book was what I had used to study calculus and is my first recommendation to those learning it. A lot of the graphs and analogies were taken from this book.

- (3) *Limits and Continuity of Function* Lecture by Prashant Jain: This lecture, part of Bounce back series on Unacademy Atoms YouTube channel, was binged by me on a bus trip home. I have been able to solve some of the most difficult questions of limits since then.
- (4) *Limits* lecture by Abhay Mahajan: These lectures (on the Vedantu JEE made EJEE channel) were the source of many of the questions
- (5) *Differentiation and Continuity* lecture by Abhay Mahajan: Again, used mainly for questions.
- (6) *Indefinite Integration* lecture by Abhay Mahajan: These lectures were the basis of most of the integration by substitution in the integration chapter.
- (7) *DI method* videos by Steve Chow (Blackpen Redpen): The DI Method was first introduced to me by Steve Chow. His other videos on competition math, calculus and math for fun are just a delight to watch.
- (8) *Differential Equations* lecture notes by Nikenasih Binatari of State University of Yogyakarta: While the notes were for the semester course in differential equations, the first few chapters were directly referenced for the section.
- (9) *Matrices and Determinants* lecture by Abhay Mahajan: These lectures were the basis for most of the linear algebra chapter.
- (10) *Lagrange Multipliers* videos by Khan Academy: The basis for the Lagrange multipliers section. There explanation was the simplest to understand and most straightforward, which allowed me to easily integrate it into the book.
- (11) *Lagrange Multipliers Done Correctly* handout by Evan Chen: Most of the Lagrange inequalities were sourced from this handout.
- (12) *Sum Uses of Calculus* by David Altizio: The basis of the summation section. Parts of it were removed as they were too complex, but its overall a great handout.

D.7 Number Theory

- (1) *Olympiad Number Theory Through Challenging Problems* book by Justin Stevens: One of the most used pieces of reference for the entire number theory part. Great explanation, even better questions. Stevens is doing all of a favour by making the text available for free.

- (2) *Mordern Olympiad Number Theory* book by Aditya Khurmi: Another major reference for the Number Theory part. Covers some of the most complex number theory concepts in a digestible manner. Khurmi has given a gift by making this book free on AOPS. The book is so detailed and so beautiful that I had to literally fight myself for every complex concept I wanted to add.
- (3) *A Decade of the Berkeley Math Circle Volume I* book by Zvezdelina Stankova and Tom Rike: The session 4 of this book was the inspiration for the name and the introductory passage for the first chapter of number theory. The general format of the number theory part is also inspired from here.
- (4) *Prime Numbers, Factors, and Division Tricks* handout by Linda Green: While more of her handouts occur in The Numbers Awaken, all of them have very creative examples and increase the question level gradually. Probably the best BMC handouts on number theory.
- (5) *Introduction to Modular Arithmetic* by David Altzio: It is so strange that David either gives me very complex questions or introductory questions. This one was had introductory questions.
- (6) *Bases Part I-II* problem set by Pratima Karpe: The source of almost all of the base questions.
- (7) *Putnam Problem Solving Seminar: Number Theory* handouts by Ravi Vakil: These handouts by Ravi Vakil from the years 2002, 2003, 2004, 2006 was the source of the Putnam or Putnam adjacent problems.
- (8) *Functional equations* handout by Maxim Li: The only functional equation handout which I was able to understand in the first read. If it were not for this handout, I would myself never have understood functional equations in the first place.
- (9) *Functional Equations & Recurrence Relations* handout by Ted Alper: While I didn't use the Reccurence relations, or the analogy between functional equation and recurrence relations, I did use it for the purpose of getting introductory questions.
- (10) *Functional Equations* handout by Igor Ganichev: This was literally a list of problems. I solved all of them and used the ones I particularly found instructive.
- (11) *Putnam Problem Solving Seminar: Functional Equations* handouts by Vin De Silva, Mark Lucianovic and Ravi Vakil: This was also from the same seminars in the years 2005, 2006. Again, they are source of the Putnam or Putnam adjacent problems.

- (12) *Introduction to Functional Equations* handout by Evan Chen: I really want to know what Evan means by "Introduction" because this was by far the most complex reference. However, it had a lot of good examples and problems.
- (13) *Diophantine Equations* by David Altizio: This one is again somewhat on the easier side. Great for the introductory problems. However, I think David doesn't like NT much.

D.8 The Numbers Awaken

- (1) *Sledgehammers in number theory* by CJ Quines: The main reason the Bazooka chapter exists. Lovely handout!
- (2) *Order's Modulo a Prime* by Evan Chen: While I complained about David's number thoery handouts being easy, I'll admit that Evan's handouts required more focus than boiling water with your minds.

E

Appendix

Appendix E: Problem Sources

This chapter talks more about the various exams we have ~~copied and pasted~~ sourced from. Note, individual authors and books have been not mentioned here. They are in Appendix F.

- AIME - American Invitatitional Math Contest
- ARML - American Regions Mathematics League
- AMC 8 - American Math Contest 8
- AMC 10 - American Math Contest 10
- AMC 12 - American Math Contest 12
- Cayley - Centre for Education in Mathematics and Computing Cayley Math Contest
- China - Chine Team Selection Test
- Canada - Canada Team Selection Test
- Fermat - Centre for Education in Mathematics and Computing Fermat Math Contest
- IOQM - Indian Olympiad Qualifiers for Mathematics
- IMO- International Math Olympiad
- IMOSL - International Math Olympiad Short list

- IrMO - Irish Math Olympiad
- ISI - Indian Statistical Institute Admission test
- Italy - Italy Team Selection Test
- Pascal - Centre for Education in Mathematics and Computing Pascal Math Contest
- PROMYS - Program in Mathematics for Young Scientists application form
- Purple Comet - Purple Comet! Math Meet
- Putnam - William Lowell Putnam Mathematical Competition
- USAMO - United States of America Mathematical Olympiad