

Linear Algebra by Sterling K. Berberian

Frosty

April 11, 2026

Contents

1	Vector Spaces	1
1.1	Motivation (Vectors in 3-space)	1
1.2	$\mathbb{R}^n$ and $\mathbb{C}^n$	2
1.3	Vector Spaces: The Axioms and some Examples	3
1.4	Vector Spaces: First Consequences of the Axioms	4
1.5	Linear Combinations of Vectors	5
1.6	Linear Subspaces	7
2	Linear Mappings	11
2.1	Linear Mappings	11
2.2	Linear Mappings and Linear Subspaces: Kernel and Range	13
2.3	Spaces of Linear Mappings: $\mathcal{L}(V, W)$ and $\mathcal{L}(V)$	14
2.4	Isomorphic Vector Spaces	19
2.5	Equivalence Relations and Quotient Sets	22
2.6	Quotient Vector Spaces	24
2.7	The First Isomorphism Theorem	27
3	Structure of Vector Spaces	29
3.1	Linear Subspace Generated by a Subset	29
3.2	Linear Independence	32
3.3	Linear Independence	35

1 Vector Spaces

1.1 Motivation (Vectors in 3-space)

Problem 1

Given the point $P = (2, -1, 3)$ and the vector $u = [-3, 4, 5]$, find the point Q such that $\overrightarrow{PQ} = u$.

Proof. We require $\vec{u} = [-3, 4, 5] = (q_1 - 2, q_2 - (-1), q_3 - 3)$. Solving componentwise, we see $(q_1, q_2, q_3) = (-1, 3, 8)$. Thus, $Q = (-1, 3, 8)$. ■

Problem 2

Given the points $P(2, -1, 3)$, $Q(3, 4, 1)$, $R(4, -3, 4)$, $S(5, 2, 2)$. True or false (explain:) $PQSR$ is a parallelogram.

Proof. For $PQSR$ to be a parallelogram, we require $\vec{PQ} = \vec{SR}$.

$$\vec{PQ} = [3 - 2, 4 - (-1), 1 - 3] = [1, 5, -2]$$

$$\vec{SR} = [4 - 5, -3 - 2, 4 - 2] = [-1, -5, 2]$$


Since

$$[1, 5, -2] \neq [-1, -5, 2],$$

it follows that $PQSR$ is not a parallelogram. ■

Problem 4

Show that for every triple points P, Q, R , $\vec{PQ} + \vec{QR} = \vec{PR}$. [Method 1: Calculate components. Method 2: Draw a picture.]



1.2 $\mathbb{R}^n$ and $\mathbb{C}^n$

Problem 2

Consider the vectors $u = (2, 1)$, $v = (-5, 3)$, $w = (3, 4)$ in $\mathbb{R}^2$. Do there exist real numbers a, b such that $au + bv = w$? What if $v = (6, 3)$.

Proof. We require $au + bv = w \iff a(2, 1) + b(-5, 3) = (3, 4) \iff (2a, a) + (-5b, 3b) = (3, 4) \iff (2a - 5b, a + 3b) = (3, 4)$. Thus

$$\begin{cases} 2a - 5b = 3 \\ a + 3b = 4 \end{cases}$$

which has the solution $a = \frac{29}{11}$, $b = \frac{5}{11}$. If $v = (6, 3)$, we require

$$a(2, 1) + b(6, 3) = (3, 4) \iff (2a + 6b, a + 3b) = (3, 4).$$

Thus

$$\begin{cases} 2a + 6b = 3 \\ a + 3b = 4 \end{cases}$$

which has no solution. To see this has no solution, note

$$a + 3b = 4 \implies 6b = 8 - 2a.$$

Plugging into the first equation gives

$$2a + (8 - 2a) = 3 \implies 8 = 3,$$

which is a contradiction. ■

Problem 3

Give a detailed proof of (8) of Theorem 1.2.3 (just checking!). [Write out all steps of the proof and give a reason for each step.]

Proof. Let $x = [x_1, x_2, \dots, x_n] \in F^n$ be a vector and let $c, d \in F$ where F is a field. Then

$$(cd)x = [(cd)x_1, (cd)x_2, \dots, (cd)x_n].$$

Since F is a field, scalar multiplication in F is associative thus

$$(cd)x_i = c(dx_i) \quad \text{for each } i = 1, 2, \dots, n.$$

Therefore,

$$(cd)x = [c(dx_1), c(dx_2), \dots, c(dx_n)] = c(dx),$$

Problem 4

In Definition 1.2.1, $n = 1$ is not ruled out. What do the elements of $\mathbb{R}^1$ look like? Describe sums and scalar multiples in $\mathbb{R}^1$.

Proof. The elements of $\mathbb{R}^1$ look like $[x_1]$ where $x_1 \in \mathbb{R}$. Sums and scalar multiples behave identically to the operations in the field $\mathbb{R}$.

1.3 Vectors Spaces: The Axioms some Examples

Problem 1

Let V be a vector space over a field F . Let T be a nonempty set and let $W = \mathcal{F}(T, V)$ be the set of all functions $x : T \rightarrow V$. Show that W can be made into a vector space over F in a natural way. [Hint: Use the definitions in Example 1.3.4 as a guide.]

Proof. For $x, y \in W$, $x = y$ means that $x(t) = y(t)$ for all $t \in T$. If $x, y \in W$ and $c \in F$, define functions $x + y$ and cx by the formulas $(x + y)(t) = x(t) + y(t)$ and $(cx)(t) = cx(t)$ for all $t \in T$. Let θ be the function defined by $\theta(t) = 0$ for all $t \in T$, and for $x \in W$, let $-x$ be the function defined by $(-x)(t) = -x(t)$ for all $t \in T$.

Problem 2

The definition of a vector space (1.3.1) can be formulated as follows. A vector space over F is a nonempty set V together with a pair of mappings $\sigma : V \rightarrow V$ and $\mu : F \times V \rightarrow V$. (σ suggests 'sum' and μ suggests 'multiple') having the following properties $\sigma(x, y) = \sigma(y, x)$ for all $x, y \in V$; $\sigma(\sigma(x, y), z) = \sigma(x, \sigma(y, z))$ for all $x, y, z \in V$ etc. The exercise: Write out the 'etc.' in detail.

Proof. We list the 9 axioms below:

1. $x, y \in V \implies \sigma(x, y) \in V$
2. $x \in V, c \in F \implies \mu(c, x) \in V$
3. $x, y \in V \implies \sigma(x, y) = \sigma(y, x)$
4. $x, y, z \in V \implies \sigma(x, \sigma(y, z)) = \sigma(\sigma(x, y), z)$
5. $\exists 0 \in V$ such that $\sigma(0, v) = v = \sigma(v, 0)$
6. $x \in V \implies -x \in V$ such that $\sigma(x, -x) = 0 = \sigma(-x, x)$
7. $x, y \in V$ and $c \in F \implies \mu(c, \sigma(x, y)) = \sigma(\mu(c, x), \mu(c, y))$
8. $\exists 1 \in F$ such that $x \in V \implies \mu(1, x) = x$
9. $a, b \in F$ and $x \in V \implies \mu(a, \mu(b, x)) = \mu(ab, x)$

Problem 3

Let V be a complex vector space (1.3.2). Show that V is also a real vector space (sums as usual, scalar multiplication restricted to real scalars). These two ways of looking at V may be indicated by writing V_c and V_r .

Proof. Clearly V_r is closed under scalar multiplication and vector addition and is a vector space.

Problem 4

Every real vector space V can be 'embedded' in a complex vector space W in the following way: let $W = V \times V$ be the real vector space constructed as in example 1.3.10 and define multiplication by complex scalars by the formula $(a + bi)(x, y) = (ax - by, bx + ay)$ for $a, b \in \mathbb{R}$ and $(x, y) \in W$. [In particular, $i(x, y) = (-y, x)$. Think of (x, y) as ' $x + iy$ '] Show that W satisfies the axioms for a complex vector space (W is called the complexification of V).

Proof. Below are the 9 vector space axioms.

1. Let $x, y \in W$. Then $x = (v_1, v_2), y = (w_1, w_2) \in V \times V$, and $(v_1, v_2) + (w_1, w_2) = (v_1 + w_1, v_2 + w_2)$. Now, $v_1 + w_1 \in V$ and $v_2 + w_2 \in V$, thus $(v_1 + w_1, v_2 + w_2) \in V \times V$.
2. Let $x \in W$ and $c \in \mathbb{C}$. Then $x = (v_1, v_2) \in V \times V, c = a + bi \in \mathbb{C}$, and $(a + bi)(v_1, v_2) = (av_1 - bv_2, bv_1 + av_2)$. Clearly $av_1 - bv_2 \in V$ and $bv_1 + av_2 \in V$, thus $(av_1 - bv_2, bv_1 + av_2) \in V \times V$.
3. Let $x, y \in W$. Then $x = (v_1, v_2), y = (w_1, w_2) \in V \times V$. Then $x + y = (v_1, v_2) + (w_1, w_2) = (v_1 + w_1, v_2 + w_2) = (w_1 + v_1, w_2 + v_2) = (w_1, w_2) + (v_1, v_2) = y + x$.
4. Let $x, y, z \in W$. Then $x = (v_1, v_2), y = (w_1, w_2), z = (s_1, s_2) \in V \times V$. Then $(x + y) + z = [(v_1, v_2) + (w_1, w_2)] + (s_1, s_2) = (v_1 + w_1, v_2 + w_2) + (s_1, s_2) = ((v_1 + w_1) + s_1, (v_2 + w_2) + s_2) = (v_1 + (w_1 + s_1), v_2 + (w_2 + s_2)) = (v_1, v_2) + (w_1 + s_1, w_2 + s_2) = (v_1, v_2) + [(w_1, w_2) + (s_1, s_2)] = x + (y + z)$.
5. Let $x \in W$. Then $x = (v_1, v_2) \in V \times V$ and $x + 0 = (v_1, v_2) + (0, 0) = (v_1 + 0, v_2 + 0) = (v_1, v_2)$.
6. Let $x \in W$. Then $x = (v_1, v_2) \in V \times V$ and $x + (-x) = (v_1, v_2) + (-v_1, -v_2) = (v_1 - v_1, v_2 - v_2) = (0, 0) = 0 = (-v_1, -v_2) + (v_1, v_2) = (-x) + x$.
7. Let $a + bi \in \mathbb{C}$ and $x, y \in W$. Then $(a + bi)((v_1, v_2) + (w_1, w_2)) = (a + bi)(v_1 + w_1, v_2 + w_2) = (a(v_1 + w_1) - b(v_2 + w_2), b(v_1 + w_1) + a(v_2 + w_2)) = (av_1 - bv_2, bv_1 + av_2) + (aw_1 - bw_2, bw_1 + aw_2) = (a + bi)(v_1, v_2) + (a + bi)(w_1, w_2)$.
8. Let $a + bi, c + di \in \mathbb{C}$ and $x = (v_1, v_2) \in W$. Then $(a + bi)((c + di)(v_1, v_2)) = (a + bi)(cv_1 - dv_2, dv_1 + cv_2) = ((ac - bd)v_1 - (ad + bc)v_2, (bc + ad)v_1 + (bd + ac)v_2) = ((a + bi)(c + di))(v_1, v_2)$.
9. Let $x = (v_1, v_2) \in W$. Then $1 \cdot (v_1, v_2) = (1v_1 - 0v_2, 0v_1 + 1v_2) = (v_1, v_2)$.

1.4 Vector Spaces: First Consequences of the Axioms

Problem 1

In a vector space, ① $x - y = \theta$ if and only if ② $x = y$; ③ $x + y = z$ if and only if ④ $x = z - y$.

Proof. (① $\rightarrow$ ②) Suppose $x - y = \theta \iff x + (-y) = \theta$. It follows that $-y = -x \iff y = x$.

(② $\rightarrow$ ①) Suppose $x = y \iff -x = -y$. Then $x + (-x) = x + (-y) = x - y = \theta$.

(③ $\rightarrow$ ④) Suppose $x + y = z \iff x + y + (-y) = z + (-y) \iff x = z - y$.

(④ $\rightarrow$ ③) Suppose $x = z - y \iff x + y = z - y + y \iff x + y = z$. ■

Problem 2

Let V be a vector space over a field F . If x is a fixed nonzero vector, then the mapping $f : F \rightarrow V$ defined by $f(c) = cx$ is injective. If c is a fixed nonzero scalar, then the mapping $g : V \rightarrow V$ defined by $g(x) = cx$ is bijective.

Proof. Suppose x is a fixed nonzero vector. Let $c, c' \in F$ such that $f(c) = f(c') \iff cx = c'x \iff c = c'$. The last step follows from Corollary 1.4.7 (ii). Thus f is injective. ■

Proof. Suppose c is a fixed nonzero scalar. The inverse function to $g(x) = cx$ is clearly $g^{-1}(x) = \frac{x}{c}$. Thus g is bijective. ■

Problem 3

If V is a vector space and y is a fixed vector, then the mapping $\tau : V \rightarrow V$ defined by $\tau(x) = x + y$ is bijective (it is called translation by the vector y).

Proof. Suppose V is a vector space and y is a fixed vector. The inverse function to $\tau(x) = x + y$ is clearly $\tau^{-1}(x) = x - y$. Thus τ is bijective. ■

Problem 4

If a is a nonzero scalar and b is vector in the space V , then the equation $ax + b = \theta$ has a unique solution x in V .

Proof. Suppose a is a nonzero scalar and b is a vector in V . The equation $ax + b = \theta$ is equivalent to $ax = -b$. Multiplying both sides by a^{-1} gives $a^{-1}(ax) = a^{-1}(-b) \iff (a^{-1}a)x = -a^{-1}b \iff x = -a^{-1}b$. ■

Problem 5

If, in a vector space, θ' is a vector such that $\theta' + x = x$ for every vector x , then $\theta' = \theta$. [Hint: Add $-x$ to both sides of the equation.]

Proof. Suppose in a vector space θ' is a vector such that $\theta' + x = x$. Adding $-x$ to both sides gives $\theta' + x + (-x) = x + (-x) \iff \theta' + \theta = \theta \iff \theta' = \theta$. ■

1.5 Linear Combinations of Vectors

Problem 1

Express the function $f(t) = \cos(t - \pi/3)$ as a linear combination of the functions $\sin t$ and $\cos t$.

Proof. We use the standard trig identity to find $f(t) = \cos(t - \pi/3) = \cos(\pi/3)\cos(t) + \sin(\pi/3)\sin(t)$. ■

Problem 2

Express the function e^t as a linear combination of the hyperbolic functions $\cosh t$ and $\sinh t$.

Proof. We have the standard identity $e^t = \cosh(t) + \sinh(t)$. ■

Problem 3

Convince yourself of the correctness of the following formulas pertaining to linear combinations in a vector space:

1. $c \sum_{i=1}^n x_i = \sum_{i=1}^n cx_i$.
2. $\sum_{i=1}^n a_i x_i + \sum_{i=1}^n b_i x_i = \sum_{i=1}^n (a_i + b_i)x_i$.
3. $\sum_{i=1}^m a_i x_i + \sum_{j=1}^n b_j y_j = \sum_{k=1}^{m+n} c_k z_k$ where $c_k = a_k$ and $z_k = x_k$ for $k = 1, 2, \dots, m$, while $c_k = b_{k-m}$ and $z_k = y_{k-m}$ for $k = m+1, m+2, \dots, m+n$.
- 4.

$$\begin{aligned} \left(\sum_{i=1}^m c_i \right) \left(\sum_{j=1}^n x_j \right) &= \sum_{i=1}^m \left(\sum_{j=1}^n c_i x_j \right) \\ &= \sum_{j=1}^n \left(\sum_{i=1}^m c_i x_j \right) \\ &= \sum_{i,j} c_i x_j \end{aligned}$$

The last expression signifying with sum, with mn terms, of the vectors $c_i x_j$ for all possible combinations of i and j .

Proof. I'm convinced. ■

Problem 5

Show that in a vector space, $x - y$ is a linear combination of x and y .

Proof. Let $\alpha = 1$ and $\beta = -1$ and it follows that $x - y = x + (-y) = \alpha x + \beta y$. ■

Problem 6

True or false (explain): In $\mathbb{R}^3$,

1. the vector $x = (0, 6, 3)$ is a linear combination of $(2, 1, 0)$ and $(-3, 2, 0)$;
2. the vector $x = (2, 1, -2)$ is a linear combination of $(2, 1, 0)$ and $(-3, 2, 0)$.

Proof. We must have

$$(0, 6, 3) = \alpha(2, 1, 0) + \beta(-3, 2, 0) = (2\alpha - 3\beta, \alpha + 2\beta, 0)$$

for $\alpha, \beta \in \mathbb{R}$. We obtain

1. $0 = 2\alpha - 3\beta$
2. $6 = \alpha + 2\beta$
3. $3 = 0$

Since $3 \neq 0$, no solution exists. Now suppose

$$(2, 1, -2) = \alpha(2, 1, 0) + \beta(-3, 2, 0) = (2\alpha - 3\beta, \alpha + 2\beta, 0).$$

We obtain

1. $2 = 2\alpha - 3\beta$
2. $1 = \alpha + 2\beta$
3. $-2 = 0$

Since $-2 \neq 0$, no solution exists. ■

1.6 Linear Subspaces

Problem 1

Let M and N be the subsets of $\mathbb{R}^2$ defined as follows:

$$M = \{(a, 0) \mid a \in \mathbb{R}\}, N = \{(0, b) \mid b \in \mathbb{R}\}$$

Show that M and N are linear subspaces of $\mathbb{R}^2$ and describe the linear subspaces $M \cap N$ and $M + N$.

Proof. First note $\theta = (0, 0) \in M, N$. Let $x, y \in M$ such that $x = (a_1, 0), y = (a_2, 0)$. Then $x + y = (a_1, 0) + (a_2, 0) = (a_1 + a_2, 0) \in M$. Let $c \in \mathbb{R}$. Then $cx = c(a_1, 0) = (ca_1, 0) \in M$. The argument for N is the same. The linear subspace $M + N$ is all of $\mathbb{R}^2$, and the linear subspace $M \cap N$ is $\{(0, 0)\}$. ■

Problem 2

Let $\mathcal{F} = \mathcal{R}$ as in Example 1.6.5, fix a nonempty subset T of $\mathbb{R}$, and let

$$M = \{x \in \mathcal{F} \mid x(t) = 0 \text{ for all } t \in T\}$$

Show that M is a linear subspace of $\mathcal{F}$.

Proof. Let $x, y \in M$. Then for all $t \in T$, $(x + y)(t) = x(t) + y(t) = 0 + 0 = 0$, so $x + y \in M$. Let $x \in M$ and $c \in \mathbb{R}$. Then for all $t \in T$, $(cx)(t) = c \cdot x(t) = c \cdot 0 = 0$, so $cx \in M$. Thus M is a linear subspace of $\mathcal{F}$. ■

Problem 3

With $\mathcal{F} = \mathcal{F}(\mathbb{R})$ as in 1.6.5, let

$$M = \{x \in \mathcal{F} \mid x = 0 \text{ on } (-\infty, 0]\}$$

$$N = \{x \in \mathcal{F} \mid x = 0 \text{ on } (0, +\infty)\}$$

Prove that $M + N = \mathcal{F}$ and $M \cap N = \{\theta\}$.

Proof. Let $f \in \mathcal{F}$. Define

$$x(t) = \begin{cases} f(t), & t > 0 \\ 0, & t \leq 0 \end{cases} \in M, \quad y(t) = \begin{cases} f(t), & t \leq 0 \\ 0, & t > 0 \end{cases} \in N.$$

Then $f = x + y$, so $M + N = \mathcal{F}$. Now, let $z \in M \cap N$. Then $z(t) = 0$ for $t \leq 0$ and $z(t) = 0$ for $t > 0$. Thus $z = \theta$ so $M \cap N = \{\theta\}$. ■

Problem 4

Let M and N be linear subspaces of a vector space V , and let

$$M \cup N = \{x \in V \mid x \in M \text{ or } x \in N\}$$

be the union of M and N . True or false (explain): $M \cup N$ is a linear subspace of V . [If true, give a proof. If false, give a counterexample (an example of V, M, N for which $M \cup N$ is not a linear subspace of V).]

Proof. Let M and N be the subsets of $\mathbb{R}^2$ defined as follows:

$$M = \{(a, 0) \mid a \in \mathbb{R}\}, \quad N = \{(0, b) \mid b \in \mathbb{R}\}.$$

We showed in Problem 1 that M and N are linear subspaces of $\mathbb{R}^2$. But notice that $(1, 0) \in M$ and $(0, 1) \in N$, yet $(1, 0) + (0, 1) = (1, 1) \notin M \cup N$. ■

Problem 6

Let V be a vector space and let A be any subset of V . There exist linear subspaces of V that contain A (for instance, V itself is such a subspace). Let $n = \{N \mid N \text{ is a linear subspace of } V \text{ containing } A\}$. Show that n contains a smallest element. [Try $\cap n$.] Thus, there exists a smallest linear subspace of V containing A .

Proof. Let P be an arbitrary linear subspace in n . Clearly, all elements in $\cap n$ are in P , thus $\cap n \subseteq P$. Now, since all linear subspaces in n contain 0, we have $0 \in \cap n$. Let $x, y \in \cap n$ and $c \in F$, where F is the field of scalars of V . Let V be an arbitrary element in n . Since $x, y \in \cap n$, it follows that $x, y \in V$, thus $x + y \in V$. Therefore, $\cap n$ is a linear subspace of V . ■

Problem 7

Let V be a vector space and let n be any set of linear subspaces of V . Let A be the union of the subspaces in n , that is,

$$A = \cup n = \{x \in V \mid x \in M \text{ for some } M \in n\}$$

Apply Exercise 6 to conclude that there exists a linear subspace of V that contains every $M \in n$.

Proof. Clearly $\cup n \subseteq V$. Let $A = \cup n$. By Problem 6, there exists a smallest linear subspace of V containing A . Since for every $M \in n$, $M \subseteq A$, this linear subspace contains every $M \in n$. ■

Problem 8

A linear subspace M of a vector of a vector space V is closed under subtraction: if $x, y \in M$ then $x - y \in M$.

Proof. Suppose $x, y \in M$, where M is a linear subspace of a vector space V . Let $c = -1$. Then $cy = -1 \cdot y = -y \in M$. Thus $x + (-y) = x - y \in M$. ■

Problem 9

Let V be a vector space and let n be a set of linear subspaces of V , having the following property: if $M, N \in n$ then there exists $P \in n$ such that $M \subset P$. (That is, any two elements of n are contained in a third.) Prove that $\cup n$ is a linear subspace of V .

Proof. Let $x, y \in \cup n$ and let c be a scalar of the vector space V . Then $x \in M$ for some $M \in n$. Since M is a linear subspace, $cx \in M$, thus $cx \in \cup n$. Now $y \in N$ for some $N \in n$. There exists $P \in n$ such that $M \subseteq P$ and $N \subseteq P$. Thus $x, y \in P$. Since P is a linear subspace, $x + y \in P$, thus $x + y \in \cup n$. It follows that $\cup n$ is a linear subspace of V . ■

Problem 11

Let M and N be linear subspaces of V . Prove that the following conditions are equivalent:

1. $M \cap N = \{\theta\}$;
2. if $y \in M, z \in N$ and $y + z = \theta$, then $y = z = \theta$;
3. if $y + z = y' + z'$, where $y, y' \in M$ and $z, z' \in N$, then $y = y'$ and $z = z'$.

Proof. (1 $\rightarrow$ 2) Suppose $M \cap N = \{\theta\}$. Furthermore, suppose $y \in M, z \in N$ and $y + z = \theta$. Then $y = -z \in M$, so $z \in M \cap N$. Since $M \cap N = \{\theta\}$, it follows that $z = \theta$. Then $y = -z = -\theta = \theta$.

(2 $\rightarrow$ 3) Suppose that if $y \in M, z \in N$ and $y + z = \theta$, then $y = z = \theta$. Suppose $y + z = y' + z'$, where $y, y' \in M$ and $z, z' \in N$. Then $(y - y') + (z - z') = \theta$. By 2 it follows that $y - y' = \theta$ and $z - z' = \theta$, so $y = y'$ and $z = z'$.

(3 $\rightarrow$ 1) Suppose that if $y + z = y' + z'$, where $y, y' \in M$ and $z, z' \in N$, then $y = y'$ and $z = z'$. Let $x \in M \cap N$. Then $x \in M$ and $x \in N$. Consider $x + \theta = \theta + x$. By 3 it follows that $x = \theta$ and $\theta = x$. Therefore $M \cap N = \{\theta\}$. ■

Problem 12

Let M and N be linear subspaces of V . If $M + N = V$ and $M \cap N = \{\theta\}$ (cf. Exercise 11), V is said to be the direct sum of M and N , written $V = M \oplus N$. Prove that the following conditions are equivalent:

1. $V = M \oplus N$;
2. for each $x \in V$, there exist unique elements $y \in M$ and $z \in N$ such that $x = y + z$.

Proof. ($\rightarrow$) Suppose $V = M \oplus N$. Let x be an arbitrary element in V . Then there exists $m \in M$ and $n \in N$ such that $x = m + n$. Suppose there also exists $m' \in M$ and $n' \in N$ such that $x = m' + n'$. Then $m + n = m' + n'$. By Problem 11, $m = m'$ and $n = n'$, thus m and n are unique.

($\leftarrow$) Suppose for each $x \in V$, there exist unique elements $y \in M$ and $z \in N$ such that $x = y + z$. Let x be an arbitrary element in V . Clearly $x = y + z$ where $y \in M$ and $z \in N$, thus $x \in M + N$. Let x be an arbitrary element in $M + N$. Thus $x = m + n$ for some $m \in M$ and $n \in N$. Therefore $V = M + N$.

Let $x \in M \cap N$. Then

$$x = x + \theta \quad \text{with } x \in M, \theta \in N,$$

and

$$x = \theta + x \quad \text{with } \theta \in M, x \in N.$$

Thus $x = \theta$. Therefore $M \cap N = \{\theta\}$. ■

Problem 13

Let V be the real vector space of all functions $x : \mathbb{R} \rightarrow \mathbb{R}$ (1.3.4). Call $y \in V$ even if $y(-t) = y(t)$ for all $t \in \mathbb{R}$, and call $z \in V$ odd if $z(-t) = -z(t)$ for all $t \in \mathbb{R}$. Let

$$M = \{y \in V \mid y \text{ is even}\}, \quad N = \{z \in V \mid z \text{ is odd}\}.$$

1. Prove that M and N are linear subspaces of V and that $V = M \oplus N$ in the sense of Exercise 12. [Hint: If $x \in V$ consider the functions of y and z defined by $y(t) = \frac{1}{2}[x(t) + x(-t)]$, $z(t) = \frac{1}{2}[x(t) - x(-t)]$.]
2. What does (i) say for $x(t) = e^t$?
3. What does (i) say for a polynomial function x ?

Proof. Clearly the zero function is in both M and N . Let $x, y \in M$ and let $c \in \mathbb{R}$. Then $x(-t) = x(t)$ and it follows that $cx(-t) = cx(t)$, thus $cx \in M$. Also,

$$(x + y)(-t) = x(-t) + y(-t) = x(t) + y(t) = (x + y)(t),$$

thus $x + y \in M$. Let $a, b \in N$ and let $c \in \mathbb{R}$. Then $a(-t) = -a(t)$ and it follows that $ca(-t) = -ca(t)$, thus $ca \in N$. Also,

$$(a + b)(-t) = a(-t) + b(-t) = -a(t) - b(t) = -(a + b)(t),$$

thus $a + b \in N$. It follows that M and N are linear subspaces of V . ■

Proof. Let $x \in V$ be arbitrary. Define functions y and z by

$$y(t) = \frac{1}{2}[x(t) + x(-t)], \quad z(t) = \frac{1}{2}[x(t) - x(-t)].$$

Notice

$$y(-t) = \frac{1}{2}[x(-t) + x(t)] = y(t),$$

so y is even and thus $y \in M$. Also

$$z(-t) = \frac{1}{2}[x(-t) - x(t)] = -\frac{1}{2}[x(t) - x(-t)] = -z(t),$$

so z is odd and thus $z \in N$. Now, for all $t \in \mathbb{R}$,

$$y(t) + z(t) = \frac{1}{2}[x(t) + x(-t)] + \frac{1}{2}[x(t) - x(-t)] = x(t).$$

Thus $x = y + z$ and therefore $V = M + N$. Suppose $x \in M \cap N$. Then x is both even and odd and

$$x(t) = x(-t) \quad \text{and} \quad x(t) = -x(-t).$$

Thus $x(t) = -x(t)$ for all t , $x(t) = 0$ for all t . Thus $M \cap N = \{\theta\}$. Therefore $V = M \oplus N$. ■

Solution (2): It says that e^t can be written uniquely as the sum of an even function and an odd function.

Solution (3): It says that any polynomial function can be written uniquely as the sum of an even polynomial and an odd polynomial.

Problem 16

Let V and W be vector spaces over the same field F and let $V \times W$ be the product vector space (1.3.11). If M is a linear subspace of V , and N is a linear subspace of W , show that $M \times N$ is a linear subspace of $V \times W$.

Proof. Suppose M is a linear subspace of V and N is a linear subspace of W . Let $x, y \in M \times N$ and let c be a scalar from the field F . Then $x = (m_1, n_1)$ and $y = (m_2, n_2)$ are elements of $M \times N$, and $cx = c(m_1, n_1) = (cm_1, cn_1)$. Now, $cm_1 \in M$ and $cn_1 \in N$, thus $cx \in M \times N$. Also $x + y = (m_1, n_1) + (m_2, n_2) = (m_1 + m_2, n_1 + n_2)$. Since $m_1 + m_2 \in M$ and $n_1 + n_2 \in N$, it follows that $x + y \in M \times N$. Since $(\theta_V, \theta_W) \in M \times N$, it follows that $M \times N$ is a linear subspace of $V \times W$. ■

Problem 17

If $V = V_1 \times V_2$ (1.3.11), $M_1 = \{(x_1, \theta) \mid x_1 \in V_1\}$ and $M_2 = \{(\theta, x_2) \mid x_2 \in V_2\}$, then $V = M_1 + M_2$ and $M_1 \cap M_2 = \{\theta\}$ (where θ stands for the zero vector of all vector spaces in sight). In the terminology of Exercise 12, V is the direct sum of its subspaces in M_1 and M_2 , that is, $V = M_1 \oplus M_2$. (Generalization?)

Definition 1. If $V = V_1 \times V_2 \times \cdots \times V_n$, and

1. $M_1 = \{(x_1, \theta_2, \dots, \theta_n) \mid x_1 \in V_1\}$
2. $M_2 = \{(\theta_1, x_2, \dots, \theta_n) \mid x_2 \in V_2\}$
3. ...
4. $M_n = \{(\theta_1, \theta_2, \dots, x_n) \mid x_n \in V_n\}$

then V is the direct sum of its subspaces $M_1, M_2, \dots, M_n$, or

$$V = M_1 \oplus M_2 \oplus \dots \oplus M_n.$$

Problem 18

Let M and N be linear subspaces of V whose union $M \cup N$ is also a linear subspace. Prove that either $M \subset N$ or $N \subset M$. [Hint: Assume to the contrary that neither of M, N is contained in the other; choose vectors $y \in M, z \in N$ with $y \notin N, z \notin M$ and try to find a home for $y + z$.]

Proof. Let $y \in M$ and $z \in N$ such that $y \notin N$ and $z \notin M$. Then $y \in M \cup N$ and $z \in M \cup N$. Since $M \cup N$ is a linear subspace, $y + z \in M \cup N$. Without loss of generality, suppose $y + z \in M$. Then $-y \in M$ by Problem 8. Since M is closed under addition $y + z + (-y) = z \in M$ which is a contradiction. ■

2 Linear Mappings

2.1 Linear Mappings

Problem 2

Let T be a set, $V = \mathcal{F}(T, \mathbb{R})$ the vector space of all functions $x : T \rightarrow \mathbb{R}$ (Example 1.3.4). Fix a point $t \in T$ and define $f : V \rightarrow \mathbb{R}$ by the formula $f(x) = x(t)$. Then f is a linear form on V .

Proof. Let $x, y \in V$ and $c \in \mathbb{R}$. Then $f(x + y) = (x + y)(t) = x(t) + y(t) = f(x) + f(y)$. and $f(cx) = (cx)(t) = cx(t) = cf(x)$. Thus f is a linear mapping. Since it maps functions in V to scalars in $\mathbb{R}$ it is a linear form. ■

Problem 3

If $T : V \rightarrow W$ is a linear mapping, then $T(x - y) = Tx - Ty$ for all $x, y \in V$.

Proof. Suppose $T : V \rightarrow W$ is a linear mapping. Let x, y be arbitrary elements in V . Then $T(x - y) = T(x + (-y)) = Tx + T(-y) = Tx + (-T(y)) = Tx - Ty$. ■

Problem 4

If V is a vector space, prove that the mapping $T : V \times V \rightarrow V$ defined by $T(x, y) = x - y$ is linear. (It is understood that $V \times V$ has the product vector space structure described in Example 1.3.10.)

Proof. Suppose V is a vector space. Let $(x, y), (a, b)$ be arbitrary elements in $V \times V$. Let $c \in \mathbb{R}$ be a scalar. Then $T((x, y) + (a, b)) = T(x + a, y + b) = (x + a) - (y + b) = (x - y) + (a - b) = T(x, y) + T(a, b)$. Also $T(c(x, y)) = T(cx, cy) = cx - cy = c(x - y) = cT(x, y)$. Thus T is linear. ■

Problem 5

With notations as in Theorem 2.1.3, an element $(a_1, \dots, a_n)$ of F^n such that $T(a_1, \dots, a_n) = \theta$ is called a (linear) relation among the vectors $x_1, \dots, x_n$. For example, suppose that $V = F^2$, $n = 3$ and $x_1 = (2, -3)$, $x_2 = (4, 1)$, $x_3 = (8, 9)$.

1. Find a formula for the linear mapping $F^3 \rightarrow F^2$ defined in Theorem 2.1.3.
2. Show that $(-2, 3, 1)$ is a relation among x_1, x_2, x_3 .

Solution (a):

$$f(a_1, a_2, a_3) = a_1x_1 + a_2x_2 + a_3x_3$$

Solution (b):

$$f(-2, 3, 1) = -2x_1 + 3x_2 - x_3 = -2(2, -3) + 3(4, 1) - (8, 9) = (-4, 6) + (12, 3) - (8, 9) = (8, 9) - (8, 9) = (0, 0) = \theta$$

Problem 6

The proof of Theorem 2.1.2 uses only the additivity of the mapping T . Give a proof using only its homogeneity. [Hint: Corollaries 1.4.3, 1.4.5]

Theorem 1. If $T : V \rightarrow W$ is a linear mapping, then $T\theta = \theta$ and $T(-x) = -(Tx)$ for all $x \in V$.

Proof. Suppose $T : V \rightarrow W$ is a linear mapping. Then $T(\theta) = T(0 \cdot x) = 0 \cdot T(x)$. Then by Corollary 1.4.3, $0 \cdot T(x) = \theta$, hence $T(\theta) = \theta$. Also $T(-x) = T((-1) \cdot x) = (-1)T(x)$. Then by Corollary 1.4.5, $(-1)T(x) = -T(x)$. Thus $T(-x) = -Tx$. ■

Problem 7

If $\mathcal{D}$ is the vector space of real polynomial functions (Example 1.3.5) and $f : \mathcal{D} \rightarrow \mathbb{R}$ is the mapping defined by $f(p) = p'(1)$ (the value of the derivative of p at 1), then f is a linear form on $\mathcal{D}$. What is the geometric meaning of $f(p) = 0$?

Proof. Let $x, y \in \mathcal{D}$ and $c \in \mathbb{R}$. Then $f(x+y) = (x+y)'(1) = x'(1) + y'(1) = f(x) + f(y)$. and $f(cx) = (cx)'(1) = c x'(1) = c f(x)$. It follows that f is a linear form on $\mathcal{D}$. ■

Solution: $f(p) = 0$ is a critical point at the value $x = 1$ of the function p .

Problem 8

Let $S : \mathcal{D} \rightarrow \mathcal{D}$ be the linear mapping of Example 2.1.5. Define $R : \mathcal{D} \rightarrow \mathcal{D}$ by $Rp = Sp + p(5)1$, where $p(5)1$ is the constant function defined by the real number $p(5)$. Then R is linear and $(Rp)' = p$. [So to speak, the constant of integration can be tailor-made for p in a linear fashion.] More generally, look at the mapping $Rp = Sp + f(p)1$, where f is any linear form on $\mathcal{D}$.

Proof. Let $x, y \in \mathcal{D}$ and $c \in \mathbb{R}$. Then $R(x+y) = S(x+y) + (x+y)(5)1 = S(x) + S(y) + x(5)1 + y(5)1 = R(x) + R(y)$. Also $R(cx) = S(cx) + (cx)(5)1 = cS(x) + cx(5)1 = cR(x)$. Thus R is linear.

For the general form notice $R(x+y) = S(x+y) + f(x+y)1 = S(x) + S(y) + f(x)1 + f(y)1 = R(x) + R(y)$. Also $R(cx) = S(cx) + f(cx)1 = cS(x) + cf(x)1 = cR(x)$. ■

2.2 Linear Mappings and Linear Subspaces: Kernel and Range

Problem 2

Let V be a vector space over F , $f : V \rightarrow F$ a linear form on V (Definition 2.1.11). Assume that f is not identically zero and choose a vector y such that $f(y) \neq 0$. Let N be the kernel of f . Prove that every vector x in V may be written $x = z + cy$ with $z \in N$ and $c \in F$, and that z and c are uniquely determined by x . [Hint: If $x \in V$, compute the value of f at the vector $x - [f(x)/f(y)]y$.]

Proof. Suppose $x, y \in V$. Notice $f\left(x - \frac{f(x)}{f(y)}y\right) = f(x) - f\left(\frac{f(x)}{f(y)}y\right) = f(x) - \frac{f(x)}{f(y)}f(y) = f(x) - f(x) = 0$. We've just shown $x - \frac{f(x)}{f(y)}y \in \ker(f)$. Thus $x = \left(x - \frac{f(x)}{f(y)}y\right) + \frac{f(x)}{f(y)}y$. Let $z = x - \frac{f(x)}{f(y)}y$ and $c = \frac{f(x)}{f(y)}$, and we see $x = z + cy$ as required.

Suppose $x = z' + c'y$ with $z' \in \ker(f)$ and $c' \in F$. Then $z + cy = z' + c'y \implies z - z' = (c' - c)y$. Applying f gives $0 = f(z - z') = (c' - c)f(y)$, so $c' - c = 0$ since $f(y) \neq 0$. Thus $c' = c$ and $z' = z$. ■

Problem 3

Let $S : V \rightarrow W$ and $T : V \rightarrow W$ be linear mappings and let $M = \{x \in V \mid Sx \in T(V)\}$. Prove that M is a linear subspace of V .

Proof. Let $x, y \in M$ and let c be a scalar of the vector space V . Since $x, y \in M$, we have $Sx \in T(V)$ and $Sy \in T(V)$. Since $T(V)$ is a linear subspace of W it follows that $S(x + y) = Sx + Sy \in T(V)$. Thus $x + y \in M$. Similarly $S(cx) = cS(x) \in T(V)$ thus $cx \in M$. Therefore M is a linear subspace of V . ■

Problem 4

If $T : V \times V \rightarrow V$ is the linear mapping $T(x, y) = x - y$ (2.1, Exercise 4), determine the kernel and range of T .

Solution: The kernel is $(x, x) \in V \times V$ and the range is V .

Problem 5

Let V be a real vector space, W its complexification (1.3, Exercise 4); write W_R for W regarded as a real vector space (1.3, Exercise 3). For $(x, y) \in W$, we have

$$(x, y) = (x, \theta) + (\theta, y) = (x, \theta) + i(y, \theta).$$

Prove that $x \mapsto (x, \theta)$ is an injective mapping $V \rightarrow W_R$. [Identifying $x \in V$ with $(x, \theta) \in W$, one can suggestively write $W = V + iV$; so to speak, V is the 'real part' of its complexification.]

Proof.

$$x(x_1) = x(x_2) \iff (x_1, 0) = (x_2, 0) \iff x_1 = x_2$$

Problem 6

Let $\mathcal{D}$ be the vector space of real polynomial functions (1.3.5) and let $T : \mathcal{D} \rightarrow \mathcal{D}$ be the linear mapping defined by $Tp = p - p'$, where p' is the derivative of p . Prove that T is injective.

Proof. Suppose $x, y \in \mathcal{D}$ such that $T(x) = T(y)$. Then $T(x) = T(y) \iff x - x' = y - y' \iff (x - y) - (x' - y') = 0$. Let $q = x - y$ then $q - q' = 0 \iff q = q'$. Thus $q = 0$ and it follows that $x = y$ and $x' = y'$. ■

2.3 Spaces of Linear Mappings: $\mathcal{L}(V, W)$ and $\mathcal{L}(V)$

Problem 1

Complete the details in the proof of Theorem 2.3.11.

Theorem 2. If V and W are vector spaces over F , then $\mathcal{L}(V, W)$ is also a vector space over F for the operations defined in 2.3.9.

Proof. Let $S, T \in \mathcal{L}(V, W)$ and let $a, b \in F$. For all $x \in V$, $(a(S + T))(x) = a((S + T)(x)) = a(S(x) + T(x)) = aS(x) + aT(x) = (aS + aT)(x)$. For all $x \in V$, $((a + b)T)(x) = (a + b)T(x) = aT(x) + bT(x) = (aT + bT)(x)$. For all $x \in V$, $((ab)T)(x) = abT(x) = a(bT(x)) = (a(bT))(x)$. For all $x \in V$, $(1T)(x) = 1 \cdot T(x) = T(x)$. ■

Problem 2

Let V be a vector space. If $R, S, T \in \mathcal{L}(V)$ and c is a scalar, the following equalities are true.

1. $(RS)T = R(ST)$;
2. $(R + S)T = RT + ST$;
3. $R(S + T) = RS + RT$;
4. $(cS)T = c(ST) = S(cT)$;
5. $TI = T = IT$; (T is the identity mapping)

Proof. For all $x \in V$.

1. $(RS)T(x) = R(S(T(x))) = R(ST(x)) = (R(ST))(x)$.
2. $(R + S)T(x) = (R + S)(T(x)) = R(T(x)) + S(T(x)) = RT(x) + ST(x) = (RT + ST)(x)$.
3. $R(S + T)(x) = R(S(x) + T(x)) = R(S(x)) + R(T(x)) = RS(x) + RT(x) = (RS + RT)(x)$.
4. $(cS)T(x) = cS(T(x)) = c(ST(x)) = (c(ST))(x) = S(cT)(x)$.
5. $TI(x) = T(I(x)) = T(x)$ and $IT(x) = I(T(x)) = T(x)$.

Problem 4

If $T : V \rightarrow W$ is a linear mapping and g is a linear form on W , show that the formula $f(x) = g(Tx)$ defines a linear form f on V . [Shortcut: $f = gT = g \circ T$] The formula $T'g = f$ defines a mapping $T' : \mathcal{L}(W, F) \rightarrow \mathcal{L}(V, F)$, where F is the field of scalars. Show that T' is linear. [T' is called the transpose (or 'adjoint' of T).]

Proof. By Theorem 2.3.13 it directly follows that f is linear.

Let $x, y \in \mathcal{L}(W, F)$ and c be a scalar in F . Then for all $v \in V$, $T'(x + y)(v) = (x + y)(T(v)) = x(T(v)) + y(T(v)) = T'(x)(v) + T'(y)(v)$. Similarly, $T'(cx)(v) = (cx)(T(v)) = c x(T(v)) = c T'(x)(v)$. Therefore, T' is linear. ■

Problem 5

If V is a vector space over F , the vector space $\mathcal{L}(V, F)$ of linear forms on V is called the dual space of V and is denoted V' . Thus, the correspondence $T \mapsto T'$ described in Exercise 4 defines a mapping $\mathcal{L}(V, W) \rightarrow \mathcal{L}(W', V')$. Show that this mapping is linear, that is, $(S + T)' = S' + T'$ and $(cT)' = cT'$ for all S, T in

$\mathcal{L}(V, W)$ and all scalars c .

Proof. Let $S, T \in \mathcal{L}(V, W)$ and let c be a scalar. Let v be a vector in V . Then $(S + T)'g(v) = g(S + T)(v) = g(S(v) + T(v)) = gS(v) + gT(v) = S'g(v) + T'g(v)$. Similarly $(cT)'g(v) = g(cT(v)) = cgT(v) = c(T'g)(v)$. Thus the correspondence $T \mapsto T'$ is linear. ■

Problem 6

With notations as in Exercise 4, prove that if T is surjective then T' is injective; in other words, show that if $T(V) = W$ then $T'g = 0 \implies g = 0$.

Proof. Suppose $T(V) = W$ and $T'g = 0$. Let $v \in V$ then $(T'g)(v) = g(T(v)) = 0$ for all $v \in V$. Thus $g = 0$ and T' is injective. ■

Problem 7

As in Example 2.1.4, let $V = \mathcal{D}$ be the real vector space of all polynomial functions $p : \mathbb{R} \rightarrow \mathbb{R}$, $D : V \rightarrow V$ the linear mapping defined by $Dp = p'$ (the derivative of p); let $D' : V' \rightarrow V'$ be the transpose of D as defined in Exercise 4. (Caution: The prime is being used with three different meanings.)

For each $t \in \mathbb{R}$ let ρ_t be the linear form on V defined by $\rho_t(p) = p(t)$. Let $[a, b]$ be a closed interval on $\mathbb{R}$ and let ρ be the linear form on V defined by

$$\rho(p) = \int_a^b p(t)dt$$

Prove : $D'\rho = \rho_b - \rho_a$. [Hint: Fundamental Theorem of Calculus.]

Proof. Let $p \in V$. Notice $(D'\rho)(p) = \rho(Dp) = \int_a^b p'(t)dt = p(b) - p(a) = \rho_b(p) - \rho_a(p)$. ■

Problem 9

Let U, V, W be vector spaces over F . Prove:

1. For fixed $T \in \mathcal{L}(U, V)$, $\psi = S \mapsto ST$ is a linear mapping $\mathcal{L}(V, W) \rightarrow \mathcal{L}(U, W)$ (see Figure 8 2.3.12).
2. For fixed $S \in \mathcal{L}(V, W)$, $\phi = T \mapsto ST$ is a mapping $\mathcal{L}(U, V) \rightarrow \mathcal{L}(U, W)$.

Proof. Let $x, y \in \mathcal{L}(V, W)$ and $v \in U$. Let c be a scalar. Then $(\psi(x + y))(v) = ((x + y)T)(v) = (x + y)(T(v)) = xT(v) + yT(v) = \psi(x)(v) + \psi(y)(v)$. Also $(\psi(cx))(v) = ((cx)T)(v) = (cx)(T(v)) = c(xT(v)) = c\psi(x)(v)$. Thus ψ is a linear mapping. ■

Proof. Let $x, y \in \mathcal{L}(U, V)$ and $v \in U$. Let c be a scalar. Then $(\phi(x + y))(v) = S(x + y)(v) = S(x(v) + y(v)) = S(x(v)) + S(y(v)) = \phi(x)(v) + \phi(y)(v)$. Also $(\phi(cx))(v) = S(cx(v)) = cS(x(v)) = c\phi(x)(v)$. Thus ϕ is a linear mapping. ■

Problem 10

If $T : U \rightarrow V$ and $S : V \rightarrow W$ are linear mappings, prove that $\text{Im}(ST) \subset \text{Im}S$ and $\text{Ker}(ST) \supset \text{Ker}T$.

Proof. Clearly T can only restrict the domain of S thus $\text{Im}(ST) \subset \text{Im}S$. Clearly any x in $\text{Ker}T$ will be in $\text{Ker}(ST)$ since $S(0) = 0$. ■

Problem 11

Let $T : V \rightarrow V$ be a linear mapping such that $\text{Im}T \subset \text{Ker}(T - I)$, where I is the identity mapping (2.3.3). Prove that $T^2 = T$. [Recall $T^2 = TT$.]

Proof. First note that $\text{Ker}(T - I) = \{v \in V \mid (T - I)(v) = 0\}$. For any $v \in V$, $(T - I)(v) = 0 \iff T(v) - I(v) = 0 \iff T(v) - v = 0 \iff T(v) = v$. Let $v \in V$ be arbitrary. Then $T(v) \in \text{Im}T$. Since $\text{Im}T \subset \text{Ker}(T - I)$, it follows that $T(v) \in \text{Ker}(T - I)$, and thus $T(T(v)) = T(v)$. Thus $T^2 = T$. ■

Problem 12

Suppose $V = M \oplus N$ in the sense of 1.6, Exercise 12. For each $x \in V$, let $x = y + z$ be its unique decomposition with $y \in M, z \in N$ and define $Px = y, Qx = z$. Prove that $P, Q \in \mathcal{L}(V)$, $P^2 = P, Q^2 = Q, P + Q = I$ and $PQ = QP = 0$.

Proof. Let $a_1, a_2 \in M$ and $b_1, b_2 \in N$. Let $x = a_1 + b_1, y = a_2 + b_2 \in V$. Then $P(x + y) = P((a_1 + a_2) + (b_1 + b_2)) = a_1 + a_2 = Px + Py$. Also $P(cx) = P(c(a_1 + b_1)) = P(ca_1 + cb_1) = ca_1 = c(Px)$. Thus $P \in \mathcal{L}(V)$. The argument that $Q \in \mathcal{L}(V)$ is similar. ■

Proof. Simply note that $\text{Im}P \subset \text{Ker}(P - I)$ thus $P^2 = P$. The argument that $Q^2 = Q$ is similar. ■

Proof. Let $a \in M$ and $b \in N$. Let $x = a + b$. Then $(P + Q)(x) = P(x) + Q(x) = a + b = x$. Thus $P + Q = I$. ■

Proof. Let $a \in M$ and $b \in N$. Let $x = a + b$. Then $(PQ)(x) = P(Q(x)) = P(Q(a + b)) = P(0 + Q(b)) = 0$. The argument for QP is similar. ■

Problem 13

Let $T : V \rightarrow V$ be a linear mapping such that $T^2 = T$, and let $M = \text{Im}T$ and $N = \text{Ker}T$. Prove that $M = \{x \in V \mid Tx = x\} = \text{Ker}(T - I)$ and that $V = M \oplus N$ in the sense of 1.6, Exercise 12.

Proof. Notice $x \in M \iff Tx = x \iff Tx - x = 0 \iff Tx - Ix = 0 \iff (T - I)(x) = 0 \iff x \in \text{Ker}(T - I)$. Let $x \in V$. Let $y = Tx \in M$ and $z = x - Tx \in N$. Notice

$$T(z) = T(x - Tx) = Tx - T^2x = Tx - Tx = 0$$

Thus $z \in \text{Ker}T = N$. Therefore $x = y + z \in M + N$. Suppose $x = y_1 + z_1 = y_2 + z_2$ with $y_i \in M, z_i \in N$. Then

$$y_1 - y_2 = z_2 - z_1 \in M \cap N.$$

But then $M \cap N = \{0\}$ because suppose $v \in M \cap N$, then $Tv = v$ and $Tv = 0$, so $v = 0$. Thus $V = M \oplus N$. ■

Problem 14

Find $S, T \in \mathcal{L}(\mathbb{R}^2)$ such that $ST \neq TS$.

Proof. Consider the linear maps $S, T \in \mathcal{L}(\mathbb{R}^2)$

$$S(x, y) = (y, 0), \quad T(x, y) = (0, x).$$

Then for any $(x, y) \in \mathbb{R}^2$

$$ST(x, y) = S(T(x, y)) = S(0, x) = (x, 0),$$

$$TS(x, y) = T(S(x, y)) = T(y, 0) = (0, y).$$

Thus $ST \neq TS$. ■

Problem 15

Let $T : U \rightarrow V$ and $S : V \rightarrow W$ be linear mappings. Prove

1. If S is injective, then $\text{Ker}(ST) = \text{Ker}(T)$.
2. If T is surjective, then $\text{Im}(ST) = \text{Im}S$.

Proof. Suppose S is injective. Let x be an arbitrary element in U . Suppose $x \in \text{Ker}(T)$ thus $T(x) = 0$. Then $(ST)(x) = S(T(x)) = S(0) = 0$. Thus $x \in \text{Ker}(ST)$. Suppose $x \in \text{Ker}(ST)$ thus $(ST)(x) = 0$. Now $S(0) = 0$ and since $S(T(x)) = 0$ and S is injective, $T(x) = 0$ thus $x \in \text{Ker}T$. It follows that $\text{Ker}(ST) = \text{Ker}(T)$. ■

Proof. Suppose T is surjective. It follows that T maps onto all elements in the domain of S . Therefore $\text{Im}(ST) = \text{Im}S$. ■

Problem 16

Let V be a real or complex vector space and let $t \in \mathcal{V}$ be such that $T^2 = I$. Define

$$M = \{x \in V \mid Tx = x\}, \quad N = \{x \in V \mid Tx = -x\}$$

Prove that M and N are linear subspaces of V such that $V = M \oplus N$. [Hint: For every vector x , $x = \frac{1}{2}(x + Tx) + \frac{1}{2}(x - Tx)$.]

Proof. Let $x \in V$ and consider $x = \frac{1}{2}(x + Tx) + \frac{1}{2}(x - Tx)$. Let $m = \frac{1}{2}(x + Tx)$ and $n = \frac{1}{2}(x - Tx)$. Then $Tm = \frac{1}{2}(Tx + T^2x) = \frac{1}{2}(Tx + x) = m$ so $m \in M$, and $Tn = \frac{1}{2}(Tx - T^2x) = \frac{1}{2}(Tx - x) = -n$ so $n \in N$. Thus $x = m + n \in M + N$.

Suppose $v \in M \cap N$. Then $Tv = v$ and $Tv = -v$, which implies $v = 0$. Thus $M \cap N = \{0\}$. It follows that $V = M \oplus N$. ■

Problem 17

Let $V = \mathcal{F}(\mathbb{R})$ be the real vector space of all functions $x : \mathbb{R} \rightarrow \mathbb{R}$ (1.3.4). For $x \in V$ define $Tx \in V$ by the formula $(Tx)(t) = x(-t)$. Analyze T in the light of Exercise 16. [Remember 1.6, Exercise 13]

Solution: The set of functions for which $x(t) = x(-t)$, i.e., the even functions, are in M . The functions for which $x(t) = -x(-t)$, i.e., the odd functions, are in N . Every function $x \in V$ can be expressed as $x = \frac{1}{2}(x + Tx) + \frac{1}{2}(x - Tx) \in M \oplus N$.

Problem 18

Let $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$, prove that $(ST)' = T'S'$. (cf. Exercise 4).

Proof. Let $f \in W'$ be a linear form on W . Then $(ST)'(f) = f \circ (ST) = (f \circ S) \circ T = T'(S'(f))$. ■

Problem 19

Let $S, T \in \mathcal{L}(V, W)$ and let $M = \{x \in V \mid Sx = Tx\}$. Prove that M is a linear subspace of V . [Hint: Consider $\text{Ker}(S - T)$.]

Proof. Note that $M = \{x \in V \mid Sx - Tx = 0\} = \{x \in V \mid (S - T)x = 0\} = \text{ker}(S - T)$. Since the kernel of a linear map is always a linear subspace, M is a linear subspace of V . ■

Problem 20

If $T : V \rightarrow W, S : W \rightarrow V$ are linear mappings such that $ST = I$, prove that $\text{Ker} T = \{0\}$ and $\text{Im} S = V$.

Proof. Let $x \in \text{Ker} T$ such that $Tx = 0$. Then $STx = Tx = 0$. Since $ST = I$, we have $x = 0$ thus $\text{Ker} T = \{0\}$.

Next, let $v \in V$. Then $v = Iv = STv = S(Tv)$. Thus $v \in \text{Im} S$. Therefore $\text{Im} S = V$. ■

Problem 21

If $V = \{\theta\}$ or $W = \{\theta\}$ then $\mathcal{L}(V, W) = \{0\}$. If $V \neq \{\theta\}$ then $I_v \neq 0$.

Proof. Suppose $V = \{\theta\}$ or $W = \{\theta\}$. Clearly there is only one linear mapping between V and W which is for all $x \in V, T(x) = 0 \in W$.

Suppose $V \neq \{\theta\}$. V has at least two elements, one of which maps to 0. The other must map to a nonzero element under I_V . ■

Problem 22

A lightning proof of Theorem 2.3.11 can be based on an earlier exercise: Since $0 \in \mathcal{L}(V, W)$, Lemma 2.3.10 shows that $\mathcal{V}, \mathcal{W}$ is a linear subspace of the vector space $\mathcal{F}(V, W)$ defined as in 1.3, Exercise 1, therefore $\mathcal{L}(V, W)$ is also a vector space (1.6.2).

Solution: OK.

Problem 23

If $T : U \rightarrow V$ and $S : V \rightarrow W$ are linear mappings, prove that $\text{Ker}(ST) = T^{-1}(\text{Ker} S)$.

Proof. Let $x \in U$. Then

$$x \in \text{Ker}(ST) \iff STx = 0 \iff S(Tx) = 0 \iff Tx \in \text{Ker} S \iff x \in T^{-1}(\text{Ker} S).$$

Therefore $\text{Ker}(ST) = T^{-1}(\text{Ker} S)$. ■

Problem 24

Let $V = \mathcal{D}$ be the vector space of real polynomial functions, $D : V \rightarrow V$ the differentiation mapping $Dp = p'$ (2.1.4). Let u be the monomial $u(t) = t$ and define another linear mapping $M : V \rightarrow V$ by the formula $Mp = up$. (So to speak, M is multiplication by t .) Prove that $DM - MD = I$ (the identity mapping). [Hint: The claim is that $(up)' - up' = p$. Remember the 'product rule'!]

Proof. By the product rule,

$$(up)' = u'p + up'.$$

Since $u(t) = t$ we know $u' = 1$. Thus

$$(up)' = p + up'.$$

Problem 25

Let $D : \mathcal{D} \rightarrow \mathcal{D}$ be the differentiating map of Exercise 24 and let $S : \mathcal{D} \rightarrow \mathcal{D}$ be the linear mapping such that $(Sp)' = p$ and $(Sp)(0) = 0$ (2.1.5). Prove:

1. If $R : \mathcal{D} \rightarrow \mathcal{D}$ is any linear mapping such that $(Rp)' = p$ for all $p \in \mathcal{D}$, then $DR = I$.
2. If $T : \mathcal{D} \rightarrow \mathcal{D}$ is a linear mapping such that $DT = 0$, then there exists a linear form f on $\mathcal{D}$ such that (in the notation of Exercise 2.3.4) $T = f \otimes 1$, where 1 is the constant function 1 . [Hint: $\text{Im} T \subset \text{Ker} D$.]
3. With R as in (i), $R = S + f \otimes 1$ for a suitable linear form f on $\mathcal{D}$. [Hint: Consider $T = R - S$.] Remember 2.1, Exercise 8?

Proof. Suppose $R : \mathcal{D} \rightarrow \mathcal{D}$ is a linear mapping such that $(Rp)' = p$ for all $p \in \mathcal{D}$. Let $p \in \mathcal{D}$. Then $(DR)(p) = D(Rp) = (Rp)' = p$. Thus $DR = I$. ■

Proof. If $T : \mathcal{D} \rightarrow \mathcal{D}$ is a linear mapping such that $DT = 0$, then for all $p \in \mathcal{D}$, Tp is a constant function. Thus $T = T \otimes 1$, where 1 is the constant function 1 . ■

Proof. Let $R : \mathcal{D} \rightarrow \mathcal{D}$ and let $T = R - S$. Then for any $p \in \mathcal{D}$, $D(Tp) = D((R - S)p) = DR(p) - DS(p) = I(p) - I(p) = 0$. Thus $DT = 0$. By part (ii), there exists a linear form f on $\mathcal{D}$ such that $T = f \otimes 1$. Therefore $R = S + T = S + f \otimes 1$. ■

Problem 26

Let $S, T \in \mathcal{L}(V)$. If $ST - I$ is injective then so is $TS - I$. [Hint: $S(TS - I) = (ST - I)S$.]

Proof. Suppose $ST - I$ is injective. Thus $\ker(ST - I) = \{0\}$. Suppose $x \in V$ such that $(TS - I)(x) = 0$. Then applying S we have $S(TS - I)(x) = 0$. By the hint we have $((ST - I)S)(x) = 0 \iff (ST - I)(S(x)) = 0$. Since $ST - I$ is injective we have $S(x) = 0$. Then $(TS - I)(x) = TS(x) - x = T0 - x = -x = 0$. Thus $x = 0$. ■

2.4 Isomorphic Vector Spaces

Problem 2

Regard $\mathbb{C}^n$ as a real vector space in the natural way (sums as usual, multiplication only by real scalars); then $\mathbb{C}^n \cong \mathbb{R}^{2n}$ as real vector spaces.

Proof. Let $r_1, r_2, \dots, r_{2n-1}, r_{2n} \in \mathbb{R}$. Consider the mapping

$$(r_1, r_2, \dots, r_{2n-1}, r_{2n}) \mapsto (r_1 + r_2i, \dots, r_{2n-1} + r_{2n}i).$$

We know that the map $(r_1, r_2) \mapsto r_1 + r_2i$ is linear and bijective over $\mathbb{R}$. Our mapping is simply an extension of this map to n components and is therefore linear and bijective. ■

Problem 3

Let X be a set, V a vector space, $f : X \rightarrow V$ a bijection. There exists a unique vector space structure on X for which f is a linear mapping (hence a vector space isomorphism). [Hint: Define sums and scalar multiples in X by the formulas $x + y = f^{-1}(f(x) + f(y))$, $cx = f^{-1}(cf(x))$. This trick is called 'transport of structure'.]

Proof. Clearly X is a vector space, since after applying f each vector space axiom becomes the corresponding axiom in V , and injectivity of f implies the axioms hold in X . ■

Problem 4

Let V be the set of n -ples $x = (x_1, \dots, x_n)$ of real numbers > 0 , that is,

$$V = \{(x_1, \dots, x_n) \mid x_i \in (0, +\infty) \text{ for } i = 1, \dots, n\}.$$

For $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n)$ in V and $c \in \mathbb{R}$, define

$$x \oplus y = (x_1 y_1, \dots, x_n y_n), c \cdot x = (x_1^c, \dots, x_n^c),$$

(here $x_i y_i$ and x_i^c are the usual product and power), Define a mapping $T : V \rightarrow \mathbb{R}^n$ by the formula $T(x_1, \dots, x_n) = (\log x_1, \dots, \log x_n)$.

1. Show that $T(x \oplus y) = Tx + Ty$ and $T(c \cdot x) = c(Tx)$ for all x, y in V and all $c \in \mathbb{R}$.
2. True or false (explain): V is a real vector space for the operations $\cdot$.

Proof. Let $x, y \in V$ and c be a scalar. Then

$$\begin{aligned} T(x \oplus y) &= T((x_1 y_1, \dots, x_n y_n)) = (\log(x_1 y_1), \dots, \log(x_n y_n)) = (\log x_1 + \log y_1, \dots, \log x_n + \log y_n) \\ &= (\log x_1, \dots, \log x_n) + (\log y_1, \dots, \log y_n) = Tx + Ty. \end{aligned}$$

Also,

$$T(c \cdot x) = T((x_1^c, \dots, x_n^c)) = (\log(x_1^c), \dots, \log(x_n^c)) = (c \log x_1, \dots, c \log x_n) = c(Tx).$$

Solution: False since scalar multiplication does not satisfy the vector space axioms.

Problem 5

Let V be a vector space, $T \in \mathcal{L}(V)$ bijective, and c a nonzero scalar. Prove that cT is bijective and that $(cT)^{-1} = c^{-1}T^{-1}$.

Proof. Let $v, v' \in V$ and suppose $cT(v) = cT(v')$. Then $cT(v) = cT(v') \iff cT(v) - cT(v') = 0 \iff c(T(v) - T(v')) = 0$. Since $c \neq 0$, we must have $T(v) = T(v')$. Since T is injective, it follows that $v = v'$. Thus cT is injective.

Let $v \in V$. Since $c \neq 0$, the scalar $\frac{1}{c}$ exists. Since V is closed under scalar multiplication, $\frac{1}{c} \cdot v = \frac{v}{c} \in V$. Since T is surjective, there exists $x \in V$ such that $T(x) = \frac{v}{c}$. Then $cT(x) = c(\frac{v}{c}) = v$. Thus cT is surjective.

Let $v \in V$, and suppose $(cT)^{-1}(v) = x$. Then $cT(x) = v \implies T(x) = \frac{v}{c} \implies x = T^{-1}(\frac{v}{c}) = T^{-1}(c^{-1}v)$. Thus $(cT)^{-1} = c^{-1}T^{-1}$.

Problem 6

Let $T : V \rightarrow W$ be a bijective linear mapping. For each $S \in \mathcal{L}(V)$, define $\rho : \mathcal{L}(V) \rightarrow \mathcal{L}(W)$ (note that the product is defined). Prove that $\rho : \mathcal{L}(V) \rightarrow \mathcal{L}(W)$ is a bijective linear mapping such that $\rho(RS) = \rho(R)\rho(S)$ for all $R, S \in \mathcal{L}(V)$.

Proof. Since the composition of linear mappings is linear, TST^{-1} is a linear mapping. Let $R, S \in \mathcal{L}(V)$. Then

$$\rho(RS) = T(RS)T^{-1} = (TRT^{-1})(TST^{-1}) = \rho(R)\rho(S).$$

Suppose $\rho(S)(v) = 0$ for some $v \in W$. Then $\rho(S)(v) = 0 \iff TST^{-1}(v) = 0$. Now $v = T(x)$ for some $x \in V$ thus $TST^{-1}(T(x)) = TS(x) = 0$. Then since T is injective $S(x) = 0$ thus ρ is injective. It is surjective because for any $U \in \mathcal{L}(W)$, setting $S = T^{-1}UT$ gives $\rho(S) = U$.

Problem 7

Let V be a vector space, $T \in \mathcal{L}(V)$. Prove:

1. If $T^2 = 0$ then $I - T$ is bijective.
2. If $T^n = 0$ for some positive integer n , then $I - T$ is bijective.

Proof. Suppose $T^2 = 0$. Let $v \in V$ and suppose $(I - T)(v) = 0 \implies T(v) - T^2(v) = 0 \implies T(v) = 0$. But $I(v) - T(v) = 0 \implies TI(v) - T^2(v) = T(0) \implies v - T(v) = 0 \implies v = T(v)$. Thus $v = 0$. It follows that $I - T$ is injective.

Let $y \in V$. We want to find $x \in V$ such that $(I - T)(x) = y$. Let $x = y + T(y)$. Then $(I - T)(x) = (I - T)(y + T(y)) = I(y + T(y)) - T(y + T(y)) = I(y) + IT(y) - T(y) - T^2(y) = y + T(y) - T(y) = y$. ■

Proof. We proceed via induction on n . The previous proof shows the base case $n = 2$. Suppose for some $n - 1$, $T^{n-1} = 0$ implies $I - T$ is bijective. Now suppose $T^n = 0$. Let $v \in V$ and suppose $(I - T)(v) = 0$. Composing with T^{n-1} , we get $T^{n-1}(v) - T^n(v) = T^{n-1}(v) = 0$. By the induction hypothesis, this implies $v = 0$, so $I - T$ is injective.

For surjectivity, let $y \in V$ and set $x = y + T(y) + \dots + T^{n-1}(y)$. Then $(I - T)(x) = y$, so $I - T$ is bijective. ■

Problem 8

Let U, V, W be vector spaces over the same field. Form the product vector spaces $U \times V, V \times W$ (Definition 1.3.11), then the product vector spaces $(U \times V) \times W$ and $U \times (V \times W)$. Prove that $(U \times V) \times W \cong U \times (V \times W)$.

Proof. Consider the mapping

$$\rho : (U \times V) \times W \longrightarrow U \times (V \times W), \quad \rho((x_1, x_2), x_3) = (x_1, (x_2, x_3)).$$

Suppose $\rho((x_1, x_2), x_3) = \rho((y_1, y_2), y_3)$. Then $(x_1, (x_2, x_3)) = (y_1, (y_2, y_3))$, so $x_1 = y_1, x_2 = y_2, x_3 = y_3$. Thus $((x_1, x_2), x_3) = ((y_1, y_2), y_3)$, and ρ is injective.

Let $(u, (v, w)) \in U \times (V \times W)$. Then $\rho((u, v), w) = (u, (v, w))$. Thus ρ is surjective. ■

Problem 9

Prove that $\mathbb{R}^2 \times \mathbb{R}^3 \cong \mathbb{R}^5$.

Proof. Consider the mapping

$$\rho : \mathbb{R}^2 \times \mathbb{R}^3 \longrightarrow \mathbb{R}^5, \quad \rho(((x_1, x_2), (x_3, x_4, x_5))) = (x_1, x_2, x_3, x_4, x_5)$$

Suppose $\rho(((x_1, x_2), (x_3, x_4, x_5))) = \rho(((y_1, y_2), (y_3, y_4, y_5)))$. Then $(x_1, x_2) = (y_1, y_2)$ and $(x_3, x_4, x_5) = (y_3, y_4, y_5)$. Thus $((x_1, x_2), (x_3, x_4, x_5)) = ((y_1, y_2), (y_3, y_4, y_5))$, and ρ is injective.

Let $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^5$. Then $\rho(((x_1, x_2), (x_3, x_4, x_5))) = (x_1, x_2, x_3, x_4, x_5)$. Thus ρ is surjective. ■

Problem 10

With notations as in Exercice 8, prove that $V \times W \cong W \times V$.

Proof. Consider the mapping

$$\rho : V \times W \longrightarrow W \times V, \quad \rho((v, w)) = (w, v).$$

Suppose $\rho((v_1, w_1)) = \rho((v_2, w_2))$. Then $(w_1, v_1) = (w_2, v_2)$, so $v_1 = v_2$ and $w_1 = w_2$. Thus $(v_1, w_1) = (v_2, w_2)$, and ρ is injective.

Let $(w, v) \in W \times V$. Then $\rho((v, w)) = (w, v)$. Thus ρ is surjective. ■

Problem 11

With notations as in Example 2.4.2, prove that $(x_1, x_2) \mapsto (x_1, x_2 + x_2, 0)$ is an isomorphism $\mathbb{R}^2 \mapsto W$.

Proof. Consider the mapping

$$\rho : \mathbb{R}^2 \longrightarrow W, \quad \rho((x_1, x_2)) = (x_1, x_2, 0).$$

Suppose $\rho((x_1, x_2)) = \rho((y_1, y_2))$. Then $(x_1, x_2, 0) = (y_1, y_2, 0)$, so $x_1 = y_1$ and $x_2 = y_2$. Thus $(x_1, x_2) = (y_1, y_2)$, and ρ is injective.

Let $(x_1, x_2, 0) \in W$. Then $\rho((x_1, x_2)) = (x_1, x_2, 0)$. Thus ρ is surjective. ■

2.5 Equivalence Relations and Quotient Sets

Problem 2

Let X be the set of all nonzero vectors in $\mathbb{R}^n$, that is, $X = \mathbb{R}^n - \{\theta\}$. For x, y in X , write $x \sim y$ if x is proportional to y , that is, if there exists a scalar c (necessarily nonzero) such that $x = cy$; this is an equivalence relation in X .

Proof. Notice that $x = 1x$, thus $x \sim x$.

Suppose $x \sim y$. Then $x = cy$ for some scalar c . Since $c \neq 0$, we have $\frac{1}{c}x = y$. Thus $y \sim x$.

Suppose $x \sim y$ and $y \sim z$. Then $x = c_1y$ and $y = c_2z$ for some scalars c_1, c_2 . Then $x = c_1y = c_1(c_2z) = (c_1c_2)z$. Thus $x \sim z$. ■

Problem 4

If λ is a nonempty set of vector spaces over the field F , then the relation of isomorphism (2.4.1) is an equivalence relation in λ .

Proof. Let $x, y, z \in \lambda$. Notice that $x \sim x$ through the identity mapping.

Suppose $x \sim y$. Then there exists an isomorphism ρ such that $\rho(x) = y$. But then $x = \rho^{-1}(y)$, thus $y \sim x$.

Suppose $x \sim y$ and $y \sim z$. Then there exist ρ_1, ρ_2 such that $\rho_1(x) = y$ and $\rho_2(y) = z$. Then $z = \rho_2(\rho_1(x))$. Thus $x \sim z$. ■

Problem 9

Let $f : X \longrightarrow Y$ be any mapping, $x \sim x'$ the equivalence relation defined by $f(x) = f(x')$ (2.5.1), $q : X \longrightarrow X/\sim$ the quotient mapping (2.5.17).

Define a mapping $g : X/\sim \longrightarrow f(X)$ as follows : if $u \in X/\sim$ define $g(u) = f(x)$, where x is any element of X such that $u = q(x)$. [If also $u = q(x')$ then $x \sim x'$, so $f(x) = f(x')$; thus $g(u)$ depends only on u not on the particular x chosen to represent the class u .] Prove (cf. Fig. 10):

1. g is bijective;
2. $f = i \circ g \circ q$, where $i : f(X) \longrightarrow Y$ is the insertion mapping (Appendix A.3.8).

In particular, if $f : X \rightarrow Y$ is surjective, then $g : X/\sim \rightarrow Y$ is bijective.

Proof. Suppose $u, u' \in X/\sim$ such that $g(u) = g(u')$. By definition, $g(u) = f(x)$ and $g(u') = f(x')$ for some $x, x' \in X$ with $u = q(x)$ and $u' = q(x')$. Then $f(x) = f(x')$, so $x \sim x'$. Thus $q(x) = q(x')$, and therefore $u = u'$. Thus g is injective.

Let y be an arbitrary element of $f(X)$. Then $y = f(x)$ for some $x \in X$. Let $u = q(x) \in X/\sim$. By definition of g , we have $g(u) = f(x) = y$. Thus g is surjective. ■

Proof. Let $x \in X$. Then $q(x) \in X/\sim$ and $g(q(x)) = f(x)$ by definition of g . Since $i : f(X) \rightarrow Y$ is the insertion mapping, $i(f(x)) = f(x)$. Thus $(i \circ g \circ q)(x) = f(x)$ so $f = i \circ g \circ q$.

Suppose $f : X \rightarrow Y$ is surjective. Then $f(X) = Y$, so $g : X/\sim \rightarrow Y$.

Suppose $u, u' \in X/\sim$ such that $g(u) = g(u')$. Then $g(u) = f(x)$ and $g(u') = f(x')$ for some $x, x' \in X$. Thus $f(x) = f(x')$, so $x \sim x'$ and thus $u = u'$. Thus g is injective.

Let $y \in Y$. Since f is surjective there exists $x \in X$ such that $f(x) = y$. Let $u = q(x) \in X/\sim$. Then $g(u) = f(x) = y$. Thus g is surjective. ■

Problem 10

For x, y in $\mathbb{R}$, write $x \sim y$ if $x - y$ is an integer (that is, $x - y \in \mathbb{Z}$); this is an equivalence relation in $\mathbb{R}$. The quotient set is denoted $\mathbb{R}/\mathbb{Z}$; the equivalence class $[x]$ of $x \in \mathbb{R}$ is the set $x + \mathbb{Z} = \{x + n \mid n \in \mathbb{Z}\}$.

1. There is a connection between functions defined on $\mathbb{R}/\mathbb{Z}$ and functions defined on $\mathbb{R}$ that are periodic of period 1; can you make it precise?
2. There exists a bijection $g : \mathbb{R}/\mathbb{Z} \rightarrow U$, where $U = \{(a, b) \in \mathbb{R}^2 \mid a^2 + b^2 = 1\}$ is the 'unit circle' in $\mathbb{R}^2$. [Hint: Apply Exercise 9 to the function $f : \mathbb{R} \rightarrow U$ defined by $f(t) = (\cos 2\pi t, \sin 2\pi t)$.]

Solution: Consider a periodic function of period 1, say $f : \mathbb{R} \rightarrow \mathbb{R}$. Then for all $x \in \mathbb{R}$ and all $n \in \mathbb{Z}$,

$$f(x + n) = f(x).$$

Thus f takes the same value on all elements of the equivalence class

$$[x] = x + \mathbb{Z}.$$

Conversely, any function defined on $\mathbb{R}/\mathbb{Z}$ determines a function on $\mathbb{R}$ that is periodic of period 1 by composition with the quotient map.

Proof. Consider the function

$$f : \mathbb{R} \rightarrow U, \quad f(t) = (\cos 2\pi t, \sin 2\pi t).$$

Notice that $f(t + n) = f(t)$ for all $n \in \mathbb{Z}$, so f is constant on equivalence classes of $\mathbb{R}/\mathbb{Z}$. By Exercise 9 there is a map

$$g : \mathbb{R}/\mathbb{Z} \rightarrow U, \quad g([t]) = f(t).$$

Suppose $g([t]) = g([s])$. Then $f(t) = f(s)$ so $t - s \in \mathbb{Z}$. Thus $[t] = [s]$. Suppose $(a, b) \in U$. There exists $t \in [0, 1)$ such that $(\cos 2\pi t, \sin 2\pi t) = (a, b)$, so $(a, b) = g([t])$. ■

Problem 11

For x, y in $\mathbb{R}$, write $x \sim y$ if $x - y$ is an integral multiple of 2π (that is, $x - y \in 2\pi\mathbb{Z}$). With an eye on Exercise 10, discuss periodic functions of period 2π and describe a bijection of $\mathbb{R}/2\pi\mathbb{Z}$ onto the unit circle U .

Solution: The equivalence class of $x \in \mathbb{R}$ in $\mathbb{R}/2\pi\mathbb{Z}$ is $[x] = \{x + 2\pi n \mid n \in \mathbb{Z}\}$. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is periodic of period 2π if $f(x + 2\pi) = f(x)$ for all $x \in \mathbb{R}$. A bijection of $\mathbb{R}/2\pi\mathbb{Z}$ onto the unit circle $U = \{(a, b) \in \mathbb{R}^2 \mid a^2 + b^2 = 1\}$ can be defined as

$$g : \mathbb{R}/2\pi\mathbb{Z} \rightarrow U, \quad g([x]) = (\cos x, \sin x),$$

2.6 Quotient Vector Spaces

Problem 1

Prove that if $V = M \oplus N$ (1.6, Exercise 12) then $V/M \cong N$. [Hint: Restrict the quotient mapping $V \rightarrow V/M$ to N (Appendix A.3.8, and calculate the kernel and range of the restricted mapping.)]

Proof. Suppose $V = M \oplus N$. Let $q : V \rightarrow V/M$ be the quotient mapping. Then consider $(q \mid N) : N \rightarrow V/M$, which is the restriction of q to N . It was shown in the proof of Theorem 2.6.1 that q is linear. Suppose $x \in N$ such that $(q \mid N)(x) = [0]$. By definition, this means $x \in M$. But $x \in N$ and $V = M \oplus N$ thus $x = 0$. Thus $(q \mid N)$ is injective. Let $[y] \in V/M$. Since $V = M \oplus N$ then $y = m + n$ with $m \in M$ and $n \in N$. Then $(q \mid N)(n) = [n] = [m + n] = [y]$. Thus $(q \mid N)$ is surjective. Therefore $q \mid N$ is bijective, and it follows that $V/M \cong N$. ■

Problem 2

Let M and N be linear subspaces of the vector spaces V and W , respectively. Form the product vector space $V \times W$ (Definition 1.3.11). Then $M \times N$ is a linear subspace of $V \times W$ and

$$(V \times W)/(M \times N) \cong (V/M) \times (W/N).$$

Prove this by showing that

$$(x, y) + M \times N \rightarrow (x + M, y + N),$$

defines a function $(V \times W)/(M \times N) \rightarrow (V/M) \times (W/N)$ and that this function is linear and bijective. [The essential first step to show that if $u = (x, y) + M \times N = (x', y') + M \times N$, then $(x + M, y + N) = (x' + M, y' + N)$, that is, the proposed functional value at u depends only on u and not on the particular ordered pair $(x, y) \in u$ chosen to represent it (cf. the proof of Theorem 2.6.1).]

Proof. Let $q : (V \times W)/(M \times N) \rightarrow (V/M) \times (W/N)$ be a mapping defined as

$$q((x, y) + M \times N) = (x + M, y + N).$$

We first show that q is well defined. Suppose $u = (x, y) + M \times N = (x', y') + M \times N$. Then

$$(x, y) - (x', y') \in M \times N.$$

Thus $(x - x', y - y') = (m, n)$ for some $m \in M$ and $n \in N$. Then $x - x' \in M$ and $y - y' \in N$ thus $x + M = x' + M$ and $y + N = y' + N$. Thus $(x + M, y + N) = (x' + M, y' + N)$.

We now show q is additive. Let $(x, y), (x', y')$ be elements in $V \times W$. Then

$$q((x, y) + (x', y')) = q((x + x', y + y')) = (x + x' + M, y + y' + N) = (x + M, y + N) + (x' + M, y' + N) = q((x, y)) + q((x', y')).$$

We now show q is homogeneous.

$$q(c(x, y)) = q((cx, cy)) = (cx + M, cy + N) = c(x + M, y + N) = cq(x, y).$$

We now show q is injective. Suppose $q((x, y)) = (0 + M, 0 + N)$. Then $x \in M$ and $y \in N$. Thus $(x, y) \in M \times N$ so $(x, y) + M \times N = M \times N$. Therefore $(x, y) = 0$ in $(V \times W)/(M \times N)$, and q is injective.

We now show q is surjective. Let $(x + M, y + N)$ be an element in $(V/M) \times (W/N)$. Then $q((x, y)) = (x + M, y + N)$, thus q is surjective. ■

Problem 3

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be the linear form $f(x) = 2x_1 - 3x_2 + x_3$ and let M be its kernel. As in Example 2.5.9, think of V/M as the set of planes consisting of M and the planes parallel to it. If u and v are two of these planes, their sum $u + v$ in V/M can be visualized as follows: choose a point P on the plane u and a point Q on the plane v ; add the arrows OP and OQ using the parallelogram rule, obtaining an arrow OR ; the plane through R parallel to M is $u + v$.

Solution: O.K.

Problem 4

Let M be a linear subspace of V , $u = x + M$ and $v = y + M$ two elements of V/M .

1. Let $A = \{s + t \mid s \in u, t \in v\}$ be the sum of the sets $x + M$ and $y + M$ in the sense of Definition 1.6.9. Then A is also a coset, namely $A = x + y + M$. [This offers a definition of $u + v$ that is obviously independent of the vectors x and y chosen to represent u and v .]
2. If c is a nonzero scalar, then the set $B = \{cs \mid s \in u\}$ is a coset, namely $B = cx + M$. What if $c = 0$?

Proof. Let a be an arbitrary element in A such that $a = s + t$. Suppose $s = x + m$ and $t = y + m'$ where $m, m' \in M$. Then taking sums we have $s + t = x + y + m + m'$. Now $m + m' \in M$ since M is closed under addition. Thus $x + y + m + m' \in x + y + M$.

Conversely, suppose $t \in x + y + M$ such that $t = x + 0 + y + m$. Then $0, m \in M$ thus $x + 0 \in x + M$ and $y + m \in y + M$ thus $t \in A$. ■

Proof. Suppose c is a nonzero scalar. Let b be an arbitrary element in B such that $b = cs$ where $s \in u$. Suppose $s = x + m$ where $m \in M$. Then $b = cs = c(x + m) = cx + cm$. Now $cm \in M$ thus $b = cx + cm \in cx + M$.

Conversely, suppose t is an arbitrary element in $cx + M$ such that $t = cx + m$ where $m \in M$. Since $m \in M$ and $c \neq 0$ we have $\frac{1}{c} \cdot m = \frac{m}{c} \in M$, so $t = cx + m \iff t = c(x + \frac{m}{c})$. Now $x + \frac{m}{c} \in u$ so $t \in B$ as required. ■

Solution: If $c = 0$ then B is the set $\{0\}$.

Problem 5

Let V be a vector space, M and N linear subspaces of V , and x, y vectors in V . Prove:

1. $x + M \subset y + N$ if and only if $M \subset N$ and $x - y \in N$.
2. $(x + M) \cap (y + N) \neq \emptyset$ if and only if $x - y \in M + N$.
3. If $z \in (x + M) \cap (y + N)$ then $(x + M) \cap (y + N) = z + M \cap N$.

Proof. ($\implies$) Suppose $x + M \subset y + N$. Let m be an arbitrary element in M . Since $x + m \in x + M$ and $x + M \subset y + N$ we have $x + m = y + n$ for some $n \in N$. Now for arbitrary $m \in M$, we have $x + m = y + n$ for some $n \in N$, so $m = n - (x - y)$. Since $n \in N$ and $x - y \in N$ we have $m \in N$. Thus $M \subset N$ as required.

($\impliedby$) Suppose $M \subset N$ and $x - y \in N$. Let $t \in x + M$ such that $t = x + m$ for some $m \in M$. Since $M \subset N$, $x - y \in N$, we have $m \in N$ thus $t = y + ((x - y) + m) \in y + N$. Therefore $x + M \subset y + N$. ■

Proof. ($\implies$) Suppose $(x + M) \cap (y + N) \neq \emptyset$. Let $t \in (x + M) \cap (y + N)$ such that $t = x + m$ and $t = y + n$ for $m \in M$ and $n \in N$. Then $x + m = y + n \iff x - y = n - m \in N - M = M + N$.

($\impliedby$) Suppose $x - y \in M + N$. Then $x - y = m + n$ for some $m \in M$ and $n \in N$. Then $x - n = y + m$. Then $x - n \in x + N$ and $y + m \in y + M$ thus $x + N \cap y + M \neq \emptyset$. ■

Proof. Suppose $z \in (x + M) \cap (y + N)$. Then $z = x + m$ and $z = y + n$ for $m \in M$ and $n \in N$. Let t be an arbitrary element in $(x + M) \cap (y + N)$. Then $t = x + m'$ and $t = y + n'$ for some $m' \in M$ and $n' \in N$. Then

$$t - z = (x + m') - (x + m) \in M,$$

and

$$t - z = (y + n') - (y + n) \in N.$$

Thus $t - z \in M \cap N$. Thus $t \in z + M \cap N$. Conversely, let $t \in z + M \cap N$. Then $t = z + m$ where $m \in M \cap N$. Thus $t \in x + M$ and $t \in y + N$ and therefore $t \in (x + M) \cap (y + N)$. ■

Problem 6

Let X be a nonempty set. Let R be a subset of the cartesian product $X \times X$, such that (i) R contains the 'diagonal' $X \times X$, that is, $(x, x) \in R$ for all $x \in X$; (ii) R is 'symmetric in the diagonal', that is, if $(x, y) \in R$ then $(y, x) \in R$; and (iii) if $(x, y) \in R$ and $(y, z) \in R$ then $(x, z) \in R$. Does this suggest a way of defining an equivalence relation $x \sim y$ in X .

Solution:

$$x \sim y \iff (x, y) \in R$$

Problem 7

Let $T : V \rightarrow W$ be a linear mapping, M a linear subspace of V such that $M \subset \text{Ker} T$, and $Q : V \rightarrow V/M$ the quotient mapping. Prove:

1. There exists a (unique) linear mapping $S : V/M \rightarrow W$ such that $T = SQ$ (Fig. 11).
2. S is injective if and only if $M = \text{Ker} T$.

Proof. Consider $S : V/M \rightarrow W$ defined

$$S(x + M) = T(x).$$

We first show S is well defined. Suppose $x + M = y + M$. Therefore $x - y \in M$. Since $M \subset \text{Ker} T$, we have $T(x - y) = 0 \implies T(x) = T(y)$. Then let $x + M, y + M$ be arbitrary elements in V/M . Then

$$S((x + M) + (y + M)) = S(x + y + M) = T(x + y) = T(x) + T(y) = S(x + M) + S(y + M).$$

Let c be a scalar. Then

$$S(c(x + M)) = S(cx + M) = T(cx) = cT(x).$$

Thus S is a linear mapping. Now let v be an arbitrary element in V . Then $(SQ)(v) = S(Q(v)) = S(v + M) = T(v)$.

Suppose there exists S' such that $S' : V/M \rightarrow W$ and $T = S'Q$. For all $v \in V$ we have $T(v) = S'(Q(v)) = S(Q(v))$. Thus S, S' agree at all cosets and therefore $S' = S$ and S is unique. ■

Proof. Suppose S is injective. We know $M \subset \text{Ker} T$ so let t be an arbitrary element in $\text{Ker} T$. Then $T(t) = (SQ)(t) = S(t + M) = 0$. Since S is injective we have $t + M = 0 + M$ thus $t \in M$. Thus $M = \text{Ker} T$.

Suppose $M = \text{Ker} T$. Let $x + M$ be an arbitrary element in V/M . Suppose $S(x + M) = 0$. Then $T(x) = 0$ thus $x \in M$ and S is injective. ■

Problem 8

Let M be a linear subspace of V , $Q : V \rightarrow V/M$ the quotient mapping, $T : V \rightarrow V$ a linear mapping such that $T(M) \subset M$. Prove that there exists a (unique) linear mapping: $S : V/M \rightarrow V/M$ such that $SQ = QT$ (Fig. 12). [Hint: Apply Exercise 7 to the linear mapping $QT : V \rightarrow V/M$.]

Proof. Let v be an arbitrary element in M . Since $T(M) \subset M$, we have $T(v) \in M$. Then $QT(v) = Q(T(v)) = 0$. Thus $M \subset \text{Ker}(QT)$. We can then apply Exercise 7 to find a linear mapping $S : V/M \rightarrow V/M$ such that $SQ = QT$. ■

2.7 The First Isomorphism Theorem

Problem 1

When there's a first, there's a second. The following result is called the Second isomorphism theorem. Let V be a vector space, M and N linear subspaces of V , $M \cap N$ their intersection and $M + N$ their sum (1.6.9). Then $M \cap N$ is a linear subspace of M , N is a linear subspace of $M + N$, and

$$M/(M \cap N) \cong (M + N)/N.$$

[Hint: Let $Q : V \rightarrow V/N$ be the quotient mapping and let $T = Q|_M$ be the restriction of Q to M (Appendix 3.8), that is, $T : M \rightarrow V/N$ and $Tx = x + N$ for all $x \in M$. Show that T has range $(M + N)/N$ and kernel $M \cap N$.]

Proof. Let $Q : V \rightarrow V/N$ be the quotient mapping and let $T = Q|_M$ be the restriction of Q to M , that is, $T : M \rightarrow V/N$ and $Tx = x + N$ for all $x \in M$.

We first show that $\text{Ran}(T) = (M + N)/N$. Notice

$$\text{Ran}(T) = T(M) = \{m + N : m \in M\}.$$

$\text{Ran}(T) \subset (M + N)/N$: Let $x \in \text{Ran}(T)$. Then $x = m + N$ for some $m \in M$. Since $m \in M \subset M + N$, we have $m + N \in (M + N)/N$. Thus $\text{Ran}(T) \subset (M + N)/N$.

$(M + N)/N \subset \text{Ran}(T)$: Let $x \in (M + N)/N$. Then $x = (m + n) + N$ for some $m \in M, n \in N$. But $(m + n) + N = m + N = Tm$, so $x \in \text{Ran}(T)$.

Thus

$$\text{Ran}(T) = (M + N)/N.$$

We now show that $\text{Ker}(T) = M \cap N$.

$\text{Ker}(T) \subset M \cap N$: Suppose $x \in \text{Ker}(T)$. Then $Tx = x + N = N$, so $x \in N$. Since $x \in M$, we have $x \in M \cap N$.

$M \cap N \subset \text{Ker}(T)$: Suppose $x \in M \cap N$. Then $x \in N$, so $Tx = x + N = N$. Thus $x \in \text{Ker}(T)$.

Thus $\text{Ker}(T) = M \cap N$.

Therefore, by the First Isomorphism Theorem,

$$M/(M \cap N) \cong \text{Ran}(T) = (M + N)/N.$$

Problem 2

Let M, N be linear subspaces of the vector spaces V, W and let

$$T : V \times W \rightarrow (V/M) \times (W/N),$$

be the linear mapping defined by $T(x, y) = (x + M, y + N)$. Determine the kernel of T and deduce another proof of 2.6, Exercise 2.

2.6 Problem 2

Let M and N be linear subspaces of the vector spaces V and W , respectively. Form the product vector space $V \times W$ (Definition 1.3.11). Then $M \times N$ is a linear subspace of $V \times W$ and

$$(V \times W)/(M \times N) \cong (V/M) \times (W/N).$$

Prove this by showing that

$$(x, y) + M \times N \longrightarrow (x + M, y + N),$$

defines a function $(V \times W)/(M \times N) \longrightarrow (V/M) \times (W/N)$ and that this function is linear and bijective.

[The essential first step to show that if $u = (x, y) + M \times N = (x', y') + M \times N$, then $(x + M, y + N) = (x' + M, y' + N)$, that is, the proposed functional value at u depends only on u and not on the particular ordered pair $(x, y) \in u$ chosen to represent it (cf. the proof of Theorem 2.6.1).]

Proof. We must find $(v, w) \in V \times W$ such that

$$T(v, w) = (0 + M, 0 + N) \in (V/M) \times (W/N).$$

This holds iff $v + M = M$ and $w + N = N$ so $v \in M$ and $w \in N$. Thus

$$\ker(T) = M \times N.$$

By the First Isomorphism Theorem

$$(V \times W)/(M \times N) \cong \text{Ran}(T) = (V/M) \times (W/N).$$

■

Problem 3

Let V be a vector space over F , $f : V \longrightarrow F$ a linear form on V (Definition 2.1.11), N the kernel of f . Prove that if f is not identically zero, then $V/N \cong F$. [Hint: If $f(z) \neq 0$ and c is any scalar, evaluate f at the vector $(c/f(z))z$.]

Proof. Suppose f is not identically zero. Thus there exists $v \in V$ such that $f(v) \neq 0$. Suppose $f(v) = t$ where $t \in F$ and let c be an arbitrary element in F . We first show f is surjective. Let $c' = \frac{c}{t}$. Then

$$f(c'v) = c'f(v) = \frac{c}{t} \cdot t = c.$$

Thus f is surjective. Now $\ker(f) = N$ by definition. Thus, by the First Isomorphism Theorem

$$V/N \cong \text{Ran}(f) = F.$$

■

Problem 4

Let V be a vector space, $V \times V$ the product vector space (1.3.11), and $\nabla = \{(x, x) \mid x \in V\}$ (called the diagonal of $V \times V$). Prove that ∇ is a linear subspace of $V \times V$ and that $(V \times V)/\nabla \cong V$. [Hint: 2.2, Exercise 4.]

Proof. Firstly, $(0, 0) \in \nabla$. Let $(x, x), (y, y) \in \nabla$ and c a scalar. Then $(x, x) + (y, y) = (x + y, x + y) \in \nabla$. Also, $c(x, x) = (cx, cx) \in \nabla$. Thus ∇ is a linear subspace of $V \times V$. Define $f : V \times V \rightarrow V$ by $f(x, y) = x - y$. Then $\ker(f) = \{(x, x) \mid x \in V\} = \nabla$. By Exercise 4 of 2.2, f is surjective. Thus by the First Isomorphism Theorem

$$(V \times V)/\nabla \cong V.$$

■

Problem 5

Let N be the set of vectors in $\mathbb{R}^5$ whose last two coordinates are zero:

$$N = \{x_1, x_2, x_3, 0, 0\} \mid x_1, x_2, x_3 \in \mathbb{R}.$$

Prove that N is a linear subspace of $\mathbb{R}^5$, $\mathbb{R}^3 \cong N$, and $\mathbb{R}^5/N \cong \mathbb{R}^2$. [Taking some liberties with the notation, $\mathbb{R}^5/\mathbb{R}^3 = \mathbb{R}^2$.]

Proof. Firstly, $(0, 0, 0, 0, 0) \in N$. Let $(x_1, x_2, x_3, 0, 0), (y_1, y_2, y_3, 0, 0) \in N$ and c be a scalar. Then $(x_1, x_2, x_3, 0, 0) + (y_1, y_2, y_3, 0, 0) = (x_1 + y_1, x_2 + y_2, x_3 + y_3, 0, 0) \in N$. Also $c(x_1, x_2, x_3, 0, 0) = (cx_1, cx_2, cx_3, 0, 0) \in N$. Thus N is a linear subspace of $\mathbb{R}^5$.

Consider the mapping $f : \mathbb{R}^3 \rightarrow N$ defined

$$f(x_1, x_2, x_3) = (x_1, x_2, x_3, 0, 0).$$

Let $(x_1, x_2, x_3, 0, 0) \in N$. Then $f(x_1, x_2, x_3) = (x_1, x_2, x_3, 0, 0)$, so f is surjective. Also, if $f(x_1, x_2, x_3) = (0, 0, 0, 0, 0)$ then $(x_1, x_2, x_3) = (0, 0, 0)$, so f is injective. Thus f is bijective, and $\mathbb{R}^3 \cong N$.

Consider the linear mapping $g : \mathbb{R}^5 \rightarrow \mathbb{R}^2$ defined by

$$g(x_1, x_2, x_3, x_4, x_5) = (x_4, x_5).$$

Then $\ker(g) = N$, and g is surjective. Thus by the First Isomorphism Theorem

$$\mathbb{R}^5/N \cong \mathbb{R}^2.$$

Problem 6

Let T be a set, A a subset of T , $V = \mathcal{F}(T, \mathbb{R})$ and $W = \mathcal{F}(A, \mathbb{R})$ the vector spaces of functions constructed in 1.3.4. Let N be the set of all functions $x \in V$ such that $x(t) = 0$ for all $t \in A$. Then N is a linear subspace of V and $V/N \cong W$. [Hint: Consider the mapping $x \mapsto x|_A$, ($x \in V$), where $x|_A$ is the restriction of the function x to A (Appendix A.3.8).]

Proof. Firstly, the zero function $0 \in V$ satisfies $0(t) = 0$ for all $t \in A$, so $0 \in N$. Let $x, y \in N$ and c be a scalar. Then for all $t \in A$

$$(x + y)(t) = x(t) + y(t) = 0 + 0 = 0,$$

and

$$(cx)(t) = cx(t) = c \cdot 0 = 0.$$

Thus $x + y \in N$ and $cx \in N$, so N is a linear subspace of V . Consider the mapping $f : V \rightarrow W$ defined by

$$f(x) = x|_A.$$

Then f is linear. Also

$$\ker(f) = \{x \in V \mid x(t) = 0 \text{ for all } t \in A\} = N.$$

Since every function in W is the restriction of some function in V , f is surjective. Thus by the First Isomorphism Theorem

$$V/N \cong W.$$

3 Structure of Vector Spaces

3.1 Linear Subspace Generated by a Subset

Problem 1

If $T : V \rightarrow W$ is linear and A is a subset of V such that $T(A)$ is generating for W , then T is surjective.

Proof. Let w be an arbitrary element in W . Since $T(A)$ generates W , w is a linear combination of elements in $T(A)$. Suppose

$$w = c_1 a_1 + \cdots + c_n a_n \text{ such that } a_1, \dots, a_n \in T(A).$$

But $a_i = T(v_i)$ for some $v_i \in V$, so

$$c_1 a_1 + \cdots + c_n a_n = c_1 T(v_1) + \cdots + c_n T(v_n) = T(c_1 v_1) + \cdots + T(c_n v_n) = T(c_1 v_1 + \cdots + c_n v_n).$$

The last two equalities hold since T is linear. Thus T is surjective. ■

Problem 2

If M and N are linear subspaces of V such that $V = M + N$ (1.6.9) and $T : V \rightarrow V/N$ is the quotient mapping, then $T(M) = V/N$ but M need not equal V ; thus we cannot conclude in Exercise 1 that A is a generating for V . Can you exhibit an example of such V, M and N ?

Proof. Consider $V = \mathbb{R}$, $M = \{0\}$, and $N = \mathbb{R}$. Then $V = M + N$, but $M \neq V$. Let $T : V \rightarrow V/N$ be the quotient map. Since $V/N = \{0\}$, we have $T(M) = \{0\} = V/N$. ■

Problem 3

(Substitution Principle) Suppose $x, x_1, \dots, x_n$ are vectors such that x is a linear combination of $x_1, \dots, x_n$ but not of $x_1, \dots, x_{n-1}$. Prove:

$$[\{x_1, \dots, x_n\}] = [\{x_1, \dots, x_{n-1}, x\}].$$

Proof. Since x is a linear combination of $x_1, \dots, x_n$ suppose

$$x = c_1 x_1 + \cdots + c_n x_n.$$

Since x is not a linear combination of $x_1, \dots, x_{n-1}$, we have $c_n \neq 0$, so

$$x_n = \frac{1}{c_n} (x - c_1 x_1 - \cdots - c_{n-1} x_{n-1}).$$

Let $v \in [\{x_1, \dots, x_n\}]$ and suppose

$$v = d_1 x_1 + \cdots + d_n x_n.$$

Substituting for x_n , we get

$$v = d_1 x_1 + \cdots + d_{n-1} x_{n-1} + d_n \frac{1}{c_n} (x - c_1 x_1 - \cdots - c_{n-1} x_{n-1}) \in [\{x_1, \dots, x_{n-1}, x\}].$$

Conversely, let $v \in [\{x_1, \dots, x_{n-1}, x\}]$ and suppose

$$v = d_1 x_1 + \cdots + d_{n-1} x_{n-1} + dx.$$

Then

$$x = c_1 x_1 + \cdots + c_n x_n \implies dx = (dc_1)x_1 + \cdots + (dc_n)x_n.$$

Thus

$$v = d_1 x_1 + \cdots + d_{n-1} x_{n-1} + (dc_1)x_1 + \cdots + (dc_n)x_n \in [\{x_1, \dots, x_n\}].$$

■

Problem 4

Let $S : V \rightarrow W$ and $T : V \rightarrow W$ be linear mappings, and let A be a subset of V such that $V = [A]$. Prove that if $Sx = Tx$ for all $x \in A$, then $S = T$. (Informally, a linear mapping is determined by its values on a generating set.)

Proof. Suppose $Sx = Tx$ for all $x \in A$. Let $x' \in V$ and since $V = [A]$ so

$$x' = c_1 a_1 + \cdots + c_n a_n.$$

Then

$$\begin{aligned} T(x') &= T(c_1 a_1 + \cdots + c_n a_n) \\ &= T(c_1 a_1) + \cdots + T(c_n a_n) \\ &= c_1 T(a_1) + \cdots + c_n T(a_n) \\ &= c_1 S(a_1) + \cdots + c_n S(a_n) \\ &= S(c_1 a_1) + \cdots + S(c_n a_n) \\ &= S(c_1 a_1 + \cdots + c_n a_n) \\ &= S(x'). \end{aligned}$$

Thus $Sx' = Tx'$ for all $x' \in V$, so $S = T$. ■

Problem 5

Let V be a vector space, A a nonempty subset of V . For each finite subset $\{x_1, \dots, x_n\}$ of A , let

$$M(x_1, \dots, x_n) = \{a_1 x_1 + \cdots + a_n x_n \mid a_1, \dots, a_n \text{ scalars}\};$$

as noted in the proof of Corollary 3.1.5, this is a linear subspace of V . Let $\mathcal{N}$ be the set of all linear subspaces of V obtained in this way (by considering all possible finite subsets of A); by definition, $[A] = \bigcup \mathcal{N}$ (the union of the subspaces belonging to $\mathcal{N}$). Prove that $[A]$ is a linear subspace of V by verifying criterion 1.6, Exercise 9.

Proof. We must show that if $M, K \in \mathcal{N}$ then there exists $P \in \mathcal{N}$ such that $M \subset P$ and $K \subset P$. Therefore suppose $M, K \in \mathcal{N}$. Then there exist finite subsets $\{x_1, \dots, x_n\} \subseteq A$ and $\{y_1, \dots, y_m\} \subseteq A$ such that

$$M = M(x_1, \dots, x_n), \quad K = M(y_1, \dots, y_m).$$

Let

$$P = M(x_1, \dots, x_n, y_1, \dots, y_m).$$

Then $P \in \mathcal{N}$, and clearly $M \subset P$ and $K \subset P$. From 1.6 Exercise 9 it follows that $\bigcup \mathcal{N}$ is a linear subspace of V . ■

Problem 6

Let V and W be vector spaces, $V \times W$ the product vector space (1.3.11); let $A \subset V$ and $B \subset W$. Prove:

1. $[A \times B] \subset [A] \times [B]$.
2. In general, $[A \times B] \neq [A] \times [B]$. **Hint:** Consider $A = \emptyset, B = W$; or $V = W = R$ and $A = B = \{1\}$.
3. $[(A \times \{\emptyset\}) \cup (\{\emptyset\} \times B)] = [A] \times [B]$. **Hint**

$$\left(\sum_{i=1}^m c_i(x_i, 0) + \sum_{j=1}^n d_j(0, y_j) \right).$$

Proof. Let (x, y) be an arbitrary element in $[A \times B]$. Then

$$(x, y) = c_1(a_1, b_1) + \cdots + c_n(a_n, b_n)$$

for some $(a_i, b_i) \in A \times B$. But then

$$(x, y) = (c_1 a_1 + \cdots + c_n a_n, c_1 b_1 + \cdots + c_n b_n).$$

Thus

$$x = c_1 a_1 + \cdots + c_n a_n \in [A], \quad y = c_1 b_1 + \cdots + c_n b_n \in [B],$$

so $(x, y) \in [A] \times [B]$. ■

Proof. Let $V = W = \mathbb{R}$ and $A = B = \{1\}$. Then

$$A \times B = \{(1, 1)\},$$

so

$$[A \times B] = \{c(1, 1) \mid c \in \mathbb{R}\} = \{(c, c) \mid c \in \mathbb{R}\}.$$

Conversely

$$[A] = \mathbb{R}, \quad [B] = \mathbb{R},$$

so

$$[A] \times [B] = \mathbb{R} \times \mathbb{R}.$$

Since

$$\{(c, c) \mid c \in \mathbb{R}\} \neq \mathbb{R} \times \mathbb{R},$$

it follows that

$$[A \times B] \neq [A] \times [B].$$
■

Proof. Let $(x, y) \in [(A \times \{0\}) \cup (\{0\} \times B)]$. Then (x, y) is a linear combination of elements of the form $(a, 0)$ and $(0, b)$ so

$$(x, y) = \sum_{i=1}^m c_i(x_i, 0) + \sum_{j=1}^n d_j(0, y_j),$$

where $x_i \in A$ and $y_j \in B$. Then

$$(x, y) = \left(\sum_{i=1}^m c_i x_i, \sum_{j=1}^n d_j y_j \right),$$

so $x \in [A]$ and $y \in [B]$. Thus

$$[(A \times \{0\}) \cup (\{0\} \times B)] \subset [A] \times [B].$$

Conversely let $(x, y) \in [A] \times [B]$. Then

$$x = \sum_{i=1}^m c_i x_i, \quad y = \sum_{j=1}^n d_j y_j,$$

with $x_i \in A, y_j \in B$. Then

$$(x, y) = \sum_{i=1}^m c_i(x_i, 0) + \sum_{j=1}^n d_j(0, y_j),$$

which shows $(x, y) \in [(A \times \{0\}) \cup (\{0\} \times B)]$. Thus

$$[(A \times \{0\}) \cup (\{0\} \times B)] = [A] \times [B].$$
■

3.2 Linear Independence

Problem 1

Let $V = \mathcal{F}(\mathbb{R})$ be the vector space of all real-valued functions of a real variable (Example 1.3.4). Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by $f(t) = \sin(t + \pi/6)$. Prove that the function f , $\sin$, $\cos$ are linearly dependent.

Proof. Simply note

$$\begin{aligned} f(t) &= \sin(t + \pi/6) \\ &= \sin(t) \cos(\pi/6) + \cos(t) \sin(\pi/6) \\ &= \frac{\sqrt{3}}{2} \sin(t) + \frac{1}{2} \cos(t). \end{aligned}$$

Problem 2

In the vector space of real polynomial function (Example 1.3.5), show that the function p_1, p_2, p_3, p_4 defined by $p_1(t) = t^2 + t + 1$, $p_2(t) = t^2 + 5$, $p_3(t) = 2t + 1$, $p_4(t) = 3t^2 + 5t$ are linearly dependent (cf. 1.5.3).

Proof. Simply note

$$\begin{aligned} -35p_1(t) + 8p_2(t) - 5p_3(t) + 9p_4(t) &= -35(t^2 + t + 1) + 8(t^2 + 5) - 5(2t + 1) + 9(3t^2 + 5t) \\ &= (-35t^2 - 35t - 35) + (8t^2 + 40) + (-10t - 5) + (27t^2 + 45t) \\ &= 0. \end{aligned}$$

Problem 3

Show that in $\mathbb{R}^3$ the vectors e_1, e_2, e_3 , $(3, -2, 4)$ are linearly dependent (cf. 1.5.3).

Proof. Simply note

$$(3, -2, 4) = 3e_1 - 2e_2 + 4e_3.$$

Problem 5

If $T : V \rightarrow W$ is a linear mapping and $x_1, \dots, x_n$ are linearly dependent vectors in V , then the vectors $Tx_1, \dots, Tx_n$ of W are linearly dependent.

Proof. Since $x_1, \dots, x_n$ are linearly dependent, there exist scalars $c_1, \dots, c_n$, not all zero, such that

$$c_1x_1 + \dots + c_nx_n = 0 \implies T(c_1x_1 + \dots + c_nx_n) = T(0) = c_1T(x_1) + \dots + c_nT(x_n) = 0.$$

Problem 6

A finite list of vectors $x_1, \dots, x_n$ is linearly dependent if and only if there exists an index j such that $x_j \in [\{x_i \mid i \neq j\}]$ (cf. 3.1.5).

Proof. ($\implies$) Suppose a finite list of vectors $x_1, \dots, x_n$ is linearly dependent. We have $c_1, \dots, c_n$ not all zero such that

$$c_1x_1 + \dots + c_nx_n = 0.$$

Wlog suppose $c_j \neq 0$ for some $1 \leq j \leq n$. Then

$$x_j = \frac{-1}{c_j}(c_1x_1 + \dots + c_{j-1}x_{j-1} + c_{j+1}x_{j+1} + \dots + c_nx_n) \in [\{x_i \mid i \neq j\}].$$

($\impliedby$) Suppose there exists an index j such that $x_j \in [\{x_i \mid i \neq j\}]$. Then we have

$$x_j = c_1x_1 + \dots + c_{j-1}x_{j-1} + c_{j+1}x_{j+1} + \dots + c_nx_n.$$

Thus

$$-x_j + c_1x_1 + \dots + c_{j-1}x_{j-1} + c_{j+1}x_{j+1} + \dots + c_nx_n = 0.$$

Thus $x_1, \dots, x_n$ is linearly dependent. ■

Problem 7

If $T : V \longrightarrow W$ is an injective linear mapping and if $x_1, \dots, x_n$ are vectors in V such that $Tx_1, \dots, Tx_n$ are linearly dependent in W , then $x_1, \dots, x_n$ are linearly dependent in V .

Proof. Suppose $T : V \longrightarrow W$ is an injective linear mapping and $Tx_1, \dots, Tx_n$ are linearly dependent in W . We have scalars $c_1, \dots, c_n$ not all zero such that

$$c_1Tx_1 + \dots + c_nTx_n = T(c_1x_1 + \dots + c_nx_n) = 0.$$

Since T is injective we have

$$c_1x_1 + \dots + c_nx_n = 0.$$

Thus $x_1, \dots, x_n$ are linearly dependent. ■

Problem 8

Let V be a vector space. A subset A of V is said to be linearly dependent if there exists a finite list of distinct vectors $x_1, \dots, x_n$ in A that is linearly dependent in the sense of Definition 3.2.1. Prove:

1. A subset of A of V is linearly dependent if and only if some vector in A is the linear span of the remaining vectors of A , that is, there exists $x \in A$ such that $x \in [A - \{x\}]$.
2. If $A \subset B \subset V$ and A is linearly dependent, then B is also linearly dependent.

Proof. This just follows from Problem 6. ■

Proof. Since B is linearly dependent there exists scalars $c_1, \dots, c_n$ not all zero such that

$$c_1x_1 + \dots + c_nx_n = 0.$$

Since $A \subset B$ we have $x_1, \dots, x_n \in A$ thus A is linearly dependent. ■

3.3 Linear Independence

Problem 1

Show that the vectors $(2, -1, 1)$, $(-1, 2, 1)$, $(1, 1, 1)$ in $\mathbb{R}^3$ are linearly independent.

Proof. Suppose there exists scalars not all zero c_1, c_2, c_3 such that

$$c_1(2, -1, 1) + c_2(-1, 2, 1) + c_3(1, 1, 1) = 0.$$

Wlog suppose $c_1 \neq 0$ so we have

$$(2, -1, 1) = \frac{c_1}{c_1}(-1, 2, 1) + \frac{c_3}{c_1}(1, 1, 1).$$

■

Problem 2

Problem 3

Problem 6

Problem 9

Problem 12