

Glaucoma Classification

Glaucoma Classification:

Dropdown 1: Cup-to-Disc Ratio (CDR)

Field ID: cup_to_disc_ratio

Type: single-select

Options:

- 0 – ≤ 0.3 (normal)
- 1 – 0.4–0.5 (borderline)
- 2 – 0.6–0.7 (suspicious)
- 3 – 0.8–0.9 (advanced cupping)
- 4 – 1.0 (total cupping)

Dropdown 2: Neuroretinal Rim

Field ID: neuroretinal_rim

Type: single-select

Options:

- 0 – Normal (ISNT rule preserved: Inferior $\geq$ Superior $\geq$ Nasal $\geq$ Temporal)
- 1 – ISNT rule violation (thinning not following normal pattern)
- 2 – Focal notching (localized rim loss)
- 3 – Diffuse thinning
- 4 – Near-total or total rim loss

Dropdown 3: Disc Hemorrhage (Drance Hemorrhage)

Field ID: disc_hemorrhage

Type: single-select

Options:

- 0 – None
- 1 – Present (splinter hemorrhage at or near disc margin)

Dropdown 4: Peripapillary Atrophy (PPA)

Field ID: peripapillary_atrophy

Type: single-select

Options:

- 0 – None
- 1 – Beta-zone PPA only (atrophy with visible sclera/choroidal vessels)
- 2 – Large or progressive beta-zone PPA

Dropdown 5: Retinal Nerve Fiber Layer (RNFL) Defect

Field ID: rnfl_defect

Type: single-select

Options:

- 0 – No visible RNFL defect
- 1 – Wedge-shaped RNFL defect (localized)
- 2 – Diffuse RNFL thinning
- 3 – Both wedge defect and diffuse thinning

Dropdown 6: Optic Disc Pallor

Field ID: disc_pallor

Type: single-select

Options:

- 0 – Normal color
- 1 – Mild pallor

2 – Significant pallor (consider other optic neuropathies)

Dropdown 7: Vessel Changes

Field ID: vessel_changes

Type: single-select

Options:

0 – Normal vessel pattern

1 – Bayoneting (vessels deflect at rim edge)

2 – Nasalization of vessels

3 – Both bayoneting and nasalization

Note: Glaucoma diagnosis from fundus photography alone has limitations. IOP measurement, visual field testing (perimetry), and OCT of the RNFL/ganglion cell layer are typically required for definitive diagnosis and staging. These labels capture structural findings visible on fundus images.