

MAT327 PROBLEM SET 1

JACK CERONI

CONTENTS

1. Problem 1	2
2. Problem 2	2
3. Problem 4	2
4. Problem 5	2

1. PROBLEM 1

Since the sequence is Cauchy, it converges to some point x . It may be the case that x is in $\{x_k\}$, or it may be the case that it is not. Either way, we can take $B = A \cup \{x\}$. Pick an open cover of A called \mathcal{U} . Let $V \in \mathcal{U}$ contain $\{x\}$. Since the sequence converges to $\{x\}$, all but finitely many of the x_k must be in V . Taking elements of \mathcal{U} that contain all other elements of x_k not in V , and we have a finite subcover, so A is compact.

Take the sequence $x_k = k$. Clearly, this sequence is not Cauchy, as $d(x_m, x_n) = |m - n| \geq 1$ for all m and n not equal. In addition, this set is not compact as it is not bounded (and it will clearly not be bounded even if we add a point).

2. PROBLEM 2

It is easy to see from the definition that $\lambda(0) = 0$.

First consider $\mu \circ \lambda$. We pick $\epsilon > 0$, and choose δ small enough such that $|\lambda(x)| \leq C|x|$, and $|x| < \delta \Rightarrow |\mu(x)| < \frac{\epsilon}{C}|x|$. We then note that:

$$|x| < \min\left(\delta, \frac{\delta}{C}\right) \Rightarrow |\lambda(x)| \leq C|x| < \delta \Rightarrow |\mu(\lambda(x))| < \frac{\epsilon}{C}|\lambda(x)| \leq \epsilon|x|$$

so $\mu \circ \lambda$ is tiny. Now, we can consider $\lambda \circ \mu$. We choose $\epsilon > 0$, and δ small enough such that $|\lambda(x)| \leq C|x|$, and $|x| < \delta$ implies $|\mu(x)| < \frac{\epsilon}{C}|x|$. It follows that:

$$|x| < \min\left(\delta, \frac{\delta C}{\epsilon}\right) \Rightarrow |\mu(x)| < \frac{\epsilon}{C}|x| \Rightarrow |\lambda(\mu(x))| \leq C|\mu(x)| < \epsilon|x|$$

where we know that $|\lambda(\mu(x))| \leq C|\mu(x)|$ as $|\mu(x)| < \delta$.

3. PROBLEM 4

Recall the axis-crawl method. Let U be an open set inside which all the partial derivatives exist and are bounded. We let $h = (h_1, \dots, h_n)$ be in U , and define the sequence of points $t_0 = (0, \dots, 0)$, $t_1 = (h_1, 0, \dots, 0)$, \dots , $t_n = h$. We then note that:

$$f(h) - f(0) = \sum_{j=1}^n f(t_j) - f(t_{j-1}) = \sum_{i=1}^n f(t_{i-1} + h_i e_i) - f(t_{i-1})$$

Note that the functions obtained by setting all h_j constant, except for the k -th spot are differentiable in U , as the partial derivatives exist. Hence, we know from the mean value theorem that there exists some $c_i \in (0, h_i)$ for which $f(t_{i-1} + h_i e_i) - f(t_{i-1}) = D_i f(c_i) h_i$. Thus:

$$f(h) - f(0) = \sum_{j=1}^n D_i f(c_i) h_i \Rightarrow |f(h) - f(0)| = \left| \sum_{j=1}^n D_i f(c_i) h_i \right| \leq \sum_{j=1}^n |D_i f(c_i) h_i| \leq M \sum_{j=1}^n |h_i|$$

as each partial derivative is bounded. Taking the limit as $h \rightarrow 0$ allows us to conclude from squeeze theorem that $\lim_{h \rightarrow 0} |f(h) - f(0)| = 0$, so by definition, f is continuous at 0.

4. PROBLEM 5

Part A

From chain rule:

$$[Dh](x, y) = [Df](x, y, g(x, y)) \circ [Dk](x, y)$$

where $k(x, y) = (x, y, g(x, y))$. Clearly:

$$[Df](x, y, g(x, y)) = (D_1f(x, y, g(x, y)) \quad D_2f(x, y, g(x, y)) \quad D_3f(x, y, g(x, y)))$$

and:

$$[Dk](x, y) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ D_1g(x, y) & D_2g(x, y) \end{pmatrix}$$

which implies that:

$$[Dh](x, y) = (D_1f(x, y, g(x, y)) + D_3f(x, y, g(x, y))D_1g(x, y) \quad D_2f(x, y, g(x, y)) + D_3f(x, y, g(x, y))D_2g(x, y))$$