

NE585

NUCLEAR FUEL CYCLES

Nuclear engineering basics

1

R. A. Borrelli

University of Idaho

Idaho Falls Center for Higher Education



University of Idaho
Department of Nuclear Engineering
and Industrial Management

Learning objectives

Demonstrating fundamental concepts in nuclear physics

Analyzing decay data to identify radionuclides

Summarizing different decay mechanisms

Book – Chapter 2

Learning nodes

Relativity

Atomic structure

Binding energy

Q-value

Radioactivity

Chart of the nuclides

Decay chains

Half life

Relativity

We apply energy from nuclear reactions and radiation interactions with matter

$$E_{REST} = m_0 c^2 \quad (1)$$

$$E = mc^2 \quad (2)$$

Rest mass energy for an electron – 0.511 *MeV*

Save that for later

Energy released or energy injected




Figure. Fission




Figure. Fusion

Determine the energy of a fusion neutron that is not at rest

$$m = \frac{m_0}{\sqrt{1 - (\frac{v^2}{c^2})}} \rightarrow E = mc^2 \quad (3)$$

With $v = 5.2 \times 10^7 \text{ m/s}$, that is about $0.17c$; not negligible

So we have to compute the not rest mass

$$m_0(n) = 1.674927211 \times 10^{-27} \text{ kg}$$

Fusion neutron at that speed has $E = 14.5 \text{ MeV}$

For relativistic calculations use MeV

Atomic structure

The shell model describes how subatomic particles are arranged




Figure. Calcium shell model

15: Phosphorus

2,8,5




Figure. Phosphorus shell model

Notation

$${}^A_Z Q_{A-Z} = {}^{235}_{92} U_{143} \quad (4)$$

Binding energy

There are four fundamental forces of interaction

Gravity – Obvious

Strong nuclear – Holds the nucleus together

Electromagnetic – Basically electrons & magnetism

Weak nuclear – Changes the flavor of quarks

To blow out a nucleus, a lot of energy is needed

Would it be constant or variable per nucleus?

Why would you want to blow out the nucleus?

$$\Delta m = (A - Z)m_n - Zm_p - M \quad (5)$$

$$\Delta m({}^{239}_{94}\text{Pu}) = 1759 \text{ MeV} = 7.40 \frac{\text{MeV}}{\text{nucleon}} \quad (6)$$

This energy corresponds to a velocity of 0.88c

$$931.5 \frac{\text{MeV}}{\text{amu}}$$

Binding energy is used to compare atomic stability, reaction energy, probability of fission/fusion




Figure. Binding energy curve

Light nuclei get more stable by fusion; heavy nuclei get more stable by fission




Figure. Inverted binding energy curve

Q-value

The Q-value for a reaction indicates energy requirements



- (1) Conservation of nucleons
- (2) Conservation of charge
- (3) Conservation of momentum
- (4) Conservation of energy

Compute the Q value for this reaction





$$Q = [(M_{\text{Li}} + M_n) - (M_{\alpha} + M_H)]c^2 \quad (10)$$

$$Q = 4.78 \text{ MeV} \quad (11)$$

$$Q > 0 \rightarrow ?$$

$$Q < 0 \rightarrow ?$$

What is the energy of fission from thorium and uranium fuel cycles?



For comparison, burning a carbon atom releases 4 eV

Radioactivity

Scientists in late 1800s made discoveries which would change the course of science, medicine, and history in the 20th Century



The Nobel Prize in Physics 1903 was divided, one half awarded to Antoine Henri Becquerel *"in recognition of the extraordinary services he has rendered by his discovery of spontaneous radioactivity"* the other half jointly to Pierre Curie and Marie Curie, née Skłodowska *"in recognition of the extraordinary services they have rendered by their joint researches on the radiation phenomena discovered by Professor Henri Becquerel"*.

Figure. 1903 Nobel Prize in physics winners

Radiation is energy as particles or electromagnetic waves, like sunshine




Figure. Chart of the electromagnetic spectrum

Radiation is energy as particles or electromagnetic waves




Figure. Radiation shielding materials




Figure. Radiation shielding materials

Radiation is energy as particles or electromagnetic waves

Charged particles are –

- (1) protons (+)
- (2) alpha (++)
- (3) beta (+/-)
- (4) heavy ions (varying)
- (5) U, Pu (3+, 4+, 5+, 6+)

Neutrons have no charge, but are also ionizing

Ionizing rays are x-rays, g-rays, cosmic rays

Ernest Rutherford discovered the three main kinds of radioactive decay (Chemistry Nobel 08)




Figure. Types of radioactive decay

Alpha particle decay is the ejection of a helium atom from a heavy nucleus



$$Q = 4.268 \text{ MeV} \quad (15)$$

Sometimes nucleus decays to excited state and then emits a gamma ray (for any kind of radiation)

Discrete energy spectrum

Some % decay at certain energy

Radium-226 decays to radon-222




Figure. Radium-226 decay

β^- particle decay occurs for unstable nuclei with excessive neutrons



The neutron is converted to a proton and a beta negative (electron) and anti-neutrino (momentum) are ejected

Sometimes nucleus decays to excited state and then emits a gamma ray (for any kind of radiation)

Gold-198 decays to mercury-198




Figure. Gold-198 decay

β^+ particle decay occurs for unstable nuclei with deficient neutrons



The proton is converted to a neutron and a beta positive (positron) and neutrino (momentum) are ejected

Sometimes nucleus decays to excited state and then emits a gamma ray (for any kind of radiation)

Sodium-22 decays to neon-22




Figure. Sodium-22 decay

In both forms of beta decay, the energies are exhibited by a continuous energy spectrum




Figure. Beta decay energy spectrum

$$\overline{E}(\beta^-) = 0.3E_{MAX} \quad (18)$$

$$\overline{E}(\beta^+) = 0.4E_{MAX} \quad (19)$$

Electron capture also occurs for unstable nuclei with deficient neutrons



The K shell electron (not free) is absorbed by the nucleus to convert the proton to a neutron

Sometimes nucleus decays to excited state and then emits a gamma ray (for any kind of radiation)

Electron is captured by the nucleus




Figure. Notional electron capture

Electron capture and β^+ are competing processes



$$Q = 2.842 \text{ MeV} \quad (23)$$

$Q > 1.022 \text{ MeV}$ positron favored

$Q < 1.022 \text{ MeV}$ electron capture favored

Why?

γ -ray emission is not a decay mode, but superfast de-excitation of the nucleus

Interactions of gamma rays with matter is a very important branch of nuclear physics

Chart of the nuclides

Chart of the nuclides wide shot




Figure. Chart of the nuclides

Chart of the nuclides example

Ra 224 3.66 d α 5.6854 5.4486... γ 241..., C14 σ 12.0	Ra 225 14.8 d β^- 0.3, 0.4 γ 40 e^-	Ra 226 1600 a α 4.7843, 4.601... γ 186... C14 σ 12.8 σ_T < 5E-5
Fr 223 21.8 m β^- 1.1... α 5.34 γ 50, 80, 235...	Fr 224 3.3 m β^- 2.6, 2.8... γ 216, 132, 837 1341...	Fr 225 4.0 m β^- 1.6... γ 182, 32, 225 200...
Rn 222 3.825 d α 5.48948... γ (510) σ 0.74	Rn 223 23.2 m β^- γ 593, 417, 636 655...	Rn 224 1.78 h β^- γ 261, 266...

Figure. Chart of the nuclides example

Decay chains

Thorium decays to lead




Figure. Thorium decay chain

Neptunium decays to thallium




Figure. Neptunium decay chain

Uranium-238 decays to lead




Figure. Uranium-238 decay chain

Uranium-235 decays to lead




Figure. Uranium-235 decay chain

Half life

A radioactive isotope decays with a unique characteristic time

Decay is stochastic, characterized by Poisson distribution

So the decay of any one radionuclide cannot be predicted

The probability per time that a nucleus will decay is a constant

Decay is described by a Poisson process

Poisson process notes

Occurrences are randomly distributed in time

What is the 'occurrence' here?

Homogeneous, long term decay rate is constant

$$\mu \equiv \lambda \Delta t \quad (24)$$

$$P(n) = e^{-\mu} \cdot \frac{\mu^n}{n!} \quad (25)$$

Defintion of expected value and variance on the OER

A radioactive isotope decays with a unique characteristic time

$n(t)$ atoms at t have decayed in dt

$$\lambda n(t)dt \quad (26)$$

The number of atoms that decay on average in the interval $t, (t + dt)$ can be expressed as –

$$-dn(t) = \lambda n(t)dt \quad (27)$$

You should be able to solve this for $n(0) = n_0$

Radionuclides are characterized by half life

$$t_{\frac{1}{2}} = \frac{\ln 2}{\lambda} \quad (28)$$

Half life can be derived from the decay law

A radionuclide is 'negligible' after 10 half-lives have elapsed




Figure. Notional half life

Why would this be important?

How was half-life discovered?

Rutherford discovered alpha and beta particle decay

In the course of, he noticed that there was a characteristic time

Basically observed decay time was the same for each atom

Experiments with thorium led to the discovery of radon

Then observing the radon gas led to the decay/half-life (65 s)

Gold-198 graphical half life




Figure. Gold-198 half life

Engineers like straight lines




Figure. Gold-198 half life on a log scale

Slope of the line is the decay constant



Acronyms I

OER Online Educational Resource.

UI University of Idaho.

UIIF Idaho Falls Center for Higher Education.