

L2 Sets and mathematical proofs

1ma406/1ma901 Linear algebra

Jonas Nordqvist

Wason selection task

As we've seen earlier for two propositions A and B we have

$$A \implies B \equiv \neg B \implies \neg A.$$


Wason selection task

As we've seen earlier for two propositions A and B we have

$$A \implies B \equiv \neg B \implies \neg A.$$

Below you see a famous problem called the Wason selection task. The exact formulation in this slide is taken from Wikipedia.

You are shown a set of four cards placed on a table, each of which has a number on one side and a colored patch on the other side. The visible faces of the cards show 3, 8, red and brown. Which card(s) must you turn over in order to test the truth of the proposition that if a card shows an even number on one face, then its opposite face is red?



- Skim
- Identify what is important
- Ask questions
- Careful reading
- Stop periodically to review
- Read statements first – proofs later

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Example: Give a proof by contraposition of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Example: Give a proof by contraposition of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by contradiction. Because the statement $r \wedge \neg r$ is a contradiction whenever r is proposition, we can prove that p is true if we can show that $\neg p \implies r \wedge \neg r$ is true for some proposition r .

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Example: Give a proof by contraposition of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by contradiction. Because the statement $r \wedge \neg r$ is a contradiction whenever r is proposition, we can prove that p is true if we can show that $\neg p \implies r \wedge \neg r$ is true for some proposition r .

Example: Give a proof by contradiction of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Example: Give a proof by contraposition of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by contradiction. Because the statement $r \wedge \neg r$ is a contradiction whenever r is proposition, we can prove that p is true if we can show that $\neg p \implies r \wedge \neg r$ is true for some proposition r .

Example: Give a proof by contradiction of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by cases. A proof by cases must cover all possible cases that arise in a theorem.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Example: Give a proof by contraposition of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by contradiction. Because the statement $r \wedge \neg r$ is a contradiction whenever r is proposition, we can prove that p is true if we can show that $\neg p \implies r \wedge \neg r$ is true for some proposition r .

Example: Give a proof by contradiction of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by cases. A proof by cases must cover all possible cases that arise in a theorem.

Example: Give a proof by cases of the theorem 'If n is an integer, then $n^2 \geq n$ '.

Introduction to mathematical proofs

Direct proof. A direct proof of a conditional statement $p \implies q$ is constructed when the first step is the assumption that p is true; subsequent steps are constructed using rules of inference, with the final step showing that q must also be true.

Example: Give a direct proof of the theorem 'If n is an odd integer, then n^2 is odd'.

Proof by contraposition. Proofs by contraposition make use of the fact that the conditional statement $p \implies q$ is equivalent to its contrapositive, $\neg q \implies \neg p$.

Example: Give a proof by contraposition of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by contradiction. Because the statement $r \wedge \neg r$ is a contradiction whenever r is proposition, we can prove that p is true if we can show that $\neg p \implies r \wedge \neg r$ is true for some proposition r .

Example: Give a proof by contradiction of the theorem 'If n is an integer and $3n + 2$ is odd, then n is odd'.

Proof by cases. A proof by cases must cover all possible cases that arise in a theorem.

Example: Give a proof by cases of the theorem 'If n is an integer, then $n^2 \geq n$ '.

Proof by induction. Left to a course in discrete mathematics.

- i) In the direct method, to show $A \implies B$ we assume A and proceed to show B .
- ii) This is often done in smaller steps where each conclusion is direct, for instance

$$A \implies A_1, \quad A_1 \implies A_2, \quad \dots \quad A_{k-1} \implies A_k, \quad A_k \implies B.$$

- i) State what you will be proving, e.g. $A \iff B$
- ii) State the assumption A and prove that this implies B , i.e. $A \implies B$.
- iii) State the assumption B and prove that this implies A , i.e. $B \implies A$.

- i) State that you will make a proof by cases.
- ii) Conclude what cases you will use, *e.g.* Case 1, ..., Case n
- iii) Conclude the proof for Case 1
- iv) Conclude the proof for Case 2
- ...) ...
- $n + ii$) Conclude the proof for Case n

- i) State that you are assuming the statement is false. Seasoned mathematicians will recognize that the proof will be by contradiction.
- ii) Write out what the statement being false means using negation.
- iii) Work out what this would imply until you find a contradiction.
- iv) Announce that a contradiction has been found.

- i) State that you will make a proof by contraposition of the statement, e.g. we want to prove that $\neg B \implies \neg A$.
- ii) Use any of the previous mentioned techniques to prove the implication.

Examples of mathematical proofs

Give a direct proof of the following proposition:

Theorem

If n is odd, then n^2 is odd.

Examples of mathematical proofs

Give a direct proof of the following proposition:

Theorem

If n is odd, then n^2 is odd.

Give a proof by contraposition of the following propositions:

Theorem

If n^2 is even, then n is even.

Theorem

If $x^2 - 6x + 5$ is even then x is odd.

Examples of mathematical proofs

Give a direct proof of the following proposition:

Theorem

If n is odd, then n^2 is odd.

Give a proof by contraposition of the following propositions:

Theorem

If n^2 is even, then n is even.

Theorem

If $x^2 - 6x + 5$ is even then x is odd.

Give a proof by contradiction of the following proposition:

Theorem

Suppose a, b be integers. If $a + b \geq 19$, then $a \geq 10$ or $b \geq 10$.

Examples of mathematical proofs

Give a direct proof of the following proposition:

Theorem

If n is odd, then n^2 is odd.

Give a proof by contraposition of the following propositions:

Theorem

If n^2 is even, then n is even.

Theorem

If $x^2 - 6x + 5$ is even then x is odd.

Give a proof by contradiction of the following proposition:

Theorem

Suppose a, b be integers. If $a + b \geq 19$, then $a \geq 10$ or $b \geq 10$.

Give a proof by cases of the following proposition:

Theorem

Let n be an integer. Then $n^3 - n$ is a multiple of 3.

Sets and notation

Definition

A *set* is an unordered collection of objects, called *elements* or *members* of the set. A set is said to *contain* its elements. We write $a \in A$ to denote that a is an element of the set A . The notation $a \notin A$ denotes that A is not an element of the set A .

Sets and notation

Definition

A *set* is an unordered collection of objects, called *elements* or *members* of the set. A set is said to *contain* its elements. We write $a \in A$ to denote that a is an element of the set A . The notation $a \notin A$ denotes that A is not an element of the set A .

Let \mathbb{N} denote the set of positive integers.

Let A be the set of all positive integers less than or equal to 10.

Sets and notation

Definition

A *set* is an unordered collection of objects, called *elements* or *members* of the set. A set is said to *contain* its elements. We write $a \in A$ to denote that a is an element of the set A . The notation $a \notin A$ denotes that A is not an element of the set A .

Let \mathbb{N} denote the set of positive integers.

Let A be the set of all positive integers less than or equal to 10.

Roster notation

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} \text{ or}$$

$$A = \{1, 2, \dots, 10\}$$

Sets and notation

Definition

A *set* is an unordered collection of objects, called *elements* or *members* of the set. A set is said to *contain* its elements. We write $a \in A$ to denote that a is an element of the set A . The notation $a \notin A$ denotes that A is not an element of the set A .

Let \mathbb{N} denote the set of positive integers.

Let A be the set of all positive integers less than or equal to 10.

Roster notation

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} \text{ or}$$

$$A = \{1, 2, \dots, 10\}$$

Set builder notation

$$A = \{x \in \mathbb{N} \mid x \leq 10\}$$

Sets and notation

Definition

A *set* is an unordered collection of objects, called *elements* or *members* of the set. A set is said to *contain* its elements. We write $a \in A$ to denote that a is an element of the set A . The notation $a \notin A$ denotes that A is not an element of the set A .

Let \mathbb{N} denote the set of positive integers.

Let A be the set of all positive integers less than or equal to 10.

Roster notation

$$A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\} \text{ or}$$

$$A = \{1, 2, \dots, 10\}$$

Set builder notation

$$A = \{x \in \mathbb{N} \mid x \leq 10\}$$

The set builder notation reads 'A is the set of all $x \in \mathbb{N}$ such that $x \leq 10$.'

Facts: $5 \in A$ and $11 \notin A$

These are also sets $\{\text{red}, \text{blue}\}$, $\{1, 2, \text{rain}, \pi\}$, $\{1, 2, \{2, 3\}, 3, \{3\}\}$.

However, we are primarily interested in number sets.

Important sets

Notation	set	elements
\emptyset	empty set	the set that contains no elements
\mathbb{Z}	integers	$\{\dots, -2, -1, 0, 1, 2, \dots\}$
\mathbb{Z}^+	non-negative integers	$\{0, 1, 2, 3, \dots\}$
\mathbb{N}	natural numbers	$\{1, 2, 3, \dots\}$
\mathbb{Q}	rational numbers	$\{\frac{p}{q} p \in \mathbb{Z}, q \in \mathbb{Z}, \text{ and } q \neq 0\}$
\mathbb{R}	real numbers	'decimal numbers' like $1, -\pi, 0.33, e, \sqrt{2}$
$\mathbb{R}_{>0}$	positive real numbers	$\{x \in \mathbb{R} x > 0\}$
\mathbb{C}	complex numbers	$\{a + ib a, b \in \mathbb{R} \text{ and } i = \sqrt{-1}\}$

Important sets

Notation	set	elements
\emptyset	empty set	the set that contains no elements
\mathbb{Z}	integers	$\{\dots, -2, -1, 0, 1, 2, \dots\}$
\mathbb{Z}^+	non-negative integers	$\{0, 1, 2, 3, \dots\}$
\mathbb{N}	natural numbers	$\{1, 2, 3, \dots\}$
\mathbb{Q}	rational numbers	$\{\frac{p}{q} p \in \mathbb{Z}, q \in \mathbb{Z}, \text{ and } q \neq 0\}$
\mathbb{R}	real numbers	'decimal numbers' like $1, -\pi, 0.33, e, \sqrt{2}$
$\mathbb{R}_{>0}$	positive real numbers	$\{x \in \mathbb{R} x > 0\}$
\mathbb{C}	complex numbers	$\{a + ib a, b \in \mathbb{R} \text{ and } i = \sqrt{-1}\}$

Note that some books reverse the definitions of \mathbb{Z}^+ and \mathbb{N} .

Important sets

Notation	set	elements
\emptyset	empty set	the set that contains no elements
\mathbb{Z}	integers	$\{\dots, -2, -1, 0, 1, 2, \dots\}$
\mathbb{Z}^+	non-negative integers	$\{0, 1, 2, 3, \dots\}$
\mathbb{N}	natural numbers	$\{1, 2, 3, \dots\}$
\mathbb{Q}	rational numbers	$\{\frac{p}{q} p \in \mathbb{Z}, q \in \mathbb{Z}, \text{ and } q \neq 0\}$
\mathbb{R}	real numbers	'decimal numbers' like $1, -\pi, 0.33, e, \sqrt{2}$
$\mathbb{R}_{>0}$	positive real numbers	$\{x \in \mathbb{R} x > 0\}$
\mathbb{C}	complex numbers	$\{a + ib a, b \in \mathbb{R} \text{ and } i = \sqrt{-1}\}$

Note that some books reverse the definitions of \mathbb{Z}^+ and \mathbb{N} .

Intervals on the real line. When a and b are real numbers such that $a < b$ we write

$[a, b] = \{x \in \mathbb{R} a \leq x \leq b\}$	the closed interval from a to b
$[a, b) = \{x \in \mathbb{R} a \leq x < b\}$	half open interval
$(a, b] = \{x \in \mathbb{R} a < x \leq b\}$	half open interval
$(a, b) = \{x \in \mathbb{R} a < x < b\}$	the open interval from a to b

The set $C[a, b]$ of all continuous real valued functions on the closed interval $[a, b]$.

Definition (Subset and set equality)

- A set A is said to be a **subset** of a set B , denoted $A \subseteq B$, if every element appearing in A also appears in B . That is,

$$x \in A \subseteq B \iff x \in A \implies x \in B.$$

For example, $\mathbb{Z} \subseteq \mathbb{Q}$

Definition (Subset and set equality)

- A set A is said to be a **subset** of a set B , denoted $A \subseteq B$, if every element appearing in A also appears in B . That is,

$$x \in A \subseteq B \iff x \in A \implies x \in B.$$

For example, $\mathbb{Z} \subseteq \mathbb{Q}$

- A set A is said to be a **equal** to a set B , denoted $A = B$, if every element appearing in A also appears in B , and the converse. That is,

$$x \in A = B \iff A \subseteq B \text{ AND } B \subseteq A.$$

For example, $\{1, 2, 3\} = \{2, 1, 3\}$.

Definition (Subset and set equality)

- A set A is said to be a **subset** of a set B , denoted $A \subseteq B$, if every element appearing in A also appears in B . That is,

$$x \in A \subseteq B \quad \Longleftrightarrow \quad x \in A \implies x \in B.$$

For example, $\mathbb{Z} \subseteq \mathbb{Q}$

- A set A is said to be **equal** to a set B , denoted $A = B$, if every element appearing in A also appears in B , and the converse. That is,

$$x \in A = B \quad \Longleftrightarrow \quad A \subseteq B \text{ AND } B \subseteq A.$$

For example, $\{1, 2, 3\} = \{2, 1, 3\}$.

- A set A is said to be a **proper subset** of a set B , denoted $A \subset B$, if every element appearing in A also appears in B but $A \neq B$. That is,

$$x \in A \subset B \quad \Longleftrightarrow \quad (x \in A \implies x \in B) \text{ AND } (A \neq B).$$

For example, $\mathbb{Z} \subset \mathbb{Q}$

Cardinality

Let A be a set. If A contains exactly n distinct elements where n is a nonnegative integer, we say that A is a **finite set** and that n is the **cardinality** of A , denoted $|A|$. That is $|A| = n$.

Cardinality

Let A be a set. If A contains exactly n distinct elements where n is a nonnegative integer, we say that A is a **finite set** and that n is the **cardinality** of A , denoted $|A|$. That is $|A| = n$.

Power sets

Given a set A , the **power set** of A is the set of all subsets of A , denoted by $\mathcal{P}(A)$. So

$$B \in \mathcal{P}(A) \iff B \subseteq A.$$

Example:

Cardinality

Let A be a set. If A contains exactly n distinct elements where n is a nonnegative integer, we say that A is a **finite set** and that n is the **cardinality** of A , denoted $|A|$. That is $|A| = n$.

Power sets

Given a set A , the **power set** of A is the set of all subsets of A , denoted by $\mathcal{P}(A)$. So

$$B \in \mathcal{P}(A) \iff B \subseteq A.$$

Example: If $X = \{1, 2\}$, then the elements of $\mathcal{P}(X)$ are \emptyset , $\{1\}$, $\{2\}$, and $\{1, 2\}$. That is

$$\mathcal{P}(X) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}.$$

Cardinality

Let A be a set. If A contains exactly n distinct elements where n is a nonnegative integer, we say that A is a **finite set** and that n is the **cardinality** of A , denoted $|A|$. That is $|A| = n$.

Power sets

Given a set A , the **power set** of A is the set of all subsets of A , denoted by $\mathcal{P}(A)$. So

$$B \in \mathcal{P}(A) \iff B \subseteq A.$$

Example: If $X = \{1, 2\}$, then the elements of $\mathcal{P}(X)$ are \emptyset , $\{1\}$, $\{2\}$, and $\{1, 2\}$. That is

$$\mathcal{P}(X) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}.$$

Proposition

Let A be a finite set of cardinality n . Then the cardinality of its power set $|\mathcal{P}(A)| = 2^n$.

Ordered n -tuples

A **ordered n -tuple** (a_1, a_2, \dots, a_n) is the ordered collection that has a_1 as its first element, a_2 as its second element, \dots , and a_n as its n th element. We say that two ordered n -tuples (a_1, a_2, \dots, a_n) and (b_1, b_2, \dots, b_n) are equal if and only if $a_i = b_i$ for every $i = 1, 2, \dots, n$.

Ordered n -tuples

A **ordered n -tuple** (a_1, a_2, \dots, a_n) is the ordered collection that has a_1 as its first element, a_2 as its second element, \dots , and a_n as its n th element. We say that two ordered n -tuples (a_1, a_2, \dots, a_n) and (b_1, b_2, \dots, b_n) are equal if and only if $a_i = b_i$ for every $i = 1, 2, \dots, n$.

Cartesian product of two sets

Let A and B be two sets. The **Cartesian product** of A and B , denoted by $A \times B$, is the set of all ordered pairs (a, b) , where $a \in A$ and $b \in B$. That is

$$A \times B = \{(a, b) | a \in A \wedge b \in B\}.$$

Ordered n -tuples

A **ordered n -tuple** (a_1, a_2, \dots, a_n) is the ordered collection that has a_1 as its first element, a_2 as its second element, \dots , and a_n as its n th element. We say that two ordered n -tuples (a_1, a_2, \dots, a_n) and (b_1, b_2, \dots, b_n) are equal if and only if $a_i = b_i$ for every $i = 1, 2, \dots, n$.

Cartesian product of two sets

Let A and B be two sets. The **Cartesian product** of A and B , denoted by $A \times B$, is the set of all ordered pairs (a, b) , where $a \in A$ and $b \in B$. That is

$$A \times B = \{(a, b) | a \in A \wedge b \in B\}.$$

Example 1: $S = \{1, 2\}$ and $T = \{a, b, c\}$. Then

$$S \times T = \{(1, a), (1, b), (1, c), (2, a), (2, b), (2, c)\}$$

The Cartesian coordinate plane \mathbb{R}^2 and in \mathbb{R}^3

Let \mathbb{R} be the set of real numbers and put $\mathbb{R}^2 := \mathbb{R} \times \mathbb{R}$. That is

$$\mathbb{R}^2 := \{(a, b) | a \in \mathbb{R} \wedge b \in \mathbb{R}\}.$$

Further, we have $\mathbb{R}^3 = \mathbb{R} \times \mathbb{R} \times \mathbb{R}$, *i.e.*

$$\mathbb{R}^3 := \{(a, b, c) | a \in \mathbb{R} \wedge b \in \mathbb{R} \wedge c \in \mathbb{R}\}.$$

Sets of the form \mathbb{R}^n for some positive integer n is the corner stone in our course in Linear algebra.

Combining Sets – Set Operations

There are several basic ways to combine sets. For example, put

$X := \{1, 2, 3\}$; $Y := \{2, 3, 4\}$.

Combining Sets – Set Operations

There are several basic ways to combine sets. For example, put

$X := \{1, 2, 3\}$; $Y := \{2, 3, 4\}$.

- The **union** of sets A and B , denoted $A \cup B$, includes exactly the elements appearing in A or B or both. That is,

$$x \in A \cup B \iff x \in A \text{ OR } x \in B.$$

So, $X \cup Y = \{1, 2, 3, 4\}$

Combining Sets – Set Operations

There are several basic ways to combine sets. For example, put

$X := \{1, 2, 3\}$; $Y := \{2, 3, 4\}$.

- The **union** of sets A and B , denoted $A \cup B$, includes exactly the elements appearing in A or B or both. That is,

$$x \in A \cup B \iff x \in A \text{ OR } x \in B.$$

So, $X \cup Y = \{1, 2, 3, 4\}$

- The **intersection** of A and B , denoted $A \cap B$, consists of all elements that appear in both A and B . That is,

$$x \in A \cap B \iff x \in A \text{ AND } x \in B.$$

So, $X \cap Y = \{2, 3\}$

Combining Sets – Set Operations

There are several basic ways to combine sets. For example, put

$X := \{1, 2, 3\}$; $Y := \{2, 3, 4\}$.

- The **union** of sets A and B , denoted $A \cup B$, includes exactly the elements appearing in A or B or both. That is,

$$x \in A \cup B \iff x \in A \text{ OR } x \in B.$$

So, $X \cup Y = \{1, 2, 3, 4\}$

- The **intersection** of A and B , denoted $A \cap B$, consists of all elements that appear in both A and B . That is,

$$x \in A \cap B \iff x \in A \text{ AND } x \in B.$$

So, $X \cap Y = \{2, 3\}$

- The **set difference** of A and B , denoted $A - B$, consists of all elements that are in A , but not in B . That is,

$$x \in A - B \iff x \in A \text{ AND } x \notin B.$$

So, $X - Y = \{1\}$.

Combining Sets – Set Operations

There are several basic ways to combine sets. For example, put

$X := \{1, 2, 3\}$; $Y := \{2, 3, 4\}$.

- The **union** of sets A and B , denoted $A \cup B$, includes exactly the elements appearing in A or B or both. That is,

$$x \in A \cup B \iff x \in A \text{ OR } x \in B.$$

So, $X \cup Y = \{1, 2, 3, 4\}$

- The **intersection** of A and B , denoted $A \cap B$, consists of all elements that appear in both A and B . That is,

$$x \in A \cap B \iff x \in A \text{ AND } x \in B.$$

So, $X \cap Y = \{2, 3\}$

- The **set difference** of A and B , denoted $A - B$, consists of all elements that are in A , but not in B . That is,

$$x \in A - B \iff x \in A \text{ AND } x \notin B.$$

So, $X - Y = \{1\}$.

- Given a set A and some universal set U such that $A \subseteq U$, the **complement** of A in U , denoted A^c , consists of all elements in $U - A$. That is,

$$A^c := U - A.$$

So, $X^c = \{4, 5\}$ is the complement of X in $U = \{1, 2, 3, 4, 5\}$.

