

SOLUTIONS SHEET 3

YANNIS BÄHNI

Exercise 1. Let $D \subseteq \mathbb{C}$ be non-empty and open in \mathbb{C} and $f_1, f_2 : D \rightarrow \mathbb{C}$ be real differentiable. Fix some $z_0 \in D$. Since f_1 and f_2 are real differentiable in z_0 there exists $\varphi_1, \varphi_2, \psi_1, \psi_2 : D \rightarrow \mathbb{C}$ continuous at z_0 such that

$$f_1(z) = f_1(z_0) + (z - z_0)\varphi_1(z) + (\bar{z} - \bar{z}_0)\psi_1(z) \quad (1)$$

$$f_2(z) = f_2(z_0) + (z - z_0)\varphi_2(z) + (\bar{z} - \bar{z}_0)\psi_2(z) \quad (2)$$

for all $z \in D$.

(i) Let $a, b \in \mathbb{C}$. Multiplying (1) by a , (2) by b and adding both equations yields

$$af_1(z) + bf_2(z) = af_1(z_0) + bf_2(z_0) + (z - z_0)(a\varphi_1(z) + b\varphi_2(z)) + (\bar{z} - \bar{z}_0)(a\psi_1(z) + b\psi_2(z)) \quad (3)$$

for all $z \in D$. Clearly, $a\varphi_1 + b\varphi_2$ and $a\psi_1 + b\psi_2$ are continuous functions in z_0 and from (3) we deduce

$$\frac{\partial(af_1 + bf_2)}{\partial z}(z_0) = a \frac{\partial f_1}{\partial z}(z_0) + b \frac{\partial f_2}{\partial z}(z_0) \quad (4)$$

and

$$\frac{\partial(af_1 + bf_2)}{\partial \bar{z}}(z_0) = a \frac{\partial f_1}{\partial \bar{z}}(z_0) + b \frac{\partial f_2}{\partial \bar{z}}(z_0). \quad (5)$$

Since $z_0 \in D$ was arbitrary, we conclude

$$\frac{\partial(af_1 + bf_2)}{\partial z} = a \frac{\partial f_1}{\partial z} + b \frac{\partial f_2}{\partial z} \quad \text{and} \quad \frac{\partial(af_1 + bf_2)}{\partial \bar{z}} = a \frac{\partial f_1}{\partial \bar{z}} + b \frac{\partial f_2}{\partial \bar{z}}. \quad (6)$$

(ii) Multiplying (1) and (2) yields

$$\begin{aligned} f_1 f_2 &= f_1(z_0)f_2(z_0) + (z - z_0)[\varphi_1 f_2(z_0) + f_1(z_0)\varphi_2 + (z - z_0)\varphi_1\varphi_2 + (\bar{z} - \bar{z}_0)\psi_1\varphi_2] \\ &\quad + (\bar{z} - \bar{z}_0)[\psi_1 f_2(z_0) + f_1(z_0)\psi_2 + (z - z_0)\psi_2\varphi_1 + (\bar{z} - \bar{z}_0)\psi_1\psi_2] \end{aligned}$$

where the argument z is omitted. Clearly, the two functions in the square brackets are continuous at z_0 and evaluating them at z_0 yields

$$\frac{\partial(f_1 f_2)}{\partial z}(z_0) = \frac{\partial f_1}{\partial z}(z_0)f_2(z_0) + f_1(z_0)\frac{\partial f_2}{\partial z}(z_0) \quad (7)$$

and

$$\frac{\partial(f_1 f_2)}{\partial \bar{z}}(z_0) = \frac{\partial f_1}{\partial \bar{z}}(z_0)f_2(z_0) + f_1(z_0)\frac{\partial f_2}{\partial \bar{z}}(z_0). \quad (8)$$

Since $z_0 \in D$ was arbitrary, we conclude

$$\frac{\partial(f_1 f_2)}{\partial z} = \frac{\partial f_1}{\partial z} f_2 + f_1 \frac{\partial f_2}{\partial z} \quad \text{and} \quad \frac{\partial(f_1 f_2)}{\partial \bar{z}} = \frac{\partial f_1}{\partial \bar{z}} f_2 + f_1 \frac{\partial f_2}{\partial \bar{z}}. \quad (9)$$

(iii) Conjugating (1) yields

$$\overline{f_1}(z) = \overline{f_1}(z_0) + (\bar{z} - \bar{z}_0)\overline{\varphi_1}(z) + (z - z_0)\overline{\psi_1}(z). \quad (10)$$

From (10) we deduce

$$\frac{\partial \overline{f_1}}{\partial \bar{z}}(z_0) = \overline{\varphi_1}(z_0) = \overline{\frac{\partial f_1}{\partial z}}(z_0) \quad (11)$$

since φ_1 and ψ_1 are also continuous at z_0 . Taking conjugates in (11) and use that $z_0 \in D$ was arbitrary finally yields

$$\overline{\frac{\partial f_1}{\partial \bar{z}}} = \frac{\partial f_1}{\partial z}. \quad (12)$$

(iv) This follows directly from

$$z = z_0 + (z - z_0) \quad \text{and} \quad \bar{z} = \bar{z}_0 + (\bar{z} - \bar{z}_0). \quad (13)$$