

YANNIS BÄHNI

SOLUTION
BOOK TO
INTRODUCTION
TO SMOOTH
MANIFOLDS BY
JOHN M. LEE

YANNIS BÄHNI

SOLUTION
BOOK TO
INTRODUCTION
TO SMOOTH
MANIFOLDS BY
JOHN M. LEE

Contents

Chapter 1. Smooth Maps	1
1 Smooth Functions and Smooth Maps	1
1.1 Smooth Functions on Manifolds	1
Chapter 2. Tangent Vectors	3
1 Tangent Vectors	3
1.1 Tangent Vectors on Manifolds	3
Chapter 3. Vector Fields	5
1 Vector Fields on Manifolds	5
Chapter 4. The Cotangent Bundle	6
1 Line Integrals	6
1.1 The Winding Number*	6
Chapter 5. Tensors	7
1 Pullbacks of Tensor Fields	7
Chapter 6. Orientations	8
1 Orientations of Vector Spaces	8
Appendix. Bibliography	9

CHAPTER 1

Smooth Maps

1. Smooth Functions and Smooth Maps

1.1. Smooth Functions on Manifolds. We follow the terminology established in [Gri07, p. 515].

Exercise 1.1. Let M be a smooth manifold. $\mathcal{C}^\infty(M)$ is an associative and commutative \mathbb{R} -algebra with identity under the usual pointwise defined operations.

Solution 1.1. First we show that $\mathcal{C}^\infty(M)$ is a real vector space. Since $\mathcal{C}^\infty(M) \subseteq \mathbb{R}^M$ it is enough to show that $\mathcal{C}^\infty(M)$ is a linear subspace of the real vector space \mathbb{R}^M . Clearly, $\mathcal{C}^\infty(M) \neq \emptyset$, since $\chi_M \in \mathcal{C}^\infty(M)$. Indeed, for $p \in M$ we find a chart (U, φ) such that $p \in U$ and the composition $\chi_M \circ \varphi^{-1} : \varphi(U) \rightarrow \mathbb{R}$ is clearly the function $\chi_{\varphi(U)}$, which is smooth since it is constant. Now let $f, g \in \mathcal{C}^\infty(M)$, $\lambda \in \mathbb{R}$ and $p \in M$. By definition, there exist charts (U, φ) , (V, ψ) such that $f \circ \varphi^{-1}$ and $g \circ \psi^{-1}$ are smooth. Now consider the chart $(U \cap V, \varphi)$. Then by

$$(\lambda f + g) \circ \varphi^{-1} = \lambda(f \circ \varphi^{-1}) + (g \circ \varphi^{-1}) = \lambda(f \circ \varphi^{-1}) + ((g \circ \psi^{-1}) \circ (\psi \circ \varphi^{-1}))$$

we have that $\lambda f + g \in \mathcal{C}^\infty(M)$. Hence $\mathcal{C}^\infty(M)$ is a real vector space.

Now define a product map $\cdot : \mathcal{C}^\infty(M) \times \mathcal{C}^\infty(M) \rightarrow \mathcal{C}^\infty(M)$ by pointwise multiplication. Indeed, similar to the previous reasoning, by

$$(f \cdot g) \circ \varphi^{-1} = (f \circ \varphi^{-1}) \cdot (g \circ \varphi^{-1}) = (f \circ \varphi^{-1}) \cdot ((g \circ \psi^{-1}) \circ (\psi \circ \varphi^{-1}))$$

we have that $f \cdot g$ is smooth. Let $f, g, h \in \mathcal{C}^\infty(M)$ and $\lambda \in \mathbb{R}$. Then for $p \in M$

$$\begin{aligned} ((\lambda f + g) \cdot h)(p) &= (\lambda f + g)(p)h(p) \\ &= (\lambda f(p) + g(p))h(p) \\ &= \lambda f(p)h(p) + g(p)h(p) \\ &= \lambda(f \cdot h)(p) + (g \cdot h)(p) \\ &= (\lambda(f \cdot h))(p) + (g \cdot h)(p) \\ &= (\lambda(f \cdot h) + (g \cdot h))(p) \end{aligned}$$

shows that \cdot is bilinear in the first argument. A similar computation shows that \cdot is bilinear. By

$$\begin{aligned} ((f \cdot g) \cdot h)(p) &= (f \cdot g)(p)h(p) \\ &= f(p)g(p)h(p) \\ &= f(p)(g \cdot h)(p) \\ &= (f \cdot (g \cdot h))(p) \end{aligned}$$

we see that \cdot is associative. Furthermore by

$$(f \cdot g)(p) = f(p)g(p) = g(p)f(p) = (g \cdot f)(p)$$

we see that \cdot is commutative. Finally, the identity element is given by χ_M since

$$(\chi_M \cdot f)(p) = \chi_M(p)f(p) = 1f(p) = f(p).$$

CHAPTER 2

Tangent Vectors

1. Tangent Vectors

1.1. Tangent Vectors on Manifolds.

Exercise 1.1. Let M be a smooth manifold and $p \in M$. The set of all derivations at p , written $T_p M$, is a real vector space under the usual pointwise defined operations.

Solution 1.1. Clearly $T_p M \subseteq L(\mathcal{C}^\infty(M); \mathbb{R})$ and thus it is enough to show that $T_p M$ is a linear subspace of $L(\mathcal{C}^\infty(M); \mathbb{R})$ (see [Lee13, p. 626]). We have $T_p M \neq \emptyset$, since $0 \in T_p M$ defined by $f \mapsto 0$. Let $u, v \in T_p M$, $\lambda \in \mathbb{R}$ and $f, g \in \mathcal{C}^\infty(M)$. Then by

$$\begin{aligned} (\lambda u + v)(fg) &= \lambda u(fg) + v(fg) \\ &= f(p)(\lambda u(g) + v(g)) + g(p)(\lambda u(f) + v(f)) \\ &= f(p)(\lambda u + v)(g) + g(p)(\lambda u + v)(f) \end{aligned}$$

we have that $\lambda u + v \in T_p M$.

Exercise 1.2. Suppose M is a smooth manifold. Let $p \in M$, $v \in T_p M$ and $f \in \mathcal{C}^\infty(M)$. If f is constant, then $v(f) = 0$.

Solution 1.2. First assume that $f = \chi_M$. Then

$$v(\chi_M) = v(\chi_M \cdot \chi_M) = f(p)v(\chi_M) + f(p)v(\chi_M) = v(\chi_M) + v(\chi_M) \quad (1)$$

implies that $v(f) = 0$. Hence if $f = \lambda \chi_M$ for $\lambda \in \mathbb{R}$, the \mathbb{R} -linearity of v implies that

$$v(f) = v(\lambda \chi_M) = \lambda v(\chi_M) = 0. \quad (2)$$

Exercise 1.3 (Properties of Differentials). Let M , N and P be smooth manifolds, let $F : M \rightarrow N$ and $G : N \rightarrow P$ be smooth maps, and let $p \in M$.

- (a) $dF_p : T_p M \rightarrow T_{F(p)} N$ is \mathbb{R} -linear.
- (b) $d(G \circ F)_p = dG_{F(p)} \circ dF_p$.
- (c) $d(id_M)_p = id_{T_p M}$.
- (d) If F is a diffeomorphism, then dF_p is an isomorphism and $(dF_p)^{-1} = d(F^{-1})_{F(p)}$.

Solution 1.3. Let $u, v \in T_p M$, $\lambda \in \mathbb{R}$ and $f \in \mathcal{C}^\infty(N)$. Then

$$\begin{aligned} dF_p(\lambda u + v)(f) &= (\lambda u + v)(f \circ F) \\ &= \lambda u(f \circ F) + v(f \circ F) \\ &= \lambda dF_p(u)(f) + dF_p(v)(f). \end{aligned}$$

This shows part (a). Let $v \in T_p M$ and $f \in \mathcal{C}^\infty(P)$. Then

$$\begin{aligned} d(G \circ F)_p(v)(f) &= v(f \circ (G \circ F)) \\ &= v((f \circ G) \circ F) \\ &= dF_p(f \circ G) \\ &= dG_{F(p)}(dF_p(v))(f) \\ &= (dG_{F(p)} \circ dF_p)(v)(f). \end{aligned}$$

This shows part (b). Part (c) is immediate by

$$d(id_M)_p(v)(f) = v(f \circ id_M) = v(f)$$

for $f \in \mathcal{C}^\infty(M)$. Finally, part (b) and (c) yields

$$dF_p \circ d(F^{-1})_{F(p)} = d(F \circ F^{-1})_{F(p)} = d(id_N)_{F(p)} = id_{T_{F(p)}N}$$

and

$$d(F^{-1})_{F(p)} \circ dF_p = d(F^{-1} \circ F)_p = d(id_M)_p = id_{T_p M}$$

which shows that dF_p is bijective with inverse $(dF_p)^{-1} = d(F^{-1})_{F(p)}$ by uniqueness. Since by part (a) dF_p is linear, we have that dF_p is an isomorphism (see [Lee13, p. 622]). This shows part (d).

CHAPTER 3

Vector Fields

1. Vector Fields on Manifolds

Exercise 1.1. Let M be a smooth manifold.

- (a) If $X, Y \in \mathfrak{X}(M)$, $f, g \in \mathcal{C}^\infty(M)$, then $fX + gY \in \mathfrak{X}(M)$.

CHAPTER 4

The Cotangent Bundle

1. Line Integrals

1.1. The Winding Number*.

Definition 1.1 (Winding Number). Let $z_0 \in \mathbb{C}$ and $\gamma : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$ be a piecewise continuously differentiable closed path. Then

$$W(\gamma, z_0) := \frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - z_0} \quad (3)$$

is called the **winding number** of γ around z_0 .

Proposition 1.1. Let $z_0 := x_0 + iy_0 \in \mathbb{C}$ and $\gamma : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$ be a piecewise continuously differentiable closed path. Then

$$W(\gamma, z_0) = \int_{\gamma} \omega \quad (4)$$

where $\omega \in \Omega^1(\mathbb{R}^2 \setminus \{z_0\})$ is given by

$$\omega := \frac{1}{2\pi} \frac{(x - x_0) dy - (y - y_0) dx}{(x - x_0)^2 + (y - y_0)^2}. \quad (5)$$

Proof. This immediately follows from

$$\begin{aligned} \int_{\gamma} \frac{dz}{z - z_0} &= \int_{\gamma} \frac{dx + i dy}{(x + iy) - (x_0 + iy_0)} \\ &= \int_{\gamma} \frac{dx + i dy}{(x - x_0) + i(y - y_0)} \frac{(x - x_0) - i(y - y_0)}{(x - x_0) - i(y - y_0)} \\ &= \int_{\gamma} \frac{(x - x_0) dx + ((x - x_0) dy - (y - y_0) dx) + (y - y_0) dy}{(x - x_0)^2 + (y - y_0)^2} \\ &= \int_{\gamma} \frac{(x - x_0) dx + (y - y_0) dy}{(x - x_0)^2 + (y - y_0)^2} + i \int_{\gamma} \frac{(x - x_0) dy - (y - y_0) dx}{(x - x_0)^2 + (y - y_0)^2} \\ &= i \int_{\gamma} \frac{(x - x_0) dy - (y - y_0) dx}{(x - x_0)^2 + (y - y_0)^2} \end{aligned}$$

by the fundamental theorem for line integrals [Lee13, p. 291] since

$$d \left(\frac{1}{2} \log((x - x_0)^2 + (y - y_0)^2) \right) = \frac{(x - x_0) dx + (y - y_0) dy}{(x - x_0)^2 + (y - y_0)^2}.$$

□

Remark 1.1. By proposition 1.1, the definition of the winding number in complex analysis given by definition 1.1 coincides with the one usually given in algebraic topology (see for example [Ful95, pp. 19–20]).

CHAPTER 5

Tensors

1. Pullbacks of Tensor Fields

Exercise 1.1 (Properties of Tensor Pullbacks). Suppose $F : M \rightarrow N$ is a smooth mapping and A, B are covariant tensor fields on N . Then

(a) $F^*(A \otimes B) = F^*A \otimes F^*B$.

Solution 1.1. Let $p \in M$. Then we have

$$\begin{aligned} (F^*(A \otimes B))_p(v_1, \dots, v_{k+l}) &= (A \otimes B)_{F(p)}(dF_p(v_1), \dots, dF_p(v_{k+l})) \\ &= (A_{F(p)} \otimes B_{F(p)})(dF_p(v_1), \dots, dF_p(v_{k+l})) \\ &= A_{F(p)}(dF_p(v_1), \dots, dF_p(v_k)) B_{F(p)}(dF_p(v_{k+1}), \dots, dF_p(v_{k+l})) \\ &= (F^*A)_p(v_1, \dots, v_k) (F^*B)_p(v_{k+1}, \dots, v_{k+l}) \\ &= ((F^*A)_p \otimes (F^*B)_p)(v_1, \dots, v_{k+l}) \\ &= (F^*A \otimes F^*B)_p(v_1, \dots, v_{k+l}) \end{aligned}$$

for all $v_1, \dots, v_{k+l} \in T_pM$.

CHAPTER 6

Orientations

1. Orientations of Vector Spaces

Exercise 1.1. Let V be a vector space of dimension $n \geq 1$. Define a relation \sim on the set of all ordered bases of V by

$$(v_1, \dots, v_n) \sim (w_1, \dots, w_n) \quad :\Leftrightarrow \quad \det B > 0 \quad (6)$$

where B denotes the transition matrix defined by $w_j = B_j^i v_i$. Show that \sim is an equivalence relation and that $|X/\sim| = 2$.

Solution 1.1. Clearly $(v_1, \dots, v_n) \sim (v_1, \dots, v_n)$ by $v_j = \delta_j^i v_i$. Assume $(v_1, \dots, v_n) \sim (w_1, \dots, w_n)$. Thus B defined by $w_j = B_j^i v_i$ has a positive determinant. But then by $\det(B^{-1}) = (\det(B))^{-1}$ also $\det(B^{-1})$ is positive and $v_j = (B^{-1})_j^i w_i$. Hence $(w_1, \dots, w_n) \sim (v_1, \dots, v_n)$. Lastly, assume that also $(w_1, \dots, w_n) \sim (u_1, \dots, u_n)$. Hence there exists a matrix A such that $u_j = A_j^i w_i$ where $\det(A) > 0$. Thus $u_j = A_j^i w_i = A_j^i (B_i^k v_k) = (A_j^i B_i^k) v_k$ and by $\det(AB) = \det(A) \det(B) > 0$ we get that $(v_1, \dots, v_n) \sim (u_1, \dots, u_n)$. Hence \sim is an equivalence relation.

By [Gri07, p. 335] every vector space has a basis. Let us denote it by (v_1, \dots, v_n) . Therefore

$$(\tilde{v}_1, \dots, \tilde{v}_n) := (-v_1, \dots, v_n)$$

is also a basis for V simply by considering the transition matrix

$$\tilde{B} := \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix}$$

defined by $v_j = \tilde{B}_j^i \tilde{v}_i$. Let (w_1, \dots, w_n) be an ordered basis for V . Let the transition matrix B be defined by $w_j = B_j^i v_i$. If $\det(B) > 0$, we have that

$$(w_1, \dots, w_n) \sim (v_1, \dots, v_n).$$

Otherwise, if $\det(B) < 0$

$$w_j = B_j^i v_i = B_j^i (\hat{B}_i^k \hat{v}_k) = (B_j^i \hat{B}_i^k) \hat{v}_k$$

together with $\det(B\hat{B}) = \det(B) \det(\hat{B}) > 0$ yields

$$(w_1, \dots, w_n) \sim (\tilde{v}_1, \dots, \tilde{v}_n).$$

Since $\det(B) \neq 0$ by the nonsingularity of B , we have that there are exactly two equivalence classes

$$[(v_1, \dots, v_n)]_\sim \quad \text{and} \quad [(-v_1, \dots, v_n)]_\sim.$$

Bibliography

- [Ful95] William Fulton. *Algebraic Topology - A first Course*. Graduate Texts in Mathematics. Springer Science + Business Media, Inc., 1995.
- [Gri07] Pierre Antoine Grillet. *Abstract Algebra*. Graduate Texts in Mathematics. Springer Science + Business Media, LLC, 2007.
- [Lee13] John M. Lee. *Introduction to Smooth Manifolds*. Second Edition. Graduate Texts in Mathematics. Springer, 2013.