

# CLASSICAL FOURIER ANALYSIS: INTERPOLATION OF $L^p$ SPACES

YANNIS BÄHNI

**Abstract.** In this written seminar work I will basically follow the section *Interpolation* in the third edition of the book *Classical Fourier Analysis* by Loukas Grafakos. I will review three basic but important theorems on interpolation of operators on  $L^p$  spaces, starting with the *Riesz-Thorin Interpolation Theorem* based on complex analysis, its generalization, the *Stein-Weiss Interpolation Theorem of Analytic Families of Operators* and finally a theorem based on real methods, the *Marcinkiewicz Interpolation Theorem*. We are mainly concerned with the notion of linear operators as well as slight generalizations of them.

## Contents

List of Figures . . . . .	1
<b>1 Introduction and Basic Definitions . . . . .</b>	<b>2</b>
<b>2 The Complex Method . . . . .</b>	<b>2</b>
2.1 Hadamard's Three Lines Lemma . . . . .	2
2.2 The Riesz-Thorin Interpolation Theorem . . . . .	5
<b>3 Interpolation of Analytic Families of Operators . . . . .</b>	<b>10</b>
3.1 Extension of Hadamard's Three Lines Lemma . . . . .	10
3.1.1 Auxiliary Lemmata . . . . .	10
3.1.2 The Lemma . . . . .	14
3.2 Stein-Weiss Interpolation Theorem of Analytic Families of Operators . . . . .	20
<b>4 The Real Method . . . . .</b>	<b>22</b>
4.1 The Marcinkiewicz Interpolation Theorem . . . . .	23
<b>Appendix A Limit superior and limit inferior revisited . . . . .</b>	<b>29</b>
<b>Appendix B Measure Theory . . . . .</b>	<b>30</b>
<b>References . . . . .</b>	<b>31</b>

## List of Figures

1 Sketch of the setting of Hadamard's three lines lemma. . . . .	3
2 Sketch of the rectangle $\overline{R}$ . . . . .	5

---

(Yannis Bähni) UNIVERSITY OF ZURICH, RÄMISTRASSE 71, 8006 ZURICH

E-mail address: [yannis.baehni@uzh.ch](mailto:yannis.baehni@uzh.ch).

I would like to thank Dr. Chiara Saffirio for many helpful suggestions, Prof. Dr. Benjamin Schlein for his brilliant Analysis I/II/III courses as well as scripts and proof hints and of course Loukas Grafakos, who helped me a lot with understanding his proofs.

**1. Introduction and Basic Definitions.** Suppose  $(p_0, q_0), (p_1, q_1) \in [1, \infty] \times [1, \infty]$  are two pairs of indices and assume that the estimates

$$\|T(f)\|_{L^{q_0}} \leq M_0 \|f\|_{L^{p_0}} \quad \text{and} \quad \|T(f)\|_{L^{q_1}} \leq M_1 \|f\|_{L^{p_1}}$$

hold, where  $T$  is an appropriately chosen operator. Does this imply that

$$\|T(f)\|_{L^q} \leq M \|f\|_{L^p} \quad \text{for other pairs } (p, q) \in [1, \infty]?$$

Those and similar questions will be answered by a tool called *interpolation*, in our case interpolation of  $L^p$  spaces. Using interpolation it is possible to reduce difficult estimates to endpoint estimates and so interpolation can (but not always does) simplify matters. Among the numerous applications of interpolation is by far the shortest proof of *Young's inequality for convolutions* [Gra14, pp. 22–23]. However, there is not *the* interpolation theorem, merely a family of theorems which can be roughly divided into two main categories: *real* and *complex* interpolation methods. Real methods use so called *cut-off* functions to divide the functions in the domain of the operator  $T$  into a bounded and unbounded part and then establish bounds on each of those parts whereas complex interpolation theorems are based upon standard results in complex analysis and are more restrictive on the operator  $T$  in question but yield more natural bounds (even continuous estimates) and will therefore be considered in this task. First we need a rigorous idea of what an appropriately chosen operator means in the context of Lebesgue spaces.

**DEFINITION 1.1.** Let  $(X, \mu)$  and  $(Y, \nu)$  be measure spaces. Further let  $T$  be an operator defined on a linear space of complex-valued measurable functions on  $X$  and taking values in the set of all complex-valued, finite almost everywhere, measurable functions on  $Y$ . Then  $T$  is called *linear* if for all functions  $f$  and  $g$  in the domain of  $T$  and all  $z \in \mathbb{C}$

$$T(f + g) = T(f) + T(g) \quad T(zf) = zT(f) \tag{1}$$

holds and *quasi-linear* if

$$|T(f + g)| \leq K (|T(f)| + |T(g)|) \quad |T(zf)| = |z| |T(f)| \tag{2}$$

holds for some real constant  $K > 0$ . If  $K = 1$ ,  $T$  is called *sublinear*.

**2. The Complex Method.** This theorem will unfortunately only be applicable to linear operators but will yield quite a natural bound of the operator on the intermediate space. The proof will make strong use of complex variables technique. A major tool will be an application of the maximum modulus principle, known as *Hadamard's three lines lemma*.

**2.1. Hadamard's Three Lines Lemma.** The proof of the Riesz-Thorin interpolation theorem heavily relies on Hadamard's three lines lemma which is itself based on a restatement of the maximum modulus theorem (see [Rud87, p. 212]) formulated in [Rud87, p. 253]. To do so, we have first to establish some common terminology. A complex-valued function  $f$  is said to be *holomorphic* in  $\Omega \subseteq \mathbb{C}$  open, if  $f'(z)$  exists for any  $z \in \Omega$ . By

a region we shall mean a nonempty connected open subset of the complex plane. The restatement reads as follows.

**THEOREM.** *Let  $\Omega \subseteq \mathbb{C}$  be a bounded region and  $f$  be a continuous function on  $\overline{\Omega}$  which is holomorphic in  $\Omega$ . Then*

$$|f(z)| \leq \sup \{|f(z)| : z \in \partial\Omega\}$$

*for every  $z \in \Omega$ . If equality holds at one point  $z \in \Omega$ , then  $f$  is constant.*

**LEMMA 2.1.** (Hadamard's three lines lemma) *Let  $F$  be a holomorphic function in the strip  $S := \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$ , continuous and bounded on  $\overline{S}$ , such that  $|F(z)| \leq B_0$  when  $\operatorname{Re} z = 0$  and  $|F(z)| \leq B_1$  when  $\operatorname{Re} z = 1$ , for some  $0 < B_0, B_1 < \infty$ . Then  $|F(z)| \leq B_0^{1-\theta} B_1^\theta$  when  $\operatorname{Re} z = \theta$ , for any  $0 < \theta < 1$ .*




FIGURE 1. Sketch of the setting of Hadamard's three lines lemma.

*Proof.* For  $z \in \overline{S}$  define

$$G(z) := \frac{F(z)}{B_0^{1-z} B_1^z} \quad G_n(z) := G(z) e^{(z^2-1)/n}, \quad n \in \mathbb{N}_{>0}$$

$G(z)$  and  $G_n(z)$  are holomorphic in  $S$  by

$$G'(z) = \frac{F'(z) - F(z) \log(B_1/B_0)}{B_0^{1-z} B_1^z} \quad G'_n(z) = G'(z) e^{(z^2-1)/n} + \frac{2}{n} z G_n(z)$$

and  $e^z \neq 0$  for every  $z \in \mathbb{C}$ . Further, we have

$$|B_0^{1-z} B_1^z| = (B_0^{1-z} B_0^{1-\bar{z}} B_1^z B_1^{\bar{z}})^{1/2} = B_0^{1-\operatorname{Re} z} B_1^{\operatorname{Re} z}$$

Consider  $0 \leq \operatorname{Re} z \leq 1$  and  $B_0 \geq 1$ . Then  $B_0^{1-\operatorname{Re} z} \geq 1$  and  $B_0^{1-\operatorname{Re} z} \geq B_0$  in the case  $B_0 < 1$ . Similarly,  $B_1^{\operatorname{Re} z} \geq 1$  if  $B_1 \geq 1$  and  $B_1^{\operatorname{Re} z} \leq B_1$  if  $B_1 < 1$ . Hence

$$|B_0^{1-z} B_1^z| \geq \min\{1, B_0\} \min\{1, B_1\} > 0 \quad (3)$$

for all  $z \in \bar{S}$ . Since  $F$  is bounded on  $\bar{S}$ , we have  $|F(z)| \leq L$  for some  $L > 0$  and all  $z \in \bar{S}$ . Thus by (3)

$$|G(z)| = \frac{|F(z)|}{|B_0^{1-z} B_1^z|} \leq \frac{L}{\min\{1, B_0\} \min\{1, B_1\}} =: M$$

for every  $z \in \bar{S}$ . Fix  $n \in \mathbb{N}_{>0}$  and write  $z := x + iy \in \bar{S}$ . Then

$$|G_n(z)| \leq M \left( e^{(x^2+2ixy-y^2-1)/n} e^{(x^2-2ixy-y^2-1)/n} \right)^{1/2} = M e^{-y^2/n} e^{(x^2-1)/n} \leq M e^{-y^2/n}$$

for  $0 \leq x \leq 1$ . Thus

$$\lim_{y \rightarrow \pm\infty} \sup\{|G_n(z)| : 0 \leq x \leq 1\} = 0$$

Hence there exist  $C_0(n), C_1(n) \in \mathbb{R}$ , such that

$$\sup\{|G_n(z)| : 0 \leq x \leq 1\} \leq 1$$

whenever  $y > C_0(n)$  or  $y < C_1(n)$ . Letting

$$C(n) := \max\{|C_0(n)| + 1, |C_1(n)| + 1\}$$

we conclude  $|G_n(z)| \leq 1$  for all  $0 \leq x \leq 1$  when  $|y| \geq C(n)$ . Now consider the rectangle  $R := (0, 1) \times (-C(n), C(n))$ . We have  $|G_n(z)| \leq 1$  on the lines  $[0, 1] \times \{\pm C(n)\}$ . By

$$|G_n(iy)| = \frac{|F(iy)|}{|B_0^{1-iy} B_1^{iy}|} e^{-(y^2+1)/n} \leq 1 \quad |G_n(1+iy)| = \frac{|F(1+iy)|}{|B_0^{-iy} B_1^{1+iy}|} e^{-y^2/n} \leq 1$$

we have  $|G_n(z)| \leq 1$  on the lines  $\{0\} \times [-C(n), C(n)]$ ,  $\{1\} \times [-C(n), C(n)]$ . Thus  $|G_n(z)| \leq 1$  on  $\partial R$ . Since  $|G_n(z)|$  is continuous on  $\bar{R}$ , holomorphic in  $R$  and  $R$  is a bounded region, the maximum modulus theorem implies

$$|G_n(z)| \leq \sup\{|G_n(z)| : z \in \partial R\} \leq 1$$

for every  $z \in R$ . Therefore  $|G_n(z)| \leq 1$  on  $\bar{R}$  and so  $|G_n(z)| \leq 1$  on  $\bar{S}$ . Since inequalities are preserved by limits and the modulus is a continuous function, we have that  $|G(z)| = \lim_{n \rightarrow \infty} |G_n(z)| \leq 1$  for  $z \in \bar{S}$ . We conclude by

$$|F(\theta + it)| = |G(\theta + it)| |B_0^{1-\theta-it} B_1^{\theta+it}| \leq B_0^{1-\theta} B_1^\theta$$

whenever  $0 < \theta < 1$ ,  $t \in \mathbb{R}$ . □




FIGURE 2. Sketch of the rectangle  $\bar{R}$ .

**2.2. The Riesz-Thorin Interpolation Theorem.** Here we will answer our question raised in the introduction for the case where  $T$  is a linear operator. The theorem as it is in [Gra14, p. 37] will be slightly generalized to not necessarily  $\sigma$ -finite spaces in the manner of [Fol99, p. 200], but not in the most general formulation possible where  $T$  is an operator defined on a larger domain. To shorten the argumentation we will introduce some notation. For two measure spaces  $(X, \mu)$ ,  $(Y, \nu)$  let  $\Sigma_X$  and  $\Sigma_Y$  denote the set of all finitely simple functions on  $X$ ,  $Y$  respectively. Furthermore recall, that a measure  $\mu$  on a measure space  $(X, \mu)$  is called *semifinite* if for each measurable set  $A$  with  $\mu(A) = \infty$  there exists a measurable set  $B \subseteq A$  with  $0 < \mu(B) < \infty$  (see [Fol99, p. 25]).

**THEOREM 2.1.** (Riesz-Thorin interpolation theorem) *Suppose that  $(X, \mu)$ ,  $(Y, \nu)$  are measure spaces and  $1 \leq p_0, p_1, q_0, q_1 \leq \infty$ . If  $q_0 = q_1 = \infty$ , suppose also that  $\nu$  is semifinite. Let  $T$  be a linear operator defined on  $\Sigma_X$  and taking values in the set of measurable functions on  $Y$ , such that for some  $0 < M_0, M_1 < \infty$  the estimates*

$$\|T(f)\|_{L^{q_0}} \leq M_0 \|f\|_{L^{p_0}} \quad \text{and} \quad \|T(f)\|_{L^{q_1}} \leq M_1 \|f\|_{L^{p_1}} \quad (4)$$

*hold for all  $f \in \Sigma_X$ . Then for all  $0 \leq \theta \leq 1$  we have*

$$\|T(f)\|_{L^q} \leq M_0^{1-\theta} M_1^\theta \|f\|_{L^p} \quad (5)$$

*for all  $f \in \Sigma_X$ , where*

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1} \quad \text{and} \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}.$$

*Proof.* The idea is to bound the quantity (see [Fol99, p. 189])

$$M_q(T(f)) = \sup \left\{ \left| \int_Y T(f)g \, d\nu \right| : g \in \Sigma_Y, \|g\|_{L^{q'}} = 1 \right\}$$

appropriately (recall  $q' := q/(q-1)$ ). If either  $\theta = 0$  or  $\theta = 1$ , the estimate (5) follows directly from the hypotheses (4) on  $T$ . Thus we may assume  $0 < \theta < 1$ . Furthermore, if  $f \in \Sigma_X$ ,  $\|f\|_{L^p} = 0$ , then  $f = 0$   $\mu$ -a.e. and either one of the hypotheses on  $T$  in (4) implies  $T(f) = 0$   $\mu$ -a.e. and thus the estimate (5) holds trivially. Therefore we can assume  $\|f\|_{L^p} \neq 0$ . Fix  $f \in \Sigma_X$ ,  $g \in \Sigma_Y$  with representation

$$f = \sum_{j=1}^n a_j e^{i\alpha_j} \chi_{A_j} \quad g = \sum_{k=1}^m b_k e^{i\beta_k} \chi_{B_k}$$

where  $a_j, b_k \neq 0$ ,  $\alpha_j, \beta_k \in \mathbb{R}$  for any  $j = 1, \dots, n$ ,  $k = 1, \dots, m$ , the sets  $A_j$  and  $B_k$  are each pairwise disjoint with  $\mu(A_j), \nu(B_k) < \infty$  and so, that  $\|g\|_{L^{q'}} \neq 0$ . Define

$$P(z) := \frac{p}{p_0}(1-z) + \frac{p}{p_1}z \quad Q(z) := \frac{q'}{q'_0}(1-z) + \frac{q'}{q'_1}z$$

for  $z \in \mathbb{C}$  (since either  $p = \infty$  implies  $p_0 = p_1 = \infty$  or  $q = 1$  implies  $q_0 = q_1 = 1$ , the functions  $P, Q$  are well-defined). Further let

$$f_z := \sum_{j=1}^n a_j^{P(z)} e^{i\alpha_j} \chi_{A_j} \quad g_z := \sum_{k=1}^m b_k^{Q(z)} e^{i\beta_k} \chi_{B_k} \quad (6)$$

and

$$F(z) := \int_Y T(f_z)g_z \, d\nu \quad (7)$$

By (6), (7) and the linearity of the operator  $T$  we have

$$F(z) = \sum_{j=1}^n \sum_{k=1}^m a_j^{P(z)} b_k^{Q(z)} e^{i\alpha_j} e^{i\beta_k} \int_Y T(\chi_{A_j}) \chi_{B_k} \, d\nu$$

Applying Hölder's inequality yields

$$\begin{aligned} \left| \int_Y T(\chi_{A_j}) \chi_{B_k} \, d\nu \right| &\leq \int_Y |T(\chi_{A_j}) \chi_{B_k}| \, d\nu \\ &= \|T(\chi_{A_j}) \chi_{B_k}\|_{L^1} \\ &\leq \|T(\chi_{A_j})\|_{L^{q_0}} \|\chi_{B_k}\|_{L^{q'_0}} \\ &\leq M_0 \|\chi_{A_j}\|_{L^{p_0}} \|\chi_{B_k}\|_{L^{q'_0}} \\ &\leq M_0 \mu(A_j)^{1/p_0} \nu(B_k)^{1/q'_0} \end{aligned} \quad (8)$$

for each  $j = 1, \dots, n$ ,  $k = 1, \dots, m$  (even in the cases where either  $p_0 = \infty$  or  $q'_0 = \infty$ , or both, by observing that  $\|\chi_A\|_{L^\infty} \leq 1$  for any measurable set  $A$ ). Thus the function  $F$  is well-defined on  $\mathbb{C}$ . Let  $t \in \mathbb{R}$ . For  $p, p_0 \neq \infty$

$$\begin{aligned} \|f_{it}\|_{L^{p_0}} &= \left( \sum_{j=1}^n \int_{A_j} |f_{it}|^{p_0} d\mu + \int_{X \setminus \bigcup_{j=1}^n A_j} |f_{it}|^{p_0} d\mu \right)^{1/p_0} \\ &= \left( \sum_{j=1}^n |a_j^{P(it)} e^{i\alpha_j}|^{p_0} \int_X \chi_{A_j} d\mu \right)^{1/p_0} \\ &= \left( \sum_{j=1}^n a_j^{p_0 \operatorname{Re} P(it)} \mu(A_j) \right)^{1/p_0} \\ &= \left( \sum_{j=1}^n a_j^p \mu(A_j) \right)^{p/(p_0 p)} \\ &= \|f\|_{L^p}^{p/p_0} \end{aligned}$$

holds. Let  $p_0 = \infty$ ,  $p \neq \infty$ . Then  $\|f_{it}\|_{L^\infty} = 1$  since  $|a_j^{P(it)}| = a_j^{p/p_0} = 1$  and that there exists some index  $j$ , such that  $\mu(A_j) \neq 0$ . If  $p = \infty$ , then  $p_0 = p_1 = \infty$  and thus  $P(it) = 1$ . By the same considerations we have  $\|g_{it}\|_{L^{q'_0}} = \|g\|_{L^{q'}}^{q'/q'_0}$ . Hence

$$\begin{aligned} |F(it)| &\leq \int_Y |T(f_{it})g_{it}| d\nu \\ &= \|T(f_{it})g_{it}\|_{L^1} \\ &\leq \|T(f_{it})\|_{L^{q_0}} \|g_{it}\|_{L^{q'_0}} \\ &\leq M_0 \|f_{it}\|_{L^{p_0}} \|g_{it}\|_{L^{q'_0}} \\ &= M_0 \|f\|_{L^p}^{p/p_0} \|g\|_{L^{q'}}^{q'/q'_0} \end{aligned}$$

by Hölder's inequality. In an analogous manner we derive

$$\|f_{1+it}\|_{L^{p_1}} = \|f\|_{L^p}^{p/p_1} \quad \|g_{1+it}\|_{L^{q'_1}} = \|g\|_{L^{q'}}^{q'/q'_1}$$

and thus

$$|F(1+it)| \leq M_1 \|f\|_{L^p}^{p/p_1} \|g\|_{L^{q'}}^{q'/q'_1}$$

Further by estimate (8)

$$\begin{aligned}
|F(z)| &\leq \sum_{j=1}^n \sum_{k=1}^m |a_j^{P(z)}| |b_k^{Q(z)}| \left| \int_Y T(\chi_{A_j}) \chi_{B_k} d\nu \right| \\
&\leq M_0 \sum_{j=1}^n \sum_{k=1}^m a_j^{\operatorname{Re} P(z)} b_k^{\operatorname{Re} Q(z)} \mu(A_j)^{1/p_0} \nu(B_k)^{1/q'_0} \\
&\leq M_0 \sum_{j=1}^n \sum_{k=1}^m \max\{1, a_j^{p/p_0+p/p_1}\} \max\{1, b_k^{q'/q'_0+q'/q'_1}\} \mu(A_j)^{1/p_0} \nu(B_k)^{1/q'_0}
\end{aligned}$$

Hence  $F$  is bounded on  $\bar{S}$  by some constant depending on  $f$  and  $g$  only. By

$$\begin{aligned}
F'(z) &= \sum_{j=1}^n \sum_{k=1}^m a_j^{P(z)} \log(a_j) \left( \frac{p}{p_1} - \frac{p}{p_0} \right) b_k^{Q(z)} e^{i\alpha_j} e^{i\beta_k} \int_Y T(\chi_{A_j}) \chi_{B_k} d\nu \\
&\quad + \sum_{j=1}^n \sum_{k=1}^m a_j^{P(z)} b_k^{Q(z)} \log(b_k) \left( \frac{q'}{q'_1} - \frac{q'}{q'_0} \right) e^{i\alpha_j} e^{i\beta_k} \int_Y T(\chi_{A_j}) \chi_{B_k} d\nu
\end{aligned}$$

it is immediate, that  $F$  is an entire function and thus holomorphic in  $S$  and continuous on  $\bar{S}$ . Therefore lemma 2.1 yields

$$|F(z)| \leq \left( M_0 \|f\|_{L^p}^{p/p_0} \|g\|_{L^{q'}}^{q'/q'_0} \right)^{1-\theta} \left( M_1 \|f\|_{L^p}^{p/p_1} \|g\|_{L^{q'}}^{q'/q'_1} \right)^\theta = M_0^{1-\theta} M_1^\theta \|f\|_{L^p} \|g\|_{L^{q'}}$$

for  $\operatorname{Re} z = \theta$ ,  $0 < \theta < 1$ . We have

$$\{T(f) \neq 0\} = \bigcup_{n=1}^{\infty} \{|T(f)| > 1/n\}$$

and by Chebychev's inequality (see [Fol99, p. 193]) either

$$\nu(\{|T(f)| > 1/n\}) \leq n^{q_0} \|T(f)\|_{L^{q_0}}^{q_0} \leq n^{q_0} M_0^{q_0} \|f\|_{L^{p_0}}^{q_0}$$

or

$$\nu(\{|T(f)| > 1/n\}) \leq n^{q_1} \|T(f)\|_{L^{q_1}}^{q_1} \leq n^{q_1} M_1^{q_1} \|f\|_{L^{p_1}}^{q_1}$$

whenever  $q_0 \neq \infty$  or  $q_1 \neq \infty$ . Therefore, the set  $\{T(f) \neq 0\}$  is  $\sigma$ -finite unless  $q_0 = q_1 = \infty$ . Further we have  $P(\theta) = Q(\theta) = 1$ . Thus by

$$\begin{aligned}
M_q(T(f)) &= \sup \left\{ \left| \int_Y T(f) g d\nu \right| : g \in \Sigma_Y, \|g\|_{L^{q'}} = 1 \right\} \\
&= \sup \{|F(\theta)| : g \in \Sigma_Y, \|g\|_{L^{q'}} = 1\} \\
&\leq M_0^{1-\theta} M_1^\theta \|f\|_{L^p}
\end{aligned}$$



we conclude

$$\|T(f)\|_{L^q} = M_q(T(f)) \leq M_0^{1-\theta} M_1^\theta \|f\|_{L^p}$$

for any  $f \in \Sigma_X$ . □

REMARK 2.1. A more general version of the Riesz-Thorin interpolation theorem can be found in [Fol99, pp. 200–202]. There, a linear map  $T : L^{p_0} + L^{p_1} \rightarrow L^{q_0} + L^{q_1}$  is considered. This follows using a density argument from the current version of the theorem and will not be proven here.

REMARK 2.2. Using the previous remark, a standard application of the Riesz-Thorin interpolation theorem is to prove Young's inequality for convolutions [Gra14, pp. 22–23]. Let  $G$  be a locally compact group and  $\lambda$  be a left invariant Haar measure on  $G$ , furthermore we assume that  $G$  is a countable union of compact subsets, hence the pair  $(G, \lambda)$  forms a  $\sigma$ -finite measure space.

THEOREM. (Young's inequality) Let  $1 \leq p, q, r \leq \infty$  satisfy

$$\frac{1}{q} + 1 = \frac{1}{p} + \frac{1}{r} \tag{9}$$

Then for all  $f \in L^p(G)$ ,  $g \in L^r(G)$  satisfying  $\|g\|_{L^r} = \|\tilde{g}\|$  we have  $f * g$  exists  $\lambda$ -a.e. and satisfies

$$\|f * g\|_{L^1} \leq \|g\|_{L^r} \|f\|_{L^p}$$

*Proof.* Fix  $g \in L^r(G)$  and let  $T(f) := f * g$  be defined on  $L^1(G) + L^{r'}(G)$ . Obviously,  $T$  is a linear operator by the linearity of the integral. By Minkowski's integral inequality (see exercise 1.1.6 [Gra14, p. 13]) we get

$$\begin{aligned} \|T(f)\|_{L^r} &= \left( \int_G \left| \int_G f(y) g(y^{-1}x) d\lambda(y) \right|^r d\lambda(x) \right)^{1/r} \\ &\leq \int_G \left( \int_G |f(y)|^r |g(y^{-1}x)|^r d\lambda(x) \right)^{1/r} d\lambda(y) \\ &= \int_G |f(y)| \left( \int_G |g(y^{-1}x)|^r d\lambda(y^{-1}x) \right)^{1/r} d\lambda(y) \\ &= \int_G |f(y)| \left( \int_G |g(z)|^r d\lambda(z) \right)^{1/r} d\lambda(y) \\ &\leq \|f\|_{L^1} \|g\|_{L^r} \end{aligned}$$

for  $f \in L^1(g, \mu)$  and  $1 \leq p < \infty$ . The case  $r = \infty$  follows from

$$|(f * g)(x)| = \left| \int_G f(y) g(y^{-1}x) d\lambda(y) \right| \leq \int_G |f(y)| |g(y^{-1}x)| d\lambda(y) \leq \|g\|_{L^\infty} \|f\|_{L^1}$$

By stipulating  $h(y) := g(y^{-1}x)$  we have

$$\begin{aligned} |(f * g)(x)| &= \left| \int_G f(y)g(y^{-1}x) d\lambda(y) \right| \leq \int_G |f(y)g(y^{-1}x)| d\lambda(y) \\ &= \|fh\|_{L^1} \leq \|f\|_{L^{r'}} \|h\|_{L^r} = \|f\|_{L^{r'}} \|\tilde{g}\|_{L^r} = \|g\|_{L^r} \|f\|_{L^{r'}} \end{aligned}$$

for  $r < \infty$  and  $f \in L^{r'}(G)$ , since

$$\|h\|_{L^r}^r = \int_G |g(y^{-1}x)|^r d\lambda(y) = \int_G |\tilde{g}(x^{-1}y)| d\lambda(y) = \|\tilde{g}\|_{L^r}^r$$

The Riesz-Thorin interpolation theorem now yields for any  $0 < \theta < 1$

$$\|f * g\|_{L^q} = \|T(f)\|_{L^q} \leq \|g\|_{L^{r'}}^{1-\theta} \|g\|_{L^r}^\theta \|f\|_{L^p} = \|g\|_{L^r} \|f\|_{L^p} \quad (10)$$

where

$$\frac{1}{p} = \frac{1-\theta}{1} + \frac{\theta}{r'} \quad \frac{1}{q} = \frac{1-\theta}{r} + \frac{\theta}{\infty}$$

□

REMARK 2.3. *The proof would be much shorter if we just used Minkowski's inequality [Gra14, pp. 21–22] instead of Minkowski's integral inequality. However, the proof given here is an alternative version of the one given already for Minkowski's inequality.*

**3. Interpolation of Analytic Families of Operators.** The generalization of the classical Riesz-Thorin interpolation theorem to analytic families of operators is due to *E. M. Stein* and *Guido Weiss*<sup>1</sup>. Crucial for its proof is again an application of advanced topics in complex analysis.

**3.1. Extension of Hadamard's Three Lines Lemma.** This lemma is inspired by a lemma originally proposed by I.I.Hirschman. I will stick for the most part to the proof given originally in the paper by Stein and Weiss and for some parts to the proof given in [Gra14, pp. 43–45].

**3.1.1. Auxiliary Lemmata.** To shorten the proof of the extension of Hadamard's three lines lemma, I will summarize the most important facts used during the proof.

LEMMA 3.1. *Let  $D := \{z \in \mathbb{C} : |z| < 1\}$  be the open unit disc and*

$$h(z) := \frac{1}{\pi i} \log \left( i \frac{1+z}{1-z} \right)$$

*for  $z \in \overline{D} \setminus \{\pm 1\}$  where we are taking that continuous branch of  $\log z$  in the complex plane slit along the negative imaginary axis,  $\mathbb{C} \setminus (\{0\} \times [0, \infty))$ . Then  $h$  is a holomorphic function in  $D$  which maps  $\overline{D} \setminus \{\pm 1\}$  bijectively onto the closure  $\overline{S}$  of the strip  $S := \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$ .*

<sup>1</sup><https://projecteuclid.org/euclid.tmj/1178244785>, last accessed November 12, 2016.

*Proof.* Consider the mapping  $\varphi : \mathbb{S}^2 \rightarrow \mathbb{S}^2$  defined by

$$\varphi(z) := \frac{1+z}{1-z}$$

This  $\varphi$  maps  $\{-1, 0, 1\}$  to  $\{0, 1, \infty\}$  and is a conformal mapping by [Rud87, pp. 278–279]. The segment  $(-1, 1)$  maps onto the positive real axis as can be verified by considering the corresponding real limits and the unit circle  $\mathbb{S}^1$  passes through  $-1$  and  $1$ , hence  $\varphi(\mathbb{S}^1)$  is a straight line through  $\varphi(-1) = 0$ . Since  $\mathbb{S}^1$  makes a right angle with the real axis at  $-1$  so does  $\varphi(\mathbb{S}^1)$  at  $0$  by the conformality of  $\varphi$ . Thus  $\varphi(\mathbb{S}^1)$  is the imaginary axis. Since  $\varphi(0) = 1$ , it follows that  $\varphi$  is a conformal one-to-one mapping of the open unit disc onto the open right half plane. Furthermore,  $\varphi(\bar{D} \setminus \{\pm 1\}) = \mathbb{H}^\times$  where  $\mathbb{H}^\times := \{z \in \mathbb{C} : \operatorname{Re} z \geq 0\} \setminus \{0\}$ . Thus  $i\varphi(\bar{D} \setminus \{\pm 1\}) = i\mathbb{H}^\times = \{z \in \mathbb{C} : \operatorname{Im} z \geq 0\} \setminus \{0\}$  and so

$$\log i\mathbb{H}^\times = \log |\mathbb{H}^\times| + i \arg i\mathbb{H}^\times = \{z \in \mathbb{C} : 0 \leq \operatorname{Im} z \leq \pi\}$$

Finally, multiplication with the preceding factor  $1/(\pi i)$  yields  $h(\bar{D} \setminus \{\pm 1\}) = \bar{S}$ . Furthermore, we have

$$h^{-1}(z) = \frac{e^{\pi iz} - i}{e^{\pi iz} + i}$$

□

LEMMA. Let  $X$  be a topological space. A function  $f : X \rightarrow [-\infty, \infty)$  is upper semicontinuous if and only if for all  $\alpha \in \mathbb{R}$  the set  $f^{-1}([-\infty, \alpha))$  is open.

*Proof.* Suppose  $f : X \rightarrow [-\infty, \infty)$  is upper semicontinuous and fix  $\alpha \in \mathbb{R}$ . We have that

$$f^{-1}([-\infty, \alpha)) = \bigcup_{x \in \{f < \alpha\}} U_x$$

where  $U_x$  is a neighbourhood of  $x$  such that  $f < \alpha$  for any element in  $U_x$ . Conversely, for  $x_0 \in X$  and  $\alpha > f(x_0)$  we have that  $f^{-1}([-\infty, \alpha))$  is open and  $x_0 \in f^{-1}([-\infty, \alpha))$ . □

LEMMA. An upper semicontinuous function  $f : X \rightarrow [-\infty, \infty)$  on a compact topological space attains its supremum. In particular it is bounded from above.

*Proof.*  $f(X)$  is bounded from above since otherwise

$$X = \bigcup_{n \in \mathbb{N}} f^{-1}([-\infty, n))$$

would not have any finite subcover. Therefore  $\sup_{x \in X} f(x)$  exists. Further we have  $f(x_0) = \sup_{x \in X} f(x)$  for some  $x_0 \in X$  since otherwise

$$X = \bigcup_{n \in \mathbb{N}} f^{-1}([-\infty, \sup_{x \in X} f(x) - 1/n))$$

would not have any finite subcover. □

LEMMA 3.2. *Let  $\Omega \subseteq \mathbb{C}$  and  $f : \Omega \rightarrow \mathbb{C}$  continuous. Then  $\log |f|$  is upper semicontinuous on  $\Omega$ .*

*Proof.* Let us consider the topological space  $(\Omega, |\cdot|)$ . Let  $z_0 \in \Omega$  so such that  $f(z_0) \neq 0$ . Then  $\log |f|$  is continuous as a composition of continuous functions. If  $M > f(z_0)$ , then  $M - \log |f(z_0)| > 0$  and thus there exists some  $\delta > 0$  such that  $z \in B_\delta(z_0)$  implies  $|\log |f(z)| - \log |f(z_0)|| < M - \log |f(z_0)|$  or equivalently  $\log |f(z)| < M$ . Now let  $z_0 \in \Omega$  so such that  $f(z_0) = 0$ . By convention  $\log |f(z_0)| = -\infty$ . Furthermore,  $M > \log |f(z_0)|$  for any  $M \in \mathbb{R}$ . The condition  $M > \log |f(z)|$  is equivalent to  $|f(z)| < e^M$ . But  $f(z_0) = 0$  and so

$$|f(z)| = |f(z) - f(z_0)| < e^M$$

Since  $f$  is continuous at  $z_0$  and  $e^M > 0$  we find  $\delta > 0$  such that  $z \in B_\delta(z_0)$  implies  $|f(z)| < e^M$ .  $\square$

LEMMA 3.3. *The mapping  $\Phi : \mathbb{R} \rightarrow (-\pi, 0)$  defined by  $\Phi(t) := -i \log(h^{-1}(it))$  is a  $C^1$ -Diffeomorphism with  $|D\Phi(t)| = \pi \operatorname{sech}(\pi t)$ . In an analogous manner we have that  $\Psi : \mathbb{R} \rightarrow (0, \pi)$ ,  $\Psi(t) := -i \log(h^{-1}(1+it))$  is a  $C^1$ -Diffeomorphism with  $|D\Psi(t)| = \pi \operatorname{sech}(\pi t)$ .*

*Proof.* It is easier to consider  $\Phi^{-1}(\varphi) = -ih(e^{i\varphi})$  and  $\Psi^{-1}(\varphi) = i - ih(e^{i\varphi})$  (this already shows that  $\Phi, \Psi$  are bijective mappings). Let us consider  $\varphi$  only since the argumentation for  $\Psi$  is similar. By  $|e^{i\varphi}| = 1$  it is immediate that  $\Phi^{-1}$  is a real valued function. Furthermore,  $\lim_{\varphi \rightarrow -\pi} \Phi^{-1}(\varphi) = \infty$ ,  $\lim_{\varphi \rightarrow 0} \Phi^{-1}(\varphi) = -\infty$  and  $\Phi^{-1}$  is clearly continuously differentiable. Using

$$h^{-1}(it) = \frac{e^{-\pi t} - i}{e^{-\pi t} + i}$$

we get

$$|D\Phi(t)| = \pi \left| \frac{e^{-\pi t}}{e^{\pi t} - i} - \frac{e^{-\pi t}}{e^{-\pi t} + i} \right| = \pi \left| \frac{2e^{-\pi t}}{e^{-2\pi t} + 1} \right| = \pi \left| \frac{2}{e^{-\pi t} + e^{\pi t}} \right| = \pi \operatorname{sech}(\pi t)$$

$\square$

LEMMA 3.4. *Let  $1/(2e - 1) \leq \rho < 1$  and  $\zeta = \rho e^{i\theta}$ . Then*

$$\left| \log \left| \frac{1 + \zeta}{1 - \zeta} \right| \right| \leq 1 + \log \frac{1}{|\cos(\theta/2)|} + \log \frac{1}{|\sin(\theta/2)|}$$

*Proof.* This proof is due to Prof. Schlein. We have on the one hand

$$|1 + \zeta| \leq 1 + |\zeta| = 1 + \rho$$

and on the other hand

$$|1 - \zeta| \geq |\operatorname{Im} \zeta| = \rho |\sin(\theta)|$$

Hence

$$\begin{aligned}
 \log \frac{|1 + \zeta|}{|1 - \zeta|} &\leq \log \frac{1 + \rho}{\rho |\sin(\theta)|} \\
 &= \log \frac{1 + \rho}{2\rho |\sin(\theta/2)| |\cos(\theta/2)|} \\
 &= \log \frac{1 + \rho}{2\rho} + \log \frac{1}{|\sin(\theta/2)|} + \log \frac{1}{|\cos(\theta/2)|} \\
 &\leq 1 + \log \frac{1}{|\sin(\theta/2)|} + \log \frac{1}{|\cos(\theta/2)|}
 \end{aligned}$$

since

$$\frac{1 + \rho}{2\rho} = \frac{1}{2} + \frac{1}{2\rho} \leq e$$

Now by

$$-\log \frac{|1 + \zeta|}{|1 - \zeta|} = \log \frac{|1 - \zeta|}{|1 + \zeta|}$$

which corresponds to considering  $-\zeta = e^{i\pi}\zeta = e^{i(\pi+\theta)}$  in the first case, yields by invoking the identities

$$\cos\left(\frac{\pi + \theta}{2}\right) = -\sin(\theta/2) \quad \sin\left(\frac{\pi + \theta}{2}\right) = \cos(\theta/2)$$

the bound

$$-\log \frac{|1 + \zeta|}{|1 - \zeta|} \leq 1 + \log \frac{1}{|\sin(\theta/2)|} + \log \frac{1}{|\cos(\theta/2)|}$$

and we are done. □

LEMMA 3.5. *Let  $0 \leq \tau_0 < \pi$ . Then*

$$\frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} \in L^1[-\pi, \pi]$$

*Proof.* The case  $\tau_0 = 0$  is trivial. We use [Els11, pp. 153–154]. By the symmetry of the integrand it is enough to consider

$$\int_0^\pi \frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} d\theta$$

Thus we may perform the splitting

$$\int_0^1 \frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} d\theta + \int_1^\pi \frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} d\theta$$

Let us consider only the first integral, the second one is similar. Then we have

$$\frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \leq 1$$

for  $0 \leq \theta \leq 1$  and

$$\lim_{\theta \searrow 0} \frac{(\theta/2)^{\tau_0/\pi}}{\sin(\theta/2)^{\tau_0/\pi}} = \lim_{\theta \searrow 0} \frac{(\theta/2)^{\tau_0/\pi}}{\left(\frac{\theta}{2} + O\left(\frac{\theta^3}{8}\right)\right)^{\tau_0/\pi}} = \lim_{\theta \searrow 0} \frac{(\theta/2)^{\tau_0/\pi}}{(\theta/2)^{\tau_0/\pi} \left(1 + O\left(\frac{\theta^2}{4}\right)\right)^{\tau_0/\pi}} = 1$$

Thus

$$\frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} \sim \frac{1}{(\theta/2)^{\tau_0/\pi}}$$

when  $\theta \searrow 0$  and since  $\tau_0 < \pi$ , the latter integral converges. Thus we conclude by the comparison theorem.  $\square$

**3.1.2. The Lemma.** Now we are ready to prove the extension of the lemma 2.1. Recall, that a function  $f : X \rightarrow [-\infty, \infty)$  defined on a topological space  $X$  is said to be *upper semicontinuous* if for every point  $x_0 \in X$  and each real  $M > f(x_0)$  a neighbourhood  $U$  of  $x_0$  exists such that  $M > f(x)$  for every  $x \in U$  (see [HS91, p. 199]).

**LEMMA 3.6.** (Hadamard's three lines lemma, extension) *Let  $F$  be a holomorphic function in the strip  $S := \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$  and continuous on  $\bar{S}$ , such that for some  $0 < A < \infty$  and  $0 \leq \tau_0 < \pi$  we have  $\log |F(z)| \leq Ae^{\tau_0 |\operatorname{Im} z|}$  for every  $z \in \bar{S}$ . Then*

$$|F(z)| \leq \exp \left( \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[ \frac{\log |F(it + iy)|}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log |F(1 + it + iy)|}{\cosh(\pi t) + \cos(\pi x)} \right] dt \right)$$

*whenever  $z := x + iy \in S$ .*

*Proof.* We will first prove the case  $y = 0$ . Assume  $F$  to be not identically zero (the case where  $F$  is identically zero is trivial). Let  $h$  be as in lemma (3.1) and let  $\zeta := \rho e^{i\theta}$ ,  $0 \leq \rho < 1$ . Since  $\zeta \in D$ , we have  $0 < \operatorname{Re} h(\zeta) < 1$  and thus the hypothesis on  $F$  and lemma (3.4) yields

$$\log |F(h(\zeta))| \leq Ae^{\frac{\tau_0}{\pi} |\log|(1+\zeta)/(1-\zeta)||} \leq Ae^{\tau_0/\pi} \frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} \quad (11)$$

for  $1/(2e-1) \leq \rho$ . Since  $0 < \tau_0 < \pi$ , inequality (11) asserts, that  $\log |F(h(\zeta))|$  is bounded from above by an integrable function of  $\theta$ , independently of  $\rho \geq 1/(2e-1)$ . Furthermore we have

$$M := \sup \{ \log |F(h(\zeta))| : \zeta \in \bar{B}_{1/(2e-1)} \} < \infty \quad (12)$$

since a upper semicontinuous function on a compact space attains its supremum (see lemma 3.1.1). Hence

$$\log |F(h(\rho e^{i\theta}))| \leq \max \left\{ M, A e^{\tau_0/\pi} \frac{1}{|\cos(\theta/2)|^{\tau_0/\pi}} \frac{1}{|\sin(\theta/2)|^{\tau_0/\pi}} \right\} =: g(\theta) \quad (13)$$

for any  $0 \leq \rho < 1$  where  $g \in L^1[-\pi, \pi]$ . Let  $0 \leq \rho < R < 1$  and  $a_1, \dots, a_n$  denote the zeros of  $F(h(\zeta))$  for  $|\zeta| < R$  (since  $F \circ h$  is holomorphic for  $|\zeta| < 1$  there are indeed only finitely many ones) multiple zeros being repeated. Then for  $F(h(\zeta)) \neq 0$  we have by the *Poisson-Jensen formula* (see [Ahl79, p. 208])

$$\log |F(h(\zeta))| = - \sum_{k=1}^n \log \left| \frac{R^2 - \bar{a}_k \zeta}{R(\zeta - a_k)} \right| + \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left[ \frac{R e^{it} + \zeta}{R e^{it} - \zeta} \right] \log |F(h(R e^{it}))| dt \quad (14)$$

Therefore by

$$\begin{aligned} \operatorname{Re} \left[ \frac{R e^{it} + \zeta}{R e^{it} - \zeta} \right] &= \operatorname{Re} \left[ \frac{R^2 - 2i \operatorname{Im} [\zeta R e^{-it}] - |\zeta|^2}{R^2 - 2 \operatorname{Re} [\zeta R e^{-it}] + |\zeta|^2} \right] \\ &= \operatorname{Re} \left[ \frac{R^2 - 2i R \rho \sin(\theta - t) - \rho^2}{R^2 - 2R \rho \cos(\theta - t) + \rho^2} \right] \\ &= \frac{R^2 - \rho^2}{R^2 - 2R \rho \cos(\theta - t) + \rho^2} \end{aligned}$$

and since  $(R^2 - |a_k|^2)(R^2 - \rho^2) \geq 0$  for all  $k = 1, \dots, n$  implies  $|R^2 - \bar{a}_k \zeta| \geq |R(\zeta - a_k)|$  the estimate

$$\log |F(h(\zeta))| \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{R^2 - \rho^2}{R^2 - 2R \rho \cos(\theta - t) + \rho^2} \log |F(h(R e^{it}))| dt \quad (15)$$

is valid for every  $|\zeta| < R$ . By [Rud87, p. 236] we have

$$\frac{R - \rho}{R + \rho} \leq \frac{R^2 - \rho^2}{R^2 - 2R \rho \cos(\theta - \varphi) + \rho^2} \leq \frac{R + \rho}{R - \rho} \quad (16)$$

for  $0 \leq \rho < R$ . Combining (13) and (16) yields

$$\frac{R^2 - \rho^2}{R^2 - 2R \rho \cos(\theta - t) + \rho^2} \log |F(h(\zeta))| \leq \frac{R + \rho}{R - \rho} g(\theta) =: G(\theta) \quad (17)$$

where  $G \in L^1[-\pi, \pi]$ . For  $0 < R < 1$  let

$$f_R(\varphi) := \frac{R^2 - \rho^2}{R^2 - 2R \rho \cos(\theta - \varphi) + \rho^2} \log |F(h(R e^{i\varphi}))|$$

and for  $\varphi \notin \{0, \pi\}$

$$f(\varphi) := \frac{1 - \rho^2}{1 - 2\rho \cos(\theta - \varphi) + \rho^2} \log |F(h(e^{i\varphi}))|$$

Since  $\log |F(h(\zeta))|$  is upper semicontinuous on  $\overline{D} \setminus \{\pm 1\}$  by lemma 3.2 we get

$$\begin{aligned} \limsup_{R \nearrow 1} f_R(\varphi) &= \limsup_{R \nearrow 1} \left[ \frac{R^2 - \rho^2}{R^2 - 2R\rho \cos(\theta - \varphi) + \rho^2} \log |F(h(Re^{i\varphi}))| \right] \\ &= \lim_{R \nearrow 1} \frac{R^2 - \rho^2}{R^2 - 2R\rho \cos(\theta - \varphi) + \rho^2} \limsup_{R \nearrow 1} \log |F(h(Re^{i\varphi}))| \\ &= \frac{1 - \rho^2}{1 - 2\rho \cos(\theta - \varphi) + \rho^2} \log |F(h(e^{i\varphi}))| = f(\varphi) \end{aligned}$$

using [Bou95, p. 363] and proposition A.1. The functions  $G - f_R$  being non-negative, an application of Fatou's lemma yields

$$\int_{-\pi}^{\pi} \liminf_{R \nearrow 1} [G(\varphi) - f_R(\varphi)] d\varphi \leq \liminf_{R \nearrow 1} \int_{-\pi}^{\pi} [G(\varphi) - f_R(\varphi)] d\varphi$$

By [Bou95, p. 354], we get

$$\limsup_{R \nearrow 1} \int_{-\pi}^{\pi} [f_R(\varphi) - G(\varphi)] d\varphi \leq \int_{-\pi}^{\pi} \limsup_{R \nearrow 1} [f_R(\varphi) - G(\varphi)] d\varphi$$

and thus

$$\begin{aligned} \limsup_{R \nearrow 1} \int_{-\pi}^{\pi} f_R(\varphi) d\varphi - \int_{-\pi}^{\pi} G(\varphi) d\varphi &= \limsup_{R \nearrow 1} \int_{-\pi}^{\pi} f_R(\varphi) d\varphi - \lim_{R \nearrow 1} \int_{-\pi}^{\pi} G(\varphi) d\varphi \\ &= \limsup_{R \nearrow 1} \int_{-\pi}^{\pi} [f_R(\varphi) - G(\varphi)] d\varphi \\ &\leq \int_{-\pi}^{\pi} \limsup_{R \nearrow 1} [f_R(\varphi) - G(\varphi)] d\varphi \\ &\leq \int_{-\pi}^{\pi} \limsup_{R \nearrow 1} f_R(\varphi) d\varphi - \int_{-\pi}^{\pi} \lim_{R \nearrow 1} G(\varphi) d\varphi \\ &= \int_{-\pi}^{\pi} \limsup_{R \nearrow 1} f_R(\varphi) d\varphi - \int_{-\pi}^{\pi} G(\varphi) d\varphi \end{aligned}$$

by [Bou95, p. 358]. Hence

$$\limsup_{R \nearrow 1} \int_{-\pi}^{\pi} f_R(\varphi) d\varphi \leq \int_{-\pi}^{\pi} \limsup_{R \nearrow 1} f_R(\varphi) d\varphi$$

and so

$$\log |F(h(\zeta))| \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1 - \rho^2}{1 - 2\rho \cos(\theta - \varphi) + \rho^2} \log |F(h(e^{i\varphi}))| d\varphi \quad (18)$$



The lemma now follows from (18) by a change of variables. By stipulating  $x := h(\zeta)$  we obtain

$$\begin{aligned}\zeta = h^{-1}(x) &= \frac{e^{\pi i x} - i}{e^{\pi i x} + i} = \frac{\cos(\pi x) + i \sin(\pi x) - i}{\cos(\pi x) + i \sin(\pi x) + i} \\ &= \frac{\cos(\pi x) + i \sin(\pi x) - i \cos(\pi x) - i \sin(\pi x) - i}{\cos(\pi x) + i \sin(\pi x) + i \cos(\pi x) - i \sin(\pi x) - i} \\ &= -i \frac{\cos(\pi x)}{1 + \sin(\pi x)} = \left( \frac{\cos(\pi x)}{1 + \sin(\pi x)} \right) e^{-i\pi/2} \quad (19)\end{aligned}$$

by

$$\begin{aligned}(\cos(\pi x) + i \sin(\pi x) - i)(\cos(\pi x) - i \sin(\pi x) - i) \\ = \cos^2(\pi x) - i \sin(\pi x) \cos(\pi x) - i \cos(\pi x) + i \sin(\pi x) \cos(\pi x) \\ + \sin^2(\pi x) + \sin(\pi x) - i \cos(\pi x) - \sin(\pi x) - 1 = -2i \cos(\pi x)\end{aligned}$$

and

$$\begin{aligned}(\cos(\pi x) + i \sin(\pi x) + i)(\cos(\pi x) - i \sin(\pi x) - i) \\ = \cos^2(\pi x) - i \sin(\pi x) \cos(\pi x) - i \cos(\pi x) + i \sin(\pi x) \cos(\pi x) \\ + \sin^2(\pi x) + \sin(\pi x) + i \cos(\pi x) + \sin(\pi x) + 1 = 2 + 2 \sin(\pi x)\end{aligned}$$

From equality (19) we deduce  $\rho = \frac{\cos(\pi x)}{1 + \sin(\pi x)}$ ,  $\theta = -\frac{\pi}{2}$  if  $0 < x \leq \frac{1}{2}$  and  $\rho = -\frac{\cos(\pi x)}{1 + \sin(\pi x)}$ ,  $\theta = \frac{\pi}{2}$  if  $\frac{1}{2} \leq x < 1$ . Let  $0 < x \leq \frac{1}{2}$ . Then we have

$$\begin{aligned}\frac{1 - \rho^2}{1 - 2\rho \cos(\theta - \varphi) + \rho^2} \\ = \frac{1 + 2 \sin(\pi x) + \sin^2(\pi x) - \cos^2(\pi x)}{1 + 2 \sin(\pi x) + \sin^2(\pi x) + 2 \cos(\pi x) \sin(\varphi)(1 + \sin(\pi x)) + \cos^2(\pi x)} \\ = \frac{\sin(\pi x) + \sin^2(\pi x)}{1 + \sin(\pi x) + \cos(\pi x) \sin(\varphi)(1 + \sin(\pi x))} = \frac{\sin(\pi x)}{1 + \cos(\pi x) \sin(\varphi)}\end{aligned}$$

and also for  $\frac{1}{2} \leq x < 1$ . Let  $\Phi$  and  $\Psi$  be defined as in lemma (3.3). We have

$$\begin{aligned}e^{i\Phi(t)} = h^{-1}(it) &= \frac{e^{-\pi t} - i e^{-\pi t} - i}{e^{-\pi t} + i e^{-\pi t} - i} = \frac{e^{-2\pi t} - 2i e^{-\pi t} - 1}{e^{-2\pi t} + 1} = \frac{e^{-2\pi t} - 1}{e^{-2\pi t} + 1} - \frac{2i e^{-\pi t}}{e^{-2\pi t} + 1} \\ &= \frac{e^{-2\pi t} - 1}{e^{-2\pi t} + 1} - \frac{2i}{e^{-\pi t} + e^{\pi t}} = \frac{1 - e^{2\pi t}}{1 + e^{2\pi t}} - \frac{2i}{e^{-\pi t} + e^{\pi t}} = -\tanh(\pi t) - i \operatorname{sech}(\pi t)\end{aligned}$$

and thus

$$\begin{aligned}
\sin(\Phi(t)) \cosh(\pi t) &= \sin(-i \log(-\tanh(\pi t) - i \operatorname{sech}(\pi t))) \cosh(\pi t) \\
&= \frac{1}{2i} \left[ -\tanh(\pi t) - i \operatorname{sech}(\pi t) + \frac{1}{\tanh(\pi t) + i \operatorname{sech}(\pi t)} \right] \cosh(\pi t) \\
&= \frac{1}{2i} \left[ \frac{\cosh(\pi t) - \tanh(\pi t) \sinh(\pi t) - 2i \tanh(\pi t) + \operatorname{sech}(\pi t)}{\tanh(\pi t) + i \operatorname{sech}(\pi t)} \right] \\
&= \frac{1}{2i} \left[ \frac{\cosh^2(\pi t) - \sinh^2(\pi t) - 2i \sinh(\pi t) + 1}{\sinh(\pi t) + i} \right] \\
&= \frac{1 - i \sinh(\pi t)}{i \sinh(\pi t) - 1} = -1
\end{aligned}$$

Therefore the transformation formula yields

$$\frac{1}{2\pi} \int_{-\pi}^0 \frac{\sin(\pi x)}{1 + \cos(\pi x) \sin(\varphi)} \log |F(h(e^{i\varphi}))| d\varphi = \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin(\pi x)}{\cosh(\pi t) - \cos(\pi x)} \log |F(it)| dt$$

and in a similar manner

$$\frac{1}{2\pi} \int_0^{\pi} \frac{\sin(\pi x)}{1 + \cos(\pi x) \sin(\varphi)} \log |F(h(e^{i\varphi}))| d\varphi = \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin(\pi x)}{\cosh(\pi t) + \cos(\pi x)} \log |F(1 + it)| dt$$

holds since

$$\begin{aligned}
\sin(\Psi(t)) \cosh(\pi t) &= \sin(-i \log(-\tanh(\pi t) + i \operatorname{sech}(\pi t))) \cosh(\pi t) \\
&= \frac{1}{2i} \left[ -\tanh(\pi t) + i \operatorname{sech}(\pi t) - \frac{1}{-\tanh(\pi t) + i \operatorname{sech}(\pi t)} \right] \cosh(\pi t) \\
&= \frac{1}{2i} \left[ \frac{-\cosh(\pi t) + \tanh(\pi t) \sinh(\pi t) - 2i \tanh(\pi t) - \operatorname{sech}(\pi t)}{-\tanh(\pi t) + i \operatorname{sech}(\pi t)} \right] \\
&= \frac{1}{2i} \left[ \frac{-\cosh^2(\pi t) + \sinh^2(\pi t) - 2i \sinh(\pi t) - 1}{i - \sinh(\pi t)} \right] \\
&= \frac{1 + i \sinh(\pi t)}{1 + i \sinh(\pi t)} = 1
\end{aligned}$$

Thus the case  $y = 0$  is proven.

The case  $y \neq 0$  follows easily from the previous one. Fix  $y \neq 0$  and define  $G(z) := F(z + iy)$  for  $z \in \bar{S}$ . Then  $G$  is a holomorphic function in  $S$  and continuous on  $\bar{S}$  as a composition of continuous and holomorphic functions. Moreover, the hypothesis on  $F$  yields

$$\log |G(z)| = \log |F(z + iy)| \leq A e^{\tau_0 |\operatorname{Im} z + y|} \leq A e^{\tau_0 |\operatorname{Im} z|} e^{\tau_0 |y|} \quad (20)$$

for all  $z \in \bar{S}$ . The previous case yields for  $G$  with  $A$  replaced by  $A e^{\tau_0 |y|}$

$$|G(x)| \leq \exp \left( \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[ \frac{\log |G(it)|}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log |G(1+it)|}{\cosh(\pi t) + \cos(\pi x)} \right] dt \right) \quad (21)$$

Now, observing  $G(x) = F(x + iy)$ ,  $G(it) = F(it + iy)$  and  $G(1+it) = F(1+it + iy)$  yields the desired result.  $\square$

REMARK 3.1. *Exercise 1.3.8. [Gra14, p. 48] shows that the name extension is an appropriate choice. For  $0 < x < 1$  consider*

$$\begin{aligned} \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \frac{1}{\cosh(\pi t) + \cos(\pi x)} dt &= \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \frac{1}{\frac{1}{2}(e^{\pi t} + e^{-\pi t}) + \cos(\pi x)} dt \\ &= \frac{\sin(\pi x)}{\pi} \int_0^{\infty} \frac{1}{s^2 + 2 \cos(\pi x)s + 1} ds \\ &= \frac{\sin(\pi x)}{\pi} \int_0^{\infty} \frac{1}{(s + \cos(\pi x))^2 + \sin^2(\pi x)} ds \\ &= \frac{1}{\pi \sin(\pi x)} \int_0^{\infty} \frac{1}{\left(\frac{s + \cos(\pi x)}{\sin(\pi x)}\right)^2 + 1} ds \\ &= \frac{1}{\pi} \int_{\cot(\pi x)}^{\infty} \frac{1}{u^2 + 1} du \\ &= \frac{1}{\pi} \left[ \frac{\pi}{2} - \arctan(\cot(\pi x)) \right] \\ &= x \end{aligned}$$

and in the same manner

$$\frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \frac{1}{\cosh(\pi t) - \cos(\pi x)} dt = 1 - x$$

Assume that  $F$  is holomorphic in  $S$ , continuous and bounded on  $\bar{S}$  with  $|F(z)| \leq B_0$  when  $\operatorname{Re} z = 0$  and  $|F(z)| \leq B_1$  when  $\operatorname{Re} z = 1$  for some  $0 < B_0, B_1 < \infty$ . If  $|F(z)| \leq M$  for  $0 < M < \infty$ ,  $F$  satisfies the hypothesis of lemma 3.6 with  $A := \log(M)$  and  $\tau_0 = 0$ . Therefore

$$\begin{aligned} |F(z)| &\leq \exp \left( \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[ \frac{\log |F(it + iy)|}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log |F(1+it + iy)|}{\cosh(\pi t) + \cos(\pi x)} \right] dt \right) \\ &\leq \exp \left( \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[ \frac{\log B_0}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log B_1}{\cosh(\pi t) + \cos(\pi x)} \right] dt \right) \\ &= \exp(x \log B_0 + (1-x) \log B_1) \\ &= B_0^x B_1^{1-x} \end{aligned}$$

whenever  $z := x + iy \in S$ . Hence lemma 3.6 reduces to lemma 2.1.

**3.2. Stein-Weiss Interpolation Theorem of Analytic Families of Operators.** Because of the complex nature of its proof, the Riesz-Thorin theorem 2.1 can be extended to appropriate families of linear operators  $(T_z)_{z \in \Omega}$  depending on a parameter  $z \in \Omega \subseteq \mathbb{C}$ . This result is due to Elias M. Stein and Guido Weiss. First we need to establish some common terminology.

**DEFINITION 3.1.** (Analytic family, admissible growth) *Let  $(X, \mu)$ ,  $(Y, \nu)$  be two  $\sigma$ -finite measure spaces and  $(T_z)_{z \in \bar{S}}$ , where  $T_z$  is defined  $\Sigma_X$  and taking values in the space of all measurable functions on  $Y$  such that*

$$\int_Y |T_z(\chi_A)\chi_B| d\nu \quad (22)$$

*whenever  $\mu(A), \nu(B) < \infty$ . The family  $(T_z)_{z \in \bar{S}}$  is said to be analytic if for all  $f \in \Sigma_X$ ,  $g \in \Sigma_Y$  we have that*

$$z \mapsto \int_Y T_z(f)g d\nu \quad (23)$$

*is analytic on  $S$  and continuous on  $\bar{S}$ . Further, an analytic family  $(T_z)_{z \in \bar{S}}$  is called of admissible growth, if there is a constant  $\tau_0 \in [0, \pi)$ , such that for all  $f \in \Sigma_X$ ,  $g \in \Sigma_Y$  a constant  $C(f, g)$  exists with*

$$\log \left| \int_Y T_z(f)g d\nu \right| \leq C(f, g) e^{\tau_0 |\operatorname{Im} z|} \quad (24)$$

*for all  $z \in \bar{S}$ .*

Now we are able to formulate the theorem.

**THEOREM 3.1.** (Stein-Weiss interpolation theorem of Analytic Families of Operators) *Let  $(T_z)_{z \in \bar{S}}$  be an analytic family of admissible growth,  $1 \leq p_0, p_1, q_0, q_1 \leq \infty$  and suppose that  $M_0, M_1$  are positive functions on the real line such that for some  $\tau_1 \in [0, \pi)$*

$$\sup_{-\infty < y < \infty} e^{-\tau_1 |y|} \log M_0(y) < \infty \quad \text{and} \quad \sup_{-\infty < y < \infty} e^{-\tau_1 |y|} \log M_1(y) < \infty. \quad (25)$$

*Fix  $0 < \theta < 1$  and define*

$$\frac{1}{p} := \frac{1-\theta}{p_0} + \frac{\theta}{p_1} \quad \text{and} \quad \frac{1}{q} := \frac{1-\theta}{q_0} + \frac{\theta}{q_1}. \quad (26)$$

*Further suppose that for all  $f \in \Sigma_X$  and  $y \in \mathbb{R}$  we have*

$$\|T_{iy}(f)\|_{L^{q_0}} \leq M_0(y) \|f\|_{L^{p_0}} \quad \text{and} \quad \|T_{1+iy}(f)\|_{L^{q_1}} \leq M_1(y) \|f\|_{L^{p_1}}. \quad (27)$$

Then for all  $f \in \Sigma_X$  we have

$$\|T_\theta(f)\|_{L^q} \leq M(\theta) \|f\|_{L^p}$$

where for  $0 < x < 1$

$$M(x) = \exp \left( \frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[ \frac{\log M_0(t)}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log M_1(t)}{\cosh(\pi t) + \cos(\pi x)} \right] dt \right).$$

*Proof.* Fix  $0 < \theta < 1$  and  $f \in \Sigma_X$ ,  $g \in \Sigma_Y$  with  $\|f\|_{L^p} = \|g\|_{L^{q'}} = 1$ . Define  $f_z, g_z$  as in (6) and for  $z \in \bar{S}$

$$F(z) := \int_Y T_z(f_z) g_z d\nu$$

Since the family  $(T_z)_{z \in \bar{S}}$  is of admissible growth we have that there exist constants  $c(\chi_{A_j}, \chi_{B_k})$  for any  $j = 1, \dots, n$  and  $k = 1, \dots, m$  such that

$$\log \left| \int_{B_k} T_z(\chi_{A_j}) d\nu \right| \leq c(\chi_{A_j}, \chi_{B_k}) e^{\tau_0 |\operatorname{Im} z|}$$

For shortness we will denote these constants simply by  $c(A_j, B_k)$  and get

$$\begin{aligned} \log |F(z)| &= \log \left| \sum_{j=1}^n \sum_{k=1}^m a_j^{P(z)} b_j^{Q(z)} e^{i\alpha_j} e^{i\beta_k} \int_Y T_z(\chi_{A_j})(y) \chi_{B_k}(y) d\nu(y) \right| \\ &\leq \log \left[ \sum_{j=1}^n \sum_{k=1}^m \max \{1, a_j^{p/p_0+p/p_1}\} \max \{1, b_k^{q'/q'_0+q'/q'_1}\} \left| \int_{B_k} T_z(\chi_{A_j}) d\nu \right| \right] \\ &\leq \log \left[ \sum_{j=1}^n \sum_{k=1}^m (1+a_j)^{p/p_0+p/p_1} (1+b_k)^{q'/q'_0+q'/q'_1} e^{c(A_j, B_k) e^{\tau_0 |\operatorname{Im} z|}} \right] \\ &\leq \log \left[ \sum_{j=1}^n \sum_{k=1}^m e^{\log((1+a_j)^{p/p_0+p/p_1} (1+b_k)^{q'/q'_0+q'/q'_1}) + c(A_j, B_k) e^{\tau_0 |\operatorname{Im} z|}} \right] \\ &\leq \log (mne^{\sum_{j=1}^n \sum_{k=1}^m \log((1+a_j)^{p/p_0+p/p_1} (1+b_k)^{q'/q'_0+q'/q'_1}) + c(A_j, B_k) e^{\tau_0 |\operatorname{Im} z|}}) \\ &= \log(mn) + \sum_{j=1}^n \sum_{k=1}^m \log((1+a_j)^{p/p_0+p/p_1} (1+b_k)^{q'/q'_0+q'/q'_1}) + c(A_j, B_k) e^{\tau_0 |\operatorname{Im} z|} \end{aligned}$$

since  $\tau_0 \in [0, \pi)$  and thus  $e^{\tau_0 |\operatorname{Im} z|} \geq 1$ ,  $F$  satisfies the hypotheses of the extension of Hadamard's three lines lemma 3.6 with

$$A = \log(mn) + \sum_{j=1}^n \sum_{k=1}^m \left( \frac{p}{p_0} + \frac{p}{p_1} \right) \log(1+a_j) + \left( \frac{q'}{q'_0} + \frac{q'}{q'_1} \right) \log(1+b_k) + c(A_j, B_k)$$

The same calculations as in the proof of the Riesz-Thorin interpolation theorem 2.1 yield for  $y \in \mathbb{R}$

$$\|f_{iy}\|_{L^{p_0}} = \|f\|_{L^p}^{p/p_0} = 1 = \|g\|_{L^{q'}}^{q'/q'_0} = \|g_{iy}\|_{L^{q'_0}}$$

and

$$\|f_{1+iy}\|_{L^{p_1}} = \|f\|_{L^p}^{p/p_1} = 1 = \|g\|_{L^{q'}}^{q'/q'_1} = \|g_{1+iy}\|_{L^{q'_1}}$$

Further

$$\|F(iy)\| \leq \|T_{iy}(f_{iy})\|_{L^{q_0}} \|g_{iy}\|_{L^{q'_0}} \leq M_0(y) \|f_{iy}\|_{L^{p_0}} \|g_{iy}\|_{L^{q'_0}} = M_0(y)$$

and

$$\|F(1+iy)\| \leq \|T_{1+iy}(f_{1+iy})\|_{L^{q_1}} \|g_{1+iy}\|_{L^{q'_1}} \leq M_1(y) \|f_{1+iy}\|_{L^{p_1}} \|g_{1+iy}\|_{L^{q'_1}} = M_1(y)$$

by Hölder's inequality and the hypotheses on the analytic family  $(T_z)_{z \in \bar{S}}$ . Therefore the extension of Hadamard's three lines lemma 3.6 yields

$$|F(x)| \leq \exp\left(\frac{\sin(\pi x)}{2} \int_{-\infty}^{\infty} \left[ \frac{\log M_0(t)}{\cosh(\pi t) - \cos(\pi x)} + \frac{\log M_1(t)}{\cosh(\pi t) + \cos(\pi x)} \right] dt\right) = M(x)$$

for every  $0 < x < 1$ . Furthermore observe that

$$F(\theta) = \int_Y T_\theta(f) g \, d\nu$$

and thus by [Fol99, p. 189]

$$\begin{aligned} M_q(T_\theta(f)) &= \sup \left\{ \left| \int_Y T_\theta(f) g \, d\nu \right| : g \in \Sigma_Y, \|g\|_{L^{q'}} \right\} \\ &= \sup \{ |F(\theta)| : g \in \Sigma_Y, \|g\|_{L^{q'}} \} \\ &\leq M(\theta) \end{aligned}$$

Since  $M(\theta)$  is an absolutely convergent integral (this is immediate by the growth conditions (25)) for any  $0 < \theta < 1$ ,  $M_q(T_\theta(f)) < \infty$  and thus  $M_q(T_\theta(f)) = \|T_\theta(f)\|_{L^q}$ . The general statement follows by replacing  $f$  with  $f/\|f\|_{L^p}$  when  $\|f\|_{L^p} \neq 0$ . The theorem is trivially true when  $\|f\|_{L^p} = 0$ .  $\square$

**4. The Real Method.** In this last section we are concerned with one interpolation theorem which uses real variables techniques for its proof. This stands in contrast with the complex variables techniques used for the previous two theorems. Proofs by real variables techniques are often more straight forward and do not make use of advanced theorems. Therefore the proofs are likely longer and less natural.

**4.1. The Marcinkiewicz Interpolation Theorem.** This theorem applies to sublinear operators (as well as for quasilinear operators by a slight change of the constant) which is in comparison to the linearity assumed by the other interpolation theorems less restrictive on the nature of the operator in question. Unfortunately, this theorem will not provide an exact answer to the question raised in the introduction since the constant will differ as well as the continuity property of the estimate.

**THEOREM 4.1.** (The Marcinkiewicz interpolation theorem) *Let  $(X, \mu)$ ,  $(Y, \nu)$  be two  $\sigma$ -finite measure spaces and  $0 < p_0 < p_1 \leq \infty$ . Further let  $T$  be a sublinear operator defined on*

$$L^{p_0} + L^{p_1} := \{f_0 + f_1 : f_0 \in L^{p_0}(X, \mu), f_1 \in L^{p_1}(X, \mu)\}$$

*and taking values in the space of measurable functions on  $Y$ . Assume that there exist  $0 \leq A_0, A_1 < \infty$  such that*

$$\|T(f)\|_{L^{p_0}, \infty} \leq A_0 \|f\|_{L^{p_0}} \quad \text{and} \quad \|T(f)\|_{L^{p_1}, \infty} \leq A_1 \|f\|_{L^{p_1}} \quad (28)$$

*for all  $f \in L^{p_0}$ ,  $f \in L^{p_1}$  respectively. Then for all  $p_0 < p < p_1$  and for all  $f \in L^p$  we have the estimate*

$$\|T(f)\|_{L^p} \leq A \|f\|_{L^p} \quad (29)$$

*where*

$$A := 2 \left( \frac{p}{p - p_0} + \frac{p}{p_1 - p} \right)^{1/p} A_0^{\frac{\frac{1}{p} - \frac{1}{p_1}}{\frac{1}{p_0} - \frac{1}{p_1}}} A_1^{\frac{\frac{1}{p_0} - \frac{1}{p}}{\frac{1}{p_0} - \frac{1}{p_1}}}. \quad (30)$$

*Proof.* Let us first consider the case  $p_1 < \infty$ . Fix  $f \in L^p(X, \mu)$ ,  $\alpha > 0$  and  $\delta > 0$  ( $\delta$  will be determined later). We split  $f$  using so-called *cut-off* functions, by stipulating  $f \equiv f_0(\cdot; \alpha, \delta) + f_1(\cdot; \alpha, \delta)$ , where  $f_0(\cdot; \alpha, \delta)$  is the *unbounded part* of  $f$  and  $f_1(\cdot; \alpha, \delta)$  is the *bounded part* of  $f$ , defined by

$$\begin{aligned} f_0(x; \alpha, \delta) &:= \begin{cases} f(x), & |f(x)| > \delta\alpha, \\ 0, & |f(x)| \leq \delta\alpha. \end{cases} \\ f_1(x; \alpha, \delta) &:= \begin{cases} f(x), & |f(x)| \leq \delta\alpha, \\ 0, & |f(x)| > \delta\alpha. \end{cases} \end{aligned} \quad (31)$$

for  $x \in X$ . To facilitate reading I will omit the dependency of  $f_0(\cdot; \alpha, \delta)$  and  $f_1(\cdot; \alpha, \delta)$  upon the parameters  $\alpha$  and  $\delta$  in what follows and simply write  $f_0, f_1$  respectively.

**LEMMA 4.1.** *The functions  $f_0$  and  $f_1$  defined above satisfy  $f_0 \in L^{p_0}(X, \mu)$  and  $f_1 \in L^{p_1}(X, \mu)$  respectively.*

*Proof.* Since  $p_0 < p$  we have

$$\begin{aligned}
\|f_0\|_{L^{p_0}}^{p_0} &= \int_X |f_0|^{p_0} d\mu = \int_X |f|^{p_0} \cdot \chi_{\{|f|>\delta\alpha\}} d\mu \stackrel{(\dagger)}{=} \int_{\{|f|>\delta\alpha\}} |f|^{p_0} d\mu \\
&= \int_{\{|f|>\delta\alpha\}} |f|^p |f|^{p_0-p} d\mu = \int_{\{|f|>\delta\alpha\}} \frac{|f|^p}{|f|^{p-p_0}} d\mu \\
&\leq \frac{1}{(\delta\alpha)^{p-p_0}} \int_{\{|f|>\delta\alpha\}} |f|^p d\mu = (\delta\alpha)^{p_0-p} \int_X |f|^p \cdot \chi_{\{|f|>\delta\alpha\}} d\mu \\
&\leq (\delta\alpha)^{p_0-p} \int_X |f|^p d\mu = (\delta\alpha)^{p_0-p} \|f\|_{L^p}^p < \infty
\end{aligned} \tag{32}$$

Thus  $f_0 \in L^{p_0}(X, \mu)$ . Analogously it can be checked, that  $f_1 \in L^{p_1}(X, \mu)$  by the estimate  $\|f_1\|_{L^{p_1}}^{p_1} \leq (\delta\alpha)^{p_1-p} \|f\|_{L^p}^p$ .

*Proof of the equality*  $(\dagger)$ . Assume  $\mu$  is defined on the  $\sigma$ -algebra  $\mathcal{A}$ . We have to prove that  $\{|f| > \delta\alpha\} \in \mathcal{A}$ <sup>2</sup>. Since  $f$  is complex-valued, we may write  $f \equiv \operatorname{Re} f + i\operatorname{Im} f$  and thus  $|f|^2 \equiv \operatorname{Re}^2 f + \operatorname{Im}^2 f$ . Since  $f$  is measurable by hypothesis this implies that  $\operatorname{Re} f$  and  $\operatorname{Im} f$  are measurable<sup>3</sup>. Further for measurable real-valued functions  $f, g : (X, \mathcal{A}) \rightarrow (\overline{\mathbb{R}}, \overline{\mathfrak{B}})$ <sup>4</sup> the functions  $f+g$  and  $f \cdot g$  are measurable<sup>5</sup> and thus  $|f|^2$  is measurable. Hence  $\{\operatorname{Re}^2 f + \operatorname{Im}^2 f > \lambda\} \in \mathcal{A}$ <sup>6</sup> for any  $\lambda \in \mathbb{R}$ . So especially for  $\lambda := (\delta\alpha)^2$  we have  $\{|f| > \delta\alpha\} \in \mathcal{A}$ <sup>7</sup>. In a similar manner it can also be proven that  $\{|f| \leq \delta\alpha\} \in \mathcal{A}$ . By Lemma 4.2 and the fact that  $f \cdot g$  is measurable for two measurable functions  $f, g : (X, \mathcal{A}) \rightarrow (\mathbb{C}, \mathfrak{B}^2)$ <sup>8</sup>,  $f_0$  and  $f_1$  are measurable since  $f_0 \equiv f \cdot \chi_{\{|f|>\delta\alpha\}}$  and  $f_1 \equiv f \cdot \chi_{\{|f|\leq\delta\alpha\}}$ .

One subtlety is left to clear: the  $\mu$ -integrability of either  $|f_1|^{p_0}$  or  $|f_1|^{p_1}$  requires that  $|f_0|^{p_0}$  and  $|f_1|^{p_1}$  are measurable functions. By the fact that any continuous map  $g : (X, d_X) \rightarrow (Y, d_Y)$  between metric spaces is Borel-measurable (see [Els11, p. 86]) and that the composition of measurable functions is again measurable (see [Els11, p. 87]), the measurability of either  $f_0$  or  $f_1$  follows by  $|f_0|^{p_0} \equiv \cdot^{p_0} \circ |f \cdot \chi_{\{|f|>\delta\alpha\}}|$  and  $|f_1|^{p_1} \equiv \cdot^{p_1} \circ |f \cdot \chi_{\{|f|\leq\delta\alpha\}}|$  by stipulating  $\cdot^p : (\mathbb{R}_{\geq 0}, |\cdot|) \rightarrow (\mathbb{C}, |\cdot|)$ ,  $x^p := \exp(p \log(x))$  for  $p > 0$  and  $x \in \mathbb{R}_{>0}$  and  $x^p := 0$  if  $x = 0$ .  $\square$

By lemma (4.1) we therefore have  $f \equiv f_0 + f_1 \in L^{p_0} + L^{p_1}$ .

<sup>2</sup> For  $Y \in \mathcal{A}$  the  $\mu$ -integral of  $f : X \rightarrow \mathbb{C}$  over  $Y$  is defined to be  $\int_Y f d\mu := \int_X f \cdot \chi_Y d\mu$ . For more details see [Els11, pp. 135–136].

<sup>3</sup>For a proof see [Els11, p. 106]

<sup>4</sup> $\overline{\mathfrak{B}} := \sigma(\mathbb{R})$  and  $\overline{\mathfrak{B}} = \{B \cup E : B \in \mathfrak{B}, E \subseteq \{\pm\infty\}\}$ .

<sup>5</sup>For a proof see [Els11, p. 107].

<sup>6</sup>For a proof see [Els11, pp. 105–106]

<sup>7</sup>This follows from the fact that  $x < y$  if and only if  $x^n < y^n$  for  $n \in \mathbb{N}_{>0}$  and some real numbers  $x, y > 0$  (see [Zor04, p. 119]).

<sup>8</sup>Els11, p. 107.



LEMMA 4.2. For fixed  $\alpha > 0$ , the distribution function  $d_{T(f)}(\alpha)$  obeys an upper bound of the form

$$d_{T(f)}(\alpha) \leq \left(\frac{A_0}{\alpha/2}\right)^{p_0} \|f_0\|_{L^{p_0}}^{p_0} + \left(\frac{A_1}{\alpha/2}\right)^{p_1} \|f_1\|_{L^{p_1}}^{p_1}$$

*Proof.* Since  $T$  is a sublinear operator we have  $|T(f)| = |T(f_0 + f_1)| \leq |T(f_0)| + |T(f_1)|$ . Thus for any  $y \in Y$  with  $|T(f)(y)| > \alpha$  we therefore have either  $|T(f_0)(y)| > \alpha/2$  or  $|T(f_1)(y)| > \alpha/2$ <sup>9</sup>. Hence

$$\{|T(f)| > \alpha\} \subseteq \{|T(f_0)| > \alpha/2\} \cup \{|T(f_1)| > \alpha/2\}$$

and so by the monotonicity and subadditivity property of the measure  $\mu$  we have

$$\begin{aligned} d_{T(f)}(\alpha) &= \mu(\{|T(f)| > \alpha\}) \\ &\leq \mu(\{|T(f_0)| > \alpha/2\} \cup \{|T(f_1)| > \alpha/2\}) \\ &\leq \mu(\{|T(f_0)| > \alpha/2\}) + \mu(\{|T(f_1)| > \alpha/2\}) \\ &= d_{T(f_0)}(\alpha/2) + d_{T(f_1)}(\alpha/2) \end{aligned} \tag{33}$$

Now by hypothesis (??) we can estimate  $d_{T(f_0)}(\alpha/2)$  as follows

$$\begin{aligned} d_{T(f_0)}(\alpha/2) &= \left(\frac{\alpha/2}{\alpha/2}\right)^{p_0} d_{T(f_0)}(\alpha/2) \\ &\leq \left(\frac{1}{\alpha/2}\right)^{p_0} \left[ \sup \left\{ \gamma d_{T(f_0)}(\gamma)^{1/p_0} : \gamma > 0 \right\} \right]^{p_0} \\ &= \left(\frac{1}{\alpha/2}\right)^{p_0} \|T(f_0)\|_{L^{p_0, \infty}}^{p_0} \\ &\leq \left(\frac{A_0}{\alpha/2}\right)^{p_0} \|f_0\|_{L^{p_0}}^{p_0} \end{aligned} \tag{34}$$

Analogously, we get  $d_{T(f_1)}(\alpha/2) \leq \left(\frac{A_1}{\alpha/2}\right)^{p_1} \|f_1\|_{L^{p_1}}^{p_1}$  by hypothesis (??).  $\square$

By

$$\int_0^{\frac{1}{\delta}|f|} \alpha^{p-p_0-1} d\lambda = \begin{cases} \frac{1}{p-p_0} \frac{1}{\delta^{p-p_0}} |f|^{p-p_0}, & p \geq p_0 + 1 \\ = \lim_{\omega \rightarrow 0^+} \int_{\omega}^{\frac{1}{\delta}|f|} \alpha^{p-p_0-1} d\lambda \\ = \lim_{\omega \rightarrow 0^+} \left[ \frac{1}{p-p_0} \alpha^{p-p_0} \right]_{\omega}^{\frac{1}{\delta}|f|} \\ = \frac{1}{p-p_0} \left[ \frac{1}{\delta^{p-p_0}} |f|^{p-p_0} - \lim_{\omega \rightarrow 0^+} \omega^{p-p_0} \right] \\ = \frac{1}{p-p_0} \frac{1}{\delta^{p-p_0}} |f|^{p-p_0}, & p_0 < p < p_0 + 1 \end{cases} \tag{35}$$

<sup>9</sup>Without loss of generality assume  $|T(f_0)(y)| \leq |T(f_1)(y)|$ . Then we have  $\alpha < |T(f)(y)| \leq |T(f_0)(y)| + |T(f_1)(y)| \leq 2|T(f_1)(y)|$  (this is possible since  $\mathbb{R}$  is an ordered field).

and

$$\begin{aligned}
 \int_{\frac{1}{\delta}|f|}^{\infty} \alpha^{p-p_1-1} d\lambda &= \lim_{\omega \rightarrow \infty} \left[ \frac{1}{p-p_1} \alpha^{p-p_1} \right]_{\frac{1}{\delta}|f|}^{\omega} \\
 &= \frac{1}{p-p_1} \left[ \lim_{\omega \rightarrow \infty} \omega^{p-p_1} - \frac{1}{\delta^{p-p_1}} |f|^{p-p_1} \right] \\
 &= \frac{1}{p_1-p} \frac{1}{\delta^{p-p_1}} |f|^{p-p_1}
 \end{aligned} \tag{36}$$

and the representation  $\|f\|_{L^p}^p = p \int_0^{\infty} \alpha^{p-1} d_f(\alpha) d\lambda$  for  $0 < p < \infty$  we get

$$\begin{aligned}
 \|T(f)\|_{L^p}^p &= p \int_0^{\infty} \alpha^{p-1} d_{T(f)} d\lambda \\
 &\leq p (2A_0)^{p_0} \int_0^{\infty} \alpha^{p-p_0-1} \int_{\{|f|>\delta\alpha\}} |f|^{p_0} d\mu d\lambda \\
 &\quad + p (2A_1)^{p_1} \int_0^{\infty} \alpha^{p-p_1-1} \int_{\{|f|\leq\delta\alpha\}} |f|^{p_1} d\mu d\lambda \\
 &= p (2A_0)^{p_0} \int_{\{|f|>0\}} |f|^{p_0} \int_0^{\frac{1}{\delta}|f|} \alpha^{p-p_0-1} d\lambda d\mu \\
 &\quad + p (2A_0)^{p_0} \int_{\{|f|=0\}} |f|^{p_0} \int_0^{\frac{1}{\delta}|f|} \alpha^{p-p_0-1} d\lambda d\mu \\
 &\quad + p (2A_1)^{p_1} \int_X |f|^{p_1} \int_{\frac{1}{\delta}|f|}^{\infty} \alpha^{p-p_1-1} d\lambda d\mu \\
 &= p (2A_0)^{p_0} \int_X |f|^{p_0} \int_0^{\frac{1}{\delta}|f|} \alpha^{p-p_0-1} d\lambda d\mu \\
 &\quad + p (2A_1)^{p_1} \int_X |f|^{p_1} \int_{\frac{1}{\delta}|f|}^{\infty} \alpha^{p-p_1-1} d\lambda d\mu \\
 &= \frac{p (2A_0)^{p_0}}{p-p_0} \frac{1}{\delta^{p-p_0}} \int_X |f|^{p_0} |f|^{p-p_0} d\mu \\
 &\quad + \frac{p (2A_1)^{p_1}}{p_1-p} \frac{1}{\delta^{p-p_1}} \int_X |f|^{p_1} |f|^{p-p_1} d\mu \\
 &= p \left( \frac{(2A_0)^{p_0}}{p-p_0} \frac{1}{\delta^{p-p_0}} + \frac{(2A_1)^{p_1}}{p_1-p} \delta^{p_1-p} \right) \|f\|_{L^p}^p
 \end{aligned} \tag{37}$$

We pick  $\delta > 0$  such that  $(2A_0)^{p_0} \delta^{p_0-p} = (2A_1)^{p_1} \delta^{p_1-p}$ . Solving for  $\delta$  yields

$$\delta = \frac{1}{2} \left( \frac{A_0}{A_1} \right)^{p_1/(p_1-p_0)} \quad (38)$$

Substituting this in estimate (37) leads to

$$\begin{aligned} \|T(f)\|_{L^p}^p &\leq p \left( \frac{(2A_0)^{p_0}}{p-p_0} \frac{2^{p-p_0} A_1^{\frac{p_1(p-p_0)}{p_1-p_0}}}{A_0^{\frac{p_0(p-p_0)}{p_1-p_0}}} + \frac{(2A_1)^{p_1}}{p_1-p} \frac{A_0^{\frac{p_0(p_1-p)}{p_1-p_0}}}{2^{p_1-p} A_1^{\frac{p_1(p_1-p)}{p_1-p_0}}} \right) \|f\|_{L^p}^p \\ &= 2^p p \left( \frac{A_0^{\frac{p_0(p_1-p)}{p_1-p_0}} A_1^{\frac{p_1(p-p_0)}{p_1-p_0}}}{p-p_0} + \frac{A_0^{\frac{p_0(p_1-p)}{p_1-p_0}} A_1^{\frac{p_1(p-p_0)}{p_1-p_0}}}{p_1-p} \right) \|f\|_{L^p}^p \end{aligned} \quad (39)$$

And taking the  $p$ -th power further

$$\begin{aligned} \|T(f)\|_{L^p} &\leq 2 \left( \frac{p}{p-p_0} + \frac{p}{p_1-p} \right)^{1/p} A_0^{\frac{p_0(p_1-p)}{p(p_1-p_0)}} A_1^{\frac{p_1(p-p_0)}{p(p_1-p_0)}} \|f\|_{L^p} \\ &= 2 \left( \frac{p}{p-p_0} + \frac{p}{p_1-p} \right)^{1/p} A_0^{\frac{p_0(p_1-p)}{p(p_1-p_0)} \frac{p_1}{p_1}} A_1^{\frac{p_1(p-p_0)}{p(p_1-p_0)} \frac{p_0}{p_0}} \|f\|_{L^p} \\ &= 2 \left( \frac{p}{p-p_0} + \frac{p}{p_1-p} \right)^{1/p} A_0^{\frac{\frac{p_1-p}{p p_1}}{\frac{p_1-p_0}{p_0 p_1}}} A_1^{\frac{\frac{p-p_0}{p_0 p}}{\frac{p_1-p_0}{p_0 p_1}}} \|f\|_{L^p} \\ &= 2 \left( \frac{p}{p-p_0} + \frac{p}{p_1-p} \right)^{1/p} A_0^{\frac{\frac{1}{p} - \frac{1}{p_1}}{\frac{1}{p_0} - \frac{1}{p_1}}} A_1^{\frac{\frac{1}{p_0} - \frac{1}{p}}{\frac{1}{p_0} - \frac{1}{p_1}}} \|f\|_{L^p} \end{aligned} \quad (40)$$

Assume  $p_1 = \infty$ . We again use the cut-off functions defined in (31) to decompose  $f$ . Since  $\{|f_1| > \delta\alpha\} = \emptyset$ , we have

$$\|T(f_1)\|_{L^\infty} \leq A_1 \|f_1\|_{L^\infty} = A_1 \inf \{B > 0 : \mu(\{|f_1| > B\}) = 0\} \leq A_1 \delta\alpha = \alpha/2$$

Provided we stipulate  $\delta := 1/(2A_1)$ . Therefore the set  $\{|T(f_1)| > \alpha/2\}$  has measure zero (this is immediate since  $\|T(f_1)\|_{L^\infty} = \inf \{B > 0 : \mu(\{|T(f_1)| > B\}) = 0\} \leq \alpha/2$  and any subset of a set with measure zero has itself measure zero). Thus similar to part **b.** of (i.) we get  $d_{T(f)}(\alpha) \leq d_{T(f_0)}(\alpha/2)$ .

Hypothesis (??) yields the estimate  $d_{T(f_0)}(\alpha/2) \leq \left( \frac{A_0}{\alpha/2} \right)^{p_0} \int_{\{2A_1|f|>\alpha\}} |f|^{p_0} d\mu$ .

Thus by **a.** and **b.**

$$\begin{aligned}
\|T(f)\|_{L^p}^p &= p \int_0^\infty \alpha^{p-1} d_{T(f)} \lambda \\
&\leq p(2A_0)^{p_0} \int_0^\infty \alpha^{p-p_0-1} \int_{\{2A_1|f|>\alpha\}} |f|^{p_0} d\mu d\lambda \\
&= p(2A_0)^{p_0} \int_X |f|^{p_0} \int_0^{2A_1|f|} \alpha^{p-p_0-1} d\lambda d\mu \\
&= \frac{2^p p A_0^{p_0} A_1^{p-p_0}}{p-p_0} \int_X |f|^p d\mu \\
&= \frac{2^p p A_0^{p_0} A_1^{p-p_0}}{p-p_0} \|f\|_{L^p}^p
\end{aligned} \tag{41}$$

That the constant  $2^p p A_0^{p_0} A_1^{p-p_0} / (p-p_0)$  found in (41) is the  $p$ -th power of the one stated in the theorem can be seen by passing the constant (30) to the limit  $p_1 \rightarrow \infty$ :

$$\begin{aligned}
\lim_{p_1 \rightarrow \infty} A &= \lim_{p_1 \rightarrow \infty} \left[ 2 \left( \frac{p}{p-p_0} + \frac{p}{p_1-p} \right)^{1/p} A_0^{\frac{\frac{1}{p}-\frac{1}{p_1}}{\frac{1}{p_0}-\frac{1}{p_1}}} A_1^{\frac{\frac{1}{p_0}-\frac{1}{p}}{\frac{1}{p_0}-\frac{1}{p_1}}} \right] \\
&= 2 \exp \left[ \frac{1}{p} \log \left( \frac{p}{p-p_0} + \lim_{p_1 \rightarrow \infty} \frac{1}{p_1} \frac{p}{1-p \lim_{p_1 \rightarrow \infty} \frac{1}{p_1}} \right) \right] \\
&\quad \cdot \lim_{p_1 \rightarrow \infty} A_0^{\frac{\frac{1}{p}-\frac{1}{p_1}}{\frac{1}{p_0}-\frac{1}{p_1}}} \cdot \lim_{p_1 \rightarrow \infty} A_1^{\frac{\frac{1}{p_0}-\frac{1}{p}}{\frac{1}{p_0}-\frac{1}{p_1}}} \\
&= 2 \left( \frac{p}{p-p_0} \right)^{1/p} \exp \left[ \frac{\frac{1}{p} - \lim_{p_1 \rightarrow +\infty} \frac{1}{p_1}}{\frac{1}{p_0} - \lim_{p_1 \rightarrow \infty} \frac{1}{p_1}} \log(A_0) \right] \\
&\quad \cdot \exp \left[ \frac{\frac{1}{p_0} - \frac{1}{p}}{\frac{1}{p_0} - \lim_{p_1 \rightarrow \infty} \frac{1}{p_1}} \log(A_1) \right] \\
&= 2 \left( \frac{p}{p-p_0} \right)^{1/p} A_0^{\frac{p_0}{p}} A_1^{1-\frac{p_0}{p}}
\end{aligned}$$

□

## Appendix A. Limit superior and limit inferior revisited

DEFINITION A.1. Let  $(X, d)$  a metric space,  $E \subseteq X$ ,  $f : E \rightarrow \mathbb{R}$  and  $a \in X$  be a limit point of  $E$ . Then we define the upper limit of  $f$  at  $a$  as

$$\limsup_{x \rightarrow a} f(x) := \lim_{\varepsilon \downarrow 0} \left[ \sup \left\{ f(x) : x \in E \cap \dot{B}_\varepsilon(a) \right\} \right]$$

and the lower limit of  $f$  at  $a$  as

$$\liminf_{x \rightarrow a} f(x) := - \limsup_{x \rightarrow a} (-f)(x)$$

PROPOSITION A.1. Let  $(X, d)$  a metric space,  $E \subseteq X$ ,  $f, g : E \rightarrow \mathbb{R}$ , where  $f$  is bounded and  $a \in X$  be a limit point of  $E$ . Then

$$\limsup_{x \rightarrow a} (fg)(x) = \limsup_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$$

whenever both sides exist and  $\lim_{x \rightarrow a} g(x) \geq 0$ .

*Proof.* Write

$$fg = f \lim_{x \rightarrow a} g(x) + f \left[ g - \lim_{x \rightarrow a} g(x) \right]$$

By [Bou95, p. 358] we have

$$\begin{aligned} \limsup_{x \rightarrow a} (fg)(x) &= \limsup_{x \rightarrow a} \left( f(x) \lim_{x \rightarrow a} g(x) + f(x) \left[ g(x) - \lim_{x \rightarrow a} g(x) \right] \right) \\ &= \limsup_{x \rightarrow a} \left( f(x) \lim_{x \rightarrow a} g(x) \right) + \lim_{x \rightarrow a} \left( f(x) \left[ g(x) - \lim_{x \rightarrow a} g(x) \right] \right) \\ &= \limsup_{x \rightarrow a} \left( f(x) \lim_{x \rightarrow a} g(x) \right) \end{aligned}$$

since  $\lim_{x \rightarrow a} [g(x) - \lim_{x \rightarrow a} g(x)] = 0$  and  $f$  is bounded. Fix  $\varepsilon > 0$ . Further by [Bou95, p. 357] and  $\lim_{x \rightarrow a} g(x) \geq 0$

$$\sup \left\{ f(x) \lim_{x \rightarrow a} g(x) : x \in E \cap \dot{B}_\varepsilon(a) \right\} = \sup \left\{ f(x) : x \in E \cap \dot{B}_\varepsilon(a) \right\} \lim_{x \rightarrow a} g(x)$$

Hence

$$\begin{aligned}
\limsup_{x \rightarrow a} (fg)(x) &= \limsup_{x \rightarrow a} \left( f(x) \lim_{x \rightarrow a} g(x) \right) \\
&= \lim_{\varepsilon \downarrow 0} \left[ \sup \left\{ f(x) \lim_{x \rightarrow a} g(x) : x \in E \cap \dot{B}_\varepsilon(a) \right\} \right] \\
&= \lim_{\varepsilon \downarrow 0} \left[ \sup \left\{ f(x) : x \in E \cap \dot{B}_\varepsilon(a) \right\} \right] \lim_{x \rightarrow a} g(x) \\
&= \limsup_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)
\end{aligned} \tag{42}$$

□

## Appendix B. Measure Theory

Let  $(X, \mu)$  be a measure space. Recall, that if for each measurable set  $E$  with  $\mu(E) = \infty$  there exists a measurable set  $F \subseteq E$  and  $0 < \mu(F) < \infty$ ,  $\mu$  is called *semifinite*.

LEMMA B.1. *Every  $\sigma$ -finite measure is semifinite.*

*Proof.* Let  $X = \bigcup_{n \in \mathbb{N}} X_n$  where  $\mu(X_n) < \infty$  and  $E$  be measurable with  $\mu(E) = \infty$ . By letting  $Y_n := \bigcup_{k \leq n} X_k$ ,  $Y_n$  is an increasing sequence. Then  $E \cap Y_n$  is measurable and since  $E \cap Y_n \subseteq Y_n$ ,  $\mu(E \cap Y_n) < \infty$  for each  $n \in \mathbb{N}$ . By the continuity from below (see [Coh13, p. 10] or [Fol99, p. 26]) we have

$$\infty = \mu(E) = \mu(E \cap X) = \mu \left( E \cap \left( \bigcup_{n \in \mathbb{N}} Y_n \right) \right) = \mu \left( \bigcup_{n \in \mathbb{N}} (E \cap Y_n) \right) = \lim_{n \rightarrow \infty} \mu(E \cap Y_n)$$

Hence for every  $C > 0$  there exists  $N \in \mathbb{N}$ , such that  $\infty > \mu(E \cap Y_n) > C$  for  $n > N$ . □

## References

- [Ahl79] Lars V. Ahlfors. *Complex Analysis*. Third Edition. Mc Graw Hill Education, 1979.
- [Bou95] Nicolas Bourbaki. *General Topology - Chapters 1-4*. Elements of Mathematics. Springer-Verlag Berlin Heidelberg, 1995.
- [Coh13] Donald L. Cohn. *Measure Theory*. Second edition. Springer, 2013.
- [Els11] Jürgen Elstrodt. *Mass- und Integrationstheorie*. 7., korrigierte und aktualisierte Auflage. Springer Verlag, 2011.
- [Fol99] Gerald B. Folland. *Real Analysis*. Second Edition. John Wiley & Sons, Inc., 1999.
- [Gra14] Loukas Grafakos. *Classical Fourier Analysis*. Third Edition. Springer Science + Business Media New York, 2014.
- [HS91] Murali Rao and Henrik Stetkær. *Complex Analysis - An Invitation*. World Scientific, 1991.

- [Rud87] Walter Rudin. *Real and Complex Analysis*. Third Edition. McGraw-Hill Book Company, 1987.
- [Zor04] Vladimir A. Zorich. *Mathematical Analysis I*. Springer-Verlag Berlin Heidelberg, 2004.