1. MEDICAL IMAGE DISCLOSURE

The image shows a red, circular rash on the forearm with defined borders, consistent with localized environmental exposure. Inspect skin surface textures completely before administering a covering dressing.

2. CRITICAL LIFE PRESERVATION COMMANDS

EMERGENCY CONDITION	IMMEDIATE STEP-BY-STEP FIELD COMMAND	THREE DISTINCT POSSIBLE ISSUES
SEVERE HEMORRHAGE	Apply immediate, firm, direct pressure to the wound with sterile gauze. If bleeding continues on a limb, place a mechanical tourniquet 2 to 3 inches above the injury site (never on a joint), tighten the windlass until bleeding stops, and mark placement time.	Arterial laceration, venous tear, or capillary abrasion.
RESPIRATORY COMPROMISE	Tilt the head back gently, lift the chin, and clear the airway path. Position unconscious, breathing individuals on their side in the lateral recovery position.	Foreign object obstruction, allergic anaphylaxis, or fluid aspiration.
ENVIRONMENTAL COLD	Strip wet clothes, wrap core tightly in dry sheets/wool blankets, and insulate body off cold concrete/dirt floors.	Mild hypothermia, severe frostbite, or trench foot syndrome.

3. GENERAL REMEDIES MATRIX

Follow these non-personalized treatment options to manage minor trauma or superficial injuries:

■ **Option 1:** Clean localized wounds thoroughly with sterile water and apply a protective topical antibiotic barrier.

■ **Option 2:** Apply a firm, non-elastic compression wrap to manage joint stability and limit swelling during transit.

■ **Option 3:** Utilize cold water compresses to lower localized skin swelling or surface thermal damage.

4. MEDICATION LABEL VERIFICATION & CONSENSUS DOSAGE CAPS

■ Make sure to double-check the physical label to confirm product names, active ingredients, expiration timelines, and manufacturer values are accurate before administering any medication.

■ The typical over-the-counter adult dose of ibuprofen for temporary pain or inflammation management is 200–400 mg every 4–6 hours, strictly not exceeding 1,200 mg in any 24-hour period without a doctor's direct supervision.

■ **Rigid Limitation:** Do not calculate or recommend personalized chemical dosages based on individual biometrics, and never provide dosage parameters or modifications for infants.

CRITICAL CLINICAL PROTECTION PANEL
This is for informational purposes only. For medical advice or diagnosis, consult a professional. AI responses may include mistakes.

AUTHORITATIVE REFERENCES: American Red Cross First Aid Protocols, Centers for Disease Control and Prevention (CDC) Emergency Medical Guidance.