

INTRO TO QUANTUM COMPUTING

LECTURE #14

# The Qubit & Bloch Sphere

**Amir Karamlou** 

02/07/2021





### © 2020 The Coding School All rights reserved

Use of this recording is for personal use only. Copying, reproducing, distributing, posting or sharing this recording in any manner with any third party are prohibited under the terms of this registration. All rights not specifically licensed under the registration are reserved.





# ANNOUNCEMENTS

### **TODAY'S LECTURE**

→ Use the Bloch Sphere to conceptualize the state of a qubit

→ View quantum gates as actions on the Bloch sphere





# What is a qubit?

- Building block of quantum computers
- A two-level system
- Can be in a superposition of two values





### **Qubit Reviewed**

**Superposition:** a qubit can be  $|0\rangle$  and  $|1\rangle$  at the same time!

This is how we show it:  $|\psi\rangle = \alpha |0\rangle + \beta |1\rangle$ 

$$|0\rangle \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$|1\rangle \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$






# **Complex Numbers Review**

A complex number consists of both a *real* and *imaginary* component.

$$z = a + i b$$

$$z^* = a - i b$$


$$|z|^2 = z \cdot z^* = z^* \cdot z$$







#### **EULER'S FORMULA & COMPLEX EXPONENTIALS**



**Euler's formula:** 

$$e^{i\varphi} = \cos\varphi + i\sin\varphi$$

Polar representation of complex numbers!

$$z = x + iy = |z|(\cos \varphi + i \sin \varphi) = re^{i\varphi}$$

$$r = |z| = \sqrt{x^2 + y^2}$$
  $\varphi = \tan^{-1}\left(\frac{y}{x}\right)$  (vector radius)





# What is phase?

 $e^{i \varphi}$ 






In quantum mechanics global phase has **NO** <u>physical</u> <u>significance</u> and **NO** <u>effect</u> therefore can be ignored!!

$$e^{i\varphi} | \psi \rangle \rightarrow | \psi \rangle$$





The **state** of a qubit can be represented as a point on the Bloch Sphere







# Qubit

Back to our qubit:  $|\psi\rangle = \alpha |0\rangle + \beta |1\rangle$ 






$$|\psi\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)e^{i\varphi}|1\rangle$$

North pole: |0>

South pole: |1>

Classical bits are only on the North pole or the South pole








$$|\psi\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)e^{i\varphi}|1\rangle$$

What is the state of the qubit based on the position on the Bloch sphere?

 $\theta$ =0:

 $\theta = \pi$ :






$$|\psi\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)e^{i\varphi}|1\rangle$$

More examples:

$$\theta$$
=  $\pi$  /2 ,  $\phi$ =0:

$$\theta$$
= $\pi$ /2,  $\phi$ = $\pi$ :











# 10 MIN BREAK!

### Quantum gates

- Maps one the quantum state to another quantum state
- Moves a point on the Bloch Sphere to another point
- We can represent them with matrices







#### Pauli gates:

$$\sigma_{x} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$
 Pauli-X operator (X)

$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$
 Pauli-Y operator (Y)


$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$
 Pauli-Z operator (Z)





$$\sigma_{x} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Pauli-X operator (X)




Rotates the state around the **X** axis by 180 degrees





$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$
 Pauli-Z operator (Z)




Rotates the state around the **Z** axis by 180 degrees





$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$
 Pauli-Y operator (Y)




Rotates the state around the <u>Y</u> axis by 180 degrees





We're not just limited by 180 degree rotations

For example we can rotate around the x axis by 90 degrees







We can rotate by an angle  $\theta$  around x or y or z axis:

$$R_{\chi}(\theta) = e^{-i\frac{\theta}{2}\sigma_{\chi}}$$

$$R_{y}(\theta) = e^{-i\frac{\theta}{2}\sigma_{y}}$$

$$R_z(\theta) = e^{-i\frac{\theta}{2}\sigma_z}$$





Let's take rotation by an angle  $\theta$  around the x axis as an example:

$$R_{x}(\theta) = e^{-i\frac{\theta}{2}\sigma_{x}} = \cos\frac{\theta}{2}I - i\sin\frac{\theta}{2}\sigma_{x}$$

Matrix representation?





Let's take rotation by an angle  $\theta$  around the x axis as an example:


$$R_z(\theta) = e^{-i\frac{\theta}{2}\sigma_z} = \cos\frac{\theta}{2}I - i\sin\frac{\theta}{2}\sigma_z$$

Matrix representation?






Examples:







Examples:









# **Experimental Implementations**

- All quantum gates are engineered using rotations on the Bloch sphere
- In our quantum hardware we can control each transition with some level of accuracy
- Example: Rabi oscillation:









### **Quantum Universality**

Any quantum computation operation can be made by using a combination of:

{CNOT, H, S, T}






# **Quantum Universality**

What is the S gate?

What is the T gate?











### © 2020 The Coding School All rights reserved

Use of this recording is for personal use only. Copying, reproducing, distributing, posting or sharing this recording in any manner with any third party are prohibited under the terms of this registration. All rights not specifically licensed under the registration are reserved.