


© 2020 The Coding School
All rights reserved

Use of this recording is for personal use only. Copying, reproducing, distributing, posting or sharing this recording in any manner with any third party are prohibited under the terms of this registration. All rights not specifically licensed under the registration are reserved.




INTRO TO QUANTUM COMPUTING

LECTURE #15

QUANTUM COMPUTATION PT. 3 : QUANTUM CIRCUIT MATHEMATICS

FRANCISCA VASCONCELOS

2/28/2021



ANNOUNCEMENTS

QUANTUM COMPUTATION LECTURE SERIES

Lecture 1 – The Quantum Circuit Model



How can we perform computation with quantum systems?

CONCEPTS

Lecture 2 – Qiskit Tutorial



How can we program quantum circuits?

PROGRAMMING

Lecture 3 – Quantum Circuit Mathematics



How can we represent quantum circuits mathematically?

MATH


Lectures 4-6 – Introductory Quantum Protocols and Algorithms

How can we leverage quantum for cryptography, teleportation, and algorithms?

APPLICATION

TODAY'S LECTURE

- 1. Quantum Circuit Model Review
- 2. Single-Qubit Circuit Math
 - a) Single-Qubit States Math
 - b) Single-Qubit Gates Math
 - c) Single-Qubit Measurement Math
 - d) Putting it All Together!
- 3. Multi-Qubit Circuit Math
 - a) The Tensor Product
 - b) Multi-Qubit States Math
 - c) Multi-Qubit Gates Math
 - d) Multi-Qubit Measurement Math
 - e) Putting it All Together!



QUANTUM CIRCUIT MODEL REVIEW

THE QUANTUM CIRCUIT MODEL


The quantum circuit is a theoretical model for quantum computation. It has three key components:

**STATES,
GATES,
& MEASUREMENT**



Now, how do we put these all together to create a quantum circuit?

THE QUANTUM CIRCUIT MODEL



PROGRAMMING CIRCUITS IN QISKIT...


Last class, you learned how to program these circuits in Qiskit!

Construct Bell state

In [17]:

```
1 bell = QuantumCircuit(2, 2)
2 bell.h(1)
3 bell.cx(1, 0)
4 bell.measure(range(2), range(2))
5 bell.draw()
```


Out[17]:




What math are we learning today and why do we need it ???



SINGLE-QUBIT CIRCUIT MATH



THE SINGLE-QUBIT CIRCUIT



```
1 qc_z = QuantumCircuit(1, 1)
2 qc_z.h(0)
3 qc_z.z(0)
4 qc_z.h(0)
5 qc_z.measure(0, 0)
6 qc_z.draw()
```




REMEMBER, MATHEMATICALLY...

Quantum states are complex vectors!

$$|\psi\rangle = \begin{pmatrix} |0\rangle \\ |1\rangle \end{pmatrix} = \alpha|0\rangle + \beta|1\rangle$$

Quantum gates are complex (unitary) matrices!



$$|\psi\rangle = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad \psi_1, \psi_2 \in \mathbb{C}$$

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi} \sin\left(\frac{\theta}{2}\right)|1\rangle = \begin{pmatrix} \cos(\theta/2) \\ e^{i\phi} \sin(\theta/2) \end{pmatrix}$$

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad a_{..} \in \mathbb{C}$$

$$A = \begin{pmatrix} \cos(\theta/2) & -e^{i\lambda} \sin(\theta/2) \\ e^{i\phi} \sin(\theta/2) & e^{i\lambda+i\phi} \cos(\theta/2) \end{pmatrix}$$

REMEMBER, MATHEMATICALLY...

Quantum states are complex vectors!

$$|\psi\rangle \text{———}$$

Quantum gates are complex (unitary) matrices!

$$\text{——} \boxed{A} \text{——}$$

$$|\psi\rangle = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}, \quad \psi_1, \psi_2 \in \mathbb{C}$$

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi} \sin\left(\frac{\theta}{2}\right)|1\rangle = \begin{pmatrix} \cos(\theta/2) \\ e^{i\phi} \sin(\theta/2) \end{pmatrix}$$

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad a_{..} \in \mathbb{C}$$

$$A = \begin{pmatrix} \cos(\theta/2) & -e^{i\lambda} \sin(\theta/2) \\ e^{i\phi} \sin(\theta/2) & e^{i\lambda+i\phi} \cos(\theta/2) \end{pmatrix}$$

REVIEW OF SINGLE QUBIT STATES

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$|+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$


$$|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$|-\rangle = \frac{|0\rangle - |1\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$


REVIEW OF SINGLE QUBIT GATES




$$X = \begin{matrix} \rightarrow |0\rangle\langle 1| + |1\rangle\langle 0| \\ = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$



$$Z = \begin{matrix} \rightarrow |0\rangle\langle 0| - |1\rangle\langle 1| \\ = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \end{matrix}$$




$$Y = \begin{matrix} \rightarrow i|0\rangle\langle 1| + i|1\rangle\langle 0| \\ = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \end{matrix}$$



$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$\begin{aligned} H|0\rangle &= |\leftarrow\rangle \\ H|1\rangle &= |\rightarrow\rangle \end{aligned}$$

IN A CIRCUIT...



$$|\Phi\rangle = A |\psi\rangle$$

$$|\Phi\rangle = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

```

1 qc_z = QuantumCircuit(1, 1)
2 qc_z.h(0)

```


$|q\rangle = |0\rangle$

$$H|q\rangle = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = |+\rangle$$

$H|0\rangle = |+\rangle$

AND IN A LARGER CIRCUIT...



$$C(B(A|\psi\rangle))$$

$|\Phi\rangle$

$|B\rangle$

$|\psi'\rangle = \underline{CBA} |\psi\rangle$

```

1 qc_z = QuantumCircuit(1, 1)
2 qc_z.h(0)
3 qc_z.z(0)
4 qc_z.h(0)

```


$q - \boxed{H} - \boxed{Z} - \boxed{H} - |\Phi\rangle$

$HZH|q\rangle \leftarrow |q\rangle = |0\rangle$

$= \left(\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right) \left[\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right] \left(\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right) \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

$|\Phi\rangle = \begin{bmatrix} A \\ B \end{bmatrix}$

ORDER MATTERS!




$$\begin{aligned} & 5 \times 4 \times 1 \\ & = 1 \times 4 \times 5 \\ & = 4 \times 1 \times 5 \end{aligned}$$

$CBA \neq BAC$
generally

Heisenberg Uncertainty Prin.
Commutators

QUANTUM PRACTICE TIME!




$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$


$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

QUANTUM PRACTICE SOLUTION!



$$\begin{aligned} H \times |0\rangle &= \frac{1}{\sqrt{2}} \left(\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &= \frac{1}{\sqrt{2}} \left(\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) \\ &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = |-\rangle \end{aligned}$$

SO, WHAT IS MEASUREMENT?



Assume we are measuring
in the Z-basis!

$$|\phi\rangle = CBA |\psi\rangle$$

$$|\phi\rangle = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

prob. amp. of measuring
the state $|0\rangle$

prob. amp. of measuring
the state $|1\rangle$





$$P(\text{meas. } |0\rangle) = |\alpha|^2$$

$$P(\text{meas. } |1\rangle) = |\beta|^2$$



QUANTUM PRACTICE TIME!


Solve for the likelihoods of measuring $|0\rangle$ and $|1\rangle$ for each of the following quantum circuits...



QUANTUM PRACTICE SOLUTIONS!

Solve for the likelihoods of measuring $|0\rangle$ and $|1\rangle$ for each of the following quantum circuits...


(1)



$|0\rangle \rightarrow 100\%$

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \rightarrow \begin{aligned} P(\text{meas. } |0\rangle) &= |1|^2 = 1 \rightarrow 100\% \\ P(\text{meas. } |1\rangle) &= |0|^2 = 0 \rightarrow 0\% \end{aligned}$$

(2)



$$|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rightarrow \begin{aligned} P(\text{meas. } |1\rangle) &= 1 (100\%) \\ P(\text{meas. } |0\rangle) &= 0 (0\%) \end{aligned}$$

QUANTUM PRACTICE SOLUTIONS!

Solve for the likelihoods of measuring $|0\rangle$ and $|1\rangle$ for each of the following quantum circuits...

(3) $|0\rangle \xrightarrow{H} |+\rangle$

$H|0\rangle = |+\rangle$

$|+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \rightarrow P(\text{meas. } |0\rangle) = \left|\frac{1}{\sqrt{2}}\right|^2 = \frac{1}{2} \text{ 50\%}$

\downarrow 50%

$P(\text{meas. } |1\rangle) = \left|\frac{1}{\sqrt{2}}\right|^2 = \frac{1}{2} \text{ 50\%}$

(4) $|0\rangle \xrightarrow{X} |1\rangle \xrightarrow{H} |-\rangle$

$H|1\rangle = |-\rangle$

$|-\rangle = \frac{|0\rangle - |1\rangle}{\sqrt{2}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \rightarrow P(\text{meas. } |0\rangle) = \left|\frac{1}{\sqrt{2}}\right|^2 = \frac{1}{2}$

\downarrow 50%

$P(\text{meas. } |1\rangle) = \left|-\frac{1}{\sqrt{2}}\right|^2 = \frac{1}{2} \text{ 50\%}$

SOME OBSERVATIONS...

- Since matrix multiplication results in a new matrix, a series of gates can be represented by a single matrix, which is just the product of all the gate matrices...



$$|\psi\rangle \xrightarrow{\text{--- A --- B --- C ---}} = |\psi\rangle \xrightarrow{\boxed{D = CBA}}$$

- The “wire” without a gate is equivalent to applying the identity matrix/operation!

$$|\psi\rangle \xrightarrow{\text{---}} = |\psi\rangle \xrightarrow{\boxed{I}} \quad \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

- Experimentally, circuits generally have to start in the $|0\rangle$ state. So, while you can start your circuit in any state theoretically, it is good to keep in mind that experimentally this will involve applying gates.

$$|1\rangle \xrightarrow{\text{---}} = |0\rangle \xrightarrow{\boxed{X}} \xrightarrow{\uparrow} |1\rangle$$



BREAK TIME!

3:05 PM EST

$$|\Psi\rangle = \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \rightarrow |+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}} \leftarrow \text{SUPERPOSITION}$$



$$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \cancel{|0\rangle\langle 1| + |1\rangle\langle 0|}$$

$$\langle 0|0\rangle = 1 \rightarrow (|0\rangle)(\cancel{|0\rangle}) = 1(1) + 0(0) = 1$$

$$\langle 1|1\rangle = 1$$

$$\langle 0|1\rangle = 0 \rightarrow (|0\rangle)(\cancel{|1\rangle}) = 1(0) + 0(1) = 0$$

$$\langle 1|0\rangle = 0$$

$\cancel{|0\rangle}$

$$\begin{aligned} X|0\rangle &= (|0\rangle\langle 1| + |1\rangle\langle 0|)|0\rangle \\ &= \cancel{|0\rangle\langle 1|} + |1\rangle\langle 0| \\ &= \cancel{|0\rangle\langle 1|} + |1\rangle\cancel{\langle 0|} \\ &= \cancel{|0\rangle} + |1\rangle \\ &= |1\rangle \end{aligned}$$

$|0\rangle\langle 0| = \begin{pmatrix} 1 \\ 0 \end{pmatrix}(\cancel{|0\rangle}) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$



$$|1\rangle\langle 1| = \begin{pmatrix} 0 \\ 1 \end{pmatrix}(\cancel{|1\rangle}) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$|0\rangle\langle 1| = \cancel{\begin{pmatrix} 1 \\ 0 \end{pmatrix}}(\cancel{|1\rangle}) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$


$$|1\rangle\langle 0| = \cancel{\begin{pmatrix} 0 \\ 1 \end{pmatrix}}(\cancel{|0\rangle}) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

$$\begin{aligned}
 X \Psi &= (|11\rangle\langle 01| + |10\rangle\langle 11|) (\alpha|10\rangle + \beta|11\rangle) \\
 &= \alpha|11\rangle\cancel{\langle 01|0\rangle} + \cancel{\beta|11\rangle\langle 01|1\rangle} + \cancel{\alpha|10\rangle\langle 11|0\rangle} + \beta|10\rangle\cancel{\langle 11|1\rangle}
 \end{aligned}$$


$$\begin{aligned}
 &= \alpha|11\rangle + \beta|10\rangle \\
 \Psi &= \alpha|10\rangle + \beta|11\rangle
 \end{aligned}$$



$$\begin{aligned}
 H|10\rangle &= |+\rangle \\
 H|+\rangle &= |0\rangle \\
 H|-> &= |1\rangle
 \end{aligned}$$




$$\begin{aligned}
 |\Psi\rangle &-> H -> \alpha \\
 |+\rangle &\rightarrow |0\rangle \\
 |-> &\rightarrow |1\rangle
 \end{aligned}$$



MULTI-QUBIT CIRCUIT MATH

MANY SINGLE-QUBIT CIRCUITS


What if we have a collection of n single-qubit circuits?



There are 2 different ways we can approach solving for the output n - bit string.


MANY SINGLE-QUBIT CIRCUITS

Method #1: Individually solve each of the n single-qubit circuits.



MANY SINGLE-QUBIT CIRCUITS

Method #2: Simultaneously solve all n single-qubit circuits.



$$T_n \dots T_2 T_1 |\Psi\rangle = |\Psi'\rangle$$

$(2^n \times 2^n) \dots (2^n \times 2^n)$

$(2^n \times 2^n) \quad (2^n \times 1)$

$(2^n \times 1)$

$|\Psi\rangle - T_1 - T_2 - \dots - T_n - \alpha$


A handwritten note on the right side of the diagram shows the expression $T_n \dots T_2 T_1 |\Psi\rangle = |\Psi'\rangle$. Above this, a purple bracket groups the terms $(2^n \times 2^n) \dots (2^n \times 2^n)$. To the right, a blue bracket groups the terms $(2^n \times 1)$ and $(2^n \times 1)$. A blue arrow points from the bracketed term $(2^n \times 1)$ towards the final state $|\Psi'\rangle$. Below the diagram, the state $|\Psi\rangle$ is followed by a sequence of boxes labeled $T_1, T_2, \dots, T_n, \alpha$, with a minus sign between each box, indicating the decomposition of the total unitary into individual components.

MULTI-QUBIT CIRCUITS

Are all multi-qubit circuits just collections of single-qubit circuits?

To answer this, let's think back to our 3 key quantum properties:

SUPERPOSITION




H!

QUANTUM INTERFERENCE



$$H|0\rangle = |+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}$$
$$H|+\rangle = |0\rangle$$

ENTANGLEMENT






MULTI-QUBIT GATES

To achieve entanglement, our qubits need to interact,
meaning we need 2-qubit (or larger) gates!

Construct Bell state

```
In [17]: 1 bell = QuantumCircuit(2, 2)
2 bell.h(1)
3 bell.cx(1, 0)
4 bell.measure(range(2), range(2))
5 bell.draw()
```


Out[17]:



Note that these are $(2^2 \times 2^2) = (4 \times 4)$ matrices.



MULTI-QUBIT GATES

However, with 2-qubit gates, we can no longer use **Method #1** (individually calculating each single-qubit circuit) for calculating the output of the multi-qubit circuit...




So now let's learn **Method #2!**

THE MULT-QUBIT CIRCUIT MODEL



TENSOR PRODUCT

We use the *tensor product* to solve multi-qubit circuits!



MULTI-QUBIT STATES & GATES

Mathematically, a quantum bit string is simply a ***tensor product*** of qubits!


$$|\psi_1\rangle \text{---} \\ |\psi_2\rangle \text{---} \\ \vdots \\ |\psi_n\rangle \text{---}$$
$$= |\Psi\rangle \text{---} \\ |\psi_1\rangle \otimes |\psi_2\rangle \otimes \dots \otimes |\psi_n\rangle$$

Mathematically, gates acting on different qubits are a ***tensor product*** of all the gates!

$$|\psi_1\rangle \xrightarrow{A_1} \text{---} \\ |\psi_2\rangle \xrightarrow{A_2} \text{---} \\ \vdots \\ |\psi_n\rangle \xrightarrow{A_n} \text{---}$$
$$= |\Psi\rangle \xrightarrow{A} \text{---}$$
$$A = A_1 \otimes A_2 \otimes \dots \otimes A_n$$

TENSOR

A tensor is essentially an n-dimensional vector... In quantum computing, we care about vectors (states) and matrices (gates), which are 1-D and 2-D tensors!



Dude, that sounds
in-tense...



imgflip.com

THE TENSOR (Kronecker) PRODUCT

If \mathbf{A} is an $(n \times m)$ matrix and \mathbf{B} is a $(p \times q)$ matrix, their tensor product is the $(pn \times mq)$ matrix:

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_{11}\mathbf{B} & a_{12}\mathbf{B} & \cdots & a_{1m}\mathbf{B} \\ a_{21}\mathbf{B} & a_{22}\mathbf{B} & \cdots & a_{2m}\mathbf{B} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1}\mathbf{B} & a_{n2}\mathbf{B} & \cdots & a_{nm}\mathbf{B} \end{pmatrix}$$

$$\mathbf{A} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$$

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_1 \otimes \mathbf{B} \\ a_2 \otimes \mathbf{B} \end{pmatrix} = \begin{pmatrix} a_1 \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \\ a_2 \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \end{pmatrix} = \begin{pmatrix} a_1 b_{11} & a_1 b_{12} \\ a_1 b_{21} & a_1 b_{22} \\ a_2 b_{11} & a_2 b_{12} \\ a_2 b_{21} & a_2 b_{22} \end{pmatrix}$$

QUANTUM PRACTICE TIME!

Write out the full matrix for the following tensor products...

$$(1) \quad \begin{pmatrix} 1 \\ 2 \end{pmatrix} \otimes \begin{pmatrix} 3 \\ 4 \end{pmatrix} =$$

$$(3) \quad \begin{pmatrix} 1 \\ -1 \end{pmatrix} \otimes \begin{pmatrix} 2 & 3 \\ 4 & 5 \end{pmatrix} =$$

$$(2) \quad \begin{pmatrix} 3 \\ 4 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 2 \end{pmatrix} =$$

$$(4) \quad \begin{pmatrix} 2 & 3 \\ 4 & 5 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ -1 \end{pmatrix} =$$

QUANTUM PRACTICE SOLUTIONS!

Write out the full matrix for the following tensor products...

$$(1) \quad \begin{pmatrix} 1 \\ 2 \end{pmatrix} \otimes \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 1 & \begin{pmatrix} 3 \\ 4 \end{pmatrix} \\ 2 & \begin{pmatrix} 3 \\ 4 \end{pmatrix} \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \\ 6 \\ 8 \end{pmatrix}$$

↑
order matters!

$$(2) \quad \begin{pmatrix} 3 \\ 4 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 & \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ 4 & \begin{pmatrix} 1 \\ 2 \end{pmatrix} \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \\ 4 \\ 8 \end{pmatrix}$$

$| \psi_1 \rangle - \boxed{A} -$

$| \psi_2 \rangle - \boxed{B} - \neq$

$| \psi_1 \rangle - \boxed{B} -$
 $| \psi_2 \rangle - \boxed{A} -$

$$(3) \quad \begin{pmatrix} 1 \\ -1 \end{pmatrix} \otimes \begin{pmatrix} 2 & 3 \\ 4 & 5 \end{pmatrix} = \begin{pmatrix} 1 & \begin{pmatrix} 2 & 3 \\ 4 & 5 \end{pmatrix} \\ -1 & \begin{pmatrix} 2 & 3 \\ 4 & 5 \end{pmatrix} \end{pmatrix}$$


\neq

$$= \begin{pmatrix} 2 & 3 \\ 4 & 5 \\ -2 & -3 \\ -4 & -5 \end{pmatrix}$$

$$(4) \quad \begin{pmatrix} 2 & 3 \\ 4 & 5 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 2 & \begin{pmatrix} 1 \\ -1 \end{pmatrix} & 3 & \begin{pmatrix} 1 \\ -1 \end{pmatrix} \\ 4 & \begin{pmatrix} 1 \\ -1 \end{pmatrix} & 5 & \begin{pmatrix} 1 \\ -1 \end{pmatrix} \end{pmatrix}$$
$$= \begin{pmatrix} 2 & -2 & 3 & -3 \\ 4 & -4 & 5 & -5 \end{pmatrix}$$

MULTI-QUBIT STATES

Mathematically, a quantum bit string is simply a tensor product of qubits!



$$\begin{aligned} |01\rangle &= |0\rangle \otimes |1\rangle \\ |11\rangle &= |1\rangle \otimes |1\rangle \quad \left(\begin{matrix} 1 \\ 0 \end{matrix}\right) \quad \left(\begin{matrix} 1 \\ 1 \end{matrix}\right) \quad \left(\begin{matrix} 0 \\ 1 \end{matrix}\right) \end{aligned}$$

$$|\psi_1 \psi_2 \dots \psi_n\rangle = |\psi_1\rangle \otimes |\psi_2\rangle \otimes \dots \otimes |\psi_n\rangle$$

n qubits are represented mathematically by a vector of length 2^n

$(2^n \times 1)$

```
bell = QuantumCircuit(2, 2)
```

WORKED EXAMPLES

(1)


$$\begin{aligned}
 |010\rangle &= |0\rangle \otimes |1\rangle \otimes |0\rangle \\
 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \left(\begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) \\
 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \\
 &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} = 2^3
 \end{aligned}$$

(2)

$$\begin{aligned}
 |\pm\pm\rangle &= \frac{1}{\sqrt{2}}(|+)\otimes\frac{1}{\sqrt{2}}(|+|-|-) \\
 &= \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \\
 &= \frac{1}{2} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}
 \end{aligned}$$

MULTI-QUBIT GATES

If we want to calculate how a set of quantum gates will affect our multi-qubit state, we will need to take the tensor product of the gate matrices.




$$\underbrace{A_1 \otimes A_2 \otimes \dots \otimes A_n}_{\text{n single qubit gates on different wires}} \underbrace{|\psi_1 \ \psi_2 \dots \ \psi_n\rangle}_{\text{mathematically by a matrix of dimension $(2^n \times 2^n)$}} \\ = \underbrace{A_1}_{|\phi_1\rangle} |\psi_1\rangle \otimes \underbrace{A_2}_{|\phi_2\rangle} |\psi_2\rangle \otimes \dots \otimes \underbrace{A_n}_{|\phi_n\rangle} |\psi_n\rangle$$

n single qubit gates on different wires are represented mathematically by a matrix of dimension $(2^n \times 2^n)$

```
bell = QuantumCircuit(2, 2)
bell.i(0)
bell.h(1)
```

Note that wherever there is not a gate, we must insert an identity matrix.

WORKED EXAMPLE



$|\Phi\rangle =$


$$\begin{aligned}
 |\Phi\rangle &= (|0\rangle \otimes |0\rangle \otimes |0\rangle) = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \\
 &= (|0\rangle \otimes I \otimes H) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} \end{pmatrix} \otimes \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \\
 &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \otimes \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\
 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \\
 &\quad \text{The probability amplitudes are } \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}. \\
 &\quad \text{The probability of measuring each state is } \left(\frac{1}{\sqrt{2}}\right)^2 = \frac{1}{2} \approx 50\%.
 \end{aligned}$$

MULTI-QUBIT MEASUREMENT

In single-qubit measurement, we were simply measuring 1 quantum state.

```
qc_z.measure(0, 0)
```


This state is in a superposition (linear combination) of $|0\rangle$ and $|1\rangle$:



In multi-qubit measurement, we are now measuring n states.

```
bell.measure(range(2), range(2))
```


These n states are in a superposition of all 2^n possible bit strings! :



$$|\phi\rangle = \begin{pmatrix} \alpha \\ \beta \\ \gamma \\ \vdots \\ \tau \end{pmatrix} \rightarrow |\alpha|^2 P(\text{meas } 0) \\ |\beta|^2 P(\text{meas } 1) \\ |\gamma|^2 P(\text{meas } 2) \\ \vdots \\ |\tau|^2 P(\text{meas } n-1) \end{pmatrix}$$

MULTI-QUBIT MEASUREMENT


Example: 2-qubit circuit measurement




UNIFORM SUPERPOSITION?

Want to have equa) likelihood of measuring every possible quantum bitstring.

ex) 2-qubit circuit : $P(|00\rangle) = P(|01\rangle) = P(|10\rangle) = P(|11\rangle)$



WORKED EXAMPLE



$$|\phi\rangle = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$$




P(|00>) = |\frac{1}{2}|^2 = \frac{1}{4} 25%
 P(|01>) = |\frac{1}{2}|^2 = \frac{1}{4} 25%
 P(|10>) = |\frac{1}{2}|^2 = \frac{1}{4} 25%
 P(|11>) = |\frac{1}{2}|^2 = \frac{1}{4} 25%

$$|0> \otimes |0> = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1(1) \\ 0(1) \\ 0(0) \\ 0(0) \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$H \otimes H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

$$= \frac{1}{2} \begin{pmatrix} 1(1,1) & 1(1,-1) \\ 1(-1,1) & 1(-1,-1) \end{pmatrix}$$

WORKED EXAMPLE




ENTANGLED STATES?


Bell's states (Bell's experiment)

$$\frac{|00\rangle + |11\rangle}{\sqrt{2}}, \quad \frac{|01\rangle + |10\rangle}{\sqrt{2}}, \quad \frac{|01\rangle - |10\rangle}{\sqrt{2}}, \quad \frac{|00\rangle - |11\rangle}{\sqrt{2}}$$

ANTI



WORKED EXAMPLE




$$|0\rangle \otimes |0\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{aligned} T_1: H \otimes I &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \end{pmatrix} \\ T_2: CNOT &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \end{aligned}$$

$$|\psi\rangle = T_2 T_1 |\psi\rangle = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \end{pmatrix} \right) \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \frac{\sqrt{2}}{2} \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{2}}{2} \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

WORKED EXAMPLE




$$\left(\begin{array}{l} \frac{1}{2} \\ 0 \\ 0 \\ \frac{1}{2} \end{array} \right) \rightarrow \begin{array}{l} P(|00\rangle) = |\frac{1}{2}|^2 = \frac{1}{2} \quad 50\% \\ P(|01\rangle) = |0|^2 = 0 \\ P(|10\rangle) = |0|^2 = 0 \\ P(|11\rangle) = |\frac{1}{2}|^2 = \frac{1}{2} \quad 50\% \end{array}$$

$|00\rangle$


$|11\rangle$

SOME MULTI-QUBIT CIRCUITS TO COME...


DEUTSCH-JOZSA




QUANTUM TELEPORTATION




GROVER'S



QAOA





© 2020 The Coding School
All rights reserved

Use of this recording is for personal use only. Copying, reproducing, distributing, posting or sharing this recording in any manner with any third party are prohibited under the terms of this registration. All rights not specifically licensed under the registration are reserved.