

the **nature** EDUCATION

## Knowledge Project

Powered by  
LEARN MORE

ADVANCED

EVOLUTION | Lead Editor: Nick Bisceglia



## Geographic Mosaics of Coevolution

By: Jason D. Hoeksema © 2012 Nature Education

Citation: Hoeksema, J. D. (2012) Geographic Mosaics of Coevolution. *Nature Education Knowledge* 3(10):19

Species constantly engage in strong interactions with other species - parasites, predators, prey, and mutualists. As a result, their traits may coevolve and diversify in geographic mosaics.

Aa Aa Aa



When G. E. Hutchinson argued that evolution is a play taking place on an ecological stage (1965), his point was that ecological interactions — between species and the living (biotic) and non-living (abiotic) aspects of their environment — are the drivers of adaptive evolution. His illustrative examples demonstrate the ubiquity of **interspecific interactions** that dominate the ecology species: he showed that the foraging niches of organisms as diverse as rhinoceros, birds, cone shells, copepods, and rotifers seem to have diversified in response to competition among similar species within those groups. Beyond competition, most species also interact with other species as a consumer or parasite, a food resource or prey item, or both. Finally, most species are engaged in several kinds of mutualisms with other species, in which both species typically benefit; for example, plants may engage simultaneously in mutually beneficial interactions with fruit-dispersing birds, pollinating insects, nitrogen-fixing bacteria, and mycorrhizal fungi that act like extended root systems in the soil. One potentially dramatic consequence of these interactions can be **coevolution**, in which the traits of each of two species evolve in response to the traits of the other, resulting in relentlessly reciprocal evolutionary changes in the traits of those interacting species.

Coevolution has been studied for almost as long as the theory of evolution by natural selection has been around. Throughout their writings, Darwin and Wallace both recognized the importance of the coevolutionary process in evolutionary diversification, emphasizing the

primacy of species interactions as drivers of adaptive evolutionary change in species. Indeed, it has been argued that Darwin's book on orchids provided the catalyst for all subsequent studies of coevolution and the evolution of extreme specialization (Ghiselin 1984, Thompson 1994). Since Darwin and Wallace, studies on coevolution have shown that species interactions can drive rapid and sustained evolutionary change in species at multiple spatial and temporal scales, generating genetic diversity within populations, leading to adaptive differentiation among populations, and often leading to **ecological speciation** (Schluter 2009). It has been argued that much of the diversity on earth is a consequence of coevolutionary diversification in species interactions (Thompson 1994, 2005, Ehrlich & Raven 1964). Clearly, studies of coevolution in species interactions can lend insight into the fundamental processes generating and maintaining biodiversity, including genetic and phenotypic diversity within and between species.

## What is Coevolution?

Coevolutionary selection between two species within a single local community (i.e., local coevolutionary selection) has the potential to act as a potent evolutionary force on the traits of species by virtue of the feedback driven by its inherent reciprocity. The reciprocity of coevolutionary selection means that the fitnesses of two interacting species depend not only on their own genotypes (and associated traits), but also on each other's genotypes (and traits). This means that there is a genotype by genotype ( $G \times G$ ) interaction on the fitnesses of both species. As a result of this reciprocity, adaptive changes in the traits of one species may trigger subsequent adaptive changes in the traits of the second species, which in turn feed back to cause further adaptive evolution of the first species, and so on. Put another way, the relative fitness of genotypes in one species is context-dependent and the context itself can also evolve (Wade 2003).

For example, the relative fitnesses of different genotypes of a hummingbird and a hummingbird-pollinated plant may depend not only on hummingbird bill shape (which differs among hummingbird genotypes), or on flower morphology (which differs among plant genotypes), but also on the match between the two, generating a  $G \times G$  interaction for both hummingbird fitness and plant fitness. If hummingbird bill shape adapts in response to selection driven by the average flower morphology in the plant population, the nature of the reciprocal selection on flower morphology (driven by the newly evolved average hummingbird bill shape) changes as a result.

## The Crucial Spatial Perspective: Coevolution Occurs in Geographic Mosaics

Although local coevolutionary selection is an important building block of coevolutionary dynamics, studies conducted in the last few decades on coevolution in a wide range of species interactions have demonstrated that coevolution is inherently geographically structured. Specifically, coevolution between two species can progress along very different trajectories in different places, causing trait differences to evolve among different populations of the same species. Because most species are collections of genetically variable populations distributed among environments, and because those environments often differ in abiotic factors and/or **community composition**, the pattern and strength of natural selection imposed by species on each others' traits are also expected to be highly variable among those environments. This recognition that coevolution is an inherently geographically-structured process has been synthesized by John Thompson as the **Geographic Mosaic Theory of Coevolution** (GMTC, Thompson 2005).

Specifically, the GMTC proposes that coevolution exhibits three inherently geographic characteristics, beyond local coevolutionary selection: coevolutionary hotspots and coldspots, geographic selection mosaics, and trait remixing (Figure 1a). Coevolutionary hotspots are communities in which the pairwise interaction between two species exhibits coevolutionary selection (i.e., there is a  $G \times G$  interaction on the fitnesses of both species) and these hotspots are expected to be embedded in a geographic matrix of coldspots. In coldspots, selection is not reciprocal, including communities in which only one of the species occurs (i.e., structural coldspots). Geographic selection mosaics occur when a fitness function of coevolutionary selection (dependence of fitness on  $G \times G$ ) differs among environments ( $E$ ) (i.e., there is a genotype by genotype by environment interaction [ $G \times G \times E$ ] for fitness of at least one of the two species). Trait remixing is the suite of processes that potentially influence the geographic distributions of **alleles at gene loci** underlying coevolving traits (i.e., the geographic structure of  $G$  for both species), including **mutations**, **gene flow**, **genetic drift**, and population extinction/recolonization dynamics.


## Testing the Importance of Geographic Mosaics of Coevolution

Mathematical modeling studies of geographic mosaics of coevolution have shown how the three key elements of the GMTC can interact to generate a wide variety of outcomes of coevolution, including patterns of **specificity**, **local adaptation/maladaptation**, and patterns of genetic variation within and between populations (e.g., Nuismer *et al.* 2000, Gomulkiewicz *et al.* 2000, Nuismer 2006). Moreover, empirical studies across a wide range of species interactions — conifers and crossbills (Benkman *et al.* 2001), snails and trematodes (Lively *et al.* 2004), camellias and weevils (Toju & Sota 2006), garter snakes and newts (Brodie *et al.* 2002), and *Greya* moths that act as both pollinators and seed parasites of *Lithophragma* plants (Thompson & Cunningham 2002), to name just a few — have provided support for key elements of the theory (reviewed by Thompson 2005). For example, Johnson *et al.* (2010) studied interactions between big bluestem

grass (*Andropogon gerardii*) and its arbuscular mycorrhizal (AM) fungal symbionts from three different populations. They estimated plant and fungal performance in all reciprocal combinations of plant populations ( $G_P$ ), whole AM fungal guilds of species ( $G_F$ ), and sterile soils (E) in a greenhouse experiment. They found a significant  $G_P \times G_F \times E$  interaction for a key trait of the interaction: the formation of **arbuscules**, which are the fungal structures in AM plant roots through which nutrients pass. In fact, arbuscule formation was highest in local combinations of plants, fungi, and soils, suggesting that a selection mosaic (different  $G_P \times G_F$  interactions for different E) has resulted in local adaptation of plants and fungi to each other in that interaction.


However, it remains challenging to perform such rigorous tests of key phenomena central to the GMTCC, especially coevolutionary selection mosaics (Gomulkiewicz *et al.* 2007), since ideal tests require that all combinations of genotypes of all important interacting species are replicated and studied across multiple field sites or artificial gradients of key environmental variables (Nuismer & Gandon 2008). As a result, many of the necessary experiments have not yet been performed to test the importance of ongoing coevolution and geographic mosaics of coevolution in dominant species interactions, such as those between plants and mycorrhizal fungi (Hoeksema 2010). Furthermore, non-reciprocal adaptation of species to both biotic and abiotic environmental factors has been repeatedly demonstrated to be an important force driving diversification in species (e.g., Bohrer *et al.* 2003, Saenz-Romero *et al.* 2006). Thus, a key question that ultimately needs to be addressed in a comprehensive fashion is, what are the relative roles of non-reciprocal selection from biotic and abiotic factors versus the three elements of the GMTCC (selection mosaics, hotspots and coldspots, and trait remixing) in driving trait diversification of species?

Broadly speaking, there are at least three hypothetical answers to this question for a particular species interaction (Figure 1, Hoeksema 2010). One hypothesis is that all three of the GMTCC elements — hotspots/coldspots, selection mosaics, and trait remixing—are important and interact to play prominent roles in driving diversification within and across communities in which two species occur (Figure 1a). A second hypothesis is that coevolutionary selection is relatively important but is consistent among populations (i.e., true geographic selection mosaics [ $G \times G \times E$  interactions] and hotspots/coldspots are relatively unimportant or uncommon [Figure 1b]). A third hypothesis is that local coevolutionary selection is consistently less important compared to non-reciprocal selection on species by abiotic environmental factors, or by one partner species on the other (Figure 1c). It will be exciting to see whether new individual studies, or perhaps **meta-analyses** of sets of existing studies, can be designed that distinguish among these possibilities, ultimately helping us to better understand what kinds of drama tend to dominate the evolutionary plays taking place on Hutchinson's ecological stage.



**Figure 1: Alternative hypotheses on the relative importance of the three elements of the geographic mosaic theory of coevolution (GMTC) and non-reciprocal selection.**

Each circle represents a separate local community. Arrows inside circles represent selection among plants (P), mycorrhizal fungi (F), and the abiotic environment (E). Arrows outside circles represent gene flow among communities. a) Hypothesis 1: All three of the GMTC elements — hotspots/coldspots, selection mosaics, and trait remixing—are important, and interact to play prominent roles in driving diversification within and across communities in which two species occur. Reciprocal selection is shown in some communities (hotspots) and not in others (coldspots). The dashed arrow indicates that plant selection on fungi differs, compared to the first community, implying a selection mosaic. The main effect of the abiotic environment is to modify interspecific selection exerted by plants on mycorrhizal fungi. b) Hypothesis 2: Coevolutionary selection is important but is consistent among populations (i.e., true geographic selection mosaics [G x G x E interactions] and hotspots/coldspots are relatively unimportant or uncommon). Coevolution can still be geographically variable, if the distribution of genotypes differs among communities due to trait remixing processes. c) Hypothesis 3: Ongoing local coevolutionary selection is consistently less important compared to non-reciprocal selection on species by abiotic environmental factors, or by one partner species on the other.

© 2012 **Nature Education** Modified from Hoeksema 2010. All rights reserved. 

## References and Recommended Reading

Benkman, C. W. *et al.* The influence of a competitor on the geographic mosaic of coevolution between crossbills and lodgepole pine. *Evolution* **55**, 282–294 (2001).

Bohrer, G. *et al.* Effects of different Kalahari-desert VA mycorrhizal communities on mineral acquisition and depletion from the soil by host plants. *Journal of Arid Environments* **55**, 193–208 (2003).

Brodie, E. D. Jr. *et al.* The evolutionary response of predators to dangerous prey: Hotspots and coldspots in the geographic mosaic of coevolution between newts and snakes. *Evolution* **56**, 2067–2082 (2002).

Ehrlich, P. R. & Raven, P. H. Butterflies and plants: A study in coevolution. *Evolution* **18**, 586–608 (1964).

Ghiselin, M. T. "Pages xi–xix" in *The Various Contrivances by Which Orchids are Fertilized by Insects*, by C. Darwin, Reprint of 2nd Edition, 1877. Chicago, IL: University of Chicago Press, 1984.

Gomulkiewicz, R. *et al.* Hot spots, cold spots, and the geographic mosaic theory of coevolution. *The American Naturalist* **156**, 156–174 (2000).

——— Dos and don'ts of testing the geographic mosaic theory of coevolution. *Heredity* **98**, 249–258 (2007).

Hoeksema, J. D. Tansley Review. Ongoing coevolution in mycorrhizal interactions. *New Phytologist* **187**, 286–300 (2010).

Johnson, N. C. *et al.* Resource limitation is a driver of local adaptation in mycorrhizal symbioses. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 2093–2098 (2010).

Lively, C. M. *et al.* Host sex and local adaptation by parasites in a snail-trematode interaction. *The American Naturalist* **164**, S6–S18 (2004).

Nuismer, S. L. Parasite local adaptation in a geographic mosaic. *Evolution* **60**, 24–30 (2006).

Nuismer, S. L. & Gandon, S. Moving beyond common-garden and transplant designs: Insight into the causes of local adaptation in species interactions. *The American Naturalist* **171**, 658–668 (2008).

Nuismer, S. L. *et al.* Coevolutionary clines across selection mosaics. *Evolution* **54**, 1102–1115 (2000).

Saenz-Romero, C. *et al.* Altitudinal genetic variation among *Pinus oocarpa* populations in Michoacan, Mexico: Implications for seed zoning, conservation, tree breeding and global warming. *Forest Ecology and Management* **229**, 340–350 (2006).

Schluter, D. Evidence for ecological speciation and its alternative. *Science* **323**, 737–741 (2009).

Thompson, J. N. *The Coevolutionary Process*. Chicago, IL: University of Chicago Press, 1994.

Thompson, J. N. *The Geographic Mosaic of Coevolution*. Chicago, IL: University of Chicago Press, 2005.

Thompson, J. N. & Cunningham, B. M. Geographic structure and dynamics of coevolutionary selection. *Nature* **417**, 735–738 (2002).

Toju, H. & Sota, T. Imbalance of predator and prey armament: Geographic clines in phenotypic interface and natural selection. *The American Naturalist* **167**, 105–117 (2006).

Wade, M. J. Community genetics and species interactions. *Ecology* **84**, 583–585 (2003).

[Outline](#) | [Keywords](#)

[FEEDBACK](#)



## Explore This Subject

### BASIC

Evolution Is Change in the Inherited Traits  
of a Population through Successive  
Generations

The Hardy-Weinberg Principle

Using Molecular Techniques to Answer  
Ecological Questions

Evolution Introduction

Game Theory, Evolutionary Stable  
Strategies and the Evolution of Biological  
Interactions

Life History Evolution

Mutations Are the Raw Materials of  
Evolution

Speciation: The Origin of New Species

## ADVANCED

---

Avian Egg Coloration and Visual Ecology

The Ecology of Avian Brood Parasitism

The Geography and Ecology of  
Diversification in Neotropical Freshwaters

The Maintenance of Species Diversity

Natural Selection, Genetic Drift, and Gene  
Flow Do Not Act in Isolation in Natural  
Populations

Neutral Theory of Species Diversity

Population Genomics

Semelparity and Iteroparity

Geographic Mosaics of Coevolution

Milankovitch Cycles, Paleoclimatic  
Change, and Hominin Evolution

Paleoecology and Paleoenvironment: a  
Case Study of Plio-Pleistocene Mammals  
from Laetoli

Testing Models of Modern Human Origins  
with Archaeology and Anatomy

## INTERMEDIATE

---

Why Are Life Histories So Variable?

Comparative Genomics

Case Study: The Glorious, Golden, and  
Gigantic Quaking Aspen

Cybertaxonomy and Ecology

Molecular Genetic Techniques and  
Markers for Ecological Research

Ecological Opportunity: Trigger of  
Adaptive Radiation

Evidence for Meat-Eating by Early  
Humans

Resource Partitioning and Why It Matters

The Evolution of Aging

