

PH4209 Assignment — 03

22MS037

Problem 1 — Moran Process with constant selection but without mutation:

Take a population of $N=100$ individuals one of which consist of type A and the remaining of type B, initially. Assume that the fitness of type A is r and the fitness of type B is 1 and neither type can mutate to the other.

1. Write a program to obtain the time-evolution of the frequencies of the two types in the population when evolution occurs according to the Moran Process i.e. in every generation only one individual is picked at random for death and another individual is picked for reproduction with a probability proportional to its fitness. Use $r=1.01$ Run the simulation for as long as it takes for any one of the two types to get fixed in the population.
2. **Obtaining the Fixation Probability:** Repeat the above simulation for $N_t=1000$ trials and find out the fraction of times each of the two types A and B get fixed ? Check whether the invasion probability you obtain from the simulation matches with the theoretical value of invasion probability.

The fitness of type A is $r=1.01$ and the fitness of type B is 1. We simulate the process until one of the types gets fixed in the population. Let x_i be the fixation probability of type A starting from the initial state $x = \frac{i}{N}$. The theoretical invasion probability for type A can be calculated using the formula:

$$x_i = \frac{1 - r^{-i}}{1 - r^{-N}} \quad (1)$$

For our case, with $i = 1$ and $N = 100$, the theoretical invasion probability for type A is:

$$x_1 = \frac{1 - 1.01^{-1}}{1 - 1.01^{-100}} \approx 0.016 \quad (2)$$




Figure 1: Invasion of type A

We simulated the Moran process for $N_t = 1000$ trials. The fraction of times type A got fixed in the population was 0.011 and the fraction of times type B got fixed was 0.989.




Figure 2: Fixation of type A with 1000 trials

The convergence can be observed as the number of trials increases, and the simulated fixation probability approaches the theoretical value of approximately 0.016 for type A. Here is the simulated probability of fixation for type A for $N_t = 5000$ trials.




Figure 3: Fixation of type A with 5000 trials. The simulated fixation probability is approximately 0.016.

Problem 2 — Moran Process with selection

If **half** the initial population consists of type A, write a program to obtain the time-evolution of the frequencies of the two types in the population when individuals making up the population in the next generation are chosen from members of the current generation with a probability proportional to their fitness. Use $r=0.99$. Run the simulation for as long as it takes for any one of the two types to get fixed in the population.

1. **Obtaining the Fixation Probability:** Repeat the above simulation for $N_t=1000$ trials and find out the fraction of times each of the two types A and B get fixed ?
2. For any one trial, plot the evolution of frequency of type A and type with time.

We simulate the Moran process with selection for $N = 100$ individuals, where initially 50 individuals are of type A and 50 individuals are of type B. The fitness of type A is $r = 0.99$ and the fitness of type B is 1. We simulate the process until one of the types gets fixed in the population. The theoretical invasion probability for type A can be calculated using the formula:

$$x_i = \frac{1 - r^{-i}}{1 - r^{-N}} \quad (3)$$

For our case, with $i = 50$ and $N = 100$, the theoretical invasion probability for type A is:

$$x_{\{50\}} = \frac{1 - 0.99^{-50}}{1 - 0.99^{-100}} \approx 0.377 \quad (4)$$

Invasion Probabilities from Moran Process




Figure 4: Fixation of type A and type B in the Moran process with selection for $r=0.99$

One can actually observe the convergence of the simulated fixation probabilities to the theoretical values as the number of trials increases. For $N_t = 5000$ trials, the simulated fixation probabilities are,




Figure 5: Fixation of type A and type B in the Moran process with selection for 5000 trials. The simulated fixation probabilities are approximately 0.37 for type A and 0.63 for type B.

Below is the plot of the evolution of frequencies of type A and type B over time for one trial.




Figure 6: Trajectory of frequencies of type A and type B over time for one trial.