
1	measufficie à mathamaticar intation
	Page No:
	Logistic Regression
2.10	11 rab 10 2500 r 1 2 250000 900 1 19 19 19 19 19 19
*	To some classification reproblem!
	A to the first of
	NO OF PRIVING TRAIN 2000 100 (0
	A ITHOUGH TOO PAIN
119751	120 Hour 100 7 (10) 7 (10) 6 (10) 4 (10) 5 (10) 4
	ACCUMENT TO LONG TO 1229 1898
	77290 117223119 98
*	Logistic Regression can be use for Binary
50/2.19ix	Problem solving
	Problem; solving in home in grinvose; maldorf
	Possifair
	1. O.5 Thresold value
(ross)	111 xxx priderouse to suppedicted point >0.5
	L) Poss
	0.5 - P.P. 4 0.5
	L) Fuit
05.37.40=63	Distriction of the Property of the simple of the state of
(Fe	ii) 0 1 2 3 4 5 6 7 hours madel
	model model
M [*]	- The same of the
-1	
	what is The Problem to solve Binary Problem
	using lineur Regression model
(4)	9.01111110 C
	The second secon
	J XXXX
1 PICIT, FIQUES	01 1 2 3 4 6 5 7 12
	0.5 -
1230 50	The state of the s
11	3 2 3 4 5 8 7 8 9 10 11 12
1	



(;(b)ad .;	use sigmoid Dipunction factor su arto 1 infloutcome.
della	"sigmoid fictivation function Responsible for soughing"
	It e = 00+come between 0 to I
	(D) DH) go! - (CO, O) T
	Hence $Z = ho(\alpha) + Go + O_{I}\alpha_{I}$
	0 2 P (41 10 ((20) mit -1) POI - (11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	A ST STORY OF HET IS A STORY OF THE STORY OF
	$ho(\alpha) = G(0o + O_1 \alpha_1).$
to the	((x) sd -1) gol) (x-E) - ((x) sd) gol y - = (20,00) t
. Id 11	ho(x) = 1
	(c)
	(c/ct 0 + 00) = 5 + 1 = 3 + 1
3 4	I+ c(ω0+02α2)
€0/2	They are the cost in of Foreson Regards
9.0	I+eOTO (CONVEX FUNCTION) + I
->	timeum Regnessian cost for of logistic Regnessian
THE	(α);)2 (α; - ho (α);)2
	$J(00.01) = I \int (4i - \hat{4})^2$
	$2n \qquad \qquad ho(\alpha) = I = I$
104	(,x,0) = + x (0) (00 (00) (00) (00) (00) (00) (00)
	tonvex / (convergence of a pare wow-convex
	Function Algo 3 1001
	1 / Y
	(1000) 6 /1 10 10 10 10 10 10 10 10 10 10 10 10 10
	gradient minima
	+ + + + + + + + + + + + + + + + + + + +
	310 bic minima 310 bic minima
	Nec 1

	Date:	
June 2 FUE	That's why we don't use I(0.02)	1 / (A! - pe(x)!) 5
prite up. E	or ania count caira mer coino ciere es	'iomgia"
1 3	we use "log los" cost function.	
/ '	2(00,02) - (-103(40(x)) if y=1	2:0311
· · · · · · · · · · · · · · · · · · ·	-10g (1-no(x)) if 4=0	
70 Y 20	J(00,03) = -4103 (ho(a)) - (1-4) (103 (1-h	
87 /s	ho(o))	
77	Here $ho(\infty) = \frac{1}{1+e^{-(00+0)}}$	(3)
*	- in the second of the local of	Regression
F 101213	Hega virginos - de pamo moizea egall	कारक क्षेत्र हैं-
	(Rewice) Costo Function,	0.00)
(. D. O + 10) * 9	J(09,03) by chamaing value	06(0) (0. 601
5 34/1	Recite (1/40) A STANT	
V	1 1 1	
	pretice . Francis &	
pariting brother	paginera 3 July 2 July	