



CHAPTER 1

Factoring Polynomials

We factor a polynomial into two or more polynomials of lower degree. For example, let's say that you wanted to factor $5x^3 - 45x$. You would note that you can factor out $5x$ from every term. Thus,

$$5x^3 - 45x = (5x)(x^2 - 9)$$

And then, you might notice that the second factor looks like the difference of squares, so

$$5x^3 - 45x = (5x)(x + 3)(x - 3)$$


That is as far as we can factorize this polynomial.

Why do we care? The factors make it easy to find the roots of the polynomial. This polynomial evaluates to zero if and only if at least one of the factors is zero. Here we see that

- The factor $(5x)$ is zero when x is zero.
- The factor $(x + 3)$ is zero when x is -3 .
- The factor $(x - 3)$ is zero when x is 3 .

So looking at the factorization, you can see that $5x^3 - 45x$ is zero when x is 0, -3, or 3.

This is a graph of that polynomial with its roots circled:



1.1 How to factor polynomials

The first step when you are trying to factor a polynomial is to find the greatest common divisor for all the terms, and pull that out. In this case, the greatest common divisor will also be a monomial: its degree is the least of the degrees of the terms, its coefficient will be the greatest common divisor of the coefficients of the terms.

For example, what can you pull out of this polynomial?

$$12x^{100} + 30x^3 + 42x^7$$

The greatest common divisor of the coefficients (12, 30, and 42) is 6. The least of the degrees of terms (100, 31, and 17) is 17. So you can pull out $6x^{17}$:

$$12x^{100} + 30x^3 + 42x^7 = (6x^{17})(2x^{83} + 5x^{14} + 7)$$

Exercise 1 Factoring out the GCD monomial*Working Space**Answer on Page 5*

So, now you have the product of a monomial and a polynomial. If you are lucky, the polynomial part looks familiar, like the difference of squares or a row from Pascal's triangle.

Often you are trying factor a quadratic like $x^2 + 5x + 6$ in a pair of binomials. In this case, the result would be $(x + 3)(x + 2)$. Let's check that:

$$(x + 3)(x + 2) = (x)(x) + (3)(x) + (2)(x) + (3)(2) = x^2 + 5x + 6$$

Notice that 3 and 2 multiply to 6 and add to 5. If I were trying to factor $x^2 + 5x + 6$, I would ask myself "What are two numbers that when multiplied equal 6 and when added equal 5?" And I would might guess wrong a couple of times. For example, I might say to myself "Well, 6 times 1 is 6. Maybe those work. But 6 and 1 add 7. So those don't work."

Solving these sorts of problems are like solving a Sudoku puzzle: you try things and realize they are wrong, so you backtrack and try something else.

The numbers are sometimes negative. For example, $x^2 + 3x - 10$ factors into $(x + 5)(x - 2)$.

Exercise 2 Factoring quadratics*Working Space**Answer on Page 5*

This is a draft chapter from the Kontinua Project. Please see our website (<https://kontinua.org/>) for more details.



APPENDIX A

Answers to Exercises

Answer to Exercise 1 (on page 3)

Answer to Exercise 2 (on page 3)



INDEX

factoring polynomials, 1