



PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

CS40032: Principles of Programming Languages

Module 03: λ -Calculus: Semantics

Partha Pratim Das

Department of Computer Science and Engineering
Indian Institute of Technology, Kharagpur

ppd@cse.iitkgp.ac.in

Jan 22, 27: 2020



PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

Semantics of λ -Expressions

Source:

λ - Calculus Overview

Operational Semantics of Pure Functional Languages



Free and Bound Variable

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- An occurrence of a variable x is said to be *bound* when it occurs in the body M of an abstraction $\lambda x.M$
- We say that λx is a *binder* whose scope is M
- An occurrence of x is *free* if it appears in a position where it is *not bound* by an enclosing abstraction on x
- For example,
 - Occurrences of x in xy and $\lambda y.xy$ are *free*
 - Occurrences of x in $\lambda x.x$ and $\lambda z.\lambda x.\lambda y.x(yz)$ are *bound*
 - In $(\lambda x.x)x$ the first occurrence of x is *bound* and the second is *free*
- In a loose parallel to C functions, consider the *bound* variables as *local* (including *parameters*) and *free* variables as *global* or *non-local*



Free and Bound Variable

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- In an abstraction, the variable named is referred to as the **bound** variable and the associated λ -expression is the **body** of the abstraction
- In an expression of the form:

$$\lambda v. e$$

occurrences of variable v in expression e are **bound**

- All occurrences of other variables are **free**
- Example:

$$((\lambda x. \lambda y. (xy))(yw))$$

- x , and y are **bound** in first part
- y , and w are **free** in second part



Free and Bound Variable: Other Contexts

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- $\int_0^1 x^2 dx; \int_0^1 A * x^2 dx$
- $\sum_{x=1}^{10} \frac{1}{x}; \sum_{x=1}^{10} K * \frac{1}{x}$
- $\lim_{x \rightarrow \infty} e^{-x}; \lim_{x \rightarrow \infty} (M + e^{-x})$
- `int succ(int x) { return x + 1; }`
- $\forall x \in \mathbb{R}, x > 1 \Rightarrow \frac{1}{x} < 1$



Free and Bound Variable

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* An occurrence of a variable v in a λ -expression is called **bound** if it is within the scope of a λv ; otherwise it is called **free**
 - A variable may occur both bound and free in the same λ -expression – for example, in $\lambda x. y \lambda y. y x$ the first occurrence of y is free and the other two are bound



Set of Free Variables

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* The **set of free variables** in an expression E , denoted by $FV(E)$, is defined as follows:
 - a. $FV(c) = \Phi$ for any constant c
 - b. $FV(x) = \{x\}$ for any variable x
 - c. $FV(E1\ E2) = FV(E1) \cup FV(E2)$
 - d. $FV(\lambda x. E) = FV(E) - \{x\}$
- A λ -expression E with no free variables ($FV(E) = \Phi$) is called **closed**



Substitution

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- The notation $E[v \rightarrow E_1]$ refers to the λ -expression obtained by replacing each free occurrence of the variable v in E by the λ -expression E_1
- Naive Rules of Substitution:
 - a. $v[v \rightarrow E_1] = E_1$ for any variable v
 - b. $x[v \rightarrow E_1] = x$ for any variable $x \neq v$
 - c. $(\lambda v. E)[v \rightarrow E_1] = \lambda v. (E[v \rightarrow E_1])$
 - d. $(E_{rator} E_{rand})[v \rightarrow E_1] = ((E_{rator}[v \rightarrow E_1])(E_{rand}[v \rightarrow E_1]))$
- Does it work?

$$(\lambda y. x)[x \rightarrow (\lambda z. zw)] = \lambda y. \lambda z. zw$$

YES!



Unsafe Substitution: Example

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Consider:

$$(\lambda x. x)[x \rightarrow y] = \lambda x. (x[x \rightarrow y]) = \lambda x. y$$

conflicts with a basic understanding that the names of bound variables (that is, parameters) do not matter.

- The identity function is the same whether we write it as $\lambda x.x$ or $\lambda z.z$ or $\lambda fred.fred$.
- If these do not behave the same way under substitution they would not behave the same way under evaluation and that seems wrong
- The mistake is that the substitution should only apply to free variables and not bound ones**
- Here x is bound in the term so we should not substitute it
- That seems to give us what we want:

$$(\lambda x.x)[x \rightarrow y] = \lambda x.x$$



Unsafe Substitution: Example

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Again, the naive substitution

$$(\lambda x. (mul\ y\ x))[y \rightarrow x] \Rightarrow (\lambda x. (mul\ x\ x))$$

is unsafe since the result represents a squaring operation whereas the original lambda expression does not

- A substitution is **valid** or **safe** if no free variable in $E1$ becomes bound as a result of the substitution $E[v \rightarrow E1]$
- An invalid substitution involves a **variable capture** or **name clash**
- Correct way would be:

$$(\lambda x. (mul\ y\ x))[y \rightarrow x] \Rightarrow (\lambda z. (mul\ y\ z))[y \rightarrow x]$$

$$(\lambda z. (mul\ y\ z))[y \rightarrow x] \Rightarrow (\lambda z. (mul\ x\ z))$$

- Unsafe substitutions change in semantics!



Substitution

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- **Definition:** The **substitution** of an expression for a (*free*) variable in a λ -expression is denoted by $E[v \rightarrow E_1]$ and is defined as follows:
 - a. $v[v \rightarrow E_1] = E_1$ for any variable v
 - b. $x[v \rightarrow E_1] = x$ for any variable $x \neq v$
 - c. $c[v \rightarrow E_1] = c$ for any constant c
 - d. $(E_{rator} E_{rand})[v \rightarrow E_1] = ((E_{rator}[v \rightarrow E_1])(E_{rand}[v \rightarrow E_1]))$
 - e. $(\lambda v. E)[v \rightarrow E_1] = (\lambda v. E)$ // v is not free in E
 - f. $(\lambda x. E)[v \rightarrow E_1] = \lambda x. (E[v \rightarrow E_1])$ when $x \neq v$ and $x \notin FV(E_1)$
 - g. $(\lambda x. E)[v \rightarrow E_1] = \lambda z. (E[x \rightarrow z][v \rightarrow E_1])$ when $x \neq v$ and $x \in FV(E_1)$, where $z \neq v$ and $z \notin FV(E E_1)$
- In part (g), the first substitution $E[x \rightarrow z]$ replaces the bound variable x that will capture the free x 's in E_1 by an entirely new bound variable z . Then the intended substitution can be performed safely.



Substitution Example

PoPL

Partha Pratim Das

Semantics

Free and Bound Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and Applicative Order

$$\begin{array}{ll}
(\lambda y. (\lambda f. f x) y) [x \rightarrow f y] & \Rightarrow_{\alpha} \\
\lambda z. ((\lambda f. f x) z) [x \rightarrow f y] & \Rightarrow \text{by g) since } y \in FV(f y) \\
\lambda z. ((\lambda f. f x) [x \rightarrow f y] z[x \rightarrow f y]) & \Rightarrow \text{by d)} \\
\lambda z. ((\lambda f. f x) [x \rightarrow f y] z) & \Rightarrow \text{by b)} \\
\lambda z. (\lambda g. (g x) [x \rightarrow f y]) z & \Rightarrow \text{by g) since } f \in FV(f y) \\
\lambda z. (\lambda g. g (f y)) z & \Rightarrow \text{by d), b), and a)}
\end{array}$$

Rules

- $v[v \rightarrow E_1] = E_1$ for any variable v
- $x[v \rightarrow E_1] = x$ for any variable $x \neq v$
- $c[v \rightarrow E_1] = c$ for any constant c
- $(E_{rator} E_{rand})[v \rightarrow E_1] = ((E_{rator}[v \rightarrow E_1])(E_{rand}[v \rightarrow E_1]))$
- $(\lambda v. E)[v \rightarrow E_1] = (\lambda v. E)$
- $(\lambda x. E)[v \rightarrow E_1] = \lambda x. (E[v \rightarrow E_1])$ when $x \neq v$ and $x \notin FV(E_1)$
- $(\lambda x. E)[v \rightarrow E_1] = \lambda z. (E[x \rightarrow z][v \rightarrow E_1])$ when $x \neq v$ and $x \in FV(E_1)$, where $z \neq v$ and $z \notin FV(E E_1)$



Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- A λ -expression has as its meaning the λ -expression that results after all its function applications (combinations) are carried out
- Evaluating a λ -expression is called **reduction**
- Four rules of reduction
 - α -Reduction: Renaming rule
 - β -Reduction: Substitution rule
 - η -Reduction: Function Equality rule
 - δ -Reduction: Pre-defined Constants' rule



α -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* If v and w are variables and E is a λ -expression,

$$\lambda v. E \Rightarrow_{\alpha} \lambda w. E[v \rightarrow w]$$

provided that w does not occur at all in E , which makes the substitution $E[v \rightarrow w]$ safe

- The equivalence of expressions under α -reduction is what makes part g) of the definition of substitution correct
- The α -reduction rule simply allows the changing of bound variables as long as there is no capture of a free variable occurrence
- The two sides of the rule can be thought of as variants of each other, both members of an equivalence class of *congruent* λ -expressions



α -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- The last example contains two α -reductions:

$$\lambda y. (\lambda f. f \ x) \ y \Rightarrow_{\alpha} \lambda y. ((\lambda f. f \ x) \ y)[y \rightarrow z] \Rightarrow_{\alpha} \lambda z. (\lambda f. f \ x) \ z$$

$$\lambda z. (\lambda f. f \ x) \ z \Rightarrow_{\alpha} \lambda z. ((\lambda f. f \ x) \ z)[f \rightarrow g] \Rightarrow_{\alpha} \lambda z. (\lambda g. g \ x) \ z$$

- Now that we have a justification of the substitution mechanism, the main simplification rule can be formally defined



β -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* If v is a variable and E and E_1 are λ -expressions,

$$(\lambda v. E) E_1 \Rightarrow_{\beta} E[v \rightarrow E_1]$$

provided that the substitution $E[v \rightarrow E_1]$ is carried out according to the rules for a safe substitution

- This β -reduction rule describes the function application rule in which the actual parameter or argument E_1 is *passed to* the function $(\lambda v. E)$ by substituting the argument for the formal parameter v in the function



β -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* If v is a variable and E and E_1 are λ -expressions,

$$(\lambda v. E) E_1 \Rightarrow_{\beta} E[v \rightarrow E_1]$$

provided that the substitution $E[v \rightarrow E_1]$ is carried out according to the rules for a safe substitution

- The left side $(\lambda v. E) E_1$ of a β -reduction is called a **β -redex** – derived from *reduction expression* – and meaning an expression that can be β -reduced
- β -reduction is the main rule of evaluation in the λ -calculus
- α -reduction makes the substitutions for variables valid



β -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and

Applicative Order

- The evaluation of a λ -expression consists of a series of β -reductions, possibly interspersed with α -reductions to change bound variables to avoid confusion
- Take $E \Rightarrow F$ to mean $E \Rightarrow_{\beta} F$ or $E \Rightarrow_{\alpha} F$ and let \Rightarrow^* be the reflexive and transitive closure of \Rightarrow
- Hence:
 - For any expression E , $E \Rightarrow^* E$ and
 - For any three expressions, $(E_1 \Rightarrow^* E_2$ and $E_2 \Rightarrow^* E_3)$ implies $E_1 \Rightarrow^* E_3$
- The goal of evaluation in the λ -calculus is to reduce a λ -expression via \Rightarrow until it contains no more β -redexes
- To define an *equality* relation on λ -expressions, we also allow a β -reduction rule to work backward



β -Abstraction

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* Reversing β -reduction produces the β -**abstraction** rule,

$$E[v \rightarrow E_1] \Rightarrow_{\beta} (\lambda v. E) E_1$$

and the two rules taken together give β -**conversion**, denoted by \Leftrightarrow_{β}

- Therefore $E \Leftrightarrow_{\beta} F$ if $E \Rightarrow_{\beta} F$ or $F \Rightarrow_{\beta} E$
- Take $E \Leftrightarrow F$ to mean $E \Leftrightarrow_{\beta} F$, $E \Rightarrow_{\alpha} F$ or $F \Rightarrow_{\alpha} E$ and let \Leftrightarrow^* be the reflexive and transitive closure of \Leftrightarrow
- Two λ -expressions E and F are **equivalent** or **equal** if $E \Leftrightarrow^* F$



β -Abstraction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Reductions (both α and β) are allowed to sub-expressions in a λ -expression by three rules:
 - 1 $E_1 \Rightarrow E_2$ implies $E_1 E \Rightarrow E_2 E$
 - 2 $E_1 \Rightarrow E_2$ implies $E E_1 \Rightarrow E E_2$
 - 3 $E_1 \Rightarrow E_2$ implies $\lambda x. E_1 \Rightarrow \lambda x. E_2$



η -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:* If v is a variable and E is a λ -expression (denoting a function), and v has no free occurrence in E ,

$$\lambda v. (E \ v) \Rightarrow_{\eta} E$$

- Example:

$$\lambda x. (sqr \ x) \Rightarrow_{\eta} sqr$$

$$\lambda x. (add \ 5 \ x) \Rightarrow_{\eta} (add \ 5)$$

Note: $(add \ 5 \ x)$ abbreviates $(add \ 5)$

- Take $E \Leftrightarrow_{\eta} F$ to mean $E \Rightarrow_{\eta} F$ or $F \Rightarrow_{\eta} E$



η -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- The requirement that x should have no free occurrences in E is necessary to avoid an invalid reduction such as

$$\lambda x. (add\ x\ x) \Rightarrow (add\ x)$$

- This rule fails when E represents some constants; for example, if 5 is a predefined constant numeral, $\lambda x. (5\ x)$ and 5 are not equivalent or even related
- η -reduction, justifies an extensional view of functions; that is, two functions are equal if they produce the same values when given the same arguments

$$\forall x, f(x) = g(x) \Rightarrow f = g$$



η -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- **Extensionality Theorem:** If $F_1 x \Rightarrow^* E$ and $F_2 x \Rightarrow^* E$ where $x \notin FV(F_1 F_2)$, then $F_1 \Leftrightarrow^* F_2$ where \Leftrightarrow^* includes η -reductions.

Proof.

$$F_1 \Leftrightarrow_{\eta} \lambda x. (F_1 x) \Leftrightarrow_{\eta} \lambda x. E \Leftrightarrow_{\eta} \lambda x. (F_2 x) \Leftrightarrow_{\eta} F_2$$

- The rule is not strictly necessary for reducing λ -expressions and may cause problems in the presence of constants, but included for completeness



δ -Reduction

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and

Applicative Order

- *Definition:* If the λ -calculus has predefined constants (that is, if it is not pure), rules associated with those predefined values and functions are called *delta* rules:

- Example:

$$(add\ 3\ 5) \Rightarrow_{\delta} 8$$

and

$$(not\ true) \Rightarrow_{\delta} false$$

- Example:

$$\begin{aligned} twice &= \lambda f. \lambda x. f\ (f\ x) \\ twice\ (\lambda n. (add\ n\ 1))\ 5 &\Rightarrow_{\beta} \\ (\lambda f. \lambda x. (f\ (f\ x))) (\lambda n. (add\ n\ 1))\ 5 &\Rightarrow_{\beta} \\ (\lambda x. ((\lambda n. (add\ n\ 1)) ((\lambda n. (add\ n\ 1))\ x)))\ 5 &\Rightarrow_{\beta} \\ (\lambda n. (add\ n\ 1)) ((\lambda n. (add\ n\ 1))\ 5) &\Rightarrow_{\beta} \\ (add\ ((\lambda n. (add\ n\ 1))\ 5)\ 1) &\Rightarrow_{\beta} \\ (add\ (add\ 5\ 1)\ 1) &\Rightarrow_{\delta} 7 \end{aligned}$$



Evaluation Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Call-by-Value (CBV)
 - C / C++: the argument expression is evaluated, and the resulting value is bound to the corresponding variable in the function (frequently by copying the value into a new memory region)
- Call-by-Reference (CBR)
 - C++: a function receives an implicit reference to a variable used as argument, rather than a copy of its value
 - CBR may be simulated in languages that use CBV by making use of references, such as pointers (Call-by-Address or CBA)
- Call-by-Copy-Restore (CBCR) / Value-Result
 - Fortran (old): a special case of call by reference where the provided reference is unique to the caller (Copy-in-Copy-out)
- Call-by-Name (CBN)
 - C / C++ Macro: the arguments to a function are not evaluated before the function is called – rather, they are substituted directly into the function body
 - Lazy Evaluation
 - Call-by-Need: a memorized variant of CBN where, if the function argument is evaluated, that value is stored for subsequent uses



Evaluation Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

```
#include <iostream>
using namespace std;

void f(int a, int b) { a++; b--; return; }           // CBV
void g(int& a, int& b) { a++; b--; return; }         // CBR
void h(int* pa, int* pb) { (*pa)++; (*pb)--; return; } // CBA
#define m_f(a, b) ( a * b )                         // CBN

int main() {
    int x = 3, y = 4, z = 5;
    f(x, y);
    cout << x << " " << y << endl;                 // CBV = 3 4
    g(x, y);
    cout << x << " " << y << endl;                 // CBR = 4 3
    h(&x, &y);
    cout << x << " " << y << endl;                 // CBA = 5 2
    g(z, z);
    cout << z << endl;                             // CBR = 5, CBCR = 6 or 4
    cout << m_f(x + 1, y + 1) << endl;              // CBN = x + y + 1 = 8

    return 0;
}
```



Reduction Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

Definition: A λ -expression is in **normal form** if it contains no β -redexes (and no δ -rules in an applied λ calculus), so that it cannot be further reduced using the β -rule or the δ -rule.

An expression in normal form has no more function applications to evaluate.



Reduction Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

Questions:

- 1 Can every λ -expression be reduced to a normal form?
- 2 Is there more than one way to reduce a particular λ -expression?
- 3 If there is more than one reduction strategy, does each one lead to the same normal form expression?
- 4 Is there a reduction strategy that will guarantee that a normal form expression will be produced?



Reduction Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

1. Can every λ -expression be reduced to a normal form?

No. Consider:

$$(\lambda x. x x)(\lambda x. x x) \Rightarrow$$

$$(\lambda x. x x)(\lambda x. x x) \Rightarrow$$

$$(\lambda x. x x)(\lambda x. x x) \Rightarrow$$

...



Reduction Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

2. Is there more than one way to reduce a particular λ -expression?

Yes. Consider:

$$(\lambda x. \lambda y. (add\ y\ ((\lambda z. (mul\ x\ z))\ 3)))\ 7\ 5$$

Path 1: OUTERMOST

$$(\lambda x. \lambda y. (add\ y\ ((\lambda z. (mul\ x\ z))\ 3)))\ 7\ 5 \Rightarrow_{\beta}$$

$$(\lambda y. (add\ y\ ((\lambda z. (mul\ 7\ z))\ 3)))\ 5 \Rightarrow_{\beta}$$

$$(add\ 5\ ((\lambda z. (mul\ 7\ z))\ 3)) \Rightarrow_{\beta} (add\ 5\ (mul\ 7\ 3)) \Rightarrow_{\delta} (add\ 5\ 21) \Rightarrow_{\delta} 26$$

Path 2: INNERMOST

$$(\lambda x. \lambda y. (add\ y\ ((\lambda z. (mul\ x\ z))\ 3)))\ 7\ 5 \Rightarrow_{\beta}$$

$$(\lambda x. \lambda y. (add\ y\ (mul\ x\ 3)))\ 7\ 5 \Rightarrow_{\beta} (\lambda x. (add\ 5\ (mul\ x\ 3)))\ 7 \Rightarrow_{\beta}$$

$$(add\ 5\ (mul\ 7\ 3)) \Rightarrow_{\delta} (add\ 5\ 21) \Rightarrow_{\delta} 26$$

Path 3: MIXED

$$(\lambda x. \lambda y. (add\ y\ ((\lambda z. (mul\ x\ z))\ 3)))\ 7\ 5 \Rightarrow_{\beta}$$

$$(\lambda x. \lambda y. (add\ y\ (mul\ x\ 3)))\ 7\ 5 \Rightarrow_{\beta} (\lambda y. (add\ y\ (mul\ 7\ 3)))\ 5 \Rightarrow_{\delta}$$

$$(\lambda y. (add\ y\ 21))\ 5 \Rightarrow_{\beta} (add\ 5\ 21) \Rightarrow_{\delta} 26$$



Reduction Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- 3. If there is more than one reduction strategy, does each one lead to the same normal form expression?**

No. Consider:

$$(\lambda y. 5)((\lambda x. x x)(\lambda x. x x))$$

Path 1:

$$(\lambda y. 5)((\lambda x. x x)(\lambda x. x x)) \Rightarrow 5$$

Path 2:

$$(\lambda y. 5)((\lambda x. x x)(\lambda x. x x)) \Rightarrow$$

$$(\lambda y. 5)((\lambda x. x x)(\lambda x. x x)) \Rightarrow$$

$$(\lambda y. 5)((\lambda x. x x)(\lambda x. x x)) \Rightarrow$$

...



Reduction Strategies

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

4. Is there a reduction strategy that will guarantee that a normal form expression will be produced?

Mathematician Curry proved that if an expression has a normal form, then it can be found by leftmost reduction.

A normal order reduction can have either of the following outcomes:

- 1 It reaches a unique (up to α -conversion) normal form λ -expression
- 2 It never terminates

Unfortunately, there is no algorithmic way to determine for an arbitrary λ -expression which of these two outcomes will occur



Reduction Strategies: Normal and Applicative Order

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

Two important orders of rewriting:

- **Normal Order** – rewrite the outermost (leftmost) occurrence of a function application.
 - *This is equivalent to call by name.*
- **Applicative Order** – rewrite the innermost (leftmost) occurrence of a function application first.
 - *This is equivalent to call by value.*

Normal order evaluation always gives the same results as lazy evaluation, but may end up evaluating an expression more times.



Order of Evaluation

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Example:

$$\text{double } x = x + x$$

$$\text{average } x \ y = (x + y)/2$$

- Using prefix notation:

$$\text{double } x = \text{plus } x \ x$$

$$\text{average } x \ y = \text{divide } (\text{plus } x \ y) \ 2$$

- Evaluate:

$$\text{double } (\text{average } 2 \ 4)$$



Order of Evaluation

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and

Applicative Order

- Evaluate:

double (average 2 4)

- Using normal order of evaluation:

*double (average 2 4) \Rightarrow plus (average 2 4) (average 2 4) \Rightarrow
plus (divide (plus 2 4) 2) (average 2 4) \Rightarrow
plus (divide 6 2) (average 2 4) \Rightarrow plus 3 (average 2 4) \Rightarrow
plus 3 (divide (plus 2 4) 2) \Rightarrow plus 3 (divide 6 2) \Rightarrow
plus 3 3 \Rightarrow 6*

- Notice that (average 2 4) was evaluated twice ... lazy evaluation would cache the results of the first evaluation

- Using applicative order of evaluation:

*double (average 2 4) \Rightarrow double (divide (plus 2 4) 2) \Rightarrow
double (divide 6 2) \Rightarrow double 3 \Rightarrow plus 3 3 \Rightarrow 6*



Order of Evaluation

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Consider:
 $my_if\ True\ x\ y = x$
 $my_if\ False\ x\ y = y$
- Evaluate:
 $my_if\ (less\ 3\ 4)\ (plus\ 5\ 5)\ (divide\ 1\ 0)$
- Using normal order of evaluation:
 $my_if\ (less\ 3\ 4)\ (plus\ 5\ 5)\ (divide\ 1\ 0) \Rightarrow$
 $my_if\ True\ (plus\ 5\ 5)\ (divide\ 1\ 0) \Rightarrow (plus\ 5\ 5) \Rightarrow$
 10
- Using applicative order of evaluation:
 $my_if\ (less\ 3\ 4)\ (plus\ 5\ 5)\ (divide\ 1\ 0) \Rightarrow$
 $my_if\ True\ (plus\ 5\ 5)\ (divide\ 1\ 0) \Rightarrow$
 $my_if\ True\ 10\ (divide\ 1\ 0) \Rightarrow$
 $DIVIDE\ BY\ ZERO\ ERROR$



Properties of Order of Evaluation: Strictness

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and

Applicative Order

Two important properties of evaluation order:

- If there is any evaluation order that will terminate and that will not generate an error, normal order evaluation will terminate and will not generate an error.
- ANY evaluation order that terminates without error will give the same result as any other evaluation order that terminates without error.

Definition: A function f is *strict* in an argument if that argument is always evaluated whenever an application of f is evaluated.

If a function is strict in an argument, we can safely evaluate the argument first if we need the value of applying the function.



Lazy Evaluation and Strictness Analysis

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

We can use lazy evaluation on an ad-hoc basis (e.g. for *if*), for all arguments.

For all arguments, for some implementations of functional languages we can improve efficiency using strictness analysis.

plus $a\ b$ is strict in both arguments

if $x\ y\ z$ is strict in x , but not in y and z

We can do some analysis and sometimes decide if a user-defined function is strict in some of its arguments:



Lazy Evaluation and Strictness Analysis

PoPL

Partha Pratim
Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

Examples:

- *double* x is strict in x
- *squid* $n\ x = \text{if } n = 0 \text{ then } x + 1 \text{ else } x - n$ is strict in n and x
- *crab* $n\ x = \text{if } n = 0 \text{ then } x + 1 \text{ else } n$ is strict in n but not x

If a function is strict in an argument x , it is correct to pass x by value, even with normal order evaluation semantics.

It is not always decidable whether a function is strict in an argument – if we do not know, pass using lazy evaluation.



Reduction Strategies: Normal and Applicative Order

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- *Definition:*

A **normal order reduction** always reduces the *leftmost outermost* β -redex (or δ -redex) first.

An **applicative order reduction** always reduces the *leftmost innermost* β -redex (or δ -redex) first.

- *Definition:*

For any λ -expression of the form $E = ((\lambda x. B) A)$, we say that β -redex E is outside any β -redex that occurs in B or A and that these are inside E .

A β -redex in a λ -expression is outermost if there is no β -redex outside of it, and it is innermost if there is no β -redex inside of it.

- Use AST for detection



AST of λ -expression

PoPL

Partha Pratim Das

Semantics

Free and Bound

Variables

Substitution

Reduction

α -Reduction


β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order



β -redexes in $((((\lambda x. \lambda y. (add\ x\ y)) ((\lambda z. (succ\ z))\ 5)) ((\lambda w. (sqr\ w))\ 7)))$



Order of Evaluation

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and
Applicative Order

- Applicative Order (*leftmost innermost*)
 $((\lambda n. (add\ 5\ n))\ 8) \Rightarrow$
 $((\lambda n. (add5\ n))\ 8) \Rightarrow -\ add5 : N \rightarrow N$ is curried
 $(add5\ 8) \Rightarrow 13$
 - Lazy Evaluation
 - Call-by-Name (CBN)
 - Curried functions $(f\ x\ y\ z)$ use lazy reduction
- Normal Order (*leftmost outermost*)
 $((\lambda n. (add\ 5\ n))\ 8) \Rightarrow (add\ 5\ 8) \Rightarrow 13$
 - Eager Evaluation
 - Call-by-Value (CBV)
 - Function $f(x, y, z)$ use eager reduction



Order of Evaluation

PoPL

Partha Pratim
Das

Semantics

Free and Bound
Variables

Substitution

Reduction

α -Reduction

β -Reduction

η -Reduction

δ -Reduction

Order of Evaluation

Normal and

Applicative Order

