
Pairs that share a molecule have more similar yield profiles in every screen
Analysis screens, product-forming pairs; additive residual yield profiles
pairs sharing only the role-1 molecule
pairs sharing only the role-2 molecule
pairs sharing neither molecule
C–N (Pd ligand)
C–N (Pd/Cu/Ni)
C–O (Pd/Cu)
Heck (solvent/base)
Hiyama (Pd)
Kumada (Pd/Ni/Fe)
SNAr (base/solvent)
Sonogashira (solvent/base)
Suzuki (Pd)
Ullmann C–O (Cu)
Amide coupling
Chan–Lam (Cu)
Esterification
Grignard addition
Reductive amination
Median over screens
1: N–H partner · 2: aryl (pseudo)halide
1: N–H partner · 2: aryl halide
1: phenol or alcohol · 2: aryl halide
1: alkene · 2: aryl halide
1: organosilane · 2: aryl halide
1: aryl or acyl halide · 2: Grignard reagent
1: N–H nucleophile · 2: aryl halide
1: alkyne · 2: aryl halide
1: aryl halide · 2: boronic acid
1: phenol · 2: aryl halide
1: N–H partner · 2: carboxylic acid
1: N–H partner · 2: boronic acid
1: alcohol · 2: carboxylic acid
1: Grignard reagent · 2: carbonyl compound
1: carbonyl compound · 2: amine
either above neither in 15 of 15; both in 13
−0.2
0.0
0.2
0.4
0.6
Mean correlation of yield profiles between product-forming pairs