

a
Amide > C–N mean: 6/8
Grey: two reverse cases
Condition-preference share, T
0.4
0.0
−0.4
Amide coupling
C–N ligand library
C–N metal library
b
Owner context: 0.26 vs 0.06
Higher owner share: 17/23
Owner
Non-owner
3-(methylthio)aniline
Amine
3,4-dimethoxyaniline
2-ethoxypyridin-4-amine
3-(4-methylimidazol-1-yl)-5-(trifluoromethyl)aniline
2,4-dimethylaniline
5-methylpyridin-2-amine
4-methoxyaniline
4-methoxy-N-methylaniline
3-ethynylaniline
4-(methylthio)aniline
4-methylaniline
6-methoxypyridin-3-amine
1-butylpiperazine
ethyl 3-aminobenzoate
4-methoxy-3-methylaniline
N-ethylaniline
2-methylaniline
quinoxalin-6-amine
methyl 4-amino-2-methoxybenzoate
3,4,5-trimethoxybenzoic acid
Acid
5-(trifluoromethyl)picolinic acid
5-bromo-2-methoxynicotinic acid
5-fluoropicolinic acid
−0.4
0.0
0.4
0.8
Condition-preference share, T
c
SNAr: halide 45.6%, N–H 0.3%
Base/acid and solvent
Fixed halide
Fixed N–H partner
H1
H2
H3
H4
N1
N2
N3
N4
N5
N6
N7
N8
Condition preference (pp)
−30
0
30
47 conditions; common order