

Two gravitational counting conditions for three generations are the same condition

Justin Pulford^{1,*}

¹*Unaffiliated*

(Dated: August 2, 2026)

This note is a comment on existing literature, not a new constraint or discovery. Two conditions of gravitational origin have been used to constrain the field content of the Standard Model. Cancelling the conformal anomaly requires both trace-anomaly coefficients to vanish, and has been shown to select three generations together with three right-handed neutrinos. Separately, Pauli’s finiteness requirement for the one-loop contribution to Newton’s constant in induced gravity asks that a supertrace of heat-kernel coefficients vanish, $\text{str}[k_1] = 0$, and is satisfied by the same content. These are not independent constraints. Eliminating the auxiliary scalar count from the two anomaly conditions leaves $N_{1/2} = 4N_1 - N_0/2$, and $\text{str}[k_1] = 0$ evaluates to $N_{1/2} = 4N_1$; the two coincide whenever conventional scalars decouple from the count. Both are then solved by $N_{1/2} = 48$ at $N_1 = 12$. Agreement between them is therefore arithmetic rather than evidential, and should not be read as two independent gravitational arguments for the observed generation number. The conditions are nevertheless distinguishable, and they are distinguished by the Higgs: anomaly cancellation forbids a fundamental Higgs doublet and requires thirty-six additional scalars, whereas the finiteness condition admits the doublet only if one assumes conformal coupling, $\xi = 1/6$ — an extra assumption, not fixed by the Standard Model — and then requires nothing new. We record the reduction explicitly and note that no calculation selects between the two conditions — only their differing scalar sectors do.

I. INTRODUCTION

That the Standard Model contains three generations is not explained within it. Constraints of gravitational origin are an attractive place to look for an explanation, because they are sensitive to the total field content rather than to any one sector, and because they involve no free parameters beyond the spins and multiplicities already measured.

Two such constraints have been used. The first is the cancellation of the conformal anomaly. Requiring both trace-anomaly coefficients to vanish, Navarro-Salas [1] finds a solution at $N_{1/2} = 48$ Weyl fields with $N_1 = 12$ gauge fields — three generations including three right-handed neutrinos — at the cost of removing conventional scalars from the spectrum and adding thirty-six fields of a different type.

The second descends from Pauli and from Sakharov’s induced gravity [2]. If the Einstein–Hilbert term is generated by matter loops, its coefficient is a sum over the spectrum of heat-kernel coefficients, and Visser [3] records that the one-loop contribution to Newton’s constant is finite when the supertrace $\text{str}[k_1]$ vanishes. Evaluated on Standard Model content with three right-handed neutrinos, it does — provided one takes the Higgs to be conformally coupled, $\xi = 1/6$. That value is an extra assumption: the Standard Model does not fix ξ , and conformal coupling is chosen for naturalness, not measured.

The coincidence invites a natural reading: two independent gravitational arguments, from different one-loop

coefficients, converging on the same field content. That reading is wrong. This note is a clarification of the existing literature, not a new counting constraint. The conditions reduce to the same counting relation; what looks like corroboration is an identity.

II. THE TWO CONDITIONS

Write N_0 , $N_{1/2}$, N_1 for the numbers of real scalar, Weyl fermion and gauge fields.

A. Conformal anomaly cancellation

The trace anomaly of a conformally invariant theory in four dimensions is fixed by two coefficients. Requiring both to vanish gives [1]

$$a \propto N_0 + \frac{11}{2}N_{1/2} + 62N_1 - 28N_\xi = 0, \quad (1)$$

$$c \propto N_0 + 3N_{1/2} + 12N_1 - 8N_\xi = 0, \quad (2)$$

where N_ξ counts fields of a fourth-order conformally invariant type, which contribute to the anomaly with the opposite sign to ordinary scalars and so make cancellation possible at all.

B. Finiteness of the induced Newton constant

In induced gravity the inverse Newton constant is a one-loop sum over the spectrum, $1/G \simeq -\text{str}[k_1] \kappa^2/2\pi$ with κ the cutoff, and the coefficients k_1 are those of the R term in the Seeley–DeWitt expansion [3, 4]. Per field

* pulfordj420@gmail.com

they are $1/6 - \xi$ for a real scalar with curvature coupling ξ , $-1/6$ for a Weyl spinor and $-2/3$ for a massless vector, with the supertrace supplying the fermionic sign. The finiteness condition $\text{str}[k_1] = 0$ is then

$$\frac{1}{6}N_{1/2} + \left(\frac{1}{6} - \xi\right)N_0 - \frac{2}{3}N_1 = 0. \quad (3)$$

III. THE REDUCTION

Equations (1) and (2) contain the auxiliary count N_ξ , which is not observable and is fixed by the conditions themselves. Solving (2) for it, $N_\xi = (N_0 + 3N_{1/2} + 12N_1)/8$, and substituting into (1):

$$-\frac{5}{2}N_0 - 5N_{1/2} + 20N_1 = 0 \quad \Longleftrightarrow \quad N_{1/2} = 4N_1 - \frac{1}{2}N_0. \quad (4)$$

Equation (3), meanwhile, gives $N_{1/2} = 4N_1$ whenever the scalar term drops out — that is, if one assumes $\xi = 1/6$, or trivially at $N_0 = 0$. The conformal value is not free: it is an input, motivated by naturalness of conformal coupling, not by any measurement of the Higgs nonminimal coupling.

So both conditions are the single relation

$$\boxed{N_{1/2} = 4N_1} \quad (5)$$

once conventional scalars are removed from the count, whether by setting $N_0 = 0$ or by assuming conformal coupling. With $N_1 = 12$ for $SU(3) \times SU(2) \times U(1)$ this is $N_{1/2} = 48$: sixteen Weyl fields per generation, which is fifteen plus a right-handed neutrino, three times over.

The agreement between the two conditions is therefore an identity, not a corroboration. A spectrum satisfying one satisfies the other automatically, and the fact that both select the observed generation number is a single fact rather than two.

IV. WHAT STILL SEPARATES THEM

The conditions are not the same physics, and the difference sits entirely in the scalar sector.

Anomaly cancellation cannot tolerate conventional scalars: solving (1) and (2) together requires $N_0 = 0$, from which Navarro-Salas concludes that the Higgs doublet “*cannot be considered as [a] fundamental [entity]*”, and a solution then exists only with $N_\xi = 36$ additional fourth-order fields [1]. Had the doublet been retained at $N_0 = 4$, (4) would demand $N_{1/2} = 46$, which the Standard Model with three right-handed neutrinos misses by two.

The finiteness condition takes the other route. Equation (3) removes the scalar contribution not by deleting the field but by taking $\xi = 1/6$, the conformal value, at which $1/6 - \xi$ vanishes identically. That is an extra assumption: ξ is not fixed by the Standard Model Lagrangian as usually written, and conformal coupling is

TABLE I. The two conditions agree on the counting relation and disagree on everything else about the scalar sector.

	anomaly cancellation	$\text{str}[k_1] = 0$
relation	$N_{1/2} = 4N_1 - N_0/2$	$N_{1/2} = 4N_1$
Higgs doublet	excluded	retained if $\xi = 1/6$ (assumed)
extra fields	$N_\xi = 36$	none
$N_{1/2}$ at $N_1 = 12$	48	48

chosen because it is natural in the absence of a dimensional scale that would pick another value. With that assumption the doublet is kept, nothing is added, and the price is a condition on the curvature coupling rather than on existence.

Table I is the whole note in miniature. The interesting rows are not the last one, where the two agree and are often quoted together, but the two above it, where they are incompatible. A fundamental Higgs doublet is forbidden by one condition and admitted by the other only under the extra assumption $\xi = 1/6$; thirty-six new fields are required by one and by the other not at all.

V. REMARKS

This is a comment on existing literature, not a new result. The generation-counting result is Navarro-Salas’s [1]; the finiteness condition is Pauli’s, in the form Visser records [3]. The only contribution here is the observation that the two counting relations are the same relation once scalars are handled, and that the scalar sector is what still distinguishes the conditions physically.

Imposing $\text{str}[k_1] = 0$ removes the quadratically divergent term that would generate $1/G$. It does not induce gravity; it makes the one-loop correction to a tree-level G finite. Visser notes that this is “*completely at odds with Sakharov’s original version*” [3]. The condition should be read as a naturalness statement about Newton’s constant, not as an account of its origin. The further assumption $\xi = 1/6$ for the Higgs is of the same kind: naturalness of conformal coupling, not a free output of the counting.

Pauli’s mass sum rules $\sum_n (-1)^{2S_n} g_n m_n^{2k} = 0$ are a separate matter: they are unweighted mass relations and do not reduce to (5). (Evaluated on the Standard Model with right-handed neutrinos they instead give $N_{\text{BSM}} = 68$ bosonic degrees of freedom beyond the Standard Model [5].) This note concerns only the two curvature-weighted conditions.

Nothing in the counting separates the conditions observationally. The scalar sector does: a fundamental Higgs doublet is compatible with $\text{str}[k_1] = 0$ only under the assumed $\xi = 1/6$, and not with anomaly cancellation at all, and the thirty-six fields the latter requires are a spectrum, not a bookkeeping device. Evidence for either is therefore evidence between them — a distinction that

matters precisely because the counting agreement is not.

- [1] J. Navarro-Salas, Black holes, conformal symmetry, and fundamental fields, *Class. Quantum Grav.* **41** (2024); arXiv:2403.13201.
- [2] A. D. Sakharov, Vacuum quantum fluctuations in curved space and the theory of gravitation, *Sov. Phys. Dokl.* **12** (1968) 1040.
- [3] M. Visser, Sakharov's induced gravity: a modern perspective, *Mod. Phys. Lett. A* **17** (2002) 977; arXiv:gr-qc/0204062.
- [4] S. M. Christensen and S. A. Fulling, Trace anomalies and the Hawking effect, *Phys. Rev. D* **15** (1977) 2088.
- [5] M. Visser, The Pauli sum rules imply BSM physics, *Phys. Lett. B* **791** (2019) 43; arXiv:1808.04583.