

The Normal-Gamma conjugate family

Dr. Merlise Clyde



prior for mean & variance

- ▶ conditional prior $\mu | \sigma^2 \sim N(m_0, \sigma^2/n_0)$

prior for mean & variance

- ▶ conditional prior $\mu | \sigma^2 \sim N(m_0, \sigma^2/n_0)$
- ▶ precision $\phi = 1/\sigma^2$

prior for mean & variance

- ▶ conditional prior $\mu | \sigma^2 \sim N(m_0, \sigma^2/n_0)$
- ▶ precision $\phi = 1/\sigma^2$
- ▶ conjugate prior for ϕ is a Gamma distribution

$$\phi \sim \text{Gamma}(v_0/2, s_0^2 v_0/2)$$

prior for mean & variance

- ▶ conditional prior $\mu | \sigma^2 \sim N(m_0, \sigma^2/n_0)$
- ▶ precision $\phi = 1/\sigma^2$
- ▶ conjugate prior for ϕ is a Gamma distribution

$$\phi \sim \text{Gamma}(v_0/2, s_0^2 v_0/2)$$

or

$$1/\sigma^2 \sim \text{Gamma}(v_0/2, s_0^2 v_0/2)$$

prior for mean & variance

- ▶ conditional prior $\mu | \sigma^2 \sim N(m_0, \sigma^2/n_0)$
- ▶ precision $\phi = 1/\sigma^2$
- ▶ conjugate prior for ϕ is a Gamma distribution

$$\phi \sim \text{Gamma}(v_0/2, s_0^2 v_0/2)$$

or

$$1/\sigma^2 \sim \text{Gamma}(v_0/2, s_0^2 v_0/2)$$

- ▶ together $(\mu, \phi) \sim \text{NormalGamma}(m_0, n_0, s_0^2, v_0)$

conjugate posterior distribution

- ▶ prior $(\mu, \phi) \sim \text{NormalGamma}(m_0, n_0, s_0^2, v_0)$

conjugate posterior distribution

- ▶ prior $(\mu, \phi) \sim \text{NormalGamma}(m_0, n_0, s_0^2, v_0)$
- ▶ posterior $(\mu, \phi) | \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$

conjugate posterior distribution

- ▶ prior $(\mu, \phi) \sim \text{NormalGamma}(m_0, n_0, s_0^2, v_0)$
- ▶ posterior $(\mu, \phi) | \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$
- ▶ updated parameters:

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0}$$

$$n_n = n_0 + n$$

conjugate posterior distribution

- ▶ prior $(\mu, \phi) \sim \text{NormalGamma}(m_0, n_0, s_0^2, v_0)$
- ▶ posterior $(\mu, \phi) | \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$
- ▶ updated parameters:

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0}$$

$$n_n = n_0 + n$$

$$v_n = v_0 + n$$

conjugate posterior distribution

- ▶ prior $(\mu, \phi) \sim \text{NormalGamma}(m_0, n_0, s_0^2, v_0)$
- ▶ posterior $(\mu, \phi) | \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$
- ▶ updated parameters:

$$m_n = \frac{n\bar{Y} + n_0 m_0}{n + n_0}$$

$$n_n = n_0 + n$$

$$v_n = v_0 + n$$

$$s_n^2 = \frac{1}{v_n} \left[s_0^2 v_0 + s^2 (n - 1) + \frac{n_0 n}{n_n} (\bar{Y} - m_0)^2 \right]$$

inference about μ

- ▶ joint distribution

$$(\mu, \phi) \mid \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$$

inference about μ

- ▶ joint distribution

$$(\mu, \phi) \mid \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$$

- ▶ equivalent to

$$\mu \mid \text{data}, \sigma^2 \sim N(m_n, \sigma^2/n_n)$$

inference about μ

- ▶ joint distribution

$$(\mu, \phi) \mid \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$$

- ▶ equivalent to

$$\mu \mid \text{data}, \sigma^2 \sim \mathcal{N}(m_n, \sigma^2/n_n)$$

$$1/\sigma^2 \mid \text{data} \sim \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

inference about μ

- ▶ joint distribution

$$(\mu, \phi) \mid \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$$

- ▶ equivalent to

$$\mu \mid \text{data}, \sigma^2 \sim N(m_n, \sigma^2/n_n)$$

$$1/\sigma^2 \mid \text{data} \sim \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

- ▶ marginal distribution student t distribution

$$\mu \mid \text{data} \sim t(v_n, m_n, s_n^2/n_n)$$

inference about μ

- ▶ joint distribution

$$(\mu, \phi) \mid \text{data} \sim \text{NormalGamma}(m_n, n_n, s_n^2, v_n)$$

- ▶ equivalent to

$$\mu \mid \text{data}, \sigma^2 \sim N(m_n, \sigma^2/n_n)$$

$$1/\sigma^2 \mid \text{data} \sim \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

- ▶ marginal distribution student t distribution

$$\mu \mid \text{data} \sim t(v_n, m_n, s_n^2/n_n) \Leftrightarrow t = \frac{\mu - m_n}{s_n / \sqrt{n_n}} \sim t(v_n, 0, 1)$$

total trihalomethane in tapwater (TTHM)

prior $\text{NormalGamma}(35, 25, 156.25, 24)$



total trihalomethane in tapwater (TTHM)

prior $\text{NormalGamma}(35, 25, 156.25, 24)$

data $\bar{Y} = 55.5$, $s^2 = 540.7$, $n = 28$



total trihalomethane in tapwater (TTHM)

prior $\text{NormalGamma}(35, 25, 156.25, 24)$

data $\bar{Y} = 55.5$, $s^2 = 540.7$, $n = 28$

$$n_n = 25 + 28 = 53$$

$$m_n = \frac{28 \times 55.5 + 25 \times 35}{53} = 45.8$$

$$v_n = 24 + 28 = 52$$


$$\begin{aligned}s_n^2 &= \frac{(n-1)s^2 + v_0 s_0^2 + n_0 n(m_0 - \bar{Y})^2 / n_n}{52} \\ &= \frac{1}{52} [27 \times 540.7 + 24 \times 156.25 + \frac{25 \times 28}{53} \times (35 - 55.5)^2] = 459.6\end{aligned}$$

posterior $\text{NormalGamma}(45.8, 53, 459.6, 52)$




credible interval for μ

95% credible interval: (L, U) where $P(L \leq \mu \leq U | \text{data}) = 0.95$



credible interval for μ

95% credible interval: (L, U) where $P(L \leq \mu \leq U | \text{data}) = 0.95$



$$L = t_{0.025} \sqrt{s_n^2/n_n} + m_n$$

$$U = t_{0.975} \sqrt{s_n^2/n_n} + m_n$$

using R

```
m_0 = 35; n_0 = 25; s2_0 = 156.25; v_0 = n_0 - 1
data(tapwater); Y = tapwater$tthm
ybar = mean(Y); s2 = var(Y); n = length(Y)
n_n = n_0 + n
m_n = (n*ybar + n_0*m_0)/n_n
v_n = v_0 + n
s2_n = ((n-1)*s2 + v_0*s2_0 + n_0*n*(m_0 - ybar)^2/n_n)/v_n
# CI
L = qt(.025, v_n)*sqrt(s2_n/n_n) + m_n
U = qt(.975, v_n)*sqrt(s2_n/n_n) + m_n
c(L, U)

## [1] 39.93192 51.75374
```

using R

```
m_0 = 35; n_0 = 25; s2_0 = 156.25; v_0 = n_0 - 1
data(tapwater); Y = tapwater$tthm
ybar = mean(Y); s2 = var(Y); n = length(Y)
n_n = n_0 + n
m_n = (n*ybar + n_0*m_0)/n_n
v_n = v_0 + n
s2_n = ((n-1)*s2 + v_0*s2_0 + n_0*n*(m_0 - ybar)^2/n_n)/v_n
# CI
L = qt(.025, v_n)*sqrt(s2_n/n_n) + m_n
U = qt(.975, v_n)*sqrt(s2_n/n_n) + m_n
c(L, U)

## [1] 39.93192 51.75374
```

there is a 95% chance that the mean TTHMs are between
39.9 ppb and 51.8 ppb

summary

- ▶ introduced conjugate Normal-Gamma family and updating

summary

- ▶ introduced conjugate Normal-Gamma family and updating
- ▶ introduced Student t distribution for inference about μ unconditionally

summary

- ▶ introduced conjugate Normal-Gamma family and updating
- ▶ introduced Student t distribution for inference about μ unconditionally
- ▶ compute credible intervals in R using Student t

summary

- ▶ introduced conjugate Normal-Gamma family and updating
- ▶ introduced Student t distribution for inference about μ unconditionally
- ▶ compute credible intervals in R using Student t

next: using Monte Carlo simulation to explore prior and posterior distributions