

inference via Monte Carlo sampling

Dr. Merlise Clyde

joint posterior distribution

- ▶ conditional posterior $\mu \mid \text{data}, \sigma^2 \sim \text{N}(m_n, \sigma^2/n_n)$

joint posterior distribution

▶ conditional posterior $\mu \mid \text{data}, \sigma^2 \sim \mathbf{N}(m_n, \sigma^2/n_n)$

▶ marginal posterior

$$1/\sigma^2 = \phi \mid \text{data} \sim \text{Gamma}(v_n/2, v_n s_n^2/2)$$

joint posterior distribution

- ▶ conditional posterior $\mu \mid \text{data}, \sigma^2 \sim \mathbf{N}(m_n, \sigma^2/n_n)$
- ▶ marginal posterior
 $1/\sigma^2 = \phi \mid \text{data} \sim \text{Gamma}(v_n/2, v_n s_n^2/2)$
- ▶ marginal posterior $\mu \mid \text{data} \sim \mathbf{t}(v_n, m_n, s_n^2/n_n)$

joint posterior distribution

- ▶ conditional posterior $\mu \mid \text{data}, \sigma^2 \sim \mathbf{N}(m_n, \sigma^2/n_n)$
- ▶ marginal posterior
 $1/\sigma^2 = \phi \mid \text{data} \sim \text{Gamma}(v_n/2, v_n s_n^2/2)$
- ▶ marginal posterior $\mu \mid \text{data} \sim \mathbf{t}(v_n, m_n, s_n^2/n_n)$
- ▶ inference about σ or other transformations of the parameters?

Monte Carlo simulation



<https://pixabay.com/en/gambling-roulette-game-bank-2001079/>

Monte Carlo inference for the precision

- ▶ draw S samples of ϕ from its posterior distribution

$$\phi^{(1)}, \phi^{(2)}, \dots, \phi^{(S)} \stackrel{\text{iid}}{\sim} \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

Monte Carlo inference for the precision

- ▶ draw S samples of ϕ from its posterior distribution

$$\phi^{(1)}, \phi^{(2)}, \dots, \phi^{(S)} \stackrel{\text{iid}}{\sim} \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

- ▶ empirical distribution of the samples approximates the posterior distribution

Monte Carlo inference for the precision

- ▶ draw S samples of ϕ from its posterior distribution

$$\phi^{(1)}, \phi^{(2)}, \dots, \phi^{(S)} \stackrel{\text{iid}}{\sim} \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

- ▶ empirical distribution of the samples approximates the posterior distribution
- ▶ estimate means, quantiles, probabilities or other functions of ϕ from the samples

Monte Carlo inference for the precision

- ▶ draw S samples of ϕ from its posterior distribution

$$\phi^{(1)}, \phi^{(2)}, \dots, \phi^{(S)} \stackrel{\text{iid}}{\sim} \text{Gamma}(v_n/2, s_n^2 v_n/2)$$

- ▶ empirical distribution of the samples approximates the posterior distribution
- ▶ estimate means, quantiles, probabilities or other functions of ϕ from the samples
- ▶ improved approximation as S increases

$$\frac{\sum_{i=1}^S g(\phi^{(i)})}{S} \rightarrow \mathbf{E}[g(\phi) \mid \text{data}]$$

tap water example


R code

```
set.seed(8675309)
phi = rgamma(1000, shape = v_n/2, rate=s2_n*v_n/2)
```

tap water example

R code

```
set.seed(8675309)
phi = rgamma(1000, shape = v_n/2, rate=s2_n*v_n/2)
```



inference about σ

R code

```
phi = rgamma(1000, shape = v_n/2, rate=s2_n*v_n/2)
sigma = 1/sqrt(phi)
mean(sigma)      # posterior mean of sigma

## [1] 21.79296

quantile(sigma, c(0.025, .975))

##      2.5%|      97.5%
## 18.13778 26.60354
```

summary

- ▶ introduced Monte Carlo sampling

summary

- ▶ introduced Monte Carlo sampling
- ▶ compute means, quantiles, probabilities, etc.

summary

- ▶ introduced Monte Carlo sampling
- ▶ compute means, quantiles, probabilities, etc.
- ▶ transformations of parameters

summary

- ▶ introduced Monte Carlo sampling
- ▶ compute means, quantiles, probabilities, etc.
- ▶ transformations of parameters

next: predictive distributions and choice of hyper-parameters