

The complexity of 3-colouring H -colourable graphs. Andrei Krokhin and Jakub Oprsal, 2019

Szymon Wojtulewicz

Uniwersytet Jagielloński

20.11.2023

Main result

Theorem 3.1

Let \mathbf{H} be a 3-colourable non-bipartite graph. Then $\text{PCSP}(\mathbf{H}, \mathbf{K}_3)$ is NP-hard.

Dichotomy

If \mathbf{H} is bipartite then the problem can be solved in polynomial time by using an algorithm for 2-colouring.

Reduction

$\text{PCSP}(\mathbf{H}, \mathbf{K}_3)$ admits a reduction from $\text{PCSP}(\mathbf{C}_k, \mathbf{K}_3)$ where k is the size of an odd cycle in \mathbf{H} .

Theorems 2.13 and 3.3

Theorem 2.13

Let \mathbf{H}, \mathbf{G} be graphs such that $\mathbf{H} \rightarrow \mathbf{G}$. Assume that there exists a minion homomorphism $\xi: \text{Pol}(\mathbf{H}, \mathbf{G}) \rightarrow \mathcal{M}$ for some minion \mathcal{M} on a pair of sets such that \mathcal{M} has bounded essential arity and does not contain a constant function. Then $\text{PCSP}(\mathbf{H}, \mathbf{G})$ is NP-hard.

Definition

Let N be an odd number, we define a minion $\mathcal{Z}_{\leq N}$ to be the set of all functions $f: \mathbb{Z}^n \rightarrow \mathbb{Z}$ such that $f(x_1, \dots, x_n) = c_1x_1 + \dots + c_nx_n$ for some $c_1, \dots, c_n \in \mathbb{Z}$ with $\sum_{i=1}^n |c_i| \leq N$ and $\sum_{i=1}^n c_i$ odd.

Theorem 3.3

Let $k \geq 3$ be odd and let N be the largest odd number such that $N \leq k/3$. Then there is a minion homomorphism from $\text{Pol}(\mathbf{C}_k, \mathbf{K}_3)$ to $\mathcal{Z}_{\leq N}$.

Vertex chains

$\mathbf{G} = (V(G), E(G))$. Let $\Delta_V(\mathbf{G})$ denote the free Abelian group with generators $[v], v \in V(G)$.

Edge chains

For an edge (u, v) let $[u, v]$ denote an orientation of the edge from u to v . Let $\Delta_E(\mathbf{G})$ denote the free Abelian group with generators $[u, v], (u, v) \in E(G)$, where $[u, v] = -[v, u]$ for every edge.

Definition

For any graph homomorphism $f: \mathbf{H} \rightarrow \mathbf{G}$ we define group homomorphisms $f_V: \Delta_V(\mathbf{H}) \rightarrow \Delta_V(\mathbf{G})$ and $f_E: \Delta_E(\mathbf{H}) \rightarrow \Delta_E(\mathbf{G})$ defined by

$$f_V(\sum_i c_i[v_i]) = \sum_i c_i[f(v_i)],$$

and

$$f_E(\sum_i c_i[u_i, v_i]) = \sum_i c_i[f(u_i), f(v_i)].$$

Map on a group homomorphisms

Definition 2.15

For a graph \mathbf{G} , we define a map $\partial: \Delta_E(\mathbf{G}) \rightarrow \Delta_V(\mathbf{G})$ as the group homomorphism such that $[u, v] \mapsto [v] - [u]$. for every $[u, v] \in \Delta_E(\mathbf{G})$

Lemma 2.16

For each graph homomorphism $f: \mathbf{H} \rightarrow \mathbf{G}$ and each $P \in \Delta_E(\mathbf{H})$, we have $f_V(\partial P) = \partial f_E(P)$.

Degree of a homomorphism

Lemma 5.1

Let $m, l \geq 3$, and let $f: C_m \rightarrow C_l$ be a homomorphism. Then there is an integer d such that $f_E(O_m) = d \cdot O_l$. Denote this integer d as $\deg f$.

Lemma 5.3

Let $m, l \geq 3$, assume that l is odd, and let $f: C_m \rightarrow C_l$ be a homomorphism. Then

- ① $|\deg f| \leq m/l$,
- ② the parity of $\deg f$ is the same as the parity of m , and
- ③ if $m = 4$ then $\deg f = 0$.

Lemma 5.6

Let $n \geq 2$, $f: \mathbf{C}_k^n \rightarrow \mathbf{C}_3$ be a polymorphism, and let $i \in \{1, \dots, n\}$. Then

- 1 for each edge e of \mathbf{C}_k^{n-1} , there is an integer d such that $f_E(e \times_i O_k) = 2d \cdot O_3$;
- 2 the above d does not depend on the choice of e ;
- 3 $d = \deg_i f$.

Definition

Let \mathcal{L} denote the minion of all linear maps over \mathbb{Z} , i.e., of the functions of the form $\sum c_i x_i$ where all $c_i \in \mathbb{Z}$. Define a mapping $\delta: \text{Pol}(\mathbf{C}_k, \mathbf{C}_3) \rightarrow \mathcal{L}$ by

$$\delta(f): (x_1, \dots, x_n) \mapsto \deg_1 f \cdot x_1 + \dots + \deg_n f \cdot x_n.$$

Lemma 5.7

The map δ is a minion homomorphism.

Lemma 5.7

Let $n \geq 2$. If $f \in \text{Pol}(\mathbf{C}_k, \mathbf{C}_3)$ is n -ary and g is obtained from f by identifying the first two variables, i.e.,

$$g(y, x_3, \dots, x_n) = f(y, y, x_3, \dots, x_n)$$

then $\deg_1 g = \deg_1 f + \deg_2 f$.

Lemma 5.11

Let $f: \mathbf{C}_k^n \rightarrow \mathbf{C}_3$ be a polymorphism, and $i \in \{1, \dots, n\}$. If the coordinate i in f is dummy, then $\deg_i f = 0$.

Lemma 5.12

Let N be the largest odd number such that $N \leq k/3$. Then we have $\delta(f) \in \mathcal{Z}_{\leq N}$ for all $f \in \text{Pol}(\mathbf{C}_k, \mathbf{C}_3)$.