for rats 20 mg/kg diet (1.5 mg/kg daily); for dogs 200 mg/kg (6.6 mg/kg daily). **Toxicity class** WHO III; EPA III.

ECOTOXICOLOGY

Birds LC_{50} (8 d) for Japanese quail > 10 000 mg/kg. Fish LC_{50} (96 h) for rainbow trout 2.2, common carp 37 mg/l. Bees Not toxic to bees; LD_{50} (oral) > 197 µg/bee, (topical) > 101 µg/bee. Daphnia EC_{50} (48 h) 45 mg/l. Other beneficial spp. LC_{50} (14 d) for earthworms 160 mg/kg soil.

ENVIRONMENTAL FATE

Animals Major pathways in animal metabolism are hydrolysis of the methylthio group to form the hydroxy derivative, sulfoxidation of the methylthio group, and side-chain dealkylation. Plants Major pathways in plant metabolism are the hydrolysis of the methylthio group to form the hydroxy derivative, conjugation with glutathione and side-chain dealkylation. Soil and water Degradation involves oxidation of the methylthio group to sulfoxide and sulfone, hydrolysis with the introduction of a 2-hydroxy group, dealkylation at the substituted amino groups, cleavage of the amine moiety, and ring opening. Duration of residual activity in soil is c. 3 months. K_{∞} 150.

200 diafenthiuron

Insecticide Acaricide

$$CH(CH_3)_2$$
 $-O$
 $-NHCSNHC(CH_3)_3$
 $CH(CH_3)_2$

NOMENCLATURE

Common name diafenthiuron (BSI, draft E-ISO).

IUPAC name 1-tert-butyl-3-(2,6-di-isopropyl-4-phenoxyphenyl)thiourea. C.A. name N-[2,6-bis(1-methylethyl)-4-phenoxyphenyl]-N'-(1,1-dimethylethyl)thiourea.

C.A. Hame 175[2,00013(1 Hierary) Production of Production

PHYSICO-CHEMICAL PROPERTIES

Mol. wt. 384.6 Mol. formula $C_{23}H_{32}N_2OS$

Form White powder. M.p. 144.6-147.7 °C (OECD 102) V.p. $< 2 \times 10^{-6}$ Pa (25 °C) (OECD 104) SG/density 1.09 (20 °C) (OECD 109) $K_{ow} \log P = 5.76$ (OECD 107) Solubility In water 0.06 mg/l (25 °C). In ethanol 43, acetone 320, toluene 330, *n*-hexane 9.6, *n*-octanol 26 (all in g/l, 25 °C). Stability Stable in air and water and to light.

COMMERCIALISATION

History Insecticide and acaricide reported by H. P. Streibert et al. (Proc. 1988 Br. Crop Prot. Conf., 1, 25). Introduced by Ciba-Geigy AG. Patents GB 2060626; DE 3034905 Manufacturer Ciba-Geigy.

294 diafenthiuron