

G-Cluster States, and SPT Order

Tycho Van Camp

October 14, 2025

Abstract

We study explicitly a generalization of the cluster state for some finite group G . Using an MPS representation, we study its symmetries and a stabilizing Hamiltonian. We show that the G-cluster state, which displays SPT order, is distinct from the (trivial) symmetric product state through a duality/gauging argument.

1 INTRODUCTION

We study a generalization of symmetry operators $\{O_a\}_a$ which no longer multiply according to the structure of a group, but which multiply with respect to the structure of a fusion category:

$$O_a O_b = \sum_c N_{a,b}^c O_c \quad (1)$$

We have replaced the conventional presentation of a symmetry operator U by O to emphasize the possibility of non-unitarity of the operators O . Note that the fusion of two objects equal a sum of objects, which ultimately leads to the existence of non-invertible objects- being symmetries in our case.

Here, we construct an explicit microscopic model with fusion category SPTO in 1D, generalizing the $\mathbb{Z}_2 \times \mathbb{Z}_2$ Abelian SPTO. Specifically, we replace the qubits by G -valued basis states. When G is non-Abelian the symmetry becomes $G \times \text{Rep}(G)$, where $\text{Rep}(G)$ is the fusion category of finite-dimensional representations of the group G .

2 NOTATIONS AND CONVENTIONS

We follow the notations and conventions of Feiguin et. al.. Denote group elements as $g \in G$. and their inverses as $\bar{g} \in G$. Irreducible representations (irreps) are defined via the map $\Gamma : G \rightarrow GL_{d_\Gamma}(\mathbb{C})$, and respect the group structure of G . Denote the trivial irrep as $\mathbf{1}$. The tensor product of irreps satisfy:

$$\Gamma_i \otimes \Gamma_j \cong \bigoplus_k N_{\Gamma_i, \Gamma_j}^{\Gamma_k} \Gamma_k \quad (2)$$

We represent our states as a MPS, written as:

$$|\psi_A\rangle = \sum_{\{g_i\}} \text{Tr}[BA_{g_1}^{(1)} \cdots A_{g_N}^{(N)}] |g_1, \dots, g_N\rangle \quad (3)$$

wherein A_g^i is the matrix evaluated at site i when $g_i = g$, and B encodes the choice of boundary conditions. Similarly, we write MPOs as:

$$O_a = \sum_{\{g_i, g'_i\}} \text{Tr}[BA_{g_1, g'_1}^{(1)} \cdots A_{g_N, g'_N}^{(N)}] |g_1, \dots, g_N\rangle \langle g'_1, \dots, g'_N| \quad (4)$$

Both the MPSs and the MPOs can be represented diagrammatically using Penrose notation as 3-valent tensors and 4-valent tensors, respectively. Joining the ends of the string of either of these objects amounts to taking the trace. Gluing a bra or a ket to the left or the right, respectively, one can select a particular matrix element.

We label local Hilbert space basis kets by elements of the finite group G , and which follow the group algebra $\mathbb{C}[G]$. An arbitrary single-site state is written as a superposition of these basis kets as $|\psi\rangle = \sum_{g \in G} c_g |g\rangle$ with $c_g \in \mathbb{C}$. The generalization of qubit Pauli operators to group-based Pauli operators are used extensively. The group-based X -type operators are labeled by group elements $g \in G$:

$$\overrightarrow{X}_g = \sum_{h \in G} |gh\rangle \langle h| \iff \overrightarrow{X}_g |h\rangle = |gh\rangle \quad (5)$$

$$\overleftarrow{X}_g = \sum_{h \in G} |h\bar{g}\rangle \langle h| \iff \overleftarrow{X}_g |h\rangle = |h\bar{g}\rangle \quad (6)$$

The left- X -type and the right- X type can be combined into a conjugation operator \overleftrightarrow{X}_g . The group-based Z -type operators are labeled by irreps Γ :

$$Z_\Gamma = \sum_{g \in G} \Gamma(g) \otimes |g\rangle \langle g| \iff Z_\Gamma |g\rangle = \Gamma(g) \otimes |g\rangle \quad (7)$$

$$Z_\Gamma^\dagger = \sum_{g \in G} \Gamma(\bar{g}) \otimes |g\rangle \langle g| \iff Z_\Gamma^\dagger |g\rangle = \Gamma(\bar{g}) \otimes |g\rangle \quad (8)$$

The delta function on the group is given by:

$$\delta_{g,h}^G = \sum_{\Gamma \in \text{Rep}(G)} \frac{d_\Gamma}{|G|} \text{Tr}[\Gamma(\bar{g}h)] \quad (9)$$

Since in general G is non-Abelian, $\Gamma(g_i)$ and $\Gamma(g_j)$ do not necessarily commute which means we must be careful in the ordering of operators. The commutation relations follow:

$$[\overrightarrow{X_g^{(i)}}, \overleftarrow{X_h^{(j)}}] = 0 \quad (10)$$

$$[\overrightarrow{X_g^{(i)}}, \overrightarrow{X_h^{(j)}}] \propto [\overleftarrow{X_g^{(i)}}, \overleftarrow{X_h^{(j)}}] \propto \delta_{i,j} \quad (11)$$

$$\overrightarrow{X_g} Z_\Gamma = \Gamma(\bar{g}) \cdot Z_\Gamma \overrightarrow{X_g} \quad (12)$$

$$\overleftarrow{X_g} Z_\Gamma = Z_\Gamma \cdot \Gamma(g) \overleftarrow{X_g} \quad (13)$$

Swapping of X - and Z -type operators on the same site yields a matrix $\Gamma(h)$ which acts on the virtual space. Similar to the case where we generalized the computational basis to G -valued basis states, we generalize the qubit basis $|+\rangle$ and $|-\rangle$ via basis states labeled by matrix elements of irreps:

$$|\Gamma_{\alpha\beta}\rangle := \sqrt{\frac{d_\Gamma}{|G|}} \sum_{g \in G} [\Gamma(g)]_{\alpha\beta} |g\rangle \quad (14)$$

3 THE GENERALIZED CLUSTER STATE

For a finite group G , there does not exist a natural isomorphism between group elements $g \in G$ and irreps Γ of G . This complicates matters when generalizing the stabilizers, as the ZXZ -cluster state is not a CSS stabilizer code. This caveat is circumvented by using a Hadamard gate on even sites, such that we obtain a state called the CSS-cluster state. This can be seen from the fact that ZXZ -operators are either transformed into ZZZ - or XXX - operators. Alternatively, the CSS-cluster state may also be prepared by acting with a circuit of CX -gates on the product state $|+\rangle|0\rangle|+\rangle\cdots$. This method readily generalizes for the state preparation of the G -cluster state, where one should instead use $|\mathbf{1}\rangle|e\rangle|\mathbf{1}\cdots\rangle$, where $e \in G$ denotes the identity and $\mathbf{1} \in \text{Rep}(G)$ is the trivial irrep. In contrast to the qubit-based cluster state presented as a simple graph, the G -cluster state necessitates a bipartite and directed graph. The bipartiteness stems from the fact that CX -gates (and subsequent generalizations thereof) are not symmetric in their controls and targets, such that we need to assign a particular subset of the lattice as controls. The requirement for being directed is due to the fact that there exist different group-based CX -operators, being $C\overrightarrow{X}^{(i,j)}$ and $C\overleftarrow{X}^{(i,j)}$ in our case:

$$C\overrightarrow{X}^{(i,j)} |g_i, g_j\rangle := |g_i, g_i g_j\rangle \quad C\overleftarrow{X}^{(i,j)} |g_i, g_j\rangle := |g_i, g_j \bar{g}_i\rangle \quad (15)$$

Here, we study a particular group-based cluster state specified by choosing odd-sites as controls, with all edges oriented to the left, henceforth referred to as the G -cluster state. Defining a 1D chain with n sites, each of which have a local Hilbert space $\mathbb{C}[G]$, the G -cluster state reads:

$$|\mathcal{C}\rangle = \mathcal{N} \sum_{\{g_i\} \in G} |g_1\rangle |g_1 \bar{g}_2\rangle |g_2\rangle \cdots |g_{N-1} \bar{g}_N\rangle |g_N\rangle \quad (16)$$

Alternatively, G -cluster state preparation may be implemented by acting on the product state $|\psi_0\rangle := |\mathbf{1}\rangle|e\rangle|\mathbf{1}\rangle\cdots$ with the finite depth circuit $U_{\mathcal{C}}$:

$$U_{\mathcal{C}} = \prod_{i \text{ odd}} C \vec{X}^{(i,i+1)} C \overleftarrow{X}^{(i,i-1)} \quad (17)$$

Regrouping of indices of the tensors defining the G -cluster state allows a graphical representation of the state. Furthermore, these single-site tensors respect a set of "pulling-through" identities, which allow transferring operators on the physical indices to the virtual indices.

A Hamiltonian with ground state $|\mathcal{C}\rangle$ can be constructed as a sum of commuting stabilizers:

$$H_{\mathcal{C}} = -\frac{1}{|G|} \sum_{i \text{ odd}} \left(\sum_{\Gamma \in \text{Rep}(G)} \text{Tr}[Z_{\Gamma}^{\dagger(i)} \cdot Z_{\Gamma}^{(i+1)} Z_{\Gamma}^{(i+2)}] d_{\Gamma} + \sum_{g \in G} \overleftarrow{X}_g^{(i+1)} \vec{X}_g^{(i+2)} \vec{X}_g^{(i+3)} \right) \quad (18)$$

wherein the sum over $\Gamma \in \text{Rep}(G)$ denotes a sum over irreps Γ of G (i.e. simple objects in $\text{Rep}(G)$). The group-based Pauli terms all commute with one another, and the ground-state $|\mathcal{C}\rangle$ is the joint eigenspace of all the operators with maximum eigenvalue ($1/|G|$ for X -type terms, and $d_{\Gamma}^2/|G|$ for the Z -type terms). The cluster-state Hamiltonian has four independent families of global symmetries, which in turn are respected by the ground state $|\mathcal{C}\rangle$:

$$G_R : \vec{A}_g := \prod_{i \text{ odd}} \overleftarrow{X}_g^{(i)}, \quad (19)$$

$$G_L : \vec{A}_g := \prod_{i \text{ odd}} \vec{X}_g^{(i)} \overleftarrow{X}_g^{(i+1)}, \quad (20)$$

$$\text{Rep}(G) : \hat{B}_{\Gamma} := \text{Tr} \left[\prod_{i \text{ even}} Z_{\Gamma}^{(i)} \right] \quad (21)$$

$$\text{Inn}(G) : \hat{C}_g := \prod_{i \text{ even}} \left(\overleftarrow{X}_{g'g\bar{g}'}^{(i)} \delta_{g'}^{(i+1)} \vec{X}_{g'g\bar{g}'}^{(i+2)} \right) \quad (22)$$

We thus have: two G symmetries, a $\text{Rep}(G)$ symmetry, and an $\text{Inn}(G)$ symmetry corresponding to the inner automorphism group of G . For Abelian groups, the two G symmetries are identified and the $\text{Inn}(G)$ trivializes. What remains are the two symmetries, $G \times \text{Rep}(G)$, that protect the SPTO. For the non-Abelian case, it is analytically and numerically proved that $G_R \times \text{Rep}(G)$ is the minimal subgroup also protecting the SPTO. Since \overleftarrow{A}_g and \hat{B}_{Γ} are supported on different sublattices, they commute. Whilst the \overleftarrow{A}_g operators form a representation of a group, the \hat{B}_{Γ} operators multiply according to the fusion rules of $\text{Rep}(G)$:

$$\hat{B}_{\Gamma_i} \hat{B}_{\Gamma_j} = \sum_k N_{\Gamma_i, \Gamma_j}^{\Gamma_k} \hat{B}_{\Gamma_k} \quad (23)$$

We refer to this symmetry as a fusion category symmetry, whilst high-energy nomenclature refers to these as non-invertible symmetries. See Appendix ?? This stems from the fact that the Z_Γ operators in general do not have an inverse. This is again rooted in the fact that an object and its dual, yields a sum of objects including the trivial irrep, implying that the fusion to the trivial object is not the only fusion channel.

If G is Abelian, we have that $\text{Rep}(G) \cong G$ such that the character group is isomorphic to G .

The fusion category symmetry $G \times \text{Rep}(G)$ is labelled by pair (g, Γ) and multiply as:

$$(g_i, \Gamma_i) \otimes (g_j, \Gamma_j) = \bigoplus_k N_{\Gamma_i, \Gamma_j}^{\Gamma_k} (gh, \Gamma_k) \quad (24)$$

From which the \overleftarrow{A}_g and \hat{B}_Γ symmetry operators follow the same multiplication. Notice that the multiplication is non-commutative since $gh \neq hg$, regardless of the fact that $N_{\Gamma_i, \Gamma_j}^{\Gamma_k}$ is nonzero for multiple triples (i, j, k) . This implies that $G \times \text{Rep}(G)$ is not the fusion algebra of irreps of some other finite group G' : $G \times \text{Rep}(G) \neq \text{Rep}(G')$. However, as long as the fusion category is nonanomalous, we can relate $G \times \text{Rep}(G)$ to $\text{Rep}(H)$ for some semisimple Hopf algebra H , such that: $G \times \text{Rep}(G) \cong \text{Rep}(H)$. This is satisfied for $H = \mathbb{C}[G]^* \otimes \mathbb{C}[G]$.

4 DISTINCTNESS FROM THE SYMMETRIC PRODUCT STATE: STACKING VS DUALITY

5 SIGNATURES OF SPT ORDERS: EDGES AND STRINGS

References

- Noninvertible Symmetry-Protected Topological Order in a Group-Based Cluster State; C. Feuchis, N. Tantivasadakarn, V. Albert; 2025; DOI: <https://doi.org/10.1103/PhysRevX.15.011058>

A MULTIPLICATION OF MPOs

Here we explicitly multiply the MPOs of the $\text{Rep}(G)$ symmetry. First we introduce the notation for the fusion of two irreps of the group G , which we refer to as generalized Clebsch-Gordan coefficients. This is presented by a tuning fork type diagram with two incoming legs and one outgoing leg [insert a diagram], which corresponds to the generalized Clebsch-Gordan coefficient:

$$\begin{pmatrix} \Gamma_i & \Gamma_j & | & \Gamma_k & \mu \\ \alpha & \beta & & \gamma & \end{pmatrix} \quad (25)$$

As a reminder we also write out the explicit form of the Z_Γ MPO [insert Z MPO here]:

$$alpha - MPO(Z_\Gamma) - beta = \sum_{g \in G} [\Gamma(g)]_{\alpha\beta} |g\rangle \langle g| \quad (26)$$

This allows the multiplication of two Z_Γ 's, which is represented by so called 'pulling-through'- or 'zipper' diagrams [insert pulling-through/zipper diagram here]:

Explicitely, by choosing basis vectors we can write:

$$\sum_{\epsilon, \delta} [Z_{\Gamma_i}]_{\epsilon\alpha} [Z_{\Gamma_j}]_{\delta\beta} \begin{pmatrix} \Gamma_i & \Gamma_j & | & \Gamma_k & \mu \\ \epsilon & \delta & & \gamma & \end{pmatrix} = \sum_{\Gamma_k} \sum_{\kappa} [Z_{\Gamma_k}]_{\kappa\gamma} \begin{pmatrix} \Gamma_i & \Gamma_j & | & \Gamma_k & \mu \\ \alpha & \beta & & \kappa & \end{pmatrix} \quad (27)$$