On (0+1)d SPT interfaces

Tycho Van Camp

February 2026

Abstract

We study the $(0+1)$d interfaces of symmetry-protected topological (SPT) chains in the tensor network (TN) framework. By analysing the symmetry fractionalization on the entanglement degrees of freedom, we obtain an algebra with possible anomaly. The representations of this algebra possibly possesses measurable fractional statistics,

Introduction The discovery of edge modes in symmetry protected topological (SPT) phases by Haldane [Hal83], and the subsequent explanation by Affleck, Kennedy, Lieb, and Tasaki [AKLT87], has since garnered increasing attention. The boundary of an SPT is particularly important as it serves as the symmetry-protected logical qubit for measurement-base quantum computation (MBQC) [RBB02] [RBB03]. The presence of edge modes in these SPTs [CGLW13] leads to degeneracy in the entanglement spectrum [PTBO10], and can be interpreted as a manifestation of *bulk-boundary correspondence*. In this note we study the so-called *interface algebra*, introduced in [LXY25] and[IO26a], which yields information on both the degeneracy of the interface and the braiding statistics of interface modes [RVHB19] [GRS25] [FTA25]. By leveraging the tensor network (TN) framework [VMC08] [BC17] [CPGSV21], we analyse explicit examples of models on the lattice.

Boundaries of generalized SPT chains

[Note to self: the following concepts are already introduced in the preliminaries sections: - MPO symmetries: first describe how generalized symmetries are presented in the tensor network framework by possible non-local MPOs, the multiplication of MPOs - fusion tensors: fusion tensors are used for the multiplication of MPOs, specifically the multiplication does not have to be group-like. - Action tensors: encode how MPO symmetries act on the entanglement of a state - L symbols: encode the multiplication of symmetry operators on the entanglement of a state, i.e. how does the multiplication of the two action tensors decompose as a sum of action tensors?]

Having introduced the tensor network formalism, specifically the representation of symmetry operators, and states, we now proceed to investigate the $(0+1)$d interface of two adjacent SPTs with generalized symmetry encoded in $\mathcal{C}$. Consider two contigous $\mathcal{C}$-SPTs, SPT_A and SPT_B, with corresponding uniform $\mathcal{C}$-symmetric injective MPSs, A and B. Neither the physical- or the virtual Hilbert spaces of A are necessarily isomorphic to that of B. Label the sites of the infinite chain by $i \in \mathbb{Z}$, with $i < 0$ furnished with A tensors, and with $i > 0$ furnished with B tensors. Graphically, we consider the following situation:



(1)

[Note to self: I have postponed the mention of the action tensors and corresponding L-symbols, since it seems a better 'story'/ 'buildup' of first 'just doing it' and then reintroducing the previously developed 'tools'.]

Consider the action of an MPO $\mathcal{O}_x$ on this interface:



(2)

Recall that action tensors are used for the local application of an MPO $\mathcal{O}_x$ to an MPS [GRLM23]. Given the action tensors, $({}^A\phi_x)_i$ and $({}^B\phi_x)_j$, for each of the SPT phases A and B, this simplifies to:



(3)

The operators on the boundary form what is called an *interface algebra* [LXY25] [IO26a]. In the present case, we have a $(0+1)$d interface algebra. Specifically, when SPT_A occupies the left-half-, and SPT_B occupies the right-half of the infinite chain, we have

$$ {}^{A|B}\mathfrak{I} := \{({}^{A|B}\mathcal{I}_x)_{ij} | x \in \mathcal{C}, i, j = 0, 1, ..., \dim(x)\} \tag{4} $$

The identity operator $\mathcal{O}_\mathbb{I}$ provides the unit element of ${}^{A|B}\mathfrak{I}$, whereas the muliplication is given by:



(5)



(6)



(7)

It was found in [LXY25] [IO26a] that SPT_A and SPT_B belong to the same phase[1] if and only if the interface algebra ${}^{A|B}\mathfrak{I}$ has 1-dimensional representations. More interestingly however, is the case when SPT_A is not in the same phase as SPT_B, as this is the case where degenerate interface modes occur [LXY25] [**?**]. The existence of these degenerate interface modes[2] is recognized as a hallmark signature of SPTs [FTA25]. This can again be traced back to the fractionalization of the symmetry on the entanglement of the state in question [PBTO12]. These interface modes can obey fractional statistics, and a detection procedure has been outlined in [RVHB19] [GRS25]. This is also found in [SS24] for a $\mathbb{Z}_2$-cluster state and more generally in [FTA25] for the G-cluster state. [Note to self: in [LXY25] a identification between interface algebras and funcotrs between two module is made. Does this somehow relate to duality operators?]

Example 1. As a first example, we consider the case where the $\mathbb{Z}_2$-cluster state [SAV11] [Bre15] occupies the left-half of an infinite chain, and the product state occupies the right-half of the infinite chain. Previously, we found that both states have symmetry described by $\mathbf{Vec}_{\mathbb{Z}_2} \boxtimes \mathbf{Rep}(\mathbb{Z}_2) \cong \mathbf{Vec}_{\mathbb{Z}_2} \boxtimes \mathbf{Vec}_{\mathbb{Z}_2}$ [FTA25] [IO26a].
The one-dimensional action tensors of the $\mathbb{Z}_2$-cluster state are: $({}^{\text{cluster}}\phi_g) = V_g$, with V_g given in table 1. The L-symbols evaluate to a normalised representative ω in the non-trivial second cohomology class of $\mathbb{Z}_2 \times \mathbb{Z}_2$ in $U(1)$), denoted

[1] Two systems are said to be in the same phase if their corresponding Hamiltonians can be connected via a smooth gapped path of local and symmetric Hamiltonians [GRLM23].

[2] We distinguish between edge modes when the protecting symmetry is group-based, versus interface modes when the protecting symmetry is a generalized symmetry.

$H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, U(1))$: ${}^{\text{cluster}}L_{gh} = \omega(g,h)$. The action tensors of the product state are trivial. Similarly, the L-symbols of the product state are trivial.

g	V_g
(0,0)	$\mathbb{I}_2\mathbb{I}_2$
(0,1)	$\mathbb{I}_2 Z$
(1,0)	$X\mathbb{I}_2$
(1,1)	XZ

Table 1: Projective representations for the $\mathbb{Z}_2$-cluster state. $\mathbb{I}_2$ is the 2×2 identity matrix, and X, Y, Z are the usual Pauli matrices.

Using the fact that the $\mathbb{Z}_2$-cluster state and the trivial product state are not in the same phase (which should be clear from the inequivalence[3] of L-symbols), and that the interface algebra ${}^{\text{cluster}|\text{triv}}\mathfrak{I}$ has total dimension $|\mathbb{Z}_2 \times \mathbb{Z}_2| = 4$, we conclude that ${}^{\text{cluster}|\text{triv}}\mathfrak{I} \cong \text{Mat}_2(\mathbb{C})$. The algebra of 2×2 matrices over $\mathbb{C}$, $\text{Mat}_2(\mathbb{C})$, is spanned by the usual Pauli matrices: $\{\mathbb{I}_2, \sigma^x, \sigma^y, \sigma^z\}$ [note to self: notice the different notation used for the interface algebra and the action tensors, or the bulk symmetry operators]. Multiplication of operators in the interface algebra ${}^{\text{cluster}|\text{triv}}\mathfrak{I}$ reads:

$$ {}^{\text{cluster}|\text{triv}}\mathcal{I}_g \times {}^{\text{cluster}|\text{triv}}\mathcal{I}_h = \omega(g,h)^{\text{cluster}|\text{triv}}\mathcal{I}_{gh} \tag{8} $$

Specific operators ${}^{\text{cluster}\ |\text{triv}}\mathcal{I}_g$ in the interface algebra ${}^{\text{cluster}|\text{triv}}\mathfrak{I}$ for $g \in \mathbb{Z}_2 \times \mathbb{Z}_2$ are given in table 2.

g	${}^{\text{cluster}\ |\text{triv}}\mathcal{I}_g$
$(0,0)$	$\mathbb{I}_2$
$(0,1)$	σ^z
$(1,0)$	σ^x
$(1,1)$	$i\sigma^y$

Table 2: Representation of interface algebra operators ${}^{\text{cluster}\ |\text{triv}}\mathcal{I}_g$ in ${}^{\text{cluster}|\text{triv}}\mathfrak{I}$ for $g \in \mathbb{Z}_2 \times \mathbb{Z}_2$. The $\sigma^x, \sigma^y, \sigma^z$ are the usual 2×2 Pauli matrices, and $\mathbb{I}_2$ is the 2×2 identity matrix.

Next, we provide an example of a $(0+1)$d interface algebra between SPT phases protected by a non-invertible symmetry. In contrast to the case of invertible SPT phases where the identification of a trivial product state is clear-cut, the notion of a trivial product state for non-invertible SPT phase does not exist [SS24].

[3] We say that two sets of L-symbols are equivalent if they are related by gauge transformations on the action tensors whence they are defined [GRLM23] [IO26a].

Example 2. In this example we consider the interface algebras between different $\mathbf{Rep}(D_4)$ SPT phase where D_4 is the group describing the symmetries of a square. We provide the fusion of irreducible representations of D_4 in table 3.

$\otimes$	A_1	A_2	B_1	B_2	E
A_1	A_1	A_2	B_1	B_2	E
A_2		A_1	B_2	B_1	E
B_1			A_1	A_2	E
B_2				A_1	E
E					$A_1 \oplus A_2 \oplus B_1 \oplus B_2$

Table 3: Fusion rules of irreps of the dihedral group D_4. The empty slots can be found by transposing the table. Notice that all of the fusion channels yield multiplicity-free products.

It is noteworthy to mention that $\mathbf{Rep}(D_4)$ is a group-theoretical fusion 1-category [Nat17], such that explicit lattice models of these phases can be constructed [SS24]. Specifically, these models are obtained by starting from a D_4-symmetric Hamiltonian, for which the ground state (GS) has spontaneous symmetry breaking (SSB) pattern $G \to K$, and subsequently applying (twisted) gauging [HVAS+15] [GRK23] [CGRV+25] of the subgroup $H \subseteq G$, such that K and H provide an exact factorization of G [SS24][IO26b]. Alternatively, starting from the different $\mathbf{Rep}(D_4)$-SPT models and subsequently applying the (twisted) gauging in reverse, one observes different SSB patterns of the resulting SSB models [LXY25]. It is know that $\mathbf{Rep}(D_4)$ admits 3 distinct non-invertible SPT phases[4] [Tam00]. We will label the distinct $\mathbf{Rep}(D_4)$-SPT phases by $\mathrm{SPT}_{\mathsf{A}}$, $\mathrm{SPT}_{\mathsf{B}}$, and $\mathrm{SPT}_{\mathsf{C}}$. [Note to self: it might be interesting to write down all three SPT models in TN framework following [SS24], and subsequently extracting the action tensors. For now, we'll trust [IO26a].] The L-symbols for the $\mathbf{Rep}(G)$-SPT phases boil down to (twisted) generalized Clebsch-Gordan coefficients[5] (GCGCs) [IO26a]. We list the nontrivial ${}^{\mathsf{A}}L$-symbols for $\mathrm{SPT}_{\mathsf{A}}$ in table 4.

For the other two SPT phases, $\mathrm{SPT}_{\mathsf{B}}$ and $\mathrm{SPT}_{\mathsf{C}}$, we have a *twisted* copy of the ${}^{\mathsf{A}}L$-symbols of $\mathrm{SPT}_{\mathsf{A}}$, wherein another one of the L-symbols regarding the fusion of a one-dimensional irrep (A_i or B_i) and the two-dimensional irrep (E), changes sign when interchanging the reactants. The ${}^{\mathsf{B}}L$-symbols for $\mathrm{SPT}_{\mathsf{B}}$ are provided in table 5, whilst the ${}^{\mathsf{C}}L$-symbols for $\mathrm{SPT}_{\mathsf{C}}$ are provided in table 6.

With the L-symbols at hand, we can now investigate the interface algebras between these non-invertible SPTs. Let L denote the SPT phase occupying the left-half of the infinite chain, and R denote the right-half of the infinite chain.

[4] Distinct SPT phases with symmetry encoded in the fusion 1-category $\mathcal{C}$ are provided by non-isomorphic fiber 1-functors [TW19].

[5] Generalized Clebsch-Gordan coefficients of small groups can be found in [KDWS63], and a formula for computing these for groups enjoying multiplicity-free fusion of unitary irreducible representation is given in [Aro26]. A numerical algorithm applicable to compact groups can be found in [ILYM17].

$({}^{\mathsf{A}}L^{E}_{EA_1})^{k}_{1j} = ({}^{\mathsf{A}}L^{E}_{A_1E})^{k}_{1j} = \left[\begin{smallmatrix} A_1 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\mathbb{I}_2)_{jk}$	$({}^{\mathsf{A}}L^{A_1}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} A_1 \\ 1 \end{smallmatrix}\right] = (\mathbb{I}_2)_{ij}$
$-({}^{\mathsf{A}}L^{E}_{EA_2})^{k}_{1j} = ({}^{\mathsf{A}}L^{E}_{A_2E})^{k}_{1j} = \left[\begin{smallmatrix} A_2 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (-\mathsf{i}\sigma^y)_{jk}$	$({}^{\mathsf{A}}L^{A_2}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} A_2 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\mathsf{i}\sigma^y)_{ij}$
$({}^{\mathsf{A}}L^{E}_{EB_1})^{k}_{1j} = ({}^{\mathsf{A}}L^{E}_{B_1E})^{k}_{1j} = \left[\begin{smallmatrix} B_1 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\sigma^z)_{jk}$	$({}^{\mathsf{A}}L^{B_1}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} B_1 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\sigma^z)_{ij}$
$({}^{\mathsf{A}}L^{E}_{EB_2})^{k}_{1j} = ({}^{\mathsf{A}}L^{E}_{B_2E})^{k}_{1j} = \left[\begin{smallmatrix} B_2 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\sigma^x)_{jk}$	$({}^{\mathsf{A}}L^{B_2}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} B_2 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\sigma^x)_{ij}$

Table 4: Nontrivial ${}^{\mathsf{A}}L$-symbols for the $\mathbf{Rep}(D_4)$-SPT phase $\mathrm{SPT}_{\mathsf{A}}$ [KDWS63] [IO26a]. $\sigma^x, \sigma^y, \sigma^z$ are the usual 2×2 Pauli matrices. Note that $({}^{\mathsf{B}}L^{E}_{A_2E})^{k}_{1j} = -({}^{\mathsf{B}}L^{E}_{EA_2})^{k}_{1j}$. [Note to self: We identify $\mathrm{SPT}_{\mathsf{A}}$ with the ν_3 SPT phase in [IO26a]][Note to self: although we have table overflow, this isn't really disturbing in my opinion.]

$({}^{\mathsf{B}}L^{E}_{EA_1})^{k}_{1j} = ({}^{\mathsf{B}}L^{E}_{A_1E})^{k}_{1j} = \left[\begin{smallmatrix} A_1 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\mathbb{I}_2)_{jk}$	$({}^{\mathsf{B}}L^{A_1}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} A_1 \\ 1 \end{smallmatrix}\right] = (\sigma^x)_{ij}$
$({}^{\mathsf{B}}L^{E}_{EA_2})^{k}_{1j} = ({}^{\mathsf{B}}L^{E}_{A_2E})^{k}_{1j} = \left[\begin{smallmatrix} A_2 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (-\mathsf{i}\sigma^y)_{jk}$	$({}^{\mathsf{B}}L^{A_2}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} A_2 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\sigma^z)_{ij}$
$-({}^{\mathsf{B}}L^{E}_{EB_1})^{k}_{1j} = ({}^{\mathsf{B}}L^{E}_{B_1E})^{k}_{1j} = \left[\begin{smallmatrix} B_1 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\sigma^z)_{jk}$	$({}^{\mathsf{B}}L^{B_1}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} B_1 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\mathsf{i}\sigma^y)_{ij}$
$({}^{\mathsf{B}}L^{E}_{EB_2})^{k}_{1j} = ({}^{\mathsf{B}}L^{E}_{B_2E})^{k}_{1j} = \left[\begin{smallmatrix} B_2 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\sigma^x)_{jk}$	$({}^{\mathsf{B}}L^{B_2}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} B_2 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\mathbb{I}_2)_{ij}$

Table 5: Nontrivial ${}^{\mathsf{B}}L$-symbols for the $\mathbf{Rep}(D_4)$-SPT phase $\mathrm{SPT}_{\mathsf{B}}$ [IO26a]. $\sigma^x, \sigma^y, \sigma^z$ are the usual 2×2 Pauli matrices. Note that $({}^{\mathsf{B}}L^{E}_{B_1E})^{k}_{1j} = -({}^{\mathsf{B}}L^{E}_{EB_1})^{k}_{1j}$. [Note to self: We identify $\mathrm{SPT}_{\mathsf{B}}$ with the ν_1 SPT phase in [IO26a]]

The eight-dimensional interface algebra ${}^{\mathsf{L}|\mathsf{R}}\mathfrak{I}$ is spanned by:

$$\{{}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C_i}, ({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_E)_{ij} |\ \ C = A, B, \ \ i, j = 1, 2\}. \tag{9}$$

When L is not in the same SPT phase as R, we again use the result from [IO26a] which determines that there exist no one-dimensional representations of ${}^{\mathsf{L}|\mathsf{R}}\mathfrak{I}$, to conclude that it must be the case that ${}^{\mathsf{L}|\mathsf{R}}\mathfrak{I} \cong \mathrm{Mat}_2(\mathbb{C}) \oplus \mathrm{Mat}_2(\mathbb{C})$. This also implies a four-fold degeneracy at the interface between different $\mathbf{Rep}(D_4)$-SPTs. Multiplication of operators in ${}^{\mathsf{L}|\mathsf{R}}\mathfrak{I}$ is given by

$({}^{\mathsf{C}}L^{E}_{EA_1})^{k}_{1j} = ({}^{\mathsf{C}}L^{E}_{A_1E})^{k}_{1j} = \left[\begin{smallmatrix} A_1 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\mathbb{I}_2)_{jk}$	$({}^{\mathsf{C}}L^{A_1}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} A_1 \\ 1 \end{smallmatrix}\right] = (\sigma^z)_{ij}$
$({}^{\mathsf{C}}L^{E}_{EA_2})^{k}_{1j} = ({}^{\mathsf{C}}L^{E}_{A_2E})^{k}_{1j} = \left[\begin{smallmatrix} A_2 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (-\mathsf{i}\sigma^y)_{jk}$	$({}^{\mathsf{C}}L^{A_2}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} A_2 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(-\sigma^x)_{ij}$
$({}^{\mathsf{C}}L^{E}_{EB_1})^{k}_{1j} = ({}^{\mathsf{C}}L^{E}_{B_1E})^{k}_{1j} = \left[\begin{smallmatrix} B_1 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\sigma^z)_{jk}$	$({}^{\mathsf{C}}L^{B_1}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} B_1 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(\mathbb{I}_2)_{ij}$
$-({}^{\mathsf{C}}L^{E}_{EB_2})^{k}_{1j} = ({}^{\mathsf{C}}L^{E}_{B_2E})^{k}_{1j} = \left[\begin{smallmatrix} B_2 & E \\ 1 & j \end{smallmatrix}\middle\vert\begin{smallmatrix} E \\ k \end{smallmatrix}\right] = (\sigma^x)_{jk}$	$({}^{\mathsf{C}}L^{B_2}_{EE})^{1}_{ij} = \left[\begin{smallmatrix} E & E \\ i & j \end{smallmatrix}\middle\vert\begin{smallmatrix} B_2 \\ 1 \end{smallmatrix}\right] = \frac{1}{\sqrt{2}}(-\mathsf{i}\sigma^y)_{ij}$

Table 6: Nontrivial ${}^{\mathsf{C}}L$-symbols for the $\mathbf{Rep}(D_4)$-SPT phase $\mathrm{SPT}_{\mathsf{C}}$ [IO26a]. $\sigma^x, \sigma^y, \sigma^z$ are the usual 2×2 Pauli matrices. Note that $({}^{\mathsf{C}}L^{E}_{B_2E})^{k}_{1j} = -({}^{\mathsf{C}}L^{E}_{EB_2})^{k}_{1j}$. [Note to self: We identify $\mathrm{SPT}_{\mathsf{C}}$ with the ν_2 SPT phase in [IO26a]]

$$ {}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C_i} \times {}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C_j} = {}^{\mathsf{L}}L_{C_iC_j}\,{}^{\mathsf{R}}\bar{L}_{C_iC_j}\,{}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C_iC_j} \tag{10} $$

$$ {}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C} \times ({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{D})_{ij} = \sum_{kk'} ({}^{\mathsf{L}}L_{CD})_{ik}({}^{\mathsf{R}}\bar{L}_{CD})_{k'j}({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{D})_{kk'} \tag{11} $$

$$ ({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{D})_{ij} \times {}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C} = \sum_{kk'} ({}^{\mathsf{L}}L_{DC})_{ik}({}^{\mathsf{R}}\bar{L}_{DC})_{k'j}({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{D})_{kk'} \tag{12} $$

$$ ({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{D})_{ij} \times ({}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{D})_{kl} = \sum_{C_i} ({}^{\mathsf{L}}L^{C_i}_{DD})_{ik}({}^{\mathsf{R}}\bar{L}^{C_i}_{DD})_{jl}\,{}^{\mathsf{L}|\mathsf{R}}\mathcal{I}_{C_i} \tag{13} $$

Explicit multiplication of operators in ${}^{\mathsf{L}|\mathsf{R}}\mathfrak{I}$ is obtained upon substition of relevant L-symbols provided in tables 4, 5, and 6. Importantly, as can already be seen from the fact that some of the L-symbols are the usual Pauli matrices, some of the operators in ${}^{\mathsf{L}|\mathsf{R}}\mathfrak{I}$ anticommute- a manifestation of an anomaly at the boundary[6]

Summary We have introduced the notion of $(0+1)$d interface algebras that arise when considering two contiguous $(1+1)$d SPT phases. Specifically, we have mobilized the TN framework in which the interpretation of the ideas covered here become (unreasonably) simple. Examples of $(0+1)$d interface algebras for both invertible- and non-invertible SPT phases were provided. It is observed that the interface carries an anomaly when the adjacent SPT phases are distinct, and that the interface modes obey fractional statistics.

References

[AKLT87] Ian Affleck, Tom Kennedy, Elliott H. Lieb, and Hal Tasaki. Rigorous results on valence-bond ground states in antiferromagnets.

[6] In the literature this is also referred to as *anomaly inflow* [TW19] [TvK15] or *bulk-boundary correspondence* [IO26a]

Phys. Rev. Lett., 59:799–802, Aug 1987.

[Aro26] Daniel P. Arovas. Lecture notes on group theory in physics (a work in progress). Department of Physics, University of California, San Diego, 2026.

[BC17] Jacob C Bridgeman and Christopher T Chubb. Hand-waving and interpretive dance: an introductory course on tensor networks. *Journal of Physics A: Mathematical and Theoretical*, 50(22):223001, May 2017.

[Bre15] Courtney G Brell. Generalized cluster states based on finite groups. *New Journal of Physics*, 17(2):023029, February 2015.

[CGLW13] Xie Chen, Zheng-Cheng Gu, Zheng-Xin Liu, and Xiao-Gang Wen. Symmetry protected topological orders in interacting bosonic systems, 2013.

[CGRV+25] Bram Vancraeynest-De Cuiper, José Garre-Rubio, Frank Verstraete, Kevin Vervoort, Dominic J. Williamson, and Laurens Lootens. From gauging to duality in one-dimensional quantum lattice models, 2025.

[CPGSV21] J. Ignacio Cirac, David Pérez-García, Norbert Schuch, and Frank Verstraete. Matrix product states and projected entangled pair states: Concepts, symmetries, theorems. *Reviews of Modern Physics*, 93(4), December 2021.

[FTA25] Christopher Fechisin, Nathanan Tantivasadakarn, and Victor V. Albert. Noninvertible symmetry-protected topological order in a group-based cluster state. *Physical Review X*, 15(1), March 2025.

[GRK23] José Garre-Rubio and Ilya Kull. Gauging quantum states with nonanomalous matrix product operator symmetries. *Physical Review B*, 107(7), February 2023.

[GRLM23] José Garre-Rubio, Laurens Lootens, and András Molnár. Classifying phases protected by matrix product operator symmetries using matrix product states. *Quantum*, 7:927, February 2023.

[GRS25] José Garre-Rubio and Norbert Schuch. Fractional domain wall statistics in spin chains with anomalous symmetries. *SciPost Physics*, 18(2), February 2025.

[Hal83] F. D. M. Haldane. Nonlinear field theory of large-spin heisenberg antiferromagnets: Semiclassically quantized solitons of the one-dimensional easy-axis néel state. *Phys. Rev. Lett.*, 50:1153–1156, Apr 1983.

[HVAS+15] Jutho Haegeman, Karel Van Acoleyen, Norbert Schuch, J. Ignacio Cirac, and Frank Verstraete. Gauging quantum states: From global to local symmetries in many-body systems. *Physical Review X*, 5(1), February 2015.

[ILYM17] A. Ibort, A. López Yela, and J. Moro. A new algorithm for computing branching rules and clebsch–gordan coefficients of unitary representations of compact groups. *Journal of Mathematical Physics*, 58(10), October 2017.

[IO26a] Kansei Inamura and Shuhei Ohyama. 1+1d spt phases with fusion category symmetry: interface modes and non-abelian thouless pump, 2026.

[IO26b] Kansei Inamura and Shuhei Ohyama. Generalized cluster states in 2+1d: non-invertible symmetries, interfaces, and parameterized families, 2026.

[KDWS63] George F. Koster, John O. Dimmock, Robert G. Wheeler, and Hermann Statz. *Properties of the Thirty-Two Point Groups*, volume 24 of *M.I.T. Press Research Monograph*. M.I.T. Press, Cambridge, Massachusetts, 1963.

[LXY25] Da-Chuan Lu, Fu Xu, and Yi-Zhuang You. Strange correlator and string order parameter for non-invertible symmetry protected topological phases in 1+1d, 2025.

[Nat17] Sonia Natale. On the equivalence of module categories over a group-theoretical fusion category. *Symmetry, Integrability and Geometry: Methods and Applications*, June 2017.

[PBTO12] Frank Pollmann, Erez Berg, Ari M. Turner, and Masaki Oshikawa. Symmetry protection of topological phases in one-dimensional quantum spin systems. *Physical Review B*, 85(7), February 2012.

[PTBO10] Frank Pollmann, Ari M. Turner, Erez Berg, and Masaki Oshikawa. Entanglement spectrum of a topological phase in one dimension. *Physical Review B*, 81(6), February 2010.

[RBB02] Robert Raussendorf, Daniel Browne, and Hans Briegel. The one-way quantum computer–a non-network model of quantum computation. *Journal of Modern Optics*, 49(8):1299–1306, July 2002.

[RBB03] Robert Raussendorf, Daniel E. Browne, and Hans J. Briegel. Measurement-based quantum computation on cluster states. *Physical Review A*, 68(2), August 2003.

[RVHB19] Gertian Roose, Laurens Vanderstraeten, Jutho Haegeman, and Nick Bultinck. Anomalous domain wall condensation in a modified ising chain. *Physical Review B*, 99(19), May 2019.

[SAV11] W. Son, L. Amico, and V. Vedral. Topological order in 1d cluster state protected by symmetry. *Quantum Information Processing*, 11(6):1961–1968, December 2011.

[SS24] Sahand Seifnashri and Shu-Heng Shao. Cluster state as a non-invertible symmetry-protected topological phase. *Physical Review Letters*, 133(11), September 2024.

[Tam00] Daisuke Tambara. Representations of tensor categories with fusion rules of self-duality for abelian groups. *Israel Journal of Mathematics*, 118(1):29–60, 2000.

[TvK15] Ryan Thorngren and Curt von Keyserlingk. Higher spt's and a generalization of anomaly in-flow, 2015.

[TW19] Ryan Thorngren and Yifan Wang. Fusion category symmetry i: Anomaly in-flow and gapped phases, 2019.

[VMC08] F. Verstraete, V. Murg, and J.I. Cirac. Matrix product states, projected entangled pair states, and variational renormalization group methods for quantum spin systems. *Advances in Physics*, 57(2):143–224, March 2008.