The Exponential Triad

Three Constrained Decompositions of Nested Exponential Constants

Lord Soumadeep Ghosh with ChatGPT (OpenAI)

Kolkata, India

Abstract

This treatise studies the equation

$$1 + e + e^e = e^{x+y+z}, \tag{1}$$

under equal weighting, ascending exponential weights, and descending exponential weights. Taking logarithms reduces the nonlinear equation to the affine constraint

$$x + y + z = S, \qquad S := \ln(1 + e + e^e). \tag{2}$$

A general normalized-weight theorem then yields unique closed-form solutions whenever the three positive weights are fixed. The equal-weight solution is $S/3$ in each coordinate; the two exponential-weighted solutions are reversals of one another. The treatise proves existence and uniqueness, derives sensitivity formulas, proposes numerical verification procedures, and examines carefully delimited analogies to economics, financial planning, econometrics, law, physics, chemistry, biology, linguistics, psychology, psychiatry, philosophy, religion, political science, international relations, welfare economics, warfare economics, statistics, computer science, artificial intelligence, and cybersecurity.

The central methodological distinction is between a mathematical theorem and an application analogy. The algebraic results are exact; applications to empirical or normative fields require independently justified variables, units, data, assumptions, and validation. The equation by itself does not establish a scientific law, a financial recommendation, a legal rule, a clinical instrument, or a geopolitical prediction.

The treatise ends with "The End"

Contents

1 Introduction and scope **2**
- 1.1 The motivating equation . . . 3

2 Mathematical preliminaries **3**
- 2.1 Exponentials, logarithms, and equivalence . . . 3
- 2.2 Why one equation is insufficient . . . 3
- 2.3 A general normalized-weight theorem . . . 3

3 The three constrained decompositions **4**
- 3.1 Decomposition I: Equal weighting . . . 4
- 3.2 Decomposition II: Increasing exponential weighting . . . 4
- 3.3 Decomposition III: Decreasing exponential weighting . . . 4
- 3.4 Comparison and reversal symmetry . . . 5

4 Geometric interpretation **5**

5 Logic, proof obligations, and mathematical interpretation **6**
- 5.1 Existence, uniqueness, and domain . . . 6
- 5.2 What the equation does not prove . . . 6

6 Sensitivity analysis and uncertainty propagation **6**

7 Economics and financial planning and analysis **7**

7.1 An allocation template . . . 7
7.2 Opportunity cost and the limits of proportional allocation . . . 7
7.3 Welfare economics and distribution . . . 7

8 Econometrics and statistical identification 7

9 Optimization, statistics, and risk analysis 8

10 Physics, chemistry, and biology 8
10.1 Physics: dimensionless exponents . . . 8
10.2 Chemistry: a structural analogy, not kinetics . . . 8
10.3 Biology: growth and population models . . . 9

11 Linguistics, psychology, and psychiatry 9
11.1 Linguistics . . . 9
11.2 Psychology and choice architecture . . . 9
11.3 Psychiatry and clinical limits . . . 9

12 Philosophy, religion, and the logic of analogy 9
12.1 Mathematical truth and interpretation . . . 9
12.2 Ethical reflection and religious traditions . . . 9

13 Law, political science, diplomacy, and international relations 10
13.1 Law and proportionality . . . 10
13.2 Political science and institutional priorities . . . 10
13.3 Diplomacy and international relations . . . 10
13.4 Geopolitics, warfare economics, and warfare . . . 10

14 Computing, data science, and AI/ML 10
14.1 Algorithmic specification . . . 10
14.2 Numerical stability . . . 11
14.3 Reference implementation and unit tests . . . 11
14.4 AI and machine learning . . . 11
14.5 Cybersecurity and model governance . . . 11

15 A reusable generalized theorem 12

16 Limitations and research agenda 12

17 Conclusion 12

List of Figures

1 Schematic simplex picture. The drawing illustrates the normalized-weight geometry, not the exact plotted coordinates of the three numerical solutions. . . . 5

List of Tables

1 The three constrained solutions. Decimal values are rounded; the closed forms above are exact. . . . 5
2 Illustrative model-governance questions for a weighted decomposition. . . . 8

1 Introduction and scope

The expression $1 + e + e^e$ combines a constant, the base of the natural logarithm, and an iterated exponential. The equation (1) asks for three real numbers whose sum, after exponentiation, reproduces this positive constant. By itself, one equation in three unknowns does not determine a unique triple. A supplementary constraint is therefore not a technical afterthought: it is the information that selects a point from a two-dimensional set of possibilities.

The exposition follows four rules. First, every algebraic claim is stated with its assumptions. Second, exact formulas are distinguished from decimal approximations. Third, a model's interpretation is kept separate from its algebra. Fourth, cross-disciplinary connections are presented as possible modelling templates rather than claims that the same equation already explains each field. This distinction is consistent with the philosophical literature on analogical reasoning, which treats analogies as guides to inquiry rather than as deductive proofs of target-domain claims [22].

1.1 The motivating equation

Because the right-hand side of (1) is positive and the left-hand side is positive, taking the natural logarithm is legitimate. The logarithm is the inverse of the exponential function on the positive reals [1, 2, 3]. Thus (1) is equivalent over real x, y, z to (2). The quantity S is the total to be decomposed.

$$S = \ln(1 + e + e^e) \approx 2.937708171169375. \tag{3}$$

The important point is that S is fixed while its decomposition is not. We must add a second layer of structure to distinguish one triple from another.

2 Mathematical preliminaries

2.1 Exponentials, logarithms, and equivalence

For every real number u, $e^u > 0$; conversely, every positive real number has a unique natural logarithm. Hence

$$\begin{aligned} e^{x+y+z} = 1 + e + e^e &\iff \ln\big(e^{x+y+z}\big) = \ln(1 + e + e^e) && (4)\\ &\iff x + y + z = \ln(1 + e + e^e) && (5)\\ &\iff x + y + z = S. && (6) \end{aligned}$$

This is an equivalence, not merely a one-way implication. It establishes that solving the original equation over the reals is exactly the same as decomposing the scalar S into three real summands.

2.2 Why one equation is insufficient

The solution set of $x + y + z = S$ is a plane in three-dimensional real space. Choosing arbitrary real x and y gives

$$z = S - x - y. \tag{7}$$

Consequently, infinitely many triples solve the original equation. A unique solution appears only after adding enough independent restrictions. The three constraints studied here are alternative restrictions; they are not imposed simultaneously.

2.3 A general normalized-weight theorem

Theorem 1 (normalized-weight decomposition). Let $S \in \mathbb{R}$ and let w_1, w_2, w_3 be fixed positive real numbers. The system

$$\begin{aligned} x + y + z &= S, && (8)\\ x : y : z &= w_1 : w_2 : w_3 && (9) \end{aligned}$$

has exactly one real solution, namely

$$x = S\frac{w_1}{W}, \qquad y = S\frac{w_2}{W}, \qquad z = S\frac{w_3}{W}, \tag{10}$$

where $W = w_1 + w_2 + w_3 > 0$.

Proof. The ratio constraint means that there exists a scalar c with $x = cw_1$, $y = cw_2$, and $z = cw_3$. Substituting into the total gives $cW = S$. Since $W > 0$, it follows that $c = S/W$, yielding (10). This value of c is unique, so the triple is unique. Substitution also verifies that the three coordinates sum to S. □

The theorem is a simple linear-algebra result. Its usefulness is organisational: it separates the total S from the rule governing its allocation. The proportions w_i/W sum to one and can therefore be interpreted as normalized weights. This is related to the general use of normalized nonnegative coefficients in optimization, although no optimization objective is needed for the present theorem [4].

3 The three constrained decompositions

3.1 Decomposition I: Equal weighting

Impose the constraint

$$x = y = z. \tag{11}$$

Writing all three coordinates as c, the total equation becomes $3c = S$. Therefore,

$$\boxed{x = y = z = \frac{1}{3}\ln(1 + e + e^e)}. \tag{12}$$

Numerically,

$$x = y = z \approx 0.979236057056458. \tag{13}$$

This is the unique equal-weight allocation. Equality of the coordinates is an imposed symmetry condition; it is not a consequence of the original equation alone.

3.2 Decomposition II: Increasing exponential weighting

Impose

$$x = \frac{y}{e} = \frac{z}{e^2}. \tag{14}$$

The condition is equivalent to $y = ex$ and $z = e^2x$, so the ratio is

$$x : y : z = 1 : e : e^2. \tag{15}$$

Substitution into $x + y + z = S$ yields

$$x(1 + e + e^2) = S, \tag{16}$$

$$x = \frac{S}{1 + e + e^2}. \tag{17}$$

The complete solution is

$$x_\uparrow = \frac{\ln(1 + e + e^e)}{1 + e + e^2}, \tag{18}$$

$$y_\uparrow = \frac{e\ln(1 + e + e^e)}{1 + e + e^2}, \tag{19}$$

$$z_\uparrow = \frac{e^2\ln(1 + e + e^e)}{1 + e + e^2}. \tag{20}$$

The arrows are labels indicating the ascending sequence of weights; they do not denote a time derivative. The numerical values are

$$x_\uparrow \approx 0.264483550457689, \tag{21}$$

$$y_\uparrow \approx 0.718940829135467, \tag{22}$$

$$z_\uparrow \approx 1.954283791576219. \tag{23}$$

3.3 Decomposition III: Decreasing exponential weighting

Now impose the different condition

$$x = ey = e^2z. \tag{24}$$

It follows that $y = x/e$ and $z = x/e^2$; hence

$$x : y : z = e^2 : e : 1. \tag{25}$$

The sum condition gives

$$x\left(1+\frac{1}{e}+\frac{1}{e^2}\right)=S. \tag{26}$$

Multiplying numerator and denominator by e^2 gives the closed-form solution

$$x_{\downarrow}=\frac{e^2\ln(1+e+e^e)}{e^2+e+1}, \tag{27}$$

$$y_{\downarrow}=\frac{e\ln(1+e+e^e)}{e^2+e+1}, \tag{28}$$

$$z_{\downarrow}=\frac{\ln(1+e+e^e)}{e^2+e+1}. \tag{29}$$

Numerically,

$$x_{\downarrow}\approx 1.954283791576219, \tag{30}$$

$$y_{\downarrow}\approx 0.718940829135467, \tag{31}$$

$$z_{\downarrow}\approx 0.264483550457689. \tag{32}$$

3.4 Comparison and reversal symmetry

Table 1: The three constrained solutions. Decimal values are rounded; the closed forms above are exact.

Case	Constraint	Ratio $x:y:z$	Approximate (x,y,z)
I	$x=y=z$	$1:1:1$	(0.979236, 0.979236, 0.979236)
II	$x=y/e=z/e^2$	$1:e:e^2$	(0.264484, 0.718941, 1.954284)
III	$x=ey=e^2z$	$e^2:e:1$	(1.954284, 0.718941, 0.264484)

Cases II and III are reversals. In particular,

$$x_{\uparrow}=z_{\downarrow}, \tag{33}$$

$$y_{\uparrow}=y_{\downarrow}, \tag{34}$$

$$z_{\uparrow}=x_{\downarrow}. \tag{35}$$

This follows because the weights in Case III are the reverse of those in Case II while their normalizing denominator, $1+e+e^2$, is unchanged. The two solutions are therefore related by the coordinate permutation $(x,y,z)\mapsto(z,y,x)$.

4 Geometric interpretation

The unrestricted solution set $x+y+z=S$ is a plane. If nonnegativity is also imposed, its intersection with the nonnegative orthant is a triangle, the two-dimensional simplex scaled by S. Each positive weighting selects a single point on that triangle. The three points correspond to the equal-weight centre and two points reflected across the plane $x=z$ within the triangle.




Figure 1: Schematic simplex picture. The drawing illustrates the normalized-weight geometry, not the exact plotted coordinates of the three numerical solutions.

The barycentric coordinates of a point in this simplex are precisely the normalized weights $(w_1/W, w_2/W, w_3/W)$. This provides a concise geometric meaning for the theorem: the total S fixes the size of the simplex, while the constraint fixes a location within it.

5 Logic, proof obligations, and mathematical interpretation

The reasoning uses the following chain:

1. Positivity makes the logarithm of the left-hand side well-defined.
2. Injectivity of the exponential and logarithm transforms the original equation into a sum constraint.
3. Each additional condition specifies a positive ratio of coordinates.
4. The normalization theorem turns that ratio into a unique triple.
5. Substitution checks the total and the ratio independently.

The proof does not depend on numerical approximation. Decimal calculations are verification aids, not premises. This distinction is central to mathematical practice: a numerical result can reveal an arithmetic error but does not replace a proof of the exact claim [1, 2].

5.1 Existence, uniqueness, and domain

For all three cases, the weights are strictly positive and $S > 0$. The denominator in each solution is therefore positive. The solution exists and is unique over the reals. In addition, each coordinate is positive, and the values sum to S. If the variables were allowed to be complex, the interpretation of the original equation and the inverse logarithm would require additional care because the complex logarithm is multivalued. This treatise confines the main theorem to real variables.

5.2 What the equation does not prove

The equation alone does not imply the equal-weight condition, either exponential ratio, or any empirical theory. Nor does the use of e establish a connection to a particular natural process. A mathematical structure can be reused in another field only after a defensible mapping from the mathematical objects to the target domain is supplied. Analogical reasoning may suggest hypotheses, but those hypotheses require independent testing [22].

6 Sensitivity analysis and uncertainty propagation

The general solution can be written compactly as

$$a_i = q_i S, \qquad q_i := \frac{w_i}{W}, \qquad \sum_{i=1}^{3} q_i = 1. \tag{36}$$

For fixed weights, each allocation changes linearly with the total. Therefore,

$$\frac{\partial a_i}{\partial S} = q_i. \tag{37}$$

A one-unit change in S is distributed across the three components in their normalized proportions. This is a sensitivity statement, not a claim that the underlying constant is uncertain.

To make input sensitivity explicit, let

$$S(a, b, c) = \ln(a + b + c), \qquad a, b, c > 0. \tag{38}$$

Then

$$\frac{\partial S}{\partial a} = \frac{1}{a + b + c}, \tag{39}$$

$$\frac{\partial S}{\partial b} = \frac{1}{a + b + c}, \tag{40}$$

$$\frac{\partial S}{\partial c} = \frac{1}{a + b + c}. \tag{41}$$

For fixed weights, $a_i = q_i S(a, b, c)$, so

$$\frac{\partial a_i}{\partial a} = \frac{\partial a_i}{\partial b} = \frac{\partial a_i}{\partial c} = \frac{q_i}{a + b + c}. \tag{42}$$

At the current values $(a, b, c) = (1, e, e^e)$, this derivative gives a first-order response to perturbations in the three summands. It should not be mistaken for a causal interpretation of those constants.

If an empirical process supplied an estimate $\widehat{S}$ with variance σ_S^2, then under fixed weights the delta-method variance and covariance propagation would be

$$\mathrm{Var}(\widehat{a}_i) = q_i^2 \sigma_S^2, \tag{43}$$

$$\mathrm{Cov}(\widehat{a}_i, \widehat{a}_j) = q_i q_j \sigma_S^2. \tag{44}$$

These relations are conditional on uncertainty being present in S. In the purely mathematical problem, all inputs are exact definitions and no sampling variance exists.

7 Economics and financial planning and analysis

7.1 An allocation template

The weights can be interpreted abstractly as priority coefficients and S as a fixed budget, resource index, or bookkeeping total. The normalized solution allocates that total in fixed proportions. In financial planning and analysis (FP&A), the template could serve as a simple scenario-allocation worksheet when the analyst has already justified the total and weights. It does not, by itself, determine a firm's budget, cost of capital, risk tolerance, liquidity requirement, or investment policy. Those inputs require accounting definitions, forecasts, constraints, and governance [6].

An economically meaningful implementation must check dimensional consistency. Here S is dimensionless. A dollar budget cannot equal this mathematical number without a separate scale parameter and a mapping from the model to currency units. In practice one might introduce a monetary total B and define $A_i = Bq_i$, but this is a new model rather than a conclusion of (1).

7.2 Opportunity cost and the limits of proportional allocation

Proportional allocation assumes that the weights are appropriate and that the same percentage rule should be used across components. It does not incorporate marginal returns, uncertainty, correlation, taxes, transaction costs, liquidity constraints, or intertemporal discounting. Portfolio theory, for example, studies trade-offs between expected return and risk, not merely normalized weights [5]. A financial analyst should compare a simple allocation baseline against a model that represents the actual objective and constraints.

7.3 Welfare economics and distribution

The three decompositions illustrate distinct distribution rules: equal shares, ascending priority weights, and descending priority weights. They do not establish which rule is socially optimal. A welfare comparison requires a social objective, a description of whose welfare counts, distributional information, and constraints. Social choice theory demonstrates that aggregating individual preferences raises issues that cannot be resolved by arithmetic normalization alone [8, 9]. Equality of allocations, equality of welfare, and fairness are distinct concepts.

8 Econometrics and statistical identification

The distinction between a constraint and an estimated relationship is essential. The equations here provide a deterministic mapping from S and the weights to (x, y, z); they do not estimate a parameter from observed data. A single observation of the total $x + y + z$ cannot identify three free components. The constraints supply the missing structure by assumption.

For an empirical analogue, suppose an analyst observes component outcomes a_{it} across units i and periods t, and proposes

$$a_{it} = q_i S_t + \varepsilon_{it}, \tag{45}$$

where q_i is a prespecified allocation share, S_t is a period-specific total, and ε_{it} is a residual. This is a proposed statistical model, not a consequence of the original identity. Estimation would require a sampling design, a definition of the observed variables, assumptions about errors, and appropriate diagnostics. Econometric inference depends on identification and assumptions rather than on algebraic resemblance alone [7].

Possible diagnostics include checking whether components add to the recorded total, testing whether estimated shares are stable across time, checking residual dependence, and comparing the constrained model with an unconstrained benchmark. A better in-sample fit would not automatically justify a causal interpretation or reliable out-of-sample predictions.

9 Optimization, statistics, and risk analysis

The fixed-weight problem can be viewed as a feasible-set selection: among all triples with a fixed total, the ratio restriction selects one point. It is not necessary to define an objective function. If an objective were added, the solution might differ; for example, minimizing a quadratic loss subject to the total constraint can yield different shares depending on the loss coefficients. This is why an allocation rule should not be confused with an optimality theorem [4].

For risk analysis, the three cases can be treated as structural scenarios. The equal-weight case is a neutral benchmark; the two weighted cases test how a reversed ordering of priorities changes the components while preserving the total. Scenario analysis is useful for revealing dependence on assumptions, but scenario selection is not probabilistic forecasting. Probabilities must be supplied or estimated separately.

Table 2: Illustrative model-governance questions for a weighted decomposition.

Question	Required check
Total consistency	Verify that the components sum to the specified total within numerical tolerance.
Constraint consistency	Verify that ratios match the declared weights.
Sensitivity	Vary totals and weights and record the resulting allocations.
Interpretability	Define what each variable means, including units and direction of preference.
Validation	Compare model outputs against an independent benchmark or observed data.
Governance	Record assumptions, version, decision owner, and conditions under which the model must not be used.

10 Physics, chemistry, and biology

10.1 Physics: dimensionless exponents

In physical equations, the argument of an exponential must be dimensionless. The present variables appear in e^{x+y+z}, so the sum $x+y+z$ is dimensionless; the individual variables are naturally treated as dimensionless as well unless a scale has been introduced explicitly. This is a useful dimensional-analysis check, not evidence that the equation describes a physical system. Standard physics texts stress the role of units, dimensions, and consistency in formulating quantitative laws [12].

An exponential response occurs in many physical settings, but the presence of an exponential alone does not identify the governing mechanism. A physical application must supply the state variables, parameters, boundary conditions, and empirical or theoretical derivation.

10.2 Chemistry: a structural analogy, not kinetics

Chemical kinetics and temperature-dependent rate models often use exponential functions. For example, the Arrhenius equation involves an exponential of the ratio between activation energy and thermal energy [13, 14]. That exponent is dimensionless because the units in the numerator and denominator cancel. The present expression has no activation energy, temperature, rate constant, or reaction pathway; it should therefore not be presented as an Arrhenius model or as a chemical law.

10.3 Biology: growth and population models

Exponential growth models describe processes in which the instantaneous rate of increase is proportional to the quantity present under specified assumptions. The IUPAC definition describes growth of this kind as proportional to the amount already present [15]. The triad equation does not include time, a population state, a growth rate, births, deaths, or resource limitations. Its relevance is methodological: it shows how an aggregate scalar can be decomposed under explicit allocation constraints. Any biological model would need its own dynamics and data [16].

11 Linguistics, psychology, and psychiatry

11.1 Linguistics

A linguist could use normalized weights as a bookkeeping device for allocating a corpus statistic among three predefined categories. Such a use would require a reproducible annotation scheme and operational definitions. Language frequency distributions, semantic interpretation, and linguistic change involve more than a three-component identity; observed patterns such as word-frequency regularities must be studied with corpus evidence and explicit assumptions [17, 18]. No claim about language follows from (1) alone.

11.2 Psychology and choice architecture

As a decision-making illustration, the three cases can represent three alternative ways of presenting priorities: equal weighting, front-loaded weighting, and reverse weighting. This can motivate an experiment about framing or preference elicitation, but it does not predict human behaviour. A behavioural study would need a preregistered hypothesis, a suitable sample, validated measures, ethical review where required, and statistical analysis. Psychological judgments can be affected by many contextual factors, and a mathematical allocation rule is not a substitute for observed choices [19].

11.3 Psychiatry and clinical limits

There is no clinical, diagnostic, or therapeutic content in this equation. The variables are abstract real numbers, not symptoms, diagnoses, biomarkers, or treatment outcomes. They cannot be used to infer any person's mental state or psychiatric condition. Clinical instruments require evidence of validity, reliability, appropriate use, and professional interpretation; diagnostic frameworks such as DSM-5-TR have a distinct clinical purpose and are not reducible to this algebraic problem [20].

12 Philosophy, religion, and the logic of analogy

12.1 Mathematical truth and interpretation

The proof establishes a conditional mathematical statement: if the equation and a specified constraint hold over the stated domain, then the displayed solution follows. Interpretation requires additional premises. This is a useful example of the distinction between formal validity and the truth of a model's empirical premises. The philosophy of analogy likewise warns that similarity in one respect does not prove similarity in every relevant respect [22].

12.2 Ethical reflection and religious traditions

The equal and weighted decompositions can be used as neutral prompts for discussing equality, priority, stewardship, or obligation. They do not demonstrate a religious doctrine, settle theological questions, or establish which distribution is morally right. Religious traditions contain diverse interpretive, historical, and ethical arguments; a mathematical ratio cannot stand in for that interpretive work [21]. The appropriate conclusion is modest: formal models can clarify the consequences of a declared rule, while moral justification remains a separate task.

13 Law, political science, diplomacy, and international relations

13.1 Law and proportionality

A legal decision-maker may use quantitative models as evidence or as administrative aids only within the applicable legal framework. The fact that a weighted rule is mathematically consistent does not make it lawful, procedurally fair, or proportionate. Legal interpretation also depends on jurisdiction, authoritative texts, precedent where applicable, due process, and reasoned application. The concept of a legal rule is not equivalent to a numerical constraint [10].

13.2 Political science and institutional priorities

The weights can serve as a stylized representation of three institutional priorities. Reversing the weights reverses the allocation while preserving the aggregate. This is a transparent comparative-static result, but it does not predict electoral choices, bargaining outcomes, institutional behaviour, or distributional coalitions. Those claims need a political mechanism and evidence.

13.3 Diplomacy and international relations

In a diplomatic simulation, the weight vector might represent the negotiated priority given to three policy areas, subject to a fixed index. The normalized solution would make the consequences of a stated prioritization rule easy to audit. Actual diplomacy also involves bargaining, commitments, information asymmetry, credibility, domestic politics, and legal obligations. International relations theories offer competing accounts of these processes, none of which follows solely from the triad identity [23, 24].

13.4 Geopolitics, warfare economics, and warfare

Resource-allocation arithmetic can support high-level budgeting and scenario comparison, but it cannot determine force requirements, predict conflict, or establish a lawful course of action. Any real defence or security application would require validated data, operational context, civilian-harm analysis, oversight, and compliance with applicable law. The Charter of the United Nations establishes, among other principles, obligations concerning peaceful settlement and restrictions on the threat or use of force in international relations [11]. Historical work on state formation and coercion also illustrates that the relationship between fiscal resources and military power is institutional and historical, not a universal scalar identity [25].

The proper role of the triad in these domains is therefore limited to transparent arithmetic: holding a declared total fixed, one can show how alternative weights redistribute that total. No operational military recommendation or geopolitical prediction is derived here.

14 Computer science, algorithms, data science, and AI/ML

14.1 Algorithmic specification

For positive weights, the deterministic algorithm has constant-time complexity with respect to the number of coordinates when the dimension is fixed at three. The corresponding general n-component implementation uses standard linear-time summation and allocation passes [26]. A direct implementation follows the proof:

```
INPUT: total S, weights w1, w2, w3
REQUIRE: S is finite; each weight is finite and positive
W = w1 + w2 + w3
x = S * w1 / W
y = S * w2 / W
z = S * w3 / W
RETURN x, y, z
```

For a general number n of components, summing the weights takes $O(n)$ time and the allocation pass takes $O(n)$ time; storage is $O(n)$ if the vector is retained, or $O(1)$ additional storage if emitted as a stream.

14.2 Numerical stability

The given constants are modest, so direct evaluation is safe in ordinary floating-point arithmetic. In a generalized problem with very large positive summands, computing $\ln(a+b+c)$ directly may overflow before the logarithm is applied. The log-sum-exp identity can avoid this problem. For real u_i, define $m = \max_i u_i$; then

$$\ln\left(\sum_{i=1}^{n} e^{u_i}\right) = m + \ln\left(\sum_{i=1}^{n} e^{u_i - m}\right). \tag{46}$$

Here $1 + e + e^e = e^0 + e^1 + e^e$, so S can be computed as the log-sum-exp of $(0, 1, e)$. Stable numerical methods are part of reliable scientific computation; they do not change the exact symbolic result.

14.3 Reference implementation and unit tests

A minimal implementation can calculate the three cases and assert the defining properties. The following language-neutral pseudocode emphasizes checks rather than a particular programming language:

```
S = log(1 + exp(1) + exp(exp(1)))
near(a, b) = abs(a - b) < tolerance

case_equal = (S/3, S/3, S/3)
W = 1 + exp(1) + exp(2)
case_increasing = (S/W, exp(1)*S/W, exp(2)*S/W)
case_decreasing = (exp(2)*S/W, exp(1)*S/W, S/W)

for case in [case_equal, case_increasing, case_decreasing]:
assert near(sum(case), S)

assert near(case_increasing[0], case_decreasing[2])
assert near(case_increasing[1], case_decreasing[1])
assert near(case_increasing[2], case_decreasing[0])
```

A production implementation should validate finite inputs, document its precision, and choose a tolerance appropriate to the numeric representation. Tests should check both the sum and the declared ratio. Unit tests verify the implementation relative to the specification; they do not prove that the specification is appropriate for an external use case.

14.4 AI and machine learning

The normalized-weight rule could be a deterministic baseline in a larger system, or a feature transformation where domain experts have justified the weights. It is not, by itself, a trained machine-learning model: there is no training set, loss function, learned parameter, validation sample, or deployment context. If embedded in an AI system, the rule should be documented, tested against the intended use, and monitored for failure modes. Risk-management frameworks emphasize context, measurement, governance, and ongoing management rather than treating a formula as self-validating [27, 28, 29, 31].

14.5 Cybersecurity and model governance

From a software-security perspective, the core calculation is simple, but surrounding systems can still fail through malformed input, unsafe serialization, dependency risk, logging of sensitive information, or inappropriate access controls. Risk assessment should cover the system boundary and the consequences of misuse, not just the arithmetic routine. NIST SP 800-30 provides a structured risk-assessment framework for information systems and organizations [30]. No security guarantee follows from the mathematical correctness of this function.

15 A reusable generalized theorem

The same result extends to n components. Let $S \in \mathbb{R}$, and let $w_i > 0$ for $i = 1, \ldots, n$. Under the conditions

$$\sum_{i=1}^{n} a_i = S, \tag{47}$$

$$a_1 : a_2 : \cdots : a_n = w_1 : w_2 : \cdots : w_n, \tag{48}$$

there is exactly one solution,

$$a_i = S \frac{w_i}{\sum_{j=1}^{n} w_j}, \qquad i = 1, \ldots, n. \tag{49}$$

The proof is identical: write $a_i = cw_i$, sum, solve for c, and substitute. The fixed-dimensional triad is the case $n = 3$. This extension is a mathematical statement about positive weights, not an empirical theorem about any particular class of institutions or natural systems.

16 Limitations and research agenda

The present result is deliberately small and exact. Several research directions are possible, but each introduces new assumptions:

1. **Endogenous weights:** allow the weights to depend on observed variables or on the components themselves, then study existence and uniqueness.
2. **Uncertain total:** treat S as an estimator or random variable and derive uncertainty intervals under explicit distributional assumptions.
3. **Optimization:** add a stated objective and constraints, then compare the resulting optimum with the proportional solution.
4. **Empirical validation:** define a real target domain, collect data, predefine benchmarks, and test out-of-sample performance.
5. **Normative assessment:** state the social or institutional objective and evaluate distributional consequences separately from algebraic feasibility.

None of these extensions follows automatically from the original equation. Each is a potential next step, not a result claimed in this treatise.

17 Conclusion

The equation $1 + e + e^e = e^{x+y+z}$ is equivalent over the reals to the single linear constraint $x + y + z = S$, where $S = \ln(1 + e + e^e)$. The constraint alone leaves infinitely many solutions. Equal weighting, increasing exponential weighting, and decreasing exponential weighting each select one unique triple. The weighted solutions are exact reversals, and all three preserve the same total.

The most general result is the normalized-weight theorem: a fixed total and fixed positive ratios determine one and only one allocation. That theorem is useful as an algebraic and computational template. Its application to economics, science, law, public policy, or AI depends on independently specifying what the variables mean and validating the model in context. The mathematical accomplishment is the derivation and verification of the decompositions; the broader interdisciplinary value is a disciplined framework for distinguishing an exact result from the hypotheses one might build around it.

References

[1] T. M. Apostol, *Calculus, Volume 1: One-Variable Calculus, with an Introduction to Linear Algebra*, 2nd ed., Wiley, New York, 1967.

[2] W. Rudin, *Principles of Mathematical Analysis*, 3rd ed., McGraw–Hill, New York, 1976.

[3] NIST Digital Library of Mathematical Functions, National Institute of Standards and Technology, *DLMF*, especially the chapters on algebraic and analytic methods and elementary functions, version 1.2.8, 2026.

[4] S. Boyd and L. Vandenberghe, *Convex Optimization*, Cambridge University Press, Cambridge, 2004.

[5] H. Markowitz, "Portfolio Selection," *The Journal of Finance*, vol. 7, no. 1, pp. 77–91, 1952.

[6] R. A. Brealey, S. C. Myers, and F. Allen, *Principles of Corporate Finance*, 12th ed., McGraw–Hill Education, New York, 2017.

[7] J. M. Wooldridge, *Introductory Econometrics: A Modern Approach*, 7th ed., Cengage, Boston, 2020.

[8] K. J. Arrow, *Social Choice and Individual Values*, 2nd ed., Yale University Press, New Haven, 1963.

[9] A. K. Sen, *Collective Choice and Social Welfare*, Holden-Day, San Francisco, 1970.

[10] H. L. A. Hart, *The Concept of Law*, 3rd ed., with a postscript edited by P. A. Bulloch and J. Raz, Oxford University Press, Oxford, 2012.

[11] United Nations, *Charter of the United Nations*, signed 26 June 1945, entered into force 24 October 1945.

[12] D. Halliday, R. Resnick, and J. Walker, *Fundamentals of Physics*, 10th ed., Wiley, Hoboken, 2014.

[13] P. Atkins and J. de Paula, *Physical Chemistry*, 10th ed., Oxford University Press, Oxford, 2014.

[14] International Union of Pure and Applied Chemistry, "Arrhenius Equation," *Compendium of Chemical Terminology (the Gold Book)*, 5th ed., online version 5.0.0, 2025, DOI: 10.1351/goldbook.A00446.

[15] International Union of Pure and Applied Chemistry, "Exponential Growth," *Compendium of Chemical Terminology (the Gold Book)*, 5th ed., online version 5.0.0, 2025, DOI: 10.1351/goldbook.14672.

[16] N. J. Gotelli, *A Primer of Ecology*, 4th ed., Sinauer Associates, Sunderland, MA, 2008.

[17] F. de Saussure, *Course in General Linguistics*, translated by W. Baskin, Philosophical Library, New York, 1959; originally published in French in 1916.

[18] G. K. Zipf, *Human Behavior and the Principle of Least Effort*, Addison-Wesley, Cambridge, MA, 1949.

[19] D. Kahneman, *Thinking, Fast and Slow*, Farrar, Straus and Giroux, New York, 2011.

[20] American Psychiatric Association, *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed., text revision (DSM-5-TR), American Psychiatric Association Publishing, Washington, DC, 2022.

[21] N. Smart, *The World's Religions*, 2nd ed., Cambridge University Press, Cambridge, 1998.

[22] P. Bartha, "Analogy and Analogical Reasoning," in *The Stanford Encyclopedia of Philosophy*, substantive revision 2019.

[23] K. N. Waltz, *Theory of International Politics*, Addison-Wesley, Reading, MA, 1979.

[24] T. C. Schelling, *Arms and Influence*, Yale University Press, New Haven, 1966.

[25] C. Tilly, *Coercion, Capital, and European States, AD 990–1990*, Basil Blackwell, Oxford, 1990.

[26] T. H. Cormen, C. E. Leiserson, R. L. Rivest, and C. Stein, *Introduction to Algorithms*, 4th ed., MIT Press, Cambridge, MA, 2022.

[27] C. M. Bishop, *Pattern Recognition and Machine Learning*, Springer, New York, 2006.

[28] T. Hastie, R. Tibshirani, and J. Friedman, *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*, 2nd ed., Springer, New York, 2009.

[29] I. Goodfellow, Y. Bengio, and A. Courville, *Deep Learning*, MIT Press, Cambridge, MA, 2016.

[30] Joint Task Force Transformation Initiative, *Guide for Conducting Risk Assessments*, NIST Special Publication 800-30, Revision 1, National Institute of Standards and Technology, Gaithersburg, MD, 2012.

[31] National Institute of Standards and Technology, *Artificial Intelligence Risk Management Framework (AI RMF 1.0)*, NIST AI 100-1, Gaithersburg, MD, 2023.

Glossary

Allocation share
The normalized weight $q_i = w_i/W$ assigned to component i.

Constraint
A condition restricting the set of feasible triples, such as $x = y = z$ or a specified ratio.

Decomposition
A representation of the total S as a sum of components.

Dimensionless quantity
A quantity with no physical units; exponential arguments must be dimensionless.

Identification
The conditions under which a model's unknown quantities can be uniquely determined from its assumptions and data.

Log-sum-exp
The identity used to compute the logarithm of a sum of exponentials in a numerically stable way.

Normalized weights
Positive weights divided by their sum, so that the resulting shares sum to one.

Proportional allocation
An allocation in which each component is a fixed share of the total.

Reversal symmetry
The transformation swapping the first and third coordinates while leaving the middle coordinate unchanged.

Simplex
The set of nonnegative component vectors whose coordinates sum to a fixed total.

Structural assumption
A restriction imposed by a model rather than estimated directly from the data.

Uniqueness
The property that exactly one solution satisfies all stated conditions.

The End