

Further Explorations of the $\Psi(s)$ Function and Its Applications

Lord Soumadeep Ghosh et al. with ChatGPT (OpenAI)

Kolkata, India

Abstract

This paper develops further analytical and computational consequences of the transcendental function $\Psi(s)$ introduced through the geometry of intersecting exponential curves associated with a regular pentagon. The central object is the inverse function of a scalar parametrization

$$S(W) = \ln(c - \sigma W e^W) + (c - 1)W + \sigma e^{-W}, \quad (1)$$

where

$$c = \cos \frac{2\pi}{5}, \quad \sigma = \sin \frac{2\pi}{5}. \quad (2)$$

Thus $\Psi(s) = S^{-1}(s)$. Earlier work established the geometric construction, monotonicity, inverse-function interpretation, fixed-point formulation, Newton iteration, and large- s asymptotics of Ψ . The present paper develops a concise approximation architecture combining Padé approximation, logarithmic-rational approximation, asymptotic expansion, Newton refinement, and residual-based error certification.

The mathematical construction is then situated within a broader interdisciplinary framework involving numerical analysis, statistics, econometrics, economics, finance, physics, chemistry, biology, psychology, philosophy, religion, political science, geopolitics, diplomacy, international relations, welfare economics, warfare economics, computer science, data science, and artificial intelligence. These applications are interpreted as mathematical analogies or potential modelling directions rather than as claims that the present scalar function has already been empirically validated in all such domains.

The paper ends with “The End”

1 Introduction

The function $\Psi(s)$ arises from a geometric construction involving exponential curves associated with a regular pentagon. The original construction reduces a geometric intersection problem to a scalar equation and defines Ψ as the inverse of the resulting monotone parametrization [1].

The fundamental relation is

$$s = \ln(c - \sigma \Psi e^\Psi) + (c - 1)\Psi + \sigma e^{-\Psi}. \quad (3)$$

The important mathematical point is that the equation contains both Ψe^Ψ and $e^{-\Psi}$ together with a logarithm. Consequently, ordinary Lambert- W transformations provide useful partial reductions but do not immediately produce an elementary closed form [1, 2].

The objective here is therefore not to force an elementary expression. Instead, we construct an explicit and numerically certifiable representation of Ψ .

2 The Exact Inverse Problem

Define

$$S(W) = \ln(c - \sigma W e^W) + (c - 1)W + \sigma e^{-W}. \quad (4)$$

The function of interest is

$$\Psi(s) = S^{-1}(s). \quad (5)$$

For the pentagonal case,

$$c = \frac{\sqrt{5} - 1}{4}, \quad \sigma = \sin \frac{2\pi}{5}. \quad (6)$$

The derivative is

$$S'(W) = -\frac{\sigma e^W(1+W)}{c - \sigma W e^W} + (c-1) - \sigma e^{-W}. \quad (7)$$

The second derivative is

$$S''(W) = -\frac{\sigma e^W(W+2)}{c - \sigma W e^W} - \frac{\sigma^2 e^{2W}(1+W)^2}{(c - \sigma W e^W)^2} + \sigma e^{-W}. \quad (8)$$

The original construction establishes the relevant monotonicity and therefore the uniqueness of the inverse on the intended branch [1].

3 A Useful Global Derivative Bound

For the relevant branch, write

$$x = -W > 0. \quad (9)$$

Then

$$c - \sigma W e^W = c + \sigma x e^{-x} > c. \quad (10)$$

A direct examination of the two regions $0 < x \leq 1$ and $x > 1$ gives the useful lower bound

$$|S'(W)| > 1 - c + \sigma. \quad (11)$$

Numerically,

$$1 - c + \sigma = 1.642039521920206\dots \quad (12)$$

This inequality is especially useful because it converts an equation residual directly into an error bound.

Let $w(s)$ be any approximation to $\Psi(s)$ and define

$$R(s) = S(w(s)) - s. \quad (13)$$

By the mean value theorem,

$$|w(s) - \Psi(s)| \leq \frac{|R(s)|}{1 - c + \sigma}. \quad (14)$$

Hence a small residual is itself a mathematical certificate of proximity to the exact inverse.

4 Asymptotic Structure

The large- s behavior obtained in the original work is

$$\Psi(s) \sim -\ln \frac{s}{\sigma}. \quad (15)$$

Let

$$L = \ln \frac{s}{\sigma}. \quad (16)$$

A refined expansion is

$$\Psi(s) = -L + \frac{a}{s} + \frac{b}{s^2} + \frac{d}{s^3} + O\left(\frac{(\ln s)^4}{s^4}\right), \quad (17)$$

where

$$a = (1 - c)L + \ln c, \quad (18)$$

$$b = \frac{a^2}{2} + (c - 1)a + \frac{\sigma^2}{c}L, \quad (19)$$

$$d = -\frac{\sigma^4 L^2}{2c^2} + \frac{\sigma^2 La}{c} - \frac{a^3}{6} + ab - \frac{\sigma^2 a}{c} + (c - 1)b. \quad (20)$$

Thus an explicit third-order asymptotic approximation is

$$W_{\text{AS3}}(s) = -L + \frac{a}{s} + \frac{b}{s^2} + \frac{d}{s^3}. \quad (21)$$

The existence of an asymptotic expansion is important computationally: the special function can be evaluated without solving the defining transcendental equation from scratch for every large value of s [1, 3].

5 Local Padé Approximation

Near $s = 1$, put

$$x = s - 1. \quad (22)$$

The local expansion obtained numerically is

$$\begin{aligned} \Psi(s) = & -0.3563813483622252 - 0.3528906435696236x - 0.0276989605880154x^2 \\ & + 0.0221737726077293x^3 - 0.0000909192373878x^4 - 0.0026394065658282x^5 + \dots \end{aligned} \quad (23)$$

A corresponding Padé approximation is

$$P_{5,5}(s) = \frac{p_0 + p_1x + p_2x^2 + p_3x^3 + p_4x^4 + p_5x^5}{1 + q_1x + q_2x^2 + q_3x^3 + q_4x^4 + q_5x^5}, \quad (24)$$

where

$$p_0 = -0.3563813483622252, \quad p_1 = -0.4578445565692827, \quad (25)$$

$$p_2 = -0.2265116389938571, \quad p_3 = -0.09268654807694569, \quad (26)$$

$$p_4 = -0.01770577172118884, \quad p_5 = -0.001914385696193794, \quad (27)$$

and

$$q_1 = 0.2944988941817016, \quad q_2 = 0.2662506737335640, \quad (28)$$

$$q_3 = 0.03576403724665172, \quad q_4 = 0.01164298574205669, \quad (29)$$

$$q_5 = 0.0001477575131530266. \quad (30)$$

The coefficients are numerical approximations. A publication-grade error bound should therefore be established using interval arithmetic rather than inferred solely from floating-point sampling.

6 Logarithmic-Rational Approximation

For intermediate values of s , define

$$u = \frac{1}{s}, \quad L = \ln \frac{s}{\sigma}. \quad (31)$$

A useful explicit approximation is

$$W_{\text{LR}}(s) = -L + \frac{A(u)L + B(u)}{s}, \quad (32)$$

where

$$A(u) = \frac{0.6909830056 + 4.30027266u + 7.20412554u^2 + 0.54484507u^3}{1 - 0.09672448u + 2.84142873u^2 + 0.75879977u^3}, \quad (33)$$

and

$$B(u) = \frac{-1.174359 - 6.71993505u + 3.13047005u^2 + 2.74554685u^3}{1 - 0.09672448u + 2.84142873u^2 + 0.75879977u^3}. \quad (34)$$

For $2 \leq s$, the denominator satisfies

$$1 - 0.09672448u + 2.84142873u^2 + 0.75879977u^3 > 0.95163776 \quad (35)$$

whenever $u \leq 1/2$.

Thus this approximation has no denominator singularity on its intended domain.

7 Piecewise Approximation

Numerical comparison of the residuals

$$R_{\text{LR}}(s) = S(W_{\text{LR}}(s)) - s \quad (36)$$

and

$$R_{\text{AS3}}(s) = S(W_{\text{AS3}}(s)) - s \quad (37)$$

gives the approximate transition

$$s_* \approx 132.5772467247115. \quad (38)$$

The resulting initial approximation is

$$W_0(s) = \begin{cases} P_{5,5}(s), & 1 \leq s \leq 2, \\ W_{\text{LR}}(s), & 2 < s < s_*, \\ W_{\text{AS3}}(s), & s \geq s_*. \end{cases} \quad (39)$$

The construction is summarized schematically in Figure 1.




Figure 1: Piecewise approximation, Newton refinement, and residual certification.

8 Newton Refinement

For any initial approximation $W_0(s)$ define

$$W_1(s) = W_0(s) - \frac{S(W_0(s)) - s}{S'(W_0(s))}. \quad (40)$$

Because $S(\Psi(s)) = s$, Taylor's theorem gives the familiar quadratic Newton error structure

$$|W_1 - \Psi| \leq \frac{M}{2m} |W_0 - \Psi|^2, \quad (41)$$

where

$$m = \inf |S'|, \quad M = \sup |S''| \quad (42)$$

over the relevant local interval.

An especially useful a posteriori certificate avoids the need for a global bound on S'' :

$$E(s) = \frac{|S(W_1(s)) - s|}{1 - c + \sigma}. \quad (43)$$

Consequently,

$$|\Psi(s) - W_1(s)| \leq E(s). \quad (44)$$

This produces a self-certifying numerical evaluator.

9 Numerical Evidence and Rigorous Certification

High-precision numerical experiments indicate an initial residual envelope of approximately

$$\max_{s \geq 1} |S(W_0(s)) - s| \approx 3.24 \times 10^{-4}. \quad (45)$$

After one Newton correction, the observed residual decreases to approximately

$$\max_{s \geq 1} |S(W_1(s)) - s| \approx 4.0 \times 10^{-10}. \quad (46)$$

Using the derivative bound gives the corresponding numerical error estimate

$$|\Psi(s) - W_1(s)| \lesssim 2.44 \times 10^{-10}. \quad (47)$$

These figures are numerical evidence rather than a formal global interval proof. A publication-grade certificate requires outward-rounded interval arithmetic over a finite covering of the bounded intervals and an analytic tail estimate for the unbounded asymptotic region.

The distinction is important:

$$\text{numerical agreement} \neq \text{formal global error proof}. \quad (48)$$

10 Numerical Analysis and Computer Science

The construction can be interpreted as a validated numerical special function.

10.1 Algorithm

$$\text{PSIEVALUATE}(s, \varepsilon) \quad (49)$$

may be represented by the following pseudocode.

```

INPUT: s >= 1, tolerance epsilon

IF 1 <= s <= 2:
    w <- Pade55(s)
ELSE IF 2 < s < s_star:
    w <- LogRational(s)
ELSE:
    w <- Asymptotic3(s)

REPEAT:
    r <- S(w) - s
    dp <- S_prime(w)
    w <- w - r/dp
    E <- abs(S(w) - s)/(1-c+sigma)
UNTIL E <= epsilon

RETURN w, E

```

This algorithm illustrates an important distinction between a symbolic closed form and a computable special function. A function can possess a well-defined mathematical identity while requiring numerical inversion, just as many classical special functions are evaluated algorithmically rather than through elementary expressions [4, 5].

11 Statistics and Data Science

The residual

$$R(s) = S(\widehat{\Psi}(s)) - s \quad (50)$$

can be interpreted as a deterministic model discrepancy.

For a numerical experiment with observations

$$\{(s_i, \widehat{\Psi}_i)\}_{i=1}^n, \quad (51)$$

one can construct residual diagnostics such as

$$\overline{R} = \frac{1}{n} \sum_{i=1}^n R_i, \quad (52)$$

$$\text{RMSE} = \left(\frac{1}{n} \sum_{i=1}^n R_i^2 \right)^{1/2}, \quad (53)$$

$$R_{\max} = \max_i |R_i|. \quad (54)$$

These are useful for empirical validation but should not be confused with mathematical error bounds. Statistics estimates properties of sampled computations; interval analysis can certify properties over continuous domains.

The same distinction is fundamental in data science: training error, test error, approximation error, numerical error, and formal verification are different objects [6, 8].

12 Economics and Finance

The scalar equation

$$s = S(W) \quad (55)$$

can be interpreted abstractly as an equilibrium mapping between an observable or control variable s and a latent state W .

An inverse function therefore provides a latent-state recovery mechanism:

$$W = \Psi(s). \quad (56)$$

This structure is analogous to inverse problems in economics, finance, and econometrics, where an observed equilibrium quantity is used to recover an unobserved structural parameter.

For example, an abstract economic state equation can be written as

$$y_t = F(\theta_t, x_t), \quad (57)$$

with an inverse representation

$$\theta_t = F^{-1}(y_t, x_t). \quad (58)$$

The analogy suggests possible applications in nonlinear equilibrium inversion, latent-variable models, asset-pricing systems, monetary-policy models, and macroeconomic regime identification. It does not establish that Ψ is itself an empirically estimated economic law.

The same inverse structure is relevant to the author's broader work on Ghosh's M Measure and monetary-policy systems [11, 12, 13, 14].

13 Econometrics and Identification

A useful econometric interpretation is obtained from

$$s_i = S(W_i) + \varepsilon_i. \quad (59)$$

If S is strictly monotonic, the deterministic model is identified through

$$W_i = \Psi(s_i) \quad (60)$$

when $\varepsilon_i = 0$.

With stochastic disturbances, however, the inverse estimator becomes

$$\widehat{W}_i = \Psi(s_i - \widehat{\varepsilon}_i), \quad (61)$$

and uncertainty must be propagated through the nonlinear derivative

$$\Psi'(s) = \frac{1}{S'(\Psi(s))}. \quad (62)$$

Thus

$$|\Psi'(s)| < \frac{1}{1 - c + \sigma}. \quad (63)$$

The derivative bound consequently has an econometric interpretation as a global Lipschitz-type stability bound for the inverse mapping.

14 Physics and Other Natural Sciences

The mathematical structure is not restricted to economics.

In physics, inverse functions of nonlinear equilibrium maps arise in thermodynamics, statistical mechanics, nonlinear optics, transport problems, and dynamical systems. The abstract structure

$$\text{observable} = F(\text{latent state}) \quad (64)$$

followed by

$$\text{latent state} = F^{-1}(\text{observable}) \quad (65)$$

is common throughout mathematical physics.

In chemistry, analogous inverse mappings occur when experimentally observed quantities are used to recover equilibrium parameters, reaction coordinates, or nonlinear constitutive parameters.

In biology, inverse nonlinear maps arise in parameter identification, population dynamics, biochemical networks, and systems biology.

These are structural analogies rather than empirical applications of Ψ .

15 Psychology and Psychiatry

The inverse-map perspective can also be used metaphorically in psychology and psychiatry.

An observed behavioral variable can be represented abstractly as

$$y = F(z, \eta), \quad (66)$$

where z denotes a latent state and η denotes contextual variables.

An inverse model seeks

$$z = F^{-1}(y, \eta). \quad (67)$$

The mathematical lesson from Ψ is that inverse identification can be well behaved when monotonicity and derivative bounds are available, while non-identification can arise when several latent states produce the same observable.

No clinical or diagnostic conclusion follows from this analogy.

16 Philosophy and Epistemology

The Ψ construction also illustrates a distinction between representation and explanation.

The defining equation

$$S(\Psi(s)) = s \quad (68)$$

specifies the mathematical object.

The approximation

$$\Psi(s) \approx W_0(s) \quad (69)$$

provides a computational representation.

The residual certificate

$$|S(W_0) - s| \quad (70)$$

provides evidence about the quality of that representation.

Thus three epistemic levels should be distinguished:

$$\text{definition} \neq \text{approximation} \neq \text{verification}. \quad (71)$$

This distinction is important in mathematical modelling because numerical success does not by itself establish a theorem.

17 Religion and Conceptual Metaphor

Religious and philosophical traditions frequently distinguish between an underlying object and its representation, manifestation, or interpretation. The present mathematical framework can be used as a metaphor for that distinction:

$$\text{latent structure} \longrightarrow \text{observable representation} \longrightarrow \text{inverse interpretation}. \quad (72)$$

This is a conceptual analogy rather than a theological proposition. The mathematical function Ψ should not be treated as evidence for any religious doctrine.

18 Political Science, Geopolitics, and International Relations

Political systems can similarly be represented as nonlinear mappings between latent institutional states and observable outcomes.

An abstract model is

$$Y_t = F(X_t, Z_t), \quad (73)$$

where X_t denotes institutional variables and Z_t denotes environmental or strategic conditions.

When the mapping is identifiable, inverse analysis takes the form

$$X_t = F^{-1}(Y_t, Z_t). \quad (74)$$

Such a framework can be applied descriptively to institutional analysis, diplomacy, international relations, strategic interaction, and geopolitical systems.

It does not determine political preferences or prescribe political choices. Its role is analytical: to clarify mappings, assumptions, identification, and uncertainty.

19 Welfare Economics and Warfare Economics

The same mathematical architecture can represent a switching system.

Let

$$q_t \in \{D, W\} \quad (75)$$

denote developmental and warfare-economic regimes. An abstract state transition can be written as

$$q_{t+1} = \mathcal{T}(q_t, X_t, \varepsilon_t). \quad (76)$$

A welfare objective might be

$$\max \mathbb{E}_0 \sum_{t=0}^{\infty} \beta^t U(C_t, G_t, S_t), \quad (77)$$

where C_t represents consumption, G_t productive expenditure, and S_t a security state.

A warfare-economics extension would introduce resource constraints such as

$$Y_t = C_t + G_t + M_t + D_t, \quad (78)$$

where M_t represents security expenditure and D_t represents other defensive or strategic resource uses.

The Ψ framework contributes here as an example of how a nonlinear inverse map can recover an underlying scalar state from an observed aggregate.

No military operational recommendation follows from this abstract model.

20 Artificial Intelligence and Machine Learning

The fixed-point formulation of the original Ψ construction provides a natural connection to equilibrium machine learning.

Suppose

$$z = T_s(z). \quad (79)$$

Then a fixed point satisfies

$$F_s(z) = T_s(z) - z = 0. \quad (80)$$

This structure resembles equilibrium neural networks and implicit layers, where the model output is defined as the solution to an implicit nonlinear system rather than by a finite feed-forward computation [9, 10].

The original Ψ construction gives the scalar example

$$T_s(z) = \frac{c}{\sigma} e^{z+s} - \frac{1}{\sigma} e^{cz+\sigma e^{z+s}}. \quad (81)$$

Newton iteration then supplies

$$z_{n+1} = z_n - \frac{F_s(z_n)}{F'_s(z_n)}. \quad (82)$$

This creates a direct conceptual bridge between transcendental equations, fixed-point computation, implicit machine learning, and numerical optimization.

21 Data Compression, Security, and Computational Systems

The function Ψ can also serve as a compact nonlinear transformation within a larger computational pipeline:

$$\text{raw data} \longrightarrow \text{feature representation} \longrightarrow \Psi \longrightarrow \text{compressed state} \longrightarrow \text{secure transmission}. \quad (83)$$

Such a use would require empirical validation of numerical stability, precision requirements, computational cost, and sensitivity.

The broader data-science perspective is consistent with the standard pipeline of data collection, cleaning, modelling, deployment, and feedback [6, 7].

22 Interdisciplinary Synthesis

The common structure across the disciplines discussed above can be written as

$$\boxed{\text{observable state} \longleftrightarrow \text{nonlinear mapping} \longleftrightarrow \text{latent state}.} \quad (84)$$

The Ψ function provides a mathematically explicit example:

$$s \longleftrightarrow S(W) \longleftrightarrow W = \Psi(s). \quad (85)$$

The interdisciplinary interpretation is summarized in Table 1.

These interpretations should be regarded as modelling possibilities, not as empirical validation.

23 Generalization to Regular n -gons

The original construction suggests a family of functions indexed by the number of polygonal sides. Let

$$\theta_n = \frac{2\pi}{n}, \quad c_n = \cos \theta_n, \quad \sigma_n = \sin \theta_n. \quad (86)$$

The generalized scalar parametrization is

$$s_n(W) = \ln(c_n - \sigma_n W e^W) + (c_n - 1)W + \sigma_n e^{-W}. \quad (87)$$

Table 1: Potential interdisciplinary interpretations of the inverse-map structure.

Field	Observable	Latent or recovered quantity
Mathematics	s	$W = \Psi(s)$
Statistics	sample statistic	model parameter
Econometrics	equilibrium outcome	structural state
Finance	market observable	latent nonlinear factor
Economics	aggregate outcome	structural regime
Physics	measured quantity	physical state
Chemistry	equilibrium observable	reaction parameter
Biology	population signal	latent system state
Psychology	behavior	latent state model
Computer science	output	implicit computational state
AI/ML	network output	equilibrium representation
Political science	institutional outcome	latent configuration
Geopolitics	strategic observable	system state
Welfare economics	welfare outcome	allocation state
Warfare economics	security output	resource regime

Define

$$\Psi_n(s) = s_n^{-1}(s). \quad (88)$$

The pentagonal function is then the special case

$$\Psi = \Psi_5. \quad (89)$$

This produces a hierarchy

$$\{\Psi_3, \Psi_4, \Psi_5, \dots\}, \quad (90)$$

opening a broader research programme in special-function theory [1].

24 Research Programme

The remaining mathematical work can be organized into four stages.

1. Establish rigorous interval bounds for the Padé residual on $1 \leq s \leq 2$.
2. Establish rigorous interval bounds for the logarithmic-rational residual on $2 < s < s_*$.
3. Establish an analytic asymptotic remainder bound for W_{AS3} on $s \geq s_*$.
4. Combine the three bounds with the derivative inequality

$$|S'(W)| > 1 - c + \sigma \quad (91)$$

to obtain a global certified approximation theorem.

A further direction is the construction of arbitrary-order coefficients $A_k(L)$ in

$$\Psi(s) = -\ln \frac{s}{\sigma} + \sum_{k=1}^N \frac{A_k(\ln(s/\sigma))}{s^k} + R_N(s). \quad (92)$$

This would provide a systematic asymptotic library for Ψ .

25 Conclusion

The $\Psi(s)$ function can be viewed simultaneously as a geometric inverse, a transcendental special function, an implicit numerical problem, and a prototype inverse map for nonlinear systems.

The central computational result is the piecewise representation

$$W_0(s) = \begin{cases} P_{5,5}(s), & 1 \leq s \leq 2, \\ W_{LR}(s), & 2 < s < s_*, \\ W_{AS3}(s), & s \geq s_*, \end{cases} \quad (93)$$

followed by

$$W_1(s) = W_0(s) - \frac{S(W_0(s)) - s}{S'(W_0(s))}. \quad (94)$$

The residual

$$R_1(s) = S(W_1(s)) - s \quad (95)$$

then gives the practical error certificate

$$|\Psi(s) - W_1(s)| \leq \frac{|R_1(s)|}{1 - c + \sigma}. \quad (96)$$

High-precision numerical experiments indicate an error scale of roughly 10^{-10} after one Newton correction over the tested domain. A fully rigorous global bound remains a matter of validated interval computation, not of further curve fitting.

The broader significance is therefore methodological. A newly defined inverse transcendental function does not require an elementary closed form to become mathematically useful. An exact definition, asymptotic theory, explicit approximations, numerical algorithms, and rigorous residual certificates together constitute a viable theory of computation for the function.

References

- [1] S. Ghosh, *The $\Psi(s)$ Function and Its Applications*, Kolkata, India.
- [2] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth, “On the Lambert W Function,” *Advances in Computational Mathematics*, Vol. 5, 1996, pp. 329–359.
- [3] N. G. de Bruijn, *Asymptotic Methods in Analysis*, Dover Publications.
- [4] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes*, Cambridge University Press.
- [5] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, eds., *NIST Handbook of Mathematical Functions*, Cambridge University Press.
- [6] S. Ghosh, *Introspection and Retrospection on the Age of Data Science, Neural Networks and AI*, Kolkata, India.
- [7] S. Ghosh, *The State-of-the-Art in Data Compression and Data Security circa 2025*, Kolkata, India.
- [8] I. Goodfellow, Y. Bengio, and A. Courville, *Deep Learning*, MIT Press.
- [9] Z. Bai, J. Kolter, and V. Koltun, “Deep Equilibrium Models,” *Advances in Neural Information Processing Systems*, 2019.
- [10] S. Bai, J. Z. Kolter, and V. Koltun, “Multiscale Deep Equilibrium Models,” arXiv preprint.
- [11] S. Ghosh, *Non-Trivial Functional Solutions to Ghosh’s M Measure*, Kolkata, India.
- [12] S. Ghosh, *The Implications of Ghosh’s M Measure for Macroeconomics*, Kolkata, India.
- [13] S. Ghosh, *Optimal Monetary Policy with Ghosh’s M Measure*, Kolkata, India.
- [14] S. Ghosh, *Optimal Monetary Policy with Ghosh’s M Measure in Open Economies*, Kolkata, India.

Glossary

$\Psi(s)$

The inverse transcendental function defined by $S(\Psi(s)) = s$.

$S(W)$

The scalar parametrization

$$S(W) = \ln(c - \sigma W e^W) + (c - 1)W + \sigma e^{-W}. \quad (97)$$

Inverse function

A function that recovers the input of a one-to-one mapping.

Residual

The discrepancy

$$R(s) = S(\hat{\Psi}(s)) - s. \quad (98)$$

Padé approximation

A rational approximation constructed to reproduce local series information.

Asymptotic expansion

An expansion whose successive terms describe behavior in a limiting regime, here $s \rightarrow \infty$.

Newton refinement

An iterative root-finding procedure based on the local derivative.

Implicit model

A model in which a state is defined through an equation rather than an explicit forward formula.

Residual certificate

An error bound obtained by combining a computable residual with a lower bound on the derivative.

Validated numerics

Numerical computation in which rounding and approximation errors are controlled sufficiently to produce mathematically certified enclosures.

Latent state

An unobserved quantity inferred from observable variables through a model.

Equilibrium neural network

A neural architecture whose state is defined implicitly as a fixed point.

Structural identification

Recovery of model parameters or latent states from observable implications under specified assumptions.

Welfare economics

The study of economic allocations and their consequences for social welfare.

Warfare economics

The study of economic resource allocation under security, conflict, or war-related constraints.

The End