

alphafold analysis

shivani

```
library(bio3d)
```

PDB File names of models

```
files <- list.files("hiv_monomer_94b5b/", pattern = ".pdb",  
  full.names=T)
```

Align and superpose

```
pdbbs <- pdbaln(files, fit=TRUE, exefile="msa")
```

Reading PDB files:

```
hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_001_alphafold2_ptm_model_5_seed_000.pdb  
hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_002_alphafold2_ptm_model_1_seed_000.pdb  
hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_003_alphafold2_ptm_model_4_seed_000.pdb  
hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_004_alphafold2_ptm_model_3_seed_000.pdb  
hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_005_alphafold2_ptm_model_2_seed_000.pdb  
.....
```

Extracting sequences

```
pdb/seq: 1   name: hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_001_alphafold2_ptm_mo  
pdb/seq: 2   name: hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_002_alphafold2_ptm_mo  
pdb/seq: 3   name: hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_003_alphafold2_ptm_mo  
pdb/seq: 4   name: hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_004_alphafold2_ptm_mo  
pdb/seq: 5   name: hiv_monomer_94b5b//hiv_monomer_94b5b_unrelaxed_rank_005_alphafold2_ptm_mo
```

RMSD analysis

```
rd <- rmsd(pdbbs)
```

Warning in rmsd(pdbbs): No indices provided, using the 99 non NA positions


```
mean(rd)
```

```
[1] 0.54368
```

```
source("http://tinyurl.com/newviewngl")  
library(NGLViewerR)  
#view.pdbbs(pdbbs)
```

#PCA

```
pc <- pca(pdbbs)  
plot(pc)
```



Residue conservaation from alignment file

```
aln_file <- list.files(path="hiv_monomer_94b5b/",
                      pattern = ".a3m$",
                      full.names = TRUE)
aln_file
```

```
[1] "hiv_monomer_94b5b//hiv_monomer_94b5b.a3m"
```

```
aln <- read.fasta(aln_file[1], to.upper = TRUE)
```

```
[1] " ** Duplicated sequence id's: 101 **"
```

```
dim(aln$ali)
```

```
[1] 5378 132
```

score residue conservation:


```
sim <- conserv(aln)
```

```
con <- consensus(aln, cutoff = 0.9)
con$seq
```

```
[1] "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-"
[19] "-" "-" "-" "-" "-" "-" "D" "T" "G" "A" "-" "-" "-" "-" "-" "-" "-" "-"
[37] "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-"
[55] "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-"
[73] "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-"
[91] "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-"
[109] "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-" "-"
[127] "-" "-" "-" "-" "-" "-"
```

plot the conservation along

```
plot(sim, typ="h")
```



conserved positions in structure:

```
pdb <- read.pdb(files[1])  
#view.pdb(pdb, backgroundColor="pink", highlight=atom.select(pdb, resno=25:28), highlight.styl
```

Instead of HIV Dimer analysis, Barry let us analyze our own novel protein sequence in Alphafold