

unite: Unified liNe Integration Turbo Engine

Raphael Erik Hviding¹, Anna de Graaff^{2,1}, Alberto Torralba³,
Helena Treiber⁴, and Morgan Fouesneau¹

¹ Max-Planck-Institut für Astronomie ² Center for Astrophysics Harvard & Smithsonian ³ Institute of Science and Technology Austria ⁴ Princeton University ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#)).

Summary

Astronomical spectroscopy, whereby light from celestial sources is dispersed into its constituent wavelengths (colors), is a cornerstone of modern astrophysical research. In particular the characterization of spectral lines, which arise from atomic and molecular transitions in astrophysical gas, allows astronomers to measure fundamental physical properties such as redshift, chemical composition, temperature, density, and kinematics of the emitting or absorbing material. Therefore, the flexibility, speed, and accuracy of spectral line-fitting tools directly impact the scientific return of spectroscopic observations.

unite (Unified liNe Integration Turbo Engine) is a Python package for fast and accurate Bayesian inference of spectral line features from one or more spectra simultaneously. It is built primarily on JAX ([Bradbury et al., 2018](#)) (for speed and automatic differentiation), NumPyro ([Bingham et al., 2019](#); [Phan et al., 2019](#)) (for probabilistic programming), and Astropy ([Astropy Collaboration et al., 2013, 2018, 2022](#)) (for units and FITS handling), and is designed to be flexible and extensible to a wide range of spectroscopic applications.

Statement of need

The Near-Infrared Spectrograph (NIRSpec; [Jakobsen et al. \(2022\)](#)) on the James Webb Space Telescope (JWST; [Gardner et al. \(2023\)](#)) is the frontier instrument for near-infrared spectroscopy and has already provided revolutionary insights across a wide range of astrophysical topics, from the characterization of exoplanet atmospheres to the discovery of the most distant galaxies known. However, a defining challenge of NIRSpec spectroscopy is that the detector critically undersamples the instrumental line spread function (LSF) across all gratings and observing modes. Therefore, evaluating the model at pixel centers rather than integrating over the pixel domains introduces systematic errors in recovered fluxes and line widths, which can bias scientific inferences. When existing spectral line fitting tools can account for this, they typically do so via computationally and memory expensive super-sampling and convolution. In addition, typical JWST/NIRSpec programs observe targets with multiple gratings to leverage their complementary strengths (i.e. high-resolution to measure kinematics, low-resolution to constrain fluxes and continuum shapes), so a statistically optimal analysis must fit all gratings simultaneously with shared astrophysical parameters and grating-specific calibration offsets. unite is designed to address both of these challenges while additionally providing a flexible, extensible framework for spectroscopic analysis that can be applied to any instrument with user-supplied resolving power and pixel-scale functions. By relying on existing optimized libraries for probabilistic programming and automatic differentiation, unite delivers scientifically rigorous Bayesian inference without compromising on speed, enabling the analysis of large spectroscopic datasets with accurate uncertainty quantification. The target audience is observational astronomers, especially those working with JWST/NIRSpec data, that want to do inference over large spectroscopic samples, deal with undersampled data, and/or fit

multiple spectra simultaneously with shared parameters.

Software design

When fitting spectroscopic data sets, fitting routines typically assume that the model evaluated at the pixel center is a good representation of the average of the model over the pixel domain, which is what the instrument actually measures. This approximation is well justified when the spectrum is critically sampled or over-sampled, i.e. when the signal changes slowly over the pixel domain, but breaks down when the spectrum is undersampled and the signal changes rapidly over the pixel domain. In the case for NIRSpec, which for a point source is undersampled by a factor of ~ 1.5 (de Graaff et al., 2024), this can lead to a bias of 10 – 20% in recovered line widths if not properly accounted for. This can be addressed integrating the model over the pixel domain, providing the exact solution for the observed signal regardless of the degree of undersampling. We rely on the assumption that the LSF is well approximated by a Gaussian kernel, which is a good approximation for NIRSpec (Shajib et al., 2025) and many other spectrographs, especially in the undersampled regime.

unite computes the integrals of continua and line models analytically where possible. However, analytic pixel integration is not possible for all model setups, in particular in the presence of optical-depth parametrized absorption lines where the nonlinear transmission $e^{-\tau\phi}$ couples the line depth and profile shape in a way that prevents closed-form solutions for the pixel integrals. In these cases, unite provides a numerical convolution mode which supersamples the intrinsic model on a fine wavelength grid and convolves with the wavelength-dependent LSF kernel to produce a pixel-convolved model. This model accurately captures the nonlinear coupling of line depth and profile shape, but is more computationally expensive than the analytic integration mode; users can choose the appropriate mode for their application.

Despite implementing integration, quadrature, and convolution modes, unite is fast and efficient thanks to its JAX backend, which provides just-in-time (JIT) compilation and native GPU support. At its core, unite is a domain-specific language for building probabilistic models of spectroscopic data. Users build a declarative configuration of line and continuum components and assign priors to physical parameters via token instances which can be shared across multiple model components with arithmetic combinations. The prior system is expressive: priors on any parameter can depend on other parameters through arithmetic expressions and topologically sorted dependency chains, enabling physical constraints such as requiring a broad component's velocity width to exceed that of a narrow component by a certain amount, or fixing flux ratios between doublet lines.

In addition, users specify the instrumental configuration carrying empirical calibrations of the wavelength-dependent resolving power, pixel scale, and flux normalization for each disperser, which can be shared across instruments. One aspect that sets unite apart from other spectral fitting tools is that it treats instrumental calibration parameters as first-class citizens in the inference process; pixel offsets, resolution scales, and flux normalizations can be directly incorporated into the model with priors and sampled jointly with astrophysical parameters, allowing for instrumental uncertainties to directly propagate to the inferred properties. For example, when fitting NIRSpec data, users can fold in the empirically observed offsets in wavelength solutions and flux scales derived in de Graaff, Brammer, et al. (2025) to obtain realistic uncertainties on line fluxes and kinematics that account for systematic uncertainties in the instrument calibration.

All configurations are serializable to human-readable YAML for reproducibility and sharing. unite assembles a NumPyro probabilistic model for inference with any compatible sampler, including SVI for quick exploratory fits, NUTS for full posterior sampling, and nested sampling for model comparison and evidence calculation. Finally, unite provides convenience functions for extracting results as parameter tables and per-spectrum model predictions into domain-appropriate FITS HDU lists.

93 The package is publicly available on GitHub and PyPI under the GPL-3.0-or-later license,
94 with a DOI minted via Zenodo (Hviding, 2026) and accompanied by Sphinx documentation
95 including narrative guides, an API reference, and executable tutorials. CI/CD workflows ensure
96 that the code is tested and documented with each update, and the project is open to feedback
97 and contributions from the community.

98 State of the field

99 Spectral line analysis is among the most common operations in observational astronomy, and
100 the landscape of fitting software is correspondingly rich.

101 The dominant paradigm for emission-line fitting is non-linear least squares, with most Python
102 codes built on LMFIT (Newville et al., 2014) or `scipy.optimize` (Virtanen et al., 2020) directly.

103 pPXF (Cappellari, 2017, 2023; Cappellari & Emsellem, 2004) is the standard tool for stellar
104 kinematics and stellar-population recovery, fitting observed galaxy spectra as non-negative linear
105 combinations of SSP templates and emission lines convolved with a parametric line-of-sight
106 velocity distribution with optional supersampling.

107 PySpecKit (Ginsburg et al., 2022) is a general-purpose spectral fitting toolkit supporting a
108 range of profile shapes and optional MCMC uncertainty estimation.

109 LiMe (Fernández et al., 2024) is a modern library designed for large and complex datasets
110 including JWST spectra, with batch fitting across many spectra and flexible profile shapes.

111 Q3DFIT (Rupke, 2014; Rupke et al., 2021, 2023) targets IFU spectroscopy of quasar host
112 galaxies, fitting multi-component Gaussian profiles via least squares with the stellar continuum
113 pre-subtracted using pPXF.

114 BADASS (Sexton et al., 2021, 2024) is a comprehensive Bayesian emission-line code to
115 simultaneously infer AGN power-law continuum, FeII pseudo-continuum, stellar LOSVD (via
116 pPXF), and multi-component Gaussian emission lines in a single posterior.

117 These codes occupy various niches but none provide analytic pixel integration or first-class
118 joint multi-grating fitting. Many of these codes provide either least-squares optimization,
119 which does not provide posterior distributions, or Bayesian inference via `emcee` (Foreman-
120 Mackey et al., 2013) or `pymc` (Abril-Pla et al., 2023). `unite` is more closely comparable to
121 the latter category of Bayesian tools and builds natively on JAX and NumPyro, inheriting JIT
122 compilation, automatic differentiation, and GPU support, enabling scalable inference across
123 large spectroscopic samples.

124 Research impact statement

125 `unite` has already been used in several published JWST/NIRSpec analyses, demonstrating its
126 utility for emission line characterization.

- 127 ■ **Accurate Line Fluxes and Kinematics:** With the accurate accounting for undersampling,
128 `unite` has been used to robustly characterize fluxes and kinematics for both emission
129 lines and absorption features in high-redshift galaxies, providing insights into the physical
130 processes driving star formation and black hole growth. (de Graaff, Hviding, et al., 2025;
131 Naidu et al., 2025; Sun et al., 2026; Wang et al., 2025, 2026)
- 132 ■ **Multi-Component Line Fitting:** `unite` has been used to identify broad Balmer emission
133 (broad $H\alpha$, $H\beta$) in high-redshift ($z \gtrsim 4$) galaxies observed with NIRSpec, providing
134 evidence for active supermassive black hole growth in the early universe. Multi-grating
135 joint fitting is critical as grating spectroscopy often lacks in signal-to-noise but by coupling
136 physical constraints on flux from low-resolution gratings, `unite` enables robust detections
137 of these components. (de Graaff, Hviding, et al., 2025; Hviding et al., 2025, 2026)

- **Redshift Precision:** unite was used in the analysis of MoM-z14, which at the time of publication was (and remains) the most distant spectroscopically confirmed galaxy known. Coupling line parameters and properly accounting for undersampling allowed for a factor of 5 improvement in redshift precision than from continuum estimates alone even with faint, marginally detected, emission lines. (Naidu et al., 2026)

AI usage disclosure

Claude Code (Anthropic) was used as an AI coding assistant during the development of unite, its documentation, and the drafting of this paper. It was used to generate new code, refactor existing code, and write certain sections of the documentation. All scientific decisions, algorithmic design choices, and final content were made and verified by the authors.

Acknowledgements

REH acknowledges support by the German Aerospace Center (DLR) and the Federal Ministry for Economic Affairs and Energy (BMWi) through program 50OR2403 'RUBIES'. AdG acknowledges support from a Clay Fellowship awarded by the Smithsonian Astrophysical Observatory.

We acknowledge the support of the Data Science Group at the Max Planck Institute for Astronomy (MPIA) and especially Morgan Fouesneau and Ivelina Momcheva for their invaluable assistance providing high level feedback for the development of the core unite routines.

References

- Abril-Pla, O., Andreani, V., Carroll, C., Dong, L., Fonnesbeck, C. J., Kochurov, M., Kumar, R., Lao, J., Luhmann, C. C., Martin, O. A., Osthege, M., Vieira, R., Wiecki, T., & Zinkov, R. (2023). PyMC: A modern and comprehensive probabilistic programming framework in Python. *PeerJ Computer Science*, 9(e1516). <https://doi.org/10.7717/peerj-cs.1516>
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., Earl, N., Starkman, N., Bradley, L., Shupe, D. L., Patil, A. A., Corrales, L., Brasseur, C. E., Nöthe, M., Donath, A., Tollerud, E., Morris, B. M., Ginsburg, A., Vaher, E., Weaver, B. A., Tocknell, J., Jamieson, W., ... Astropy Project Contributors. (2022). The Astropy Project: Sustaining and Growing a Community-oriented Open-source Project and the Latest Major Release (v5.0) of the Core Package. 935(2), 167. <https://doi.org/10.3847/1538-4357/ac7c74>
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., Lim, P. L., Crawford, S. M., Conseil, S., Shupe, D. L., Craig, M. W., Dencheva, N., Ginsburg, A., VanderPlas, J. T., Bradley, L. D., Pérez-Suárez, D., de Val-Borro, M., Paper Contributors, (., Aldcroft, T. L., Cruz, K. L., Robitaille, T. P., ... Contributors, (. (2018). The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package. 156, 123. <https://doi.org/10.3847/1538-3881/aabc4f>
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., Greenfield, P., Droettboom, M., Bray, E., Aldcroft, T., Davis, M., Ginsburg, A., Price-Whelan, A. M., Kerzendorf, W. E., Conley, A., Crighton, N., Barbary, K., Muna, D., Ferguson, H., Grollier, F., Parikh, M. M., Nair, P. H., ... Streicher, O. (2013). Astropy: A community Python package for astronomy. 558, A33. <https://doi.org/10.1051/0004-6361/201322068>
- Bingham, E., Chen, J. P., Jankowiak, M., Obermeyer, F., Pradhan, N., Karaletsos, T., Singh, R., Szerlip, P. A., Horsfall, P., & Goodman, N. D. (2019). Pyro: Deep universal probabilistic programming. *J. Mach. Learn. Res.*, 20, 28:1–28:6. <http://jmlr.org/papers/v20/18-403.html>

- 181 Bradbury, J., Frostig, R., Hawkins, P., Johnson, M. J., Katariya, Y., Leary, C., Maclaurin,
182 D., Necula, G., Paszke, A., VanderPlas, J., Wanderman-Milne, S., & Zhang, Q. (2018).
183 JAX: Composable transformations of Python+NumPy programs (Version 0.3.13). <http://github.com/jax-ml/jax>
184
- 185 Cappellari, M. (2017). Improving the full spectrum fitting method: accurate convolution with
186 Gauss-Hermite functions. *MNRAS*, 466(1), 798–811. [https://doi.org/10.1093/mnras/](https://doi.org/10.1093/mnras/stw3020)
187 [stw3020](https://doi.org/10.1093/mnras/stw3020)
- 188 Cappellari, M. (2023). Full spectrum fitting with photometry in PPXF: stellar population
189 versus dynamical masses, non-parametric star formation history and metallicity for 3200
190 LEGA-C galaxies at redshift $z \approx 0.8$. *MNRAS*, 526(3), 3273–3300. [https://doi.org/10.](https://doi.org/10.1093/mnras/stad2597)
191 [1093/mnras/stad2597](https://doi.org/10.1093/mnras/stad2597)
- 192 Cappellari, M., & Emsellem, E. (2004). Parametric Recovery of Line-of-Sight Velocity
193 Distributions from Absorption-Line Spectra of Galaxies via Penalized Likelihood. *PASP*,
194 116(816), 138–147. <https://doi.org/10.1086/381875>
- 195 de Graaff, A., Brammer, G., Weibel, A., Lewis, Z., Maseda, M. V., Oesch, P. A., Bezanson, R.,
196 Boogaard, L. A., Cleri, N. J., Cooper, O. R., Gottumukkala, R., Greene, J. E., Hirschmann,
197 M., Hviding, R. E., Katz, H., Labbé, I., Leja, J., Matthee, J., McConachie, I., ... Williams,
198 C. C. (2025). RUBIES: A complete census of the bright and red distant Universe with
199 JWST/NIRSpec. *A&A*, 697, A189. <https://doi.org/10.1051/0004-6361/202452186>
- 200 de Graaff, A., Hviding, R. E., Naidu, R. P., Greene, J. E., Miller, T. B., Leja, J., Matthee,
201 J., Brammer, G., Katz, H., Bezanson, R., Boogaard, L. A., Bose, S., Chisholm, J.,
202 Cleri, N. J., Dayal, P., Feldmann, R., Fudamoto, Y., Fujimoto, S., Furtak, L. J., ...
203 van der Wel, A. (2025). Little Red Dots host Black Hole Stars: A unified family of
204 gas-reddened AGN revealed by JWST/NIRSpec spectroscopy. *arXiv*, arXiv:2511.21820.
205 <https://doi.org/10.48550/arXiv.2511.21820>
- 206 de Graaff, A., Rix, H.-W., Carniani, S., Suess, K. A., Charlot, S., Curtis-Lake, E., Arribas,
207 S., Baker, W. M., Boyett, K., Bunker, A. J., Cameron, A. J., Chevillard, J., Curti,
208 M., Eisenstein, D. J., Franx, M., Hainline, K., Hausen, R., Ji, Z., Johnson, B. D., ...
209 Willott, C. (2024). Ionised gas kinematics and dynamical masses of $z \gtrsim 6$ galaxies from
210 JADES/NIRSpec high-resolution spectroscopy. *A&A*, 684, A87. [https://doi.org/10.1051/](https://doi.org/10.1051/0004-6361/202347755)
211 [0004-6361/202347755](https://doi.org/10.1051/0004-6361/202347755)
- 212 Fernández, V., Amorín, R., Firpo, V., & Morisset, C. (2024). LIME: A Line MEasuring library
213 for large and complex spectroscopic data sets. I. Implementation of a virtual observatory
214 for JWST spectra. *A&A*, 688, A69. <https://doi.org/10.1051/0004-6361/202449224>
- 215 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. (2013). emcee: The MCMC
216 Hammer. *PASP*, 125(925), 306. <https://doi.org/10.1086/670067>
- 217 Gardner, J. P., Mather, J. C., Abbott, R., Abell, J. S., Abernathy, M., Abney, F. E., Abraham,
218 J. G., Abraham, R., Abul-Huda, Y. M., Acton, S., Adams, C. K., Adams, E., Adler, D.
219 S., Adriaensen, M., Aguilar, J. A., Ahmed, M., Ahmed, N. S., Ahmed, T., Albat, R., ...
220 Dellagatta, G. (2023). The James Webb Space Telescope Mission. *PASP*, 135(1048),
221 068001. <https://doi.org/10.1088/1538-3873/acd1b5>
- 222 Ginsburg, A., Sokolov, V., de Val-Borro, M., Rosolowsky, E., Pineda, J. E., Sipőcz, B. M., &
223 Henshaw, J. D. (2022). Pyspeckit: A Spectroscopic Analysis and Plotting Package. *AJ*,
224 163(6), 291. <https://doi.org/10.3847/1538-3881/ac695a>
- 225 Hviding, R. E. (2026). *Unite: Unified liNe integration turbo engine* (Version v2.9.2). Zenodo.
226 <https://doi.org/10.5281/zenodo.20158737>
- 227 Hviding, R. E., de Graaff, A., Liu, H., Goulding, A. D., Ma, Y., Greene, J. E., Boogaard, L.
228 A., Bunker, A. J., Cleri, N. J., Franx, M., Hirschmann, M., Leja, J., Matthee, J., Naidu,
229 R. P., Setton, D. J., Übler, H., Venturi, G., & Wang, B. (2026). The X-Ray Dot: Exotic

- 230 Dust or a Late-stage Little Red Dot? *ApJL*, 1000(1), L18. [https://doi.org/10.3847/2041-](https://doi.org/10.3847/2041-231 8213/ae4c88)
- 232 Hviding, R. E., de Graaff, A., Miller, T. B., Setton, D. J., Greene, J. E., Labbé, I., Brammer,
233 G., Bezanson, R., Boogaard, L. A., Cleri, N. J., Leja, J., Maseda, M. V., McConachie, I.,
234 Matthee, J., Naidu, R. P., Oesch, P. A., Wang, B., Whitaker, K. E., & Williams, C. C.
235 (2025). RUBIES: A spectroscopic census of little red dots: All point sources with v-shaped
236 continua have broad lines. *A&A*, 702, A57. <https://doi.org/10.1051/0004-6361/202555816>
- 237 Jakobsen, P., Ferruit, P., Alves de Oliveira, C., Arribas, S., Bagnasco, G., Barho, R., Beck, T. L.,
238 Birkmann, S., Böker, T., Bunker, A. J., Charlot, S., de Jong, P., de Marchi, G., Ehrenwinkler,
239 R., Falcolini, M., Fels, R., Franx, M., Franz, D., Funke, M., ... Zincke, C. (2022). The
240 Near-Infrared Spectrograph (NIRSpec) on the James Webb Space Telescope. I. Overview
241 of the instrument and its capabilities. *A&A*, 661, A80. [https://doi.org/10.1051/0004-](https://doi.org/10.1051/0004-242 6361/202142663)
- 243 Naidu, R. P., Matthee, J., Katz, H., de Graaff, A., Oesch, P., Smith, A., Greene, J. E.,
244 Brammer, G., Weibel, A., Hviding, R., Chisholm, J., Labbé, I., Simcoe, R. A., Witten,
245 C., Atek, H., Baggen, J. F. W., Belli, S., Bezanson, R., Boogaard, L. A., ... Watson, D.
246 (2025). A “Black Hole Star” Reveals the Remarkable Gas-Enshrouded Hearts of the Little
247 Red Dots. *arXiv*, arXiv:2503.16596. <https://doi.org/10.48550/arXiv.2503.16596>
- 248 Naidu, R. P., Oesch, P. A., Brammer, G., Weibel, A., Li, Y., Matthee, J., Chisholm, J., Pollock,
249 C. L., Heintz, K. E., Johnson, B. D., Shen, X., Hviding, R. E., Leja, J., Tacchella, S.,
250 Ganguly, A., Witten, C., Atek, H., Belli, S., Bose, S., ... Whitaker, K. E. (2026). A Cosmic
251 Miracle: A Remarkably Luminous Galaxy at $z_{\text{spec}} = 14.44$ Confirmed with JWST. *OJAp*,
252 9, 56033. <https://doi.org/10.33232/001c.156033>
- 253 Newville, M., Stensitzki, T., Allen, D. B., & Ingargiola, A. (2014). *LMFIT: Non-linear*
254 *least-square minimization and curve-fitting for python* (Version 0.8.0). Zenodo. <https://doi.org/10.5281/zenodo.11813>
- 256 Phan, D., Pradhan, N., & Jankowiak, M. (2019). Composable effects for flexible and accelerated
257 probabilistic programming in NumPyro. *arXiv Preprint arXiv:1912.11554*.
- 258 Rupke, D. S. N. (2014). *IFSFIT: Spectral Fitting for Integral Field Spectrographs* (p.
259 ascl:1409.005).
- 260 Rupke, D. S. N., Schweitzer, M., Viola, V., Lutz, D., Sturm, E., Spoon, H., Veilleux, S., &
261 Kim, D.-C. (2021). *QUESTFIT: Fitter for mid-infrared galaxy spectra* (p. ascl:2112.002).
- 262 Rupke, D. S. N., Wylezalek, D., Zakamska, N., Veilleux, S., Vayner, A., Bertemes, C., Ishikawa,
263 Y., Liu, W., Lim, H. X. G., Murphree, G., Whitesell, L., McCrory, R., & Anicetti, C. (2023).
264 *q3dfit: PSF decomposition and spectral analysis for JWST-IFU spectroscopy*. Astrophysics
265 Source Code Library, record ascl:2310.004.
- 266 Sexton, R. O., Doan, S. M., Reefe, M., & Matzko, W. (2024). *BADASS: Bayesian AGN*
267 *Decomposition Analysis for SDSS Spectra*. Astrophysics Source Code Library, record
268 ascl:2412.004.
- 269 Sexton, R. O., Matzko, W., Darden, N., Canalizo, G., & Gorjian, V. (2021). Bayesian
270 AGN Decomposition Analysis for SDSS spectra: a correlation analysis of [O III] $\lambda 5007$
271 outflow kinematics with AGN and host galaxy properties. *MNRAS*, 500(3), 2871–2895.
272 <https://doi.org/10.1093/mnras/staa3278>
- 273 Shajib, A. J., Treu, T., Melo, A., Roberts-Borsani, G., Knabel, S., Cappellari, M., & Frieman, J.
274 A. (2025). An accurate measurement of the spectral resolution of the JWST Near Infrared
275 Spectrograph. *A&A*, 702, L12. <https://doi.org/10.1051/0004-6361/202556281>
- 276 Sun, W. Q., Naidu, R. P., Matthee, J., de Graaff, A., Chisholm, J., Greene, J. E., Oesch,
277 P. A., Torralba, A., Hviding, R. E., Brammer, G., Simcoe, R. A., Bose, S., Bouwens,

- 278 R., Dayal, P., Eilers, A.-C., Fei, Q., Furtak, L. J., Gottumukkala, R., Goulding, A., ...
279 Williams, C. C. (2026). Little Red Dot — Host Galaxy = Black Hole Star: A Gas-
280 Enshrouded Heart at the Center of Every Little Red Dot. *arXiv*, arXiv:2601.20929. <https://doi.org/10.48550/arXiv.2601.20929>
281
- 282 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
283 Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson,
284 J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., ... SciPy
285 1.0 Contributors. (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in
286 Python. *Nature Methods*, 17, 261–272. <https://doi.org/10.1038/s41592-019-0686-2>
- 287 Wang, B., Leja, J., Katz, H., Inayoshi, K., Cleri, N. J., de Graaff, A., Hviding, R. E., van
288 Dokkum, P., Greene, J. E., Labbé, I., Matthee, J., McConachie, I., Naidu, R. P., & Nelson,
289 E. J. (2025). The Missing Hard Photons of Little Red Dots: Their Incident Ionizing Spectra
290 Resemble Massive Stars. *arXiv*, arXiv:2508.18358. [https://doi.org/10.48550/arXiv.2508.](https://doi.org/10.48550/arXiv.2508.18358)
291 [18358](https://doi.org/10.48550/arXiv.2508.18358)
- 292 Wang, B., Leja, J., Labbe, I., Greene, J. E., Liu, H., de Graaff, A., Hviding, R. E., Matthee,
293 J., Quataert, E., Bezanson, R., Boogaard, L. A., Brammer, G., Burgasser, A. J., Chen,
294 Y.-X., Cleri, N. J., Cutler, S. E., Dayal, P., Furtak, L. J., Fujimoto, S., ... Zitrin, A. (2026).
295 Water absorption confirms cool atmospheres in two little red dots. *arXiv*, arXiv:2602.06024.
296 <https://doi.org/10.48550/arXiv.2602.06024>